ARTEM: a method for RNA tertiary motif identification with backbone permutations, and its example application to kink-turn-like motifs

Eugene F. Baulin[1,*], Davyd R. Bohdan[1], Dawid Kowalski[1], Milena Serwatka[1], Julia Świerczyńska[1,2], Zuzanna Żyra[1,3], Janusz M. Bujnicki[1]

1. International Institute of Molecular and Cell Biology in Warsaw, ul. Ks. Trojdena 4, 02-109 Warsaw, Poland;

2. University of Warsaw, Faculty of Biology, ul. Ilji Miecznikowa 1, 02-096 Warsaw, Poland;

3. Medical University of Gdańsk, ul. Marii Skłodowskiej-Curie 3a, 80-210 Gdańsk, Poland;

corresponding author: ebaulin@iimcb.gov.pl, efbaulin@gmail.com

ABSTRACT

The functions of non-coding RNAs are largely defined by their three-dimensional structures. RNA 3D structure is organized hierarchically and consists of recurrent building blocks called tertiary motifs. The computational problem of RNA tertiary motif search remains largely unsolved, as standard approaches are restrained by sequence, interaction network, or backbone topology. We developed the ARTEM superposition algorithm, which is free from these limitations. Here, we present a version of ARTEM that allows automated searches of RNA structure databases to identify 3D structure motifs. We exemplify it by a search of motifs isosteric to the kink-turn motif. This widespread motif plays a role in many aspects of RNA function, and its mutations are known to cause several human syndromes. With ARTEM, we discovered two new kink-turn topologies, multiple no-kink variants of the motif, and showed that a ribosomal junction in bacteria forms either a kink or a no-kink variant depending on the species. Additionally, we identified kink-turns in the catalytic core of group II introns, whose structures have not previously been characterized as containing kink-turns. ARTEM opens a fundamentally new way to study RNA 3D folds and motifs and analyze their correlations and variations.

KEYWORDS

Kink-turn motif, k-junction motif, A-minor junction, RNA tertiary motif, motif search, structure superposition, backbone topology, isostericity

INTRODUCTION

The functions of non-coding RNAs are largely defined by their three-dimensional structures [1]. RNA 3D structure is organized hierarchically and can be seen as a set of building blocks, RNA 3D modules [2]. The recurrent modules that preserve their features in various RNAs across different structural contexts are called RNA tertiary motifs [3-5]. Commonly, RNA tertiary motifs are classified into two major classes: local loop motifs, nested within elements of RNA secondary structure [6], such as kink-turns [7], and long-range motifs formed between distant structural segments, such as D-loop/T-loop interaction motifs [8]. Long-range motifs remain much less explored [9, 10] compared to loop motifs [6, 11, 12].

Among the main reasons for the lag in long-range motif exploration are the limitations of existing computational methods designed for the motif search problem, which is formulated as the task of identifying instances of a given 3D motif within a given RNA 3D structure or a set of structures [11, 13-18]. The commonly used methods are limited by at least one of three major restraints: RNA sequence (types of bases) [14, 15], annotations of interactions (e.g., types of base pairs [19]) [13, 15, 16, 18, 20], and backbone topology (types of loops [21, 22]) [11, 12, 14, 16, 17, 23-25]. Sequence restraints may hide rare non-canonical motif variants, such as the U-minor variant of the A-minor interaction motif [26]. Searching by interaction network relies on pairwise interaction annotations that differ significantly depending on the tool used [10]. The topology restraints pose the greatest limitation, resulting in just a few examples of structural similarities identified to date between motifs of different topologies [10, 26-28]. Although tools for topology-independent RNA 3D structure superposition exist, they are designed for the global RNA 3D structure alignment problem and are not suitable for motif search [29-31]. These limitations cause current approaches to favor loop motifs over long-range motifs and substantially hinder the identification of structural similarities between the motifs formed within different backbone topology contexts.

The kink-turn motif [32], one of the most well-studied RNA tertiary motifs [7, 33-35], exemplifies the consequences of the aforementioned limitations. Kink-turns are typically characterized by a canonical stem (C-stem) followed by a bulge that forms a sharp kink in the backbone and a non-canonical stem (NC-stem) starting with *trans-Sugar-Hoogsteen* (*tSH*) *G-A* base pairs [7] (Figure 1). The kink-turn architecture of an internal loop was discovered in one of the first experimentally resolved ribosomal RNA 3D structures in 2001 [32]. The kink-turn architecture of a three-way junction loop was discovered only in 2011 [27] and later described as the k-junction motif in 2014 [28], despite its multiple instances in the same 3D structures of ribosomal RNAs [28]. It was shown that kink-turns are extremely widespread, with examples in spliceosomal RNAs [36], group I introns [37], and riboswitches [28], playing roles in many aspects of RNA function, such as serving as a protein binding site [38]. Mutations disrupting the

kink-turn structure in the spliceosomal U4atac RNA have been shown to cause various diseases, such as Taybi–Linder syndrome [36] and Roifman syndrome [39]. Although the resemblance between kink-turns and diverse A-minor junctions was briefly mentioned as early as 2011 [27], the variety of discovered kink-turns remains limited to the two mentioned topologies [7].

Recently, we developed ARTEM [10], a method for sequentially unconstrained superposition of RNA 3D modules. Here, we present a new version of the ARTEM approach adapted to the tertiary motif search problem. We demonstrate the practical utility of the algorithm to identify motifs structurally similar to the kink-turn motif. We identified two new kink-turn topologies: a five-way junction and an external loop. We also discovered several kink-turn-like motifs, including A-minor junctions [27] and long-range motifs [10], that preserve the core interactions of a kink-turn but do not involve the kink. Finally, we demonstrate that the 23S rRNA J94/99 junction, previously identified as a k-junction in *H.marismortui* [28], adopts the no-kink junction topology in other bacteria with a slight distortion of the core kink-turn interactions. Additionally, we identified the coordination loop in group II introns as a kink-turn, although group II intron structures have not been previously characterized as containing kink-turns. Using a benchmark set of group II intron structures, we demonstrated that ARTEM outperforms existing motif search tools. ARTEM can be uniformly applied to search for both local and long-range motifs, and its advantages open a fundamentally new way to study RNA 3D folds and motifs, and to analyze their correlations and variations.

RESULTS

We developed ARTEM 2.0 (https://github.com/david-bogdan-r/ARTEM), a new version of the ARTEM algorithm tailored to the RNA tertiary motif search problem. This tool enables automated searches of RNA structure databases against multiple query modules. ARTEM is equipped with a rich set of user-defined parameters to specify particular RNA 3D structure regions of interest, impose restraints on the identified motifs, and save the superimposed matches or entire RNA-containing input complexes in PDB or mmCIF format. ARTEM advances beyond the limitations of the existing tools by exclusively relying on motif isostericity. It applies to any chosen reference motif, without differentiation between loop motifs and long-range motifs. We exemplified its capabilities by searching for kink-turn-like RNA tertiary motifs.

New backbone topology variants of the kink-turn motif

To search for instances of the kink-turn motif, we selected the representatives of its two known backbone topology variants (Figure 1, Table 1): an instance of the kink-turn internal loop from a 23S rRNA [40] (module #1) and an instance of the k-junction variant

from a TPP riboswitch [41] (module #2). We executed ARTEM for the two instances against all RNA-containing entries in the Protein Data Bank [42], see the Methods section for details. The results included 26,564 matches (Supplementary Table S1). We annotated the matches with their backbone topology characteristics to facilitate subsequent analysis.

Table 1. Representative instances of the kink-turn motif variants.

#	PDB entry, chain	Description	Topology	RNA molecule	L1 residue	1n residue	Variant	Figure
1	1FFK chain 0	kink-turn Kt7 reference	internal loop	23S rRNA	G94	A80	kink	1A, 1C
2	3D2G chain A	k-junction reference	3-way junction	TPP riboswitch	C38	A72	kink	1B, 1D
3	7P7T chain A	J4/5	5-way junction	23S rRNA	G48	A117	kink	2A, 2B
4	7PKT chain 8	[J99/101]	external loop	L8 rRNA	A453	A481	kink	3A, 3C
5	7ZHG chain 2	no kink but stem	3-way junction	16S rRNA	G1073	A1032	no-kink	4A, 4C
6	8EUY chain 1	no kink but two stems	5-way junction	28S rRNA	G973	A685	no-kink	4B, 4D
7	6ME0 chain A	no kink but stem	4-way junction	group II intron	C141	A181	no-kink	5A, 5C
8	7PWO chain 1	long-range motif	helix-helix interface	28S rRNA	C1572	G2651	no-kink	5B, 5D
9	4V8O chain BA	J99/101	3-way junction	23S rRNA	G2830	A2883	kink	3B, 3D
10	3CC2 chain 0	J94/99	complex pseudoknotted junction	23S rRNA	G2823	A2914	kink	6A, 6C
11	4V8O chain BA	J94/99	complex pseudoknotted junction	23S rRNA	C2788	A2892	no-kink	6B, 6D

Figure 1. Reference kink-turn motifs. *(A) A representative 3D structure of the canonical Kt7 kink-turn [7], LSU rRNA, PDB entry 1FFK, chain 0. (B) A representative 3D structure of the k-junction [7], TPP riboswitch, PDB entry 3D2G, chain A. The interaction schemes of the (C) kink-turn and (D) the k-junction follow the commonly used nomenclature [7]: the canonical stem (C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues of the kink in purple, the third stem (T-stem) and other stems in teal, and the looped-out residues in dark turquoise. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its number in the chain*

(in bold, left) and its named position in the kink-turn (right). The 3D representations were prepared with ChimeraX (https://www.cgl.ucsf.edu/chimerax/). The 2D representations were prepared with Inkscape (https://inkscape.org/).

Via manual inspection of the results, we identified representative instances of two new backbone topology variants of the kink-turn motif (Table 1): a five-way junction in a 23S rRNA [43] (module #3, Figure 2) and an external loop in a mitochondrial LSU rRNA fragment [44] (module #4, Figure 3A,C). Below, we use the commonly accepted nomenclature of the kink-turn structure [7], see the Methods section for details.

Figure 2. Representative J4/5 five-way k-junction module. *(A) A 3D structure of the J4/5 five-way k-junction, LSU rRNA, PDB entry 7P7T, chain A. (B) A 2D interaction scheme of the junction: the canonical stem (C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues of the kink in purple, and the other stems in teal. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its*

number in the chain (in bold, left) and its named position in the kink-turn (right). The 3D representation was prepared with ChimeraX (https://www.cgl.ucsf.edu/chimerax/). The 2D representation was prepared with Inkscape (https://inkscape.org/).

We mapped the identified five-way junction (module #3) to the *J4/5* region of the 23S rRNA structure. The J4/5 region was previously reported to form a k-junction [28], but it was incorrectly identified as a three-way junction. The interaction network of the J4/5 region is closer to the reference module #1, with a *tSH G-A* base pair in position *3n-3b,* but it has a non-linear matching of residues *1n* and *2n,* resulting in a *tHH 2n-2b* base pair.

The external loop (module #4) was mapped to the *J99/101* region of the 23S rRNA, which had not been previously reported as a kink-turn or a k-junction. The *J99/101* region forms a canonical three-way k-junction architecture in non-mitochondrial 23S rRNAs (module #9, Figure 3B,D). Its interaction network is very close to that of the reference module #2, with one substantial difference: the *2n-2b* base pair is a canonical *cWW* base pair, which is incompatible with the *tSW* interaction formed between residues *-1n* and *2b* in the reference modules.

Figure 3. Representative J99/101 k-junction modules. *(A) A 3D structure of the J99/101 k-junction with an external loop architecture, L8 rRNA fragment, PDB entry 7PKT, chain 8. (B) A 3D structure of the J99/101 k-junction with a three-way junction architecture, LSU rRNA, PDB entry 4V8O, chain BA. The interaction schemes of (C) the external loop and (D) the three-way junction: the canonical stem (C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues of the kink in purple, the*

No-kink variants of the kink-turn motif

We observed matches of several unique backbone topologies that lacked the kink while exhibiting the C-stem/NC-stem arrangement isosteric to that of the kink variants (modules #5-#8, Table 1, Figure 4, Figure 5). Three of these modules (#5-#7) were of different junction loop architectures, all of which can be classified as A-minor junctions, whose resemblance to kink-turns was reported previously [27]. In contrast, module #8 is not a loop but a helix-helix interface, showcasing the long-range kink-turn-like motif for the first time.

The no-kink three-way junction (module #5, Figure 4A,C) and the no-kink five-way junction (module #6, Figure 4B,D) both have a pyrimidine in position *1b* instead of the commonly found guanosine. While this is the only notable difference for module #6, module #5 also features a pyrimidine in position *2n* and a non-linear matching of residues *1b* and *2n,* with *1b* belonging to the non-bulge strand.

The no-kink four-way junction (module #7, Figure 5A,C) is the closest module to the reference variants in terms of interactions. The long-range motif (module #8, Figure 5B,D) is the only match that features a non-adenosine (guanosine) *1n* residue. It also includes a uridine in *syn* conformation at position *2b* that does not interact with the *-1n* residue. Notably, there is a striking 3D structure similarity between modules #7 and #8 (Figure 5A,B). Their four RNA strands are arranged similarly but form canonical base pairs with different partners (see the top two base pairs in Figure 5), resulting in distinct topologies: a four-way junction and a long-range helix-helix interface.

Figure 4. Representative no-kink variants #5 and #6. *(A) A 3D structure of the three-way A-minor junction, 16S rRNA, PDB entry 7ZHG, chain 2. (B) A 3D structure of the five-way A-minor junction, 28S rRNA, PDB entry 8EUY, chain 1. The interaction schemes of (C) the three-way junction and (D) the five-way junction: the canonical stem*

(C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues matching the kink in purple, the looped-out residues in dark turquoise, and the other residues in teal. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its number in the chain (in bold, left) and its named position in the kink-turn (right). The 3D representations were prepared with ChimeraX (https://www.cgl.ucsf.edu/chimerax/). The 2D representations were prepared with Inkscape (https://inkscape.org/).

Figure 5. Representative no-kink variants #7 and #8. (A) A 3D structure of the four-way A-minor junction, group II intron, PDB entry 6ME0, chain A. (B) A 3D structure of the long-range helix-helix interface, 28S rRNA, PDB entry 7PWO, chain 1. The interaction schemes of (C) the four-way junction and (D) the long-range motif: the canonical stem (C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues matching the kink in purple, and the other residues in teal. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its number in the chain (in bold, left) and its named position in the kink-turn (right). The 3D representations were prepared with ChimeraX (https://www.cgl.ucsf.edu/chimerax/). The 2D representations were prepared with Inkscape (https://inkscape.org/).

The identified kink/no-kink switch

We discovered that the *J94/99* region of 23S rRNA, another ribosomal k-junction identified previously in *H.marismortui* [28] (module #10, Figure 6A,C), adopts the no-kink junction topology in other bacteria (module #11, Figure 6B,D) with a slight distortion of the core kink-turn interactions (Figure 6D).

In module #11, Helix 98, absent from the *H.marismortui* 23S rRNA [45], is inserted between residues *L1.1* and *L2* that form the kink in module #10. The *1n* adenosine in the no-kink variant is pushed out of its position by residue *1b* and interacts with residue *L1.1* instead of *L1* (Figure 6D). Also, both modules #10 and #11 formally belong to complex pseudoknotted junction loops, which may be considered yet another backbone topology variant.

Figure 6. Representative J94/99 region modules. *(A) A 3D structure of the J94/99 k-junction in H.marismortui, 23S rRNA, PDB entry 3CC2, chain 0. (B) A 3D structure of the J94/99 no-kink variant, LSU rRNA, PDB entry 4V8O, chain BA. The interaction schemes of (C) the J94/99 k-junction and (D) the no-kink variant: the canonical stem (C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues of the kink in purple, and the other residues in teal. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its number in the chain (in bold, left) and its named position in the kink-turn (right). The 3D representations were prepared with ChimeraX (https://www.cgl.ucsf.edu/chimerax/). The 2D representations were prepared with Inkscape (https://inkscape.org/).*

Conservation analysis of ribosomal k-junctions

We analyzed the sequence conservation of the three k-junctions identified in 23S rRNAs (Figure 7). The conservation patterns differ substantially between the junction regions but align well with the individual characteristics of the motifs.

The *J4/5* region exhibits the largest number of strictly conserved residues, including the *tHS A-G* base pair in position *3b-3n,* which is also characteristic of the canonical Kt7 kink-turn (module #1). The *1n* residue in the *J94/99* region is the least conserved *1n* residue among the junctions, consistent with its distorted position in the presence of *Helix 98* (module #11). Surprisingly, the *2b-2n cWW U-A* base pair is highly conserved in the *J99/101* region, while the *J99/101* C-stem base pairs are the least conserved among the junctions.

Figure 7. Motif logos of the 23S rRNA k-junctions. *The J4/5 junction logo follows the scheme of module #3, PDB entry 7P7T, chain A. The J94/99 junction logo follows the scheme of module #10, PDB entry 3CC2, chain 0. The J99/101 junction logo follows the scheme of module #9, PDB entry 4V8O, chain BA. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its number in the chain (in bold, left) and its named position in the kink-turn (right). The 2D representations were prepared with Inkscape (https://inkscape.org/) and WebLogo (https://weblogo.berkeley.edu/).*

The coordination loop in group II introns is a kink-turn

While manually inspecting the kink-turn matches obtained with ARTEM, we identified a kink-turn of the canonical internal loop architecture in a 3D structure of group IIC intron [46] (Figure 8A). To the best of our knowledge, occurrences of kink-turns in group II intron structures were never previously reported. To close this gap, we selected a set of seven group II introns from the representative set of RNA structures [47] and surveyed them for kink-turns, both manually and using ARTEM and other existing motif search tools [13-18] (Supplementary Table S2), see the Methods section for details.

Figure 8. Group IIC intron coordination loop. *(A) A representative 3D structure of the coordination loop kink-turn, group IIC intron, PDB entry 6ME0, chain A. (B) The interaction scheme of the kink-turn: the canonical stem (C-stem) in gold, the non-canonical stem (NC-stem) in green and gray, the residues of the kink in purple, and the other residues in teal. The base pair representation follows the Leontis-Westhof (LW) classification [19]. The backbone connectivity is depicted with arrows. Each base is marked with its number in the chain (in bold, left) and its named position in the kink-turn (right). The tertiary interactions (Greek letters) and the intron regions are marked according to the conventional naming [46, 48]. (C) ARTEM benchmarking against other tools that identified at least one of the seven kink-turns in the seven representative group II intron structures. The 3D representation was prepared with ChimeraX (https://www.cgl.ucsf.edu/chimerax/). The 2D representation was prepared with Inkscape (https://inkscape.org/). The barchart was prepared with Matplotlib (https://matplotlib.org/).*

We identified seven kink-turn modules in the seven group II intron structures (Table 2). Five of the seven kink-turn modules were found in the functional intron region of domain I (DI) known as the coordination loop [48, 49], present in the structures of group IIB and group IIC introns. As implied by its name, the loop coordinates the formation of the intron's catalytic core by interacting with the Κ extension bulge and the exon binding site 1 (EBS1) loop [48]. Additionally, the coordination loop encompasses the single-residue exon binding site 3 (EBS3). Subsequently, exon-intron recognition is facilitated through interactions with the intron binding sites IBS1 and IBS3 (Figure 8B). The remaining two kink-turns were identified as instances of a single Domain III (DIII) kink-turn module in group IIC introns.

Table 2. Kink-turn characteristics of representative group II intron structures.

PDB entry chain	3IGI chain A	5G2X chain A	6CHR chain A	6ME0 chain A	7UIN chain B	8H2H chain A	8T2S chain B
Intron	group IIC	group IIA	group IIB	group IIC	group IIC	group IIA	group IIC
Coordination loop	C148-C157 G220-G226	G203-U210 A339-C348	A186-C197 G293-U299	A243-C253 G362-U368	C151-C160 G228-G234	G203-U210 A339-C348	C151-C160 G228-G234
L1-1n base pair	U150-A224(syn)		G188-A297	G245-A366	U153-A232		U153-A232
К extension bulge	A134-A144	A185-A201	A171-A182	A228-A239	A137-A147	A185-A201	A137-A147
К extension - coordination loop interactions	tHW A138-U150(L1) stack A137-A151(L2)	stack U195-A344 stack U193-G207	ribose-base A175-U189 (L1.1)	N4-O6 H-bond C232-G245	tHW A141-U153(L1) stack G140-A154(L2)	stack U195-A344 stack U193-G207	tHW A141-U153(L1) stack G140-A154(L2)
σ-σ′ BWE				A771 A248(L3) A769			
δ′-δ base pair	G155-C180		G195-C255	A251-U326	G158-C183		G158-C183
EBS3	A223		G296	C365	A231		A231
IBS3	A7, chain B		C623	DG15, chain B	DT36, chain A		G1
Coordination loop residues exposed to reverse transcriptase				C247(L2) G249(1b) C365(EBS3)	U155(L3) A231(EBS3)		U155(L3) A231(EBS3)
DIII kink-turn					G343-U352 A366-C372		G343-U352 A366-C372
Coordination loop kink-turn was identified by	ARTEM ≥12nt ARTEM ≥11nt		ARTEM ≥11nt	ARTEM ≥12nt ARTEM ≥11nt RNAMotifScanX	ARTEM ≥12nt ARTEM ≥11nt		ARTEM ≥12nt ARTEM ≥11nt
DIII kink-turn was identified by					ARTEM ≥11nt DSSR RNAMotifScanX		DSSR RNAMotifScanX

The kink-turn architecture is crucial for the functioning of the coordination loop (Table 2). The residues *L1, L1.1*, and *L2* interact with the К extension. The *L3* residue forms a base wedged element (BWE) [50] with a Domain IV (*DIV*) loop prior to DNA integration [46]. In the group IIC intron structures, the residues *L2, L3, 1b*, and *EBS3* are exposed to the reverse transcriptase protein. Furthermore, the unusual long-range δ′-δ base pair in position *3b-3n* facilitates the stacking between *EBS3* and *EBS1*. Notably, the two representative group *IIA* intron structures lack *EBS3*, and their coordination loops do not form a kink-turn.

Surprisingly, the coordination loop was never identified as a kink-turn previously. Possibly, the long-range δ′-δ base pair replacing the local *3b-3n* base pair and the looped-out *EBS3* residue hid the motif from identification via sequence- and backbone-restrained kink-turn searches. Among the previously existing motif search tools, only DSSR and RNAMotifScanX were able to identify at least one kink-turn in the seven selected 3D structures. DSSR, performing the sequence-based search, identified only the two DIII kink-turn modules. RNAMotifScanX, searching with the consensus kink-turn interaction network, identified the two DIII kink-turns and a single coordination loop kink-turn. In contrast, ARTEM, searching with a single Kt7 reference module, identified four of the five coordination loop kink-turns with the 12nt threshold on the match size. With the 11nt threshold, ARTEM identified all five coordination loop kink-turns and one of the DIII kink-turns, along with eight false positive matches. Consequently, ARTEM outperformed the existing tools, demonstrating an impressive recall of 57% at 100% precision with the 12nt threshold, and 86% recall at 43% precision with the 11nt threshold (Figure 8C).

DISCUSSION

In this work, we developed ARTEM 2.0, an adaptation of the ARTEM algorithm suitable for tackling the long-standing problem of RNA tertiary motif search. ARTEM relies exclusively on motif isostericity and enables the identification of topology-independent structural similarities between the motifs. Therefore, ARTEM overcomes the "*clear shortcoming in classifying RNA motifs simply based on their secondary structures*" [27]. This method applies to any chosen RNA 3D module, without differentiation between loop motifs and long-range motifs.

We employed ARTEM to explore the landscape of modules isosteric to the kink-turn motif. We identified several new topological kink-turn variants, including a five-way junction and an external loop. We showed that certain motifs preserve the same core interactions of the kink-turn but lack the kink. We demonstrated that the ribosomal region *J94/99* can form either a kink or a no-kink variant of the motif depending on the species, albeit with a distortion in residue arrangement in the no-kink variant. Furthermore, we identified kink-turns with a complex network of long-range base pairs in the catalytic core of group II introns, whose structures were never previously reported to involve kink-turns. Analyzing the identified representative kink-turn modules, we found that their only consensus feature is the *1n* purine (almost always adenosine) forming the type I A-minor interaction [51] with the residue *L1* (and its partner *L1p* if present) and donating its Hoogsten edge for base-pairing with the residue *1b*. Strikingly, a 3D module can even lack this essential feature and still be largely isosteric to kink-turns, as evidenced by module #11 (Figure 6B,D).

A notable limitation of ARTEM is that it may require some fine-tuning based on the particular motif of interest and the desired *FP/FN* ratio. In turn, ARTEM offers high flexibility and enables a rich set of user-defined restraints. Furthermore, ARTEM outperformed existing tools in the search for kink-turns among group II introns, even without the introduction of additional restraints on matches beyond a single match-size threshold (Figure 8C). Nevertheless, we intend to develop a separate tool, tailored for the annotation of the most prevalent RNA tertiary motifs, that can be used by non-experts as a closed user-friendly solution.

ARTEM introduces a fundamentally new approach to investigating RNA 3D folds and motifs and analyzing their correlations and variations. It extends the concept of isostericity from RNA base pairs [52] to more complex RNA tertiary motifs. By employing a purely isosteric-based RNA tertiary motif search, ARTEM promises to unveil numerous hidden similarities among recurrent RNA 3D modules, significantly advancing our understanding of the principles governing RNA 3D structure organization.

METHODS

Definitions

The definitions of RNA secondary structure elements used in this work follow our previous study [10]. A *stem* is a helical region formed by at least two consecutive stacked base pairs, of Watson-Crick G-C/A-U or wobble G-U type, and a *loop* is a set of single-stranded regions confined by a stem. A loop and a stem are called *adjacent* if at least one residue of the loop is a direct neighbor in sequence of a residue of the stem. Two elements (stems or loops) are called *distant* if they are not adjacent and don't share common adjacent elements. An *internal loop* includes two strands and is adjacent to two stems. An *N-way junction loop* involves *N > 2* strands and is adjacent to *N* stems. A loop of any type is called *external* if it includes dangling ends among its strands.

An *RNA 3D module* is loosely defined as a set of interacting residues that can be represented with a connected graph. A module is called *long-range* if it involves at least one pair of residues that belong to distant RNA secondary structure elements. Otherwise, a module is called *local*. An *RNA tertiary motif* is defined as a recurrent RNA 3D module. A module is called a *motif instance* if it's identified as the motif representative. We defined the computational problem of RNA tertiary motif search as the task of identifying instances of a given motif in a query RNA 3D structure having an instance of the motif as the reference.

In this work, we adhere to the established nomenclature of the kink-turn structure [7] and Leontis-Westhof base pair classification [19] (Figure 1). A residue forms a base pair

with another residue via one of its three edges: *Watson-Crick* (*W*, circle), *Hoogsteen* (*H*, square), or *Sugar* (*S*, triangle), in either *cis* (filled) or *trans* (empty) orientation. A canonical kink-turn comprises two helical regions, a canonical stem (C-stem, in gold) and a non-canonical stem (NC-stem, in green and gray), connected by an internal loop. The longer strand of the loop (bulge, in purple) involves the sharp kink in its backbone. The NC-stem starts with two characteristic *trans-Sugar-Hoogsteen* (*tSH*) *G-A* base pairs (in green). The residues of the bulge are labeled as *L1*, *L2*, and *L3*. The base pairs of NC-stem are labeled with positive numbers starting from the base pair adjacent to the bulge, with lowercase *b* (bulge side) or *n* (non-bulge side) letters: *1n/1b* base pair, *2n/2b* base pair, and so on. The C-stem base pairs are labeled with negative numbers: *-1b/-1n* base pair, *-2b/-2n*, and so on. The k-junction variant (Figure 1B, D) involves an additional *L1.1* bulge residue and the third stem (T-stem, in teal). The *L1* residue of the k-junction forms a canonical base pair with the residue that we labeled *L1p*.

We should note that the NC-stem, as defined in [7], deviates from the strict stem definition, with the *tSH G-A* base pairs formally falling within the internal loop. To maintain consistency, we adhere to the conventional NC-stem definition, while employing the strict stem definition for RNA secondary structure annotation.

ARTEM tool improvement

As the original ARTEM tool [10] reports all local structural similarities identified between the two RNA 3D modules, we modified it to make it more suitable for the RNA tertiary motif search. ARTEM 2.0 introduces three new features compared to its predecessor (https://github.com/david-bogdan-r/ARTEM). First, the input structures (both reference and query) can be specified with a folder or mask instead of a single file, allowing one to search against a set of RNA structures with one command. Second, we added a *nosub* option to remove sub-matches (matches involved in larger matches), allowing one to filter out partial hits. Third, we introduced an *rst* option, enabling users to specify additional restraints of four different types: (i) filtering out matches missing specified residues, (ii) requiring the specified residues to match with particular base types (allowing the IUPAC nomenclature [53]), (iii) requiring the specified residues to match under a given RMSD threshold, and (iv) requiring the specified residues to match with a continuous RNA strand. The continuity of an RNA strand is verified by O3′-P atom distances under 2.0 Å between neighboring residues.

Kink-turn motif search with ARTEM

To conduct the kink-turn motif search with ARTEM, we selected one representative for each of the two known backbone topology variants (Figure 1, Table 1): the canonical kink-turn internal loop (LSU rRNA, PDB entry 1FFK [40], chain 0, *L1* residue *G94*) and the k-junction variant (TPP riboswitch, PDB entry 3D2G [41], chain A, *L1* residue *C38*).

For the search, we confined the motif to the following 14 residues: *L1, L1p, -1b, -1n, -2b, -2n, 1b, 1n, 2b, 2n, 3b, 3n, 4b, 4n*. Other labeled residues were excluded due to their dynamic positioning between the kink-turn instances.

To determine the thresholds for the search, we executed ARTEM using the two reference instances as input. ARTEM identified a correct (consistent with labels) match of 13 residues (*L1p* is absent in the canonical kink-turn) at *RMSD* of *1.837 Å* between the two instances. Accordingly, we set the search for the matches of at least 12 residues (*sizemin = 12*) at RMSD under 2.0 Å (*rmsdmax = 2.0*). Additionally, we imposed a restraint requiring the matches to include the counterparts of all of the following residues: *L1, -1b, -1n, 1b, 1n, 2b, 2n*, as they are deemed most essential for the kink-turn motif.

We then executed ARTEM for each of the two references against all RNA-containing structures in the Protein Data Bank [42] (7,256 PDB entries as of December 7th, 2023). The search concluded within 24 hours on an AMD Ryzen 9 5950X machine equipped with 32 CPU cores and 128 GB RAM. ARTEM identified 11,025 matches for the kink-turn and 18,727 matches for the k-junction. Combining the matches identified against both references resulted in 26,564 unique matches (Supplementary Table S1). Finally, to facilitate subsequent analysis, we used the *urslib2* Python library [10] to annotate RNA secondary structure elements and additional characteristics (e.g., local/long-range, kink/no-kink) of the hits' residues based on the base pair annotations obtained with DSSR [15] (version 2.0). The kink/no-kink distinction was approximated by the distance in RNA sequence between the *2b* and *L1* residues: the kink label was assigned to the matches with a distance under ten residues.

Analysis of the kink-turn motif search results

We manually inspected the motif matches identified with ARTEM, targeting novel backbone topologies, as annotated with *urslib2*. The analysis was performed using ChimeraX [54]. The boundaries of the identified representative modules were defined manually, and the modules were then saved as separate coordinate files (https://github.com/febos/ARTEM-KT).

The naming of the ribosomal k-junctions *J4/5* and *J94/99* followed the original k-junction paper [28]. The name for the *J99/101* k-junction was derived analogously using the RibosomeGallery [45] (http://apollo.chemistry.gatech.edu/RibosomeGallery/). The mapping of the identified motifs to the named ribosomal regions was done based on the original structure papers [43, 44, 55]. We should note that modules #9 (PDB entry 4V8O, *J99/101*) and #10 (PDB entry 3CC2, *J94/99*) were missing from the original list of ARTEM matches (Supplementary Table S1). They could be identified with ARTEM only with relaxed thresholds: 11-residue hit at *RMSD = 1.975 Å* against module #2 and

12-residue hit at *RMSD = 1.958 Å* missing *-1n* counterpart against module #1, respectively.

For the conservation analysis, we mapped the three k-junctions to the Rfam seed alignment of the 23S rRNA family RF02541 [56] (https://github.com/febos/ARTEM-KT/tree/main/sto_logo). The relevant sequence regions were then excised from the alignment and used as input for the WebLogo application [57] (https://weblogo.berkeley.edu/) to construct the motif logos.

Group II intron survey for kink-turns

For the benchmarking of the motif search tools, we selected all representative RNA structures assigned with the "group II" keyword in the representative set [47] (version 3.312 with the resolution cutoff of 4.0 Å, http://rna.bgsu.edu/rna3dhub/nrlist/release/3.312). We excluded the 70-nt structure (PDB entry 1KXK) as it is a short intron fragment. The remaining set included seven group II intron structures (Table 2). We then manually identified seven kink-turn 3D modules in the seven structures: five coordination loops and two DIII kink-turns, all of them with the internal loop architecture.

Overall, we utilized seven different RNA tertiary motif search tools (Supplementary Table S2). Although direct comparisons between the tools are often unfeasible, we approached the analysis from the final user perspective, where we have a reference motif instance and a query RNA structure, aiming to identify motif instances within the structure. The 13 residues (*L1, -1b, -1n, -2b, -2n, 1b, 1n, 2b, 2n, 3b, 3n, 4b, 4n*) of the Kt7 kink-turn module (PDB entry *1FFK*, chain *0*, *L1* residue *G94*) were used as the reference motif instance. For all the tools, we considered the identified match to be a true positive (*TP*) if it overlapped with the correct kink-turn by at least 50% of its residues; otherwise, it was classified as a false positive (*FP*). A correct kink-turn instance not identified by the tool was labeled as a false negative (*FN*). Different matches of the same module were merged together before tallying the *TP, FP*, and *FN* values. The quality metrics were subsequently calculated as follows: precision = *TP / (TP + FP)*, recall = *TP / (TP + FN)*, and *F-score = 2*TP / (2*TP + FP + FN)*.

We ran ARTEM with no RMSD threshold but with two different minimum match-size thresholds: 12 residues or 11 residues. For DSSR [15] (version 2.0, http://forum.x3dna.org/rna-structures/) we derived the kink-turn annotations with the "*Normal kink-turn*" label. For RNAMotifScanX [18] (http://genome.ucf.edu/RNAMotifScanX/RNAMotifScanX-release_v0.0.5_x86-64_rhel.tar.gz) we utilized the built-in kink-turn consensus structure (*models/k-turn_consensus.struct*). WebFR3D [13] (http://rna.bgsu.edu/webfr3d/fr3d.php) was initially executed as a purely geometric search for the reference module, and then

with *cWW* symbolic restraints added. JAR3D [14] (http://rna.bgsu.edu/jar3d/) was executed with the sequence input matching the reference kink-turn (*GGGAG*CGAAGAAC*), as it does not accept 3D modules. LocalSTAR3D [16] (http://genome.ucf.edu/LocalSTAR3D/) and CircularSTAR3D [17] (https://github.com/ucfcbb/CircularSTAR3D) were run with the compatible DSSR version 1.5.3. LocalSTAR3D reported multiple 8-residue single-stem matches not involved in the correct kink-turns, which we chose to ignore and not to assign as false positives. Consequently, four tools reported zero matches overall: WebFR3D, JAR3D, LocalSTAR3D, and CircularSTAR3D.

DATA AVAILABILITY

ARTEM 2.0 is available at https://github.com/david-bogdan-r/ARTEM (DOI: 10.5281/zenodo.11243805). The data on kink-turn modules are available at https://github.com/febos/ARTEM-KT (DOI: 10.5281/zenodo.11406085).

ACKNOWLEDGMENTS

The authors acknowledge the GIFTED Programme of the Polish Children's Fund (https://fundusz.org/en/gifted-programme/) for facilitating the visit of talented students to IIMCB in Warsaw, which contributed to this work.

FUNDING

E.F.B. was supported by the European Molecular Biology Organization [EMBO fellowship ALTF 525-2022 to E.F.B]. J.M.B. was supported by the Polish National Science Centre [NCN grant 2017/25/B/NZ2/01294 to J.M.B.]. Funding for the open access charges: IIMCB funds.

REFERENCES

1. Ganser, L. R., Kelly, M. L., Herschlag, D., & Al-Hashimi, H. M. (2019). The roles of structural dynamics in the cellular functions of RNAs. *Nature reviews Molecular cell biology*, 20(8), 474-489. DOI: 10.1038/s41580-019-0136-0

2. Hendrix, D. K., Brenner, S. E., & Holbrook, S. R. (2005). RNA structural motifs: building blocks of a modular biomolecule. *Quarterly reviews of biophysics*, 38(3), 221-243. DOI: 10.1017/S0033583506004215

3. Butcher, S. E., & Pyle, A. M. (2011). The molecular interactions that stabilize RNA tertiary structure: RNA motifs, patterns, and networks. *Accounts of chemical research*, 44(12), 1302-1311. DOI: 10.1021/ar200098t

4. Leontis, N. B., & Westhof, E. (2003). Analysis of RNA motifs. *Current opinion in structural biology*, 13(3), 300-308. DOI: 10.1016/S0959-440X(03)00076-9

5. Xin, Y., Laing, C., Leontis, N. B., & Schlick, T. (2008). Annotation of tertiary interactions in RNA structures reveals variations and correlations. *Rna*, 14(12), 2465-2477. DOI: 10.1261/rna.1249208

6. Parlea, L. G., Sweeney, B. A., Hosseini-Asanjan, M., Zirbel, C. L., & Leontis, N. B. (2016). The RNA 3D Motif Atlas: Computational methods for extraction, organization and evaluation of RNA motifs. *Methods*, 103, 99-119. DOI: 10.1016/j.ymeth.2016.04.025

7. Huang, L., & Lilley, D. M. (2016). The kink turn, a key architectural element in RNA structure. *Journal of molecular biology*, 428(5), 790-801. DOI: 10.1016/j.jmb.2015.09.026

8. de Smit, M. H., Gultyaev, A. P., Hilge, M., Bink, H. H., Barends, S., Kraal, B., & Pleij, C. W. (2002). Structural variation and functional importance of a D-loop–T-loop interaction in valine-accepting tRNA-like structures of plant viral RNAs. *Nucleic acids research*, 30(19), 4232-4240. DOI: 10.1093/nar/gkf539

9. Reinharz, V., Soulé, A., Westhof, E., Waldispühl, J., & Denise, A. (2018). Mining for recurrent long-range interactions in RNA structures reveals embedded hierarchies in network families. *Nucleic acids research*, 46(8), 3841-3851. DOI: 10.1093/nar/gky197

10. Bohdan, D. R., Voronina, V. V., Bujnicki, J. M., & Baulin, E. F. (2023). A comprehensive survey of long-range tertiary interactions and motifs in non-coding RNA structures. *Nucleic Acids Research*, 51(16), 8367-8382. DOI: 10.1093/nar/gkad605

11. Richardson, K. E., Kirkpatrick, C. C., & Znosko, B. M. (2020). RNA CoSSMos 2.0: an improved searchable database of secondary structure motifs in RNA three-dimensional structures. *Database*, 2020, baz153. DOI: 10.1093/database/baz153

12. Chojnowski, G., Waleń, T., & Bujnicki, J. M. (2014). RNA Bricks—a database of RNA 3D motifs and their interactions. *Nucleic acids research*, 42(D1), D123-D131. DOI: 10.1093/nar/gkt1084

13. Petrov, A. I., Zirbel, C. L., & Leontis, N. B. (2011). WebFR3D—a server for finding, aligning and analyzing recurrent RNA 3D motifs. *Nucleic acids research*, 39(suppl_2), W50-W55. DOI: 10.1093/nar/gkr249

14. Roll, J., Zirbel, C. L., Sweeney, B., Petrov, A. I., & Leontis, N. (2016). JAR3D Webserver: Scoring and aligning RNA loop sequences to known 3D motifs. *Nucleic acids research*, 44(W1), W320-W327. DOI: 10.1093/nar/gkw453

15. Lu, X. J., Bussemaker, H. J., & Olson, W. K. (2015). DSSR: an integrated software tool for dissecting the spatial structure of RNA. *Nucleic acids research*, 43(21), e142-e142. DOI: 10.1093/nar/gkv716

16. Chen, X., Khan, N. S., & Zhang, S. (2020). LocalSTAR3D: a local stack-based RNA 3D structural alignment tool. *Nucleic acids research*, 48(13), e77-e77. DOI: 10.1093/nar/gkaa453

17. Chen, X., & Zhang, S. (2023). CircularSTAR3D: a stack-based RNA 3D structural alignment tool for circular matching. *Nucleic Acids Research*, 51(9), e53-e53. DOI: 10.1093/nar/gkad222

18. Zhong, C., & Zhang, S. (2015). RNAMotifScanX: a graph alignment approach for RNA structural motif identification. *RNA*, 21(3), 333-346. DOI: 10.1261/rna.044891.114

19. Leontis, N. B., & Westhof, E. (2001). Geometric nomenclature and classification of RNA base pairs. *Rna*, 7(4), 499-512. DOI: 10.1017/S1355838201002515

20. Islam, S., Ge, P., & Zhang, S. (2017). CompAnnotate: a comparative approach to annotate base-pairing interactions in RNA 3D structures. *Nucleic acids research*, 45(14), e136-e136. DOI: 10.1093/nar/gkx538

21. Tinoco, I., Uhlenbeck, O. C., & Levine, M. D. (1971). Estimation of secondary structure in ribonucleic acids. *Nature*, 230(5293), 362-367. DOI: 10.1038/230362a0

22. Baulin, E., Yacovlev, V., Khachko, D., Spirin, S., & Roytberg, M. (2016). URS DataBase: universe of RNA structures and their motifs. *Database*, 2016, baw085. DOI: 10.1093/database/baw085

23. Zahran, M., Sevim Bayrak, C., Elmetwaly, S., & Schlick, T. (2015). RAG-3D: a search tool for RNA 3D substructures. *Nucleic acids research*, 43(19), 9474-9488. DOI: 10.1093/nar/gkv823

24. Duarte, C. M., Wadley, L. M., & Pyle, A. M. (2003). RNA structure comparison, motif search and discovery using a reduced representation of RNA conformational space. *Nucleic Acids Research*, 31(16), 4755-4761. DOI: 10.1093/nar/gkg682

25. Popenda, M., Szachniuk, M., Blazewicz, M., Wasik, S., Burke, E. K., Blazewicz, J., & Adamiak, R. W. (2010). RNA FRABASE 2.0: an advanced web-accessible database with the capacity to search the three-dimensional fragments within RNA structures. *BMC bioinformatics*, 11, 1-12. DOI: 10.1186/1471-2105-11-231

26. Shalybkova, A. A., Mikhailova, D. S., Kulakovskiy, I. V., Fakhranurova, L. I., & Baulin, E. F. (2021). Annotation of the local context of RNA secondary structure improves the

classification and prediction of A-minors. *RNA*, 27(8), 907-919. DOI: 10.1261/rna.078535.120

27. Geary, C., Chworos, A., & Jaeger, L. (2011). Promoting RNA helical stacking via A-minor junctions. *Nucleic acids research*, 39(3), 1066-1080.DOI: 10.1093/nar/gkq748

28. Wang, J., Daldrop, P., Huang, L., & Lilley, D. M. (2014). The k-junction motif in RNA structure. *Nucleic acids research*, 42(8), 5322-5331. DOI: 10.1093/nar/gku144

29. Bohdan, D. R., Bujnicki, J. M., & Baulin, E. F. (2024). ARTEMIS-a method for topology-independent superposition of RNA 3D structures and structure-based sequence alignment. *bioRxiv*, 2024-04. DOI: 10.1101/2024.04.06.588371

30. Nguyen, M. N., Sim, A. Y., Wan, Y., Madhusudhan, M. S., & Verma, C. (2017). Topology independent comparison of RNA 3D structures using the CLICK algorithm. *Nucleic acids research*, 45(1), e5-e5. DOI: 10.1093/nar/gkw819

31. Zhang, C., Shine, M., Pyle, A. M., & Zhang, Y. (2022). US-align: universal structure alignments of proteins, nucleic acids, and macromolecular complexes. *Nature methods*, 19(9), 1109-1115. DOI: 10.1038/s41592-022-01585-1

32. Klein, D. J., Schmeing, T. M., Moore, P. B., & Steitz, T. A. (2001). The kink-turn: a new RNA secondary structure motif. *The EMBO journal*. DOI: 10.1093/emboj/20.15.4214

33. Matoušková, E., Dršata, T., Pfeifferová, L., Šponer, J., Réblová, K., & Lankaš, F. (2022). RNA kink-turns are highly anisotropic with respect to lateral displacement of the flanking stems. *Biophysical Journal*, 121(5), 705-714. DOI: 10.1016/j.bpj.2022.01.025

34. Li, M., Deng, J., Peng, X., Wang, J., Wilson, T. J., Huang, L., & Lilley, D. M. (2023). Structure and ion-dependent folding of k-junctions. *RNA*, 29(9), 1411-1422. DOI: 10.1261/rna.079678.123

35. Antonioli, A. H., Cochrane, J. C., Lipchock, S. V., & Strobel, S. A. (2010). Plasticity of the RNA kink turn structural motif. *Rna*, 16(4), 762-768. DOI: 10.1261/rna.1883810

36. Edery, P., Marcaillou, C., Sahbatou, M., Labalme, A., Chastang, J., Touraine, R., ... & Leutenegger, A. L. (2011). Association of TALS developmental disorder with defect in minor splicing component U4atac snRNA. *Science*, 332(6026), 240-243. DOI: 10.1126/science.1202205

37. Sklenovsky, P., Florová, P., Banas, P., Reblova, K., Lankas, F., Otyepka, M., & Sponer, J. (2011). Understanding RNA flexibility using explicit solvent simulations: the ribosomal and group I intron reverse kink-turn motifs. *Journal of Chemical Theory and Computation*, 7(9), 2963-2980. DOI: 10.1021/ct200204t

38. Li, B., Liu, S., Zheng, W., Liu, A., Yu, P., Wu, D., ... & Yang, J. (2024). RIP-PEN-seq identifies a class of kink-turn RNAs as splicing regulators. *Nature Biotechnology*, 42(1), 119-131. DOI: 10.1038/s41587-023-01749-0

39. Merico, D., Roifman, M., Braunschweig, U., Yuen, R. K., Alexandrova, R., Bates, A., ... & Scherer, S. W. (2015). Compound heterozygous mutations in the noncoding

RNU4ATAC cause Roifman Syndrome by disrupting minor intron splicing. *Nature communications*, 6(1), 8718. DOI: 10.1038/ncomms9718

40. Ban, N., Nissen, P., Hansen, J., Moore, P. B., & Steitz, T. A. (2000). The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution. *Science*, 289(5481), 905-920. DOI: 10.1126/science.289.5481.905

41. Thore, S., Frick, C., & Ban, N. (2008). Structural basis of thiamine pyrophosphate analogues binding to the eukaryotic riboswitch. *Journal of the American Chemical Society*, 130(26), 8116-8117. DOI: 10.1021/ja801708e

42. Burley, S. K., Bhikadiya, C., Bi, C., Bittrich, S., Chen, L., Crichlow, G. V., ... & Zardecki, C. (2022). RCSB Protein Data Bank: Celebrating 50 years of the PDB with new tools for understanding and visualizing biological macromolecules in 3D. *Protein Science*, 31(1), 187-208. DOI: 10.1002/pro.4213

43. Crowe-McAuliffe, C., Murina, V., Turnbull, K. J., Huch, S., Kasari, M., Takada, H., ... & Hauryliuk, V. (2022). Structural basis for PoxtA-mediated resistance to phenicol and oxazolidinone antibiotics. N*ature Communications*, 13(1), 1860. DOI: 10.1038/s41467-022-29274-9

44. Waltz, F., Salinas-Giegé, T., Englmeier, R., Meichel, H., Soufari, H., Kuhn, L., ... & Hashem, Y. (2021). How to build a ribosome from RNA fragments in Chlamydomonas mitochondria. *Nature Communications*, 12(1), 7176. DOI: 10.1038/s41467-021-27200-z

45. Petrov, A. S., Bernier, C. R., Gulen, B., Waterbury, C. C., Hershkovits, E., Hsiao, C., ... & Williams, L. D. (2014). Secondary structures of rRNAs from all three domains of life. *PLoS One*, 9(2), e88222. DOI: 10.1371/journal.pone.0088222

46. Haack, D. B., Yan, X., Zhang, C., Hingey, J., Lyumkis, D., Baker, T. S., & Toor, N. (2019). Cryo-EM structures of a group II intron reverse splicing into DNA. *Cell*, 178(3), 612-623. DOI: 10.1016/j.cell.2019.06.035

47. Leontis, N. B., & Zirbel, C. L. (2012). Nonredundant 3D structure datasets for RNA knowledge extraction and benchmarking. *RNA 3D structure analysis and prediction*, 281-298. DOI: 10.1007/978-3-642-25740-7_13

48. Toor, N., Keating, K. S., Fedorova, O., Rajashankar, K., Wang, J., & Pyle, A. M. (2010). Tertiary architecture of the Oceanobacillus iheyensis group II intron. *Rna*, 16(1), 57-69. DOI: 10.1261/rna.1844010

49. Toor, N., Keating, K. S., Taylor, S. D., & Pyle, A. M. (2008). Crystal structure of a self-spliced group II intron. *Science*, 320(5872), 77-82. DOI: 10.1126/science.1153803

50. Baulin, E., Metelev, V., & Bogdanov, A. (2020). Base-intercalated and base-wedged stacking elements in 3D-structure of RNA and RNA–protein complexes. *Nucleic acids research*, 48(15), 8675-8685. DOI: 10.1093/nar/gkaa610

51. Nissen, P., Ippolito, J. A., Ban, N., Moore, P. B., & Steitz, T. A. (2001). RNA tertiary interactions in the large ribosomal subunit: the A-minor motif. *Proceedings of the National Academy of Sciences*, 98(9), 4899-4903. DOI: 10.1073/pnas.081082398

52. Westhof, E. (2014). Isostericity and tautomerism of base pairs in nucleic acids. *FEBS letters*, 588(15), 2464-2469. DOI: 10.1016/j.febslet.2014.06.031

53. Cornish-Bowden, A. (1985). Nomenclature for incompletely specified bases in nucleic acid sequences: recommendations 1984. *Nucleic acids research*, 13(9), 3021. DOI: 10.1093/nar/13.9.3021

54. Pettersen, E. F., Goddard, T. D., Huang, C. C., Meng, E. C., Couch, G. S., Croll, T. I., ... & Ferrin, T. E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein science*, 30(1), 70-82. DOI: 10.1002/pro.3943

55. Jin, H., Kelley, A. C., & Ramakrishnan, V. (2011). Crystal structure of the hybrid state of ribosome in complex with the guanosine triphosphatase release factor 3. *Proceedings of the National Academy of Sciences*, 108(38), 15798-15803. DOI: 10.1073/pnas.1112185108

56. Kalvari, I., Nawrocki, E. P., Ontiveros-Palacios, N., Argasinska, J., Lamkiewicz, K., Marz, M., ... & Petrov, A. I. (2021). Rfam 14: expanded coverage of metagenomic, viral and microRNA families. *Nucleic Acids Research*, 49(D1), D192-D200. DOI: 10.1093/nar/gkaa1047

57. Crooks, G. E., Hon, G., Chandonia, J. M., & Brenner, S. E. (2004). WebLogo: a sequence logo generator. *Genome research*, 14(6), 1188-1190. DOI: 10.1101/gr.849004