Dynamic regulation of NAT10-mediated mRNA acetylation at synapses by learning and memory in mouse hippocampus

Hai-Qian Zhou[1,2#], Jia-Wei Zhang[1#], Zhen Zhu[1#], Wei-Peng Lin[1,2#], Hao-JY Jin[1], Yang-Yang Ding[1], Dong-Sheng Chen[3*] and Dong-Min Yin[1,2,4,5*]

1 Key Laboratory of Brain Functional Genomics, Ministry of Education and Shanghai, School of Life Science, East China Normal University, Shanghai, 200062, China

2 Joint Center for Translational Medicine, Shanghai Fifth People's Hospital, Fudan University and School of Life Science, East China Normal University, Shanghai, 200240, China

3 State Key Laboratory of Common Mechanism Research for Major Diseases, Suzhou Institute of Systems Medicine, Chinese Academy of Medical Sciences & Peking Union Medical College, Suzhou, 215123, China

4 NYU-ECNU Institute of Brain and Cognitive Science at NYU Shanghai, Shanghai, 200062, China.

5 Lead contact

[#]H-Q Z, J-W Z, Z Z and W-P L contributed equally to this work.

[*]Correspondence should be addressed to Prof. Dong-Sheng Chen,

Tel: 86-512-62873780

Fax: 86-512-62873779

Email: cds@ism.pumc.edu.cn

or

Prof. Dong-Min Yin,

Tel: 86-21-62232016

Fax: 86-21-62601953

Email: dmyin@brain.ecnu.edu.cn

Abstract

Post-transcriptional modification of RNA plays important roles in regulating brain functions including learning and memory. N4-acetylcytidine (ac4C), the only known form of RNA acetylation in eukaryotes, has recently been shown to regulate mRNA stability and translation efficiency. However, the function and regulation of mRNA acetylation in the brain remain largely unknown. Here we performed ac4C dot-blots, liquid chromatography tandem mass spectrometry (LC-MS/MS), and ac4C-RNA immunoprecipitation followed by next-generation sequencing (acRIP-seq) to investigate the dynamic change of ac4C mRNA in mouse hippocampus during learning and memory. This study focused on Morris water maze (MWM), a behavioral protocol commonly used for studying hippocampus-dependent spatial learning and memory. Interestingly, the levels of ac4C mRNA were upregulated in the synaptosomes (SYN) after memory formation but returned to normal levels after memories were naturally forgotten. The protein levels of N-acetyltransferase 10 (NAT10), the ac4C writer, were also increased in the SYN after memory formation and returned to control levels after memory forgetting. Strikingly, genetic ablation of Nat10 from hippocampal neurons significantly reduced the ac4C modification of Arc, Camk2α and Grin2b mRNA and their protein levels in the SYN of training mice, which was accompanied by deficits in synaptic plasticity and behaviors of learning and memory. These results demonstrate the dynamics and roles of synaptic mRNA acetylation in learning and memory, offering insights into the epitranscriptomic regulation of brain functions. The dataset of ac4C mRNA from mouse hippocampus is accessible via the website (http://ac4catlas.com/).

Introduction

Precise regulation of protein synthesis at the right place and right time is important for various brain functions including learning and memory (Besse and Ephrussi, 2008; Shrestha and Klann, 2022). For instance, local protein synthesis near stimulated synapses plays critical roles in synaptic plasticity and memory consolidation(Martin et al., 2000; Sutton and Schuman, 2006). Mechanistically, epigenetic regulations on DNA and histone proteins have been widely studied in the context of learning and memory (Day and Sweatt, 2011; Levenson and Sweatt, 2005). However, the roles of chemical modifications on mRNA in learning and memory have only become appreciated in recent years (Fischer et al., 2007; Kouzarides, 2007; Levenson et al., 2004). Especially, N6-methyladenosine (m6A), a well-studied form of mRNA methylation has been shown to regulate synaptic plasticity as well as learning and memory (Merkurjev et al., 2018; Shi et al., 2018; Zhuang et al., 2023).

N4-acetylcytidine (ac4C) is the only known form of RNA acetylation in eukaryotes (Boccaletto et al., 2018). Initially, ac4C was identified in the transfer RNA (tRNA) (Kowalski et al., 1971; Kruppa and Zachau, 1972; Oashi et al., 1972; Stern and LH, 1978). Later studies revealed that ac4C also occurred in the 18S ribosomal RNA (rRNA) (Ito et al., 2014a). More recently, ac4C has been found in mRNA from mammalian cells, where it has been shown to promote mRNA stability and translational efficiency (Arango et al., 2018; Arango et al., 2022). N-acetyltransferase 10 (NAT 10) is the only known ac4C writer in mammalian cells (Ito et al., 2014b). NAT10-mediated mRNA acetylation has been reported to regulate tumorigenesis (Liu et al., 2023b; Wang et al., 2022), stem cell differentiation (Hu et al., 2024a), cardiac remodeling (Shi et al., 2023a), germ cell maturation (Jiang et al., 2023) and pain signaling in the spinal cord (Xu et al., 2023; Zhang et al., 2023a). However, the function and regulatory mechanisms of mRNA acetylation in the brain remain poorly understood.

The hippocampus plays an important role in learning and memory (Jarrard, 1993). Pyramidal neurons in the CA3 region of hippocampus establish direct projection to CA1 pyramidal neurons through the Schaffer collateral (SC) pathway, known as the SC-CA1

synapse. The glutamatergic transmission at SC-CA1 synapses was found to be enhanced following training in the MWM (Moser et al., 1994), a behavioral paradigm for learning and memory (Morris, 1984). Long-term potentiation (LTP), a widely-studied form of synaptic plasticity in hippocampus, is most closely linked to learning and memory (Bliss and Collingridge, 1993). Several studies showed that mRNAs and ribosomes were present in the synaptosomes, contributing to protein synthesis at synapses (Biever et al., 2020). In principle, neural activities can regulate local protein synthesis through post-transcriptional modification of synaptic mRNAs (Goldie and Cairns, 2012). However, whether synaptic mRNAs were subjected to NAT10-mediated acetylation and whether this process contributed to synaptic plasticity and memory remain elusive.

Given that eukaryotic mRNAs are polyadenylated, we first conducted ac4C dot-blots and liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis to examine the acetylation of polyadenylated (polyA) RNA from mouse hippocampus during and after training in the MWM. Our findings revealed that the acetylation of polyA RNA was increased in the synaptosomes (SYN), rather than in the homogenates (HOM), following memory formation, and subsequently returned to baseline after memory forgetting. The dynamic changes of mRNA acetylation at synapses throughout the processes of learning and memory was further demonstrated by ac4C-RNA immunoprecipitation followed by next-generation sequencing (acRIP-seq). Moreover, the increased levels of ac4C mRNA in the SYN after learning and memory were accompanied with the increased protein levels of NAT10 at synapses. Genetic ablation of Nat10 from hippocampal neurons in adult mice using the Cre/Loxp strategy significantly reduced the acetylation of Arc, Camk2α and Grin2b mRNA and their protein levels at synapses, and thus impaired learning and memory in the MWM. Collectively these results demonstrate the dynamic regulation of NAT10-mediated acetylation of synaptic mRNAs during learning and memory. The dataset of ac4C epitranscriptome from mouse hippocampus is freely accessible at the website (http://ac4catlas.com/).

Results

Presence of ac4C modification of polyA RNA in mouse hippocampus

Recent studies raised the debate about the presence of ac4C in mRNA from mammalian cells (Beiki et al., 2024; Georgeson and Schwartz, 2024). We first investigate whether ac4C is a detectable modification of mRNA from mouse hippocampus. To this end, we first purified polyA RNA from the HOM of mouse hippocampus and determine their ac4C levels using LC-MS/MS. The purity of polyA RNA was verified by the bioanalyzer analysis (Figure 1A and 1B) and dramatic reduction of 18S rRNA relative to total RNA in reverse transcription followed by quantitative PCR (RT-qPCR) (Figure 1C). An ac4C standard curve was generated to calculate the stoichiometry levels of ac4C in polyA RNA samples (Figure 1D). Both the ac4C and m6A peaks can be detected by the chromatograms in the standard samples but not in the blank solution (Figure 1E and 1F). The polyA RNA purified from the HOM of mouse hippocampus showed ac4C peaks with a stoichiometry level of $0.426 \pm 0.049\%$ (n = 3 biological replicates) (Figure 1G and 1H). Likewise, the polyA RNA purified from the HOM of mouse hippocampus showed m6A peaks with a stoichiometry level of $0.173 \pm 0.018\%$ (n = 3 biological replicates) (Figure 1G and 1I), which is consistent with the previous study form mouse brain (Li et al., 2023). When the polyA RNA samples were incubated with 50 mM NH_2OH for 1 h before LC-MS/MS, a protocol of chemical deacetylation of ac4C (Sinclair et al., 2017), the stoichiometry levels of ac4C were significantly reduced (Figure 1H), indicating the validity of the LC-MS/MS analysis. By contrast, the stoichiometry levels of m6A in polyA RNA samples remained unchanged after NH_2OH treatment (Figure 1I), supporting the specificity of the deacetylation method. Altogether these results demonstrate the presence of ac4C modification in polyA RNA from mouse hippocampus. Given that polyA RNA consists of mRNA and long noncoding RNA (lncRNA) (Yu et al., 2020), we further performed acRIP-seq to analyze the ac4C mRNAs in mouse hippocampus (see below).

Characterization of ac4C mRNA in the HOM of mouse hippocampus after learning and memory

We used the behavioral protocol of MWM (Vorhees and Williams, 2006) to study how learning and memory regulate mRNA acetylation in mouse hippocampus. The control mice were placed in the MWM with a visible platform once a day for 2 days, and thus could reach the visible platform in the target region without training (Figure S1A and S1B). However, the control mice could not remember the target region during the probe test at day 6 (Figure S1A and S1C). In contrast, the learning mice were subjected to training in the MWM with a hidden platform (Figure S1D and S1E). The learning mice took 5 days of training to remember the target region during the probe test at day 6 (Figure S1E and S1F), indicative of memory formation. Consistent with the previous study (Hoz et al., 2004), the learning mice could not remember the target region during the probe test at day 20, i.e., after staying in home cage for 2 weeks without training (Figure S1D and S1G), indicating that memories were naturally forgotten (Figure S1H).

We next performed acRIP-seq (Arango *et al.*, 2018) to identify ac4C mRNA in the HOM of mouse hippocampus. Two libraries of RNA (acRIP and input) were subjected to the next-generation sequencing (NGS) for each sample (Table S1). The acRIP-seq reads were mapped to mouse hippocampal transcriptome to identify ac4C enriched regions relative to input, and the fold enrichment (FE) of an ac4C peak was defined as acRIP / input (Figure S2A). An ac4C peak was considered positive when its FE > 2 and P-value < 0.05 (Zhang et al., 2023b). When we analyzed the cumulative distribution curve of the levels of ac4C mRNA, the FE of ac4C mRNA was the sum of the FE of all ac4C peaks in one particular mRNA (see details in the Method).

We performed acRIP-seq for three HOM replicates from control and learning mice one hour after probe tests in the MWM (i.e., at day 6). The ac4C peak and ac4C mRNA identified in three biological replicates of control mice were largely overlapped (Figure S2B and S2C). Moreover, the acRIP-seq revealed the same consensus motif AGCAGCTG in the ac4C peak regions from three biological replicates of control mice (Figure S2D). These results indicate the consistency of the acRIP-seq for mouse hippocampus. The ac4C peak that were identified from at least two of the three biological replicates were considered reliable (Figure S2E and S2F). Only mRNAs

corresponding to these reliable ac4C peaks are defined as reliably ac4C mRNAs. We

identified 14516 ac4C peaks and 8303 ac4C mRNAs from the HOM of control mice

(Figure S2E and Table S2), and 13348 ac4C peaks and 8107 ac4C mRNAs from the

HOM of learning mice (Figure S2F and Table S2). As a control of the acRIP-seq, only

1800 ac4C peaks were identified from the HOM of control mice when we performed

IgG immunoprecipitation of RNA followed by NGS, indicating the specificity of

acRIP-seq.

When we compared the transcripts with ac4C modifications to the entire

transcriptome, we found that $49.22 \pm 2.52\%$ (n = 3 biological replicates) of the mRNAs

in the mouse hippocampus were modified by ac4C. In line with the previous study from

Hela cells (Arango et al., 2018), most ac4C peaks were identified in the coding

sequence (CDS) of mRNA in mouse hippocampus (Figure S2G). On average, every

acetylated transcript had 1.83 ± 0.05 (n = 3 biological replicates) ac4C peaks in mouse

hippocampus. The numbers of ac4C peak per acetylated mRNA were comparable

across three biological replicates and between control and learning mice (Figure S2H).

Likewise, the ac4C dot-blots of polyA RNA, and the cumulative distribution of FE of

ac4C mRNA in the HOM were similar between the control and learning mice (Figure

S2I and S2J). The cumulative distribution of mRNA counts in the HOM that were

revealed by the NGS of input RNA were also similar between control and learning mice

(Figure S2K). Altogether these results suggest that the total levels of ac4C mRNA in

the HOM of mouse hippocampus may not be altered by learning and memory in the

MWM.

Dynamic change of ac4C polyA RNA in the SYN of mouse hippocampus during

learning and memory in the MWM

We next purified the SYN fraction from mouse hippocampus, enriched for synaptic

proteins (PSD95 and synapsin 1) and depleted of cytoplasm (the S2 fraction)-restricted

protein (α-tubulin) (Figure 2A). We then analyzed the levels of ac4C polyA RNA

purified from the SYN with the ac4C dot-blots. The control and learning mice showed

similar levels of ac4C polyA RNA in the SYN at day 1 (Figure 2B). Strikingly, the

levels of ac4C polyA RNA showed a significant increase in the SYN of learning mice at day 6, compared with controls (Figure 2C). However, the levels of ac4C polyA RNA returned to control levels in the SYN of learning mice at day 20 (Figure 2D). These results suggest that acetylation of polyA RNA was upregulated in the SYN of mouse hippocampus during memory formation but returned to control levels after memory forgetting. The dynamic change of ac4C polyA RNA in the SYN of mouse hippocampus during learning and memory was verified by the assay of LC-MS/MS (Figure 2E).

Characterization of ac4C mRNA in the SYN of mouse hippocampus

Given that the amount of RNA from SYN (100 ng per mouse hippocampus) is much less than that from HOM (50 ug per mouse hippocampus), we used the protocol of low-input acRIP-seq to identify ac4C mRNA in the SYN (see details in the Methods). Due to the limited yield of SYN RNA, we purified the SYN RNA from the hippocampus of 10 adult mice and combined them into one sample. Each sample from control and learning mice at days 1, 6, and 20 was subjected to acRIP-seq, and thus 6 total samples which contained SYN RNA purified from 60 adult mice were sequenced to reveal the ac4C mRNA regulated by learning and memory in the SYN. Two libraries (acRIP and input) were sequenced for each biological sample (Table S3).

The transcripts and ac4C mRNAs identified from the SYN of control mice at days 1, 6 and 20 were largely overlapped (Figure S3A and S3B). Moreover, the low-input acRIP-seq revealed the same consensus motif AGCAGCTG in the ac4C peak regions of SYN mRNA from control mice at days 1, 6 and 20 (Figure S3C). In line with the results from HOM, most ac4C peaks were identified in CDS of SYN mRNA from both control and learning mice (Figure S3D and S3E). These results indicate the consistency of the low-input acRIP-seq for SYN mRNA.

Dynamic change of ac4C mRNA in the SYN of mouse hippocampus during learning and memory

The cumulative distribution of FE of ac4C mRNA in the SYN did not differ between control and learning mice at day 1 (Figure 2F). In contrast, the cumulative distribution curve of FE of ac4C mRNA in the SYN of learning mice had a significant rightward

shift at day 6, compared with that of control mice (Figure 2G). Intriguingly, the cumulative distribution of FE of ac4C mRNA in the SYN of learning mice became indistinguishable from that of control mice at day 20 (Figure 2H). The cumulative distribution of mRNA counts in the SYN remained similar between control and learning mice at days 1, 6, and 20 (Figure 2I-K). These results demonstrate that acetylation of mRNA rather than the total levels of mRNA was increased in the SYN after memory formation but returned to control levels after memory forgetting, further supporting the dynamic change of acetylation of polyA RNA in the SYN during learning and memory (Figure 2B-E).

To identify the learning-induced synaptic ac4C (LISA) mRNA, we analyzed the fold change (FC) of ac4C mRNA in the SYN of learning mice versus control mice (L/C) at days 1, 6, 20 (Figure S3F to S3H). Here we analyzed the average FE of all ac4C peaks to reflect the acetylation level of individual LISA mRNA. Ideally, acetylation of LISA mRNA should be upregulated after memory formation and return to control levels after memory forgetting. In other words, the ac4C of LISA mRNA is specifically upregulated at day 6 (the cut off value of upregulation is FC > 2) but not at days 1 or 20. Per these criteria, we identified 1237 LISA mRNAs whose ac4C levels were increased at day 6 rather than at days 1 or 20 (Figure S3I and Table S4). In contrast, the mRNA levels of LISA were similar at days 1, 6, and 20 (Figure S3I). Gene ontology (GO) analysis indicated that LISA mRNA were significantly enriched in the biological processes (BP) related with synaptic plasticity and learning or memory (Figure 3A and Table S4).

A representative LISA mRNA for Arc

Protein-protein interaction (PPI) network analysis of LISA mRNA involved in the BP of learning or memory revealed several core genes such as Arc, presenilin 1 (Psen1), and Bdnf (Figure 3B). Arc mRNA encodes activity-regulated cytoskeleton associated protein (ARC) and is unique among immediately-early genes (IEG). Arc mRNA can be transported into dendrites and translated near activated postsynaptic sites after neural activities (Lyford et al., 1995; Steward et al., 1998). Different from the regular IEG, Arc

underwent repetitive cycles of transcription and local translation, which is critical for memory consolidation (Das et al., 2023). Although Arc is a promising "master regulator" of protein synthesis-dependent forms of learning and memory, the mechanisms underlying the regulation of local translation of Arc in the SYN remained to be elucidated (Bramham et al., 2010). Analysis of the acRIP-seq data indicate that learning mice have elevated ac4C of Arc mRNA in the SYN at day 6 rather than days 1 or 20, and similar levels of ac4C Arc mRNA in the HOM at day 6, compared with control mice (Figure 3C). The genome browser tracks for the ac4C of all LISA mRNA were accessible at the ac4C website (http://ac4catlas.com/) to show altered ac4C in the SYN, and similar lack of change for those same RNAs in the HOM at day 6.

Next, we used an orthogonal method to verify the dynamic change of ac4C Arc mRNA in the SYN during learning and memory. To this end, performed RT-qPCR analysis for Arc mRNA from the input and acRIP samples purified from the SYN. The levels of input and acRIP-Arc mRNA were similar in the SYN of control and learning mice at days 1 and 20 (Figure 3D, 3E, 3G and 3H). Likewise, the protein levels of ARC were also comparable in the SYN of control and learning mice at days 1 and 20 (Figure 3F and 3I). Although the levels of input Arc mRNA were not altered in the SYN of learning mice at day 6 (Figure 3J), the levels of acRIP-Arc mRNA and ARC proteins were significantly increased in the SYN of learning mice at day 6, compared with controls (Figure 3K and 3L). In contrast, the levels of Arc mRNA and ARC proteins were similar in the HOM of control and learning mice at day 6 (Figure 3M and 3N). Note that Arc mRNA levels in the IgG-IP samples were much lower than that in the acRIP samples (Figure 3E, 3H and 3K), indicating the specificity of RT-qPCR for acRIP. Altogether the results from acRIP-seq and acRIP-qPCR demonstrate that acetylation of Arc mRNA is specifically increased in the SYN after memory formation but returned to control levels after memory forgetting.

Expression of Nat10 in the mature neurons of mouse hippocampus

Since the commercially available NAT10 antibodies were not suitable for immunostaining in the hippocampal slices, we took use of the strategy of genetic

labeling (Madisen et al., 2010; Yu et al., 2019) to investigate the expression pattern of Nat10. To this end, a heterozygous knockin mouse line (Nat10$^{Cre/+}$) that expressed the Cre recombinase right after the start codon (ATG) of Nat10 gene was generated using the CRISPR/Cas9 techniques (see details in the Method). Since the Cre recombinase and Nat10 share the same regulatory elements for gene expression in the Nat10$^{Cre/+}$ mice, the Nat10 expression pattern can be revealed through the Cre-reporting adeno-associated virus (AAV) that express EGFP in a Cre-dependent manner. In this way, the Nat10-positive cells would express EGFP in the Nat10 reporter mice (Figure S4A).

The dorsal hippocampus integrates multiple synaptic inputs from other brain regions, and is critical for learning and memory (Biane et al., 2023; Fanselow and Dong, 2010; Moser et al., 1993). To study the expression pattern of Nat10 in the dorsal hippocampus of adult mice, we injected Cre-reporting AAV into the dorsal hippocampus of 7-week-old Nat10$^{Cre/+}$ mice, and the fluorescent protein EGFP was analyzed 3 weeks after AAV injection. Analysis of the EGFP-positive cells revealed that Nat10 was mainly expressed in the pyramidal cell layer (sp) of CA1 region and granule cell layer (gcl) of dentate gyrus (DG) (Figure S4B). Around 80% of Nat10-expressing cells are mature neurons, evidenced by the overlay between EGFP and NeuN, a protein marker for mature neurons (Figure S4C and S4D). Note that about 20% Nat10-positive cells are non-neuronal cells and probably glia cells. Moreover, about 75% of Nat10-positive cells are excitatory pyramidal neurons in the CA1 region of mouse hippocampus, indicated by the co-localization of EGFP with Neurogranin (NG), a protein marker for excitatory pyramidal neurons (Figure S4C and S4E). These results indicate that Nat10 is mainly expressed by mature neurons in adult mouse hippocampus.

To verify that the expression of Cre-reporting AAV has no infection preference toward neurons, we injected the AAV into the dorsal hippocampus of 7-week-old ALDH1L1$^{Cre/ERT2}$ mice, a tamoxifen-induced transgenic Cre line for selectively targeting astrocytes (Srinivasan et al., 2016). After AAV injection for 3 week and tamoxifen gavage for 1 week, we analyzed the expression of the fluorescent protein EGFP in the astrocyte reporter mice (Figure S4F). In the CA1 region of astrocyte

reporter mice, EGFP was not expressed in neurons and all EGFP-positive cells show astrocyte-like morphology (Figure S4G). Collectively these results verify the notion that the expression of Cre-reporting AAV has no neuronal preference and strictly depends on the expression pattern of Cre.

Increase of NAT10 proteins in the SYN after learning and memory

The above-mentioned data showed that the levels of ac4C mRNA were increased in the SYN of mouse hippocampus after learning and memory. Next, we aim to investigate whether the protein levels of NAT10, the only known ac4C writer (Liu et al., 2023a), are also regulated by learning and memory. Toward this aim, we purified the SYN fraction from the hippocampus of control and learning mice at days 1, 6 and 20, and analyzed the protein levels of NAT10 by western blots (Figure 4A). The protein levels of NAT10 in the SYN were similar between control and learning mice at day 1 (Figure 4B). Interestingly, the protein levels of NAT10 were significantly increased in the SYN of learning mice at day 6, compared with controls (Figure 4C), but returned to control levels at day 20 (Figure 4D). These results indicate that protein levels of NAT10 are increased in the SYN after memory formation but return to normal levels after memory forgetting (Figure 4E).

The increase of NAT10 proteins in the SYN at day 6 could be due to the upregulation of NAT10 in the HOM (i.e., the increase of de novo protein synthesis). However, the protein levels of NAT10 were not altered in HOM of learning mice at day 6, compared to controls (Figure 4F), suggesting that learning and memory did not upregulate the de novo protein synthesis of NAT10 in mouse hippocampus. To further explore the distribution of NAT10 proteins in other subcellular fractions, we purified the nucleus (N), cytoplasm (S2) and post-synaptic density (PSD) fractions from the hippocampus of control and learning mice at day 6, and analyzed the protein levels of NAT10 in different subcellular fractions (Figure 4G). The nuclear fraction was enriched for histone but was depleted of cytoplasmic protein α-tubulin or synaptic protein SYN1 and PSD95, and vice versa for the S2 and PSD fraction (Figure 4H). The protein levels of NAT10 in the nucleus did not change at day 6 after learning and memory (Figure 4I).

Strikingly, the NAT10 proteins were significantly reduced in the cytoplasm (S2 fraction)

but increased in the PSD fraction at day 6 after learning and memory (Figure 4J and

4K). These results suggest that NAT10 proteins might be relocated from the cytoplasm

to synapses in mouse hippocampus after memory formation.

NAT10 belongs to the p300/CBP-associated factor (PCAF) subfamily of protein

acetyltransferase (Lau et al., 2000). Next, we aim to investigate whether learning and

memory also changed protein levels of PCAF in the subcellular fractions of mouse

hippocampus. The protein levels of PCAF did not change in the nucleus, cytoplasm or

PSD fractions of learning mice at day 6, compared with controls (Figure 4L-N),

indicating that learning and memory may not alter the subcellular distribution of PCAF.

Next, we performed immunofluorescent staining to study the synaptic distribution

of NAT10 proteins before and after chemical LTP stimulation in cultured hippocampal

neurons (Figure S5A) – a cellular model of learning and memory (Lu et al., 2001; Shahi

and Baudry, 1993). In control neurons treated with vehicle, NAT10 was mainly

expressed in the nucleus and cytoplasm (Figure S5B), which is consistent with the

results of western blots (Figure 4I and 4J). However, 60 minutes after chemical LTP

stimulation, the colocalization of NAT10 with PSD95, a protein marker for PSD

fraction, was significantly increased, compared with control neurons (Figure S5C and

S5D). These results indicated that distribution of NAT10 in the PSD were increased in

cultured hippocampal neurons after chemical LTP stimulation, which support the

increased protein levels of NAT10 in the PSD fraction of mouse hippocampus after

learning and memory (Figure 4K).

NAT10-dependent ac4C modification of polyA RNA in mouse hippocampus

To demonstrate the NAT10-dependent ac4C modification of poly A RNA, we used the

Cre/Loxp strategy (Tsien et al., 1996) to downregulate Nat10 expression in adult mouse

hippocampus through injection of AAV-EGFP (as a control) or AAV-EGFP-P2A-Cre

into the hippocampus of Nat10$^{flox/flox}$ mice. Given that Nat10 is mainly expressed in the

mature neurons (Figure S4), the expression of EGFP or EGFP-2A-Cre in the AAV was

driven by the human synapsin 1 (Syn1) promoter, a neuron-specific promoter

(Hedegaard et al., 2013). The Nat10$^{flox/flox}$ mice injected with AAV-EGFP and AAV-

EGFP-P2A-Cre were named control and Nat10 conditional knockout (cKO) mice

hereafter. Both the control and Nat10 cKO mice received stereotaxic AAV injection into

the dorsal hippocampus, and the HOM of dorsal hippocampus were collected for

biochemical assay 3 weeks after AAV injection (Figure S6A).

The fluorescent protein EGFP was expressed mainly in the CA1-3 regions and DG

of mouse hippocampus 3 weeks after AAV injection (Figure S6B), indicating the

successful infection of AAV. The protein levels of NAT10 were significantly reduced in

the hippocampus of Nat10 cKO mice, compared with controls (Figure S6C and S6D),

thereby verifying the effective knockdown of NAT10 by the Cre/Loxp strategy. NAT10

was previously described as a protein acetyltransferase against α-tubulin and histones

(Lv et al., 2003; Shen et al., 2009). However, immunoblotting with acetylation-specific

antibodies showed no significant change of protein acetylation in Nat10 cKO

hippocampus, compared to controls (Figure S6C, S6E and S6F). These results suggest

the redundancy of acetyltransferases for α-tubulin and histones in mouse hippocampus.

In contrast, the ac4C modification of polyA RNA was significantly decreased in the

Nat10 cKO mice, compared with controls (Figure S6G and S6H). As a control

experiment, downregulation of NAT10 did not affect the m6A modification of polyA

RNA (Figure S6I and S6J). The LC-MS/MS analysis of polyA RNA indicated that the

stoichiometry levels of ac4C rather than m6A were significantly reduced in the

hippocampus of Nat10 cKO mice, compared with controls (Figure S6K and S6L).

Altogether these results demonstrate that NAT10 specifically regulates ac4C rather than

m6A of polyA RNA in mouse hippocampus.

NAT10-dependent ac4C mapping of SYN mRNAs in learning mice

Given that learning and memory increased ac4C mRNA specifically in the SYN, we

next aim to identify NAT10-dependent ac4C modification of synaptic mRNA. To this

end, we performed low-input acRIP-seq for SYN RNA purified from the hippocampus

of control and Nat10 cKO mice at day 6 after learning and memory (Figure 5A). Due

to the limited yield of SYN RNA, we purified SYN RNA from the hippocampus of 10

mice and combined them into one sample. Two biological samples from the respective

control and Nat10 cKO mice at day 6 were subjected to low-input acRIP-seq to

demonstrate the NAT10-dependent ac4C modification of SYN mRNA. Two libraries

(acRIP and input) were sequenced for each biological sample (Table S3).

382   We identified 7097 and 1933 ac4C peaks from the SYN of control and Nat10 cKO

mice, respectively (Figure 5B). These results suggest a reduction of the a4C peak

number by conditional knockout of Nat10 in mature neurons of adult mouse

hippocampus. Out of 7097 ac4C peaks identified in control mice, 6331 peaks were

abolished and 408 peaks were downregulated in Nat10 cKO mice, and these ac4C peaks

(total 6739) were named NAT10-dependent ac4C peaks (Figure 5C) which were mainly

localized in the CDS region of mRNA (Figure 5D).

389   The Nat10 cKO mice also showed a reduction of ac4C mRNA number identified

in the SYN, compared with controls (Figure 5E). In contrast, the number of mRNA

identified in the SYN was similar between control and Nat10 cKO mice (Figure 5F).

Likewise, the total levels of ac4C mRNA were significantly reduced in the SYN of

Nat10 cKO mice, compared with controls, as evidenced by the leftward-shifted

cumulative distribution curve of FE of ac4C mRNA in the Nat10 cKO mice (Figure

5G). In contrast, the cumulative distribution curve of mRNA counts were similar in the

SYN of control and Nat10 cKO mice (Figure 5H). Out of 3778 ac4C mRNAs identified

in the SYN of control mice, 2507 ac4C mRNAs were diminished and 839 ac4C mRNAs

were downregulated in the Nat10 cKO mice, and these ac4C mRNAs (total 3346) were

named <u>NA</u>T10-dependent <u>s</u>ynaptic <u>ac</u>4C (NASA) mRNA (Figure 5I and Table S5).

400   We next analyzed the cellular expression pattern of NASA mRNA through the

single-cell RNA-seq atlas of mouse hippocampus (dropviz.org) (Saunders et al., 2018).

The results indicated that the expression of NASA mRNA was mostly enriched in

excitatory neurons (Figure 5J), which is in line with the cellular expression pattern of

NAT10 in mouse hippocampus (Figure S4). In addition, about 13% of NASA mRNAs

were relatively enriched in glia cells (Figure 5J), which is consistent with the notion

that processes of glia cells contribute to formation of the tripartite synapses (Eroglu and

Barres, 2010; Perea et al., 2009). Gene ontology (GO) analysis showed that NASA

mRNAs were significantly enriched in the biological processes (BP) related with

synaptic plasticity and learning or memory (Figure 5K and Table S5).

Regulation of ac4C and protein levels of Arc, Camk2α and Grin2b in the SYN by NAT10

Twenty-seven NASA mRNAs implicated in the BP of synaptic plasticity and learning

or memory also belong to LISA mRNA (Figure 6A and Table S4), which we named

class 1 mRNA. In contrast, 212 NASA mRNAs related with synaptic plasticity and

learning or memory did not belong to LISA mRNA (Figure 6A and Table S5), which

we named class 2 mRNA. Analysis of the protein-protein interaction (PPI) network

revealed significant interaction between the mRNAs in both class 1 and class 2.

Remarkably, 14 out of the 27 mRNAs in class 1 were found to be interconnected (Figure

6B), and 199 out of the 212 mRNAs in class 2 formed a PPI network (Figure 6C).

Arc, a representative LISA mRNA (Figure 3), was found in the PPI network of

class 1 mRNA. The mRNA of Ca^{2+}/calmodulin-dependent protein kinase 2α (Camk2α)

and Grin 2b encode the protein kinase CaMK2α and the GluN2B subunit of NMDA

receptor, respectively. CaMK2α and GluN2B are synapse-enriched proteins and play

important roles in LTP as well as learning and memory (Lisman et al., 2012; Tang et al.,

1999). Recent studies demonstrate that CaMK2α interaction with GluN2B is critical

for the induction and maintenance of LTP (Chen et al., 2024; Tullis et al., 2023). Both

Camk2α and Grin 2b were identified in the PPI network of class 2 mRNA. Analysis of

the low-input acRIP-seq data indicated the reduction of ac4C peaks of Arc,

Camk2α and Grin 2b mRNA in the SYN of Nat10 cKO mice, compared with controls

(Figure 6D-F). The genome browser tracks for the ac4C of all NASA mRNAs were

accessible at the ac4C website (http://ac4catlas.com/) to show reduced or diminished

ac4C peaks in the SYN of Nat10 cKO mice, compared with controls.

Next, we performed RT-qPCR analysis for input and acRIP RNAs purified from

the SYN of control and Nat10 cKO mice at day 6 after learning and memory to verify

the NAT10-dependent ac4C modification of Arc, Camk2α and Grin2b mRNA (Figure

6G). As shown in Figure 6H, the input mRNA levels of Arc, Camk2α and Grin2b in the SYN were comparable between control and Nat10 cKO mice. In contrast, the acRIP mRNA levels of Arc, Camk2α and Grin2b in the SYN were significantly reduced in Nat10 cKO mice, compared with controls (Figure 6I). Given that ac4C modification of mRNA has been shown to enhance translation efficiency (Arango et al., 2018; Arango et al., 2022), we sought to determine whether Nat10 influences the protein levels of ARC, CaMK2α and GluN2B in the SYN by western blot analysis (Figure 6G). The protein levels of ARC, CaMK2α and GluN2B were significantly decreased in the SYN at day 6 from Nat10 cKO mice, compared with controls (Figure 6J). Altogether, these results demonstrate that Nat10 regulates ac4C and protein levels, rather than mRNA levels of Arc, Camk2α and Grin2b in the SYN of mouse hippocampus at day 6 in the MWM.

Regulation of LTP as well as learning and memory by NAT10 in mouse hippocampus

In the following study, we investigate whether Nat10 in mouse hippocampus is important for LTP as well as learning and memory. The body and brain weight were similar between control and Nat10 cKO mice (Figure S7A and S7B). The number and intrinsic excitability of CA1 pyramidal neurons were also comparable between control and Nat10 cKO mice (Figure S7C and S7D). To address whether Nat10 regulates LTP, we performed electrophysiology recording of fEPSP (field excitatory postsynaptic potential) at the SC-CA1 synapses before and after high frequency stimulation (HFS) (Figure 7A). The induction of LTP was comparable between control and Nat10 cKO mice (Figure 7B-D). However, the maintenance of LTP especially the late phase of LTP was severely impaired in Nat10 cKO mice, compared with controls (Figure 7B, 7C, 7E and 7F).

To further study whether Nat10 is important for synaptic plasticity after learning and memory, we performed fEPSP recording to investigate glutamatergic transmission at the SC-CA1 synapses in the control and Nat10 cKO mice at day 6 after learning and memory (Figure 7G). Consistent with the previous report (Su et al., 2023), the

glutamatergic transmission at the SC-CA1 synapses was significantly enhanced at day

6 in learning mice, compared with mice without learning (Figure 7H and 7I). The

glutamatergic transmission at the SC-CA1 synapses was similar between control and

Nat10 cKO mice without learning, which indicate that Nat10 may be dispensable for

the glutamatergic transmission at the SC-CA1 synapses at basal conditions. However,

the learning and memory-induced enhancement of glutamatergic transmission at the

SC-CA1 synapses was significantly impaired in Nat10 cKO mice, compared with

controls (Figure 7H and 7I). Altogether these results indicate that NAT10 is important

for synaptic plasticity related with learning and memory in mouse hippocampus.

To investigate whether NAT10 is critical for learning and memory, both control and

Nat10 cKO mice were subjected to training in the MWM for 5 days, and the probe test

was performed at day 6 (Figure 7G). Both groups exhibited an overall decline in the

latency to reach the platform at the early stage (day 1-2) of training (Figure 7J).

However, the Nat10 cKO mice failed to improve performance during the late training

stage (day 3-5), compared with controls (Figure 7J). During the probe tests, the control

and Nat10 cKO mice showed similar distance travelled (Figure 7K and 7L). However,

the time spent in the target area and the number of platform crossing were significantly

reduced during the probe test in Nat10 cKO mice, compared with controls (Figure 7L-

N). These results demonstrate that NAT10 in mouse hippocampus is important for

spatial learning and memory.

Discussion

In this study, we demonstrate the dynamic regulation of synaptic mRNA acetylation by

learning and memory in mouse hippocampus. We identified ~1200 ac4C-modified

mRNAs (i.e., LISA mRNA) that were increased in the SYN of mouse hippocampus

after memory formation but returned to normal levels after memories were naturally

forgotten. In contrast, LISA mRNAs were not increased in the HOM of mouse

hippocampus after learning and memory. Interestingly, the protein levels of NAT10

were also increased in the SYN after memory formation but returned to control levels

after forgetting. We further showed NAT10-dependent ac4C mapping of SYN mRNAs through the acRIP-seq from control and Nat10 cKO mice. In addition, we performed RT-qPCR analysis of acRIP and input RNA to verify the NAT10-dependent ac4C modification of synaptic Arc, Camk2α and Grin2b mRNAs. Genetic ablation of NAT10 in adult mouse hippocampus also reduced the protein rather than mRNA levels of Arc and Camk2α and Grin2b in the SYN at day 6 in the MWM. Lastly, the maintenance of LTP at the SC-CA1 synapses as well as the behaviors of learning and memory were severely impaired in the Nat10 cKO mice. In sum, these results demonstrate the potential contribution of NAT10-mediated ac4C modification of mRNA to synaptic plasticity and memory.

The dynamic regulation of protein acetylation in the SYN by learning and memory has been shown in our previous studies (Zhang et al., 2021a; Zhang et al., 2021b; Zhang et al., 2021c; Zhang et al., 2024). In contrast to the well-characterized acetyltransferases and deacetylases for protein acetylation (Drazic et al., 2016), the writer and eraser of mRNA acetylation are less well understood. The NAT10 acetyltransferase is the only known ac4C writer in the mammalian cells (Cui et al., 2023; Hu *et al.*, 2024a). Recently, the deacetylase SIRT7 has been shown to erase the ac4C modification of ribosomal RNA (rRNA) and small nucleolar RNA (snoRNA) (Kudrin et al., 2021; Xu et al., 2021). However, the mechanisms underlying the deacetylation of mRNA needs further exploration.

How the ac4C modification of mRNA is regulated by neural activities remains largely unknown. Our previous studies showed that overnight starvation downregulated the ac4C modification of tyrosine hydroxylase (TH) mRNA in the sympathetic nerve terminus that innervate the olfactory mucosa in mice, which reduced TH protein levels and dopamine synthesis in the mouse olfactory mucosa after overnight starvation (Zhou et al., 2022). In this study, we demonstrate that learning and memory regulate the dynamic change of ac4C modification of mRNA at synapses of mouse hippocampus. Altogether these studies provide evidence that neural activities regulate acetylation of mRNA in a spatiotemporally specific manner.

NAT10 proteins were increased in the SYN but reduced in the cytoplasm of mouse hippocampus at day 6 in the MWM. These results suggest a relocation of NAT10 from the cytoplasm to synapses after learning and memory, which was corroborated by the immunostaining study of NAT10 distribution in cultured mouse hippocampal neurons after cLTP stimulation. Previous proteomic studies showed that NAT10 were detected in the interactome with several motor proteins such as kinesin proteins (Rouillard et al., 2016), which raised the possibility that NAT10 might be transported by the motor proteins in hippocampal neurons during learning and memory. NAT10 was highly expressed in the nucleus of hippocampal neurons, but protein levels of nuclear NAT10 were not altered at day 6 in the MWM. NAT10 has been shown to be important for regulating nuclear functions such as nucleolar assembly (Shen et al., 2009) and nuclear architecture (Larrieu et al., 2014). The function of NAT10 in the nucleus of neurons waits for future investigations.

The subcellular change of NAT10 proteins at day 6 apparently may not be caused by stress during the training in the MWM. The mice were well handled before and during the training, and thus the stress should be minor if any. The control mice were exposed to the training using visible platform at day 1, the stress levels in the control mice may be vary with the learning mice. If mild stress during the training in the MWM could cause the relocation of NAT10 to the SYN, the protein levels of NAT10 in the SYN would be increased at day 1. However, the protein levels of NAT10 in the SYN were similar at day 1 between control and learning mice. In addition, chronic mild stress has been shown to increase the total protein levels of NAT10 in mouse hippocampus (Guo et al., 2022). In contrast, the total protein levels of NAT10 in mouse hippocampus did not alter at day 6 in the MWM. These results suggest that the increased NAT10 proteins in the SYN at day 6 in the MWM may be related with learning and memory rather than stress.

The stoichiometry levels of ac4C in polyA RNA has been reported to be ~0.2% in HeLa cells (Arango *et al.*, 2018), ~0.1% in human embryonic stem cells (Hu et al., 2024b), ~0.2% in liver (Hao et al., 2022), and ~0.3% in heart (Shi et al., 2023b). Here

we showed that ~0.4% of cytidine was acetylated in mouse hippocampus. Although the total protein levels of NAT10 has been reported to be relatively lower in the whole brain tissue compared with other tissues (Jiang *et al.*, 2023), the NAT10 was shown to be highly expressed in hippocampal neurons (Gao et al., 2024). Future studies are warranted to investigate how the acetyltransferase activity of NAT10 toward mRNA was regulated in different brain regions and different tissues.

The basal glutamatergic transmission at the SC-CA1 synapses were unaltered in the Arc knockout mice (Plath et al., 2006), which is consistent with the phenotypes in the Nat10 cKO mice. On the other hand, ARC has been shown to accelerate AMPA receptor endocytosis (Chowdhury et al., 2006), and thus the induction of LTP at the SC-CA1 synapses was enhanced in the Arc knockout mice (Plath *et al.*, 2006). Although ARC protein levels were decreased in the SYN of Nat10 cKO mice, the induction of LTP at the SC-CA1 synapses was not increased in the Nat10 cKO mice. These results might be due to the reduction of other LTP-promoting proteins such as CaMK2α and GluN2b in the SYN of Nat10 cKO mice.

In line with the phenotypes of Nat10 cKO mice presented in this study, the Arc knockout mice showed deficits in the late phase of LTP at SC-CA1 synapses and impairment of memory consolidation in the MWM (Plath *et al.*, 2006). Consistent with the function of Arc in memory consolidation, the acetylation of Arc mRNA was increased in the SYN at day 6 rather than day 1 following MWM training, a period during which memories were consolidated. Moreover, the increase of ac4C and protein levels of Arc in the SYN at day 6 was NAT10-dependent, which might underlie the contribution of NAT10 in learning and memory. However, the function of NAT10 in regulating tRNA or rRNA might also contribute to learning and memory. New techniques for overexpression and erasure of site-specifically acetylated mRNA need to be developed to precisely study the function of mRNA acetylation. Nonetheless, the results presented here demonstrate the dynamic regulation and potential importance of synaptic mRNA acetylation in learning and memory.

References

Arango, D., Sturgill, D., Alhusaini, N., Dillman, A.A., Sweet, T.J., Hanson, G., Hosogane, M., Sinclair, W.R., Nanan, K.K., and Mandler, M.D. (2018). Acetylation of cytidine in mRNA promotes translation efficiency. Cell *175*, 1872-1886.

Arango, D., Sturgill, D., Yang, R., Kanai, T., Bauer, P., Roy, J., Wang, Z., Hosogane, M., Schiffers, S., and Oberdoerffer, S. (2022). Direct epitranscriptomic regulation of mammalian translation initiation through N4-acetylcytidine. *Mol Cell 82*, 2797-2814.

Beiki, H., Sturgill, D., Arango, D., Relier, S., Schiffers, S., and Oberdoerffer, S. (2024). Detection of ac4C in human mRNA is preserved upon data reassessment. *Mol Cell 84*, 1611-1625. e1613.

Besse, F., and Ephrussi, A. (2008). Translational control of localized mRNAs: restricting protein synthesis in space and time. *Nat Rev Mol Cell Biol 9*, 971-980.

Biane, J.S., Ladow, M.A., Stefanini, F., Boddu, S.P., Fan, A., Hassan, S., Dundar, N., Apodaca-Montano, D.L., Zhou, L.Z., and Fayner, V. (2023). Neural dynamics underlying associative learning in the dorsal and ventral hippocampus. *Nat Neurosci 26*, 798-809.

Biever, A., Glock, C., Tushev, G., Ciirdaeva, E., Dalmay, T., Langer, J.D., and Schuman, E.M. (2020). Monosomes actively translate synaptic mRNAs in neuronal processes. *Science 367*, eaay4991.

Bliss, T.V., and Collingridge, G.L. (1993). A synaptic model of memory: long-term potentiation in the hippocampus. *Nature 361*, 31-39.

Boccaletto, P., Machnicka, M.A., Purta, E., Piątkowski, P., Bagiński, B., Wirecki, T.K., de Crécy-Lagard, V., Ross, R., Limbach, P.A., and Kotter, A. (2018). MODOMICS: a database of RNA modification pathways. 2017 update. *Nucleic Acids Res 46*, D303-D307.

Bramham, C.R., Alme, M.N., Bittins, M., Kuipers, S.D., Nair, R.R., Pai, B., Panja, D., Schubert, M., Soule, J., Tiron, A., and Wibrand, K. (2010). The Arc of synaptic memory. *Exp Brain Res 200*, 125-140.

Cajigas, I.J., Tushev, G., Will, T.J., tom Dieck, S., Fuerst, N., and Schuman, E.M. (2012). The local transcriptome in the synaptic neuropil revealed by deep sequencing and high-resolution imaging. *Neuron 74*, 453-466.

Chen, X., Cai, Q., Zhou, J., Pleasure, S.J., Schulman, H., Zhang, M., and Nicoll, R.A. (2024). CaMKII autophosphorylation is the only enzymatic event required for synaptic memory. *Proc Natl Acad Sci U S A 121*, e2402783121.

Chowdhury, S., Shepherd, J.D., Okuno, H., Lyford, G., Petralia, R.S., Plath, N., Kuhl, D., Huganir, R.L., and Worley, P.F. (2006). Arc/Arg3.1 interacts with the endocytic machinery to regulate AMPA receptor trafficking. *Neuron 52*, 445-459.

Cui, Z., Xu, Y., Wu, P., Lu, Y., Tao, Y., Zhou, C., Cui, R., Li, J., and Han, R. (2023). NAT10 promotes osteogenic differentiation of periodontal ligament stem cells by regulating VEGFA-mediated PI3K/AKT signaling pathway through ac4C modification. *Odontology 111*, 870-882.

Das, S., Lituma, P.J., Castillo, P.E., and Singer, R.H. (2023). Maintenance of a short-lived protein required for long-term memory involves cycles of transcription and local translation. *Neuron 111*, 2051-2064

Day, J.J., and Sweatt, J.D. (2011). Epigenetic mechanisms in cognition. *Neuron 70*, 813-829.

Drazic, A., Myklebust, L.M., Ree, R., and Arnesen, T. (2016). The world of protein acetylation.

Biochim Biophys Acta 1864, 1372-1401.

Eroglu, C., and Barres, B.A. (2010). Regulation of synaptic connectivity by glia. *Nature 468*, 223-231.

Fanselow, M.S., and Dong, H.-W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron 65*, 7-19.

Fischer, A., Sananbenesi, F., Wang, X., Dobbin, M., and Tsai, L.-H. (2007). Recovery of learning and memory is associated with chromatin remodelling. *Nature 447*, 178-182.

Gao, S., Shen, R., Li, J., Jiang, Y., Sun, H., Wu, X., Li, X., Miao, C., He, M., Wang, J., and Chen, W. (2024). N-acetyltransferase 10 mediates cognitive dysfunction through the acetylation of GABABR1 mRNA in sepsis-associated encephalopathy. *Proc Natl Acad Sci U S A 121*, e2410564121.

Georgeson, J., and Schwartz, S. (2024). No evidence for ac4C within human mRNA upon data reassessment. *Mol Cell 84*, 1601-1610.

Goldie, B.J., and Cairns, M.J. (2012). Post-transcriptional trafficking and regulation of neuronal gene expression. *Mol Neurobiol 45*, 99-108.

Guca, E., Alarcon, R., Palo, M.Z., Santos, L., Alonso-Gil, S., Davyt, M., de Lima, L.H., Boissier, F., Das, S., and Zagrovic, B. (2024). N6-methyladenosine in 5′ UTR does not promote translation initiation. *Mol Neurobiol 84*, 584-595.

Guo, X.-F., Wang, X.-H., Fu, Y.-L., Meng, Q., Huang, B.-Y., Yang, R., Guo, Y., Du, Y.-R., Wang, X., and Gao, Y. (2022). Elevation of N-acetyltransferase 10 in hippocampal neurons mediates depression-and anxiety-like behaviors. *Brain Res Bull 185*, 91-98.

Hafner, A.S., Donlin Asp, P.G., Leitch, B., Herzog, E., and Schuman, E.M. (2019). Local protein synthesis is a ubiquitous feature of neuronal pre-and postsynaptic compartments. *Science 364*, eaau3644.

Hao, H., Liu, W., Miao, Y., Ma, L., Yu, B., Liu, L., Yang, C., Zhang, K., Chen, Z., Yang, J., et al. (2022). N4-acetylcytidine regulates the replication and pathogenicity of enterovirus 71. *Nucleic Acids Res 50*, 9339-9354.

Hedegaard, C., Kjaer-Sorensen, K., Madsen, L.B., Henriksen, C., Momeni, J., Bendixen, C., Oxvig, C., and Larsen, K. (2013). Porcine synapsin 1: SYN1 gene analysis and functional characterization of the promoter. *FEBS Open Bio 3*, 411-420.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell 38*, 576-589.

Hoz, L.d., Martin, S.J., and Morris, R.G.M. (2004). Forgetting, reminding, and remembering: the retrieval of lost spatial memory. *PLoS Biol 2*, e225.

Hu, Z., Lu, Y., Cao, J., Lin, L., Chen, X., Zhou, Z., Pu, J., Chen, G., Ma, X., and Deng, Q. (2024a). N-acetyltransferase NAT10 controls cell fates via connecting mRNA cytidine acetylation to chromatin signaling. *Sci Adv 10*, eadh9871.

Ito, S., Akamatsu, Y., Noma, A., Kimura, S., Miyauchi, K., Ikeuchi, Y., Suzuki, T., and
Suzuki, T. (2014a). A single acetylation of 18 S rRNA is essential for biogenesis of the
small ribosomal subunit in Saccharomyces cerevisiae. *J Biol Chem 289*, 26201-26212.
Ito, S., Horikawa, S., Suzuki, T., Kawauchi, H., Tanaka, Y., Suzuki, T., and Suzuki, T.
(2014b). Human NAT10 is an ATP-dependent RNA acetyltransferase responsible for
N4-acetylcytidine formation in 18 S ribosomal RNA (rRNA). *J Biol Chem 289*, 35724-
35730.
Jarrard, L.E. (1993). On the role of the hippocampus in learning and memory in the rat.
Behav Neural Biol 60, 9-26.
Jiang, X., Cheng, Y., Zhu, Y., Xu, C., Li, Q., Xing, X., Li, W., Zou, J., Meng, L., and
Azhar, M. (2023). Maternal NAT10 orchestrates oocyte meiotic cell-cycle progression
and maturation in mice. *Nat Commun 14*, 3729.
Kouzarides, T. (2007). Chromatin modifications and their function. *Cell 128*, 693-705.
Kowalski, S., Yamane, T., and Fresco, J.R. (1971). Nucleotide sequence of the
"denaturable" leucine transfer RNA from yeast. *Science 172*, 385-387.
Kruppa, J., and Zachau, H.G. (1972). Multiplicity of serine-specific transfer RNAs of
brewer's and baker's yeast. *Biochim Biophys Acta 277*, 499-512.
Kudrin, P., Meierhofer, D., Vågbø, C.B., and Ørom, U.A.V. (2021). Nuclear RNA-
acetylation can be erased by the deacetylase SIRT7. BioRxiv, 2021.04.06.438707.
Larrieu, D., Britton, S., Demir, M., Rodriguez, R., and Jackson, S.P. (2014). Chemical
inhibition of NAT10 corrects defects of laminopathic cells. *Science 344*, 527-532.
Lau, O.D., Courtney, A.D., Vassilev, A., Marzilli, L.A., Cotter, R.J., Nakatani, Y., and
Cole, P.A. (2000). p300/CBP-associated factor histone acetyltransferase processing of
a peptide substrate: kinetic analysis of the catalytic mechanism. *J Biol Chem 275*,
21953-21959.
Levenson, J.M., O'Riordan, K.J., Brown, K.D., Trinh, M.A., Molfese, D.L., and Sweatt,
J.D. (2004). Regulation of histone acetylation during memory formation in the
hippocampus. *J Biol Chem 279*, 40545-40559.
Levenson, J.M., and Sweatt, J.D. (2005). Epigenetic mechanisms in memory formation.
Nat Rev Neurosci 6, 108-118.
Li, B., Xi, W., Bai, Y., Liu, X., Zhang, Y., Li, L., Bian, L., Liu, C., Tang, Y., Shen, L., et
al. (2023). FTO-dependent m(6)A modification of Plpp3 in circSCMH1-regulated
vascular repair and functional recovery following stroke. *Nat Commun 14*, 489.
Lisman, J., Yasuda, R., and Raghavachari, S. (2012). Mechanisms of CaMKII action in
long-term potentiation. *Nat Rev Neurosci 13*, 169-182.
Liu, R., Wubulikasimu, Z., Cai, R., Meng, F., Cui, Q., Zhou, Y., and Li, Y. (2023a).
NAT10-mediated N4-acetylcytidine mRNA modification regulates self-renewal in
human embryonic stem cells. *Nucleic Acids Res 51*, 8514-8531.
Liu, Y., Wang, X., Liu, Y., Yang, J., Mao, W., Feng, C., Wu, X., Chen, X., Chen, L., and
Dong, P. (2023b). N4-acetylcytidine-dependent GLMP mRNA stabilization by NAT10
promotes head and neck squamous cell carcinoma metastasis and remodels tumor
microenvironment through MAPK/ERK signaling pathway. *Cell Death Dis 14*, 712.
Lu, W., Man, H., Ju, W., Trimble, W.S., MacDonald, J.F., and Wang, Y.T. (2001).
Activation of synaptic NMDA receptors induces membrane insertion of new AMPA

receptors and LTP in cultured hippocampal neurons. *Neuron 29*, 243-254.

Lv, J., Liu, H., Wang, Q., Tang, Z., Hou, L., and Zhang, B. (2003). Molecular cloning of a novel human gene encoding histone acetyltransferase-like protein involved in transcriptional activation of hTERT. *Biochem Biophys Res Commun 311*, 506-513.

Lyford, G.L., Yamagata, K., Kaufmann, W.E., Barnes, C.A., Sanders, L.K., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Lanahan, A.A., and Worley, P.F. (1995). Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron 14*, 433-445.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., and Jones, A.R. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat Neurosci 13*, 133-140.

Martin, K.C., Barad, M., and Kandel, E.R. (2000). Local protein synthesis and its role in synapse-specific plasticity. *Curr Opin Neurobiol 10*, 587-592.

Merkurjev, D., Hong, W.-T., Iida, K., Oomoto, I., Goldie, B.J., Yamaguti, H., Ohara, T., Kawaguchi, S.-y., Hirano, T., and Martin, K.C. (2018). Synaptic N 6-methyladenosine (m6A) epitranscriptome reveals functional partitioning of localized transcripts. *Nat Neurosci 21*, 1004-1014.

Morris, R. (1984). Developments of a water-maze procedure for studying spatial learning in the rat. *J Neurosci Methods 11*, 47-60.

Moser, E., Moser, M.-B., and Andersen, P. (1993). Spatial learning impairment parallels the magnitude of dorsal hippocampal lesions, but is hardly present following ventral lesions. *J Neurosci 13*, 3916-3925.

Moser, M.-B., Trommald, M., and Andersen, P. (1994). An increase in dendritic spine density on hippocampal CA1 pyramidal cells following spatial learning in adult rats suggests the formation of new synapses. *Proc Natl Acad Sci U S A 91*, 12673-12675.

Newman, A.M., Liu, C.L., Green, M.R., Gentles, A.J., Feng, W., Xu, Y., Hoang, C.D., Diehn, M., and Alizadeh, A.A. (2015). Robust enumeration of cell subsets from tissue expression profiles. *Nat Methods 12*, 453-457.

Niu, M., Cao, W., Wang, Y., Zhu, Q., Luo, J., Wang, B., Zheng, H., Weitz, D.A., and Zong, C. (2023). Droplet-based transcriptome profiling of individual synapses. *Nat Biotechnol 41*, 1332-1344.

Oashi, Z., Murao, K., Yahagi, T., Von Minden, D.L., McCloskey, J.A., and Nishimura, S. (1972). Characterization of C + located in the first position of the anticodon of Escherichia coli tRNA Met as N 4 -acetylcytidine. *Biochim Biophys Acta 262*, 209-213.

Perea, G., Navarrete, M., and Araque, A. (2009). Tripartite synapses: astrocytes process and control synaptic information. *Trends Neurosci 32*, 421-431.

Plath, N., Ohana, O., Dammermann, B., Errington, M.L., Schmitz, D., Gross, C., Mao, X., Engelsberg, A., Mahlke, C., and Welzl, H. (2006). Arc/Arg3. 1 is essential for the consolidation of synaptic plasticity and memories. *Neuron 52*, 437-444.

Rouillard, A.D., Gundersen, G.W., Fernandez, N.F., Wang, Z., Monteiro, C.D., McDermott, M.G., and Ma'ayan, A. (2016). The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database (Oxford) 2016*, baw100.

Saunders, A., Macosko, E.Z., Wysoker, A., Goldman, M., Krienen, F.M., de Rivera, H., Bien, E., Baum, M., Bortolin, L., Wang, S., et al. (2018). Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. *Cell 174*, 1015-1030

Shahi, K., and Baudry, M. (1993). Glycine-induced changes in synaptic efficacy in hippocampal slices involve changes in AMPA receptors. *Brain Res 627*, 261-266.

Shen, Q., Zheng, X., McNutt, M.A., Guang, L., Sun, Y., Wang, J., Gong, Y., Hou, L., and Zhang, B. (2009). NAT10, a nucleolar protein, localizes to the midbody and regulates cytokinesis and acetylation of microtubules. *Exp Cell Res 315*, 1653-1667.

Shi, H., Zhang, X., Weng, Y.-L., Lu, Z., Liu, Y., Lu, Z., Li, J., Hao, P., Zhang, Y., Zhang, F., et al. (2018). m6A facilitates hippocampus-dependent learning and memory through YTHDF1. *Nature 563*, 249-253.

Shi, J., Yang, C., Zhang, J., Zhao, K., Li, P., Kong, C., Wu, X., Sun, H., Zheng, R., and Sun, W. (2023a). NAT10 is involved in cardiac remodeling through ac4C-mediated transcriptomic regulation. *Circ Res 133*, 989-1002.

Shrestha, P., and Klann, E. (2022). Spatiotemporally resolved protein synthesis as a molecular framework for memory consolidation. *Trends Neurosci 45*, 297-311.

Sinclair, W.R., Arango, D., Shrimp, J.H., Zengeya, T.T., Thomas, J.M., Montgomery, D.C., Fox, S.D., Andresson, T., Oberdoerffer, S., and Meier, J.L. (2017). Profiling cytidine acetylation with specific affinity and reactivity. *ACS Chem Biol 12*, 2922-2926.

Srinivasan, R., Lu, T.Y., Chai, H., Xu, J., Huang, B.S., Golshani, P., Coppola, G., and Khakh, B.S. (2016). New Transgenic Mouse Lines for Selectively Targeting Astrocytes and Studying Calcium Signals in Astrocyte Processes In Situ and In Vivo. *Neuron 92*, 1181-1195.

Stern, L., and LH, S. (1978). The role of the minor base N4-acetylcytidine in the function of the Escherichia coli noninitiator methionine transfer RNA. *J Biol Chem 17,6132-6139.*

Steward, O., Wallace, C.S., Lyford, G.L., and Worley, P.F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron 21*, 741-751.

Su, J., Huang, F., Tian, Y., Tian, R., Qianqian, G., Bello, S.T., Zeng, D., Jendrichovsky, P., Lau, C.G., and Xiong, W. (2023). Entorhinohippocampal cholecystokinin modulates spatial learning by facilitating neuroplasticity of hippocampal CA3-CA1 synapses. *Cell Rep 42,113467.*

Sutton, M.A., and Schuman, E.M. (2006). Dendritic protein synthesis, synaptic plasticity, and memory. *Cell 127*, 49-58.

Tan, G.H., Liu, Y.Y., Hu, X.L., Yin, D.M., Mei, L., and Xiong, Z.Q. (2011). Neuregulin 1 represses limbic epileptogenesis through ErbB4 in parvalbumin-expressing interneurons. *Nat Neurosci 15*, 258-266.

Tang, Y.P., Shimizu, E., Dube, G.R., Rampon, C., Kerchner, G.A., Zhuo, M., Liu, G., and Tsien, J.Z. (1999). Genetic enhancement of learning and memory in mice. *Nature 401*, 63-69.

Thalalla Gamage, S., Sas-Chen, A., Schwartz, S., and Meier, J.L. (2021). Quantitative nucleotide resolution profiling of RNA cytidine acetylation by ac4C-seq. *Nat Protoc 16*, 2286-2307.

Ting, A.K., Chen, Y., Wen, L., Yin, D.M., Shen, C., Tao, Y., Liu, X., Xiong, W.C., and Mei, L. (2011). Neuregulin 1 promotes excitatory synapse development and function in GABAergic interneurons. *J Neurosci 31*, 15-25.

Tsien, J.Z., Chen, D.F., Gerber, D., Tom, C., Mercer, E.H., Anderson, D.J., Mayford, M., Kandel, E.R., and Tonegawa, S. (1996). Subregion-and cell type–restricted gene knockout in mouse brain. *Cell 87*, 1317-1326.

Tullis, J.E., Larsen, M.E., Rumian, N.L., Freund, R.K., Boxer, E.E., Brown, C.N., Coultrap, S.J., Schulman, H., Aoto, J., Dell'Acqua, M.L., and Bayer, K.U. (2023). LTP induction by structural rather than enzymatic functions of CaMKII. *Nature 621*, 146-153.

Vorhees, C.V., and Williams, M.T. (2006). Morris water maze: procedures for assessing spatial and related forms of learning and memory. *Nat Protoc 1*, 848-858.

Wang, G., Zhang, M., Zhang, Y., Xie, Y., Zou, J., Zhong, J., Zheng, Z., Zhou, X., Zheng, Y., and Chen, B. (2022). NAT10-mediated mRNA N4-acetylcytidine modification promotes bladder cancer progression. *Clin Transl Med 12*, e738.

Wu, D., Bacaj, T., Morishita, W., Goswami, D., Arendt, K.L., Xu, W., Chen, L., Malenka, R.C., and Südhof, T.C. (2017). Postsynaptic synaptotagmins mediate AMPA receptor exocytosis during LTP. *Nature 544*, 316-321.

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., and Zhan, L. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb) 2,100141*.

Xu, C., Zhang, J., Zhang, J., and Liu, B. (2021). SIRT7 is a deacetylase of N4-acetylcytidine on ribosomal RNA. *Genome Instab Dis 2*, 253-260.

Xu, T., Wang, J., Wu, Y., Wu, J.Y., Lu, W.C., Liu, M., Zhang, S.B., Xie, D., Xin, W.J., and Xie, J.D. (2023). Ac4C Enhances the Translation Efficiency of Vegfa mRNA and Mediates Central Sensitization in Spinal Dorsal Horn in Neuropathic Pain. *Adv Sci 10*, 2303113.

Yin, D.-M., Chen, Y.-J., Lu, Y.-S., Bean, J.C., Sathyamurthy, A., Shen, C., Liu, X., Lin, T.W., Smith, C.A., and Xiong, W.-C. (2013). Reversal of behavioral deficits and synaptic dysfunction in mice overexpressing neuregulin 1. *Neuron 78*, 644-657.

Yu, F., Zhang, Y., Cheng, C., Wang, W., Zhou, Z., Rang, W., Yu, H., Wei, Y., Wu, Q., and Zhang, Y. (2020). Poly (A)-seq: A method for direct sequencing and analysis of the transcriptomic poly (A)-tails. *PLoS One 15*, e0234696.

Yu, Q., Liu, Y.-Z., Zhu, Y.-B., Wang, Y.-Y., Li, Q., and Yin, D.-M. (2019). Genetic labeling reveals temporal and spatial expression pattern of D2 dopamine receptor in rat forebrain. *Brain Struct Funct 224*, 1035-1049.

Zhang, H.-L., Han, W., Du, Y.-Q., Zhao, B., Yang, P., and Yin, D.-M. (2021a). SRC3 acetylates calmodulin in the mouse brain to regulate synaptic plasticity and fear learning. *J Biol Chem 297,* 101044.

Zhang, H.-L., Zhao, B., Han, W., Sun, Y.-B., Yang, P., Chen, Y., Ni, D., Zhang, J., and Yin, D.-M. (2021b). Acetylation of calmodulin regulates synaptic plasticity and fear learning. *J Biol Chem 297,* 101034.

Zhang, H.-L., Zhao, B., Yang, P., Du, Y.-Q., Han, W., Xu, J., and Yin, D.-M. (2021c). Steroid receptor coactivator 3 regulates synaptic plasticity and hippocampus-dependent

memory. *Neurosci Bull 37*, 1645-1657.

Zhang, H.L., Hu, S., Yang, P., Long, H.C., Ma, Q.H., Yin, D.M., and Xu, G.Y. (2024).

HDAC9-mediated calmodulin deacetylation induces memory impairment in

Alzheimer's disease. *CNS Neurosci Ther 30*, e14573.

Zhang, M., Yang, K., Wang, Q.-H., Xie, L., Liu, Q., Wei, R., Tao, Y., Zheng, H.-L., Lin,

N., and Xu, H. (2023a). The cytidine N-acetyltransferase NAT10 participates in

peripheral nerve injury-induced neuropathic pain by stabilizing SYT9 expression in

primary sensory neurons. *J Neurosci 43*, 3009-3027.

Zhang, W., Gao, J., Fan, L., Wang, J., He, B., Wang, Y., Zhang, X., and Mao, H. (2023b).

ac4C acetylation regulates mRNA stability and translation efficiency in osteosarcoma.

Heliyon *9*.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,

C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based Analysis of

ChIP-Seq (MACS). *Genome Biol 9*, R137.

Zhou, H.-Q., Zhuang, L.-J., Bao, H.-Q., Li, S.-J., Dai, F.-Y., Wang, P., Li, Q., and Yin,

D.-M. (2022). Olfactory regulation by dopamine and DRD2 receptor in the nose. *Proc*

Natl Acad Sci U S A 119, e2118570119.

Zhuang, M., Geng, X., Han, P., Che, P., Liang, F., Liu, C., Yang, L., Yu, J., Zhang, Z.,

Dong, W., and Ji, S.-J. (2023). YTHDF2 in dentate gyrus is the m6A reader mediating

m6A modification in hippocampus-dependent learning and memory. *Mol Psychiatry 28*,

1679-1691.

Methods

Animals

Adult male mice were used in this study to eliminate the possible effects of the hormone

cycle. For stereotaxic injection of AAV, six-week-old mice were used. For behavioral

analysis, nine-week-old mice were used. For ac4C dot-blots, LC-MS/MS, acRIP-seq,

mice aged between nine and ten weeks were used. For immunofluorescence, western

blots and electrophysiology, ten-week-old mice were used. Animals were housed in

rooms at 23°C and 50% humidity in a 12 h light/dark cycle and with food and water

available ad libitum. Animal experimental procedures were approved by the

Institutional Animal Care and Use Committee of East China Normal University. The

astrocyte-specific Cre (Aldh1l1-CreER) was purchased from Jackson lab (Stock No.

029655).

Generation of Nat10$^{Cre/+}$ knockin mice

The mouse Nat10 gene has 29 exons and the translational start codon (ATG) is localized

in exon 2. We employed CRISPR/Cas9 technology in conjunction with homologous

recombination to insert a Cre-WPRE-polyA expression cassette at the translational start

codon of the Nat10 gene in C57BL/6J mice. The process commenced with the synthesis of Cas9 mRNA and a guide RNA (gRNA) targeting the start codon of Nat10, with the gRNA sequence being 5′-catcatgaatcggaagaaggtgg-3′. A donor vector was then constructed using In-Fusion cloning, containing 2.6 kb 5′ and 3′ homology arms flanking the Cre-WPRE-pA cassette, designed for Cre recombinase expression, enhanced by the WPRE element and concluded with a polyadenylation signal. The cocktail of Cas9 mRNA, gRNA, and the donor vector was microinjected into fertilized C57BL/6J mouse embryos, leading to the birth of F0 generation mice. These mice were screened for the precise genomic integration of the cassette through PCR amplification and sequencing. Successful modifications were observed in F0 mice, which were subsequently bred with C57BL/6J mice to secure F1 progeny harboring the modification.

The resulting offspring were genotyped using PCR, with primers designed to detect the Cre within the Nat10 gene. The primers for genotyping wt Nat10 allele are as follows: forward: 5'tgtctttcctgtggtggtgt3'; reverse: 5'tgcgcactgaaaccctacaa3'. the primers for genotyping Nat10-cre allele are as follows: forward: 5'gaaatcatgcaggctggtgg3'; reverse: 5'aaagtcccggaaaggagctg3'. The PCR products for wt Nat10 and Nat10-Cre alleles were 392 and 393 bp, respectively.

Generation of Nat10^{flox/flox} mice

The generation of Nat10^{flox/flox} mice involved targeted modification of exon 4 within the Nat10 gene, utilizing CRISPR/Cas9-mediated gene editing. Two guide RNAs (gRNAs) were specifically designed to target and flank exon 4, enabling a precise floxed modification. The sequences for these gRNAs were as follows: gRNA1: 5′-tcagcagaccactttaaagatgg-3′ and gRNA2: 5′-tgcctgacatagtaaggtctggg-3′. Both Cas9 mRNA and gRNAs were produced through in vitro transcription techniques. A donor vector for homologous recombination was constructed using In-Fusion cloning. This vector included a 3.0 kb 5′ homology arm, a 0.7 kb loxp-flanked region (flox), and a 3.0 kb 3′ homology arm to ensure accurate genomic insertion at the Nat10 locus. The mixture of Cas9 mRNA and the two gRNAs was injected into the cytoplasm of one-cell stage embryos. These embryos were subsequently implanted into pseudopregnant females and carried to term.

The resulting offspring were genotyped using PCR, with primers designed to detect

the floxed exon 4 of Nat10 gene. The primers for genotyping the wt and floxed Nat10

allele are as follows: forward: 5'tgccgaggggatgtactcat3'; reverse:

5'agaaccccacgaactgtcct3'. The PCR products for wt Nat10 and floxed Nat10 allele were

302 and 356 bp, respectively.

Morris water maze (MWM)

The Morris Water Maze (MWM), a widely recognized method for assessing spatial

learning and memory (Vorhees and Williams, 2006), was implemented with the

following protocol: mice were acclimated to the testing environment over a period of

three days through 10-minute daily handling sessions. To aid navigation, visual cues

were strategically positioned around the perimeter of the maze. To ensure consistency

and minimize the impact of external variables, all experimental sessions were

conducted simultaneously each day. During the core phase of the experiment, spanning

five days, mice were subjected to four trials daily. In each trial, they were released from

different quadrants into the maze, with the escape platform being fixed in the first

quadrant. A trial was deemed complete when a mouse successfully located the platform

within a 60-second timeframe. If unsuccessful, the mouse was gently guided to the

platform at the 60-second mark. The mice were handled for 10 minutes each day

following the completion of the four trials. On the sixth day, a probe test was conducted,

during which the platform was removed from the maze. During this test, mice were

released from the third quadrant, and the number of times they crossed the former

platform location during a 60-second trial was recorded as a measure of their retained

spatial memory. Following the probe test, the mice were handled for a further 10

minutes before being returned to their home cages. After a two-week period in the home

cages, the mice were subjected to the probe test again.

Chemical and reagents

All chemicals and reagents utilized in this study were of analytical grade. RNA was

extracted using TRIzol® Reagent (15596-026, Invitrogen, Carlsbad, CA, USA)

following the manufacturer's protocol. Chloroform and other general laboratory

chemicals were obtained from Merck (109634, Darmstadt, Germany). The ACQUITY

HSS T3 column (2.1×100 mm, 1.8 μm) for LC-MS/MS analysis was purchased from

Waters Corporation (186008499, Milford, MA, USA). Phosphodiesterase was obtained

from Sigma (P3243-1VL, St. Louis, MO, USA). Nuclease P1 and alkaline phosphatase, used for RNA digestion, were obtained from Takara (2410A and 2250A). Hydroxylamine was from Aladdin Bio-Chem Technology Co., LTD (H164487). RNase inhibitor, which is critical for preserving RNA integrity during manipulations, was purchased from New England Biolabs (M0314S, NEB, CA, USA). Methylene blue, which is used for nucleic acid visualization, was obtained from Abcam (M9140). Poly-D-lysine was purchased from Sigma (Cat No. P1149). Tamoxifen was purchased from Sigma (Cat T5648).

Subcellular fractionation

The procedure of subcellular fractionation was adapted from the methodologies in our previous studies (Zhang *et al.*, 2021a; Zhang *et al.*, 2021b). Mouse brain tissues were processed starting with homogenization in Buffer A, which consists of 0.32 M sucrose, 1 mM $MgCl_2$, 1 mM PMSF, and a protease inhibitor cocktail to ensure comprehensive cell lysis while preserving protein integrity. The resultant homogenates were then filtered to remove cell debris and subsequently subjected to centrifugation at 500g for 5 minutes using a fixed-angle rotor. This step separates the homogenate into the pellet (P1) and supernatant (S1) fractions. The P1 fraction, containing primarily nuclei and unbroken cells, was further purified by resuspension in Buffer B (10 mM KCl, 1.5 mM $MgCl_2$, and 10 mM Tris-HCl at pH 7.4) followed by a repeat centrifugation under the same conditions to remove any remaining debris. The final pellet was resuspended in Buffer C (20 mM HEPES, pH 7.9, 25% glycerol, 1.5 mM $MgCl_2$, 1.4 M KCl, 0.2 mM EDTA, 0.2 mM PMSF, and 0.5 mM DTT), which is designed for nuclear protein extraction and incubated with agitation at 4 °C for 30 minutes to facilitate the release of nuclear proteins. A final centrifugation at 12,000g for 10 minutes was performed, and the supernatant, containing the nuclear protein fraction, was collected. The initial supernatant (S1), following the removal of P1, was further centrifuged at 10,000g for 10 minutes to partition it into the P2 fraction, which included the crude synaptosomes, and the cytosolic S2 fraction.

Western blot

Homogenates of hippocampus tissue were prepared in the RIPA buffer containing 50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 2 mM EDTA, 1% sodium deoxycholate, 1%

SDS, 1 mM PMSF, 50 mM sodium fluoride, 1 mM sodium vanadate, 1 mM DTT, and

protease inhibitors cocktails. All the protein samples were boiled at 100 °C water bath

for 10 minutes before western blot. Homogenates were resolved on SDS/PAGE and

then transferred to nitrocellulose membranes, which were incubated in the TBS buffer

(0.1% Tween-20 and 5% milk) for 1 hour at room temperature followed by incubation

of primary antibodies overnight at 4 °C. After washing, the membranes were incubated

with HRP-conjugated secondary antibody in the same TBS buffer for 2 hours at room

temperature. Immunoreactive bands were visualized by ChemiDocTM XRS + Imaging

System (BIO-RAD) using enhanced chemiluminescence (Pierce) and analyzed with

Image J (NIH). The following primary antibodies were used: anti-Histone H3 (1:1000,

Cell Signaling, 9715), anti-SYN1 (1:1000, Cell Signaling, 2312), anti-α-tubulin

(1:10000, Cell Signaling, 3873), anti-NAT10 (1:1000, Abcam, ab194297), anti-Acetyl-

Histone H3 (1:1000, Cell Signaling, 9649), anti-ARC (1:1000, Proteintech, 66550-1-

Ig), anti-PSD95 (1:1000, Cell Signaling, 3450), anti-PCAF (1:1000, Cell Signaling,

3378), and anti-GAPDH (1:5000, Abways, ab0037). The following secondary

antibodies were used: HRP-conjugated secondary antibody (goat-anti-mouse, G-21040;

goat-anti-rabbit, G-21234; 1:2000, Thermo Fisher).

1000

Immunofluorescence

The mouse brains were fixed in 4% paraformaldehyde at room temperature for 1 hour

and dehydrated in 20% sucrose at 4 °C overnight. Then the tissues were placed in an

embedding mold and frozen in OCT media at 0 °C. The embedded tissues were cut into

slices with the thickness of 30 μm using a Leica microtome (CM3050S). The brain

slices were permeabilized with 0.3% Triton-X 100 and 5% BSA in PBS and incubated

with primary antibodies at 4 °C overnight. After washing with PBS for three times,

samples were incubated with secondary antibodies for 2 hours at room temperature.

Samples were mounted with Vectashield mounting medium (Vector Labs) and images

were taken by Leica TCS SP8 confocal microscope. The following primary antibodies

were used: rabbit anti-NeuN (1:1000, Abcam, ab177487), rabbit anti-Neurogranin

(1:500, Abcam, ab217672). The following secondary antibodies were employed:

Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A32732, 1:500,

Thermo Fisher). Unbiased stereology TissueFAX Plus ST (Tissue Gnostics, Vienna,

Austria) was applied to count the number of Nat10-positive neurons. The detailed

methods for cell number counting is available on the website
(https://tissuegnostics.com/products/single-cell-analysis/tissuequest).

1018

Detection of ac4C and m6A modifications by LC-MS/MS

The analysis of LC-MS/MS followed previous studies with some modification (Arango
et al., 2018). Briefly, the RNA samples starting with 2 μg, were digested into
nucleosides using alkaline phosphatase and nuclease P1 at 37°C for 3 hours. The
digested samples were then subjected to chloroform extraction, with the aqueous phase
being utilized for LC-MS/MS analysis in an AB SCIEX QTRAP 6500+ system,
featuring an ACQUITY HSS T3 column (2.1×100 mm, 1.8 μm). Chromatographic
separation was achieved with a mobile phase gradient of 2 mM ammonium bicarbonate
in ultrapure water (A) and methanol (B), starting with 95% A and transitioning to 5%
A over 8 minutes. A calibration curve spanning 0.0005 to 1000 pmol standard samples
facilitated the accurate quantification of the epitranscriptomic modifications.

1030

ac4C and m6A dot-blot

The total RNA of mouse hippocampus was extracted with Trizol according to the
manufacturer's instructions. The polyA RNA was then purified using the Dynabeads^{TM}
polyA RNA purification kit (61006, Thermo Fisher). The dot-blot was performed
according to the previous study (Zhang et al., 2023a). We used 0.5 μg of polyA RNA
that fall in the linear range for each blot. For each HOM replicate, we purified 0.5 μg
of polyA RNA from the hippocampus of one mouse. In contrast, for each SYN replicate,
we purified 0.5 μg of polyA RNA from the hippocampus of five mice. In brief, the
polyA RNA was denatured at 95 °C for 5 min using a heat block, immediately kept on
ice for 1 min, loaded onto Amersham Hybond-N+ membrane and crosslinked with
UV254 for 30 min. The membranes were blocked with 5% nonfat milk for 2 h at room
temperature (RT), and incubated with the rabbit anti-m6A (1:2000, Synaptic System,
#202003) or anti-ac4C antibody (1:500, Abcam, ab252215) at 4°C overnight. After
wash with 0.05% PBST for three times, the membranes were incubated with the HRP-
conjugated secondary antibody (goat-anti-rabbit, G-21234, 1:2000, Thermo Fisher) at
RT for 2 h. The signals were visualized by ChemiDocTM XRS + Imaging System
(BIO-RAD) using ECL and analyzed with Image J (NIH). RNA loading was verified
by staining the membranes with 0.2% methylene blue. The intensity of each dot was

normalized to total RNA.

1050

Primary neuronal culture

The primary culture of mouse hippocampal neurons were performed according to our

previous studies (Ting et al., 2011). Hippocampal explants from embryonic day 16 (E16)

mouse embryos were digested with 0.125% trypsin solution for 30 min at 37°C,

followed by trituration with pipette in plating medium (DMEM with 10% fetal bovine

serum, 10% F-12, and 1% penicillin-streptomycin). The dissociated neurons were

plated onto poly-D-lysine-coated coverslips at a density of 1×10^5 cells per 35 mm dish.

Following a 4-hour incubation period, the plating medium was replaced with

Neurobasal^{TM} medium (Thermo Fisher, 21103049) supplemented with 2% B-27^{TM}

(Thermo Fisher, 17504044), 1% GlutaMax^{TM} (Thermo Fisher, 35050061), and 1%

penicillin-streptomycin (Invitrogen). Cytarabine (Sigma, C1768-100MG) was added

after 2 days to a final concentration of 1 μM, to inhibit the proliferation of dividing non-

neuronal cells. Half volume of the culture medium was replaced every 3-4 days with

freshly prepared medium. The neurons were harvested at 14 days in vitro (DIV 14) for

immunofluorescence experiments.

1066

Chemical LTP (cLTP) and immunofluorescence

For cLTP, neurons were incubated with 300 μM glycine in a cell incubator set at 37°C

with 5% CO_2 and 95% humidity for 3 minutes, followed by an additional 60 minutes

without glycine at 37°C (Wu et al., 2017). Neurons were then fixed with 4%

paraformaldehyde for 30 minutes on ice to preserve cell membranes. After washing

with PBS, the neurons were penetrated with 0.3% Triton-X100 and blocked with 1%

BSA in PBS for 30 min at RT. After that, neurons were incubated with primary

antibodies (chicken anti-MAP2 antibody, 1:2000, Abcam, ab5392; mouse monoclonal

anti-PSD95 antibody, 1:800, Abcam, ab192757; and rabbit monoclonal anti-NAT10

antibody, 1:800, Abcam, ab194297) overnight at 4°C. Neurons were then washed by

PBS for three times before incubation with fluorescent secondary antibodies for 2 hours

at 37°C. The following secondary antibodies were employed: Chicken IgY (H+L)

Secondary Antibody (A-11039, 1:500, Thermo Fisher), Rabbit IgG (H+L) Cross-

Adsorbed Secondary Antibody (A-21244, 1:500, Thermo Fisher), Rabbit IgG (H+L)

Highly Cross-Adsorbed Secondary Antibody (A32732, 1:500, Thermo Fisher), and

Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A32727, 1:500,

Thermo Fisher). After washing with PBS, the neurons were placed on mounting

solution with DAPI (Beyotime, P0131). Confocal stacks were acquired from 3-6

individual fields on multiple coverslips per culture using a Leica SP8 confocal

microscope. Two independent cultures were analyzed for each condition, including

vehicle-treated control and glycine-treated neurons.

1088

Regular acRIP-seq for HOM

The acRIP-seq was performed following the previous studies (Arango *et al.*, 2018;

Thalalla Gamage et al., 2021). The total RNA was extracted from the HOM of mouse

hippocampus using TRIzol, the quality of the RNA was evaluated through gel

electrophoresis in order to ascertain its integrity and purity, and the concentration and

purity of the RNA were detected. Total RNA samples, initially 100 μg, were treated

with the rRNA depletion kit (New England Biolabs, Cat. No. E6310) to remove rRNA.

After rRNA depletion, the RNA was subjected to immunoprecipitation using the anti-

ac4C antibodies and the ac4C RIP kit (GenSeq, Cat No. GS-ET-005) according to the

manufacturer's instructions. Briefly, total RNAs were randomly fragmented to ~200 nt

by RNA Fragmentation Reagents (NEB, Cat No. M0348S). Protein A/G Dynabeads

(Share-Bio, SB-PR001) were coupled to the anti-ac4C antibody by rotating at room

temperature for 1 h. The RNA fragments were then incubated with the beads-linked

anti-ac4C antibodies (2 μg) in the 100 μl acRIP buffer (PBS containing 1% Triton X-

100, 2 mM BSA, and 1 mM RNase inhibitor) and rotated at 4°C for 4 h. After incubation,

the RNA/antibody complexes were washed for four times with the acRIP buffer, and

then RNAs were eluted by elution buffer (containing 0.1 M Glycine, 0.5% Triton X-

100, PH 2.5-3.1) at room temperature for 10 minutes. Finally, add NaOH solution to

adjust the pH of the elution product to neutral. Both the immunoprecipitated RNA and

1% input RNA are reverse transcribed with random hexamers, followed by RNA library

preparation with Ultra™ II Directional RNA Library Prep Kit (New England Biolabs,

Cat No. E7760) according to the manufacturer's instructions.

The quality of RNA libraries were assessed using Agilent 2100 bioanalyzer and

then subjected to the next-generation sequencing through the NovaSeq 6000 platform

(Illumina). The quality of the sequencing data is measured using Q30 scores, and low-

quality reads and 3' adaptors are removed using Cutadapt (v1.9.3). High-quality reads

are then aligned to the reference genome using Hisat2 (v2.0.4). Raw counts are generated using HTSeq (v0.9.1) and normalized using edgeR (v3.16.5). Gene expression was quantified by logged transcript per million mapped reads (log$_2$TPM).

Low-input acRIP-seq for SYN

The low-input acRIP-seq included some modifications, compared with the regular acRIP-seq. First, the RNA amount is relatively smaller (~ 1 µg, purified from the hippocampus of 10 mice) for low-input acRIP-seq, compared with regular acRIP-seq (~ 100 µg, purified from the hippocampus of 2 mice). Second, the total RNA purified from the SYN was not treated by the rRNA depletion kit before acRIP due to the low amount of rRNA in the SYN. Third, the RNA libraries for low-input acRIP-seq were generated by a different kit - SMARTer Stranded Total RNA-Seq Kit v2-Pico Input Mammalian (Clontech, Cat No. 050619). To increase the data reliability for low-input acRIP-seq, transcripts detected in at least two of the three databases of synaptic transcriptome (Cajigas et al., 2012; Hafner et al., 2019; Niu et al., 2023) were considered to be reliable mRNA expressed in the SYN.

Identification of ac4C peaks

Following RNA integrity assessments, acRIP enrichment verification, and library quality checks, raw sequencing reads from both acRIP and input samples underwent rigorous quality control. Initially, sequencing quality was evaluated using Q30 scores and FastQC to ensure base-level accuracy. Low-quality reads and 3' adaptor sequences were identified and removed using Cutadapt software (v1.9.3), resulting in high-quality clean reads for further analysis.

Post-alignment filtering was applied to exclude reads that mapped to mitochondrial DNA (chrM) and non-concordant mate pairs, which often arise from sequencing artifacts or alignment errors. The remaining clean reads were then aligned to the mouse genome (mm10) using Hisat2 software (v2.0.4), with parameters allowing a maximum of five mismatches per read to ensure alignment accuracy above 95%. Ensembl gene annotations (Release 75) were incorporated to guide accurate mapping to annotated regions. To account for transcript variability, canonical transcript sequences from the UCSC genome browser were used to generate a Bowtie2 index, enabling transcriptome mapping. Reads were aligned in local mode, which is well-suited for fragmented RNA

sequences.

Acetylation peaks (ac4C peaks) were identified using MACS software (v1.4.2)

(Zhang et al., 2008), optimized for transcriptomic data. The shifting model and local

lambda options were disabled to accommodate the unique characteristics of

transcriptome data, and the genome size was set according to the total transcript base

count. Input samples were used as controls to distinguish true ac4C-enriched peaks

from background noise. The key aspect of this process was comparing the enrichment

of clean reads between acRIP and input samples. Regions where acRIP samples

exhibited significantly higher read enrichment compared to input samples were

identified as potential acetylation peaks, reflecting areas where ac4C modifications are

likely to occur. We established strict screening criteria in the peak calling process:

adjusted $P < 0.05$ and a fold enrichment (FE) > 2. The ac4C peaks can be generated and

visualized on the UCSC genome browser (https://genome.ucsc.edu/cgi-bin/hgGateway)

using IGV (http://www.igv.org/). The FE was calculated as the ratio of acRIP to input.

$$FE = \frac{acRIP}{Input}$$

1162

1163

Sequence motif analysis

1164

1165 Sequence motifs were identified using HOMER (Heinz et al., 2010) and visualized

1166 using ggseqlogo package from R software. HOMER identifies motifs that are

1167 specifically enriched on learning set of sequences relative to control (e.g., peaks

1168 upstream and downstream sequences relative to the entire genome sequence) based

1169 primarily on the principle of ZOOPS scoring with hypergeometric enrichment.

1170

Gene ontology and functional category analysis

1171

Gene ontology (GO) analysis to detect enrichment for GO terms in differential

acetylation genes in R using the clusterProfiler (Wu et al., 2021), and the possible roles

of these differential acetylation genes were annotated according to MF, BP and CC. GO

terms with a false-discovery rate adjusted P-value of < 0.05 were visualized using R

scripts to plot. The R package clusterProfiler to visualize the genes composing KEGG

pathways, and P-value < 0.05 was used as the threshold of significant enrichment.

1178

Analysis of ac4C fold enrichment (FE) for individual mRNA

1179

1180 The fold enrichment of ac4C mRNA was reflected by the cumulative value of multiple

ac4C peaks from one particular mRNA molecule. Aggregating the cumulative sum of these peaks offers a more comprehensive overview of the total abundance of ac4C modifications across the mRNA molecule, avoiding the narrow focus on a potentially non-representative, single maximum modification site, similar with the previous study of m6A (Guca et al., 2024). Additionally, in instances where ac4C modifications on mRNA are more dispersed or present in low abundance, a single prominent peak may not be observable; instead, multiple low-abundance peaks might emerge. In such scenarios, the accumulation of multiple peaks can enhance detection sensitivity, better capture dynamic changes, and mitigate biases inherent in individual experiments. The following quantification index was used to calculate the ac4C levels of individual mRNA:

$$FE \ of \ ac4C = \Sigma peak \ (1 + 2 + 3 \ldots \ldots + n)$$

Analysis of ac4C fold change (FC) for individual mRNA

The fold change of ac4C mRNA is representative of the degree of change in the overall acetylation level of mRNA in the learning group (L) compared with the control group (C). To prevent a value of 0 in the denominator in the FC calculation, we added 1 to the FE of ac4C. The FC of ac4C mRNA was calculated using the following quantification index:

$$FC \ of \ ac4C = \frac{(FE \ of \ ac4C + 1) \ in \ L}{(FE \ of \ ac4C + 1) \ in \ C}$$

Analysis of PPI network

The STRING (https://string-db.org/) (version 12.0) knowledge base was used to create a protein-protein interaction (PPI) network. The PPI network was generated with a confidence score of 0.7 and visualized using Cytoscape (version 3.10.1). Molecular Complex Detection (MCODE) (v2.0.3) algorithm was employed to identify the components of the densely connected network.

The following software and algorithms were used for the quantification and statistical analysis of ac4C epitranscriptome.

Software and algorithms, application, authors		
R v4.3.2	Statistical computing and graphics.	R Core Team. R Foundation for

		Statistical Computing 2023.
Python v3.11	Object-oriented programming language.	Van R et al. CreateSpace 2009.
ggplot2	Creating graphics, based on The Grammar of Graphics.	Hadley W et al, Springer-Verlag New York 2016.
ggseqlogo	Generating publication-ready sequence logos using ggplot2.	Wagih et al. Bioinformatics 2018.
tidyverse	Collection of R packages designed for data science.	Wickham et al. Journal of Open Source Software 2019.
ComplexHeatmap	Arrange multiple heatmaps and supports annotation graphics.	Gu Z et al. Bioinformatics 2016.
HOMER v4.11	Motif Discovery and gene sequencing analysis.	Heinz S et al. Mol Cell 2010.
bedtools	Compare large sets of genomic features.	Aaron Q et al. Bioinformatics 2010.
cutadapt v1.9.3	Remove sequencing data adaptor.	Martin M. EMBnet 2011.
Hisat2 v2.04	Mapping sequencing reads to a single reference genome.	Zhang Y et al. Genome Research 2021.
MACS v1.4.2	Identifying transcript factor binding sites.	Yong Z et al. Genome Biology 2008.
Pandas v2.1.4	Cleaning data.	The pandas development team. Zenodo 2020.
IGV	Interactive genome visualization application.	James T et al. Bioinformatics 2023.
HTSeq (v0.9.1)	Analysing high-throughput sequencing data.	Simon A et al. Bioinformatics 2014.
edgeR (v3.16.5)	Empirical analysis of digital gene expression data.	McCarthy DJ et al. Nucleic Acids Research 2012.
CIBERSORT	Cell-type identification by estimating relative subsets of RNA transcripts.	Aaron M et al. Nature methods 2015.

1211

Identification of relative abundance of genes in different cell types

We applied the CIBERSORT (Cell-type Identification by Estimating Relative Subsets of RNA Transcripts) algorithm (Newman et al., 2015) in R to determine the enriched cell types where the genes of NASA mRNAs were expressed. To construct the reference gene signature matrices, we used Dataset C from McCarroll Lab (http://dropviz.org/) (Saunders *et al.*, 2018) to extract gene expression features for various hippocampal cell types, including excitatory neurons, inhibitory neurons, astrocytes, and microglia. These reference matrices were then used to deconvolve the gene expression data and estimate the contributions from each cell type.

RT-qPCR

The cDNA was obtained through reverse transcription using the HiScript III RT SuperMix for qPCR kit (Vazyme, R323-01), followed by amplification using the ChamQ Universal SYBR qPCR Master Mix kit (Vazyme, Q711-02). Gene expression levels measured with RT-PCR were normalized to Gapdh. The primers used in qPCR for this study were as follows: Gapdh-forward, 5'gggtgtgaaccacgagaaat3'; Gapdh-reverse 5'actgtggtcatgagcccttc3'; Arc-forward, 5'ggagggaggtcttctaccgtc3'; Arc-reverse 5'cccccacacctacagagaca3'; Grin2b-forward, 5'ctggtgaccaatggcaagcatg3'; Grin2b-reverse, 5'ggcacagagaagtcaaccacct3'; Camk2α-forward, 5'agccatcctcaccactatgctg3'; Camk2α-reverse 5'gtgtcttcgtcctcaatggtgg3'.

Stereotaxic injection of AAV into mouse hippocampus

AAV2/9-Syn1-EGFP and AAV2/9-Syn1-EGFP-P2A-Cre were generated by Hanbio Technology (Shanghai) Corp., Ltd., with a titer of 10^{12}/μl. We injected 0.3 μl AAV into each side of the mouse hippocampus. Six-week-old Nat10$^{\text{flox/flox}}$ mice received an intraperitoneal injection of alphaxalone at a dosage of 40 mg/kg for anesthesia and then were immobilized in a stereotaxic frame provided by RWD Life Science. Similarly, AAV2/9-CAG-DIO-EGFP, generated by Hanbio Technology (Shanghai) Corp., Ltd., with a titer of 10^{12}/μl, was injected into the hippocampus of seven-week-old Nat10$^{\text{Cre/+}}$ and ALDH1L1$^{\text{Cre/ERT2}}$ mice. The AAVs were delivered into mouse hippocampus through a Stoelting stereotaxic injector at a flow rate of 0.1 μl/min. The precise coordinates for the stereotaxic injection were set to anteroposterior (AP) −2.3 mm, dorsoventral (DV) −1.8 mm, and mediolateral (ML)±1.75 mm, relative to the Bregma.

1245 To induce Cre enzyme expression, mice were fed a custom-made tamoxifen-infused

1246 diet over a period of one week, with the administration occurring on an intermittent

1247 schedule (Tan et al., 2011).

1248

Electrophysiological recording

Hippocampal slices were prepared as previously described (Yin et al., 2013). Briefly,

mice (10 weeks old, male) were decapitated and transverse hippocampal slices (400 μm)

were prepared using a Vibroslice (VT 1000S; Leica) in ice-cold ACSF. For field

potential recording, the same ACSF was used for slice preparation and recording: 120

mM NaCl, 2.5 mM KCl, 1.2 mM NaH_2PO_4, 2.0 mM $CaCl_2$, 2.0 mM $MgSO_4$, 26 mM

$NaHCO_3$ and 10 mM glucose. After sectioning, hippocampal slices were allowed to

recover in the chamber for 30 min at 34°C and then for a further 2-8 h at room

temperature (25 ± 1°C). All solutions were saturated with 95% O_2/5% CO_2 (vol/vol).

The slices were placed in the recording chamber, which was superfused (3 mL/min)

with ACSF at 32-34°C. fEPSPs were evoked in the CA1 stratum radiatum by

stimulating the SC with a bipolar stimulating electrode (FHC) and recorded in current

clamp using a HEKA EPC 10 amplifier (HEKA Elektronik) with ACSF-filled glass

pipettes (1-3 MΩ). Test stimuli consisted of monophasic 100-μs pulses of constant

current (intensity adjusted to produce 25% of the maximum response) at a frequency of

0.033 Hz. The strength of synaptic transmission was determined by measuring the

initial amplitude of fEPSPs. L-LTP was induced by 4 trains of 100 Hz stimuli of 50

pulses each at 100 Hz, 10 s apart, at the same intensity as the test stimulus. The level of

E-LTP / L-LTP was determined at an average of 55-60 / 115-120 min after HFS

stimulation. To study glutamatergic transmission at SC-CA1 synapses, we recorded

fEPSP input/output (I/O) curves at SC-CA1 synapses. To construct an I/O curve,

fEPSPs were evoked by a series of incremental intensities (0.02-0.2 mA in 0.02 mA

increments, 100 μs duration, at 0.033 Hz). Whole-cell patch-clamp recordings from

CA1 neurons were visualized with infrared optics using an upright microscope

equipped with a 40× water-immersion objective (BX51WI; Olympus) and an infrared-

sensitive CCD camera. To analyze the intrinsic excitability of CA1 pyramidal neurons,

the current clamp was used and neurons were held at 0 pA and then injected with step

currents (duration: 500 ms; increments: 20 pA; range: -100 to 400 pA; interval: 10 s).

The input-output relationship was determined by plotting the injected currents against

the number of action potentials. All data were acquired at a sampling rate of 10 kHz

using PATCHMASTER version 2 x 90.1 software (HEKA Elektronik) and filtered off-

line at 1 kHz.

Generation and use of the ac4C website

The ac4Catlas website was constructed using HTML (a scripting language for front-

end development), PHP (a scripting language for backend development), and MySQL

(a structured query language for data operation). All the website codes were deployed

in a Linux server equipped with Centos environment. The raw data, processed data and

annotated files were stored in "Data Downloads" section. JBrowse (https://jbrowse.org)

was integrated to visualize the ac4C epitranscriptomic data on genomic regions of

interest. Briefly, ac4Catlas provides user-friendly interactive interfaces which allow

researchers to freely browse and perform extended operations, such as searching,

comparing, and downloading. Besides, the browse module stores detailed information

of samples and corresponding datasets as well as ac4C levels of mRNA and

differentially acetylated regions between groups.

Statistical analysis

The cumulative distribution curve was analyzed by Kolmogorov-Smirnov test in R. The

statistical analysis of percentage of ac4C summits occurring within CDS or UTRs was

performed by chi-square test in R. The regular quantification data were shown as mean

$\pm$ SEM unless otherwise mentioned. The sample size justification was based on the

previous studies and met the requirement of statistics power. For normally distributed

data with 1 experimental variable, statistical analyses were performed by parametric

analysis: unpaired (two-tailed) student's t-test for two groups and one-way ANOVA

with the Tukey's multiple-comparison test for 3 groups. Data with >1 experimental

variable were evaluated by two-way ANOVA, with post hoc Tukey, Dunnett or Sidak's multiple-comparison tests. All these statistical analyses were performed by the software of GraphPad Prism 10. Statistically significant difference was indicated as follows: **** $P < 0.0001$, *** $P < 0.001$, ** $P < 0.01$ and * $P < 0.05$.

Acknowledgements

We thank CloudSeq Inc. (Shanghai, China) for providing us with technical support on acRIP-seq. This work was supported by STI2030-Major Projects (Grant No. 2021ZD0202500) and the National Natural Science Foundation of China (Grant No. 32300560), and the Fundamental Research Funds for the Central Universities (Grant No. YBNLTS2022-011).

Author contributions

H.-Q.Z. and H.J. performed biochemical and behavioral analysis, H.-Q.Z., J.-W.Z., and Z.Z. analyzed the data, W.-P.L. and Y.-Y.D. performed electrophysiology. S.C. generated the ac4C website, D.-M.Y. conceived the idea, supervised the work and wrote the paper. All authors approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Data and materials availability

All data are available in the main text, supplementary or source data files. The original data of acRIP-seq is accessible via the website (http://ac4Catlas.com/) or GEO database (GSE261922). The source data files are provided with the paper, which contain high-resolution figures, uncropped western blot images, and original statistical data. All unique materials used are readily available from the corresponding author upon reasonable requests.

 Code availability

 Custom code used in this paper is available at https://github.com/YinDMlab/ac4CRIP-

 seq-analysis.

 Figures and Legends

 Figure 1 The presence of ac4C modification in PolyA RNA of mouse hippocampus.

 A-B, Verification of the purity of PolyA RNA through the bioanalyzer profiling. The x-

 axis represents the size of nucleotides, and the y-axis represents the fluorescent signal

 from the bioanalyzer. A, total RNA. B, PolyA RNA. **C**, Diminished levels of 18S rRNA

 in the polyA RNA samples, compared with total RNA samples, which was revealed by

 RT-qPCR analysis. The levels of 18S rRNA were normalized by that of Gapdh. ***** P**

 < 0.0001, two-tailed t-test, n = 6 biological replicates. **D**, Standard ac4C curve for LC-

 MS/MS analysis. The standard samples consisting of varying concentrations of ac4C

 nucleotide (1 pmol, 2 pmol, 5 pmol, 10 pmol, 20 pmol, and 50 pmol) were analyzed by

 MS/MS to generate a calibration curve. The x-axis represents the concentration of the

 ac4C standard samples, and the y-axis represents the signal intensity from the MS/MS

 analysis. The linear regression analysis yielded a correlation coefficient (r) of 0.998,

 indicating a strong linear relationship between ac4C concentration and LC-MS/MS

 signal. **E-F**, The chromatograms of blank samples and ac4C (E) and m6A (F) standards

 in the LC-MS/MS analysis. **G**, The representative chromatograms of ac4C and m6A

 peaks identified in the polyA RNA purified from the HOM of mouse hippocampus. **H**,

 Reduction of ac4C levels by the chemical deacetylation protocol. The stoichiometry

levels of ac4C in polyA RNA from control and NH$_2$OH-treated samples were quantified. ** P = 0.008, two-tailed t-test, n = 3 biological replicates. **I**, Unchanged m6A levels by the chemical deacetylation protocol. The stoichiometry levels of m6A in polyA RNA from control and NH$_2$OH-treated samples were quantified. NS, not significant, two-tailed t-test, n = 3 biological replicates. Quantification data are expressed as mean ± SEM.

Figure 2 Increased mRNA acetylation in the SYN of mouse hippocampus after learning and memory. A, The purified synaptosomes (SYN) of mouse hippocampus expressed high levels of PSD95 and synapsin 1 (SYN1), two synaptic protein markers, but very low levels of α-tubulin, a cytoplasmic protein marker. The homogenates (HOM), cytoplasm (S2) and SYN of mouse hippocampus were subjected to western blots with the indicated antibodies. Left panel, representative images of western blots; right panel, Ponceau staining to show equal protein loading. **B,** Acetylation of polyA RNA was not altered in SYN of mouse hippocampus at day 1. The polyA RNA purified from the SYN of control and learning mice at day 1 after MWM training were subjected to ac4C dot-blots and methylene blue staining. Left, the representative images; right, quantification data. NS, not significant, two-tailed t-test, n = 3 biological replicates. Data were normalized by controls. **C,** Acetylation of polyA RNA was increased in the

SYN of mouse hippocampus at day 6. The polyA RNA purified from the SYN of control and learning mice at day 6 after MWM training were subjected to ac4C dot-blots and methylene blue staining. Left, the representative images; right, quantification data. ** P = 0.0093, two-tailed t-test, n = 4 biological replicates. Data were normalized by controls. **D**, Acetylation of polyA RNA was not altered in the SYN of mouse hippocampus at day 20. The polyA RNA purified from the SYN of control and learning mice at day 20 after MWM training were subjected to ac4C dot-blots and methylene blue staining. Left, the representative images; right, quantification data. NS, not significant, two-tailed t-test, n = 4 biological replicates. Data were normalized by controls. **E,** Acetylation of mRNA in the SYN was increased during memory formation but returned to normal levels after forgetting. The stoichiometry levels of ac4C of polyA RNA purified from the learning (L) mice were shown at days 1, 6 and 20 after MWM training. ** P = 0.0023, one-way-ANOVA, n = 3 biological replicates. **F-H**, Acetylation of mRNA was increased after memory formation but returned to normal levels after forgetting in the SYN of mouse hippocampus. Shown are cumulative distribution curves of FE of ac4C mRNA purified from the SYN of control (C) and learning (L) mice at days 1 (**F**), 6 (**G**), and 20 (**H**) after MWM training. The FE of ac4C mRNA was quantified in the quartile boxplots, where the solid line in the box is the median and the dashed line is the mean, where the maximum and minimum values are identified. NS, not significant, *** adj-P = 9.9814e-13, Kolmogorov-Smirnov test. **I-K,** The total mRNA levels were not significantly altered in the SYN of mouse hippocampus during learning and memory. Shown are cumulative distribution curves of normalized mRNA counts for SYN in control (C) and learning (L) mice at days 1 (I), 6 (J) and 20 (K) after MWM training. The normalized mRNA counts were quantified in the quartile boxplots, where the solid line in the box is the median and the dashed line is the mean, where the maximum and minimum values are identified. NS, not significant, Kolmogorov-Smirnov test. C, control; L, learning. FE, Fold Enrichment. Quantification data are expressed as mean ± SEM.

1403

Figure 3 A representative LISA mRNA for Arc. A, The biological processes (BP) in

which LISA mRNAs are significantly overrepresented. The x-axis represents the

negative logarithm of the adjusted p-value, reflecting the significance of the enrichment,

while the y-axis shows the biological processes. **B,** The protein-protein interaction (PPI)

network of LISA mRNA implicated in the BP of learning or memory. P < 1.0e-16,

Hypergeometric Test**. C,** Increased ac4C modification of Arc in the SYN rather than in

the HOM at day 6. Also note the unaltered ac4C Arc mRNA at days 1 or 20 in the SYN.

Shown are the IGV maps of the ac4C peaks from the CDS region of Arc mRNA. The

ac4C peaks were represented by acRIP/input (See details in the Method). **D**, **G**, The

Arc mRNA levels from the input of SYN were not altered at days 1 (D) or 20 (G) after

MWM training. The RNA purified from the SYN was subjected to the RT-qPCR

analysis. NS, not significant, two-tailed t-test, n = 4 biological replicates. The levels of

Arc mRNA were normalized by that of Gapdh in each sample, and then the ratio of Arc

to Gapdh were normalized by controls. **E, H**, The ac4C modification of Arc mRNA was

not altered in the SYN at days 1 (E) or 20 (H) after MWM training. The RNA purified

through the acRIP and IgG-IP products of SYN was subjected to the RT-qPCR analysis.

NS, not significant, two-way-ANOVA followed by Tukey's multiple comparison test, n

= 4 biological replicates. Data were normalized by input. **F, I**, The protein levels of

ARC did not change in the SYN at days 1 (F) or 20 (I) after MWM training. Top,

representative western blot images; bottom, quantification data. NS, not significant, two-tailed t-test, n = 4 biological replicates. The protein levels of ARC were normalized by that of SYN1 in each sample, and then the ratio of ARC to SYN1 were normalized by controls. **J**, The Arc mRNA levels from the input of SYN were not altered at day 6 after MWM training. The RNA purified from the SYN was subjected to the RT-qPCR analysis. NS, not significant, two-tailed t-test, n = 4 biological replicates. The levels of Arc mRNA were normalized by that of Gapdh in each sample, and then the ratio of Arc to Gapdh were normalized by controls. **K**, The ac4C modification of Arc mRNA was significantly increased in the SYN at day 6 after MWM training. The RNA purified from the acRIP and IgG-IP products of SYN was subjected to the RT-qPCR analysis. *** $P < 0.001$, two-way-ANOVA followed by Tukey's multiple comparison test, n = 4 biological replicates. Data were normalized by input. **L**, The protein levels of ARC were significantly increased in the SYN at day 6 after MWM training. Top, representative western blot images; bottom, quantification data. ** $P = 0.0032$, two-tailed t-test, n = 4 biological replicates. The protein levels of ARC were normalized by that of SYN1 in each sample, and then the ratio of ARC to SYN1 were normalized by controls. **M**, The Arc mRNA levels from the HOM were not altered at day 6 after MWM training. The Arc mRNA levels were revealed by the $\log_2$TPM from the next-generation sequencing of the HOM RNA. NS, not significant, two-tailed t-test, n = 3 biological replicates. **N**, The protein levels of ARC did not change in the HOM at day 6 after MWM training. Top, representative western blot images; bottom, quantification data. NS, not significant, two-tailed t-test, n = 4 biological replicates. The protein levels of ARC were normalized by that of GAPDH in each sample, and then the ratio of ARC to GAPDH were normalized by controls. C, control; L, learning. Quantification data are expressed as mean ± SEM.

Figure 4 Increased NAT10 proteins in the SYN of mouse hippocampus after learning and memory. A, Experimental design. The SYN fractions were purified from the hippocampus of control and learning mice at days 1, 6 and 20. The NAT10 protein levels in the SYN were then assessed by western blots. **B-D**, Protein levels of NAT10 were increased in the SYN after memory formation but returned to normal levels after forgetting. The SYN proteins from control and learning mice at days 1 (B), 6 (C), and 20 (D) were subjected to western blots with the indicated antibodies. The protein levels of NAT10 were normalized by that of SYN1 in each sample, and then the ratio of NAT10 to SYN1 was normalized by controls. Top, representative western blot images; bottom, quantification data. NS, not significant, ** $P = 0.0023$, two-tailed t-test, n = 4 biological replicates. **E**, The fold change (FC) of NAT10 proteins in panel B to D was quantified. ** $P = 0.0043$, one-way-ANOVA, n = 4 biological replicates. **F**, Protein levels of NAT10 were not altered in the HOM at day 6. The HOM proteins from control and learning mice at day 6 were subjected to western blots with the indicated antibodies. Top, representative western blot images; bottom, quantification data. NS, not significant, two-tailed t-test. Data were normalized by controls, n = 6 biological replicates at each

condition. The protein levels of NAT10 were normalized by that of α-tubulin in each sample, and then the ratio of NAT10 to α-tubulin was normalized by controls. **G**, Experimental design. Proteins were extracted from nucleus (N), cytoplasm (S2), and postsynaptic density (PSD) of control and learning mouse hippocampus at day 6, and then NAT10 protein levels were assessed by western blots. **H**, Expression of different protein markers in the nuclear (N), cytoplasmic (S2), P2, SYN, and PSD fractions of mouse hippocampus. Shown are representative images of western blots. Protein samples purified from different subcellular fractions of mouse hippocampus were subjected to western blots with the indicated antibodies. **I**, Protein levels of NAT10 were not altered in the nuclear fraction of mouse hippocampus after memory formation. The nuclear protein samples from control and learning mice at day 6 were subjected to western blots with the indicated antibodies. Top, representative western blot images; bottom, quantification data. NS, not significant, two-tailed t-test, n = 4 biological replicates. The protein levels of NAT10 were normalized by that of Histone H3 proteins in each sample, and then the ratio of NAT10 to Histone H3 was normalized by controls. **J**, Protein levels of NAT10 were significantly reduced in the S2 fraction of mouse hippocampus after memory formation. The cytoplasmic protein samples from control and learning mice at day 6 were subjected to western blots with the indicated antibodies. Top, representative western blot images; bottom, quantification data. ** $P = 0.0052$, two-tailed t-test, n = 6 biological replicates. The protein levels of NAT10 were normalized by that of α-tubulin in each sample, and then the ratio of NAT10 to α-tubulin was normalized by controls. **K**, Protein levels of NAT10 were significantly increased in the PSD fraction of mouse hippocampus after memory formation. The PSD protein samples from control and learning mice at day 6 were subjected to western blots with the indicated antibodies. Top, representative western blot images; bottom, quantification data. *** $P = 0.0008$, two-tailed t-test. n = 6 biological replicates. The protein levels of NAT10 were normalized by that of PSD95 in each sample, and then the ratio of NAT10 to PSD95 was normalized by controls. **L-N**, Protein levels of PCAF were not altered in the nuclear (L), S2 (M), or PSD fraction (N) of mouse hippocampus

after memory formation. The protein samples purified from the different subcellular fractions of the hippocampus of control and learning mice at day 6 were subjected to western blots with the indicated antibodies. Top, representative western blot images; bottom, quantification data. NS, not significant, two-tailed t-test, n = 6 biological replicates. The protein levels were normalized by the protein markers in different subcellular fractions and then were normalized by controls. C, control; L, learning. Quantification data are expressed as mean ± SEM.

Figure 5 NAT10-dependent of ac4C mapping in the SYN of learning mice. A, Experimental design. Three weeks after AAV injection, the control (Ctrl) and Nat10 conditional knockout (cKO) mice were subjected to MWM training and probe test, and then the SYN fractions were purified from hippocampal tissues for low-input acRIP-seq. **B**, Reduction of ac4C peak number in the SYN of Nat10 cKO mice, compared with controls. 7097 and 1933 ac4C peaks were identified in the SYN of control and Nat10 cKO mice, respectively. **C**, Definition of NAT10-dependent ac4C peaks through acRIP summits displaying higher FE values in control relative to Nat10 cKO mice. Shown are the density scatter plot of ac4C peaks. Each dot represents an ac4C peak. 6631 ac4C

peaks are specifically identified in the SYN of control but not Nat10 cKO mice. The FE of 408 ac4C peaks was higher in the SYN of control mice than Nat10 cKO mice. **D**, NAT10-dependent ac4C peaks are predominantly located in the CDS region of mRNA. The plot shows the density of ac4C peaks along various mRNA regions, including the 5'UTR, CDS, and 3'UTR. **E**, Reduction of ac4C mRNA number in the SYN of Nat10 cKO mice, compared with controls. 3778 and 1680 ac4C mRNAs were identified in the SYN of control and Nat10 cKO groups, respectively. **F**, Similar mRNA numbers identified in the SYN of control and Nat10 cKO mice. 6041 and 6047 mRNAs were identified in the SYN of control and Nat10 cKO mice, respectively. **G**, Acetylation of mRNA in the SYN of Nat10 cKO mice was significantly reduced, compared with controls. Shown are cumulative distribution curves of FE of ac4C mRNA purified from the SYN of control and Nat10 cKO mice at day 6 after MWM training. The FE of ac4C mRNA was quantified in the quartile boxplots, where the solid line in the box is the median and the dashed line is the mean, where the maximum and minimum values are identified. *** adj-P = $1.20076e-12$, Kolmogorov-Smirnov test. **H**, The total mRNA levels were not significantly altered in the SYN of Nat10 cKO mice, compared to controls. Shown are cumulative distribution curves of normalized mRNA counts for SYN in control and Nat10 cKO mice at day 6 after MWM training. The normalized mRNA counts were quantified in the quartile boxplots, where the solid line in the box is the median and the dashed line is the mean, where the maximum and minimum values are identified. NS, not significant, Kolmogorov-Smirnov test. **I**, Definition of NAT10-dependent synaptic ac4C (NASA) mRNA through acRIP summits displaying higher FE values in control relative to Nat10 cKO mice. Shown are the density scatter plot of ac4C mRNA. Each dot represents an ac4C mRNA. 2507 ac4C mRNAs are specifically identified in the SYN of control but not Nat10 cKO mice. The FE of 839 ac4C mRNA was higher in the SYN of control mice than Nat10 cKO mice. **J**, The expression of NASA mRNA is mostly enriched in excitatory neurons of mouse hippocampus. Shown are the percentage of NASA mRNA whose expression was relatively enriched in different cell types of mouse hippocampus through analysis of the single-cell RNA-

 sequencing dataset (http://dropviz.org). **K**, The biological processes (BP) in which

1542 NASA mRNAs are significantly overrepresented. The y-axis represents the negative

1543 logarithm of the adjusted p-value, reflecting the significance of the enrichment.

1544 Quantification data are expressed as mean ± SEM.

1545

1546

Figure 6 Reduction of ac4C modification and protein levels of Arc, Camk2α and

Grin2b in the SYN of Nat10 cKO mice, compared with controls. A, Classification

of NASA mRNAs overrepresented in the BP related to synaptic plasticity and learning

or memory (Figure 5K). 27 NASA mRNAs overlap with LISA mRNAs, which were

named class 1 mRNA. In contrast, 212 NASA mRNAs do not overlap with LISA

mRNAs, which were named class 2 mRNA. **B**, **C**, The protein-protein interaction (PPI)

network of class 1 (B) and 2 (C) mRNA. Class 1: P < 3.9e-11; Class 2: P < 1.0e-16,

Hypergeometric test. **D-F**, Reduced ac4C modification of Arc (D), Camk2α (E), and

Grin2b (F) mRNA in the SYN of Nat10 cKO mice at day 6, compared with controls.

Shown are the IGV maps of the ac4C peaks from the 3' UTR to the 5' UTR of transcripts.

The ac4C peaks were represented by acRIP/input (See details in the Method). **G**,

Experimental design. Three weeks after AAV injection, the control (Ctrl) and Nat10

conditional knockout (cKO) mice were subjected to MWM training and probe test, and

then the SYN fractions were purified from hippocampal tissues for RT-qPCR and western blot analysis. **H**, The mRNA levels of Arc, Camk2α and Grin2b from the input of SYN were similar between control and Nat10 cKO mice. The RNA purified from the SYN was subjected to the RT-qPCR analysis. NS, not significant, two-tailed t-test, n = 4 biological replicates. The levels of mRNA were normalized by that of Gapdh in each sample, and then the ratio of mRNA to Gapdh were normalized by controls. **I**, The ac4C modification of Arc, Camk2α, and Grin2b mRNA was significantly reduced in the SYN of Nat10 cKO mice at day 6, compared with controls. The RNA purified from the acRIP and IgG-IP products of SYN was subjected to the RT-qPCR analysis. *** P < 0.001, two-way-ANOVA followed by Tukey's multiple comparison test, n = 4 biological replicates. Data were normalized by input. **J**, The protein levels of ARC, CaMK2α and GluN2B were significantly increased in the SYN of Nat10 cKO mice at day 6, compared with controls. Left, representative western blot images; right, quantification data. *** P < 0.001, two-tailed t-test, n = 4 biological replicates. The protein levels were normalized by that of PSD95 in each sample, and then the ratio of protein to PSD95 were normalized by controls. Quantification data are expressed as mean ± SEM.

Figure 7 Impaired LTP as well as learning and memory in the Nat10 cKO mice. A, A representative image showing recording LTP at the SC-CA1 synapses from Nat10

cKO mice using the protocol of high-frequency stimulation (HFS). Scale bar, 500 μm. **B**, The representative fEPSP traces from baseline (BSL), post-tetanus potentiation (PTP), early phase of LTP (E-LTP, 55-60 min after HFS) and late phase of LTP (L-LTP, 115-120 min after HFS). **C**, Impaired maintenance of L-LTP at the SC-CA1 synapses in Nat10 cKO mice, compared with controls. Normalized fEPSP amplitudes were plotted every 1 min for hippocampal slices from control and Nat10 cKO mice. *** $P < 0.001$, two-way ANOVA, n = 8 slices from 4 mice for each group. **D**, Normal LTP induction at the SC-CA1 synapses in Nat10 cKO mice, compared to controls. The PTP in panel C was quantified. NS, not significant, two-tailed t-test. n = 8 slices from 4 mice for each group. **E**, Unaltered E-LTP at the SC-CA1 synapses in Nat10 cKO mice, compared to controls. The E-LTP in panel C was quantified. NS, not significant, two-tailed t-test. n = 8 slices from 4 mice for each group. **F**, Impaired L-LTP at the SC-CA1 synapses in Nat10 cKO mice, compared with controls. The L-LTP in panel C was quantified. ** $P=0.008$, two-tailed t-test. n = 8 slices from 4 mice for each group. **G**, Experimental design. The control and Nat10 cKO mice were subjected to fEPSP recording at the SC-CA1 synapses after training and probe test in the MWM. **H-I**, Learning and memory-induced enhancement of glutamatergic transmission at SC-CA1 synapses was significantly impaired in Nat10 cKO mice, compared with controls. H, Representative fEPSP traces from control and Nat10 cKO mice. I, Quantification of the fEPSP amplitudes in panel H. *** Genotype $F_{(3, 680)} = 128.2$, $P < 0.0001$, two-way ANOVA followed by Sidak's multiple comparison test, n = 18 slices from 5 mice each group. **J**, Impaired learning process in Nat10 cKO mice, compared with controls. Shown are the learning curves of control and Nat10 cKO mice in the MWM. n = 14 mice per group, ** Genotype $F_{(1, 130)} = 24.08$, ** $P = 0.0014$, two-way ANOVA followed by Sidak's multiple comparison test. **K**, Similar distance travelled during probe tests between control and Nat10 cKO mice. NS, not significant, two-tailed t-test, n = 14 mice per group. **L**, Representative swimming traces of control and Nat10 cKO mice during probe tests. **M**, Time spent in the target quadrant was significant reduced in Nat10 cKO groups during probe tests, compared with controls. * $P = 0.0198$, two-

way ANOVA followed by Sidak's multiple comparison with post hoc t-test, n = 14 mice per group. **N**, Number of platform crossing was significantly reduced in Nat10 cKO mice during probe tests, compared with controls. *** P < 0.0001, two-tailed t-test, n = 14 mice per group. Quantification data are expressed as mean ± SEM.

Supplementary Figures and Legends

Figure S1 Protocols of Morris water maze (MWM) for control and learning mice.
A, MWM protocol for control mice. The control mice were placed in MWM with a visible platform at days 1 and 2, after which they returned to home cage and were subjected to probe tests at day 6. **B**, Control mice were able to discern the visible platform above the water surface and consequently locate it rapidly. Shown are the latency to find the visible platform at days 1 and 2. **C**, Control mice were not able to remember the target quadrant during probe tests at day 6. Shown are the percentage of time staying in each quadrant during probe tests. NS, not significant, one-way ANOVA, n = 9 mice. **D**, MWM protocol for learning mice. The learning mice were trained in MWM with a hidden platform from day 1 to 5, and then were subjected to probe tests at day 6 and 20. **E**, Learning mice showed improvement in their ability to find the hidden platform during the training process. Shown are the learning curves. **F**, Learning mice could remember the target quadrant during probe tests at day 6. Shown are the percentage of time staying in each quadrant during probe tests. *** P < 0.001, one-way ANOVA, n = 9 mice. **G**, Learning mice could not remember the target quadrant during probe tests at day 20. Shown are the percentage of time staying in each quadrant during probe tests. NS, not significant, one-way ANOVA, n = 6 mice. **H**, Summary of learning

score of control and learning mice at days 1, 6, and 20. Memories were formed at day

6 but were forgotten at day 20.

1636

Figure S2 No significant change of total levels of ac4C mRNA in the homogenates (HOM) of mouse hippocampus after learning and memory. A, Experimental design of acRIP-seq. The RNA samples from both input and acRIP were subjected to next-generation sequencing. Among them, input was used to eliminate the background of ac4C peaks. The process of ac4C peak calling identifies reliable ac4C peaks and calculates the FE of each ac4C peak. **B**, **C**, The ac4C peaks and ac4C mRNAs are largely overlapping across the three biological replicates of control HOM samples. Shown are the number of overlapping and non-overlapping ac4C peaks (B) and ac4C mRNAs (C) across the three biological replicates of control HOM samples. **D**, The same consensus motif, AGCAGCTG, was identified in ac4C peak regions across the three biological replicates of control HOM samples. Consensus motif and its P value were generated by HOMER (see details in the Method). 99% confidence intervals: Rep 1 target 57.20%; Rep 2 target 54.23%; Rep 3 target 55.32%. **E**, **F**, Reliable ac4C peaks were those identified in at least two of the three biological replicates. Only mRNAs corresponding to these reliable ac4C peaks were defined as reliable ac4C mRNAs.

Shown are the Venn diagram of ac4C peaks for the three biological replicates of control (E) and learning (F) HOM samples. **G**, A large proportion of ac4C peaks in the HOM of mouse hippocampus are located within the CDS of mRNA. Shown are the relative density of ac4C peaks identified in different regions of transcripts for the three biological replicates of control HOM samples. CDS, coding sequence; UTRs, untranslated regions. **H**, The number of ac4C peaks for each ac4C mRNA is comparable across the three biological replicates in both control and learning HOM samples. **I**, Acetylation of polyA RNA was not altered in the HOM of mouse hippocampus after memory formation. The polyA RNA samples used for each ac4C dot-blot were 0.5 μg that falls into the linear range. The HOM polyA RNA samples of control and learning mice at day 6 after MWM training were subjected to ac4C dot-blots and methylene blue staining. Left, representative images; right, quantification data. NS, not significant, two-tailed t-test, n = 3 biological replicates at each condition. Data were normalized by controls. **J**, Acetylation of ac4C mRNA did not significantly change in the HOM of mouse hippocampus after memory formation. Shown are cumulative distribution curves of FE of ac4C mRNA in control and learning mice at day 6 after MWM training. The FE of ac4C mRNA was quantified in the quartile boxplots, where the solid line in the box is the median and the dashed line is the mean, where the maximum and minimum values are identified. NS, not significant, Kolmogorov-Smirnov test. **K,** The total mRNA levels did not significantly change in the HOM of mouse hippocampus after memory formation. Shown are cumulative distribution curves of normalized mRNA counts in the HOM of control and learning mice at day 6 after MWM training. The normalized mRNA counts of each group were quantified in the quartile boxplots, where the solid line in the box is the median and the dashed line is the mean, where the maximum and minimum values are identified. NS, not significant, Kolmogorov-Smirnov test. FE, Fold Enrichment. C, control; L, learning.

Figure S3 Identification of synaptosome (SYN) ac4C epitranscriptome and learning-induced synaptic ac4C (LISA) mRNAs in mouse hippocampus. A, **B**, The mRNA and ac4C mRNAs are largely overlapping across the SYN control group at days 1, 6, and 20. Shown are the number of overlapping and non-overlapping mRNAs (A) and ac4C mRNAs (B) across the control SYN samples at days 1, 6 and 20 after MWM training. **C**, The same consensus motif, AGCAGCTG, was identified in ac4C peak regions across the control SYN samples at days 1, 6 and 20. Consensus motif and its P value were generated by HOMER (see details in the Method). 99% confidence intervals: Day 1 target 55.49%; Day 6 target 54.65%; Day 20 target 54.91%. **D**, **E**, A large proportion of ac4C peaks in the SYN of mouse hippocampus are located within the CDS of mRNA. Shown are the relative density of ac4C peaks identified in different regions of transcripts for control (D) and learning (E) SYN samples at days 1, 6 and 20. CDS, coding sequence; UTRs, untranslated regions. **F-H**, Scatter plot showing the fold change (FC) of mRNA and ac4C mRNA in the SYN of mouse hippocampus at days 1, 6, and 20 after MWM training. The x-axis represents the $\log_2$ FC of ac4C mRNA, while the y-axis shows the $\log_2$ FC of mRNA. The red color represents upregulated ac4C mRNAs, the blue color indicates downregulated ac4C mRNAs, and the black color represents unchanged ac4C mRNAs. The cut off values for upregulation and downregulation are FC >2 and FC < 0.5, respectively. **I**, The ac4C rather than total levels of LISA mRNA are upregulated following memory formation and return to control levels after memory forgetting. The FC of total and ac4C levels for LISA mRNA

 were quantified at days 1, 6 and 20 after MWM training.

1704 **Figure S4 Expression of Nat10 in the mature neurons of mouse hippocampus. A**,

1705 Experimental design. Cre-reporting AAV was injected into the dorsal hippocampus of

1706 7-week-old Nat10$^{Cre/+}$ mice. Three weeks after AAV injection, the fluorescent protein

1707 EGFP was analyzed in the Nat10 reporter mice. **B**, Co-localization of EGFP with NeuN

1708 in the hippocampus of Nat10 reporter mice. The hippocampal slices from Nat10

1709 reporter mice were immunostained with anti-NeuN antibodies. so, stratum oriens; sp,

1710 stratum pyramidale; sr, stratum radiatum; slm, stratum lacunosum-moleculare; sm,

1711 stratum moleculare; gcl, granule cell layer; DG: dentate gyrus. Scale bar, 100 μm. **C**,

1712 Co-localization of EGFP and NeuN (top images) or Neurogranin (NG) (bottom images)

1713 in the CA1 region of Nat10 reporter mice. The hippocampal slices from Nat10 reporter

1714 mice were immunostained with anti-NeuN or anti-NG antibodies. The yellow arrows

1715 indicate cells coexpressing Nat10 and NeuN or NG. The red arrows represent cells

1716 expressing NeuN or NG but not Nat10. The green arrows indicate cells expressing

1717 Nat10 but not NeuN or NG. Scale bar, 25 μm. **D**, Quantification of the percentage of

1718 NeuN-positive cells expressing Nat10 (Nat10^{+} / NeuN^{+} = 65.61 ± 3.245%) and the

1719 percentage of Nat10-positive cells expressing NeuN (NeuN^{+} / Nat10^{+} = 81.31 ±

1720 2.029%). n = 9 slices from 3 mice. **E**, Quantification of the percentage of NG-positive

1721 cells expressing Nat10 (Nat10^{+} / NG^{+} = 81.87 ± 2.765%) and the percentage of Nat10-

positive cells expressing NG (NG$^+$ / Nat10$^+$ = 75.82 ± 1.882%). n = 9 slices from 3 mice.

F, Experimental design. Cre-reporting AAV was injected into the dorsal hippocampus

of 7-week-old ALDH1L1$^{Cre/ERT2}$ mice. Three weeks after AAV injection, the fluorescent

protein EGFP was analyzed in the astrocyte reporter mice. **G**, Astrocytes in the CA1

region of astrocyte reporter mice do not co-localize with NeuN (white arrow). The

hippocampal slices from astrocyte reporter mice were immunostained with anti-NeuN

antibodies. Scale bar, 50 μm.

1729

1730

Figure S5 Increased colocalization of NAT10 and PSD95 in cultured mouse

hippocampal neurons after cLTP stimulation. A, Experimental design. The primary

neurons were taken from the E16 mouse hippocampus, and the immunostaining was

performed at DIV 14. **B**, Fluorescent images of DAPI, PSD95, NAT10 and MAP2 in

cultured hippocampal neurons treated with vehicle. The bottom images were enlarged

from the rectangle in the top images. Scale bar, 25 μm (top) and 100 μm (bottom). C,

Fluorescent images of DAPI, PSD95, NAT10 and MAP2 in cultured hippocampal

neurons 60 min after cLTP stimulation. The bottom images were enlarged from the

rectangle in the top images. Scale bar, 25 μm (top) and 100 μm (bottom). **D**, Increased

colocalization of NAT10 and PSD95 in the dendrites of cultured hippocampal neurons.

The number of PSD95 puncta colocalized with NAT10 was quantified. ** P = 0.0014,

two-tailed t-test, n = 30 neurons from 3 independent experiments. DIV, days in vitro.

DAPI, a nuclear marker; PSD95, a protein marker for postsynaptic density; MAP2

(microtubule-associated protein 2), a protein marker for soma and dendrites.

1745

1746

Figure S6 NAT10-dependent ac4C modification of polyA RNA in mouse hippocampus. A, Experimental design. Three weeks after AAV injection, the homogenates of hippocampal tissues from control (Ctrl) and Nat10 cKO mice were collected for western blots, ac4C and m6A dot-blots, and LC-MS/MS analysis. **B**, Infection of AAV-Syn1-EGFP-P2A-Cre in the hippocampus of 9-week-old mice. Shown are fluorescent images of the mouse hippocampus expressing EGFP. Scale bar, 500 μm. **C-F**, Downregulation of NAT10, rather than acetylated tubulin or histone H3 proteins in the hippocampus of Nat10 cKO mice, compared with controls. The protein samples from control and Nat10 cKO mouse hippocampus were probed with the indicated antibodies. C, representative western blot images, D-F, Quantification of NAT10/tubulin, acetylated-tubulin/tubulin and acetylated-H3/H3 in panel C. NS, not significant, *** P < 0.0001, two-tailed t-test, n = 6 biological replicates. Data were normalized by controls. **G-H**, Reduced ac4C modification of polyA RNA in the hippocampus of Nat10 cKO mice, compared with controls. The polyA RNA samples purified from mouse hippocampus were subjected to ac4C dot-blots and methylene blue staining. G, Representative images. H, Quantification data. *** P = 0.0006, two-tailed t-test, n = 6 biological replicates. Data were normalized by controls. **I-J**, Similar m6A modification of polyA RNA in the hippocampus of control and Nat10 cKO mice. I, Representative images. J, Quantification data. NS, not significant, two-tailed t-test, n =

4 biological replicates. **K**, Reduced ac4C modification of polyA RNA in the

hippocampus of Nat10 cKO mice, compared with controls. The stoichiometry levels of

ac4C from LC-MS/MS analysis were quantified. ** P = 0.0077, two-tailed t-test, n = 3

biological replicates. **L**, Similar m6A modification of polyA RNA in the hippocampus

of control and Nat10 cKO mice. The stoichiometry levels of m6A from LC-MS/MS

analysis were quantified. NS, not significant, two-tailed t-test, n = 3 biological

replicates.

1773

1774

Figure S7 Downregulation of NAT10 in mouse hippocampus does not affect body

or brain weight, neuronal number or intrinsic excitability. A, Similar body weight

between 10-week-old control and Nat10 cKO mice. Left, representative images of the

mice, right, quantification of body weight. NS, not significant, two-tailed t-test, n = 12

mice for each group. **B**, Similar brain weight between 10-week-old control and Nat10

cKO mice. Left, representative images of mouse brain, right, quantification of brain

weight. NS, not significant, two-tailed t-test, n = 12 mice for each group. **C**, Similar

number of neurons expressing EGFP in the CA1 region of 10-week-old control and

Nat10 cKO mice. Left, representative images of CA1 neurons expressing EGFP, right,

quantification of EGFP-expressing cells in the CA1 region. NS, not significant, two-

tailed t-test, n = 8 slices from 4 mice for each group. Data were normalized by control.

D, Similar intrinsic excitability of CA1 pyramidal neurons between 10-week-old

control and Nat10 cKO mice. Left, representative traces of action potentials (AP) of CA1 pyramidal neurons; right, quantification of AP number after injection of different currents into CA1 pyramidal neurons. NS, not significant, two-way ANOVA followed by Sidak's multiple comparison test, n = 20 cells from 4 mice for each group.

Supplementary Tables and Legends

Table S1. Library Information of acRIP-seq.

Table S2. List of mRNAs identified by regular acRIP-seq (HOM).

Table S3. List of mRNAs identified by low-input acRIP-seq (SYN).

Table S4. The information of LISA mRNA.

Table S5. The information of NASA mRNA.