Chromosome-level reference genome assembly for the protected

resource plant, *Zenia insignis*

Authors
Si-Yun Chen[1,2,4], Zhi-Yun Yang[1,4], Rong Zhang[1,2,3], Hui Liu[1,2,3], Ting-Shuang Yi[1,2]

Affiliations
1. Germplasm Bank of Wild Species, Yunnan Key Laboratory of Crop Wild Relatives Omics,
Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China
2. University of Chinese Academy of Sciences, Beijing 100049, China
3. Key Laboratory of Plant Diversity and Specialty Crops, Chinese Academy of Sciences,
Beijing 100093, China
4. these authors contributed equally: Si-Yun Chen and Zhi-Yun Yang.
corresponding author(s): Hui Liu (liuhui1@mail.kib.ac.cn), Ting-Shuang Yi
(tingshuangyi@mail.kib.ac.cn)

Abstract

Zenia insignis Chun, from the subfamily Dialioideae of Fabaceae, is a tree species of
significant economic and ecological values. It is a threatened species as per the IUCN Red
List. The absence of a reference genome has impeded research on *Z. insignis* despite its
importance. In this study, we generated a reference genome for *Z. insignis*, with a contig
N50 of 6.02Mb, a total length of 361.11 Mb, and 97.71% of the sequences assembled into 14
pseudo-chromosomes. The BUSCO assessment score for completeness is 97.30%, and the
LAI index assessment score for continuity is 14.57. To our knowledge, this is the first
reported nuclear genome of a species belonging to subfamily Dialioideae. The reference
genome will provide a valuable resource for the phylogenomic studies of the Fabaceae
family and facilitate further research on *Z. insignis*.

Background & Summary

Zenia insignis Chun, a member of the family Fabaceae, is a tree species recognized as one of
the Grade-II Wild Plants of National Priority Protection in China[1] and is listed as Near
Threatened (NT) species on the International Union for Conservation of Nature (IUCN) Red
List[2]. This species was named by the renowned botanist Woon-Young Chun in honor of
Hongjun Ren (H.C. Zen)[3]. The species is primarily distributed in southern and southwestern
China at an altitude range of 200–1000 meters and northern Vietnam[4]. *Zenia insignis* boasts
a robust root system and strongly resists drought and infertile conditions[5]. In the karst
regions of southern and southwestern China, it has been utilized as an outstanding tree
species for afforestation on rocky terrain and as a pioneering species for the control of rocky
desertification[5-7]. Additionally, it has significant economic value. The wood of *Z. insignis* is

versatile and suitable for producing doors, boards, agricultural implements, and furniture[8,9].
The plant's tender leaves serve as livestock feed and green manure in rice fields[7-9].

Research on *Z. insignis* has predominantly concentrated on seedling[10,11], reforestation
technologies[5,6], and phylogenetic studies[12-19]. Though partial and complete plastome
sequences[12-14,16] and nuclear sequences[15,17-19] have been utilized in constructing the
phylogenetic tree of the Fabaceae family, the deep phylogenetic relationships within
Fabaceae remain to be fully elucidated. A chromosome-level genome sequence of *Z. insignis*
would be instrumental in clarifying these phylogenetic relationships. In this study, we have
successfully obtained the reference genome of *Z. insignis* and performed comparative
analyses with other representative plant genomes to further our understanding of the
evolutionary history of the Fabaceae.

In this study, we integrated multiple sequencing data types to assemble and annotate the
genome of *Z. insignis*. Specifically, we used PacBio continuous long reads (CLR) (174.14 Gb,
363.43× coverage), next-generation sequencing (NGS) reads (44.14 Gb, 91.96× coverage), Hi-
C data (59.63 Gb, 124.45× coverage), and full-length transcriptome reads. The resulting
reference genome consists of 14 pseudo-chromosomes and spans a total length of 352.84
Mb with a contig N50 of 6.02 Mb. The genome's LAI index score is 14.57. The GC content of
the genome is 34.38%, and the proportion of repeat sequences is 35.47%, with long terminal
repeat (LTR) elements being the most prevalent, accounting for 9.03%. We also obtained the
chloroplast genome sequence of *Z. insignis* with a length of 159,390 bp.

We predicted 33,322 gene models, of which 90.46% (30,143) were functionally
annotated. Analysis of orthologous genes between *Z. insignis* and ten other representative
plant species revealed 608 orthogroups, with at least 72.72% of species having single-copy
genes in any given orthogroup. Additionally, 1,625 gene families underwent significant
expansion, and 741 gene families underwent significant contraction in *Z. insignis*. In the
whole genome duplication analysis, two collinear block *Ks* peaks were observed at
0.245±0.001 and 1.693±0.001, suggesting that *Z. insignis* has undergone an additional whole

genome duplication (WGD) event after the γ WGT event shared by all core eudicots.

Methods

Genome size estimation and sequencing

Fresh young leaves of *Z. insignis* were collected from the Kunming Botanical Garden (KBG),
Kunming Institute of Botany, Chinese Academy of Sciences (KIB), Yunnan Province, China.
The voucher specimen (Collection Number: Yi20005) had been deposited at the Herbaria
(KUN) of Kunming Institute of Botany, Chinese Academy of Sciences, Yunnan Province,
China. The leaves were immediately flash-frozen in liquid nitrogen to preserve their cellular
structure and molecular content. Subsequently, the samples were transferred to an ultra-
low temperature freezer maintained at −80°C for storage.

DNA/RNA extraction, library construction, and sequencing processes were provided by
Wuhan Frasergen Bioinformatics Co., Ltd. The major steps are summarized below:

1. Genomic DNA was extracted from the tender leaves of *Z. insignis* using the modified
cetyltrimethylammonium bromide (CTAB) method[20,21]. Total RNA was isolated using a Trizol
reagent (Invitrogen, CA, USA). The quantity and quality of the DNA/RNA were determined
using a NanoDrop 2000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA).

2. The quality-compliant DNA was further tested for purity and integrity using 0.8%
agarose gel electrophoresis and a Qubit dsDNA HS Assay Kit on a Qubit 3.0 Fluorometer (Life
Technologies, Carlsbad, CA, USA). RNA contamination was identified using a 1.5% agarose
gel, and purity and integrity were evaluated using a NanoDrop 2000 and a Bioanalyzer 2100
system (Agilent Technologies, CA, USA).

3. According to the manufacturer's instructions, a short-read library was constructed
using the VAHTS Universal DNA Library Prep Kit (Vazyme, Nanjing, China) for MGI. The
library's quantification and sizing were measured using a Qubit 3.0 Fluorometer and a
Bioanalyzer 2100 system (Agilent Technologies, CA, USA). Paired-end sequencing of the
short-read library was conducted on an MGI-SEQ 2000 platform (BGI, China).

4. To perform long-read sequencing, the library was prepared in accordance with the
PacBio single-molecule real-time (SMRT) protocol and then sequenced on a Pacific
Biosciences Sequel II System in CLR mode.

5. The construction of the Hi-C library commenced with the cross-linking of fresh leaves
of *Z. insignis* under vacuum infiltration with 3% formaldehyde. Following lysis, digestion,
labelling, ligation, purification, adaptor addition, and PCR amplification, the Hi-C libraries
were quantified and sequenced on the MGI-SEQ 2000 platform.

6. For short-read transcriptome sequencing, RNA was isolated from Z. insignis young
leaves using the QIAGEN kit according to the manufacturer's instructions. The short-read
library was subsequently constructed and sequenced on the MGI-SEQ 2000 platform using
paired-end sequencing.

7. The full-length transcriptome sequencing was conducted using the Pacific Biosciences
DNA Template Prep Kit 2.0 for library construction (Pacific Biosciences, CA, USA).
Subsequently, the SMRT sequencing was performed on a Pacific Biosciences Sequel II
platform.
Our study generated 44.14 Gb of raw data for the whole genome, 174.14 Gb of PacBio
CLR data, 59.63 Gb of Hi-C data, and 81.92 Mb of full-length transcriptome data (Table 1).

The genome size of *Z. insignis* was assessed using flow cytometry analysis conducted on a
BD FACScalibur flow cytometer (BD Biosciences). The test sample preparation was
conducted in accordance with the manufacturer's guidelines and aligned with established

methodologies from previous literature[22,23]. As a standard reference, we utilized *Zea mays*
(maize)[24], a species with a well-documented genome size of 2.3 Gb. The analysis yielded a
result indicating that the genome size of *Z. insignis* is approximately 480 ± 4.8 Mb (Figure
1A).

A *K-mer* analysis was conducted on *Z. insignis* to estimate its genome size. Next-
generation sequencing (NGS) data was obtained from Frasergen and processed Fastp
v0.19.3[25]. This involved the removal of adapter sequences, the filtration of reads deemed to
be of insufficient length, and the elimination of those with low quality. Subsequently,
Jellyfish v2.2.10[26] was employed to calculate the frequency distribution of the cleaned data
using 21-*mers*. Subsequently, GenomeScope v2.0[27] was employed to estimate the
fundamental characteristics of the genome. The genome size of *Z. insignis* was estimated to
be approximately 364.94 Mb, with a heterozygosity rate of 0.842% and a repeat content of
1.17% (Figure 1B).

De novo genome assembly

Figure 2 illustrates the methodology employed for the chromosome-level genome assembly
of *Z. insignis*. The objective was to utilize three distinct assemblers, compare their resulting
assemblies, and achieve a high-quality assembly. The assemblers employed were Falcon
v0.3.0[28], Canu v2.1.1[29], and Flye v2.9[30]. The following section provides a summary of the
assembly process.

1. The Falcon software was employed to assemble the CLR long reads. The resulting
contigs were then polished using long reads with GCpp v2.0.0[31]. Subsequently, the processes
above yielded the assembly contig, designated as Assembly_Contig1. Subsequently,
Assembly_Contig1 was subjected to further refinement with short reads, employed
NextPolish v1.31[32]. The output from the process was designated as the 'Falcon contig'. To
eliminate redundancy, the Falcon contig was processed with Purge_Dups v1.2.5[33], and the
resulting non-redundant contigs were designated as the 'Falcon_purge contigs'.

2. The Canu software was employed to assemble the CLR long reads, and the result was
subsequently polished with long reads by GCpp. Through these processes, we obtained
Assembly_Contig2. Assembly_Contig2 was then polished with short reads using NextPolish,
with the output from this step designated as the 'Canu_contig'. The Canu_contig was
subsequently subjected to further processing to eliminate redundancy, with the result that
the non-redundant contigs were designated as the 'Canu_purged contigs'. Purge_Dups also
employed the process of eliminating redundancy.

3. The third assembler employed was Flye, which accepted the error-corrected reads
generated by Canu as input. The output from Flye was polished with long reads by GCpp,
resulting in the generation of Assembly_Contig3. Subsequently, the contig was polished with
short reads using NextPolish, resulting in the 'Flye_contig'. The Flye_contig was subjected to

a process of redundancy elimination using Purge_Dups, resulting in the generation of a set
of non-redundant contigs designated as the 'Flye_purge contigs'.

4. Six distinct results were obtained after completing the three steps above. The
following contigs were identified: Falcon contig, Canu contig, Flye_contig, Falcon_purge
contig, Canu_purge contig, and Flye_purge contig.

BUSCO v5.4.5[34] with 1614 embryophyta single-copy orthologs was employed to assess
the completeness of the assembly, the N50[35] metric provided insights into the distributions
of contigs, and the LTR Assembly Index (LAI) v2.9.0[36] was employed to evaluate the
contiguity of the assembled genome. The results are presented in Tables 2–4. Notably, the
Canu_purged assembly exhibited the highest N50 value, the Flye_purged assembly achieved
the optimal BUSCO score, and the Falcon_purged contig assembly exhibited the highest LAI
score. In other words, the result generated by Purge_Dups exhibited a better contiguity.
Manual inspection using Hi-C reads revealed a low redundancy in Falcon_purge assembly
(Supplementary Figure 1). Consequently, the results generated by Purge_Dups were
selected for further comparison. A comparison of the Falcon_purge assembly with other
results generated by Purge_Dups revealed that it exhibited high contiguity with N50 of 6.02
Mb, BUSCO completeness of 97.30%, and LAI score of 14.57.

During the chromosome assembly phase, we integrated Hi-C data with the previously
assembled contigs to achieve chromosome-level scaffolding. Specifically, Juicer was
employed to align the Hi-C reads to the contig genome. Subsequently, the 3D-DNA pipeline[37]
was used to map the contigs onto chromosome-level scaffolds. For a visual representation of
the Hi-C data, Juicebox v1.22[37] was employed. The results of the chromosome assembly also
demonstrated that the Purge_Dups results exhibited lower redundancy, compared to the
rest two assemblies (Supplementary Figure 1). The Purge_Dups result was chosen as the
optimal choice, and subsequently, the heatmap of the result was manually adjusted (Figure
3A). The final outcome of this process was the generation of 14 pseudo-chromosomes with a
total length of 352.84 Mb, as illustrated in Figures 3A and 3B.

The GetOrganelle v1.7.3.3[38] software was employed to assemble the chloroplast genome,
successfully acquiring the complete chloroplast genome sequence depicted in Figure 3C. The
newly assembled chloroplast genome spans 159,390 base pairs (bp) and is annotated using
the PGA tool[39], as shown in Figure 3C, drawn by OGDRAW v1.3.1[40]. For comparative analysis
of the chloroplast genome, we employed the MAFFT plugin within the Geneious Prime
software suite[41] to align our sequence with a published chloroplast genome of *Z. insignis*[16]
(NC_045299.1) obtained from the National Center for Biotechnology Information(NCBI). As
illustrated in Figure 3D, the alignment result revealed that apart from some single nucleotide
polymorphisms (SNPs) in non-coding regions, there were no significant differences in large-
scale sequences.

Repeat annotation

The RepeatModeler v2.0.1[42] software was employed to predict the occurrence of repeat
sequences within the genome of Z. insignis. RepeatModeler integrates three de novo repeat-
finding programs: RECON v1.08[43], RepeatScout v1.0.6[44], and LTRHarvest v5.9[45]/LTR_retriever
v2.9.0[46]. For our analysis, we employed rmblast v2.13.0[47] as the search engine.
Subsequently, the output from RepeatModeler was processed using RepeatMasker v4.1.4[48]
to identify and mask the repetitive elements within the Z. insignis genome.

Table 5 presents the results of the repeat sequence analysis for Z. insignis. A total of
128.08 Mb was identified as repetitive sequences, constituting 35.47% of the pseudo-
chromosome sequences. The analysis revealed four primary classes of transposable
elements within the genome: 9.03% for long terminal repeats (LTRs), 3.08% for DNA
elements, 1.81% for long interspersed nuclear elements (LINEs), and 0.08% for short
interspersed nuclear elements (SINEs).

Gene prediction and functional annotation

A hybrid approach was implemented to achieve high-quality gene prediction, incorporating
de novo prediction, homology-based prediction, and RNA-Seq-assisted prediction. We
employed a similar pipeline to the AnnoSmk v1.0[49] pipeline for gene prediction. Firstly, the
RepeatMasker v4.1.4[48] software was employed to generate repeat evidence. For homology-
based prediction, protein data for Arabidopsis thaliana[50] was sourced from The Arabidopsis
Information Resource (TAIR) and employed to construct homologous gene models using
genBlastG v1.38[51] as protein evidence. For transcriptome evidence, HISAT2 v2.2.1[52] was
employed to map paired-end short transcriptome reads to the Z. insignis genome. Then, the
mapped reads were assembled using Trinity v2.11.0[53]. Concurrently, we processed full-
length transcriptome data with the IsoSeq3 v3.4.0[54] pipeline to produce high-quality
consensus transcript sequences. Minimap2 v2.22[55] was employed to align both short and
long-read data, from which the transcriptome evidence was derived. The integrated repeat,
protein, and transcript evidence were then subjected to gene prediction using MAKER2
v3.01.03[56]. Three rounds of MAKER2 analysis were executed. In the second and third rounds,
the output generated through de novo gene prediction using AUGUSTUS v3.4.0[57] was also
included. BUSCO was employed to evaluate the gene prediction results after each round of
MAKER2, yielding scores of 46.60%, 96.60%, and 96.80%, respectively (Table 6). Following
the third round of MAKER2, the results were input into the PASA pipeline v2.5.0[58], resulting
in the final output. The final BUSCO score for the AnnoSmk pipeline was 97.20% (Table 6). In
conclusion, a total of 33,322 gene models were predicted.

The functional annotation of protein-coding genes was conducted using the eggNOG-
mapper website[59], which employs the eggNOG-mapper v2.1.12[60] and the eggNOG 5
database. This tool utilizes several comprehensive databases, including Pfam[61], InterPro[62],
Gene Ontology (GO)[63], and the Kyoto Encyclopedia of Genes and Genomes (KEGG)[64], to
annotate protein sequences. This analysis resulted in the successful functional annotation of
30,143 protein-coding genes, representing 90.46% of the total gene models.

Comparative genomics and phylogenetic analysis

The identification of orthologous genes represents a central objective in the fields of comparative genomics and phylogenetics. In this study, we compared the protein sequences of *Z. insignis* with those of ten other representative plant species, including *Arabidopsis thaliana*[50](GCA_000001735.2), *Cercis chinensis*[65](PRJCA007288), *Eriobotrya japonica*[66](GWHAAZU00000000.1), *Glycine max*[67](GCA_000004515.5), *Lysidice rhodostegia*[68], *Medicago truncatula*[69](GCA_003473485.2), *Senna tora*[70](GCA_014851425.1), *Quercus mongolica*[71](GCA_011696235.1), *Vitis vinifera*[72](GCA_030704535.1), and *Polygala tenuifolia*[73](TCMPG20196). These were employed to discern orthologs and paralogs. The OrthoFinder v2.5.4[74] software was used to predict orthologous and paralogous gene families. The protein sequences from the eleven species were input into OrthoFinder, resulting in the clustering of 39,196 genes into 30,977 gene families. The analysis revealed 8,698 orthogroups that are shared by all species. Of these, 40 orthogroups comprise single-copy genes, while 608 orthogroups have at least 72.72% of species exhibiting single-copy genes within each orthogroup.

We used the 608 orthogroups obtained from our analysis to construct phylogenetic trees. Initially, the seqkit tool v0.16.1[75] was employed to extract the corresponding nucleic acid sequences. Subsequently, the DNA sequences were aligned using MUSCLE v3.81511[76], and the sequences were concatenated using seqkit to generate a supermatrix of genes. The supermatrix was subsequently employed as the input for constructing the phylogenetic tree, which was generated using RAxML v8.2.12[77] with the maximum likelihood method. The most appropriate model for RAxML, GTR+F+R3, was determined using IQTree v2.1.4-beta[78]. Subsequently, the divergence times at the nodes of the phylogenetic tree using R8s v1.81, with calibration time points derived from the TimeTree database[79]. The calibration points included the divergence times between *M. truncatula* and *G. max* (55 million years ago, MYA), *M. truncatula* and *L. rhodostegia* (68 MYA), and *G. max* and *L. rhodostegia* (68 MYA). Finally, the CAFE v5.0[80] program was employed to understand the expansion and contraction of gene families across the phylogenetic tree. The results indicate that *Z. insignis* (subfamily: Dialioideae) diverged from *S. tora* (subfamily: Caesalpinioideae) and *M. truncatula* (subfamily: Papilionoideae) around 51 MYA. The analysis of gene family dynamics identified 1,625 gene families with significant expansion (LRT test, $p \leq 0.05$) and 741 gene families with significant contraction (LRT test, $p \leq 0.05$) in the genome of *Z. insignis* (Figure 4A).

We conducted GO and KEGG enrichment analyses to understand the biological significance of the gene families that exhibited significant expansion and contraction with *a P- value* cutoff of 0.05. The GO and KEGG analyses were performed using the eggNOG-mapper web platform[59]. We employed TBtools v2.086[81] to carry out the enrichment analysis of the identified gene families. The GO enrichment analysis revealed that the families that exhibited significant expansion were predominantly involved in molecular function and biological process (Supplementary Figure 2A). The terms of the molecular function are

mainly 'metal ion transmembrane transporter activity', 'inorganic cation transmembrane
transporter activity', and 'cellulose synthase activity' (Supplementary Figure 2A). And the
terms of the biological process are mainly 'hydrocarbon metabolic process' and 'olefin
metabolic process' (Supplementary Figure 2A). Furthermore, the KEGG enrichment analysis
revealed that these families are significantly enriched in pathways such as 'alpha-Linolenic
acid metabolism', 'MAPK signaling pathway-plant', 'metabolism of terpenoids and
polyketides', 'biosynthesis of other secondary metabolites', and 'ion channels'
(Supplementary Figure 2C). Conversely, for the families that exhibited a significant
contraction, the GO enrichment analysis was predominantly involved in the biological
process of 'embryo development ending in seed dormancy', 'embryo development', 'seed
development', 'fruit development', and 'proteolysis involved in protein catabolic process'
(Supplementary Figure 2B). The corresponding KEGG enrichment analysis identified
pathways including 'protein families: metabolism', 'ubiquitin mediated proteolysis',
'chromosome and associated proteins', 'protein processing in endoplasmic reticulum', and
'brite hierarchies' (Supplementary Figure 2D).

Identification of whole genome duplication events

To elucidate the whole genome duplication (WGD) events in *Z. insignis*, a comprehensive
analysis was conducted using the WGDI toolkit v0.6.5[82] with various parameters. We used *V.*
vinifera[72] as a reference genome. First, we used BLAST+ v2.15.0[83] to identify homologous
genes (e-value 10^{-5}). We then drew a dot plot of collinear genes with the parameter -d
(Figure 5A and B). Next, we conducted a collinear analysis with the parameter -icl, calculated
Ks values with the parameter -ks, combined the results of the collinear analysis and *Ks* values
with the parameter -bi, and fitted *Ks* peaks with the parameter -pf. Finally, we plotted a *Ks*
dot plot with the parameter -bk (Figure 5C and D) and a *Ks* peak plot with the parameter -kf
(Figure 4B). The dot-plot and *Ks* dot-plot results for *Z. insignis* showed a high degree of
similarity to those shown in Figure 4A and C. They were cross-validated with the results from
V. vinifera in Figure 5B and D. The *Ks* peak analysis revealed two peaks at 0.245±0.001 and
1.693±0.001, suggesting that *Z. insignis* likely experienced a lineage-specific WGD event, as
illustrated in Figure 4B.

Technical Validation

As part of our sequencing pipeline, we conducted a rigorous sample quality assessment for
both DNA and RNA samples used in the study. The specific procedures and criteria for this
assessment are detailed in our paper's 'Methods' section, ensuring that only high-quality
samples were selected for subsequent sequencing and analysis.

In our study, we employed two distinct methodologies to estimate the genome size of *Z.*
insignis. The first method involved Flow cytometry analysis, which provided a quick and
accurate measurement of the relative DNA content within the cells of *Z. insignis*. The second
method was *K-mer* analysis, which leverages the frequency distribution of DNA sequences to
predict genome size. The details have been mentioned in the 'Method' section. Both

approaches were crucial in validating the genome size and ensuring the reliability of our genomic data.

In our assessment of the *de novo* genome assembly for *Z. insignis*, we utilized a suite of metrics, including N50, BUSCO, and LAI, to evaluate the performance of various assembly software tools. The results of these evaluations are presented in Tables 2, 3, and 4, which compare the assembly quality, completeness, and continuity across different assemblers. Based on these assessments, Falcon-purged output demonstrated the most favorable balance of completeness and continuity, making it the choice for subsequent downstream analyses. In summary, we have successfully generated a high-quality reference genome for *Z. insignis*, which may promote further biological and comparative genomic studies of this species.

Code Availability

We used all software or pipelines, depending on the tools' manuals or protocols. The software's version is described in the methods section, and the parameters are given if not default.

Acknowledgements

This research was supported by the Major Program for Basic Research Project of Yunnan Province (202401BC070001), the Youth Talents Special Project of Yunnan Province 'Xingdian Talents Support Program' (XDYC-QNRC-2022-0423), the National Natural Science Foundation of China, key international (regional) cooperative research project (No. 31720103903), the Yunnan Provincial Basic Research Program Youth Project (2019FD058), and the National Natural Science Foundation of China (No. 32270247). We are grateful to the Germplasm Bank of Wild Species, particularly the Molecular Biology Experiment Center and the iFlora High Performance Computing Center, the Kunming Botanical Garden, Ms. Ding-Jie Wang, and Mr. Zuo-Ying Xiahou.

Author contributions

T.-S. Y. and H. L. Designed the experiments. S.-Y. C. and Z.-Y. Y. performed the experiments. S.-Y. C., H. L., and R. Z. analyzed the data. S.-Y. C., T.-S. Y. and H. L. wrote the paper. All Authors read and approved the manuscript.

Competing interests

The authors declare no competing interests.

References

1   The State Council. *List of Wild Plants of National Priority Protection(First batch)*, <https://www.gov.cn/gongbao/content/2000/content_60072.htm> (2000).

2   IUCN. *The IUCN Red List of Threatened Species*, <http://www.iucnredlist.org/>
(2024).

3   Chun, W.-Y. A new genus in the Chinese flora. *Sunyatsenia* **6**, 196–198 (1946).

4   Xu, L.-R. *et al. Fabaceae(Leguminosae)* In *Flora of China* Vol. 10. Wu, Z.-Y., Raven, P.
H., & Hong, D.-Y. Eds 27 Science Press & Missouri Botanical Garden Press (2010).

5   Ao, H., He, D., Zhang, Z., Xia, H. & Yang, W. *Zenia insignis* communities in limestone
areas of Guangdong. *Tropical Geography* **17**, 275–276+278–282,
http://dx.chinadoi.cn/10.13284/j.cnki.rddl.000227 (1997).

6   Zhang, Z.-P., He, D.-Q., Ao, H.-X., Yang, W.-J. & Wu, H.-M. The characteristic of
physiological ecology of the main tree species used for afforestation at limestone
regions in North Guangdong. *Acta Phytoecol. Sin.* **17**, 133–142 (1993).

7   Zhang, Y. Growth performance on acid soil and cultivation technique of *Zenia*
insignis. *Guangdong Forestry Science and Technology* **23**, 76–79,
http://dx.chinadoi.cn/10.3969/j.issn.1006-4427.2007.01.016 (2007).

8   Hou, L., Li, Y., Li, P., Yang, Y. & Xie, Y. Study of *Zenia insignis* Chun. comprehensive
utilization. *Sci. Silvae Sin.* **37**, 139–143, http://dx.chinadoi.cn/10.11707/j.1001-
7488.20010322 (2001).

9   Yu, G. & Zhang, Y. A preliminary study on the wood fibre form of *Zenia insignis*
introduced in Fujian Province. *Journal of Fujian Forestry Science and Technology* **25**,
31–33, http://dx.chinadoi.cn/10.3969/j.issn.1002-7351.1998.01.006 (1998).

10   Qin, Y.-R., Jiang, G.-M., Cen, Z.-Y., Chen, Y.-S. & Lan, C.-Y. Research on the seed's
germination characteristic of *Zenia insignis* as the vanguard tree for afforestation in
karst areas. *Seed* **27**, 15–21, http://dx.chinadoi.cn/10.3969/j.issn.1001-
4705.2008.12.005 (2008).

11   Tang, W., Pan, B., Wei, H., Huang, Z. & Lu, Q. Comparative study on the
characteristics of seed germination and seedling growth of different *Zenia insignis*
provenances in Guangxi. *Seed* **38**, 93–98,
http://dx.chinadoi.cn/10.16590/j.cnki.1001-4705.2019.12.93 (2019).

12   Bai, H.-R., Oyebanji, O., Zhang, R. & Yi, T.-S. Plastid phylogenomic insights into the
evolution of subfamily Dialioideae (Leguminosae). *Plant Diversity* **43**, 27–34,
https://doi.org/10.1016/j.pld.2020.06.008 (2021).

13   Herendeen, P. S. *Phylogenetic relationships in Caesalpinioid Legumes: a preliminary*
analysis based on morphological and molecular data In *Advances in legume*
systematics, part 10. Klitgaard, B. B. & Bruneau, A. Eds 37–62 Royal Botanic Gardens
(2003).

14   Kajita, T., Ohashi, H., Tateishi, Y., Bailey, C. D. & Doyle, J. J. rbcL and legume
phylogeny, with particular reference to Phaseoleae, Millettieae, and allies. *Syst. Bot.*
26, 515–536, http://dx.doi.org/10.1043/0363-6445-26.3.515 (2001).

15   Koenen, E. J. M. *et al.* Large-scale genomic sequence data resolve the deepest
divergences in the legume phylogeny and support a near-simultaneous evolutionary
origin of all six subfamilies. *New Phytol.* **225**, 1355–1369,
https://doi.org/10.1016/j.molp.2021.02.006 (2020).

16   Lai, Q., Tu, T. & Zhang, D. The complete plastid genome of *Zenia insignis* Chun
382   (Leguminosae). *Mitochondrial DNA Part B-Resour.* **4**, 2926–2927,
383   https://doi.org/10.1080/23802359.2019.1660266 (2019).
384   17   The Legume Phylogeny Working Group. A new subfamily classification of the
385   Leguminosae based on a taxonomically comprehensive phylogeny. *Taxon* **66**, 44–77,
386   https://doi.org/10.12705/661.3 (2017).
387   18   Zhang, R. *et al.* Exploration of plastid phylogenomic conflict yields new insights into
388   the deep relationships of Leguminosae. *Syst. Biol.* **69**, 613–622,
389   https://doi.org/10.1093/sysbio/syaa013 (2020).
390   19   Zhao, Y. *et al.* Nuclear phylotranscriptomics and phylogenomics support numerous
391   polyploidization events and hypotheses for the evolution of rhizobial nitrogen-fixing
392   symbiosis in Fabaceae. *Mol. Plant.* **14**, 748–773,
393   https://doi.org/10.1016/j.molp.2021.02.006 (2021).
394   20   Doyle, J. J. & Doyle, J. L. A rapid DNA isolation procedure for small quantities of fresh
395   leaf tissue. *Phytochemical Bulletin* **19**, 11–15 (1987).
396   21   Allen, G. C., Flores-Vergara, M. A., Krasynanski, S., Kumar, S. & Thompson, W. F. A
397   modified protocol for rapid DNA isolation from plant tissues using
398   cetyltrimethylammonium bromide. *Nat. Protoc.* **1**, 2320–2325,
399   https://doi.org/10.1038/nprot.2006.384 (2006).
400   22   Doležel, J. & Bartoš, J. Plant DNA flow cytometry and estimation of nuclear genome
401   size. *Ann. Bot.* **95**, 99–110, https://doi.org/10.1093/aob/mci005 (2005).
402   23   Doležel, J., Greilhuber, J. & Suda, J. Estimation of nuclear DNA content in plants
403   using flow cytometry. *Nat. Protoc.* **2**, 2233–2244,
404   https://doi.org/10.1038/nprot.2007.310 (2007).
405   24   Schnable, P. S. *et al.* The B73 maize genome: complexity, diversity, and dynamics.
406   *Science* **326**, 1112–1115, https://doi.org/10.1126/science.1178534 (2009).
407   25   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ
408   preprocessor. *Bioinformatics* **34**, i884–i890,
409   https://doi.org/10.1093/bioinformatics/bty560 (2018).
410   26   Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting
411   of occurrences of *k-mers*. *Bioinformatics* **27**, 764–770,
412   https://doi.org/10.1093/bioinformatics/btr011 (2011).
413   27   Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and
414   Smudgeplot for reference-free profiling of polyploid genomes. *Nat. Commun.* **11**,
415   1432, https://doi.org/10.1038/s41467-020-14998-3 (2020).
416   28   Chin, C.-S. *et al.* Phased diploid genome assembly with single-molecule real-time
417   sequencing. *Nat. Methods* **13**, 1050–1054, https://doi.org/10.1038/nmeth.4035
418   (2016).
419   29   Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer
420   weighting and repeat separation. *Genome Res.* **27**, 722–736,
421   https://doi.org/10.1101/gr.215087.116 (2017).
422   30   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads
423   using repeat graphs. *Nat. Biotechnol.* **37**, 540–546, https://doi.org/10.1038/s41587-
424   019-0072-8 (2019).

Seifert, D. & Töpfer, A. *GitHub*, <https://github.com/PacificBiosciences/gcpp> (2021).

Hu, J., Fan, J., Sun, Z. & Liu, S. NextPolish: a fast and efficient genome polishing tool for long-read assembly. *Bioinformatics* **36**, 2253–2255, https://doi.org/10.1093/bioinformatics/btz891 (2019).

Guan, D. *et al.* Identifying and removing haplotypic duplication in primary genome assemblies. *Bioinformatics* **36**, 2896–2898, https://doi.org/10.1093/bioinformatics/btaa025 (2020).

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M. BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.* **38**, 4647–4654, https://doi.org/10.1093/molbev/msab199 (2021).

Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921, https://doi.org/10.1038/35057062 (2001).

Ou, S., Chen, J. & Jiang, N. Assessing genome assembly quality using the LTR Assembly Index (LAI). *Nucleic Acids Res.* **46**, e126–e126, https://doi.org/10.1093/nar/gky730 (2018).

Dudchenko, O. *et al. De novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science* **356**, 92–95, https://doi.org/10.1126/science.aal3327 (2016).

Jin, J.-J. *et al.* GetOrganelle: a fast and versatile toolkit for accurate *de novo* assembly of organelle genomes. *Genome Biol.* **21**, 241, https://doi.org/10.1186/s13059-020-02154-5 (2020).

Qu, X.-J., Moore, M. J., Li, D.-Z. & Yi, T.-S. PGA: a software package for rapid, accurate, and flexible batch annotation of plastomes. *Plant Methods* **15**, 50, https://doi.org/10.1186/s13007-019-0435-7 (2019).

Greiner, S., Lehwark, P. & Bock, R. OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes. *Nucleic Acids Res.* **47**, W59–W64, https://doi.org/10.1093/nar/gkz238 (2019).

Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649, https://doi.org/10.1093/bioinformatics/bts199 (2012).

Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable element families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457, https://doi.org/10.1073/pnas.1921046117 (2020).

Bao, Z. & Eddy, S. R. Automated *de novo* identification of repeat sequence families in sequenced genomes. *Genome Res.* **12**, 1269–1276, https://doi.org/10.1101/gr.88502 (2002).

Price, A. L., Jones, N. C. & Pevzner, P. A. *De novo* identification of repeat families in large genomes. *Bioinformatics* **21**, i351–i358, https://doi.org/10.1093/bioinformatics/bti1018 (2005).

Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible software for *de novo* detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18, https://doi.org/10.1186/1471-2105-9-18 (2008).

46   Ou, S. & Jiang, N. LTR_retriever: a highly accurate and sensitive program for
identification of long terminal repeat retrotransposons. *Plant Physiol.* **176**, 1410–
1422, https://doi.org/10.1104/pp.17.01310 (2017).

47   Smit, A., Hubley, R. & Rosen, J. *RMBlast*, <https://www.repeatmasker.org/rmblast/>
(2024).

48   Chen, N. Using RepeatMasker to identify repetitive elements in genomic sequences.
Curr. Protoc. Bioinf. **5**, 4.10.11–14.10.14,
https://doi.org/10.1002/0471250953.bi0410s05 (2004).

49   Yan, H. & Liu, H. *AnnoSmk*, <https://github.com/yanhui-k/AnnoSmk> (2024).

50   The Arabidopsis Genome, I. Analysis of the genome sequence of the flowering plant
Arabidopsis thaliana. Nature **408**, 796–815, https://doi.org/10.1038/35048692
(2000).

51   She, R. *et al.* genBlastG: using BLAST searches to build homologous gene models.
Bioinformatics **27**, 2141–2143, https://doi.org/10.1093/bioinformatics/btr342
(2011).

52   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**,
907–915, https://doi.org/10.1038/s41587-019-0201-4 (2019).

53   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data
without a reference genome. *Nat. Biotechnol.* **29**, 644–652,
https://doi.org/10.1038/nbt.1883 (2011).

54   Wang, B. *et al.* Unveiling the complexity of the maize transcriptome by single-
molecule long-read sequencing. *Nat. Commun.* **7**, 11708,
https://doi.org/10.1038/ncomms11708 (2016).

55   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094–3100, https://doi.org/10.1093/bioinformatics/bty191 (2018).

56   Holt, C. & Yandell, M. MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. *BMC Bioinformatics* **12**,
491, https://doi.org/10.1186/1471-2105-12-491 (2011).

57   Stanke, M., Diekhans, M., Baertsch, R. & Haussler, D. Using native and syntenically
mapped cDNA alignments to improve *de novo* gene finding. *Bioinformatics* **24**, 637–
644, https://doi.org/10.1093/bioinformatics/btn013 (2008).

58   Haas, B. J. *et al.* Improving the Arabidopsis genome annotation using maximal
transcript alignment assemblies. *Nucleic Acids Res.* **31**, 5654–5666,
https://doi.org/10.1093/nar/gkg770 (2003).

59   *eggNOG-mapper*, <http://eggnog-mapper.embl.de/> (2024).

60   Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.
eggNOG-mapper v2: functional annotation, orthology assignments, and domain
prediction at the metagenomic scale. *Mol. Biol. Evol.* **38**, 5825–5829,
https://doi.org/10.1093/molbev/msab293 (2021).

61   Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**,
D412–D419, https://doi.org/10.1093/nar/gkaa913 (2020).

62   Paysan-Lafosse, T. *et al.* InterPro in 2022. *Nucleic Acids Res.* **51**, D418–D427,
https://doi.org/10.1093/nar/gkac993 (2022).

63 Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nature*
Genet. **25**, 25–29, https://doi.org/10.1038/75556 (2000).
64 Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M. & Tanabe, M. KEGG as a
reference resource for gene and protein annotation. *Nucleic Acids Res.* **44**, D457–
D462, https://doi.org/10.1093/nar/gkv1070 (2015).
65 Li, J. *et al.* The nearly complete assembly of the *Cercis chinensis* genome and
Fabaceae phylogenomic studies provide insights into new gene evolution. *Plant*
Commun., 100422, https://doi.org/10.1016/j.xplc.2022.100422 (2022).
66 Jiang, S., An, H., Xu, F. & Zhang, X. Chromosome-level genome assembly and
annotation of the loquat (*Eriobotrya japonica*) genome. *GigaScience* **9**, giaa015,
https://doi.org/10.1093/gigascience/giaa015 (2020).
67 Schmutz, J. *et al.* Genome sequence of the palaeopolyploid soybean. *Nature* **463**,
178–183, https://doi.org/10.1038/nature08670 (2010).
68 Liu, H. & Yi, T.-S. *unplished Lysidice rhodostegia data* (Kunming Institute of Botany,
Chinese Academy of Sciences, 2024).
69 Young, N. D. *et al.* The *Medicago* genome provides insight into the evolution of
rhizobial symbioses. *Nature* **480**, 520–524, https://doi.org/10.1038/nature10625
(2011).
70 Kang, S.-H. *et al.* Genome-enabled discovery of anthraquinone biosynthesis in *Senna*
tora. Nat. Commun. **11**, 5875, https://doi.org/10.1038/s41467-020-19681-1 (2020).
71 Ai, W. *et al.* A chromosome-scale genome assembly of the Mongolian oak (*Quercus*
mongolica). *Mol. Ecol. Resour.* **22**, 2396–2410, https://doi.org/10.1111/1755-
0998.13616 (2022).
72 Jaillon, O. *et al.* The grapevine genome sequence suggests ancestral hexaploidization
in major angiosperm phyla. *Nature* **449**, 463–467,
https://doi.org/10.1038/nature06148 (2007).
73 Meng, F. *et al.* Genome assembly of *Polygala tenuifolia* provides insights into its
karyotype evolution and triterpenoid saponin biosynthesis. *Hortic. Res.-England* **10**,
uhad139, https://doi.org/10.1093/hr/uhad139 (2023).
74 Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biol.* **20**, 238, https://doi.org/10.1186/s13059-019-
1832-y (2019).
75 Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. *PLoS One* **11**, e0163962,
https://doi.org/10.1371/journal.pone.0163962 (2016).
76 Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797, https://doi.org/10.1093/nar/gkh340
(2004).
77 Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312–1313,
https://doi.org/10.1093/bioinformatics/btu033 (2014).
78 Minh, B. Q. *et al.* IQ-TREE 2: new models and efficient methods for phylogenetic
inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534,
https://doi.org/10.1093/molbev/msaa015 (2020).

79 Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819,
https://doi.org/10.1093/molbev/msx116 (2017).
80 Mendes, F. K., Vanderpool, D., Fulton, B. & Hahn, M. W. CAFE 5 models variation in
evolutionary rates among gene families. *Bioinformatics* **36**, 5516–5518,
https://doi.org/10.1093/bioinformatics/btaa1022 (2020).
81 Chen, C. *et al.* TBtools-II: A "one for all, all for one" bioinformatics platform for
biological big-data mining. *Mol. Plant.* **16**, 1733–1742,
https://doi.org/10.1016/j.molp.2023.09.010 (2023).
82 Sun, P. *et al.* WGDI: A user-friendly toolkit for evolutionary analyses of whole-
genome duplications and ancestral karyotypes. *Mol. Plant.* **15**, 1841–1851,
https://doi.org/10.1016/j.molp.2022.10.018 (2022).
83 Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**,
421, https://doi.org/10.1186/1471-2105-10-421 (2009).
84 NCBI Sequence Read Archive. *SRR29423588,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423588> (2024).
85 NCBI Sequence Read Archive. *SRR29423585,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423585> (2024).
86 NCBI Sequence Read Archive. *SRR29423587,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423587> (2024).
87 NCBI Sequence Read Archive. *SRR29423586,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423586> (2024).
88 NCBI Sequence Read Archive. *SRR29423581,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423581> (2024).
89 NCBI Sequence Read Archive. *SRR29423580,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423580> (2024).
90 NCBI Sequence Read Archive. *SRR29423579,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423579> (2024).
91 NCBI Sequence Read Archive. *SRR29423578,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423578> (2024).
92 NCBI Sequence Read Archive. *SRR29423589,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423589> (2024).
93 NCBI Sequence Read Archive. *SRR29423583,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423583> (2024).
94 NCBI Sequence Read Archive. *SRR29423582,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423582> (2024).
95 NCBI Sequence Read Archive. *SRR29423584,*
<https://www.ncbi.nlm.nih.gov/sra/SRR29423584> (2024).
96 NCBI GenBank. *JBEVYO000000000,*
<https://www.ncbi.nlm.nih.gov/nuccore/JBEVYO000000000> (2024).
97 Science Data Bank. *Genomic squences of Zenia insignis,*
<https://doi.org/10.57760/sciencedb.10931> (2024).
98 Science Data Bank. *Annotation of Zenia insignis,*
<https://doi.org/10.57760/sciencedb.07828> (2024).

99   NCBI GenBank. *PP882779*, <https://www.ncbi.nlm.nih.gov/nuccore/PP882779>
602   (2024).

Figure 1 The result of genome size estimation for *Z. insignis*. (A) Flow cytometry-based
estimation result of *Z. insignis* (left peak, the sample) and *Zea mays* (right peak, the
reference). (B) The 21-*kmer* distribution estimation result was obtained using
GenomeScope2. The distribution of *21-kmer* exhibited two major peaks, indicating certain
heterozygosity within the genome, and high-frequency *k-mers* with coverage near 1 that was
possibly generated by sequencing errors.

Figure 2 The pipeline for the chromosome-level genome assembly of *Z. insignis*.

Figure 3 The chromosome and plastome information of *Z. insignis*. (A) The interaction heat map of Hi-C among chromosomes for the *Z. insignis* genome.

The Hi-C interaction matrix shows the pairwise correlations among the 14 pseudomolecules. (B) The high-quality genomic landscape of *Z. insignis*. The

genome diagram has four circles from the outside: gene density, GC content, chromosomal length, and intraspecies collinearity. (C)The visualization of the

plastome genome of *Z. insignis* using OGDRAW. The outside circle displays genes color-coded according to their functional groups. The second circle

illustrates the large single-copy (LSC), inverted repeat (IRA, IRB), and small single-copy (SSC) regions. The third circle delineates the percentage of GC
content and genomic positions in kilobase pairs. (D) Compared analysis of plastome for *Z. insignis*. Gene(green), CDS (yellow), RNA (red), and IR regions
(grey) have been color-coded according to their functional groups. The identity of the two plastome sequences exhibited high overall similarity, except for
some regions in non-coding sequences.

621
Figure 4 The result of gene family evolution and distribution of *Ks* values. (A) The result of
gene family evolution. The plus/minus sign corresponding numbers reflect the expansion or
contraction of gene families. The color of the branch indicates different subfamilies of
Fabaceae. (B) The distribution of *Ks* values between *Z. insignis* and *V. Vinifera*. The *Ks*
distribution curve of *Z. insignis*, *V. Vinifera*, *and Z. insignis vs. V. Vinifera* exhibited a
significant overlap in the right *Ks* peak.

628
Figure 5 The result of WGDI analysis for *Z. insignis*. (A) The homologous genes dot plot of
genomic synteny between chromosomes of *Z. insignis*. The best-hit genes are indicated in
red, the second best-hit genes are in blue, and the other hit genes are in gray. (B) Dot plot of
genomic synteny between *Z. insignis* and *V. Vinifera*. The best-hit genes are indicated in red,
the second best-hit genes in blue, and the other hit genes are in gray (C) *Ks* dot plot of
genomic synteny between chromosomes of *Z. insignis*. The color of the dot indicates the *Ks*
of the gene pair. (B) *Ks* dot plot of genomic synteny between *Z. insignis* and *V. Vinifera*. The
color of the dot indicates the *Ks* of the gene pair.

637
638 Supplementary Figure 1 The heat map of the Hi-C interaction matrix generated by Purge_Dups. (A) Heat map of Hi-C for Falcon_purge. (B) Heat map of Hi-C
639 for Canu_purge. (C) The Heat map of Hi-C for Flye_purge.

640
641 Supplementary Figure 2 The result of enrichment analyses for GO and KEGG. (A)The result of GO enrichment analyses for significant expansion gene families.
642 (B) The result of GO enrichment analyses for significant contraction gene families. (C) The result of KEGG enrichment analyses for significant expansion gene
643 families. (B) The result of KEGG enrichment analyses for significant contraction gene families.

Table 1 Statistics of sequencing data

Survey data	PacBio CCS data	Hi-C data	Full-length Transcriptome data
44140.85Mb	174,139.06Mb	59,630.13Mb	81.92Mb
91.96×	363.43×	124.45×	—

Table 2 The results of the assessment for N50

Nx	Canu_contig Size	Number	Canu_purged_contig Size	Number	Falcon_contig Size	Number	Falcon_purged_contig Size	Number	Flye_contig Size	Number	Flye_purged_contig Size	Number
N90	92,660	1,400	1,431,948	43	363,551	152	1,503,027	68	172,335	253	309,802	192
N80	113,319	822	4,428,622	31	1,150,475	87	2,685,811	49	529,601	137	653,152	114
N70	156,297	367	7,093,928	23	2,280,795	59	3,819,453	38	952,759	84	1,139,158	72
N60	320,232	71	8,669,316	18	3,626,040	43	4,916,748	29	1,478,745	53	1,600,360	46
N50	4,693,084	29	10,026,892	14	4,819,753	31	6,024,120	22	2,345,742	32	2,516,555	29
N40	7,292,602	20	12,366,826	11	5,750,257	23	7,011,975	17	3,977,660	20	4,128,350	18
N30	10,196,961	13	14,068,457	8	7,098,875	16	7,310,681	12	6,920,119	12	7,140,693	11
N20	14,068,457	8	14,855,825	5	7,527,208	9	8,509,734	7	9,355,035	7	9,355,035	7
N10	15,123,523	4	15,261,855	3	9,877,742	4	11,240,351	3	9,936,040	4	10,281,987	3
TotalLength	592,944,931	2,337	377,600,313	448	457,757,591	598	360,998,718	266	371,463,918	1,303	354,656,638	744

Table 3 The results of the assessment for BUSCO

	Canu_contig	Canu_purged_contig	Falcon_contig	Falcon_purged_contig	Flye_contig	Flye_purged_contig
Complete BUSCOs (C)	98.50%	98.50%	97.50%	97.30%	98.60%	98.60%
Complete and single-copy BUSCOs (S)	44.10%	77.30%	65.80%	79.80%	76.70%	76.90%
Complete and duplicated BUSCOs (D)	54.40%	21.20%	31.70%	18.40%	21.90%	21.70%
Fragmented BUSCOs (F)	0.30%	0.60%	1.00%	1.20%	0.40%	0.60%
Missing BUSCOs (M)	1.20%	0.90%	1.50%	1.50%	1.00%	0.80%
Total BUSCO groups	1614	1614	1614	1614	1614	1614

647 Table 4 The results of the assessment for LAI

	Chromosome From	To	Intact	Total	raw_LAI	LAI
Canu_contig	1	592,944,931	0.0088	0.2003	4.38	0
Canu_purged_contig	1	377,600,313	0.0126	0.2099	5.98	11.61
Falcon_contig	1	457,757,591	0.0077	0.1684	4.57	4.62
Falcon_purged_contig	1	360,998,718	0.0119	0.1329	8.94	14.57
Flye_contig	1	371,463,918	0.0109	0.1868	5.83	11.46
Flye_purged_contig	1	354,656,638	0.0108	0.1689	6.4	12.03

648 Table 5 The result of repeat sequences annotation

Type	Length(bp)	Rate (%)
DNAs	11,129,683	3.08
LINEs	6,520,335	1.81
SINEs	298,378	0.08
LTRs	32,617,056	9.03
Rolling-circles	1,280,052	0.35
Small RNA	6,975,086	1.93
Satellites	3,559,151	0.99
Simple repeats	1,063,047	0.29
Low complexity	64,922,128	17.98
Unknown	128,083,818	35.47
Total	11,129,683	3.08

649 Table 6 The result of BUSCO for gene prediction

	R1	R2	R3	R3+PASA
Complete BUSCOs (C)	46.60%	96.60%	96.80%	97.20%
Complete and single-copy BUSCOs (S)	39.90%	44.80%	43.90%	44.20%
Complete and duplicated BUSCOs (D)	6.70%	51.80%	52.90%	53.00%
Fragmented BUSCOs (F)	3.50%	2.30%	2.30%	1.90%
Missing BUSCOs (M)	49.90%	1.10%	0.90%	0.90%
Total BUSCO groups	1614	1614	1614	1614