An Unbiased Proteomic Platform for ATE1-based Arginylation Profiling

Zongtao Lin[1,*], Yixuan Xie[1], Joanna Gongora[1], Xingyu Liu[1], Emily Zahn[1], Bibhuti Bhusana Palai[1], Daniel H.
Ramirez[1], Richard M. Searfoss[1], Francisca N. Vitorino[1], Rashmi Karki[1], Geoffrey P. Dann[2], Chenfeng Zhao[3],
Xian Han[1], Brittany MacTaggart[4], Xin Lan[5], Dechen Fu[5], Lina Greenberg[1], Yi Zhang[5], Kory J. Lavine[6],
Michael J. Greenberg[1], Dongwen Lv[7], Anna Kashina[4], Benjamin A. Garcia[1,*]

*[1]Department of Biochemistry and Molecular Biophysics, Washington University in St. Louis, St. Louis, MO
63110*

[2]Department of Biochemistry and Biophysics, University of Pennsylvania, Philadelphia, PA, 19104

[3]McKelvey School of Engineering, Washington University in St. Louis, St. Louis, MO 63110

[4]School of Veterinary Medicine, University of Pennsylvania, Philadelphia, PA 19104

[5]Department of Biochemistry, Case Western Reserve University, Cleveland, OH 44106

[6]Department of Medicine, Washington University School of Medicine, St. Louis, MO, USA

*[7]Department of Biochemistry and Structural Biology, University of Texas Health Science Center at San
Antonio, San Antonio, TX 78229*

Correspondence. **Email**: *bagarcia@wustl.edu (B.A.G.), zongtao@wustl.edu (Z.L.),* **Tel.**: *314-362-3085*

Abstract. Protein arginylation is an essential posttranslational modification (PTM) catalyzed by arginyl-
tRNA-protein transferase 1 (ATE1) in mammalian systems. Arginylation features a post-translational
conjugation of an arginyl to a protein, making it extremely challenging to differentiate from translational
arginine residues with the same mass in a protein sequence. Here we present a general ATE1-based
arginylation profiling platform for the unbiased discovery of arginylation substrates and their precise
modification sites. This method integrates isotopic arginine labeling into an ATE1 assay utilizing biological
lysates (*ex vivo*) rather than live cells, thus eliminating translational bias derived from the ribosomal activity
and enabling *bona fide* arginylation identification using isotopic features. The method has been successfully
applied to an array of peptide, protein, cell, patient, and animal tissue samples using 20 µg sample input,
with 235 unique arginylation sites revealed from human proteomes. Representative sites were validated
and followed up for their biological functions. The developed platform is globally applicable to the
aforementioned sample types and therefore paves the way for functional studies of this difficult-to-
characterize protein modification.

Keywords: arginylation, arginyltransferase, proteomics, posttranslational modification, protein profiling

Introduction

As a critical posttranslational modification (PTM), arginylation is catalyzed by arginyltransferase
(ATE1) which is the only known enzyme capable of installing arginylation in mammalian systems[1]. The
absence of arginylation in ATE1 knockout (KO) models resulted in embryonic lethality due to heart defects
in cardiovascular development and angiogenic remodeling[1]. Specifically, ATE1KO caused thinned
myocardium with immature septa, non-separation of the aorta and pulmonary artery, resulting in defects in
cardiac contractility, myofibril dysfunction and eventually embryonic death[1], demonstrating the essential
nature of arginylation[1]. Tissue-specific knockdown (KD) or deletion (KO) of arginylation resulted in impaired
myosin phosphorylation and thrombus formation[2], elevated myocardial fibrosis and progressive heart
failure[3], cardiomyocyte hypertrophy[4], and many other symptoms[5,6]. At the molecular level, the arginylation
field has long tried to understand the biological roles of arginylation in cardiovascular-associated proteins
including β-actin (Asp3 arginylation) regulating cytoskeleton and cell motility[7,8], calreticulin (CALR, Asp18
arginylation) regulating stress granules in a Ca^{2+}-dependent manner[9,10], and RGS4/5/16 (tri-oxidized Cys2
arginylation) acting as nitric oxide and oxygen sensors[11,12]. Many of these studies focused on the
implications of shortened half-lives of arginylated proteins involved in cardiovascular biology[1,7-12].

Meanwhile, many studies suggested that arginylation is also important for certain proteins to function
properly (noncanonical roles) beyond its canonical role in the *N*-degron pathway[13,14]. It is reported that
arginylation of β-amyloid guides proper α-helical shape preventing misfolding and aggregation[15].
Arginylation of α-synuclein (α-syn) facilitates brain health by preventing neurodegeneration[16,17]. Arginylation
of calreticulin promotes its association with stress granules[18,19]. Decreased arginylation of nuclear proteins
results in smaller nucleus size and architecture[20]. In addition, arginylation has been detected on histone
proteins[20,21], potentially facilitating their interaction with DNA through the positively charged Arg residue
installed post-translationally. Other prominent arginylated examples are chaperone HSPA5[22], BRCA1[23],
PDI[24], and CDC6[25]. It is becoming clear that protein arginylation has many biological functions yet to be
determined in regulating physiological processes. However, only a handful of proteins have been validated
for their arginylation sites so far, and the discovery of arginylation substrates and their modification sites is
becoming a bottleneck for the study of arginylation biology.

Identification of arginylation has been inherently challenging because both ATE1 and ribosomes use
the same arginine (Arg or R) source, arginyl-tRNA, to add an arginyl group to a protein, yielding
posttranslational and translational addition respectively of arginyl with the same mass shift (+156 Da)[13] (**Fig.**
1a). This makes unbiased differentiation of posttranslational arginylation and translational arginine residues
extremely difficult when both ATE1 and the ribosome are active (*i.e.* when cells are alive). In addition,
arginylated proteins may go through the Arg/*N*-degron pathway for rapid ubiquitin-mediated degradation,
decreasing the endogenous arginylation level for detection[13,25]. Furthermore, arginylation mostly happens
on the protein *N*-termini after proteolytic cleavage, which is poorly understood for individual proteins in the
whole proteome[13]. Additional confusion comes from trypsin miscleavage resulting in peptides starting with
an R residue in bottom-up proteomics. Disregarding those challenges, efforts have been made aiming to
systematically identify arginylation.

One early approach to screening arginylation substrates involved adding sub-divisions of a complete
cDNA library to a transcription-translation-degradation system, which assumes that arginylated substrate is
prone to the *N*-degron pathway[26]. A few arginylation substrates (*e.g.*, BiP and PDI) have been identified
from the whole proteome through the combination of ATE1 assay *ex vivo* and [³H]arginine
autoradiography[24]. The autoradiography was also used for further identification of individual proteins (*e.g.*,
CALR[10] and α-syn[16]) as ATE1 substrates. Notably, these methods heavily relied on mass spectrometry to
reveal the arginylation sites on the labeled (*ex vivo*) proteins. In contrast, proteomic profiling was attempted
to identify the endogenous (*in vivo*) arginylation sites from animal tissues using anti-arginylation antibody
enrichment and proteomics. Such an approach enabled the first proteomics-based analysis to identify 43
plausible arginylated proteins[21]. Similarly, two later studies identified 19 and 15 proteins potentially modified
by arginylation on their side-chain Asp and Glu residues[27,28], a different ATE1 catalysis mechanism from
canonical *N*-term arginylation. However, most of those sites have little overlap with existing confirmed
proteins/sites from *ex vivo* experiments and were largely postulated from search algorithms without
additional validation. Therefore, there is still a huge unmet need for the creation of an unbiased profiling
method for the arginylation field at this stage.

Inspired by unbiased labeling using isotopic Arg[24] and the screening power of mass spectrometry-
based proteomics[21], here we present a profiling technology for the unbiased discovery of ATE1 substrates
and arginylation sites from complex whole proteomes (**Fig. 1b**). We first established the workflow using
peptide and proteome-wide peptides as a proof-of-concept, then applied the technology to protein and
whole proteomes for arginylation discovery. As a result, a large catalog of unbiased arginylation sites has
been established from various cell and tissue samples. To facilitate data processing, a new computational
tool, "ArginylomePlot", was developed and made publicly available for isotopic pair analysis. To be
beneficial to the arginylation field and broader research community, we have established a publicly available
database indexing all arginylation sites discovered from this study, allowing others interested in protein
arginylation to have easy access to this growing type of data. This work could serve as the technological
foundation for studying the functions of this essential PTM, and it will have a long-lasting impact on the
arginylation field by opening new biochemical and biological frontiers.

Results

Isotopic arginylation of substrate peptide by ATE1 *in vitro*. To test the proof-of-concept of the
designed arginylation profiling strategy based on ATE1 enzymatic activity (**Fig. 1b**), we started by
isotopically labeling a standard peptide (modified from human β-actin sequence) with an unmodified Asp
(D) residue at the *N*-terminus (**Fig. 1c**, **Fig. S1**). The reconstituted ATE1 assay was slightly modified from
the well-established conditions in our previous report[29], where arginine-tRNA ligase (RARS) and ATE1
enzymes were kept at μM range. The arginylation activity of the assay was tested to be dependent on
essential components such as arginine, ATP, tRNA, RARS, ATE1, and substrate (**Fig. S2**). Replacement of
N-terminal D residue to acetylated D or Val produced no detectable arginylation products (**Fig. S3-4**). The
arginylation of the standard peptide was time-dependent (0-60 min), and most of the peptides were
arginylated with an incubation time of 30 min or longer (**Fig. S5**). Higher starting concentrations (20-300
μM) of substrate produced more arginylation product but correlated to lower overall product yields (**Fig. S5**).
The incubation time and substrate concentration were set at 30 min and 100 μM, respectively, while arginine
(2 mM) and ATP (2 mM) remained excessive.

After the ATE1 assay using isotopic Arg[10] and Arg[0] respectively, the reactions on standard peptide
were mixed (1:1) and desalted for LC-MS analysis (**Fig. 1c**). The extracted ion chromatograms of the
differentially labeled products were manually examined for hypothetical isotopic ratio (1.0) based on co-
elution peak intensities (**Fig. 1c**), demonstrating that the isotopic labeling strategy was capable of
introducing unnatural Arg[10] to arginylation sites for unbiased identification. The data was also searched by
Byonic to confirm the product identities with *N*-terminal arginylation (**Fig. S1**). The MS1 spectra of isotopic
products were extracted to validate the Arg[10]/Arg[0] pair features in doublets ("1+1" = 2 isotopic peaks, error
$\leqslant$ 10 ppm) and summarized into a boxplot for relative intensity overviews (**Fig. 1d**). The co-eluting isotopic
peaks indicated unbiased identification of arginylation. The average value of H/L intensity ratios calculated
from individual MS1 spectra was considered the H/L ratio in an LC-MS run. The isotopic labeling of the
standard peptide was replicated 4 times and produced consistent ratio profiles (**Fig. 1e**). An increasingly
stringent MS1 pair detection using co-eluting quartet ("2+2" = 4 isotopic peaks, error ≤ 10 ppm) and sextet
("3+3" = 6 isotopic peaks, error ≤ 10 ppm) yielded H/L ratios closer to hypothetical 1.0 than doublet ("1+1"
= 2 isotopic peaks) but gave decreased numbers of MS1 pairs in all replicates. To ensure all MS1 pairs are
included, we decided to use MS1 doublet as the threshold filter and include duodecet ("6+6" peaks) for pair
overview as shown in **Fig. 1d**. In addition, when the isotopically labeled peptides were mixed at other ratios
(R[10]/R[0]-peptide = 1:2 and 7:10) than 1:1 used previously, experimental H/L ratios around 0.5 and 0.7 were
observed expectedly (**Fig. S6-7**), another approach to confirm the labeled peptides. For the convenience
of experiments and data analysis, an isotopic 1:1 mix strategy with a hypothetical H/L ratio of 1.0 is used in
the rest of this study.

Figure 1. Isotopic arginine labeling and detection strategy for ATE1 substrates and arginylation sites. a, scheme of arginyl installation onto proteins by ribosomal synthesis (translational) and ATE1 (post-translational). RARS, arginine-tRNA ligase. ATE1, arginyl-tRNA-protein transferase 1. **b,** arginylation profiling platform for arginylation site and substrate discovery from biological samples. Lysate is labeled by isotopic arginine molecules (Arg^{10} and Arg^0) respectively using ATE1 assay, mixed, and digested. Peptides are fractionated and analyzed by mass spectrometry in data-dependent acquisition (DDA) mode. Proteomics data is searched to produce peptide identifications (IDs), among which peptide pairs modified by heavy and light Arg are further evaluated for MS1 isotopic features. **c**, isotopic arginylation of a peptide ATE1 substrate. EICs were extracted using monoisotopic peaks based on calculated *m/z* values. EIC in black indicates chromatography of the unmodified peptide. **d**, isotopic feature in MS1 spectra and their

summary. The first monoisotopic peak in each MS1 scan is set at 1 for normalization. Relative intensities
of other isotopic peaks are displayed. 10-ppm error is set for all MS1 isotopic peaks. **e**, ratio summary of
MS1 pairs in 4 replicates ($n = 4$) using doublet, quartet, and sextet peaks. Detailed ratios from the replicates
are provided. The numbers of pairs in quartet and sextet are normalized to the numbers of pairs in doublet
from respective replicate. EIC: extracted ion chromatogram. Nle: norleucine.

Development of in-house software for dataset analysis. Encouraged by the proof-of-concept
data, we then aimed to develop a computational tool "ArginylomePlot" (**Fig. 2a**) with several capabilities to
facilitate handling datasets from future complex samples. The software should be able to 1) filter out MS2
pairs with the same sequence modified by Arg[10] and Arg[0] (Δmass = 10.008269 Da) disregarding other
modifications (*e.g.*, Met oxidation and Cys carbamidomethylation), 2) extract their MS1 scans from raw data,
3) summarize heavy/light ratios of each arginylation peptide, and 4) export paired MS1 and MS2 scans for
data visualization and indexing. The software is publicly available to download on GitHub
(https://github.com/BeckyHan/Garcia-Lab/tree/main/ArginylomePlot). Briefly, mzXML data converted from
raw data was used as the source input data, and search results from proteomics software (*e.g.*, Byonic)
were used as peptide input. Once a co-elution MS2 pair was identified from the peptide list, their MS1
information (*e.g.*, *m/z* and charge) was used to extract pairs from mzXML for ratio calculation. To present
high-confidence proteomics data, peptide score (H or L ≥ 300) was applied as a filter before MS2 spectrum
export. In the end, the software also exports spectra as Excel files for establishing a website database. An
exemplary dataset and detailed instructions to execute the software are provided (**Supplementary Dataset**
1).

Figure 2. Arginylation analysis of whole-proteome peptides using in-house software. a, software
flowchart of customized "ArginylomePlot". Input data are mzXML and peptide identification files. Output
data are MS1, MS2, and the summary of unbiased arginylation sites. **b,** experimental workflow for
arginylation of tryptic peptides from HEK293T (ATE1 KO) cells. **c,** numbers of identified MS1 pairs,
arginylation sites, unique peptides, and unique proteins. **d,** H/L ratio distribution of all MS1 pairs and their
corresponding numbers of MS1 scans (doublets, error ≤ 10 ppm). The retention times of peptide IDs
belonging to a pair were averaged. The software will extract matching MS1 scans within 1.25 min (± 1.25
min) of the averaged retention times. **e,** analysis of the *N*-terminal residues of all unique arginylated
peptides. Q is arginylated after deamidation (Q→E). **f,** sequence logo calculated from unmodified forms of
all unique arginylated peptides. The frequency plot is generated by WebLogo. The peptide sequences were
aligned and extended to include 14 positions downstream of the arginylation sites as the P1' position. **g,**
comparison of tryptic and non-tryptic *N*-term of all arginylated peptides. Non-tryptic *N*-term may indicate
endogenously exposed *N*-termini of proteins.

Isotopic arginylation of HEK293T peptides by ATE1 *in vitro*. Assisted by the software, we next
tested the arginylation of a whole-proteome peptide mixture prepared from the whole proteome of HEK293T
cells with ATE1 knocked out to avoid potential interference from endogenous arginylation (**Fig. 2b**). ATE1
KO has been confirmed by genomic sequencing and Western blotting (section Methods, **Supplementary**
Information). After data analysis, a total of 772 arginylation pairs belonging to 167 unique arginylation sites
were detected at the MS1 level by the software (**Fig. 2c**, **Fig. S8**, **Table S1**, **Supplementary Dataset 2**).
The data showed exclusive *N*-terminal arginylation (**Supplementary Dataset 2**), even though we searched
for both *N*-terminal and sidechain arginylation at the same time. While charges 2 (83.7%) and 3 (13.8%)
account for 97.5% of all identified peptides, most of the arginylated peptide pairs (98.3%) are detected at
charge states 2 (47.3%) and 3 (51.0%) with a small portion in charge 4, indicating a charge shift for
arginylated peptides (**Fig. S9**). The distribution profile of H/L ratios (threshold: $0.1 \leq ratio \leq 10$) from all MS1
pairs is centered at 1.0 (**Fig. 2d**) which is the hypothetical ratio for each labeled peptide per experimental
design, demonstrating the success of the workflow for complex peptide sample. The doublet ratios were
also compared with the quartet and sextet profiles (**Fig. S10**). When a peptide pair was observed with the
same charge state and a Δmass of 10.008269 Da, the averaged retention time of relevant peptide IDs
containing both Arg^{10} and Arg^0 was used to create a ± 1.25 min window for MS1 paired scans extraction.
MS1 scans containing doublets (10-ppm error) were exported. The average value of H/L intensity ratios
calculated from individual MS1 scans was considered the H/L ratio of a peptide pair. The numbers of MS1
scans with the doublet feature used for generating respective H/L ratios are provided (**Fig. 2d**, **Fig. S11**).
When looking at the arginylated *N*-term residues of all unique peptides, a majority of arginylation happened
on D and E (including Q→E) residues (159 out of 165, 96.36%) (**Fig. 2e**) while the data also suggested
possible arginylation on other *N*-term residues (*e.g.*, A, G, L and V), consistent with reported specificities
and activities of ATE1[30,31]. We then generated the sequence logo of unique peptides using WebLogo 3.0[32]
(**Fig. 2f**) and the arginylation motif using pLogo[33] (**Fig. S12**). The result suggested that arginylation favors
N-term D and E residues significantly, while subsequent nonpolar residues (I, L, F, V and A) and acidic
residues (D and E) may facilitate the peptide *N*-term arginylation (**Fig. 2f**, **Fig. S12**). While most of the
unique peptides (158 out of 165, 95.76%) are tryptic after K or R cleavage, a few non-tryptic peptides were
arginylated (**Fig. 2g**). For example, the arginylation of SSBP E17 was on a non-tryptic peptide (**Fig. S13**),
this may have resulted from endogenous protein *N*-term processing and cleavage[34]. This observation
demonstrates the potential application of our peptide workflow for the discovery of the cleavage and
arginylation of endogenous protein *N*-term.

Isotopic arginylation of protein substrates by ATE1 *in vitro*. We then moved forward from
peptide to protein arginylation using CALR since it is known for *N*-term arginylation at the E18 site after
endogenous cleavage of a signaling peptide (AA 1-17)[18]. Purified 18E-CALR after overexpression in
HEK293T ATE1 KO cells was isotopically labeled and yielded the H/L ratio of 1.0 in the arginylated E18
peptide (R^{10}/R^0-E_{18}PAVYFK) (**Fig. 3a** and **Fig. S14**). The *in vitro* arginylation efficiency of CALR is
preliminarily estimated to be 64.7% for the E18 site based on MS2 spectrum counts of modified (R^{10}/R^0-
EPAVYFK, counts = 33) and unmodified (EPAVYFK, counts = 18) peptides. When counting the unmodified
peptide (EPAVYFK) in samples with and without ATE1, the arginylation efficiency was estimated to be 80.2%
(n = 3, **Fig. S15**). Replication of CALR arginylation (n = 5) showed consistent ratios near 1.0 (**Fig. 3a** and
Fig. S16). In addition, arginylation results (n = 5) of a commercial 18E-CALR (Abcam, ab276554) showed
consistent E18 labeling (**Fig. S17**). In comparison, the 18R-CALR protein was purified and tested, and the
results (n = 5) showed nearly no R^{10} arginylation due to the absence of an open *N*-term E18 residue (**Fig.**
S18). The results indicated that the isotopic arginylation strategy could be used to discover modification
sites in pure proteins.

Figure 3. Summary of arginylation sites and ATE1 substrates in human proteins and proteomes. a,
experimental workflow for CALR arginylation. **b,** numbers of identified MS1 pairs, arginylation sites, unique
peptides and unique proteins in human cells and patient tissues. **c,** overview of arginylation sites in human
cells and patient tissues. **d**, ratio plot of all MS1 pairs detected in all sample fractions. **e**, sequence logo
calculated from unmodified forms of all unique arginylated peptides. Frequency plots are generated by
WebLogo. The arrow indicates the cleavage site before arginylation. **f**, arginylation type comparison of all
unique sites. **g**, biological function analysis of ATE1 protein substrates using PANTHER. iPSC, induced
pluripotent stem cell. CF, cardiac fibroblasts. CM, cardiomyocytes. R, arginylation. RO3, Cys tri-oxidation
and arginylation. RO2, Cys di-oxidation and arginylation. R_deami, N/Q arginylation after deamidation. PD,
Parkinson's disease. PDD, PD with dementia. AD, Alzheimer's disease.

Table 1. Number of detections in isotopic pairs and MS1 scans for each arginylation site from human proteomes.

Site	n1	n2	Site	n1	n2	Site	n1	n2	Site	n1	n2	Site	n1	n2
*CALR E18 R	185	8434	LMAN2 D45 R	6	147	CNBP Q44 R_deami	3	84	NDKM K91 R	1	192	VIME D451 R	1	26
MBP D168 R	139	5153	MTAP2 Q431 R_deami	6	122	MYH7 E1902 R	3	71	ACTC E109 R	1	121	HOOK1 D659 R	1	25
*PDIA1 D18 R	135	6071	SRCA E21 R	6	108	CALU D62 R	3	70	AKAP2 D638 R	1	84	MYH7 E1489 R	1	25
*SSBP E17 R	127	4744	CBS E 4 R	6	101	KTN1 D238 R	3	61	ATAD5 V1449 R	1	70	TBA4A D69 R	1	25
*RM12 C45 RO3	115	3085	MYH7 E1461 R	6	96	HACD3 P100 R	3	58	SRRT D868 R	1	70	ACTA E318 R	1	24
NFL D468 R	66	3260	ACTB E241 R	6	79	PLEC E4050 R	3	56	SFPQ E696 R	1	68	HNRPK Q358 R_deami	1	24
SAP D405 R	52	1485	H2B1K E94 R	6	35	MYH7 D1229 R	3	54	VATD E242 R	1	63	MYH7 E1376 R	1	24
A1AT E25 R	48	1648	NFM E497 R	5	321	TBA1B D327 R	3	52	CC148 I574 R	1	54	MYH6 E73 R	1	23
CATD E64 R	38	986	MBP T200 R	5	264	NFL Q75 R_deami	3	46	NFM E884 R	1	51	NOP16 E77 R	1	23
CATD G65 R	36	1008	CH60 Q42 R_deami	5	207	CALD1 E161 R	3	42	HMGA1 E47 R	1	49	NPC2 E20 R	1	23
*ERO1A E24 R	36	984	LN28A E18 R	5	200	NT5D3 C44 RO3	3	35	RS7 E188 R	1	49	ACLY E830 R	1	21
F162A C41 RO3	26	516	LETM1 D116 R	5	188	PTBP1 D 7 R	2	124	ACTA E109 R	1	47	ATS15 D749 R	1	21
TBB5 E 3 R	25	1342	TNNI3 E165 R	5	169	FCN1 D120 R	2	111	MYOZ2 D255 R	1	44	ATP5J N33 R_deami	1	20
PDIA4 E25 R	25	756	VIME E425 R	5	166	HS90B E539 R	2	102	NCAM1 E833 R	1	43	KS6A1 D37 R	1	20
GFAP D417 R	22	722	NFL E453 R	5	145	LDB3 E257 R	2	79	BI1 N9 R_deami	1	42	ATIF1 E42 R	1	19
G3P D296 R	22	482	MYL3 E131 R	5	119	NPM D178 R	2	77	CH60 D209 R	1	41	CH60 D100 R	1	18
ODO2 D68 R	21	735	CH60 D49 R	5	111	ROA2 D161 R	2	76	RON D1232 R	1	41	HNRPL D284 R	1	18
MBP N226 R_deami	21	364	TXND5 E40 R	5	94	VIME D90 R	2	76	CH60 C447 R	1	40	MYH7 E927 R	1	18
MBP E217 R	20	582	SPTN1 E1124 R	5	61	S10AE C74 RO3	2	75	DJC10 D33 R	1	40	SNX17 E463 R	1	18
MBP G277 R	18	488	RL23A E 8 R	5	59	XRCC5 E583 R	2	66	RM12 C45 R	1	40	JTB E31 R	1	17
H4 D25 R	18	466	PGRC1 D49 R	4	201	NPM D180 R	2	63	RT18A E37 R	1	40	MITOK C37 RO3	1	17
MBP D216 R	17	550	NFL E469 R	4	193	TCAF1 V114 R	2	59	MYH9 D1846 R	1	39	RS27 D 6 R	1	17
NFM E454 R	17	466	CASQ2 E20 R	4	151	MAP1A E663 R	2	58	DREB D636 R	1	38	HBA D48 R	1	16
PGCB D23 R	16	772	TBA1B D69 R	4	146	HSP7C G372 R	2	55	NEUG C9 RO3	1	37	PCDH1 D944 R	1	16
GFAP E391 R	15	600	NFM E599 R	4	129	*GATB E42 R	2	53	DZI1L Q132 R	1	35	AT5G1 D62 R	1	15
MYH9 D1136 R	15	495	ACTC E318 R	4	119	NPM D179 R	2	53	RLA1 D19 R	1	34	F205A E1023 R	1	15
*CH60 E107 R	15	452	NFH E469 R	4	88	EFHD2 Q67 R_deami	2	50	PTBP1 G306 R	1	33	ILF2 D347 R	1	15
MLRV E22 R	14	266	NFL E507 R	4	88	TBB4A D67 R	2	50	TYB4 D6 R	1	32	MAP1B Q907 R_deami	1	14
SAP G407 R	13	172	NHRF1 E121 R	4	77	HBA N79 R_deami	2	48	CH60 G43 R	1	31	MTDC E36 R	1	14
NFM E470 R	12	354	RD23B E136 R	4	65	SCMC1 Q16 R_deami	2	45	HNRPK D279 R	1	31	MYH7 D1008 R	1	14
TBA1B D218 R	12	132	SCG1 D88 R	4	51	TFF1 E25 R	2	42	ROA1 E93 R	1	31	GLYM Q27 R_deami	1	13
AP180 N767 R_deami	12	77	EF1D E27 R	4	21	CWC15 Q45 R_deami	2	40	SPTB2 E1469 R	1	31	SF3B1 D93 R	1	13
NFM E496 R	10	770	HNRPK N306 R	3	146	AKA12 E952 R	2	37	SRRT D866 R	1	31	MBP D279 R	1	12
CLUS D23 R	10	179	MBP N226 R	3	142	MYH7 E73 R	2	35	FUBP1 N369 R_deami	1	30	NECP1 C162 RO3	1	11
NEUM E89 R	10	152	APOA1 D25 R	3	140	RL1D1 D319 R	2	32	COX17 D 6 R	1	29	RM12 C45 RO2	1	11
ATIF1 E65 R	10	148	H15 E 8 R	3	119	BACH C36 RO3	2	31	NUCL E628 R	1	29	ATIF1 D28 R	1	10
DREB D634 R	9	460	CD99 D23 R	3	115	PGCB E412 R	2	29	TBA1B D251 R	1	29	MAP1A E1580 R	1	10
MYH7 E1821 R	9	232	C99L2 D26 R	3	114	UTP18 E374 R	2	26	E2F7 D10 R	1	28	MYPT2 D816 R	1	9
ENPL D22 R	8	555	LDB3 N197 R_deami	3	112	AAK1 Q374 R_deami	2	23	IF4H D196 R	1	28	TNNT2 D108 R	1	9
NCAN E23 R	8	177	HNRPU E259 R	3	110	MLRV N154 R_deami	2	23	AQP4 E279 R	1	27	KAD4 G82 R	1	8
BCLF1 E463 R	8	160	HNRPC D123 R	3	109	TMM11 D163 R	2	20	ATP5H D129 R	1	27	NFL E509 R	1	7
RLA1 N62 R_deami	8	159	RLA1 D18 R	3	109	ACTC C259 RO2	2	17	CALX E275 R	1	27	PDLI5 E297 R	1	7
TAU N644 R_deami	8	136	PELP1 E1008 R	3	104	CDY1 D285 R	2	17	PSMD6 L84 R	1	27	DPEP3 D150 R	1	5
ACTC D53 R	7	270	CAVN2 Q161 R_deami	3	95	NFM E763 R	2	17	AKA12 E418 R	1	26	HMSD Q43 R_deami	1	5
ACTB E107 R	7	179	G3P D315 R	3	92	ANK2 E1760 R	2	15	COX41 D43 R	1	26	PTGDS E25 R	1	3
ACTB D51 R	6	257	MESD E35 R	3	92	PACN1 N330 R_deami	2	8	MPPB Q46 R_deami	1	26	PSD12 E348 R	1	1
MCFD2 E27 R	6	231	NFM E455 R	3	87	PGCB E26 R	2	7	PDLI5 I574 R	1	26	SCN1A C1588 RO2	1	1

Notes. R, arginylation (color code: black). RO3, Cys tri-oxidation and arginylation (color code: red). RO2, Cys di-oxidation and arginylation (color code: yellow). R_deami, N/Q arginylation after deamidation (color code: blue). *, validated by standard peptides. n1, the total number of fractions. n2, the total number of paired MS1 scans.

Arginylation profiling in human proteomes. Using this established platform, we profiled 14 human samples after cell/tissue lysis and ATE1 assay *ex vivo*, a key to installing isotopic Arg post-translationally under ribosome-free conditions to bypass translational Arg incorporation and bias. A total of 1876 isotopically labeled pairs belonging to 235 unique arginylation sites on 165 proteins were identified

(**Fig. 3b**, **Table 1**, **Supplementary Dataset 3**, and **Fig. S19-20**). As a negative control, the HEK293T sample without the addition of ATE1 didn't show any arginylation site (RAW data is available). Only 7 arginylation sites (ACTC C259 RO2, BI1 N9 R_deami, HBA N79 R_deami, HMSD Q43 R_deami, PSD12 E348 R, S10AE C74 RO3, and SCN1A C1588 RO2) were assigned as sidechain arginylation accounting for 3% of all unique sites, while the rest 97% of sites were identified as *N*-term arginylation. Similar to results from HEK293T peptides, most of the peptide pairs are detected at charge states 2 (45.0%) and 3 (49.9%), with small portions in charges 4 (4.9%) and 5 (0.2%) (**Fig. S21**). As cross-validation, CALR E18 and PDIA D18 are among the top high-frequency sites with 185 and 135 detections (MS1 pairs) respectively, consistent with the literature on their confirmed E18[10] and D18[24] arginylation. The number of unique arginylation sites from each sample is listed in **Fig. 3c** which shows an average of 29 sites/sample could be identified. The biological samples cover induced pluripotent stem cells (iPSC), cardiac cells derived from the iPSC, commonly cultured cell lines, and clinical patient tissues, demonstrating the potential applicability of this method to a wide range of other human proteomes. The H/L ratios (threshold used: $0.1 \leq$ ratio ≤ 10) of MS1 pairs are distributed symmetrically and perfectly centered at the hypothetical ratio of **1.0** (**Fig. 3d**). Endogenous Arg^0 at µM level may interfere with the Arg^{10} labeling, however, the final concentration of Arg^{10} is at 2 mM thus Arg^0 interference on final H/L ratios might be minimal. Since Arg^0 and Arg^{10} labeling by ATE1 was carried out separately on complex whole proteomes, deviations of H/L ratios (range: 0.3-8.5) from the hypothetical ratio of 1.0 were expected which might be introduced from the multi-step preparation procedures including ATE1 arginylation on substrates in the whole proteome, H/L mixing, digestion, and peptide fractionation (**Fig. 1b**). The doublet ratios were also compared with the quartet and sextet profiles (**Fig. S22**). The numbers of MS1 scans with the doublet, quartet, and sextet features for generating respective H/L ratios are provided, quartet and sextet pairs showed lower numbers of qualified MS1 scans than doublets based on their trendlines (**Fig. S23**). The sequence logo (**Fig. 3e, Fig. S24**) and arginylation motif (**Fig. S25**) from all unique peptides suggested that arginylation favors *N*-term D and E residues significantly, while subsequent nonpolar residues (A, L, V, and P) and acidic residues (D and E) may facilitate the peptide *N*-term arginylation. Interestingly, all four types of arginylation based on different modification masses (**Fig. S26**) have been detected including arginylation (R, 84.7%), Cys tri-oxidation arginylation (RO3, 3.4%), N/Q deamidation arginylation (R_deami, 10.6%), and Cys di-oxidation arginylation (RO2, 1.3%), (**Fig. 3f**). Comparison of arginylation sites between iPSC, cancerous cells, and patient samples showed only 32 (13+11+8) shared sites, while the remaining majority of sites are unique to their respective cell/sample groups (**Fig. S27**). Comparisons between individual samples are listed in **Fig. S28** (cardiac), **Fig. S29** (brains), and **Fig. S30** (cancerous), indicating unique and shared arginylation sites from individual samples. Replication ($n = 3$) analysis of 4 samples showed that most arginylation sites in each sample were detected in replicates (**Fig. S31**), indicating the repeatability of the profiling method. Among 165 proteins identified, 162 protein targets are involved in many key biological processes (*e.g.,* cellular process, biological regulation, and cellular metabolism) according to the PANTHER classification (**Fig. 3g**).

Validation of representative arginylation sites using standard peptides. We aimed to validate representative sites using synthetic peptides (*e.g.*, REPAVYFK for CALR E18 peptide) whose MS behaviors (MS1 spectra in **Fig. 4a** and MS2 spectra in **Fig. S32**) matched those in our MS data. The chosen sites are a mixture of known (CALR and PDIA) and unknown, and they represent high-frequency (CALR, PDIA, SSBP, and RM12), mid-frequency (ERO1A and CH60), and low-frequency (GATB) sites in our data from biological samples (**Table 1**). Such a validation added extra confidence to the Arg10 arginylation which is an unnatural process and thus can be considered an internal validation of our workflow. For example, H/L ratio = 1.0 for RM12 C45 arginylation (**Fig. 4b**) eliminated the possibility of tryptic missed cleavage between Arg44 and tri-oxidized Cys45, standard peptide confirmed this minor type of arginylation with co-presence of Cys trioxidation[1] (**Fig. S32**). It is noteworthy that the low frequency of a site doesn't necessarily mean low confidence since most of the reported sites are based on many isotopic co-eluting MS1 scans (**Fig. S23** and **Supplementary Dataset 3**) and paired MS2 spectra (**Supplementary Dataset 4**). To further confirm that these representative sites possess arginylation at the *N*-termini but not on the side chain of *N*-terminal residues, we purchased peptides containing arginylation on the side chain [*e.g.,* E(R)PAVYFK for CALR E18 peptide]. Analyzed by LCMS, side chain arginylation peptides may behave similarly (*e.g.*, CALR E18 peptide) or differently (*e.g.*, SSBP E17 peptide) with *N*-terminal arginylation peptides (**Fig. S33**). Interestingly, arginine residue on the side chain produced a signature ion at *m/z* 175.1190 when the peptide didn't end with a *C*-terminal R residue [*e.g.*, E(R)PAVYFK]. When *C*-terminal R residue was present on side

chain arginylation peptides [*e.g.*, E(R)EQPPETAAQR], both R residues could produce the same ion at *m/z*
175.1190 with a higher intensity than that from *N*-terminal arginylation peptides (*e.g.*, REEQPPETAAQR)
(**Fig. S33**). By adding the side chain arginylation peptides after ATE1 assay, the *N*-terminal arginylation was
further confirmed for standard peptide (DIAALVHSSGNIeS-NH$_2$) (**Fig. S34**), CALR pure protein (**Fig. S35**)
and CALR in HEK293T cells (**Fig. S36**).

Validation of representative arginylation sites using in-bacteria arginylation assay. We then
selected two novel sites for validating their arginylation in an ATE1-protein co-expression system in *E. coli*
reported in a recent study[35]. The two proteins are 1) a chaperone ERO1A in the endoplasmic reticulum
(ER), being similar to CALR and PDIA; and 2) SSBP being distinct from ER proteins in molecular size (17
kDa), localization (mitochondria) and function (single-stranded DNA-binding). Briefly, a protein or peptide
containing an arginylation site is fused to ubiquitin and co-expressed with human ATE1 and Ulp1 protease
in *E. coli*. The expressed protein is cleaved by protease and exposes the *N*-terminal arginylation site for
ATE1 arginylation. This system allows for in-bacteria validation of targeted arginylation. The results showed
that both ERO1A E24 and SSBP 17E were arginylated (**Fig. 4c** and **Fig. S37-38**). As a bonus, we also
detected a further leucylation on top of arginylation (**Fig. S37-38**), a unique process to modify *N*-term Arg
due to the presence of the *E. coli* leucyl/phenylalanyl-tRNA-protein transferase (LFTR)[36]. Similarly, we also
tested the in-bacteria arginylation of ERO1A and SSBP peptides, whose respective *N*-term E24 and 17E
were also arginylated in the co-expression system (**Fig. S39**). The arginylation and leucylation were also
observed by top-down proteomics (**Fig. S40-41**). The results concluded that ERO1A E24 and SSBP 17E
are arginylation target sites of ATE1.

Figure 4. Validation of representative arginylation sites. a, synthetic peptide validation of 7 arginylation
sites. **b,** representative MS2 spectra indicating RM12 arginylation on Cys45 with tri-oxidation (C45 RO3).
c, open E24 in ERO1A and E17 in SSBP are arginylated in an in-bacteria arginylation system. Protein and

ATE1 are co-expressed in E. coli for in-bacteria arginylation. The protein was purified and digested for
proteomics analysis. **d,** arginylation dependency of CALR, ERO1A, and SSBP sites on endogenous ATE1.
Anti-ATE1 was used to detect the expression of endogenous ATE1. β-tubulin was used as a loading control.
Anti-FLAG antibody was used to detect expressions of CALR, ERO1A, and SSBP proteins. Extracted ion
chromatograms (EICs) of arginylated peptide in each protein in WT and KO cells after pulldown and
proteomics are provided. **e,** relative arginylation levels of CALR, ERO1A, and SSBP sites after co-
overexpression of ATE1. Protein was purified by antibody pulldown experiment followed by proteomics
analysis. Peak areas of arginylated peptides were normalized to the sample with the highest signal and
relative ratios are displayed. Different amounts (0, 0.5, and 1 µg) of ATE1 plasmids were used for
transfection. β-tubulin was used as a loading control. Anti-FLAG antibody was used to detect expressions
of CALR, ERO1A, and SSBP proteins. std, standard peptide. KO, ATE1 KO. R, arginylation.

Validation of representative arginylation sites in cells. The unbiased nature of the profiling
method is owing to the co-elution strategy of isotopic Arg10/Arg0 peptides, however, the revealed sites are
based on an *ex vivo* ATE1 assay and thus do not necessarily conclude their occurrences *in vivo* (in cells)
where ATE1 is active. To confirm endogenous arginylation activity on the identified targets in cells, we
overexpressed the wild-type ERO1A and SSBP in HEK293T cells using CALR as a positive control. Due to
the low abundance of endogenous arginylation, arginylated peptides were not detectable in peptide
mixtures obtained from whole proteomes if overexpressed proteins were usually not purified by
immunoprecipitation. For example, the MS2 spectrum of arginylated peptide for the SSBP E17 site was
only detected once from a pulldown sample (**Fig. S42**). Nevertheless, the arginylated peptides
corresponding to all three sites (CALR E18, ERO1A E24 and SSBP E17) were identified by bottom-up
proteomics after immunoprecipitation of overexpressed proteins (**Fig. 4d** and **Fig. S42**). As a comparison,
we also performed the same experiment in HEK293T (ATE1 KO) cells (**Fig. 4d**) or on E-to-V mutants of
ERO1A E24 and SSBP E17 (**Fig. S43**), which did not show any arginylation on those sites indicating
arginylation dependency on ATE1 and arginylation selectivity on E residue. To further test their dose-
dependency on ATE1, we then co-overexpressed the ATE1 and protein substrates in HEK293T (ATE1 KO)
cells. The data show the arginylation levels of the tested sites are dose-dependent on the ATE1 expression
(**Fig. 4e**). Such dose-dependency was also observed when co-overexpressing CALR and ATE1 in WT
HEK293T cells (preliminarily estimated arginylation efficiencies between 0.025%-0.111%) (**Fig. S44**). Both
ERO1A[37] and SSBP[38] contain cleavage peptides before the arginylation sites (E24 and 17E respectively),
similar to that of CALR. Taking advantage of this feature, we further prepared 24R-ERO1A and 17R-SSBP
plasmids by inserting an R residue before E into the full-length plasmids as positive transfection controls,
the spectra of their arginylated peptides (**Fig. S45**) resulted from overexpression, affinity purification and
proteomics further confirmed the arginylation of their wild-type counterparts.

We have additionally validated the endogenous arginylation in HEK293T cells of two more sites
(proteins): A1AT E25 arginylation (**Fig. S46**) and Tau N644 arginylation after deamidation (**Fig. S47**). Similar
to CALR, ERO1A and SSBP, A1AT has a signal peptide[39] before E25 thus it is not surprising to observe its
E25 arginylation by endogenous ATE1. Tau N644 has been arginylated after deamidation in our data, which
has never been observed previously. We also confirmed the arginylation of this site using overexpression
of wild-type Tau plasmid in HEK293T cells (**Fig. S47**). We have additionally confirmed the arginylation of
the Tau E3 site (**Fig. S47**), which was proposed previously as an arginylation site of ATE1, but not validated
for its arginylation[40]. A previous study showed that purified Tau did not show detectable arginylation by ATE1
in vitro[16], possibly due to the lack of exposed arginylation sites (*e.g.*, E3 and N644 deamidation). Different
from Tau E3 as a calpains cleavage product[40], the mechanism of generating an open N644 followed by
deamidation for arginylation remains to be elucidated.

Biological functions associated with ERO1A arginylation. To investigate whether arginylated
ERO1A resides in ER, we did imaging on overexpressed ERO1A and 24R-ERO1A in HEK293T ATE1 KO
cells. Both proteins co-localized with ER marker and PDI (**Fig. 5a** and **Fig. S48**), suggesting that 24R-
ERO1A was translocated into ER after signaling peptide AA1-23 cleavage. As a positive control, we showed
that CALR-halo and its arginylation form co-localized with a GFP-tagged ER marker (**Fig. S49**). Similar to
CALR and PDIA, ERO1A is translocated into ER after signal peptide cleavage. A previous study reported
the formation of the ERO1A-PDI complex in regulating cellular function[41], our pulldown data showed that
24R-ERO1A interacted with PDI to a comparable extent with and without PDI co-overexpression (**Fig. S50-**

51). We also showed that arginylation of ERO1A maintains its enzymatic activity on oxidative protein folding
compared with WT ERO1A as indicated by the disappearance of the reduced JcM band (**Fig. S52**).

To investigate the arginylation effect on ERO1A stability and degradation using HEK293T ATE1 KO
cells, we compared the degradation patterns of ubiquitin-ERO1A fusion proteins with and without
arginylation (**Fig. S52**) according to similar experiments performed on CALR[42]. Briefly, ubiquitin is cleaved
by de-ubiquitinase after protein synthesis to expose E24 and R24 residues in ERO1A (no arginylation on
E24) and 24R-ERO1A (~100% arginylation on E24) respectively. Proteasome inhibitor MG132 treatments
after transient transfection increased the levels of ERO1A and 24R-ERO1A, suggesting that both are
involved in proteasome-mediated degradation. Similar dose dependence on MG132 was observed in co-
treatments when the ribosomal translation of both species was inhibited by cycloheximide (CHX). In addition,
24R-ERO1A showed slightly higher levels after MG132-CHX co-treatment, indicating better stability than
ERO1A against proteasomal degradation (**Fig. S52**). To take a closer look at their stabilities, we did CHX
chase experiments[42] where cells were treated with CHX after a 2-day transient transfection. The results
showed that 24R-ERO1A was less prone to degradation than ERO1A with clear degradation after CHX
treatment (**Fig. 5b, Fig. S53**). Such behavioral differences have been previously observed in cytosolic
CALR[42], suggesting that arginylation of ERO1A may also have a stabilization effect against proteasomal
degradation.

Figure 5. Biological functions of representative arginylation sites. a, imaging of ERO1A and 24R-
ERO1A compared with ER marker and co-overexpressed PDI. **b**, expression levels of cytosolic ERO1A
species in CHX chase experiment. **c,** imaging of SSBP and 17R-SSBP compared with mitochondria protein
COX4. **d**, bioenergetic profiles of HEK293T cells after SSBP and 17R-SSBP transfection measured by
Seahorse XF24. Mitochondria respiration was analyzed with basal respiration, ATP production, proton leak,
maximal respiration, spare capacity, and non-mitochondria respiration. The average values of 3 Seahorse
measurements were used for comparison.

Biological functions associated with SSBP arginylation. SSBP matures by translocation into
mitochondria after cleavage of its transit peptide (AA 1-16)[38]. We did imaging on SSBP and 17R-SSBP to
investigate whether arginylated SSBP resides in the mitochondria. Both species co-localized with
mitochondria protein COX4 (**Fig. 5c** and **Fig. S54**), suggesting that 17R-SSBP was translocated into
mitochondria after cleavage of its transit peptide AA1-16. To assess the cytosolic degradation of SSBP with
and without arginylation in the *N*-degron pathway, we evaluated its expression levels using a ubiquitin
cleavage plasmid system for transfection. As a result, we did not observe an obvious response toward
MG132 treatment (proteasome inhibitor) for both SSBP species (**Fig. S55**). CHX (protein synthesis inhibitor)
treatment reduced the expression levels of both SSBP and 17R-SSBP compared with cells without CHX
treatment, indicating that ribosomal synthesis was contributing to the cellular expression levels of both

species. 17R-SSBP expression levels were higher than WT SSBP with and without CHX treatment,
indicating a slower turnover of 17R-SSBP than SSBP. The results suggest that the major role of SSBP
arginylation might not be degradation (**Fig. S55**).

We then turned our attention to mitochondrial SSBP. SSBP KO in mouse models is embryonic
lethal whereas conditional KO in the heart results in cardiomyopathy and reduced life span (~18 weeks)
accompanied by heart respiratory chain deficiency[43]. It is noteworthy that ATE1 KO is also embryonic lethal
with heart defects[1], being similar to SSBP KO. Thus arginylation of SSBP may regulate cells through
mitochondria functions. Considering the importance of mitochondria in energy generation and heart
functions, we then investigated the functional differences between SSBP and 17R-SSBP on cellular
bioenergetics (**Fig. 5d**) based on a previous study on SSBP mutants[44]. Toward this, the Seahorse XF24
Cell Mito Stress Test was performed to measure the arginylation effects of SSBP on mitochondrial
respiration before and after sequential addition of oligomycin (oligo), carbonyl cyanide-4-
(trifluoromethoxy)phenylhydrazone (FCCP) and rotenone/antimycin A (Rot/A). Analysis of cellular oxygen
consumption rate (OCR) showed that HEK293T cells overexpressing 17R-SSBP displayed comparable
basal respiration, ATP production, and non-mitochondria respiration than WT SSBP (**Fig. 5d**). Interestingly,
17R-SSBP resulted in significantly higher maximal respiration (p = 0.0104) and spare capacity (p = 0.0076)
than cells overexpressing WT SSBP (**Fig. 5d**, **Fig. S56**, **Supplementary Dataset 5**). The relative ratios of
maximal respiration and spare capacity in 17R-SSBP are 1.6 and 8.5 compared with WT SSBP (**Fig. S56**),
indicating improved mitochondria respiration potential after SSBP arginylation. 17V-SSBP did not display
these effects on mitochondria when compared with WT SSBP (**Supplementary Dataset 6**).

Arginylation profiling in mouse tissues. Our profiling platform has been working in human cells
and tissues, to test its general applicability we next applied it to mouse tissues. We acknowledge that it is
not ideal to use human ATE1 to arginylate mouse proteomes, although such a cross-species approach has
been previously used[16]. Results from three tissues (lung, heart, and brain) showed that the platform is
applicable to arginylation discovery from mouse proteome (**Supplementary Dataset 7**). A total of 14 sites
from 69 MS1 pairs were identified (**Fig. S57**). Interestingly, CALR D18 and A1AT1 E25 are the two shared
arginylation sites across three tissues. We noticed that the mouse A1AT1 E25 site was identified by a
different peptide sequence (EDVQETDTSQK) (**Fig. S58**) from the human A1AT E25 site (EDPQGDAAQK)
(**Fig. S46**). In addition, human CALR E18 and mouse CALR D18 sites were also identified from species-
dependent sequences (EPAVYFK and DPAIYFK, respectively) (**Fig. S59**), indicating that mouse and human
sites can be used to cross-validate each other when overlapped.

Establishing a public database for arginylation sites and ATE1 substrates. We envision that
the arginylation sites discovered from the various samples will greatly advance the arginylation and *N*-
degron fields. The fact that our high-frequency hits (CALR, PDIA, and TAU *et al*)[25,40] are among the most
intensively studied arginylation proteins instilled confidence to follow the biology of new sites. This
represents a major advantage of our innovative method compared to existing profiling methods[21,28].
Meanwhile, we understand that we alone cannot follow up on all the targets/sites for biological functions, a
publicly accessible database/website (www.arginylation.com, **Fig. 6**) was thus formulated to allow everyone
to examine our data and follow the biology of their arginylation targets of interest. The isotopic MS1 and
MS2 scans discussed in this work from standard peptides, whole-proteome peptide mixtures, pure proteins,
human proteomes, and mouse proteomes are included on the website. The source code for data
visualization is publicly available to download at https://github.com/ChenfengZhao/Arginylation. The
website was designed to allow the following two capabilities, 1) users can search protein names, accession
numbers, sample types, species, and peptide sequences of their interest; 2) users can manually select
species, sample types, proteins, peptides, fractions of a sample, and charge states of the peptides from
our indexed database (**Fig. 6**). As a result, all isotopic MS1 scans, H/L ratio summaries, and H/L MS2 scans
with default settings are displayed on the website. To allow users to change our default parameters, we
enabled changes in *m/z* errors and MS2 ion types. The most important feature of our website is that users
can download our indexed data (MS1 and MS2 spectra in Excel format) based on species (*e.g.*, human
and mouse) or samples (*e.g.*, iPSC or patient brain).

Figure 6. Design and flowchart of the arginylation database and its website. Arginylation data are from peptides, whole-proteome peptides, proteins, human proteomes, and mouse proteomes. Visualization includes isotopic ratios and annotated isotopic MS1 and MS2 spectra. All indexed MS1 and MS2 scans are accessible to the public to download. A dashed arrow means a one-to-multiple inclusion relationship.

Discussion

ATE1 is the only enzyme known to catalyze arginylation in mammalian systems (*e.g.*, human and mouse), it has two isoforms in humans (ATE1-1 and ATE1-2) and four isoforms in mice (ATE1-1, ATE1-2, ATE1-3, ATE1-4). (**Fig. S60**). The expression levels and enzymatic functions vary among different isoforms[24,45]. While our work mostly focuses on establishing a general proteomic platform using human ATE1-1, using other isoforms (ATE1-2 for human proteomes, and mouse isoforms for mouse proteomes) for arginylation discovery is a necessary next step to cover the ATE1 substrate scope comprehensively. We preliminarily compared human ATE1-1 and ATE1-2 for arginylation profiling in two samples (patient brain and HeLa cells). ATE1-1 and ATE1-2 sites showed unique and shared sites, more arginylation sites were revealed by ATE1-1 than ATE1-2 for both samples (**Fig. S61-62**), indicating that ATE1-1 may have a slightly broader substrate scope for such an *ex vivo* arginylation assay. In addition, previous studies showed that the expression of ATE1 isoforms is tissue-specific and cell-dependent[24,45], thus the expression levels might be a significant consideration when choosing ATE1 isoforms for arginylation profiling experiments.

It was initially believed that arginylation occurred only on exposed *N*-terminal Asp, Glu and Cys residues resulting in protein degradation by the ubiquitin-dependent *N*-degron pathway[46]. Later studies identified proteins (*e.g.*, α-synuclein[16]) potentially modified by arginylation on their sidechain Asp and Glu residues[27], resulting from a different but minor catalysis mechanism of ATE1. Our arginylation sites were identified using a search algorithm considering both *N*-term and sidechain arginylation situations (**Fig. S26**). The resulting sites from the whole-proteome peptide mixture (**Fig. 2**) and proteomes (**Fig. 3**) are in agreement with the current understanding that ATE1 predominantly arginylate protein/peptide *N*-term C (oxidized), D, and E residues, while other residues and sidechains can be arginylated less preferably. Since the ATE1 assay was performed on proteomes, we believe most sites were arginylated as protein *N*-term (**Fig. 3** and **Supplementary Dataset 3**). A preliminary analysis of the P1 residues (protease cleavage site) in arginylated peptides showed potential protease cleavage after each of the 20 amino acid residues (**Fig. S63** and **Supplementary Dataset 3**). Cleavages at K and R sites accounting for 18.6% and 18.2% respectively might be produced from endopeptidases such as trypsin-like proteases, in contrast, how cleavages at A (13.3%) happened remains unanswered. While some sites (*e.g.*, CALR E18, ERO1A E24, and SSBP E18) were known to be protein *N*-term based on the current knowledge on cleavages of signaling peptides and transit peptides, many sites (*e.g.*, TAU N644) are left unexplained as to how the parent proteins were cleaved to expose the *N*-term sites for ATE1 arginylation. Previous studies have offered some insights into protease cleavage (*e.g.*, calpains for TAU E3[40] and caspase 3 for CDC6 D101[25]) for *N*-term

arginylation, however, the cleavage products of more than 600 proteases remain elusive. Great efforts are
needed to understand the interplay between proteolytic cleavage and arginylation.

Proteomic discovery of endogenously low-level arginylation from cells or tissues is challenging[21,28].
This is largely due to 1) the mass difference (+156 Da) introduced by arginylation being the same as the
normal arginyl residue from a protein sequence, making label-free proteomics biased to predefined
arginylation sites by nature; and 2) efficient enrichment (purification) methods are lacking. The fact that
arginyl-tRNA is the same source of arginylation and ribosomal translation makes unbiased endogenous
arginylation discovery almost impossible since one can't exclude ribosomal incorporation when cells are
alive[13]. Our initial effort using Arg10 and Arg0 for pulse labeling of arginylation in HEK293T cells yielded a
putative list of sites (**Fig. S63**). Additional efforts using cycloheximide to stop protein synthesis, MG132 to
accumulate arginylation, and peptide fractionation to boost peptide IDs did not seem to increase our
confidence. Arginylation identification from a standard proteome-wide peptide mixture is not ideal. Our
efforts on *in vivo* strategy also suggested that endogenous arginylation should have minimal contribution to
the light (Arg[0]) labeling in our *ex vivo* isotopic approach. Previous studies tried to use anti-arginylation
antibodies to reduce sample complexity and enrich endogenous arginylation[21,28], however, these efforts by
design were biased toward non-degradative ATE1 targets and high-abundance proteins. Sites from those
studies showed nearly no overlap with those previously characterized by the Varshavsky group as
degradation-inducing arginylation substrates[13]. We thus turned our attention to an *ex vivo* ATE1 enzyme
approach using cell lysates, which might be less physiologically relevant than those targeting endogenous
arginylation. In addition, arginylated proteins by *ex vivo* assay using lysates were not localized correctly in
cell compartments, thus may be false positive hits (not arginylated in cells). However, due to its unbiased
nature, this method represents a superior strategy for arginylation discovery.

This profiling method relies on deep peptide fractionation followed by LC-MS/MS analysis to
discover the low-abundance arginylated peptides, and intact peptides that are not of interest (accounting
for >99.99%) are obvious interfering signals in proteomics data. The reduced signal complexity of each
peptide fraction after fractionation is key to identifying arginylated pairs from proteome-wide samples. The
additional charge in ionization (**Fig. S9**) and polarity in chromatography of peptides (**Fig. 1c**) resulting from
arginylation may help validate peptide identity using LC-centric prediction tools such as DeepLC
(**Supplementary Dataset 8**)[47] and Chronologer[48]. Without enrichment, the sites revealed by our method
could be dependent on the expression levels of their parent proteins. We envisioned enrichment
approaches are essential to achieve improved sensitivity and efficiency in arginylation site detection. Since
ATE1 mainly installs Arg to *N*-termini of substrates, some technologies targeting *N*-term peptides might be
helpful for arginylation enrichment. Examples are TAILS (terminal amine isotopic labeling of substrates)
technology[49] and iNrich (integrated *N*-terminal peptide enrichment) method[50] which are currently under our
investigation as future directions. The anti-arginylation antibody enrichment is also under consideration,
one potential advantage of such an antibody enrichment strategy is that antibodies can be used for
enriching *in vivo* arginylated proteins. However, antibody enrichment may require scaling up the ATE1
reaction to milligram proteomes from the current 20-µg scale. Taken together, our *ex vivo* ATE1 approach
provides a robust, unbiased way to map arginylation at the proteome-wide scale, setting the stage for more
systematic and comprehensive analyses. Looking ahead, combining this strategy with targeted enrichment
techniques and in vivo validation should significantly advance our understanding of arginylation's functional
roles in diverse biological contexts.

Methods

The methods section is provided in the Supplementary Information with details available.

Acknowledgments

Z.L. thanks an NIH R21 (CA292191) and R01 (HL177113), a WUSM ICTS JIT grant (JIT1181), and support
from the Research Education Component through an NIA P30 grant (AG066444). B.A.G. thanks NIH
(HL177113, NS111997 and HD106051) and NSF (CHE 2127882) grants. D.L. thanks an NCI R21 grant
(CA286307) and the support of the Mays Cancer Center Early Career Pilot Award from CCSG (P30
CA054174). Y.Z. thanks an NIH R35 grant (GM150678). M.J.G. thanks partial support from NIH (R01
HL141086), Children's Discovery Institute of Washington University and St. Louis Children's Hospital (PM-
LI-2019-829), and the American Heart Association (970198). A. K. thanks NIH R35 (GM122505) and R01

(NS102435) grants. The authors thank Professor Natalie Niemi (WashU) for guidance on the Seahorse
assay, Professor Xia Liu and Professor Guangyong Peng (WashU) for providing mouse tissues, and
Washington University Diabetes Research Center supported by NIH Grant 5P30 DK020579.

Author contributions

Z.L. and B.A.G. conceived the project. Z.L., Y.X., and J.G. performed proof-of-concept experiments. Z.L.,
Y.X., J.G., X.Liu, E.Z., B.B.P., D.H.R., R.M.S., F.N.V., R.K., G.P.D., B.M., X.Lan, D.F., and L.G. performed
experiments. Z.L., Y.X., J.G., X.Liu, E.Z., B.B.P., D.H.R., R.K., B.M., X.Lan, D.F., L.G., Y.Z., K.J.L., M.J.G.,
D.L., A.K. and B.A.G. analyzed data. Z.L. wrote the draft. C.Z. wrote Python codes and established the
website, and X.H. wrote R codes and ArginylomePlot software. K.J.L., B.M., and A.K. provided patient
tissues. All authors wrote, revised, and approved the manuscript.

Competing interests

557 Z.L., D.L., and B.A.G. are co-founders of LasNova Therapeutics, LLC.

Data availability

Arginylation site data are publicly available to view and download from www.arginylation.com (the website
will be accessible during peer review and after publication, it will be inaccessible during revision).

RAW data are uploaded to MassIVE under dataset numbers MSV000097195, MSV000097197, and
MSV000097196. The data index is listed in **Supplementary Dataset 9**.

Log in information for reviewer: MSV000097195_reviewer, MSV000097197_reviewer, and
MSV000097196_reviewer, password: xxx.

Data supporting the findings in this study are available within the paper. Any associated data are available
from the corresponding authors upon request.

Code availability

Python codes for website construction are available to download from GitHub (link:
https://github.com/ChenfengZhao/Arginylation)

R codes for ArginylomePlot software are available to download from GitHub (link:
https://github.com/BeckyHan/Garcia-Lab/tree/main/ArginylomePlot).

The arginylation website is publicly accessible (link: www.arginylation.com) for data visualization and
download.

References

1　Kwon, Y. T., Kashina, A. S., Davydov, I. V., Hu, R. G., An, J. Y., Seo, J. W., Du, F. &
Varshavsky, A. An essential role of N-terminal arginylation in cardiovascular development.
Science **297**, 96-99, (2002). PMID: 12098698

2　Lian, L., Suzuki, A., Hayes, V., Saha, S., Han, X., Xu, T., Yates, J. R., 3rd, Poncz, M.,
Kashina, A. & Abrams, C. S. Loss of ATE1-mediated arginylation leads to impaired platelet
myosin phosphorylation, clot retraction, and in vivo thrombosis formation. *Haematologica*
99, 554-560, (2014). PMID: 24293517

3　Kurosaka, S., Leu, N. A., Pavlov, I., Han, X., Ribeiro, P. A., Xu, T., Bunte, R., Saha, S.,
Wang, J., Cornachione, A., Mai, W., Yates, J. R., 3rd, Rassier, D. E. & Kashina, A.
Arginylation regulates myofibrils to maintain heart function and prevent dilated
cardiomyopathy. *J Mol Cell Cardiol* **53**, 333-341, (2012). PMID: 22626847

4　Singh, K., Gupta, A., Sarkar, A., Gupta, I., Rana, S., Sarkar, S. & Khan, S.
Arginyltransferase knockdown attenuates cardiac hypertrophy and fibrosis through TAK1-
JNK1/2 pathway. *Sci Rep* **10**, 598, (2020). PMID: 31953451

5　Kurosaka, S., Leu, N. A., Zhang, F., Bunte, R., Saha, S., Wang, J., Guo, C., He, W. &
Kashina, A. Arginylation-dependent neural crest cell migration is essential for mouse
development. *PLoS Genet* **6**, e1000878, (2010). PMID: 20300656

6　Leite, F. S., Minozzo, F. C., Kalganov, A., Cornachione, A. S., Cheng, Y. S., Leu, N. A.,
Han, X., Saripalli, C., Yates, J. R., 3rd, Granzier, H., Kashina, A. S. & Rassier, D. E.
Reduced passive force in skeletal muscles lacking protein arginylation. *Am J Physiol Cell*
Physiol **310**, C127-135, (2016). PMID: 26511365

7　Karakozova, M., Kozak, M., Wong, C. C., Bailey, A. O., Yates, J. R., Mogilner, A., Zebroski,
H. & Kashina, A. Arginylation of beta-actin regulates actin cytoskeleton and cell motility.
Science **313**, 192-196, (2006). PMID: 16794040

8　Zhang, F., Saha, S., Shabalina, S. A. & Kashina, A. Differential arginylation of actin
isoforms is regulated by coding sequence-dependent degradation. *Science* **329**, 1534-
1537, (2010). PMID: 20847274

9　Carpio, M. A., Lopez Sambrooks, C., Durand, E. S. & Hallak, M. E. The arginylation-
dependent association of calreticulin with stress granules is regulated by calcium.
Biochem J **429**, 63-72, (2010). PMID: 20423325

10　Decca, M. B., Carpio, M. A., Bosc, C., Galiano, M. R., Job, D., Andrieux, A. & Hallak, M.
E. Post-translational arginylation of calreticulin: a new isospecies of calreticulin component
of stress granules. *J Biol Chem* **282**, 8237-8245, (2007). PMID: 17197444

11　Hu, R. G., Sheng, J., Qi, X., Xu, Z., Takahashi, T. T. & Varshavsky, A. The N-end rule
pathway as a nitric oxide sensor controlling the levels of multiple regulators. *Nature* **437**,
981-986, (2005). PMID: 16222293

12　Lee, M. J., Tasaki, T., Moroi, K., An, J. Y., Kimura, S., Davydov, I. V. & Kwon, Y. T. RGS4
and RGS5 are in vivo substrates of the N-end rule pathway. *Proc Natl Acad Sci U S A* **102**,
15030-15035, (2005). PMID: 16217033

13　Varshavsky, A. N-degron and C-degron pathways of protein degradation. *Proc Natl Acad*
Sci U S A **116**, 358-366, (2019). PMID: 30622213

14　Kim, B. H., Kim, M. K., Oh, S. J., Nguyen, K. T., Kim, J. H., Varshavsky, A., Hwang, C. S.
& Song, H. K. Crystal structure of the Ate1 arginyl-tRNA-protein transferase and
arginylation of N-degron substrates. *Proc Natl Acad Sci U S A* **119**, e2209597119, (2022).
PMID: 35878037

15　Bongiovanni, G., Fidelio, G. D., Barra, H. S. & Hallak, M. E. The post-translational
incorporation of arginine into a beta-amyloid peptide increases the probability of alpha-
helix formation. *Neuroreport* **7**, 326-328, (1995). PMID: 8742481

Wang, J., Han, X., Leu, N. A., Sterling, S., Kurosaka, S., Fina, M., Lee, V. M., Dong, D. W., Yates, J. R., 3rd & Kashina, A. Protein arginylation targets alpha synuclein, facilitates normal brain health, and prevents neurodegeneration. *Sci Rep* **7**, 11323, (2017). PMID: 28900170

Zhao, J., Pan, B., Fina, M., Huang, Y., Shimogawa, M., Luk, K. C., Rhoades, E., Petersson, E. J., Dong, D. W. & Kashina, A. alpha-Synuclein arginylation in the human brain. *Transl Neurodegener* **11**, 20, (2022). PMID: 35395956

Lopez Sambrooks, C., Carpio, M. A. & Hallak, M. E. Arginylated calreticulin at plasma membrane increases susceptibility of cells to apoptosis. *J Biol Chem* **287**, 22043-22054, (2012). PMID: 22577148

Carpio, M. A., Decca, M. B., Lopez Sambrooks, C., Durand, E. S., Montich, G. G. & Hallak, M. E. Calreticulin-dimerization induced by post-translational arginylation is critical for stress granules scaffolding. *Int J Biochem Cell Biol* **45**, 1223-1235, (2013). PMID: 23567256

Saha, S., Wong, C. C., Xu, T., Namgoong, S., Zebroski, H., Yates, J. R., 3rd & Kashina, A. Arginylation and methylation double up to regulate nuclear proteins and nuclear architecture in vivo. *Chem Biol* **18**, 1369-1378, (2011). PMID: 22118671

Wong, C. C., Xu, T., Rai, R., Bailey, A. O., Yates, J. R., 3rd, Wolf, Y. I., Zebroski, H. & Kashina, A. Global analysis of posttranslational protein arginylation. *PLoS Biol* **5**, e258, (2007). PMID: 17896865

Cha-Molstad, H., Sung, K. S., Hwang, J., Kim, K. A., Yu, J. E., Yoo, Y. D., Jang, J. M., Han, D. H., Molstad, M., Kim, J. G., Lee, Y. J., Zakrzewska, A., Kim, S. H., Kim, S. T., Kim, S. Y. *et al.* Amino-terminal arginylation targets endoplasmic reticulum chaperone BiP for autophagy through p62 binding. *Nat Cell Biol* **17**, 917-929, (2015). PMID: 26075355

Xu, Z., Payoe, R. & Fahlman, R. P. The C-terminal proteolytic fragment of the breast cancer susceptibility type 1 protein (BRCA1) is degraded by the N-end rule pathway. *J Biol Chem* **287**, 7495-7502, (2012). PMID: 22262859

Hu, R. G., Brower, C. S., Wang, H., Davydov, I. V., Sheng, J., Zhou, J., Kwon, Y. T. & Varshavsky, A. Arginyltransferase, its specificity, putative substrates, bidirectional promoter, and splicing-derived isoforms. *J Biol Chem* **281**, 32559-32573, (2006). PMID: 16943202

Yoo, Y. D., Mun, S. R., Ji, C. H., Sung, K. W., Kang, K. Y., Heo, A. J., Lee, S. H., An, J. Y., Hwang, J., Xie, X. Q., Ciechanover, A., Kim, B. Y. & Kwon, Y. T. N-terminal arginylation generates a bimodal degron that modulates autophagic proteolysis. *Proc Natl Acad Sci U S A* **115**, E2716-E2724, (2018). PMID: 29507222

Davydov, I. V. & Varshavsky, A. RGS4 is arginylated and degraded by the N-end rule pathway in vitro. *J Biol Chem* **275**, 22931-22941, (2000). PMID: 10783390

Wang, J., Han, X., Wong, C. C., Cheng, H., Aslanian, A., Xu, T., Leavis, P., Roder, H., Hedstrom, L., Yates, J. R., 3rd & Kashina, A. Arginyltransferase ATE1 catalyzes midchain arginylation of proteins at side chain carboxylates in vivo. *Chem Biol* **21**, 331-337, (2014). PMID: 24529990

MacTaggart, B., Shimogawa, M., Lougee, M., Tang, H. Y., Petersson, E. J. & Kashina, A. Global Analysis of Post-Translational Side-Chain Arginylation Using Pan-Arginylation Antibodies. *Mol Cell Proteomics* **22**, 100664, (2023). PMID: 37832787

Wang, J., Yates, J. R., 3rd & Kashina, A. Biochemical analysis of protein arginylation. *Methods Enzymol* **626**, 89-113, (2019). PMID: 31606094

Wadas, B., Piatkov, K. I., Brower, C. S. & Varshavsky, A. Analyzing N-terminal Arginylation through the Use of Peptide Arrays and Degradation Assays. *J Biol Chem* **291**, 20976-20992, (2016). PMID: 27510035

31 Wang, J., Pejaver, V. R., Dann, G. P., Wolf, M. Y., Kellis, M., Huang, Y., Garcia, B. A.,
Radivojac, P. & Kashina, A. Target site specificity and in vivo complexity of the mammalian
arginylome. *Sci Rep* **8**, 16177, (2018). PMID: 30385798

32 Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: a sequence logo
generator. *Genome Res* **14**, 1188-1190, (2004). PMID: 15173120

33 O'Shea, J. P., Chou, M. F., Quader, S. A., Ryan, J. K., Church, G. M. & Schwartz, D. pLogo:
a probabilistic approach to visualizing sequence motifs. *Nat Methods* **10**, 1211-1212,
(2013). PMID: 24097270

34 Tan, K., Fujimoto, M., Takii, R., Takaki, E., Hayashida, N. & Nakai, A. Mitochondrial SSBP1
protects cells from proteotoxic stresses by potentiating stress-induced HSF1
transcriptional activity. *Nat Commun* **6**, 6580, (2015). PMID: 25762445

35 Lan X, H. W., Kim SB, Fu D, Abeywansha T, Lou J, Balamurugan U, Kwon YT, Ji CH,
Taylor DJ, Zhang Y Oligomerization and a distinct tRNA-binding loop are important
regulators of human arginyl-transferase function. *Nature Communications* **15**, 6350,
(2024). PMID: 39068213

36 Watanabe, K., Toh, Y., Suto, K., Shimizu, Y., Oka, N., Wada, T. & Tomita, K. Protein-based
peptide-bond formation by aminoacyl-tRNA protein transferase. *Nature* **449**, 867-871,
(2007). PMID: 17891155

37 Pagani, M., Pilati, S., Bertoli, G., Valsasina, B. & Sitia, R. The C-terminal domain of yeast
Ero1p mediates membrane localization and is essential for function. *FEBS Lett* **508**, 117-
120, (2001). PMID: 11707280

38 Xu, G., Shin, S. B. & Jaffrey, S. R. Global profiling of protease cleavage sites by
chemoselective labeling of protein N-termini. *Proc Natl Acad Sci U S A* **106**, 19310-19315,
(2009). PMID: 19892738

39 Long, G. L., Chandra, T., Woo, S. L., Davie, E. W. & Kurachi, K. Complete sequence of
the cDNA for human alpha 1-antitrypsin and the gene for the S variant. *Biochemistry* **23**,
4828-4837, (1984). PMID: 6093867

40 Brower, C. S., Piatkov, K. I. & Varshavsky, A. Neurodegeneration-associated protein
fragments as short-lived substrates of the N-end rule pathway. *Mol Cell* **50**, 161-171,
(2013). PMID: 23499006

41 Zhang, Y., Li, T., Zhang, L., Shangguan, F., Shi, G., Wu, X., Cui, Y., Wang, X., Wang, X.,
Liu, Y., Lu, B., Wei, T., Wang, C. C. & Wang, L. Targeting the functional interplay between
endoplasmic reticulum oxidoreductin-1alpha and protein disulfide isomerase suppresses
the progression of cervical cancer. *EBioMedicine* **41**, 408-419, (2019). PMID: 30826359

42 Goitea, V. E. & Hallak, M. E. Calreticulin and Arginylated Calreticulin Have Different
Susceptibilities to Proteasomal Degradation. *J Biol Chem* **290**, 16403-16414, (2015).
PMID: 25969538

43 Jiang, M., Xie, X., Zhu, X., Jiang, S., Milenkovic, D., Misic, J., Shi, Y., Tandukar, N., Li, X.,
Atanassov, I., Jenninger, L., Hoberg, E., Albarran-Gutierrez, S., Szilagyi, Z., Macao, B. *et*
al. The mitochondrial single-stranded DNA binding protein is essential for initiation of
mtDNA replication. *Sci Adv* **7**, (2021). PMID: 34215584

44 Del Dotto, V., Ullah, F., Di Meo, I., Magini, P., Gusic, M., Maresca, A., Caporali, L., Palombo,
F., Tagliavini, F., Baugh, E. H., Macao, B., Szilagyi, Z., Peron, C., Gustafson, M. A., Khan,
K. *et al.* SSBP1 mutations cause mtDNA depletion underlying a complex optic atrophy
disorder. *J Clin Invest* **130**, 108-125, (2020). PMID: 31550240

45 Rai, R. & Kashina, A. Identification of mammalian arginyltransferases that modify a
specific subset of protein substrates. *Proc Natl Acad Sci U S A* **102**, 10123-10128, (2005).
PMID: 16002466

46 Bachmair, A., Finley, D. & Varshavsky, A. In vivo half-life of a protein is a function of its
amino-terminal residue. *Science* **234**, 179-186, (1986). PMID: 3018930

47   Bouwmeester, R., Gabriels, R., Hulstaert, N., Martens, L. & Degroeve, S. DeepLC can
predict retention times for peptides that carry as-yet unseen modifications. *Nat Methods*
18, 1363-1369, (2021). PMID: 34711972
48   Wilburn, D. B., Shannon, A. E., Spicer, V., Richards, A. L., Yeung, D., Swaney, D. L.,
Krokhin, O. V. & Searle, B. C. Deep learning from harmonized peptide libraries enables
retention time prediction of diverse post translational modifications. *bioRxiv*, (2023). PMID:
37398395
49   Kleifeld, O., Doucet, A., auf dem Keller, U., Prudova, A., Schilling, O., Kainthan, R. K.,
Starr, A. E., Foster, L. J., Kizhakkedathu, J. N. & Overall, C. M. Isotopic labeling of terminal
amines in complex samples identifies protein N-termini and protease cleavage products.
Nat Biotechnol **28**, 281-288, (2010). PMID: 20208520
50   Ju, S., Kwon, Y., Kim, J. M., Park, D., Lee, S., Lee, J. W., Hwang, C. S. & Lee, C. iNrich,
Rapid and Robust Method to Enrich N-Terminal Proteome in a Highly Multiplexed Platform.
Anal Chem **92**, 6462-6469, (2020). PMID: 32267142

a

ATE1, Arg[10]

18E-CALR

ATE1, Arg[0]

mix 1:1 → denature reduce alkylate digest → 505.2775[++] 510.2815[++] H/L ratio = 1.0 → H/L ratio, n = 5, replicates

b

No.

pair 1876
site 235
peptide 230
protein 165

c

No. of sites

iPSC_CF 27
iPSC 34
iPSC_CM 36
HEK293T_KO 12
HEK293T 7
MCF7 21
HCC2935 11
H1437 31
A549 16
HeLa 46
heart 47
brain_1 32
brain_2 16
brain_PD 45
brain_PDD 42
brain_AD 48

cardiac — cancerous cell — patient tissue

d

H/L ratio

pair (1876)

e

N 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 C

f

R (199)
RO3 (8)
R_deami (25)
RO2 (3)

g

cellular process (98)
bioregulation (39)
metabolism (47)
multicellular process (22)
stimuli response (18)
localization (18)
development (17)
immune (4)
interspecies interaction (2)
reproduction (2)
growth (2)
homeostasis (3)
unassigned (40)