Diptera-specific Daedalus controls Zucchini endonucleolysis in

piRNA biogenesis independent of exonucleases

Yuica Koga, Shigeki Hirakata, Mayu Negishi, Hiroya Yamazaki, and Mikiko C.

Siomi*

Department of Biological Sciences, Graduate School of Science, The University of Tokyo,

Tokyo 113-0032, Japan

*Correspondence: siomim@bs.s.u-tokyo.ac.jp

Summary

piRNAs protect germline genomes and maintain fertility by repressing transposons. Daedalus and Gasz act together as a mitochondrial scaffold for Armitage, a necessary factor for Zucchini-dependent piRNA processing. However, the mechanism underlying this function remains unclear. Here, we find that the roles of Daedalus and Gasz in this process are distinct, although both are necessary: Daedalus physically interacts with Armitage, whereas Gasz supports Daedalus to maintain its function. Daedalus binds to Armitage through two distinct regions, an extended coiled-coil identified in this study, and a SAM. The former tethers Armitage to mitochondria, while the latter controls Zucchini endonucleolysis to define the length of piRNAs in an exonuclease-independent manner. piRNAs produced in the absence of Daedalus SAM do not exhibit full transposon silencing functionality. Daedalus is Diptera specific. Unlike *Drosophila* and mosquitoes, evolutionarily more complex species, such as mice, rely on exonucleases after Zucchini processing to specify the length of piRNAs.

Keywords

Armitage, Daedalus, Gasz, piRNA, Zucchini, AmTEC, extended coiled-coil

Introduction

In *Drosophila* ovarian somatic cells (OSCs), PIWI-interacting RNAs (piRNAs) associate with one of the PIWI members, Piwi, in the cytoplasm to form the piRNA-induced silencing complex (piRISC).[1–4] piRISC then co-transcriptionally represses transposons in the nucleus. The vast majority of piRNAs in OSCs originate from the piRNA cluster, *flamenco* (*flam*).[5] The *flam* locus contains numerous transposon remnants, most of which

have an orientation opposite to the direction of transcription. Therefore, piRNAs can naturally hybridize to target mRNAs through RNA–RNA base pairing.[5,6] After nuclear processing, including splicing,[7] the *flam* transcripts are exported to the cytoplasm where they bind to female sterile (1) Yb (Yb) via *cis*-elements embedded in the RNAs. This interaction induces the assembly of non-membranous organelle Yb bodies through phase separation, which accelerates subsequent steps in the piRNA processing.[8,9]

At this point, the *flam* transcripts are coarsely digested by an unknown factor, generating numerous 5′ ends. Piwi binds one-to-one to these 5′ ends via the MID domain and becomes the Piwi-piRISC precursor (pre-Piwi-piRISC).[10,11] Concomitantly, the RNA helicase Armitage (Armi) binds to the complex, allowing pre-Piwi-piRISC to exit Yb bodies and move to the mitochondrial surface where the endonuclease Zucchini (Zuc) awaits to process the complex.[12] Zuc has a transmembrane (TM) region, which localizes Zuc to the outer membrane of mitochondria.[13–16] After Zuc RNA cleavage releases mature Piwi-piRISC from pre-Piwi-piRISC, nascent Piwi binds to the remaining piRNA precursor through its 5′ end, generating a new pre-Piwi-piRISC in Yb body-independent and Zuc-dependent manners.[17,18] This mechanism of sequential generation of Piwi-piRISC from a single piRNA precursor is known as phasing.[17,18] Piwi-piRISC is then transported to the nucleus,[19] where it associates with nascent transposon transcripts and suppresses transcription in concert with multiple cofactors, such as Gtsf1/Asterix and Maelstrom.[20–27]

An RNAi-based gene screen performed in *Drosophila* ovaries identified many factors acting in the piRNA pathway, including Gasz and Daedalus (Daed).[28] Both proteins have a TM, like Zuc, and they form a heteromeric complex that acts in concert with Armi to support Zuc cleavage in piRNA maturation.[12,29,30] However, the molecular

details of this process, especially the function of Daed, remain unclear.

In this study, we found that Daed interacts with Armi through two distinct regions,

the extended coiled-coil (eCC) identified in this study, and the sterile α motif (SAM). We

also show that Daed interacts with Gasz by forming a helical barrel using the two α-

helixes present in each of Daed and Gasz. Through this interaction, Daed stabilizes Gasz,

but not *vice versa*. Furthermore, we found that the Armi–Daed interaction through Daed

eCC tethers Armi to mitochondria, while the Armi–Daed interaction through SAM controls

Zuc endonucleolysis to specify the length of piRNAs in an exonuclease-independent

manner. piRNAs produced in the absence of Daed SAM became atypically long and their

function in transposon silencing was attenuated. Evolutionarily more complex species,

such as mice, possess exonuclease, PNLDC1, but not Daed. Mouse piRNAs, which are

atypically elongated in the absence of PNLDC1, do not exhibit full functionality in

transposon silencing. These findings indicate that piRNA length specification, which is

crucial for transposon repression, is conserved between species, but that the means of

piRNA length specification differ between species.

Results

Daed and Gasz bind with each other via their respective eCC domains

According to the current model, the Daed–Gasz heterodimer tethers Armi bound to pre-

Piwi-piRISC to mitochondria for Zuc-dependent piRNA processing.[12,29,30] In this study,

we first predicted the higher-order structure of the Daed–Gasz heterodimer using

AlphaFold2[31,32] (Figure 1A). This indicated that both Gasz and Daed have two α-helixes

and that the coiled-coil structure composed of these four α-helixes is the center of the

Daed–Gasz association (Figure 1B). Daed has SAM, coiled-coil (CC), and TM domains,

and Gasz has ankyrin repeats (ARs), SAM, and TM domains.[12,29] The predicted

structures of Daed and Gasz showed that the second of the two α-helixes (Ala191–

His235) of Daed roughly overlaps with the CC domain (Thr206–Ser234) and that the two

α-helixes in Gasz (Phe345–Glu374 and Glu383–Ser412) are present between the SAM

and TM domains (Figure 1C). The amino acid sequence-based prediction of the CC

region[33] also roughly overlaps with the tandem α-helix regions (Figure S1A), supporting

the AlphaFold2 prediction. We propose that the two tandem α-helixes present in both

Gasz and Daed are referred to as extended coiled-coil (eCC) domains (Figure 1C).

The CC domain of Daed is the region that binds to Gasz.[29] To identify the region

of Gasz responsible for binding to Daed we expressed deletion mutants of Gasz in OSCs

lacking endogenous Gasz and then performed immunoprecipitation assays. Three of the

Gasz deletion mutants, ΔAR1-3, which lacks the first three ARs, ΔAR4-6, which lacks

the second three ARs, and ΔSAM, bound to Daed as efficiently as wild-type (WT) Gasz

did, but the other mutant, Δα-H2, lacking the second α-helix of eCC did not (Figures 1D

and S1B). These results indicate that the respective eCC domains of Daed and Gasz

are responsible for the Daed–Gasz association, consistent with the predicted structure.

Daed stabilizes Gasz via protein–protein interaction

To verify whether Daed is required for piRNA biogenesis in cultured OSCs, we depleted

intracellular Daed by RNA interference (RNAi) and conducted northern blotting. In control

OSCs, both *idefix* piRNA (a representative nongenic piRNA) and *traffic jam* (*tj*) piRNA (a

representative genic piRNA) were detected as expected (Figure S2A).[12] However, in the

absence of Daed, their piRNA precursors (pre-piRNAs) accumulated aberrantly, while

mature piRNAs were barely detected (Figure S2A), as in OSCs lacking Gasz.[12] This

indicates that both Gasz and Daed are essential for piRNA maturation in OSCs.

Western blotting confirmed that RNAi was effective at depleting Daed and Gasz

in OSCs (Figure 2A). We also found that, when Daed was depleted, levels of Gasz were

also markedly reduced, but that levels of Daed were unchanged in the absence of Gasz

(Figures 2A and 2B). Such trends were not observed for their mRNA levels (Figure 2C).

These results indicate that Daed is required to stabilize Gasz at the protein level, but that

Daed is not dependent on Gasz for its stabilization.

To investigate which domain of Daed is involved in Gasz stabilization, two

deletion mutants of Daed, ΔSAM and ΔCC (part of the second α-helix of the eCC, see in

Figure 1C), were expressed in OSCs depleted of endogenous Daed. Levels of Gasz

were restored with expression of WT Daed and ΔSAM mutant Daed, but not with ΔCC

mutant Daed (Figure 2D). A previous study showed that Daed binds to Gasz via the CC

region[29] and together these findings indicate that the Daed–Gasz interaction via their

respective eCC domains is crucial for the stabilization of Gasz (Figure 2E). The Δα-H2

mutant of Gasz, which lacks the binding domain to Daed, was rather unstable in OSCs,

but this was not because of mRNA instability (Figure S2B), supporting the above notion.

Daed alone or Gasz alone is insufficient to tether Armi to mitochondria

Armi has at least two roles in piRNA biogenesis in OSCs; one is to transport nascent

Piwi to Yb bodies to promote pre-Piwi-piRISC assembly, and the other is to translocate

the complex from Yb bodies to mitochondria to promote Zuc-dependent piRNA

processing.[12,34] After this second role, Armi returns to the cytosol to repeat these

functions, although immunofluorescence shows that Armi appears to be a constant

resident of Yb bodies (Control, Figure 2F).[8,12] In the absence of Zuc, Armi signals merged

with mitochondrial signals {[I] Zuc knockdown (KD), Figure 2F}. This demonstrates

dynamic movement of Armi in OSCs and indicates that Armi remains on the

mitochondrial surface when loss of Zuc impairs piRNA maturation.[12] Results from

mitochondrial fractionation followed by western blotting were consistent with this

statement (Figures 2G and S2C).

We then depleted Daed in Zuc-lacking OSCs, where both Daed and Gasz

should be absent (Figure 2A). Under these circumstances, as expected, Armi was not

retained on mitochondria and appeared in Yb bodies ([II] Zuc KD + Daed KD, Figure 2F).

To restore the levels of both Daed and Gasz simultaneously, Flag-Daed was

exogenously expressed in the OSCs and western blotting confirmed the recovery of both

Daed and Gasz levels (Figures 2G and S2C). Immunofluorescence showed that Armi

re-localized to mitochondria ([III] Flag-Daed, Figure 2F). However, when Gasz was

forcibly expressed alone in the absence of Daed, Armi remained in Yb bodies ([IV] Flag-

Gasz, Figure 2F). A similar result was obtained in OSCs expressing only Daed ([V] Zuc

KD + Gasz KD, Figure 2F). These results indicate that Gasz alone or Daed alone is not

sufficient to tether Armi to mitochondria. In other words, both Daed and Gasz are

required to retain Armi on the mitochondrial surface for Zuc-dependent piRNA

processing. Hereafter, the Daed–Gasz heterodimer is referred to as the Armi

mitochondrial tethering complex (AmTEC). The behavior of Armi in OSCs under the

conditions described above is summarized in Figure S2D.

Daed, but not Gasz, binds to Armi via two distinct sites

We next predicted the higher-order structure of AmTEC bound with Armi (Figure 3A).

Notably, no significant interaction between Gasz and Armi was observed, but Daed and

Armi were bound at two distinct sites: The first site was through the first α-helix of eCC

(Phe144–Val177) (Figure 1C), which interacted with the N-terminal region of Armi

(Met1–Leu34) (Figures 3A, 3B, and S3A). The second interaction was between the SAM

domain of Daed (Figure 1C) and the C-terminal helicase (i.e., UPF1-like family) domain

of Armi (Figures 3A, 3C, and S3A).

To verify that the Armi–Daed interaction was via the Armi N-terminus (Figure

3B), recombinant WT Armi (WT-Flag) and its mutant lacking 34 N-terminal residues

(ΔN34-Flag) were subjected to pull-down assays with Daed (Daed ΔTM) as prey. The

Daed TM was removed because its inclusion interfered with protein preparation. Under

conditions where a negative control, Luciferase (Luc), showed no binding to Daed ΔTM,

Armi WT bound to Daed but Armi ΔN34 did not (Figure 3D). This confirms that Armi and

Daed interact through the 34 N-terminal residues of Armi. We have previously shown

that the Armi ΔN34 mutant (i.e., a shorter isoform of Armi) is unable to leave Yb bodies.[12]

The present analysis raises an alternative hypothesis; the Armi ΔN34 mutant may be

able to leave Yb bodies but not be tethered to mitochondria because of the absence of

Daed binding. This would explain why the shorter Armi isoform is inactive in piRNA

biogenesis in OSCs.

To verify the Armi–Daed interaction via the Armi UPF1-like helicase domain

(Figure 3C), three residues in the domain, Asp1015, Arg1026, and Arg1037 (D-R-R),

which appear to be key to the binding, were mutated to Lys-Glu-Glu (K-E-E) and

immunoprecipitation assays were conducted (Figure 3E). Quantification of the results

indicated that the Armi KEE mutant bound to Daed with approximately 34% of the binding

efficiency of WT Armi (Figure 3F). Meanwhile, the Armi ΔN34 mutant bound to Daed with

approximately 20% of the binding efficiency of WT Armi. Immunofluorescence showed

that the Armi ΔN34 mutant accumulated in Yb bodies more intensely than the KEE

mutant in Arm/Zuc-lacking OSCs (Figure 3G). For binding to Daed, Armi may be more

dependent on the N-terminal region than on the UPF1-like helicase domain.

A lack of Armi in OSCs severely disrupted piRNA production.[16,35] This was

almost fully rescued by the expression of WT Armi (Figure 3H). We next expressed the

ΔN34 or KEE mutant of Armi as alternatives to the WT. The ΔN34 mutant failed to rescue

the disruption to piRNA production, but the KEE mutant partially restored piRNA

biogenesis (Figures 3H and S3B). This supports the notion that the N-terminal region of

Armi is more important than the C-terminal helicase domain in piRNA biogenesis in

OSCs. The strength of the Armi–Daed interaction (Figure 3E) appears to be positively

correlated with the function of Armi.

The two Armi–Daed associations differ in their functional contributions to piRNA

biogenesis

piRNAs produced under conditions involving the Armi KEE mutant were notably

increased in length by several bases (Figure 3H). We therefore investigated whether a

similar phenomenon occurred under conditions involving the Daed ΔSAM mutant, given

that the SAM domain is the binding partner of the UPF1-like helicase domain of Armi.

Indeed, in cells expressing the Daed ΔSAM mutant as an alternative to endogenous

Daed, Piwi-bound piRNAs became several bases longer and their levels were lower than

those in cells expressing WT Daed, but higher compared with cells lacking endogenous

Daed (Figures 4A and S4A).

To examine how the lack of binding between Daed SAM and Armi UPF1-like

helicase domain actually affects piRNA biogenesis, we sequenced and analyzed Piwi-

bound piRNAs in OSCs expressing Daed ΔSAM and compared the results with those

from OSCs expressing WT Daed (Figures 4A and S4A). This length distribution of

piRNAs was indeed shifted towards the longer fractions when they were produced in

cells lacking the Daed SAM domain (Figure 4B). This trend was also observed when

piRNAs were subdivided by target transposons (Figure S4B). In contrast, piRNA-

mapped transposons were essentially unchanged (Figure S4C). This seems reasonable

given that piRNAs are produced from *flam* in both cases. Furthermore, the 1U bias[5,36]

and piRNA phasing[17,18] were similarly observed in both libraries (Figures 4C and 4D).

The β-elimination experiments showed the 3′ ends of the piRNAs to be modified with 2′-

O-methylation in both cases, unlike Argonaute 1 (Ago1)-bound miRNAs (Figure 4E).[37,38]

However, it should be noted that non-negligible amounts of Piwi immunofluorescence

remained in the cytoplasm when Daed ΔSAM was expressed in OSCs lacking

endogenous Daed (Figure 4F).

piRNA loading is essential for Piwi nuclear localization.[19,34] Therefore, we next

examined the piRNA binding status of cytoplasmic and nuclear Piwi. In contrast to

nuclear Piwi, which was bound to piRNAs even in the long fraction, cytoplasmic Piwi was

bound to few piRNAs (Figure S4D). This indicates that piRNAs produced under the

ΔSAM conditions can localize Piwi to the nucleus as piRISC, regardless of length, but

are insufficient to convert all Piwi to functional Piwi-piRISC. Consistent with this,

transposons were not fully repressed under such conditions (Figures 4G and S4E).

These results indicate that the Armi–Daed association via the UPF1-like helicase domain

of Armi and the SAM domain of Daed is influential in Piwi-piRISC production and function,

although less so than the other Armi–Daed association involving the 34 N-terminal

residues of Armi and the eCC domain of Daed.

The Armi–Daed interaction via Daed SAM optimizes Zuc endonucleolysis for

piRNA-mediated transposon repression

To explore the nature of ∆SAM piRNA extension, we selected the 50,000 piRNA 5′ end

positions common to the WT and ∆SAM and with high abundance in WT libraries (Figure

S4F) and analyzed them further. Heatmaps depicted with their 5′ end positions aligned

clearly indicate that ∆SAM piRNAs are more diverse in length. Many of them have some

extra bases at the 3′ ends (Figure 4H). This is consistent with the length distribution

shown in Figure 4A. There are also a few shorter piRNAs (Figure 4H). The relatively

small number of shorter piRNAs may be because of protection by Piwi.

To further analyze the effects of Daed SAM loss on piRNAs, we extracted the

5′-end positions that dominantly produce 26- and 27-nucleotide (nt) piRNAs in the WT

library, and then these positions were classified as "Unchanged" (51,833 positions) and

"Elongated" (53,258 positions) according to the absence or presence of 3′-end

elongation. piRNAs produced from both 5′-end groups maintained comparable phasing

and +1U bias (i.e., 1U bias of piRNAs, real and virtual, immediately following piRNAs

verified to bind to Piwi[17,18]) (Figure S4G), indicating that piRNA elongation caused by the

loss of Daed SAM is caused by Zuc endonucleolysis.

We next sorted piRNAs by their length and aligned them according to the 3′

ends of piRNAs. This showed that atypically long piRNAs of around 28–33 nt in the WT

library tended to have significantly low U bias near their 3′ ends (U depletion), just

upstream of the 5′ ends of the immediately downstream piRNAs, as reported previously

(yellow boxes in WT, Figure 4I).[18] This was interpreted as showing that when Zuc cannot

find U-rich regions in the piRNA precursors it waits to cleave the substrate until the U-

poor region has passed, thereby producing atypically long piRNAs. This also explains

why there were fewer 28–33-nt piRNAs in the WT library (Figure 4B). In contrast, there

was less notable "U depletion" in 28–33-nt ΔSAM piRNAs (yellow boxes in ΔSAM, Figure

4I). In addition, there was a substantial number of 28–33-nt piRNAs in the ΔSAM library

(Figure 4B). This indicates that when the Armi–Daed interaction is attenuated because

of a lack of Armi–Daed association via Daed SAM (and the UPF1-like helicase domain

of Armi), Zuc somehow produces these longer piRNAs more preferentially regardless of

the U content in the precursor sequences while maintaining the +1U bias.

We simulated the effect of Daed ΔSAM on piRNA production. First, we

hypothetically fragmented the sequence of the *flam* locus (21,629,891 to 21,792,731 on

the X chromosome; T was converted to U) into 163 piRNA precursors, starting at U and

into sizes of 1,000 nt or close to that size. In parallel, the data shown in Figure 4B were

fitted to normal distributions. Then we simulated Zuc-dependent piRNA production from

their 5′ ends by assuming that Zuc cleavage occurs before U, produces 18–40-nt piRNAs,

and follows the normal distributions for WT and ΔSAM. Sequence analysis of the

obtained piRNAs (25–34 nt) indicated that U depletion was observed in simulations fitting

the WT distribution, starting from 29 nt, but less so in simulations fitting the ΔSAM

distribution (Figure S4H). This suggests that Zuc has the potential to produce Piwi-

piRISC without following the U-depletion rule.

To investigate potential causes of Zuc-dependent precursor cleavage changes

in ΔSAM, molecular dynamics simulations were performed using the Armi–AmTEC

complex model (Figure 3A) and its Daed ΔSAM version. The distance between the center

of mass (CoM) of Armi helicase domain (Ser697–Cys1,148 in Figure S3A) and that of

the AmTEC was measured (n=5) and the average value was calculated at all time points

along each trajectory (Figures S5A and S5B). This analysis shows that there is little

difference in the initial Armi–AmTEC CoM distances between Daed WT and ∆SAM, but

as the simulation time progresses, the difference tends to increase for the ∆SAM version

(Figure S5C). The statistical analysis also shows that the distribution of CoM distances

at all time points for all trajectories shifts to larger values for ∆SAM (Figures S5D). These

results suggest that Daed ∆SAM tethers Armi more precariously than Daed WT. Based

on these results, we hypothesize that "proper" binding of Armi and Daed via the Daed

SAM (and the UPF1-like helicase domain of Armi) is critical for maintaining the distance

between Armi and the scaffold complex, the loss of which would obscure the location of

piRNA precursor cleavage by Zuc, negatively influencing Piwi-piRISC production.

We also sequenced and analyzed Piwi-bound piRNAs in OSCs expressing the

Armi KEE mutant or WT Armi (Figures 3H and S3B). We found that piRNAs in the Armi

KEE library were also several bases longer than the Armi WT piRNAs and retained 1U

bias (Figures S6A and S6B). Heatmaps showed that the KEE piRNAs also exhibited

characteristics of 3′ end elongation (Figure S6C). These results further support our notion

that the SAM domain of Daed and the UPF1-like helicase domain of Armi are the binding

partners and that the Armi–Daed interaction is indispensable to optimize Piwi-piRISC

maturation by Zuc and piRISC function in OSCs (Figure S6D).

Daed is Diptera-specific and controls Zuc cleavage in a Nibbler-independent manner

Armi and Gasz are widely conserved among species from invertebrates to vertebrates,

but Daed is unique to the Diptera order, which is composed of Brachycera suborder,

which includes *Drosophila* and other fly species, and Nematocera suborder which

includes mosquitoes (Figures 5A, S7A, and S7B). In contrast, other species not

belonging Diptera order do not appear to have Daed orthologs.[39]

In mice, primary piRNA biogenesis occurs in the inter-mitochondrial cement,

where Armi homolog, MOV10L1, Gasz homolog, ASZ1, Zuc homolog, PLD6, and TDRD

proteins play a role in piRNA production: MOV10L1 binds to PIWI proteins (MILI, MIWI

and MIWI2) and piRNA precursors and contributes to piRNA biogenesis through its

helicase activity.[40,41] ASZ1 is essential for the production of MILI-bound piRNAs in mice[42]

and is thought to tether MOV10L1 to mitochondria, together with TDRD proteins, such

as TDRD1, to promote MIWI-piRISC production.[40,41]

We expressed both MOV10L1 and ASZ1 in OSCs and performed

immunoprecipitation. The MOV10L1–ASZ1 interaction was detected to a similar extent

in both normal OSCs and OSCs lacking intracellular Daed (Figures 5B and 5C).

Immunofluorescence confirmed this; the expression of ASZ1 was sufficient for the

mitochondrial localization of MOV10L1, independent of the presence of intracellular

Daed (Figure 5D). This indicates that the MOV10L1–ASZ1 interaction and MOV10L1

tethering is Daed-independent, while the Armi–Gasz interaction and Armi tethering in

Drosophila required Daed. These results also indicate that the Armi–Gasz association,

at least in OSCs, does not require other factors present in mouse testis. ASZ1 was stable

in OSCs lacking endogenous Daed (Figure 5C), unlike Gasz (Figure 2A), indicating that

the molecular system that actively degrades Gasz in the absence of Daed does not act

on ASZ1.

Why did *Drosophila* acquire Daed during evolution? In mice, poly(A)-specific

ribonuclease-like domain containing 1 (PNLDC1), a homolog of poly(A)-specific

ribonuclease (PARN), trims the 3′ ends of piRNAs and piRNAs are elongated after

PNLDC1 loss of function.[43,44] Mutation in PNLDC1 reduces LINE-1-targeting piRNAs and

mislocalizes MIWI2 to the cytoplasm, leading to LINE-1 derepression. This was nearly

equal to what we observed in this study with the *Drosophila* Daed ΔSAM mutant.

However, PNLDC1 is not conserved in the majority of Diptera order species.[45] Papi is

the *Drosophila* ortholog of TDRKH, which, in mice, acts as a scaffold for PNLDC1 and

whose functional inhibition causes piRNA elongation.[46,47] However, consistent with

previous studies using Papi mutant flies,[45] its depletion had little effect on piRNA length

in OSCs (Figure 5E).

The 3′-to-5′ exonuclease, Nibbler (Nbr), trims both miRNAs and piRNAs in

Drosophila and mosquitoes,[45,48–50] but the pre-piRNAs that undergo trimming are minor

in mosquitoes, and the loss of Nbr barely affected the length of Piwi-bound piRNAs in

the *Drosophila* ovary.[45] Our study confirmed this; namely, Nbr depletion in OSCs had

little effect on the piRNA length (Figure 5F). This indicates that Nbr may function in the

trimming of miRNAs, but not piRNAs, in OSCs. The lack of a marked increase in +1U

signal in piRNAs elongated in the Daed ΔSAM mutant (Figures 4D and S4G) also

supports this view.

In contrast, Armi D-R-R, which was essential for the regulation of Zuc cleavage

in *Drosophila*, is highly conserved in the fly species that lack PNLDC1/PARN and have

Daed (Figure S8). These results indicate that *Drosophila,* and its closely related species,

may have acquired Daed after losing PNLDC1/PARN during evolution, and that they

may also have acquired the ability to produce mature piRNAs without further trimming

by exonucleases by allowing Armi to bind directly to Daed (Figure 5G).

Discussion

This study revealed that Daed and Gasz bind to each other via their eCC domains, both

of which are composed of two α-helixes that assemble into a stable helical barrel

consisting of all four of these α-helixes. When Daed lacks the second α-helix in the eCC

domain, the Daed–Gasz interaction is disrupted and Gasz becomes destabilized,

illustrating the importance of Daed in Gasz stabilization (Figure 2E). This also shows that

the intracellular level of Gasz is strictly regulated by the intracellular level of Daed,

indicating that OSCs do not tolerate the presence of Gasz alone. In contrast, OSCs do

tolerate the presence of Daed alone. However, Daed, in the absence of Gasz, failed to

anchor pre-Piwi-piRISC to mitochondria via Armi. Gasz is therefore complicit in the role

of Daed, although the contribution of Gasz to the direct binding of Daed to Armi appears

to be minimal.

Lack of binding through the Daed SAM did not completely abolish the Armi–

Daed association but resulted in aberrant Zuc function (Figure S6D). Under such

conditions, Piwi-bound piRNAs were atypically long because of extra bases at the 3′

ends, and there was no notable U depletion, a typical feature of piRNAs in normal OSCs.

In normal OSCs, atypically long piRNAs are produced only when piRNAs of typical sizes

cannot maintain the +1U bias. In contrast, in OSCs expressing Daed ∆SAM, Zuc shows

"relaxed selectivity" in terms of nucleotides and length and tends to produce longer

piRNAs, even under conditions where it can produce shorter (i.e., normal size) piRNAs

with the +1U bias, albeit in relatively small quantities. This could be because of the timing

of Zuc RNA cleavage was shifted slightly downstream of the precursor while the

preference for cleavage before U was conserved. Recombinant Zuc did not show U

selectivity,[15] indicating that, *in vivo*, Zuc may rely on a cofactor to confer U selectivity,

and that this cofactor may not require the Armi–Daed association to execute its function.

Interestingly, Piwi could accept those longer piRNAs produced under the ΔSAM

conditions and localized to the nucleus as Piwi-piRISC. At the same time, however,

nonnegligible amounts of Piwi remained in the cytoplasm because of the lack of piRNA

loading, which resulted in transposon derepression. This means that the ability of Daed

to interact with Armi via the SAM domain, in addition to the binding via the first α-helix in

the eCC domain, is essential not only to define the positions of Zuc cleavage, but also

to produce enough Piwi-piRISC for transposon repression in OSCs. It is worth noting

that no such regulatory factor has been identified to date.

Armi is unique in that it plays multiple roles in piRNA biogenesis. First, Armi

delivers unloaded Piwi to Yb bodies, facilitating the assembly of pre-Piwi-piRISC in these

organelles. In Yb bodies, Armi binds to pre-Piwi-piRISC and translocates the complex to

mitochondria. There, Armi anchors pre-Piwi-piRISC to the mitochondria by binding to

Daed. This allows Zuc to process pre-Piwi-piRISC into mature Piwi-piRISC. Armi is

therefore a factor that always accompanies Piwi and encourages it along the pathway to

develop into Piwi-piRISC.

Armi is an RNA helicase protein, and its RNA helicase activity is required to

avoid stochastic interactions with abundant mRNAs.[51] In contrast, up to this point, the

RNA helicase activity of Armi does not seem to be required for the piRNA precursors,

rather its activity would be inhibitory. However, in the absence of this activity, especially

for phasing, the sequential generation of Piwi-piRISC from a single piRNA precursor

would not occur. It is possible that an inhibitor of the RNA helicase activity of Armi is

present in OSCs and functions until phasing, but, upon binding to Armi, Daed would lose

its function allowing phasing to occur.

Animals that reproduce sexually retain the highly conserved piRNA mechanism

and use it to maintain their fertility by repressing transposons. However, the evolutionary

rate of many piRNA factors, including Armi, is rapid,[52,53] suggesting that these species

have adapted the machinery to their specific endogenous and exogenous environments.

Coevolution of piRNAs and transposons through host–pathogen arms races has been

proposed as a background to this process.[54] Although flies and mosquitoes belong to

the same Diptera order, the piRNA pathway factors are highly diverse.[55] On the other

hand, our observation that Daed is conserved among Diptera species may suggest the

importance of this factor in Dipteran piRNA biogenesis pathway where PNLDC1 is lost.

Author contributions

YK and MN performed biochemical experiments. YK, SH, and HY performed

bioinformatic analyses, computational structural prediction, simulation, and calculation.

MCS supervised and discussed the work and wrote the manuscript with the other authors.

Acknowledgments

We thank Y. Namba, Y.W. Iwasaki, H. Nishimasu, and S. Kuraku for useful discussions,

K. Kudo, T. Yasuda, and Y. Fukushima for technical assistance with the molecular

dynamics simulation, and T. Fujisawa, T. Kinoshita, and H. Ishizu for technical

assistance with the biochemical analysis. This study was supported by MEXT KAKENHI

Grant Numbers JP19H05466 (to MCS), JP22K15056 (to SH), JP23K14141 (to HY), and

JP22KJ1122 (to YK). Some computations were performed on the NIG supercomputer at

ROIS National Institute of Genetics.

Conflict of interest

The authors declare no competing interests.

Figure legends

Figure 1. Daed and Gasz interact via their respective eCC domains

(**A**) Predicted structure of the Daed–Gasz heterodimer by AlphaFold2. (**B**) Magnification

of the region indicated by the dotted line box in (**A**). The structure is shown from two

directions. (**C**) Domain structures of Daed and Gasz. Peptide sequences of eCCs of

Daed and Gasz are also shown. (**D**) Immunoprecipitation (Flag IP) and western blotting

show that one of the Gasz mutants, Δα-H2, fails to bind to Daed in OSCs lacking

endogenous Gasz. Flag-Luc: negative control. *: Heavy chains. The domain structures

Gasz WT and deletion mutants are also shown.

Figure 2. Daed stabilizes Gasz via interaction and both Daed and Gasz are

essential for the mitochondrial tethering of Armi

(**A**) Western blotting shows the high efficiency of knockdown (KD) of Daed and Gasz in

OSCs. Daed KD also destabilizes Gasz, but not *vice versa*. β-Tub: loading control. (**B**,

C) The statistics of western blotting signals (n=3; **A**) and the mRNA expression levels of

Daed and *Gasz* upon Daed KD and Gasz KD (n=3). Signal intensities (**B**) or values (**C**)

relative to control (EGFP KD) were normalized with those of β-Tub or *rp49*, respectively,

and presented as mean values±SE. **: $p <0.01$; ***: $p <0.005$; n.s.: $p >0.05$ [*t*-test

(unpaired, two-sided)]. (**D**) Western blotting showing the levels of endogenous Gasz and

Flag-Daed (WT and deletion mutants) in OSCs under each condition. β-Tub: loading

control. The domain structures of Daed WT and mutants are shown. (**E**) Model of Gasz

stabilization by Daed. (**F**) Subcellular localization of Armi (green) in OSCs under various

441 conditions, [I] to [V]. Mitochondria (mito) and Yb bodies (Yb; white arrowheads) are

442 shown in magenta. The signals of Flag-Gasz and Flag-Daed are shown in yellow. Images

443 in the same vertical row are identical cells. Control: normal OSCs. DAPI (blue): nuclei.

444 Scale bar 5 μm. (**G**) Western blotting shows the levels of endogenous Armi, Gasz, Daed,

445 Flag-Gasz (orange asterisk), and Flag-Daed (blue asterisk) in total lysates (total) and

446 mitochondrial fractions (mito) under conditions [I] to [VI]. For the efficiency of cell

447 fractionation, see Figure S2C.

449 **Figure 3. Daed interacts with Armi through two sites, each contributing differently**

450 **to piRNA biogenesis**

451 (**A**) The structure of the Armi–AmTEC complex predicted by AlphaFold2. Gray lines:

452 virtual lipid bilayers. The domain structures of Armi, Daed, and Gasz are also shown. (**B**)

453 Magnification of the "B" region in (**A**). The 34 N-terminal residues of Armi (N34) interacts

454 with the first α-helix of Daed eCC. (**C**) Magnification of the "C" region in (**A**). The Armi

455 UPF1-like helicase domain interacts with Daed SAM. Asp1015, Arg1026, and Arg1037

456 of Armi are shown in blue. (**D**) *In vitro* pull-down assays show Armi WT-Flag, but not

457 Armi ΔN34-Flag, binding to Daed (Daed ΔTM-His). Luc-Flag: negative control. *: Heavy

458 and light chains. (**E**) Immunoprecipitation (Flag IP) and western blotting show the

459 amounts of endogenous Daed co-immunoprecipitated with Armi WT-Flag, ΔN34-Flag,

460 and KEE-Flag. Luc-Flag: negative control. The domain structures of Armi WT and its

461 mutants are shown below. Asp1015, Arg1026, and Arg1037 mutated in the KEE mutant

462 are also shown. The peptide sequence of Armi is shown in Figure S3A. (**F**) The statistics

463 of western blotting signals (n=3; **E**). The signals of Daed were normalized with those of

464 Armi and presented as mean values±SE. *: $p < 0.05$; ***: $p < 0.005$ [*t*-test (unpaired,

two-sided)]. (**G**) Subcellular localization of Armi WT, ΔN34, and KEE (green) in OSCs

lacking endogenous Zuc (Zuc KD) and Armi (Armi KD). Mitochondria (mito; for WT and

KEE) and Yb bodies (Yb, white arrowhead; for ΔN34) are shown in magenta. Images in

the same vertical row are identical cells. DAPI (blue): nuclei. Scale bar: 5 μm. (**H**) Piwi-

bound piRNAs obtained under each condition. Signal intensities (n=3) are normalized as

a percentage of the total. The mean (line) and 95% confidence interval (shadow) are

shown. The amount of Piwi in each sample is shown at the bottom.

Figure 4. The loss of Armi–Daed interaction via Daed SAM obscures Zuc cleavage

and impairs Piwi-piRISC function

(**A**) Piwi-bound piRNAs obtained under each condition. Signal intensities (n=3) are

normalized as a percentage of the total. The mean (line) and 95% confidence interval

(shadow) are shown. The amount of Piwi in each sample is shown at the bottom. (**B**)

Length distribution of piRNAs (18–40 nt) in Daed WT (green) and ΔSAM (red) libraries

(n=3), presented as mean values ± SE. (**C**) Sequence logos representing 11 nt from the

5′ end of piRNAs in Daed WT and ΔSAM libraries. (**D**) Heatmaps showing 3′ end counts

of piRNAs mapped around the 10,000 most abundant piRNA 5′ ends (lower) and its

binary histograms (upper) in Daed WT (left) and ΔSAM (right). (**E**) β-elimination assays.

Piwi-bound piRNAs in Daed KD OSCs expressing Daed WT and ΔSAM are shown.

Ago1-bound miRNAs are used as positive controls. (**F**) Subcellular localization of

endogenous Piwi (green) in Daed KD OSCs expressing Daed WT and ΔSAM (magenta).

DAPI (blue): nuclei. Scale bar: 5 μm. (**G**) The expression levels of transposons (*mdg1*,

tabor, *412*, and *stalker2*) in control OSCs (dark gray), Daed-lacking OSCs (Daed KD;

light gray), Flag-Daed WT rescued (WT; green) or Flag-Daed ΔSAM rescued cells

(∆SAM; red) (n=5). Values relative to the control (EGFP KD) were normalized to *rp49*

and presented as mean values ±SE **: *p* <0.01; n.s.: *p* >0.05 [Wilcoxon rank sum exact

test (two-sided)]. (**H**) Heatmaps showing the frequency of the 3′ end positions (red)

corresponding to each 5′ end of piRNAs in Daed WT (upper) and ∆SAM (lower) libraries.

The 50,000 piRNA 5′ end positions common to both libraries and most abundant in the

WT library were used for analysis. (**I**) Sequence logos. piRNAs were sorted by their

lengths and aligned according to the 3′ end of the piRNAs (black arrowhead) and its

immediately downstream nucleotide is highlighted in green. The U depletion found in 28–

33-nt piRNAs in the WT library and the equivalent regions in ∆SAM are highlighted in

yellow.

Figure 5. Daed is Diptera-specific and controls Zuc cleavage in a Nbr-independent

manner

(**A**) Protein conservation state of Armi, Gasz, Daed, PNLDC1, Papi, and Nbr in *D.*

melanogaster and other species. (**B**, **C**) Immunoprecipitation (Flag IP) and western

blotting show Myc-ASZ1 binds to MOV10L1 in normal (**B**) and Daed-depleted OSCs (**C**).

Luc-Flag: negative control. (**D**) Subcellular localization of Myc-ASZ1 alone or MOV10L1-

Flag (green) with Myc-ASZ1 (yellow) in normal (left two row) and Daed-depleted OSCs

(right two row). Mitochondria (mito) are shown in magenta. Images in the same vertical

row are identical cells. DAPI (blue): nuclei. Scale bar: 5 μm. (**E**, **F**) The efficiency of Papi

KD (upper) and Nbr KD (lower) (n=3). Piwi-bound piRNAs in control and Papi-KD (upper)

and Nbr-KD OSCs (lower). The amount of Piwi in each sample is also shown. (**G**) Model

proposed in this study. Diptera species acquired Daed instead of PNLDC1/PARN during

evolution; Daed and Gasz form AmTEC, which stably tethers Armi to the mitochondrial

surface via two sites. This allows Zuc to define the length of piRNAs without relying on

exonucleases such as PNLDC1/PARN or Nbr. In contrast, in mice, MOV10L1/Armi is

tethered solely by ASZ1/Gasz. In this case, Zuc cleavage does not specify piRNA length;

therefore, trimming by exonucleases such as PNLDC1/PARN is required after Zuc

processing. Blue arrowhead: Zuc cleavage.

STAR★Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal antibody Anti-Daed	This study	P3F10
Mouse monoclonal antibody anti-Gasz	Yamashiro et al.[12]	1H7
Mouse monoclonal antibody anti-Armi	Saito et al.[16]	2F8A9
Mouse monoclonal antibody anti-Piwi	Saito et al.[36]	3G11
Mouse monoclonal antibody anti-Piwi	Saito et al.[36]	4D2
Mouse monoclonal antibody anti-Yb	Murota et al.[56]	1G2H3
Mouse monoclonal antibody anti-Ago1	Okamura et al.[57]	1B8
Mouse monoclonal antibody anti-β-Tub	Developmental Studies Hybridoma Bank	Cat#E7; RRID: AB_528499
HSP60 antibody, Clone LK1	StressMarq Biosciences	Cat# SMC-110 (also SMC-110B), RRID:AB_2121277
Anti-Histon H3 antibody – Nuclear Loading Control and ChIP grade	abcam	Cat# ab1791, RRID:AB_302613
Monoclonal anti-Flag M2 antibody	Sigma-Aldrich	Cat# F3165, RRID:AB_259529
Anti-Flag antibody produces in rabbit	Sigma-Aldrich	Cat# F7425, RRID:AB_439687
Anti-c-Myc antibody produces in rabbit	Sigma-Aldrich	Cat# C3956, RRID:AB_439680
anti-mouse IgG (H+L) HRP	Cappel	55558
anti-rabbit IgG HRP-linked	Cell Signaling Technology	Cat#7074, RRID: AB_2099233
Alexa Fluor 488 goat anti-mouse IgG1 antibody	Thermo Fisher Scientific	Cat#A-21121; RRID: AB_2535764

Alexa Fluor 488 goat anti-mouse IgG2b antibody	Thermo Fisher Scientific	Cat# A-21141 (also A21141), RRID:AB_141626
Alexa Fluor 555 goat anti-mouse IgG1 antibody	Thermo Fisher Scientific	Cat# A-21127, RRID: AB_2535769
Alexa Fluor 555 goat anti-mouse IgG2b antibody	Thermo Fisher Scientific	Cat# A21147, RRID:AB_1500897
Alexa Fluor 488 goat anti-rabbit IgG (H+L) antibody	Thermo Fisher Scientific	Cat# A-11008, RRID:AB_143165
Alexa Fluor 633 goat anti-rabbit IgG (H+L) antibody	Thermo Fisher Scientific	Cat# A-21070 (also A21070), RRID:AB_2535731
Bacterial and virus strains		
Escherichia coli Rosetta 2 (DE3)	Novagen	Cat#71397
Chemicals, peptides, and recombinant proteins		
Shields and Sang M3 Insect Medium w/ L-Glutamine	USBiological	Cat#S1013
Fetal Bovine Serum	NICHIREI	Cat#173012
NEBuilder HiFi DNA Assembly Master Mix	New England Biolabs	Cat#E2621X
Ligation high ver.2	Toyobo	Cat#LGK-201
Glutathione Sepharose 4B	Cytiva	Cat#17-0756-01
Phosphate-buffered saline	FUJIFILM Wako Pure Chemical	Cat#048-29805
ClearTrans SP PVDF membranes	Fujifilm Wako Pure Chemical	Cat#033-22453
Clarity Western ECL Substrate	Bio-Rad	Cat#170506
Dynabeads Protein G	Thermo Fisher Scientific	Cat#10009D
RNasin Plus Ribonuclease Inhibitor	Promega	Cat#N2615
T4 Polynucleotide Kinase	New England Biolabs	Cat#M0201L
ISOGEN II	FUJIFILM Wako Pure Chemical	Cat#311-07361
Hybond-N membrane	Cytiva	Cat#RPN303N
VECTASHIELD Antifade Mounting Medium with DAPI	VECTOR LABORATORIES	Cat#H-1200
MitoTracker Orange CMXRos	Thermo Fisher Scientific	Cat#M-7510
SYBR Gold Nucleic Acid Gel Stain	Thermo Fisher Scientific	Cat#S11494
TURBO DNase	Life Technologies	Cat#AM2239

ReverTra Ace qPCR RT master mix	Toyobo	Cat#TYFSQ201
cOmplete ULTRA Tablets, EDTA-free, Protease Inhibitor Cocktail	Roche	Cat#6538282001
RNase A	Roche	Cat#10109142001
Critical commercial assays		
Nucleofector 96-well Shuttle device	Lonza	Cat#AAF-1003S
Nucleofector 2b device	Lonza	Cat#AAB-1001
ChemiDoc XRS Plus System	Bio-Rad	Cat#1708265J1PC
LSM980 laser scanning confocal microscope	Carl Zeiss	Cat#LSM980
StepOnePlus Real-Time PCR system	Thermo Fisher Scientific	Cat#4376598
MiSeq	Illumina	Cat#SY-410-1003
SF Cell Line 96-well Nucleofector Kit	Lonza	Cat#V4SC-3096
QIAseq FastSelect-rRNA Fly Kit	QIAGEN	Cat#333262
PowerUp SYBR Green Master Mix	Thermo Fisher Scientific	Cat# A25743
MiSeq Reagent Kit v3	Illumina	Cat#MS-102-3001
Deposited data		
Raw sequencing data	This paper	GEO: GSExxxxxx
Experimental models: Cell lines		
OSC	Saito et al.[34]	N/A
Oligonucleotides		
Primer sequences	Table S1	N/A
siRNA sequences	Table S2	N/A
Probe sequences	Table S3	N/A
qPCR primer sequences	Table S4	N/A
Recombinant DNA		
pGEX	Cytiva	Cat#28-9546-48
pAc-Flag-Daed WT and mutants	This study	N/A
pAc-Flag-Armi WT	Ishizu et al.[51]	N/A
pAc-Flag-Gasz WT	Yamashiro et al.[12]	N/A
pAc-Flag-Gasz mutants	This study	N/A
pAc-Armi WT-Flag	Yamashiro et al.[12]	N/A
pAc-Armi ∆N34-Flag	Yamashiro et al.[12]	N/A
pAc-Armi KEE-Flag	This study	N/A
pAc-Myc-Luc	gift from H. Ishizu	N/A
pAc- Luc-Flag	Yamashiro et al.[12]	N/A

pAc-Flag-Luc	This study	N/A
pAc-MOV10L1-Flag	This study	N/A
pAc-Myc-ASZ1	This study	N/A
Software and algorithms		
Python (Ver. 3.8.8)	Van Rossum & Drake[58]	RRID:SCR_008394;https://www.python.org
Numpy (Ver. 1.20.1)	Harris et al.[59]	RRID:SCR_008633; https://numpy.org/ja/
Scipy (Ver. 1.6.2)	Virtanen et al.[60]	RRID:SCR_008058; https://scipy.org
Pandas (Ver. 1.2.4)	McKinney[61]	RRID:SCR_018214;https://pandas.pydata.org
R (Ver. 4.3.1)	R Development Core Team[62]	RRID:SCR_001905;https://www.r-project.org
Cutadapt (Ver. 4.0)	Martin[63]	RRID:SCR_011841;https://cutadapt.readthedocs.io/en/stable/
STAR (Ver.2.7.8a)	Dobin et al.[64]	RRID:SCR_004463;https://github.com/alexdobin/STAR
HTSeq (Ver. 0.9.1)	Anders et al.[65]	RRID:SCR_005514;https://htseq.readthedocs.io/en/latest/
bedtools (Ver. 2.30.0)	Quinlan & Hall[66]	RRID:SCR_006646;https://bedtools.readthedocs.io/en/latest/
SAMtools (Ver. 1.9)	Li et al.[67]	RRID:SCR_002105;http://www.htslib.org
deepTools (Ver. 3.3.2)	Ramírez et al.[68]	RRID:SCR_016366;https://deeptools.readthedocs.io/en/develop/
wigToBigWig (Ver. 357)	UCSC	https://www.encodeproject.org/software/wigtobigwig/
ggseqlogo	Wagih.[69]	https://omarwagih.github.io/ggseqlogo/
RSeQC (Ver. 2.6.4)	Wang et al.[70]	RRID:SCR_005275;https://rseqc.sourceforge.net
Fiji (Ver. 2.0.0-rc-69/1.52p)	Schneider et al.[71]	RRID:SCR_002285; https://fiji.sc
seaborn (Ver. 0.12.2)	Waskom et al.[72]	https://seaborn.pydata.org
AlphaFold (Ver. 2.3)	Jumper et al.[32]	https://github.com/google-deepmind/alphafold

UCSF ChimeraX (Ver. 1.6.1)	Pettersen et al.[73]	RRID:SCR_015872;https://www.cgl.ucsf.edu/chimerax/
ISOLDE (Ver. 1.6)	Croll[74]	https://tristanic.github.io/isolde/
CHARMM-GUI Martini bilayer maker (Ver. 3.8)	Jo et al.[75]; Lee et al.[76]; Qi et al.[77]	RRID:SCR_025037;https://www.charmm-gui.org
PPM 3.0 Web Server	Lomize et al.[78]	https://opm.phar.umich.edu/ppm_server
PyMOL (Ver. 2.5.0)	Schrödinger & DeLano	RRID:SCR_000305;http://www.pymol.org/
GROMACS (Version 2021.2)	Abraham et al.[79]	RRID:SCR_014565;http://www.gromacs.org
MDAnalysis (Ver. 2.7.0)	Michaud-Agrawal et al.[79]; Gowers et al. [80]	https://mdanalysis.readthedocs.io/en/latest/index.html
CLUSTAL Omega	Madeira et al.[82]	RRID:SCR_001591;https://www.ebi.ac.uk/jdispatcher/msa/clustalo
iTOL (Ver. 6.9)	Letunic & Bork[83]	RRID:SCR_018174;https://itol.embl.de
ESPript (Ver. 3.0)	Robert & Gouet[84]	https://espript.ibcp.fr
MEGA11	Tamura et al.[85]	RRID:SCR_023017;https://www.megasoftware.net/

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Mikiko C. Siomi (siomim@bs.s.u-tokyo.ac.jp).

Materials availability

Plasmids and antibodies generated in this study will be shared by the lead contact upon

request.

Data and code availability

The data supporting the findings of this study are available from the corresponding

authors upon reasonable request. The datasets produced in this study are available in

the following databases: RNA-Seq data: Gene Expression Omnibus GSExxxxxx

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSExxxxxx).

Experimental model and study participant details

OSCs were cultured at 26°C in Shields and Sang M3 medium (USBiological)

supplemented with 10% fetal bovine serum (FBS, NICHIREI), 0.6 mg/mL glutathione, 10

mU/mL insulin, and 10% fly extract.[8]

Method details

Plasmid construction

The Daed coding region was amplified from total OSC RNAs by RT-PCR and inserted

into pAc-Flag-Armi WT[51] by substituting the Armi coding region using NEBuilder HiFi

DNA Assembly Master Mix (New England Biolabs). RNAi-resistant Daed WT and its

ΔSAM and ΔCC mutants were generated by inverse PCR, according to the domain

structures described in a previous study.[29] To construct the expression vector for GST-

Daed ΔTM-His, corresponding coding regions of Daed were cloned into pGEX, after

which a His-Tag was added to the C-terminal by inverse PCR. RNAi-resistant Flag-Gasz

mutants (ΔAR1-3, ΔAR4-6, ΔSAM, and Δα-H2) were generated by inverse PCR using

RNAi-resistant Flag-Gasz WT vector[12] as a template. Vectors for RNAi-resistant Armi

WT-Flag and its ΔN34 mutant were previously constructed.[12] The Armi KEE-Flag mutant

was produced by inverse PCR using RNAi-resistant Armi WT-Flag vector as a template.

The Flag-Luc expression vector was generated from pAc-Myc-Luc (a gift from H. Ishizu)

by substituting Myc with 3×Flag. The MOV10L1 and ASZ1 coding region was amplified

from total mouse ovary RNAs by RT-PCR and inserted into pAc-Armi WT-Flag and Myc-

Daed WT by substituting original coding regions. The primers used are shown in Table

S1.

RNAi and transfection

To initiate RNAi, 5.0×10^6 OSCs were mixed with 200 pmol of siRNAs in 20 μL of Solution

SF of the Cell Line Nucleofector Kit SF (Lonza Bioscience) and electroporation was

performed using a Nucleofector 96-well Shuttle device (Lonza Bioscience). For plasmid

transfection or plasmid/siRNA co-transfection, 1.5×10^7 OSCs were mixed with 1.5–10

μg of plasmids with or without 600 pmol of siRNAs in 100 μL of transfection buffer [180

mM sodium phosphate buffer for Church and Gilbert hybridization (pH 7.2), 5 mM KCl,

15 mM $MgCl_2$, and 50 mM D-mannitol] modified from a previous report,[86] and

electroporation was performed by a Nucleofector 2b device (Lonza Bioscience). Cells

were transfected with siRNAs twice with a 48-h interval. For rescue assays, endogenous

genes were depleted by RNAi and plasmids and siRNAs were introduced into the cells

after 48 h of culture. Cells were harvested after another 48 h of culture. For

immunofluorescence in Figure 2F, 500 pmol of siRNAs were used for RNAi and

plasmid/siRNA co-transfection, and cells were harvested after another 72 h of culture.

Sequences of siRNAs are indicated in Table S2.

Production of anti-Daed monoclonal antibody

GST-Daed (Met1–Asp197) was expressed in *E. coli* (Rosetta-DE3) and the inclusion

bodies were used for mouse immunization. Production and selection of hybridomas that

produce anti-Daed monoclonal antibodies were performed as described previously.[87]

Mitochondrial fractionation

Mitochondria were fractionated as previously reported.[12] Briefly, 1.2×10^8 OSCs were

suspended in 500 μL of buffer A [30 mM Tris-HCl (pH 7.3), 225 mM D-mannitol, 75 mM

sucrose, and 0.05 mM EGTA] and the cell suspension was passed 30 times through a

30-gauge needle attached to a 1-mL syringe. The lysates were centrifuged at 600 *g* for

5 min and the supernatants were collected, and repeated twice. The lysates were then

centrifuged at 7,000 *g* for 10 min and the supernatants were collected as the cytoplasmic

fraction. This step was also repeated twice. The pellets were resuspended in 500 μL of

buffer B [30 mM Tris-HCl (pH 7.3), 225 mM D-mannitol, and 75 mM sucrose] and

centrifuged at 10,000 *g* for 10 min. The supernatant was used as the mitochondrial

fraction.

Nuclear fractionation

Nuclear fractionation was performed as previously reported with some modifications. In

brief, cells were suspended in hypotonic buffer [10 mM HEPES-KOH (pH 7.3), 10 mM

KCl, 1.5 mM $MgCl_2$, 0.5 mM dithiothreitol (DTT), 2 μg/mL leupeptin, 2 μg/mL pepstatin

A, and 0.5% aprotinin] by pipetting six times and passed 3 times through a 25-gauge

needle attached to a 1-mL syringe. The lysates were centrifuged at 400 *g* for 10 min to

collect nuclei. The supernatants were collected as the cytoplasmic fraction, then the

concentration of KCl was adjusted to 200 mM and centrifuged at 20,000 *g* for 20 min

before used for Piwi-piRISC isolation (see below). The nuclear fraction was washed with

hypotonic buffer and resuspended in chromatin co-immunoprecipitation buffer [50 mM

HEPES-KOH (pH 7.3), 200 mM KCl, 1 mM EDTA, 1% Triton X-100, 0.1% Na

deoxycholate, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]. The nuclear

fraction was sonicated and centrifuged at 20,000 *g* for 20 min. The supernatant was

collected as nuclear extract fraction for Piwi-piRISC isolation.

Western blotting

Western blotting was performed essentially as described previously.[88] Proteins

separated on SDS polyacrylamide gels were transferred to ClearTrans SP PVDF

membranes (Fujifilm Wako Pure Chemical) and incubated in PBS containing 0.1%

Tween-20 (T-PBS) supplemented with 5% skimmed milk. After washing with T-PBS, the

membranes were incubated with the following primary antibodies: anti-Daed (1:1,000

dilution, this study), anti-Gasz (1:1,000 dilution),[12] anti-Armi (2F8A9, 1:1,000 dilution),[16]

anti-Piwi (3G11, 1:1,000 dilution),[36] anti-β-Tub (1:1,000 dilution, E7, Developmental

Studies Hybridoma Bank), anti-HSP60(1:500 dilution, LK1, StressMarq Biosciences),

anti-Flag M2 (1:5,000 dilution, F3165, Sigma), anti-Histon H3 (1:5,000 dilution, ab1791,

abcam) monoclonal and anti-c-Myc (1:1,000 dilution, C3956, Sigma) polycronal

antibodies. After washing with T-PBS, the membranes were incubated with peroxidase-

conjugated anti-mouse IgG antibodies (1:5,000 dilution, 55558, Cappel) or anti-rabbit

IgG antibodies (1:5,000 dilution, 7074, Cell Signaling Technology). The membranes

were then reacted with Clarity Western ECL Substrate (Bio-Rad) and images were

collected using ChemiDoc XRS Plus System (Bio-Rad). The signal intensity was

quantified by Fiji (Ver. 2.0.0-rc-69/1.52p).[71]

Immunofluorescence

Immunofluorescence was performed as described previously.[16] OSCs were fixed with

4% formaldehyde in PBS. The cells were permeabilized with PBS containing 0.1% Triton

X-100, blocked with PBS containing 3% BSA, and incubated with the following primary

antibodies: anti-Armi (2F8A9, 1:200 dilution),[16] anti-Piwi (4D2, 1:200 dilution,[36] anti-Yb

(1G2H3, 1:200 dilution),[56] anti-Flag M2 (F3165, 1:1,000 dilution, Sigma) monoclonal and

anti-c-Myc (1:300 dilution, C3956, Sigma) polyclonal antibodies. Anti-Flag polyclonal

antibodies (F7425, 1:300 dilution, Sigma) were used to obtain the results shown in Figure

2F. After washing with PBS, cells were incubated with the following secondary antibodies

(1:1,000 dilution): Alexa 488-conjugated anti-mouse IgG1, Alexa 488-conjugated anti-

mouse IgG2b, Alexa 555-conjugated anti-mouse IgG1, Alexa 555-conjugated anti-

mouse IgG2b, Alexa 488-conjugated anti-rabbit IgG and Alexa 633-conjugated anti-

rabbit IgG (all from Thermo Fisher Scientific). Mitochondria were stained with

MitoTracker Orange CMXRos (Thermo Fisher Scientific) before fixation. Cells were

mounted with VECTASHIELD Mounting Medium with DAPI (Vector Laboratories).

Images were obtained using LSM980 laser scanning confocal microscope (Carl Zeiss).

Immunoprecipitation

Immunoprecipitation was performed as described previously.[16] Cells were suspended in

binding buffer A [30 mM HEPES-KOH (pH 7.3), 150 mM KOAc, 5 mM Mg(OAc)$_2$, 5 mM

DTT, 1% Triton X-100, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]

supplemented with 0.04 U/μL RNasin Plus RNase Inhibitor (Promega) and the cell

suspension was passed 10 times through a 30-gauge needle and centrifuged. The

obtained supernatants (cell lysates) were incubated at 4°C with Dynabeads Protein G

(Thermo Fisher Scientific) pre-bound with antibodies. After 1 h of incubation, the beads

were washed with binding buffer and the materials bound on the beads were analyzed.

Piwi-piRISC isolation from OSCs

Cells were lysed by sonication in binding buffer B [50 mM HEPES-KOH (pH 7.3), 200

mM KCl, 1 mM EDTA, 1 mM DTT, 1% Triton X-100, 0.1% Na deoxycholate, 2 μg/mL

leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin] supplemented with 0.1 U/μL RNasin

Plus RNase Inhibitor. Cell lysates were then incubated with anti-Piwi monoclonal

antibodies (3G11) pre-bound to Dynabeads Protein G at 4°C for 2 h. The beads were

washed with IP-wash buffer [20 mM HEPES-KOH (pH 7.3), 300 mM NaCl, 1 mM DTT,

0.05% NP-40, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin] and with

high-salt IP-wash buffer [20 mM HEPES-KOH (pH 7.3), 500 mM NaCl, 1 mM DTT, 0.05%

NP-40, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]. RNAs were eluted

from the beads with phenol-chloroform and precipitated with ethanol. Eluted RNAs were

separated on denaturing polyacrylamide gels and detected by SYBR Gold Nucleic Acid

Gel Stain (Thermo Fisher Scientific). Signal intensity was calculated using Fiji, and

displayed using Python3 and seaborn (Ver.0.12.2). For piRNA sequencing, cells were

lysed in binding buffer C [20 mM HEPES-KOH (pH 7.3), 150 mM NaCl, 1mM EDTA, 1

mM DTT, 0.5% NP-40, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]

supplemented with 0.04 U/μL RNasin Plus RNase Inhibitor. RNAs were separated on

denaturing polyacrylamide gels and 20–40-nt RNAs were eluted from the gels and

cloned as described previously.[8] QIAseq FastSelect-rRNA Fly Kit (QIAGEN) was used

to remove rRNA from the library.

Northern blotting

Total RNAs were isolated from OSCs using ISOGEN II (Fujifilm Wako Pure Chemical)

and treated with 0.04 U/μL TURBO DNase (Life Technologies). Northern blotting was

carried out as described previously.[89] Total RNAs were separated on denaturing

polyacrylamide gels, blotted on Hybond-N+ hybridization membranes (Cytiva), and UV

cross-linked. The membranes were incubated at 42°C for 30 min in hybridization buffer

[200 mM sodium phosphate buffer (pH 7.2), 7% SDS, and 1 mM EDTA] and probed with

^{32}P-labeled oligonucleotides at 42°C overnight. The membranes were washed with NB-

wash buffer [30 mM sodium citrate (pH 7.0), 300 mM NaCl, and 0.1% SDS] at 42°C and

scanned using Typhoon FLA 9500 (Cytiva). Sequences of oligodeoxynucleotides are

shown in Table S3.

RT-qPCR

Total RNAs from OSCs were reverse-transcribed using ReverTra Ace (Toyobo).

Resulting cDNAs were amplified with StepOnePlus (Thermo Fisher Scientific) using

PowerUp SYBR Green Master Mix (Thermo Fisher Scientific). Sequences of primers are

shown in Table S4.

Recombinant proteins

GST-Daed ΔTM-His was expressed in *E. coli* (Rosetta DE3) at 16°C overnight. *E. coli*

was harvested and lysed by sonication in lysis buffer [30 mM Tris-HCl (pH 8.0), 500 mM

NaCl, 1% Triton X-100, 10 mM imidazole, 1× cOmplete ULTRA Tablets (Roche), and 1

mM DTT]. The supernatants after centrifugation were bound to Ni-Sepharose 6 Fast

Flow (Cytiva) at 4°C for 1 h. After washing with wash buffer A [30 mM Tris-HCl (pH 8.0),

500 mM NaCl, 1% Triton X-100, 10 mM imidazole, and 1 mM DTT], proteins were eluted

from the beads with elution buffer A [30 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% Triton

X-100, 250 mM imidazole, and 1 mM DTT]. Eluted proteins were bound to Glutathione

Sepharose 4B (Cytiva) at 4°C for 1 h. After washing with wash buffer B [30 mM Tris-HCl

(pH 8.0), 500 mM NaCl, 1% Triton X-100, and 1 mM DTT], proteins were eluted from the

beads with elution buffer B [30 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% Triton X-100, 1

mM DTT, and 15 mM glutathione], dialyzed in dialysis buffer [30 mM Tris-HCl (pH 8.0),

150 mM NaCl, 1 mM DTT, 10% glycerol, and 8 μg/mL GST-HRV3C protease], and

passed through Glutathione Sepharose 4B to remove the protease and GST.

Pull-down assay

OSCs expressing Armi WT-Flag and its ΔN34 mutant were suspended with bait-

preparing buffer [30 mM HEPES-KOH (pH 7.3), 150 mM KOAc, 500 mM NaCl, 5 mM

Mg(OAc)$_2$, 5 mM DTT, 1% Triton X-100, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, 0.5%

aprotinin, and 25 μg/mL RNase A (Roche)]. The cell suspension was passed five times

through a 25-gauge needle and another five times through a 30-gauge needle, and then

centrifuged. The supernatants were incubated at 4°C for 1 h with anti-Flag antibodies

pre-bound onto Dynabeads Protein G. Beads were washed with bait-preparing buffer

and then binding buffer D [30 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM DTT, and 1%

Triton X-100]. Daed ΔTM-His was added to the beads and incubated at 4°C for 1.5 h.

After washing with binding buffer, proteins were eluted from the beads, separated on an

SDS polyacrylamide gel, and stained with CBB.

β-**elimination assay**

β-elimination was performed as described previously with some modifications.[37] piRNAs

were isolated from Piwi immunoprecipitated from OSCs. miRNAs were isolated from

Ago1 immunoprecipitated from OSCs using monoclonal anti-Ago1 antibodies[57]. RNAs

were incubated in 10 mM NaIO$_4$ for 40 min at 0°C in the dark. After ethanol precipitation,

RNAs were dissolved in 1 M L(+)-lysine-HCl (pH 8.5) and incubated for 90 min at 45°C.

After ethanol precipitation, RNAs were separated on a denaturing polyacrylamide gel

and detected by SYBR Gold Nucleic Acid Gel Stain.

piRNA sequencing and bioinformatic analysis

piRNAs were sequenced using MiSeq (Illumina) with MiSeq Reagent Kit v3 (Illumina).

piRNA-seq analysis was performed as described previously[8] with some modifications.

Adapter-trimmed reads of ≥ 18 nt in length were mapped against the *Drosophila* genome

(BDGP Release 6 + ISO1 MT/dm6) with FlyBase gene model (Release 6.21) using

STAR[64] (Ver. 2.7.8a). Uniquely mapped reads with up to one mismatch were used for

the subsequent analysis. Read length of each biological triplicate was used in the

analysis in Figures 4B and S6A. After confirming high correlations between samples

(Figures S9A and S9B), reads were merged and used for subsequent analysis. Reads

mapped to exons of coding genes (Figures S9A and S9B) were counted with HTSeq[65]

(Ver. 0.9.1) using annotation file obtained from FlyBase and normalized to RPM+1 reads.

Spearman's rho was calculated using R[62] (Ver. 4.3.1). For transposon annotation

(Figures S4B and S4C), the annotation file of repeat sequences was obtained from

RepeatMasker (dm6-Apr 2006-RepeatMasker open-4.0.6-Dfam 2.0) and transposons

were extracted without distinction between int and LTR. Reads mapped antisense to

transposons were counted using HTSeq. For the analysis in Figure S4C, reads were

normalized to RPM+1 reads. For the analysis in Figure S4B, transposons which mapped

> 5 RPM reads in the sum of WT and ΔSAM were selected, and the mean length of

mapped reads for each transposon were calculated. For 1U bias and +1U bias analyses

(Figures 4C, 4I, and S6B), the 5′ ends and their downstream sequences of piRNAs (1U

bias) or *Drosophila* genome sequences around the 3′ ends (+1U bias) were obtained

using bedtools[66] (Ver. 2.30.0) and SAMtools (Ver. 1.9).[67] Sequence logos were

generated using ggseqlogo.[69] For the phasing analysis (Figure 4D), 3′ end positions

around piRNA 5′ end position were normalized to RPM and showed in heatmaps using

RSeQC[70] (Ver. 2.6.4), deepTools[68] (Ver. 3.3.2), SAMtools[67] (Ver. 1.9) and "wigToBigWig"

from UCSC. Binary histograms were generated using the matrix of heatmaps. Values

were transformed into binary format (counts = 0 → 0, counts > 0 → 1) in accordance

with a previous study.[18] For the comparison of 3′ end position in WT and mutant libraries

of Daed or Armi (Figures 4H and S6C), 50,000 piRNA 5′ end positions, common to the

WT and mutant libraries and most abundant in the WT library, were extracted. The

percentage of 3′ end positions corresponding to each 5′ end position was calculated and

shown in heatmaps. 5′ end positions were ordered by the 3′ end position of WT library

data and the same order was maintained in the mutant library data. In Figure S4G, the

length of dominant piRNAs produced from the 5′ end was used for the 5′-end clarification.

First, the 5′ end positions that dominantly produce 26- or 27-nt piRNAs in the WT library

was extracted, and after combining, these positions were classified as "Unchanged" and

"Elongated" according to the length of dominant piRNAs in ΔSAM library. The

subsequent analysis was conducted using the piRNAs produced from each 5′-end

groups.

piRISC production simulation

The *flam* sequence was downloaded from FlyBase (Release FB2023_05) (21,629,891

to 21,792,731 on the X chromosome) and subjected to the digestion into 163 piRNA

precursors (starting at U and around 1,000 nt). The precursors were cleaved to 18–40-

nt piRNAs immediately before U. The cleavage position was determined based on the

distribution of the actual piRNA length, which was obtained by fitting a normal distribution

to the piRNA sequencing data. The simulation was performed using Python3[58] (Ver.

3.8.8) with add-on libraries [Numpy[59] (Ver. 1.20.1), Scipy[60] (Ver. 1.6.2), and Pandas[61]

(Ver. 1.2.4)].

Structure prediction

A structure model of the AmTEC (Figure 1A) and Armi–AmTEC complex (Figure 3A) was

built by structure prediction with ColabFold[90] (Ver. 1.5.5) and AlphaFold[32] (Ver. 2.3) using

protein amino acid sequences from FlyBase (Release FB2023_04). Five complex

models were built, and structural predictions and optimization were performed five times

for each complex model. The structure model with the highest pLDDT value underwent

relax steps. The N-terminal disorder region and TM domain orientations of Daed and

Gasz were appropriately changed at the linker region using ISOLDE[74] (Ver. 1.6). The

structure was visualized using UCSF ChimeraX[73] (Ver. 1.6.1).

Molecular dynamics simulation

The protein complex was embedded in a bilayer membrane in the simulation box

described below using CHARMM-GUI Martini bilayer maker[75–77] (Ver. 3.8). Orientation

of the protein complex in lipid bilayer was determined using PPM 3.0 Web Server[78] prior

to the construction of the simulation box. The structure model of the Armi–AmTEC [Daed

(ΔSAM)] complex was generated using PyMOL (2.5.0), UCSF ChimeraX, and ISOLDE

based on the Armi–AmTEC complex model. Initial structures were arranged in the

250×250 nm bilayer membrane (67% of POPC and 33% DOPE, containing roughly

2,000 lipid molecules in total) and the thickness of water at the top and bottom of the

system was set to 100 nm. Ionic strength and temperature were set to 150 mM NaCl and

303.15 K, respectively. Molecular dynamics simulations were performed using

GROMACS[79] (Version 2021.2) with Martini 3 coarse-grained model.[91] First, energy

minimization was performed using the steepest-descent method. Then, equilibration was

conducted using the V-rescale method five times (2 fs×500,000 steps, 5 fs×200,000

steps, 10 fs×100,000 steps, 15 fs×50,000 steps, and 20 fs×50,000 steps). After the

equilibration runs, simulations were performed at 20 fs for 50,000,000 steps. The

structural trajectories were calculated using gmx trjconv tool for clustering and centering

of the protein complex. The root mean square deviation (RMSD) of protein complexes

were calculated using gmx rms tool. The distances between the center of mass of Armi

helicase domain (Ser697–Cys1,148) and that of AmTEC were calculated using Python3

(Ver. 3.9.18) with add-on libraries [Numpy (Ver. 1.24.3) and MDAnalysis[80,81] (Ver. 2.7.0)],

and visualized using R.[62]

Ortholog search and protein sequence alignment

Orthologs for Armi/MOV10L1, Gasz/ASZ1, PNLDC1, Papi/TDRKH, Nbr were searched

using OrthoDB v11. Orthologs for Daed were searched using OrthoDB v11[39] and NCBI

PSI-Blast. Species taxonomy was according to NCBI and phylogenetic tree (Figure 5A)

were generated following previous studies.[92,93] Multiple sequence alignment (Figure

S5B) was conducted using CLUSTAL Omega multiple alignment[82] and visualized using

ESPript (Ver. 3.0).[84] The phylogenetic tree (Figure S5A) was generated using Neighbor-

Joining method[94] in MEGA11[85].   Multiple sequence alignment (Figure S6) was

conducted using CLUSTAL Omega multiple alignment[82] and visualized using iTOL (Ver.

6.9)[83]. The following protein sequences were acquired from NCBI: *D. melanogaster* Armi

(NP_001014556), *H. sapiens* Moloney leukemia virus 10-like protein 1 (MOV10L1)

(NP_061868.1), *M. musculus* MOV10L1 (XP_006521619.1), *G. gallus* MOV10L1

(XP_040514825.1), *X. tropicalis* MOV10L1 (NP_001072624.1), *D. rerio* MOV10L1

(NP_001070795.1), *Z. nevadensis* Armi (XP_021937793.1), *T. palmi* Armi

(XP_034251738.1), *A. echinatior* Armi (XP_011062616.1), *B. terrestris* Armi

(XP_012166280.1), *D. ponderosae* Armi (XP_048525683.1), *T. castaneum* Armi

(XP_008199251.1), *B. anynana* Armi (XP_023946786.2), *B. mori* Armi

(XP_037867032.1), *A. gambiae* Armi (XP_061506580.1), *A. aegypti* Armi

(XP_021698921.1), *B. dorsalis* Armi (XP_011214121.1), *C. capitata* Armi

(XP_004537921.1), *M. domestica* Armi (XP_058975516.1) and *S. calcitrans* Armi

(XP_013110641.2).

Quantification and statistical analysis

All experiments were performed over two independent experiments (biological

replicates). Sample sizes were predetermined following previously performed

experiments with similar setup in the field. Statistical methods, *p* value and number of

replicates are indicated in each figure and its legend. Statistical analyses were performed

in the R software environment (version 4.3.1). Shapiro-Wilk test was used for testing

normality when statistical methods based on the normal distribution was used.

Experiments were not blinded as consistent with the previous studies in the field.

Supplemental information

Document S1. Figures S1–S9 and Table S1–S4

Supplemental figure titles

Figure S1. Related to Figure 1; Coiled-coil prediction and specificity of anti-Daed

antibodies

Figure S2. Related to Figure 2; Functions and behaviors of Daed and Gasz in OSCs

Figure S3. Related to Figure 3; Effects of Armi mutants on piRNA biogenesis

Figure S4. Related to Figure 4; Effects of Daed ∆SAM on piRNA biogenesis and Piwi-

piRISC function

Figure S5. Related to Figure 4; MD simulation of Armi–AmTEC complex

Figure S6. Related to Figure 4; Effects of Armi KEE on piRNA biogenesis

Figure S7. Related to Figure 5; Sequence alignment of Daed in species

Figure S8. Related to Figure 5; Sequence alignment of Armi UPF1-like helicase domain

in species

Figure S9. Related to STAR Method "piRNA sequencing and bioinformatic analysis";

Reproducibility of piRNA-seq

Supplemental table titles

Table S1. Related to STAR Method "Plasmid construction"; Sequences of primers used

for plasmid construction

Table S2. Related to STAR Method "RNAi and transfection"; Sequences of siRNAs

Table S3. Related to STAR Method "Northern blotting"; Sequences of oligos used for

northern blotting

Table S4. Related to STAR Method "RT-qPCR"; Sequences of primers used for RT-

qPCR

References

1. Czech, B., Munafò, M., Ciabrelli, F., Eastwood, E.L., Fabry, M.H., Kneuss, E., and
Hannon, G.J. (2018). piRNA-Guided Genome Defense: From Biogenesis to
Silencing. Annual Review of Genetics *52*, 131–157.
https://doi.org/10.1146/annurev-genet-120417-031441.
2. Huang, X., Fejes Tóth, K., and Aravin, A.A. (2017). piRNA Biogenesis in Drosophila
melanogaster. Trends in Genetics *33*, 882–894.
https://doi.org/10.1016/j.tig.2017.09.002.
3. Ozata, D.M., Gainetdinov, I., Zoch, A., O'Carroll, D., and Zamore, P.D. (2019). PIWI-
interacting RNAs: small RNAs with big functions. Nat Rev Genet *20*, 89–108.
https://doi.org/10.1038/s41576-018-0073-3.
4. Yamashiro, H., and Siomi, M.C. (2018). PIWI-Interacting RNA in Drosophila:
Biogenesis, Transposon Regulation, and Beyond. Chem Rev *118*, 4404–4421.
https://doi.org/10.1021/acs.chemrev.7b00393.
5. Brennecke, J., Aravin, A.A., Stark, A., Dus, M., Kellis, M., Sachidanandam, R., and
Hannon, G.J. (2007). Discrete Small RNA-Generating Loci as Master Regulators of
Transposon Activity in Drosophila. Cell *128*, 1089–1103.
https://doi.org/10.1016/j.cell.2007.01.043.
6. Malone, C.D., Brennecke, J., Dus, M., Stark, A., McCombie, W.R., Sachidanandam,
R., and Hannon, G.J. (2009). Specialized piRNA Pathways Act in Germline and
Somatic Tissues of the Drosophila Ovary. Cell *137*, 522–535.
https://doi.org/10.1016/j.cell.2009.03.040.
7. Goriaux, C., Desset, S., Renaud, Y., Vaury, C., and Brasset, E. (2014).
Transcriptional properties and splicing of the flamenco piRNA cluster. EMBO reports
15, 411–418. https://doi.org/10.1002/embr.201337898.
8. Hirakata, S., Ishizu, H., Fujita, A., Tomoe, Y., and Siomi, M.C. (2019). Requirements
for multivalent Yb body assembly in transposon silencing in Drosophila. EMBO Rep
20, e47708. https://doi.org/10.15252/embr.201947708.
9. Ishizu, H., Iwasaki, Y.W., Hirakata, S., Ozaki, H., Iwasaki, W., Siomi, H., and Siomi,
M.C. (2015). Somatic Primary piRNA Biogenesis Driven by cis-Acting RNA
Elements and trans-Acting Yb. Cell Reports *12*, 429–440.
https://doi.org/10.1016/j.celrep.2015.06.035.
10. Cora, E., Pandey, R.R., Xiol, J., Taylor, J., Sachidanandam, R., McCarthy, A.A., and

Pillai, R.S. (2014). The MID-PIWI module of Piwi proteins specifies nucleotide- and strand-biases of piRNAs. RNA *20*, 773–781. https://doi.org/10.1261/rna.044701.114.

11. Stein, C.B., Genzor, P., Mitra, S., Elchert, A.R., Ipsaro, J.J., Benner, L., Sobti, S., Su, Y., Hammell, M., Joshua-Tor, L., et al. (2019). Decoding the 5′ nucleotide bias of PIWI-interacting RNAs. Nat Commun *10*, 828. https://doi.org/10.1038/s41467-019-08803-z.

12. Yamashiro, H., Negishi, M., Kinoshita, T., Ishizu, H., Ohtani, H., and Siomi, M.C. (2020). Armitage determines Piwi–piRISC processing from precursor formation and quality control to inter-organelle translocation. EMBO reports *21*, e48769. https://doi.org/10.15252/embr.201948769.

13. Choi, S.-Y., Huang, P., Jenkins, G.M., Chan, D.C., Schiller, J., and Frohman, M.A. (2006). A common lipid links Mfn-mediated mitochondrial fusion and SNARE-regulated exocytosis. Nat Cell Biol *8*, 1255–1262. https://doi.org/10.1038/ncb1487.

14. Ipsaro, J.J., Haase, A.D., Knott, S.R., Joshua-Tor, L., and Hannon, G.J. (2012). The structural biochemistry of Zucchini implicates it as a nuclease in piRNA biogenesis. Nature *491*, 279–283. https://doi.org/10.1038/nature11502.

15. Nishimasu, H., Ishizu, H., Saito, K., Fukuhara, S., Kamatani, M.K., Bonnefond, L., Matsumoto, N., Nishizawa, T., Nakanaga, K., Aoki, J., et al. (2012). Structure and function of Zucchini endoribonuclease in piRNA biogenesis. Nature *491*, 284–287. https://doi.org/10.1038/nature11509.

16. Saito, K., Ishizu, H., Komai, M., Kotani, H., Kawamura, Y., Nishida, K.M., Siomi, H., and Siomi, M.C. (2010). Roles for the Yb body components Armitage and Yb in primary piRNA biogenesis in Drosophila. Genes Dev *24*, 2493–2498. https://doi.org/10.1101/gad.1989510.

17. Han, B.W., Wang, W., Li, C., Weng, Z., and Zamore, P.D. (2015). Noncoding RNA. piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production. Science *348*, 817–821. https://doi.org/10.1126/science.aaa1264.

18. Mohn, F., Handler, D., and Brennecke, J. (2015). piRNA-guided slicing specifies transcripts for Zucchini-dependent, phased piRNA biogenesis. Science *348*, 812–817. https://doi.org/10.1126/science.aaa1039.

19. Yashiro, R., Murota, Y., Nishida, K.M., Yamashiro, H., Fujii, K., Ogai, A., Yamanaka, S., Negishi, L., Siomi, H., and Siomi, M.C. (2018). Piwi Nuclear Localization and Its Regulatory Mechanism in Drosophila Ovarian Somatic Cells. Cell Reports *23*, 3647–3657. https://doi.org/10.1016/j.celrep.2018.05.051.

20. Dönertas, D., Sienski, G., and Brennecke, J. (2013). Drosophila Gtsf1 is an essential component of the Piwi-mediated transcriptional silencing complex. Genes Dev *27*,

1693–1705. https://doi.org/10.1101/gad.221150.113.

21. Fabry, M.H., Ciabrelli, F., Munafò, M., Eastwood, E.L., Kneuss, E., Falciatori, I.,
Falconio, F.A., Hannon, G.J., and Czech, B. (2019). piRNA-guided co-transcriptional
silencing coopts nuclear export factors. eLife *8*, e47999.
https://doi.org/10.7554/eLife.47999.

22. Murano, K., Iwasaki, Y.W., Ishizu, H., Mashiko, A., Shibuya, A., Kondo, S., Adachi,
S., Suzuki, S., Saito, K., Natsume, T., et al. (2019). Nuclear RNA export factor
variant initiates piRNA-guided co-transcriptional silencing. The EMBO Journal *38*,
e102870. https://doi.org/10.15252/embj.2019102870.

23. Ohtani, H., Iwasaki, Y.W., Shibuya, A., Siomi, H., Siomi, M.C., and Saito, K. (2013).
DmGTSF1 is necessary for Piwi–piRISC-mediated transcriptional transposon
silencing in the Drosophila ovary. Genes Dev *27*, 1656–1661.
https://doi.org/10.1101/gad.221515.113.

24. Onishi, R., Sato, K., Murano, K., Negishi, L., Siomi, H., and Siomi, M.C. (2020). Piwi
suppresses transcription of Brahma-dependent transposons via Maelstrom in
ovarian somatic cells. Science Advances *6*, eaaz7420.
https://doi.org/10.1126/sciadv.aaz7420.

25. Sienski, G., Dönertas, D., and Brennecke, J. (2012). Transcriptional silencing of
transposons by Piwi and maelstrom and its impact on chromatin state and gene
expression. Cell *151*, 964–980. https://doi.org/10.1016/j.cell.2012.10.040.

26. Sienski, G., Batki, J., Senti, K.-A., Dönertas, D., Tirian, L., Meixner, K., and
Brennecke, J. (2015). Silencio/CG9754 connects the Piwi–piRNA complex to the
cellular heterochromatin machinery. Genes Dev *29*, 2258–2271.
https://doi.org/10.1101/gad.271908.115.

27. Yu, Y., Gu, J., Jin, Y., Luo, Y., Preall, J.B., Ma, J., Czech, B., and Hannon, G.J.
(2015). Panoramix enforces piRNA-dependent cotranscriptional silencing. Science
350, 339–342. https://doi.org/10.1126/science.aab0700.

28. Czech, B., Preall, J.B., McGinn, J., and Hannon, G.J. (2013). A transcriptome-wide
RNAi screen in the Drosophila ovary reveals factors of the germline piRNA pathway.
Mol Cell *50*, 749–761. https://doi.org/10.1016/j.molcel.2013.04.007.

29. Munafò, M., Manelli, V., Falconio, F.A., Sawle, A., Kneuss, E., Eastwood, E.L., Seah,
J.W.E., Czech, B., and Hannon, G.J. (2019). Daedalus and Gasz recruit Armitage
to mitochondria, bringing piRNA precursors to the biogenesis machinery. Genes Dev.
33, 844–856. https://doi.org/10.1101/gad.325662.119.

30. Ge, D.T., Wang, W., Tipping, C., Gainetdinov, I., Weng, Z., and Zamore, P.D. (2019).
The RNA-Binding ATPase, Armitage, Couples piRNA Amplification in Nuage to

Phased piRNA Production on Mitochondria. Molecular Cell *74*, 982-995.e6.
https://doi.org/10.1016/j.molcel.2019.04.006.

31. Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A.,
Bates, R., Blackwell, S., Yim, J., et al. (2022). Protein complex prediction with
AlphaFold-Multimer. Preprint at bioRxiv, https://doi.org/10.1101/2021.10.04.463034
https://doi.org/10.1101/2021.10.04.463034.

32. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.
https://doi.org/10.1038/s41586-021-03819-2.

33. Madeo, G., Savojardo, C., Manfredi, M., Martelli, P.L., and Casadio, R. (2023).
CoCoNat: a novel method based on deep learning for coiled-coil prediction.
Bioinformatics *39*, btad495. https://doi.org/10.1093/bioinformatics/btad495.

34. Saito, K., Inagaki, S., Mituyama, T., Kawamura, Y., Ono, Y., Sakota, E., Kotani, H.,
Asai, K., Siomi, H., and Siomi, M.C. (2009). A regulatory circuit for piwi by the large
Maf gene traffic jam in Drosophila. Nature *461*, 1296–1299.
https://doi.org/10.1038/nature08501.

35. Olivieri, D., Sykora, M.M., Sachidanandam, R., Mechtler, K., and Brennecke, J.
(2010). An in vivo RNAi assay identifies major genetic and cellular requirements for
primary piRNA biogenesis in Drosophila. EMBO J *29*, 3301–3317.
https://doi.org/10.1038/emboj.2010.212.

36. Saito, K., Nishida, K.M., Mori, T., Kawamura, Y., Miyoshi, K., Nagami, T., Siomi, H.,
and Siomi, M.C. (2006). Specific association of Piwi with rasiRNAs derived from
retrotransposon and heterochromatic regions in the Drosophila genome. Genes Dev
20, 2214–2222. https://doi.org/10.1101/gad.1454806.

37. Gunawardane, L.S., Saito, K., Nishida, K.M., Miyoshi, K., Kawamura, Y., Nagami,
T., Siomi, H., and Siomi, M.C. (2007). A Slicer-Mediated Mechanism for Repeat-
Associated siRNA 5' End Formation in Drosophila. Science *315*, 1587–1590.
https://doi.org/10.1126/science.1140494.

38. Horwich, M.D., Li, C., Matranga, C., Vagin, V., Farley, G., Wang, P., and Zamore,
P.D. (2007). The Drosophila RNA Methyltransferase, DmHen1, Modifies Germline
piRNAs and Single-Stranded siRNAs in RISC. Current Biology *17*, 1265–1272.
https://doi.org/10.1016/j.cub.2007.06.030.

39. Kuznetsov, D., Tegenfeldt, F., Manni, M., Seppey, M., Berkeley, M., Kriventseva,
E.V., and Zdobnov, E.M. (2023). OrthoDB v11: annotation of orthologs in the widest
sampling of organismal diversity. Nucleic Acids Research *51*, D445–D451.

https://doi.org/10.1093/nar/gkac998.

40. Vourekas, A., Zheng, K., Fu, Q., Maragkakis, M., Alexiou, P., Ma, J., Pillai, R.S., Mourelatos, Z., and Wang, P.J. (2015). The RNA helicase MOV10L1 binds piRNA precursors to initiate piRNA processing. Genes Dev *29*, 617–629. https://doi.org/10.1101/gad.254631.114.

41. Zheng, K., Xiol, J., Reuter, M., Eckardt, S., Leu, N.A., McLaughlin, K.J., Stark, A., Sachidanandam, R., Pillai, R.S., and Wang, P.J. (2010). Mouse MOV10L1 associates with Piwi proteins and is an essential component of the Piwi-interacting RNA (piRNA) pathway. Proceedings of the National Academy of Sciences *107*, 11841–11846. https://doi.org/10.1073/pnas.1003953107.

42. Ikeda, S., Tanaka, K., Ohtani, R., Kanda, A., Sotomaru, Y., Kono, T., and Obata, Y. (2022). Disruption of piRNA machinery by deletion of ASZ1/GASZ results in the expression of aberrant chimeric transcripts in gonocytes. Journal of Reproduction and Development *68*, 125–136. https://doi.org/10.1262/jrd.2021-146.

43. Ding, D., Liu, J., Dong, K., Midic, U., Hess, R.A., Xie, H., Demireva, E.Y., and Chen, C. (2017). PNLDC1 is essential for piRNA 3′ end trimming and transposon silencing during spermatogenesis in mice. Nat Commun *8*, 819. https://doi.org/10.1038/s41467-017-00854-4.

44. Nishimura, T., Nagamori, I., Nakatani, T., Izumi, N., Tomari, Y., Kuramochi-Miyagawa, S., and Nakano, T. (2018). PNLDC1, mouse pre-piRNA Trimmer, is required for meiotic and post-meiotic male germ cell development. EMBO reports *19*, e44957. https://doi.org/10.15252/embr.201744957.

45. Hayashi, R., Schnabl, J., Handler, D., Mohn, F., Ameres, S.L., and Brennecke, J. (2016). Genetic and mechanistic diversity of piRNA 3' end formation. Nature *539*, 588–592. https://doi.org/10.1038/nature20162.

46. Honda, S., Kirino, Y., Maragkakis, M., Alexiou, P., Ohtaki, A., Murali, R., Mourelatos, Z., and Kirino, Y. (2013). Mitochondrial protein BmPAPI modulates the length of mature piRNAs. RNA *19*, 1405–1418. https://doi.org/10.1261/rna.040428.113.

47. Saxe, J.P., Chen, M., Zhao, H., and Lin, H. (2013). Tdrkh is essential for spermatogenesis and participates in primary piRNA biogenesis in the germline. The EMBO Journal *32*, 1869–1885. https://doi.org/10.1038/emboj.2013.121.

48. Feltzin, V.L., Khaladkar, M., Abe, M., Parisi, M., Hendriks, G.-J., Kim, J., and Bonini, N.M. (2015). The exonuclease Nibbler regulates age-associated traits and modulates piRNA length in Drosophila. Aging Cell *14*, 443–452. https://doi.org/10.1111/acel.12323.

49. Joosten, J., Overheul, G.J., Van Rij, R.P., and Miesen, P. (2021). Endogenous

piRNA-guided slicing triggers responder and trailer piRNA production from viral RNA
in Aedes aegypti mosquitoes. Nucleic Acids Research *49*, 8886–8899.
https://doi.org/10.1093/nar/gkab640.

50. Wang, H., Ma, Z., Niu, K., Xiao, Y., Wu, X., Pan, C., Zhao, Y., Wang, K., Zhang, Y.,
and Liu, N. (2016). Antagonistic roles of Nibbler and Hen1 in modulating piRNA 3′
ends in Drosophila. Development *143*, 530–539. https://doi.org/10.1242/dev.128116.

51. Ishizu, H., Kinoshita, T., Hirakata, S., Komatsuzaki, C., and Siomi, M.C. (2019).
Distinct and Collaborative Functions of Yb and Armitage in Transposon-Targeting
piRNA Biogenesis. Cell Reports *27*, 1822-1835.e8.
https://doi.org/10.1016/j.celrep.2019.04.029.

52. Lee, Y.C.G., and Langley, C.H. (2012). Long-Term and Short-Term Evolutionary
Impacts of Transposable Elements on Drosophila. Genetics *192*, 1411–1432.
https://doi.org/10.1534/genetics.112.145714.

53. Obbard, D.J., Gordon, K.H.J., Buck, A.H., and Jiggins, F.M. (2008). The evolution
of RNAi as a defence against viruses and transposable elements. Philosophical
Transactions of the Royal Society B: Biological Sciences *364*, 99–115.
https://doi.org/10.1098/rstb.2008.0168.

54. Parhad, S.S., and Theurkauf, W.E. (2019). Rapid evolution and conserved function
of the piRNA pathway. Open Biology *9*, 180181.
https://doi.org/10.1098/rsob.180181.

55. Gamez, S., Srivastav, S., Akbari, O.S., and Lau, N.C. (2020). Diverse Defenses: A
Perspective Comparing Dipteran Piwi-piRNA Pathways. Cells *9*, 2180.
https://doi.org/10.3390/cells9102180.

56. Murota, Y., Ishizu, H., Nakagawa, S., Iwasaki, Y.W., Shibata, S., Kamatani, M.K.,
Saito, K., Okano, H., Siomi, H., and Siomi, M.C. (2014). Yb integrates piRNA
intermediates and processing factors into perinuclear bodies to enhance piRISC
assembly. Cell Rep *8*, 103–113. https://doi.org/10.1016/j.celrep.2014.05.043.

57. Okamura, K., Ishizuka, A., Siomi, H., and Siomi, M.C. (2004). Distinct roles for
Argonaute proteins in small RNA-directed RNA cleavage pathways. Genes Dev *18*,
1655–1666. https://doi.org/10.1101/gad.1210204.

58. Van Rossum, G., and Drake, F.L. (2009). Python 3 Reference Manual
(CreateSpace).

59. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P.,
Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array
programming with NumPy. Nature *585*, 357–362. https://doi.org/10.1038/s41586-
020-2649-2.

60. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau,
D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0:
fundamental algorithms for scientific computing in Python. Nat Methods *17*, 261–
272. https://doi.org/10.1038/s41592-019-0686-2.

61. McKinney, W. (2010). Data Structures for Statistical Computing in Python.
Proceedings of the 9th Python in Science Conference, 56–61.
https://doi.org/10.25080/Majora-92bf1922-00a.

62. R Development Core Team (2023). R : A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria.

63. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal *17*, 10–12. https://doi.org/10.14806/ej.17.1.200.

64. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

65. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work
with high-throughput sequencing data. Bioinformatics *31*, 166–169.
https://doi.org/10.1093/bioinformatics/btu638.

66. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for
comparing genomic features. Bioinformatics *26*, 841–842.
https://doi.org/10.1093/bioinformatics/btq033.

67. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup
(2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*,
2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

68. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne,
S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for
deep-sequencing data analysis. Nucleic Acids Res *44*, W160-165.
https://doi.org/10.1093/nar/gkw257.

69. Wagih, O. (2017). ggseqlogo: a versatile R package for drawing sequence logos.
Bioinformatics *33*, 3645–3647. https://doi.org/10.1093/bioinformatics/btx469.

70. Wang, L., Wang, S., and Li, W. (2012). RSeQC: quality control of RNA-seq
experiments. Bioinformatics *28*, 2184–2185.
https://doi.org/10.1093/bioinformatics/bts356.

71. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676–682.

https://doi.org/10.1038/nmeth.2019.

72. Waskom, M.L. (2021). seaborn: statistical data visualization. Journal of Open Source Software *6*, 3021. https://doi.org/10.21105/joss.03021.

73. Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Science *30,* 70–82. https://doi.org/10.1002/pro.3943.

74. Croll, T.I. (2018). ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps. Acta Crystallogr D Struct Biol *74*, 519–530. https://doi.org/10.1107/S2059798318002425.

75. Jo, S., Kim, T., Iyer, V.G., and Im, W. (2008). CHARMM-GUI: A web-based graphical user interface for CHARMM. Journal of Computational Chemistry *29*, 1859–1865. https://doi.org/10.1002/jcc.20945.

76. Lee, J., Cheng, X., Swails, J.M., Yeom, M.S., Eastman, P.K., Lemkul, J.A., Wei, S., Buckner, J., Jeong, J.C., Qi, Y., et al. (2016). CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. J Chem Theory Comput *12*, 405–413. https://doi.org/10.1021/acs.jctc.5b00935.

77. Qi, Y., Ingólfsson, H.I., Cheng, X., Lee, J., Marrink, S.J., and Im, W. (2015). CHARMM-GUI Martini Maker for Coarse-Grained Simulations with the Martini Force Field. J Chem Theory Comput *11*, 4486–4494. https://doi.org/10.1021/acs.jctc.5b00513.

78. Lomize, A.L., Todd, S.C., and Pogozheva, I.D. (2022). Spatial arrangement of proteins in planar and curved membranes by PPM 3.0. Protein Sci *31*, 209–220. https://doi.org/10.1002/pro.4219.

79. Abraham, M.J., Murtola, T., Schulz, R., Páll, S., Smith, J.C., Hess, B., and Lindahl, E. (2015). GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX *1–2*, 19–25. https://doi.org/10.1016/j.softx.2015.06.001.

80. Michaud-Agrawal, N., Denning, E.J., Woolf, T.B., and Beckstein, O. (2011). MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. Journal of Computational Chemistry *32*, 2319–2327. https://doi.org/10.1002/jcc.21787.

81. Gowers, R.J., Linke, M., Barnoud, J., Reddy, T.J.E., Melo, M.N., Seyler, S.L., Domański, J., Dotson, D.L., Buchoux, S., Kenney, I.M., et al. (2016). MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. Proceedings of the 15th Python in Science Conference, 98–105.

1154   https://doi.org/10.25080/Majora-629e541a-00e.

82. Madeira, F., Pearce, M., Tivey, A.R.N., Basutkar, P., Lee, J., Edbali, O., Madhusoodanan, N., Kolesnikov, A., and Lopez, R. (2022). Search and sequence analysis tools services from EMBL-EBI in 2022. Nucleic Acids Res *50*, W276–W279. https://doi.org/10.1093/nar/gkac240.

83. Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. Nucleic Acids Research, gkae268. https://doi.org/10.1093/nar/gkae268.

84. Robert, X., and Gouet, P. (2014). Deciphering key features in protein structures with the new ENDscript server. Nucleic Acids Research *42*, W320–W324. https://doi.org/10.1093/nar/gku316.

85. Tamura, K., Stecher, G., and Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Molecular Biology and Evolution *38*, 3022–3027. https://doi.org/10.1093/molbev/msab120.

86. Nye, J., Buster, D.W., and Rogers, G.C. (2014). The use of cultured Drosophila cells for studying the microtubule cytoskeleton. Methods Mol Biol *1136*, 81–101. https://doi.org/10.1007/978-1-4939-0329-0_6.

87. Nishida, K.M., Iwasaki, Y.W., Murota, Y., Nagao, A., Mannen, T., Kato, Y., Siomi, H., and Siomi, M.C. (2015). Respective functions of two distinct Siwi complexes assembled during PIWI-interacting RNA biogenesis in Bombyx germ cells. Cell Rep *10*, 193–203. https://doi.org/10.1016/j.celrep.2014.12.013.

88. Miyoshi, K., Tsukumo, H., Nagami, T., Siomi, H., and Siomi, M.C. (2005). Slicer function of Drosophila Argonautes and its involvement in RISC formation. Genes Dev *19*, 2837–2848. https://doi.org/10.1101/gad.1370605.

89. Saito, K., Ishizuka, A., Siomi, H., and Siomi, M.C. (2005). Processing of pre-microRNAs by the Dicer-1-Loquacious complex in Drosophila cells. PLoS Biol *3*, e235. https://doi.org/10.1371/journal.pbio.0030235.

90. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022). ColabFold: making protein folding accessible to all. Nat Methods *19*, 679–682. https://doi.org/10.1038/s41592-022-01488-1.

91. Souza, P.C.T., Alessandri, R., Barnoud, J., Thallmair, S., Faustino, I., Grünewald, F., Patmanidis, I., Abdizadeh, H., Bruininks, B.M.H., Wassenaar, T.A., et al. (2021). Martini 3: a general purpose force field for coarse-grained molecular dynamics. Nat Methods *18*, 382–388. https://doi.org/10.1038/s41592-021-01098-3.

92. Lambkin, C.L., Sinclair, B.J., Pape, T., Courtney, G.W., Skevington, J.H., Meier, R., Yeates, D.K., Blagoderov, V., and Wiegmann, B.M. (2013). The phylogenetic

relationships among infraorders and superfamilies of Diptera based on morphological evidence. Systematic Entomology *38*, 164–179. https://doi.org/10.1111/j.1365-3113.2012.00652.x.

93. Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen, P.B., Ware, J., Flouri, T., Beutel, R.G., et al. (2014). Phylogenomics resolves the timing and pattern of insect evolution. Science *346*, 763–767. https://doi.org/10.1126/science.1257570.

94. Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol Biol Evol *4*, 406–425. https://doi.org/10.1093/oxfordjournals.molbev.a040454.

A

AR
SAM
SAM
Daed
CC
α-H2
Gasz
TM
TM

B

Daed eCC
Gasz eCC Gasz eCC
CC α-H2
Daed Gasz
α-H2
Gasz
CC
Daed
Daed eCC
90°

C

Daed ⟨ SAM   CC TM ⟩ 278 aa
Daed eCC

```
140              eCC      150              160              170              180
ICNH PEAENYNNQILGNAGNHFMSLRASLAATNYKLQVSTPEDVVVGDKR
```

```
190              200       CC      210              220              230
YASCFAQETLKSVNQI TEEIAKDLAKIEANAQNARSQKKVQGNES HKKKW
```

Gasz ⟨ AR 1 2 3 4 5 6   SAM   TM ⟩ 461 aa
AR   Gasz eCC
α-H2

```
340     eCC      350              360              370              380        α-H2
DEMNN FYEILLITASHLQHLVIIQASLRFILKNQEHNKMGQPSEM QVAAM
```

```
390              400              410
KSNLQGYRDVLNQITKTVKYLGS FSPSQNP
```

━━ α-helix

D

Gasz KD
input   Flag IP
Luc WT ΔAR1-3 ΔAR4-6 ΔSAM Δα-H2   Luc WT ΔAR1-3 ΔAR4-6 ΔSAM Δα-H2
(kDa)
70 —
55 —   Flag-Luc
                                                              *
45 —   Flag-Gasz
35 —
30 —   Daed

AR   SAM   α-H2 TM
Gasz WT   [1 2 3 4 5 6] [SAM] [α-H2] 461 aa
ΔAR1-3   [4 5 6] [SAM] [α-H2] 360 aa
ΔAR4-6   [1 2 3   ] [SAM] [α-H2] 363 aa
ΔSAM   [1 2 3 4 5 6] [α-H2] 384 aa
Δα-H2   [1 2 3 4 5 6] [SAM] 432 aa

Diptera-specific Daedalus controls Zucchini endonucleolysis in piRNA biogenesis independent of exonucleases

Yuica Koga, Shigeki Hirakata, Mayu Negishi, Hiroya Yamazaki, and Mikiko C. Siomi[*]

Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo 113-0032, Japan

*Correspondence: siomim@bs.s.u-tokyo.ac.jp

Figure S1. Related to Figure 1; Coiled-coil prediction and specificity of anti-Daed antibodies

(**A**) Sequence-based CC prediction of Daed (upper) and Gasz (lower) by CoCoNat. Red line indicates the

CC region predicted by AlphaFold2. (**B**) Western blotting shows that anti-Daed monoclonal antibodies

produced in this study are specific to Daed. Daed KD: Daed knockdown by RNAi. control: EGFP KD. β-

Tub: loading control.

Figure S2. Related to Figure 2; Functions and behaviors of Daed and Gasz in OSCs

(**A**) Northern blotting using probes that detect *idefix* piRNA and *tj* piRNA. miR-310: loading control. KD: knockdown by RNAi. control: EGFP KD. pre-piRNA: piRNA precursors. (**B**) Western blotting shows the levels of Flag-Gasz WT and Flag-Gasz Δα-H2 where the same amounts of plasmids were used for transfection. β-Tub: loading control. The statistics of western blotting signals (n=3; left) and the mRNA expression levels (n=3; right). Signal intensities (left) or values (right) relative to F-Gasz WT were normalized with those of β-Tub or *rp49*, respectively, and presented as mean values±SE. ***: *p* <0.005;

n.s.: *p* >0.05 [*t*-test (unpaired, two-sided)]. (**C**) Western blotting shows HSP60 (a mitochondrial fraction marker) and β-Tub (a cytoplasmic fraction marker) in total lysates, and mitochondrial and cytoplasmic fractions in Figure 2G. (**D**) Summary of Armi subcellular localization in OSCs under each condition.

A

Armi | N34 | | DEXXQ-box | UPF1-like | 1,188 aa

```
MFTYVSKFFT NPDRNREDIL ESLDRENSFL DQKLMEEKMD QQLKANPNEI  50
NGVLSNKIAE LTHGLSEMDV SKESSCTARK GVITSLDGDR GVIDKDVLFE  100
TKVAEDIILD LHVGCVVEYL TFTTGEAMRV VKVKSILEHS WEDTSQKEIE  150
KAVDNLKNEK PTFFNTETRS VLGLISQRLA SSIDVETEYG QLTVELDNIE  200
MNFIPTNGDR VRLECNIQLD DGFVDKQGEI LEVTKLFPTR IQEGEKCIVE  250
RVYVHMVVLG PETYILKTDL PTGTDLHLGD IVLADLIECQ YSKFTRRAIK  300
ITPLEKNFGA TKLTQQSSMG SSGSGKAVTV TGVNRFITAE LWQKESVSLK  350
LTNNLNRTLR LESITVCNDS ESQLSVVSPL ESKEISSGSE ITVTFEIHTQ  400
FLGEAIEKYV LNFDLLKVRR VFTVIVCKTK EEVAEAEKRM IAAEALMAPG  450
RNSQERSRFY ANQVWCNKVD VIPGQQIVTK RRFVALRLGC FEVPKELRQI  500
CLTSERRQEM IKAIEQHYSF LTEPLSIKTY MHRFRLLLHL EEIECFVNFR  550
NYDRDRAHFL RDGEFLTLQI ENLAERRPSL VIGDTLRVIN PWSDPDSQTT  600
KSYEGIIHKV LFDRILLKFH SSFQEKYNGE DYRLEFYFSR YSFRKQHHAI  650
SKIVGVMGED FLFPSKVTKR ENPQLDVYMK DDDMYLYDSK LEWYNQSLNS  700
IQKRAVFNIL RGEAENIPYV LFGPPGSGKT MTLIETLLQL VRNLPGARIL  750
VGTPSNSSAD LVTKRLIDSK ALLQGDFIRL VSYNQVEKDL IPPEIMSYCA  800
TSDVGAVGSC EDKMMVTESG LKLRCQAKFI GTHRITISTC TTLGNFLQLG  850
FPAGHFTHVL FDEAGQCTEP ETMVPIVMLT KKRSQVVLSG DPRQLQSIVT  900
SRIASKMGFS ISFLERLLER SPYRKDLQRF PESSGYNPLV LTKLLYNYRA  950
LPSIMSIYSR LFYDDELIPV LSEKDSRESR LLSKLRCVFE SEKDIPQAHG  1000
TFFYGIIGEN RQNNDSPSWF NPQEVREVFL MTIALYRANV TADQIGIITP  1050
YQKQVKMLRS MFIGTDVVMP KIGSVEEFQG QERDIILIST VRSSEEILRM  1100
DARFSLGFVR CSKRLNVAVS RARAMMIIFG NPHLLAVDEC WRQLILFCVK  1150
NNAYFGCDLP QMVINQKDEV PVCLETFVPS LNTTDDLN              1188
```

B

Figure S3. Related to Figure 3; Effects of Armi mutants on piRNA biogenesis

(**A**) Peptide sequence and domain structure of Armi. The ATP binding residues and nucleotide binding residues are indicated in red and white, respectively. Asp1015, Arg1026, and Arg1037 are highlighted in blue. (**B**) Quantification of piRNA levels (n=3) (Figure 3H). Values relative to the control (EGFP KD) were normalized to Piwi level and presented as mean values±SE. ***: $p < 0.005$; n.s.: $p > 0.05$ [t-test (unpaired, two-sided)].

Figure S4. Related to Figure 4; Effects of Daed ΔSAM on piRNA biogenesis and Piwi-piRISC

function

(**A**) Quantification of piRNA levels (n=3) (Figure 4A). Values relative to the control (EGFP KD) were normalized to Piwi level and presented as mean values±SE. ***: p <0.005; *: p <0.05; [t-test (unpaired, two-sided)]. (**B**, **C**) Scatter plots showing the length distribution of piRNAs upon subdivision by target transposons (**B**) and the abundance of reads mapped antisense to transposons (**C**; Spearman's rho=0.98) in Daed WT and ΔSAM libraries. (**D**) Left: Piwi-bound piRNAs obtained from cytoplasmic fraction (Cyto) and nuclear extract fraction (Nuc). Immunoprecipitated Piwi was detected by western blotting. Right: Western blotting showing β-Tub (a cytoplasmic marker) and Histon H3 (a nucleus marker) in each fraction.

(**E**) The expression levels of transposons (*gypsy*, *blood*, *gypsy5*, *quasimodo*, *17.6*, and *mdg3*) in control OSCs (dark gray), Daed-lacking OSCs (Daed KD; light gray), Flag-Daed WT rescued (WT; green) or Flag-Daed ΔSAM rescued cells (ΔSAM; red) (n=5). Values relative to the control (EGFP KD) were normalized to *rp49* and presented as mean values±SE **: p <0.01; *: p <0.05; n.s.: p >0.05 [Wilcoxon rank sum exact test (two-sided)]. (**F**) Venn diagram showing the variety of positions of piRNA 5′ ends in the WT and ΔSAM libraries. (**G**) Heatmaps showing 5′ end counts of piRNAs mapped around the 10,000 most abundant piRNA 3′ ends (middle) and its binary histogram (top). In sequence logos (bottom), piRNAs were aligned according to the 3′ end of the piRNAs (black arrowhead) and its immediately downstream nucleotide is highlighted in green. Left: Daed ΔSAM piRNAs in "Unchanged" group. Right: Daed ΔSAM piRNAs in "Elongated" group. (**H**) piRNA production simulation. The U depletion in 29–34-nt piRNAs in WT and equivalent regions in ΔSAM are highlighted in yellow.

Figure S5. Related to Figure 4; MD simulation of Armi–AmTEC complex

(**A**, **B**) Root mean square deviation (RMSD) of the Armi–AmTEC complex (**A**) and its Daed ΔSAM version (**B**) in molecular dynamics simulations (n=5) indicates that the simulation reaches equilibrium state over the simulation time. (**C**) The average distance between the center of mass (CoM) of Armi helicase domain and that of AmTEC (n=5) for each time point of the trajectory is shown. (**D**) Violin plot of CoM distances between Armi helicase domain and AmTEC (blue) and its Daed ΔSAM version (red) for all time points in all trajectories. ***: $p < 0.005$ [Welch's *t*-test (two-sided)].

Figure S6. Related to Figure 4; Effects of Armi KEE on piRNA biogenesis

(**A**) Length distribution of piRNAs (18–40 nt) in Armi WT (green) and KEE (red) libraries (n=3), presented

as mean values±SE. (**B**) Sequence logos representing 11 nt from the 5′ end of piRNAs in the Armi WT

and KEE libraries. (**C**) Heatmaps showing the frequency of the 3′ end positions (red) corresponding to

each 5′ end of piRNAs in Armi WT (upper) and KEE (lower) libraries. The 50,000 piRNA 5′ end positions

common to both libraries and most abundant in the WT library were used for analysis. (**D**) The model

summarizing the effects of the perturbation of two Armi–AmTEC interactions (Armi N34–Daed eCC and Armi UPF1-like–Daed SAM). Blue arrowhead (left panel): Zuc cleavage. Red arrowhead (right panel): obscured Zuc cleavage. White arrowhead (right panel): Zuc cleavage corresponding to WT. Black dashed line with a red cross (middle and right panel): perturbated interaction.

Figure S7. Related to Figure 5; Sequence alignment of Daed in species

(**A**, **B**) Phylogenetic tree (**A**) and multiple sequence alignment (**B**) of Daed orthologs constructed and visualized using Clustal Omega, MEGA 11 and ESPript.

Figure S8. Related to Figure 5; Sequence alignment of Armi UPF1-like helicase domain in species

Multiple sequence alignment of *D. melanogaster* Armi and its orthologs. Red arrowheads: Asp1015, Arg1026, and Arg1037 (D-R-R) of *D. melanogaster* Armi.

Figure S9. Related to STAR Method "piRNA sequencing and bioinformatic analysis"; Reproducibility of piRNA-seq

(**A,B**) Scatter plots showing reproducibility of piRNA-seq of Daed WT, ΔSAM mutant (**A**), Armi WT and KEE mutant libraries (**B**). Gray dots indicate the number of normalized piRNA reads mapped to coding genes.

Table S1. Related to STAR Method "Plasmid construction"; Sequences of primers used for plasmid construction

name	primer sequence	plasmid
Daed-clonig-insert_F	CTTGGTACCGAATTCATGGCAACGAAGACGCACAC	pAc-Flag-Daed
Daed-clonig-insert_R	GACTCGAGCGGCCGCTTAAGTCTGAACGGAGATCC	pAc-Flag-Daed
Daed-clonig-vector_F	TAAGCGGCCGCTCGAGTCTA	pAc-Flag-Daed
Daed-clonig-vector_R	CATGAATTCGGTACCAAGCT	pAc-Flag-Daed
Flag-Luc_insert_F	ATGGACTACAAAGACCATGACG	pAc-Flag-Luc
Flag-Luc_insert_R	CTTGTCATCGTCATCCTTGT	pAc-Flag-Luc
Flag-Luc_vector_F	GTCTTTGTAGTCCATGGTGGCTAGCCCGAT	pAc-Flag-Luc
Flag-Luc_vector_R	GGATGACGATGACAAGATGGAAGACGCCAAAAACATAAAGAAAG	pAc-Flag-Luc
Flag-Daed (RNAi resistant)_F	GATAAGCTCATTGCCGAGAACATT	pAc-Flag-Daed (RNAi resistant)
Flag-Daed (RNAi resistant)_R	AACGCATGTACTGGCGTAGTCC	pAc-Flag-Daed (RNAi resistant)
Flag-Daed ΔSAM_F	ATTCCCAACCAGAATCCCAGTTACG	pAc-Flag-Daed ΔSAM
Flag-Daed ΔSAM_R	TGTACTGGCGTAGTCCGTTTCCACG	pAc-Flag-Daed ΔSAM
Flag-Daed ΔCC_F	CACAAGAAGAAGAAGTGGAGCCTGG	pAc-Flag-Daed ΔCC
Flag-Daed ΔCC_R	GATCTGATTCACGCTTTTTAGAGTT	pAc-Flag-Daed ΔCC
GST-Daed_full_length_insert_F	GGGCCCCTGGGATCCATGGCAACGAAGACGCACAC	pGEX-GST-Daed
GST-Daed_full_length_insert_R	ACCCGGGAATTCCGGTTAAGTCTGAACGGAGATCC	pGEX-GST-Daed
GST-Daed_full_length_vector_F	CCGGAATTCCCGGGTCGACTCGAGC	pGEX-GST-Daed
GST-Daed_full_length_vector_R	GGATCCCAGGGGCCCCTGGAACAGA	pGEX-GST-Daed
GST-Daed ΔTM_F	TGGACCAAGTTCCGCAGCGCTCCCC	pGEX-GST-Daed ΔTM
GST-Daed ΔTM_R	CTTCTTCTTCTTGTGACTCTCGTTG	pGEX-GST-Daed ΔTM
GST-Daed ΔTM-His_F	CATCATCACTAACCGGAATTCCCGG	pGEX-GST-Daed ΔTM-His
GST-Daed ΔTM-His_R	ATGATGATGAGTCTGAACGGAGATC	pGEX-GST-Daed ΔTM-His
GST-Daed (N1-197)_insert_F	GGGCCCCTGGGATCCATGGCAACGAAGACGCA	pGEX-GST-Daed (N1-197)
GST-Daed (N1-197)_insert_R	ACCCGGGAATTCCGGTTATTCCTGGGCAAAGCAG	pGEX-GST-Daed (N1-197)
GST-Daed (N1-197)_vector_F	CCGGAATTCCCGGGT	pGEX-GST-Daed (N1-197)
GST-Daed (N1-197)_vector_R	GGATCCCAGGGGCC	pGEX-GST-Daed (N1-197)
Flag-Gasz ΔAR1-3_F	CACGGAGCCGTGATCAATGTTTCGG	pAc-Flag-Gasz ΔAR1-3
Flag-Gasz ΔAR1-3_R	CTTGGAGCGTTTCACCTCATCCGAG	pAc-Flag-Gasz ΔAR1-3
Flag-Gasz ΔAR4-6_F	CCAGCCTCCACTTACCTAGTGCCGA	pAc-Flag-Gasz ΔAR4-6
Flag-Gasz ΔAR4-6_R	CCGCAAGAGCAGGCCCACAATTCTC	pAc-Flag-Gasz ΔAR4-6
Flag-Gasz ΔSAM_F	TTCTATGAGATCCTACTGATCACCG	pAc-Flag-Gasz ΔSAM
Flag-Gasz ΔSAM_R	TTGAAAATATTCCGGGCAATCGCTC	pAc-Flag-Gasz ΔSAM
Flag-Gasz ΔCC_F	TTTAGCCCCTCTCAGAATCCCTTGT	pAc-Flag-Gasz ΔCC
Flag-Gasz ΔCC_R	CTCCGAAGGTTGCCCCATTTTATTG	pAc-Flag-Gasz ΔCC
Armi D1015K-Flag_F	ATAACAAATCaCCTTCCTGGTTCAA	pAc-Armi KEE-Flag
Armi D1015K-Flag_R	TTTGCCTGTTTTCGCCAATGATTCC	pAc-Armi KEE-Flag
Armi R1026E-Flag_F	AATCCCCAAGAGGTCGAGGAGGTGT	pAc-Armi KEE-Flag
Armi R1026E-Flag_R	GAACCAGGAAGGTGAATCGTTATTT	pAc-Armi KEE-Flag
Armi R1037E-Flag_F	ACGAGGCCAATGTGACTGCCGATCA	pAc-Armi KEE-Flag
Armi R1037E-Flag_R	ACAGGGCGATGGTCATTAGGAACAC	pAc-Armi KEE-Flag
MOV10L1-clonig-insert_F	ATCGGGCTAGCCACCATGCTGCGCCTTGCTGCCAA	pAc-MOV10L1-Flag
MOV10L1-clonig-insert_R	TTTGTAGTCCTCGAGCTTTTGGAGAGCCTGGAGTT	pAc-MOV10L1-Flag
MOV10L1-clonig-vector_F	CTCGAGGACTACAAAGACCATGACG	pAc-MOV10L1-Flag
MOV10L1-clonig-vector_R	CATGGTGGCTAGCCCGATCCGGGGT	pAc-MOV10L1-Flag
ASZ1-clonig-insert_F	GGTACCGAATTCATGGCAGCGGGCACACTGAGGGG	pAc-Myc-ASZ1
ASZ1-clonig-insert_R	TCGAGCGGCCGCTTATTTTTCTCTGCAAAGTTAGCT	pAc-Myc-ASZ1
ASZ1-clonig-vector_F	TAAGCGGCCGCTCGAGTCTAGAGGG	pAc-Myc-ASZ1
ASZ1-clonig-vector_R	CATGAATTCGGTACCAAGCTTGTTC	pAc-Myc-ASZ1

Table S2. Related to STAR Method "RNAi and transfection"; Sequences of siRNAs

name	sense	antisense
siEGFP	*GGCAAGCUGACCCUGAAGU*TT	*ACUUCAGGGUCAGCUUGCC*TT
siDaed	*CCUGUGUGGACAAACUUAUUG*	*AUAAGUUUGUCCACACAGGUG*
siGasz	*CGCUCGAAACCAGCUCCUU*TT	*AAGGAGCUGGUUUCGAGCG*TT
siArmi	*CAGCUUAAGGCGAAUCCAA*TT	*UUGGAUUCGCCUUAAGCUG*TT
siZuc	*GCAUUGCCGUCAGCACUGU*TT	*ACAGUGCUGACGGCAAUGC*TT
siPapi	*GAUUACGAGACAAAGACAG*TT	*CUGUCUUUGUCUCGUAAUC*TT
siNbr	*CCUAAUGAGCAACCTGAAG*TT	*CUUCAGGUUGCUCAUUAGG*TT

*: RNA is shown in itallics

Table S3. Related to STAR Method "Northern blotting"; Sequences of oligos used for northern blotting

name	probe sequence
idefix-piRNA	AAACTACTGGCAATCGTTTGGGAA
tj-piRNA	GGTAATGGGAATGCACTTCTCTTGAA
miR-310	AAAGGCCGGGAAGTGTGCAATA

Table S4. Related to STAR Method "RT-qPCR"; Sequences of primers used for RT-qPCR

name	forward primer (5′ > 3′)	reverse primer (5′ > 3′)
rp49	CCGCTTCAAGGGACAGTATCTG	ATCTCGCCGCAGTAAACGC
Daed	GCAACGAAGACGCACACAAT	TCGGTGTGCGAGTGAAAGTT
Gasz	ACTTCCTGGGCTACAACACG	GGCCTCCAGTATGCCGTTAA
Flag-Gasz	ATGGACTACAAAGACCATGACGGTG	GAGCAGGCCCACAATTCTCTCCGCC
Nbr	GCCTTGGCAAGAAACTGAAC	ACCCCGCCTGAGAAAGTTAT
Papi	GACAGAGGTGGCACATTGGA	GTTCAGAGTTGTCGGCTGGA
mdg1	AACAGAAACGCCAGCAACAGC	CGTTCCCATGTCCGTTGTGAT
tabor	ACAACAGAAGCCGCAAACTG	CAGCTATTGTGAGTGGGCGT
412	ATGCCACTGGTCTAAGCACC	ATAGTGGTGCAAGTCCCTGC
stalker2	CACACAGAAGAACAAGCGGC	TTAACCGCAACGCTTCCTCT
gypsy	GAACCGGTGTACTCAAGAGC	CCAGGTCGGGCTGTTATAGG
blood	AAGACTGCCCTCGACCATTG	ATTCGCTGTTGTCATGTGCG
quasimodo	CCCGACAACAGACAAGAGGC	AGCCCAAGTGTTGCACTGTT
gypsy5	GCCCAGAGACAACGACAGAA	CTGTCTTTGCTGTCCCGGAT
17.6	AATTGGCGCAGTCGATGTGA	TGCCCGTTATCGTGGTACTT
mdg3	ATGCCATACGATGCGCAAAC	CGCTCTTCCTTGTGGGACAT