Diptera-specific Daedalus controls Zucchini endonucleolysis in

piRNA biogenesis independent of exonucleases

Yuica Koga, Shigeki Hirakata, Mayu Negishi, Hiroya Yamazaki, and Mikiko C.

Siomi*

Department of Biological Sciences, Graduate School of Science, The University of Tokyo,

Tokyo 113-0032, Japan

*Correspondence: siomim@bs.s.u-tokyo.ac.jp

Summary

PIWI-interacting RNAs (piRNAs) protect germline genomes and maintain fertility by repressing transposons. Daedalus and Gasz act together as a mitochondrial scaffold for Armitage, a necessary factor for Zucchini-dependent piRNA processing. However, the mechanism underlying this function remains unclear. Here, we find that the requirements of Daedalus and Gasz in this process are different: Daedalus physically interacts with Armitage, whereas Gasz supports Daedalus to maintain its function. Daedalus binds to Armitage through two distinct regions, an extended coiled-coil identified in this study, and a sterile α motif (SAM). The former tethers Armitage to mitochondria, while the latter controls Zucchini endonucleolysis to define the length of piRNAs in an exonuclease-independent manner. piRNAs produced in the absence of Daedalus SAM do not exhibit full transposon silencing functionality. Daedalus is Diptera specific. Evolutionarily more complex species, such as mice, rely on exonucleases after Zuc processing to specify the length of piRNAs.

Keywords

Armitage, Daedalus, Gasz, piRNA, Zucchini, AmTEC, extended coiled-coil

Introduction

In *Drosophila* ovarian somatic cells (OSCs), the PIWI-interacting RNA (piRNA)-induced silencing complex (piRISC) assembles with one of the PIWI members, Piwi, in the cytoplasm.[1–4] piRISC then co-transcriptionally represses transposons in the nucleus. The vast majority of piRNAs in OSCs originate from the piRNA cluster, *flamenco* (*flam*).[5] The *flam* locus contains numerous transposon remnants, most of which have an orientation

opposite to the direction of transcription. Therefore, piRNAs can naturally hybridize to target mRNAs through RNA–RNA base pairing.[5,6] After nuclear processing, including splicing,[7] the *flam* transcripts are exported to the cytoplasm where they bind to female sterile (1) Yb (Yb) via *cis*-elements embedded in the RNAs. This interaction induces the assembly of non-membranous organelle Yb bodies through phase separation, which accelerates subsequent steps in the piRNA processing.[8,9]

At this point, the *flam* transcripts are coarsely digested by an unknown factor, generating numerous 5′ ends. Piwi binds one-to-one to these 5′ ends via the MID domain and becomes the Piwi-piRISC precursor (pre-Piwi-piRISC).[10,11] Concomitantly, the RNA helicase Armitage (Armi) binds to the complex, allowing pre-Piwi-piRISC to exit Yb bodies and move to the mitochondrial surface where Zucchini (Zuc) awaits to process the complex.[12] Zuc has a transmembrane (TM) region, which localizes Zuc to the outer membrane of mitochondria.[13–16] After Zuc RNA cleavage releases mature Piwi-piRISC from pre-Piwi-piRISC, nascent Piwi binds to the remaining piRNA precursor through its 5′ end, generating a new pre-Piwi-piRISC in Yb body-independent and Zuc-dependent manners. This mechanism of sequential generation of Piwi-piRISC from a single piRNA precursor is known as phasing.[17,18] Piwi-piRISC is then transported to the nucleus,[19] where it associates with nascent transposon transcripts and suppresses transcription in concert with multiple cofactors, such as Gtsf1/Asterix and Maelstrom.[20–27]

An RNAi-based gene screen performed in *Drosophila* ovaries identified many factors acting in the piRNA pathway, including Gasz and Daedalus (Daed).[28] Both proteins have a TM, like Zuc, and they form a heteromeric complex that acts in concert with Armi to support Zuc cleavage in piRNA maturation.[12,29] However, the molecular details of this process, especially the function of Daed, remain unclear.

In this study, we found that Daed interacts with Armi through two distinct regions,

the extended coiled-coil (eCC) identified in this study, and the sterile α motif (SAM). We

also show that Daed interacts with Gasz by forming a helical barrel using the two α-

helices present in each of Daed and Gasz. Through this interaction, Daed stabilizes Gasz,

but not *vice versa*. Furthermore, we found that the Armi−Daed interaction through Daed

eCC tethers Armi to mitochondria, while the Armi−Daed interaction through SAM controls

Zuc endonucleolysis to specify the length of piRNAs in an exonuclease-independent

manner. piRNAs produced in the absence of Daed SAM became atypically long and their

function in transposon silencing was attenuated. Evolutionarily more complex species,

such as mice, possess exonuclease, PNLDC1, but not Daed. Mouse piRNAs, which are

atypically elongated in the absence of PNLDC1, do not exhibit full functionality in

transposon silencing. These findings indicate that piRNA length specification, which is

crucial for transposon repression, is conserved between species, but that the means of

piRNA length specification differ between species.

Results

Daed and Gasz bind with each other via their respective eCC domains

According to the current model, the Daed−Gasz heterodimer tethers Armi bound to pre-

Piwi-piRISC to mitochondria for Zuc-dependent piRNA processing.[12,29] In this study, we

first predicted the higher-order structure of the Daed−Gasz heterodimer using

AlphaFold2[30,31] (Figure 1A). This indicated that both Gasz and Daed have two α-helixes

and that the coiled-coil structure composed of these four α-helixes is the center of the

Daed−Gasz association (Figure 1B). Daed has SAM, coiled-coil (CC), and TM domains,

and Gasz has ankyrin repeats (ARs), SAM, and TM domains.[12,29] The predicted

structures of Daed and Gasz showed that the second of the two α-helixes (Ala191–

His235) of Daed roughly overlaps with the CC domain (Thr206–Ser234) and that the two

α-helixes in Gasz (Phe345–Glu374 and Glu383–Ser412) are present between the SAM

and TM domains (Figure 1C). The amino acid sequence-based prediction of the CC

region[32] also roughly overlaps with the tandem α-helix regions (Figure S1A), supporting

the AlphaFold2 prediction. We propose that the two tandem α-helixes present in both

Gasz and Daed are referred to as extended coiled-coil (eCC) domains (Figure 1C).

90   The CC domain of Daed is the region that binds to Gasz.[29] To identify the region

of Gasz responsible for binding to Daed we expressed deletion mutants of Gasz in OSCs

lacking endogenous Gasz and then performed immunoprecipitation assays. Three of the

Gasz deletion mutants, ΔAR1-3, which lacks the first three ARs, ΔAR4-6, which lacks

the second three ARs, and ΔSAM, bound to Daed as efficiently as wild-type (WT) Gasz

did, but the other mutant, Δα-H2, lacking the second α-helix of eCC did not (Figures 1D

and S1B). These results indicate that the respective eCC domains of Daed and Gasz

are responsible for the Daed–Gasz association, consistent with the predicted structure.

Daed stabilizes Gasz via protein–protein interaction

To verify whether Daed is required for piRNA biogenesis in cultured OSCs, we depleted

intracellular Daed by RNA interference (RNAi) and conducted northern blotting. In control

OSCs, both *idefix* piRNA (a representative nongenic piRNA) and *traffic jam* (*tj*) piRNA (a

representative genic piRNA) were detected as expected (Figure S2A).[12] However, in the

absence of Daed, their piRNA precursors (pre-piRNAs) accumulated aberrantly, while

mature piRNAs were barely detected (Figure S2A), as in OSCs lacking Gasz.[12] This

indicates that both Gasz and Daed are essential for piRNA maturation in OSCs.

Western blotting confirmed that RNAi was effective at depleting Daed and Gasz

in OSCs (Figure 2A). We also found that, when Daed was depleted, levels of Gasz were

also markedly reduced, but that levels of Daed were unchanged in the absence of Gasz

(Figures 2A and 2B). Such trends were not observed for their mRNA levels (Figure 2C).

These results indicate that Daed is required to stabilize Gasz at the protein level, but that

Daed is not dependent on Gasz for its stabilization.

To investigate which domain of Daed is involved in Gasz stabilization, two

deletion mutants of Daed, ΔSAM and ΔCC (part of the second α-helix of the eCC, see in

Figure 1C), were expressed in OSCs depleted of endogenous Daed. Levels of Gasz

were restored with expression of WT Daed and ΔSAM mutant Daed, but not with ΔCC

mutant Daed (Figure 2D). A previous study showed that Daed binds to Gasz via the CC

region[29] and together these findings indicate that the Daed–Gasz interaction via their

respective eCC domains is crucial for the stabilization of Gasz (Figure 2E). The Δα-H2

mutant of Gasz, which lacks the binding domain to Daed, was rather unstable in OSCs,

but this was not because of mRNA instability (Figure S2B), supporting the above notion.

Daed alone or Gasz alone is insufficient to tether Armi to mitochondria

Armi has at least two roles in piRNA biogenesis in OSCs; one is to transport nascent

Piwi to Yb bodies to promote pre-Piwi-piRISC assembly, and the other is to translocate

the complex from Yb bodies to mitochondria to promote Zuc-dependent piRNA

processing.[12,33] After this second role, Armi returns to the cytosol to repeat these

functions, although immunofluorescence shows that Armi appears to be a constant

resident of Yb bodies (Control, Figure 2F).[8,12] In the absence of Zuc, Armi signals merged

with mitochondrial signals {[I] Zuc knockdown (KD), Figure 2F}. This demonstrates

dynamic movement of Armi in OSCs and indicates that Armi remains on the

mitochondrial surface when loss of Zuc impairs piRNA maturation.[12] Results from

mitochondrial fractionation followed by western blotting were consistent with this

statement (Figures 2G and S2C).

We then depleted Daed in Zuc-lacking OSCs, where both Daed and Gasz

should be absent (Figure 2A). Under these circumstances, as expected, Armi was not

retained on mitochondria and appeared in Yb bodies ([II] Zuc KD + Daed KD, Figure 2F).

To restore the levels of both Daed and Gasz simultaneously, Flag-Daed was

exogenously expressed in the OSCs and western blotting confirmed the recovery of both

Daed and Gasz levels (Figures 2G and S2C). Immunofluorescence showed that Armi

re-localized to mitochondria ([III] Flag-Daed, Figure 2F). However, when Gasz was

forcibly expressed alone in the absence of Daed, Armi remained in Yb bodies ([IV] Flag-

Gasz, Figure 2F). A similar result was obtained in OSCs expressing only Daed ([V] Zuc

KD + Gasz KD, Figure 2F). These results indicate that Gasz alone or Daed alone is not

sufficient to tether Armi to mitochondria. In other words, both Daed and Gasz are

required to retain Armi on the mitochondrial surface for Zuc-dependent piRNA

processing. Hereafter, the Daed–Gasz heterodimer is referred to as the Armi

mitochondrial tethering complex (AmTEC). The behavior of Armi in OSCs under the

conditions described above is summarized in Figure S2D.

Daed, but not Gasz, binds to Armi via two distinct sites

We next predicted the higher-order structure of AmTEC bound with Armi (Figure 3A).

Notably, no significant interaction between Gasz and Armi was observed, but Daed and

Armi were bound at two distinct sites: The first site was through the first α-helix of eCC

(Phe144–Val177) (Figure 1C), which interacted with the N-terminal region of Armi

(Met1–Leu34) (Figures 3A, 3B, and S3A). The second interaction was between the SAM

domain of Daed (Figure 1C) and the C-terminal helicase (i.e., UPF1-like family) domain

of Armi (Figures 3A, 3C, and S3A).

To verify that the Armi–Daed interaction was via the Armi N-terminus (Figure

3B), recombinant WT Armi (WT-Flag) and its mutant lacking 34 N-terminal residues

(ΔN34-Flag) were subjected to pull-down assays with Daed (Daed ΔTM) as prey. The

Daed TM was removed because its inclusion interfered with protein preparation. Under

conditions where a negative control, Luciferase (Luc), showed no binding to Daed ΔTM,

Armi WT bound to Daed but Armi ΔN34 did not (Figure 3D). This confirms that Armi and

Daed interact through the 34 N-terminal residues of Armi. We have previously shown

that the Armi ΔN34 mutant (i.e., a shorter isoform of Armi) is unable to leave Yb bodies.[12]

The present analysis raises an alternative hypothesis; the Armi ΔN34 mutant may be

able to leave Yb bodies but not be tethered to mitochondria because of the absence of

Daed binding. This would explain why the shorter Armi isoform is inactive in piRNA

biogenesis in OSCs.

To verify the Armi–Daed interaction via the Armi UPF1-like helicase domain

(Figure 3C), three residues in the domain, Asp1015, Arg1026, and Arg1037 (D-R-R),

which appear to be key to the binding, were mutated to Lys-Glu-Glu (K-E-E) and

immunoprecipitation assays were conducted (Figure 3E). Quantification of the results

indicated that the Armi KEE mutant bound to Daed with approximately 34% of the binding

efficiency of WT Armi (Figure 3F). Meanwhile, the Armi ΔN34 mutant bound to Daed with

approximately 20% of the binding efficiency of WT Armi. Immunofluorescence showed

that the Armi ΔN34 mutant accumulated in Yb bodies more intensely than the KEE

mutant in Arm/Zuc-lacking OSCs (Figure 3G). For binding to Daed, Armi may be more

dependent on the N-terminal region than on the UPF1-like helicase domain.

A lack of Armi in OSCs severely disrupted piRNA production.[16,34] This was

almost fully rescued by the expression of WT Armi (Figure 3H). We next expressed the

ΔN34 or KEE mutants of Armi as alternatives to the WT. The ΔN34 mutant failed to

rescue the disruption to piRNA production, but the KEE mutant partially restored piRNA

biogenesis (Figures 3H, S3B and S3C). This supports the notion that the N-terminal

region of Armi is more important than the C-terminal helicase domain in piRNA

biogenesis in OSCs. The strength of the Armi–Daed interaction (Figure 3E) appears to

be positively correlated with the function of Armi.

The two Armi–Daed associations differ in their functional contributions to piRNA

biogenesis

piRNAs produced under conditions involving the Armi KEE mutant were notably

increased in length by several bases (Figure 3H). We therefore investigated whether a

similar phenomenon occurred under conditions involving the Daed ΔSAM mutant, given

that the SAM domain is the binding partner of the UPF1-like helicase domain of Armi.

Indeed, in cells expressing the Daed ΔSAM mutant as an alternative to endogenous

Daed, Piwi-bound piRNAs became several bases longer and their levels were lower than

those in cells expressing WT Daed, but higher compared with cells lacking endogenous

Daed (Figures 4A and S4A).

To examine how the lack of binding between Daed SAM and Armi UPF1-like

domain actually affects piRNA biogenesis, we sequenced and analyzed Piwi-bound

piRNAs in OSCs expressing Daed ΔSAM and compared the results with those from

OSCs expressing WT Daed (Figures 4A and S4A). This length distribution of piRNAs

was indeed shifted towards the longer fractions when they were produced in cells lacking

the Daed SAM domain (Figure 4B). This trend was also observed when piRNAs were

subdivided by target transposons (Figure S4B). In contrast, piRNA-mapped transposons

were essentially unchanged (Figure S4C). This seems reasonable given that piRNAs

are produced from *flam* in both cases. Furthermore, the 1U bias[5,35] and piRNA

phasing[17,18] were similarly observed in both libraries (Figures 4C and 4D). The β-

elimination experiments showed the 3′ ends of the piRNAs to be modified with 2′-*O*-

methylation in both cases, unlike Argonaute 1 (Ago1)-bound miRNAs (Figure 4E).[36,37]

It should be noted that non-negligible amounts of Piwi immunofluorescence

remained in the cytoplasm when Daed ΔSAM was expressed in OSCs lacking

endogenous Daed (Figure 4F). piRNA loading is essential for Piwi nuclear

localization.[19,33] Therefore, we next examined the piRNA binding status of cytoplasmic

and nuclear Piwi. In contrast to nuclear Piwi, which was bound to piRNAs even in the

long fraction, cytoplasmic Piwi was bound to few piRNAs (Figure S4D). This indicates

that piRNAs produced under the ΔSAM conditions can localize Piwi to the nucleus as

piRISC, regardless of length, but are insufficient to convert all Piwi to functional Piwi-

piRISC. Consistent with this, transposons were not fully repressed under such conditions

(Figures 4G and S4E). These results indicate that the Armi–Daed association via the

UPF1-like helicase domain of Armi and the SAM domain of Daed is influential in Piwi-

piRISC production and function, although less so than the other Armi–Daed association

involving the 34 N-terminal residues of Armi and the eCC domain of Daed.

The Armi–Daed interaction via Daed SAM optimizes Zuc endonucleolysis for

piRNA-mediated transposon repression

To explore the nature of ΔSAM piRNA extension, we selected the 50,000 piRNA 5′ end positions common to the WT and ΔSAM and with high abundance in WT libraries (Figure S4F) and analyzed them further. Heatmaps depicted with their 5′ end positions aligned clearly indicate that ΔSAM piRNAs are more diverse in length. Many of them have some extra bases at the 3′ ends (Figure 4H). This is consistent with the length distribution shown in Figure 4A. There are also a few shorter piRNAs (Figure 4H). The relatively small number of shorter piRNAs may be because of protection by Piwi.

To further analyze the effects of Daed SAM loss on piRNAs, we extracted abundant 5′-end positions of 26- and 27-nucleotide (nt) piRNAs in the WT library, and then these piRNAs were classified as "Unchanged" (51,833 positions) and "Elongated" (53,258 positions) according to the absence or presence of 3′-end elongation. Both piRNA groups maintained comparable phasing and +1U bias (i.e., 1U bias of piRNAs, real and virtual, immediately following piRNAs verified to bind to Piwi[17,18]) (Figure S4G), indicating that piRNA elongation caused by the loss of Daed SAM is caused by Zuc endonucleolysis.

We next sorted piRNAs by their length and aligned them according to the 3′ ends of piRNAs. This showed that atypically long piRNAs of around 28–33 nt in the WT library tended to have significantly low U bias near their 3′ ends (U depletion), just upstream of the 5′ ends of the immediately downstream piRNAs, as reported previously (yellow boxes in WT, Figure 4I).[18] This was interpreted as showing that when Zuc cannot find U-rich regions in the piRNA precursors it waits to cleave the substrate until the U-poor region has passed, thereby producing atypically long piRNAs. This also explains why there were fewer 28–33-nt piRNAs in the WT library (Figure 4B). In contrast, there

was less notable "U depletion" in 28–33-nt ΔSAM piRNAs (yellow boxes in ΔSAM, Figure

4I). In addition, there was a substantial number of 28–33-nt piRNAs in the ΔSAM library

(Figure 4B). This indicates that when the Armi–Daed interaction is attenuated because

of a lack of Armi–Daed association via Daed SAM (and the UPF1-like helicase domain

of Armi), Zuc somehow produces these longer piRNAs more preferentially regardless of

the U content in the precursor sequences while maintaining the +1U bias.

We simulated the effect of Daed ΔSAM on piRNA production. First, the data

shown in Figure 4B were fitted to normal distributions. In parallel, we hypothetically

fragmented the sequence of the *flam* locus (21,629,891 to 21,792,731 on the X

chromosome; T was converted to U) into 163 piRNA precursors, starting at U and into

sizes of 1,000 nt or close to that size, and then simulated Zuc-dependent piRNA

production from their 5′ ends by assuming that Zuc cleavage occurs before U, produces

18–40-nt piRNAs, and follows the normal distributions for WT and ΔSAM. Sequence

analysis of the obtained piRNAs (25–34 nt) indicated that U depletion was observed in

simulations fitting the WT distribution, starting from 29 nt, but less so in simulations fitting

the ΔSAM distribution (Figure S4H). This indicates that Zuc can make Piwi-piRISC

without the rule of following U depletion, but that the level of piRISC is insufficient for

transposon repression.

We also sequenced and analyzed Piwi-bound piRNAs in OSCs expressing the

Armi KEE mutant or WT Armi (Figures 3H and S3B). We found that piRNAs in the Armi

KEE library were also several bases longer than the Armi WT piRNAs and retained 1U

bias (Figures S4I and S4J). Heatmaps showed that the KEE piRNAs also exhibited

characteristics of 3′ end elongation (Figure S4K). These results further support our notion

that the SAM domain of Daed and the UPF1-like helicase domain of Armi are the binding

partners and that the Armi–Daed interaction is indispensable to optimize Piwi-piRISC

maturation by Zuc and piRISC function in OSCs.

Daed is Diptera-specific and controls Zuc cleavage in a Nibbler-independent

manner

Armi and Gasz are widely conserved among species from invertebrates to vertebrates,

but Daed is unique to the Diptera order, which is composed of Brachycera suborder,

which includes *Drosophila* and other fly species, and Nematocera suborder which

includes mosquitoes (Figure 5A, S5A and S5B). In contrast, other species not belonging

Diptera order do not appear to have Daed orthologs.[38]

In mice, primary piRNA biogenesis occurs in the inter-mitochondrial cement,

where Armi homolog, MOV10L1, Gasz homolog, ASZ1, Zuc homolog, PLD6, and TDRD

proteins play a role in piRNA production: MOV10L1 binds to PIWI proteins (MILI, MIWI

and MIWI2) and piRNA precursors and contributes to piRNA biogenesis through its

helicase activity.[39,40] ASZ1 is essential for the production of MILI-bound piRNAs in mice[41]

and is thought to tether MOV10L1 to mitochondria, together with TDRD proteins, such

as TDRD1, to promote MIWI-piRISC production.[39,40]

We expressed both MOV10L1 and ASZ1 in OSCs and performed

immunoprecipitation. The MOV10L1−ASZ1 interaction was detected to a similar extent

in both normal OSCs and OSCs lacking intracellular Daed (Figures 5B and 5C).

Immunofluorescence confirmed this; the expression of ASZ1 was sufficient for the

mitochondrial localization of MOV10L1, independent of the presence of intracellular

Daed (Figure 5D). This indicates that the MOV10L1−ASZ1 interaction and MOV10L1

tethering is Daed-independent, while the Armi−Gasz interaction and Armi tethering in

Drosophila required Daed. These results also indicate that the Armi–Gasz association,

at least in OSCs, does not require other factors present in mouse testis. ASZ1 was stable

in OSCs lacking endogenous Daed (Figure 5C), unlike Gasz (Figure 2A), indicating that

the molecular system that actively degrades Gasz in the absence of Daed does not act

on ASZ1.

Why did *Drosophila* acquire Daed during evolution? In mice, poly(A)-specific

ribonuclease-like domain containing 1 (PNLDC1), a homolog of poly(A)-specific

ribonuclease (PARN), trims the 3′ ends of piRNAs and piRNAs are elongated after

PNLDC1 loss of function.[42,43] Mutation in PNLDC1 reduces LINE-1-targeting piRNAs and

mislocalizes MIWI2 to the cytoplasm, leading to LINE-1 derepression. This was nearly

equal to what we observed in this study with the *Drosophila* Daed ΔSAM mutant.

However, PNLDC1 is not conserved in the majority of Diptera order species.[44] Papi is

the *Drosophila* ortholog of TDRKH, which, in mice, acts as a scaffold for PNLDC1 and

whose functional inhibition causes piRNA elongation.[45,46] However, consistent with

previous studies using Papi mutant flies,[44] its depletion had little effect on piRNA length

in OSCs (Figure 5E).

The 3′-to-5′ exonuclease, Nibbler (Nbr), trims both miRNAs and piRNAs in

Drosophila and mosquitoes,[44,47–49] but the pre-piRNAs that undergo trimming are minor

in mosquitoes, and the loss of Nbr barely affected the length of Piwi-bound piRNAs in

the *Drosophila* ovary.[44] Our study confirmed this; namely, Nbr depletion in OSCs had

little effect on the piRNA length (Figure 5F). This indicates that Nbr may function in the

trimming of miRNAs, but not piRNAs, in OSCs. The lack of a marked increase in +1U

signal in piRNAs elongated in the Daed ΔSAM mutant (Figures 4D and S4G) also

supports this view.

In contrast, Armi D-R-R, which was essential for the regulation of Zuc cleavage

in *Drosophila*, is highly conserved in the fly species that lack PNLDC1/PARN and have

Daed (Figure S5C). These results indicate that *Drosophila,* and its closely related

species, may have acquired Daed after losing PNLDC1/PARN during evolution, and that

they may also have acquired the ability to produce mature piRNAs without further

trimming by exonucleases by allowing Armi to bind directly to Daed (Figure 5G).

Discussion

This study revealed that Daed and Gasz bind to each other via their eCC domains, both

of which are composed of two α-helixes that assemble into a stable helical barrel

consisting of all four of these α-helixes. When Daed lacks the second α-helix in the eCC

domain, the Daed–Gasz interaction is disrupted and Gasz becomes destabilized,

illustrating the importance of Daed in Gasz stabilization (Figure 2E). This also shows that

the intracellular level of Gasz is strictly regulated by the intracellular level of Daed,

indicating that OSCs do not tolerate the presence of Gasz alone. In contrast, OSCs do

tolerate the presence of Daed alone. However, Daed, in the absence of Gasz, failed to

anchor pre-Piwi-piRISC to mitochondria via Armi. Gasz is therefore complicit in the role

of Daed, although the contribution of Gasz to the direct binding of Daed to Armi appears

to be minimal.

Lack of binding through the Daed SAM did not completely abolish the Armi–

Daed association but resulted in aberrant Zuc function (Figure S3C). Under such

conditions, Piwi-bound piRNAs were atypically long because of extra bases at the 3′

ends, and there was no notable U depletion, a typical feature of piRNAs in normal OSCs.

In normal OSCs, atypically long piRNAs are produced only when piRNAs of typical sizes

cannot maintain the +1U bias. In contrast, in OSCs expressing Daed ΔSAM, Zuc shows

"relaxed selectivity" in terms of nucleotides and length and tends to produce longer

piRNAs, even under conditions where it can produce shorter (i.e., normal size) piRNAs

with the +1U bias, albeit in relatively small quantities. This could be because of the timing

of Zuc RNA cleavage was shifted slightly downstream of the precursor while the

preference for cleavage before U was conserved. Recombinant Zuc did not show U

selectivity,[15] indicating that, *in vivo*, Zuc may rely on a cofactor to confer U selectivity,

and that this cofactor may not require the Armi–Daed association to execute its function.

Interestingly, Piwi could accept those longer piRNAs produced under the ΔSAM

conditions and localized to the nucleus as Piwi-piRISC. At the same time, however,

nonnegligible amounts of Piwi remained in the cytoplasm because of the lack of piRNA

loading, which resulted in transposon derepression. This means that the ability of Daed

to interact with Armi via the SAM domain, in addition to the binding via the first α-helix in

the eCC domain, is essential not only to define the positions of Zuc cleavage, but also

to produce enough Piwi-piRISC for transposon repression in OSCs. It is worth noting

that no such regulatory factor has been identified to date.

Armi is unique in that it plays multiple roles in piRNA biogenesis. First, Armi

delivers unloaded Piwi to Yb bodies, facilitating the assembly of pre-Piwi-piRISC in these

organelles. In Yb bodies, Armi binds to pre-Piwi-piRISC and translocates the complex to

mitochondria. There, Armi anchors pre-Piwi-piRISC to the mitochondria by binding to

Daed. This allows Zuc to process pre-Piwi-piRISC into mature Piwi-piRISC. Armi is

therefore a factor that always accompanies Piwi and encourages it along the pathway to

develop into Piwi-piRISC.

Armi is an RNA helicase protein, and its RNA helicase activity is required to

avoid stochastic interactions with abundant mRNAs.[50] In contrast, up to this point, the

RNA helicase activity of Armi does not seem to be required for the piRNA precursors,

rather its activity would be inhibitory. However, in the absence of this activity, especially

for phasing, the sequential generation of Piwi-piRISC from a single piRNA precursor

would not occur. It is possible that an inhibitor of the RNA helicase activity of Armi is

present in OSCs and functions until phasing, but, upon binding to Armi, Daed would lose

its function allowing phasing to occur.

Animals that reproduce sexually retain the highly conserved piRNA mechanism

and use it to maintain their fertility by repressing transposons. However, the evolutionary

rate of many piRNA factors, including Armi, is rapid,[51,52] suggesting that these species

have adapted the machinery to their specific endogenous and exogenous environments.

Coevolution of piRNAs and transposons through host–pathogen arms races has been

proposed as a background to this process.[53] Although flies and mosquitoes belong to

the same Diptera order, the piRNA pathway factors are highly diverse.[54] On the other

hand, our observation that Daed is conserved among Diptera species may suggest the

importance of this factor in Dipteran piRNA biogenesis pathway where PNLDC1 is lost.

Author contributions

YK and MN performed biochemical experiments. YK, SH, and HY performed

bioinformatic analyses, computational structural prediction, simulation, and calculation.

MCS supervised and discussed the work and wrote the manuscript with the other authors.

Acknowledgments

We thank Y. Namba, Y.W. Iwasaki, H. Nishimasu, S. Kuraku, Y. Fukushima, T. Yasuda

for useful discussions, and T. Fujisawa, T. Kinoshita, and H. Ishizu for technical

assistance with the biochemical analysis. This study was supported by MEXT KAKENHI

Grant Numbers JP19H05466 (to MCS), JP22K15056 (to SH), JP23K14141 (to HY), and

JP22KJ1122 (to YK). Some computations were performed on the NIG supercomputer at

ROIS National Institute of Genetics.

Conflict of interest

The authors declare no competing interests.

Figure legends

Figure 1. Daed and Gasz interact via their respective eCC domains

(**A**) Predicted structure of the Daed-Gasz heterodimer by AlphaFold2. (**B**) Magnification

of the region indicated by the dotted line box in (**A**). The structure is shown from two

directions. (**C**) Domain structures of Daed and Gasz. Peptide sequences of eCCs of

Daed and Gasz are also shown. (**D**) Immunoprecipitation (Flag IP) and western blotting

show that one of the Gasz mutants, Δα-H2, fails to bind to Daed in OSCs lacking

endogenous Gasz. Flag-Luc: negative control. *: Heavy chains. The domain structures

Gasz WT and deletion mutants are also shown.

Figure 2. Daed stabilizes Gasz via interaction and both Daed and Gasz are

essential for the mitochondrial tethering of Armi

(**A**) Western blotting shows the high efficiency of knockdown (KD) of Daed and Gasz in

OSCs. Daed KD also destabilizes Gasz, but not *vice versa*. β-Tub: loading control. (**B**,

C) The statistics of western blotting signals (n=3; **A**) and the mRNA expression levels of

Daed and *Gasz* upon Daed KD and Gasz KD. Signal intensities (**B**) or values (**C**) relative

to control (EGFP KD) were normalized with those of β-Tub or *rp49*, respectively, and

presented as mean values±SE. **: $p<0.01$; ***: $p<0.005$; n.s.: $p>0.05$ [*t*-test (unpaired,

two-sided)]. (**D**) Western blotting showing the levels of endogenous Gasz and Flag-Daed

(WT and deletion mutants) in OSCs under each condition. β-Tub: loading control. The

domain structures of Daed WT and mutants are shown. (**E**) Model of Gasz stabilization

by Daed. (**F**) Subcellular localization of Armi (green) in OSCs under various conditions,

[I] to [V]. Mitochondria (mito) and Yb bodies (Yb; white arrowheads) are shown in

magenta. The signals of Flag-Gasz and Flag-Daed are shown in yellow. Images in the

same vertical row are identical cells. Control: normal OSCs. DAPI (blue): nuclei. Scale

bar 5 μm. (**G**) Western blotting shows the levels of endogenous Armi, Gasz, Daed, Flag-

Gasz (orange asterisk), and Flag-Daed (blue asterisk) in total lysates (total) and

mitochondrial fractions (mito) under conditions [I] to [VI]. For the efficiency of cell

fractionation, see Figure S2C.

Figure 3. Daed interacts with Armi through two sites, each contributing differently

to piRNA biogenesis

(**A**) The structure of the Armi–AmTEC complex predicted by AlphaFold2. Gray lines:

virtual lipid bilayers. The domain structures of Armi, Daed, and Gasz are also shown. (**B**)

Magnification of the "B" region in (**A**). The 34 N-terminal residues of Armi (N34) interacts

with the first α-helix of Daed eCC. (**C**) Magnification of the "C" region in (**A**). The Armi

UPF1-like helicase domain interacts with Daed SAM. Asp1015, Arg1026, and Arg1037

of Armi are shown in blue. (**D**) *In vitro* pull-down assays show Armi WT-Flag, but not

Armi ΔN34-Flag, binding to Daed (Daed ΔTM-His). Luc-Flag: negative control. *: Heavy

and light chains. (**E**) Immunoprecipitation (Flag IP) and western blotting show the

amounts of endogenous Daed co-immunoprecipitated with Armi WT-Flag, ΔN34-Flag,

and KEE-Flag. Luc-Flag: negative control. The domain structures of Armi WT and its

mutants are shown below. Asp1015, Arg1026, and Arg1037 mutated in the KEE mutant

are also shown. The peptide sequence of Armi is shown in Figure S3A. (**F**) The statistics

of western blotting signals (n=3; **E**). The signals of Daed were normalized with those of

Armi and presented as mean values±SE. *: p <0.05; ***: p <0.005 [t-test (unpaired,

two-sided)]. (**G**) Subcellular localization of Armi WT, ΔN34, and KEE (green) in OSCs

lacking endogenous Zuc (Zuc KD) and Armi (Armi KD). Mitochondria (mito; for WT and

KEE) and Yb bodies (Yb, white arrowhead; for ΔN34) are shown in magenta. Images in

the same vertical row are identical cells. DAPI (blue): nuclei. Scale bar: 5 μm. (**H**) Piwi-

bound piRNAs obtained under each condition. Signal intensities (n=3) are normalized as

a percentage of the total. The mean (line) and standard deviation (shadow) are shown.

The amount of Piwi in each sample is shown at the bottom.

Figure 4. The loss of Armi–Daed interaction via Daed SAM obscures Zuc cleavage

and impairs Piwi-piRISC function

(**A**) Piwi-bound piRNAs obtained under each condition. Signal intensities (n=3) are

normalized as a percentage of the total. The mean (line) and standard deviation

(shadow) are shown. The amount of Piwi in each sample is shown at the bottom. (**B**)

Length distribution of piRNAs (18–40 nt) in Daed WT (green) and ΔSAM (red) libraries

(n=3), presented as mean values±SE. (**C**) Sequence logos representing 11 nt from the

5′ end of piRNAs in Daed WT and ΔSAM libraries. (**D**) Heatmaps showing 3′ end counts

of piRNAs mapped around the 10,000 most abundant piRNA 5′ ends (lower) and its

binary histograms (upper) in Daed WT (left) and ΔSAM (right). (**E**) β-elimination assays.

Piwi-bound piRNAs in Daed KD OSCs expressing Daed WT and ΔSAM are shown.

Ago1-bound miRNAs are used as positive controls. (**F**) Subcellular localization of

endogenous Piwi (green) in Daed KD OSCs expressing Daed WT and ΔSAM (magenta).

DAPI (blue): nuclei. Scale bar: 5 μm. (**G**) The expression levels of transposons (*mdg1*,

tabor, *412*, and *stalker2*) in control OSCs (dark gray), Daed-lacking OSCs (Daed KD;

light gray), Flag-Daed WT rescued (WT; green) or Flag-Daed ΔSAM rescued cells

(ΔSAM; red) (n=5). Values relative to the control (EGFP KD) were normalized to *rp49*

and presented as mean values ± SE **: $p < 0.01$; n.s.: $p > 0.05$ [Wilcoxon rank sum exact

test (two-sided)]. (**H**) Heatmaps showing the frequency of the 3′ end positions (red)

corresponding to each 5′ end of piRNAs in Daed WT (upper) and ΔSAM (lower) libraries.

The 50,000 piRNA 5′ end positions common to both libraries and most abundant in the

WT library were used for analysis. (**I**) Sequence logos. piRNAs were sorted by their

lengths and aligned according to the 3′ end of the piRNAs (black arrowhead) and its

immediately downstream nucleotide is highlighted in green. The U depletion found in 28–

33-nt piRNAs in the WT library and the equivalent regions in ΔSAM are highlighted in

yellow.

Figure 5. Daed is fly-specific and controls Zuc cleavage in a Nbr-independent manner

(**A**) Protein conservation state of Armi, Gasz, Daed, PNLDC1, Papi, and Nbr in *D.*

melanogaster and other species. (**B**, **C**) Immunoprecipitation (Flag IP) and western

blotting show Myc-ASZ1 binds to MOV10L1 in normal (**B**) and Daed-depleted OSCs (**C**).

Luc-Flag: negative control. (**D**) Subcellular localization of Myc-ASZ1 alone or MOV10L1-

Flag (green) with Myc-ASZ1 (yellow) in normal (left two row) and Daed-depleted OSCs

(right two row). Mitochondria (mito) are shown in magenta. Images in the same vertical

row are identical cells. DAPI (blue): nuclei. Scale bar: 5 μm. (**E**, **F**) The efficiency of Papi

KD (upper) and Nbr KD (lower) (n=3). Piwi-bound piRNAs in control and Papi-KD (upper)

and Nbr-KD OSCs (lower). The amount of Piwi in each sample is also shown. (**G**) Model

proposed in this study. Diptera species acquired Daed instead of PNLDC1/PARN during

evolution; Daed and Gasz form AmTAC, which stably tethers Armi to the mitochondrial

surface via two sites. This allows Zuc to define the length of piRNAs without relying on

exonucleases such as PNLDC1/PARN or Nbr. In contrast, in mice, MOV10L1/Armi is

tethered solely by ASZ1/Gasz. In this case, Zuc cleavage does not specify piRNA length;

therefore, trimming by exonucleases such as PNLDC1/PARN is required after Zuc

processing. Blue arrowhead: Zuc cleavage.

STAR★Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal antibody Anti-Daed	This study	P3F10
Mouse monoclonal antibody anti-Gasz	Yamashiro et al.[12]	1H7
Mouse monoclonal antibody anti-Armi	Saito et al.[16]	2F8A9
Mouse monoclonal antibody anti-Piwi	Saito et al.[35]	3G11
Mouse monoclonal antibody anti-Piwi	Saito et al.[35]	4D2
Mouse monoclonal antibody anti-Yb	Murota et al.[55]	1G2H3
Mouse monoclonal antibody anti-Ago1	Okamura et al.[56]	1B8
Mouse monoclonal antibody anti-β-Tub	Developmental Studies Hybridoma Bank	Cat#E7; RRID: AB_528499
HSP60 antibody, Clone LK1	StressMarq Biosciences	Cat# SMC-110 (also SMC-110B), RRID:AB_2121277

Anti-Histon H3 antibody – Nuclear Loading Control and ChIP grade	abcam	Cat# ab1791, RRID:AB_302613
Monoclonal anti-Flag M2 antibody	Sigma-Aldrich	Cat# F3165, RRID:AB_259529
Anti-Flag antibody produces in rabbit	Sigma-Aldrich	Cat# F7425, RRID:AB_439687
Anti-c-Myc antibody produces in rabbit	Sigma-Aldrich	Cat# C3956, RRID:AB_439680
anti-mouse IgG (H+L) HRP	Cappel	55558
anti-rabbit IgG HRP-linked	Cell Signaling Technology	Cat#7074, RRID: AB_2099233
Alexa Fluor 488 goat anti-mouse IgG1 antibody	Thermo Fisher Scientific	Cat#A-21121; RRID: AB_2535764
Alexa Fluor 488 goat anti-mouse IgG2b antibody	Thermo Fisher Scientific	Cat# A-21141 (also A21141), RRID:AB_141626
Alexa Fluor 555 goat anti-mouse IgG1 antibody	Thermo Fisher Scientific	Cat#A-21127; RRID: AB_2535769
Alexa Fluor 555 goat anti-mouse IgG2b antibody	Thermo Fisher Scientific	Cat# A21147, RRID:AB_1500897
Alexa Fluor 633 goat anti-rabbit IgG (H+L) antibody	Thermo Fisher Scientific	Cat# A-21070 (also A21070), RRID:AB_2535731
Bacterial and virus strains		
Escherichia coli Rosetta 2 (DE3)	Novagen	Cat#71397
Chemicals, peptides, and recombinant proteins		
Shields and Sang M3 Insect Medium w/ L-Glutamine	USBiological	Cat#S1013
Fetal Bovine Serum	NICHIREI	Cat#173012
NEBuilder HiFi DNA Assembly Master Mix	New England Biolabs	Cat#E2621X
Ligation high ver.2	Toyobo	Cat#LGK-201
Glutathione Sepharose 4B	Cytiva	Cat#17-0756-01
Phosphate-buffered saline	FUJIFILM Wako Pure Chemical	Cat#048-29805
Thiophilic-Superflow Resin	Clontech	Cat#Z5616N
ClearTrans SP PVDF membranes	Fujifilm Wako Pure Chemical	Cat#033-22453
Clarity Western ECL Substrate	Bio-Rad	Cat#170506
Dynabeads Protein G	Thermo Fisher Scientific	Cat#10009D
RNasin Plus Ribonuclease Inhibitor	Promega	Cat#N2615
T4 Polynucleotide Kinase	New England Biolabs	Cat#M0201L

ISOGEN II	FUJIFILM Wako Pure Chemical	Cat#311-07361
Hybond-N membrane	Cytiva	Cat#RPN303N
VECTASHIELD Antifade Mounting Medium with DAPI	VECTOR LABORATORIES	Cat#H-1200
MitoTracker Orange CMXRos	Thermo Fisher Scientific	Cat#M-7510
SYBR Gold Nucleic Acid Gel Stain	Thermo Fisher Scientific	Cat#S11494
TURBO DNase	Life Technologies	Cat#AM2239
ReverTra Ace qPCR RT master mix	Toyobo	Cat#TYFSQ201
cOmplete ULTRA Tablets, EDTA-free, Protease Inhibitor Cocktail	Roche	Cat#6538282001
RNase A	Roche	Cat#10109142001
Critical commercial assays		
Nucleofector 96-well Shuttle device	Lonza	Cat#AAF-1003S
Nucleofector 2b device	Lonza	Cat#AAB-1001
ChemiDoc XRS Plus System	Bio-Rad	Cat#1708265J1PC
LSM980 laser scanning confocal microscope	Carl Zeiss	Cat#LSM980
StepOnePlus Real-Time PCR system	Thermo Fisher Scientific	Cat#4376598
MiSeq	Illumina	Cat#SY-410-1003
SF Cell Line 96-well Nucleofector Kit	Lonza	Cat#V4SC-3096
QIAseq FastSelect-rRNA Fly Kit	QIAGEN	Cat#333262
PowerUp SYBR Green Master Mix	Thermo Fisher Scientific	Cat# A25743
MiSeq Reagent Kit v3	Illumina	Cat#MS-102-3001
Deposited data		
Raw sequencing data	This paper	GEO: GSExxxxxx
Experimental models: Cell lines		
OSC	Saito et al.[33]	N/A
Oligonucleotides		
Primer sequences	Table S1	N/A
siRNA sequences	Table S2	N/A
Probe sequences	Table S3	N/A
qPCR primer sequences	Table S4	N/A
Recombinant DNA		
pGEX	Cytiva	Cat#28-9546-48
pAc-Flag-Daed WT and mutants	This study	N/A

pAc-Flag-Armi WT	Ishizu et al.[50]	N/A
pAc-Flag-Gasz WT	Yamashiro et al.[12]	N/A
pAc-Flag-Gasz mutants	This study	N/A
pAc-Armi WT-Flag	Yamashiro et al.[12]	N/A
pAc-Armi ∆N34-Flag	Yamashiro et al.[12]	N/A
pAc-Armi KEE-Flag	This study	N/A
pAc-Myc-Luc	gift from H. Ishizu	N/A
pAc- Luc-Flag	Yamashiro et al.[12]	N/A
pAc-Flag-Luc	This study	N/A
pAc-MOV10L1-Flag	This study	N/A
pAc-Myc-ASZ1	This study	N/A
Software and algorithms		
Python (Ver. 3.8.8)	Van Rossum & Drake[57]	RRID:SCR_008394;https://www.python.org
Numpy (Ver. 1.20.1)	Harris et al.[58]	RRID:SCR_008633; https://numpy.org/ja/
Scipy (Ver. 1.6.2)	Virtanen et al.[59]	RRID:SCR_008058; https://scipy.org
Pandas (Ver. 1.2.4)	McKinney[60]	RRID:SCR_018214;https://pandas.pydata.org
R (Ver. 4.3.1)	R Development Core Team[61]	RRID:SCR_001905;https://www.r-project.org
Cutadapt (Ver. 4.0)	Martin[62]	RRID:SCR_011841;https://cutadapt.readthedocs.io/en/stable/
STAR (Ver.2.7.8a)	Dobin et al.[63]	RRID:SCR_004463;https://github.com/alexdobin/STAR
HTSeq (Ver. 0.9.1)	Anders et al.[64]	RRID:SCR_005514;https://htseq.readthedocs.io/en/latest/
bedtools (Ver. 2.30.0)	Quinlan & Hall[65]	RRID:SCR_006646;https://bedtools.readthedocs.io/en/latest/
SAMtools (Ver. 1.9)	Li et al.[66]	RRID:SCR_002105;http://www.htslib.org
deepTools (Ver. 3.3.2)	Ramírez et al.[67]	RRID:SCR_016366;https://deeptools.readthedocs.io/en/develop/
wigToBigWig (Ver. 357)	UCSC	https://www.encodeproject.org/software/wigtobigwig/

ggseqlogo	Wagih.[68]	https://omarwagih.github.io/ggseqlogo/
RSeQC (Ver. 2.6.4)	Wang et al.[69]	RRID:SCR_005275;https://rseqc.sourceforge.net
Fiji (Ver. 2.0.0-rc-69/1.52p)	Schneider et al.[70]	RRID:SCR_002285; https://fiji.sc
AlphaFold (Ver. 2.3)	Jumper et al.[31]	https://github.com/google-deepmind/alphafold
UCSF ChimeraX (Ver. 1.6.1)	Pettersen et al.[71]	RRID:SCR_015872;https://www.cgl.ucsf.edu/chimerax/
ISOLDE (Ver. 1.6)	Croll[72]	https://tristanic.github.io/isolde/
CLUSTAL Omega	Madeira et al.[73]	RRID:SCR_001591;https://www.ebi.ac.uk/jdispatcher/msa/clustalo
iTOL (Ver. 6.9)	Letunic & Bork[74]	RRID:SCR_018174;https://itol.embl.de
seaborn (Ver. 0.12.2)	Waskom et al.[75]	https://seaborn.pydata.org
ESPript (Ver. 3.0)	Robert & Gouet[76]	https://espript.ibcp.fr
MEGA11	Tamura et al.[77]	RRID:SCR_023017; https://www.megasoftware.net/

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Mikiko C. Siomi (siomim@bs.s.u-tokyo.ac.jp).

Materials availability

Plasmids and antibodies generated in this study will be shared by the lead contact upon

request.

Data and code availability

The data supporting the findings of this study are available from the corresponding

authors upon reasonable request. The datasets produced in this study are available in

the following databases: RNA-Seq data: Gene Expression Omnibus GSExxxxxx

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSExxxxxx).

Experimental model and study participant details

OSCs were cultured at 26°C in Shields and Sang M3 medium (USBiological)

supplemented with 10% fetal bovine serum (FBS, NICHIREI), 0.6 mg/mL glutathione, 10

mU/mL insulin, and 10% fly extract.[8]

Method details

Plasmid construction

The Daed coding region was amplified from total OSC RNAs by RT-PCR and inserted

into pAc-Flag-Armi WT[50] by substituting the Armi coding region using NEBuilder HiFi

DNA Assembly Master Mix (New England Biolabs). RNAi-resistant Daed WT and its

ΔSAM and ΔCC mutants were generated by inverse PCR, according to the domain

structures described in a previous study.[29] To construct the expression vector for GST-

Daed ΔTM-His, corresponding coding regions of Daed were cloned into pGEX, after

which a His-Tag was added to the C-terminal by inverse PCR. RNAi-resistant Flag-Gasz

mutants (ΔAR1-3, ΔAR4-6, ΔSAM, and Δα-H2) were generated by inverse PCR using

RNAi-resistant Flag-Gasz WT vector[12] as a template. Vectors for RNAi-resistant Armi

WT-Flag and its ΔN34 mutant were previously constructed.[12] The Armi KEE-Flag mutant

was produced by inverse PCR using RNAi-resistant Armi WT-Flag vector as a template.

The Flag-Luc expression vector was generated from pAc-Myc-Luc (a gift from H. Ishizu)

by substituting Myc with 3×Flag. The primers used are shown in Table S1. The MOV10L1

and ASZ1 coding region was amplified from total mouse ovary RNAs by RT-PCR and

inserted into pAc-Armi WT-Flag and Myc-Daed WT by substituting original coding

regions.

RNAi and transfection

To initiate RNAi, 5.0×10^6 OSCs were mixed with 200 pmol of siRNAs in 20 μL of Solution

SF of the Cell Line Nucleofector Kit SF (Lonza Bioscience) and electroporation was

performed using a Nucleofector 96-well Shuttle device (Lonza Bioscience). For plasmid

transfection or plasmid/siRNA co-transfection, 1.5×10^7 OSCs were mixed with 1.5–10

μg of plasmids with or without 600 pmol of siRNAs in 100 μL of transfection buffer [180

mM sodium phosphate buffer for Church and Gilbert hybridization (pH 7.2), 5 mM KCl,

15 mM $MgCl_2$, and 50 mM D-mannitol] modified from a previous report,[78] and

electroporation was performed by a Nucleofector 2b device (Lonza Bioscience). Cells

were transfected with siRNAs twice with a 48-h interval. For rescue assays, endogenous

genes were depleted by RNAi and plasmids and siRNAs were introduced into the cells

after 48 h of culture. Cells were harvested after another 48 h of culture. For

immunofluorescence in Figure 2F, 500 pmol of siRNAs were used for RNAi and

plasmid/siRNA co-transfection, and cells were harvested after another 72 h of culture.

Sequences of siRNAs are indicated in Table S2.

Production of anti-Daed monoclonal antibody

GST-Daed (Met1–Asp197) was expressed in *E. coli* (Rosetta-DE3) and the inclusion

bodies were used for mouse immunization. Production and selection of hybridomas that

produce anti-Daed monoclonal antibodies were performed as described previously.[79]

Mitochondrial fractionation

Mitochondria were fractionated as previously reported.[12] Briefly, 1.2×10^8 OSCs were

suspended in 500 µL of buffer A [30 mM Tris-HCl (pH 7.3), 225 mM D-mannitol, 75 mM

sucrose, and 0.05 mM EGTA] and the cell suspension was passed 30 times through a

30-gauge needle attached to a 1-mL syringe. The lysates were centrifuged at 600 g for

5 min and the supernatants were collected, and repeated twice. The lysates were then

centrifuged at 7,000 g for 10 min and the supernatants were collected as the cytoplasmic

fraction. This step was also repeated twice. The pellets were resuspended in 500 µL of

buffer B [30 mM Tris-HCl (pH 7.3), 225 mM D-mannitol, and 75 mM sucrose] and

centrifuged at 10,000 g for 10 min. The supernatant was used as the mitochondrial

fraction.

Nuclear fractionation

Nuclear fractionation was performed as previously reported[24] with some modifications.

In brief, cells were suspended in hypotonic buffer [10 mM HEPES-KOH (pH 7.3), 10 mM

KCl, 1.5 mM $MgCl_2$, 0.5 mM dithiothreitol (DTT), 2 µg/mL leupeptin, 2 µg/mL pepstatin

A, and 0.5% aprotinin] by pipetting six times and passed 3 times through a 25-gauge

needle attached to a 1-mL syringe. The lysates were centrifuged at 400 g for 10 min to

collect nuclei. The supernatants were collected as the cytoplasmic fraction, then the

concentration of KCl was adjusted to 200 mM and centrifuged at 20,000 g for 20 min

before used for Piwi-piRISC isolation (see below). The nuclear fraction was washed with

hypotonic buffer and resuspended in chromatin co-immunoprecipitation buffer [50 mM

HEPES-KOH (pH 7.3), 200 mM KCl, 1 mM EDTA, 1% Triton X-100, 0.1% Na

deoxycholate, 2 µg/mL leupeptin, 2 µg/mL pepstatin A, and 0.5% aprotinin]. The nuclear

fraction was sonicated and centrifuged at 20,000 g for 20 min. The supernatant was

collected as nuclear extract fraction for Piwi-piRISC isolation.

Western blotting

Western blotting was performed essentially as described previously.[80] Proteins

separated on SDS polyacrylamide gels were transferred to ClearTrans SP PVDF

membranes (Fujifilm Wako Pure Chemical) and incubated in PBS containing 0.1%

Tween-20 (T-PBS) supplemented with 5% skimmed milk. After washing with T-PBS, the

membranes were incubated with the following primary antibodies: anti-Daed (1:1,000

dilution, this study), anti-Gasz (1:1,000 dilution),[12] anti-Armi (2F8A9, 1:1,000 dilution),[16]

anti-Piwi (3G11, 1:1,000 dilution),[35] anti-β-Tub (1:1,000 dilution, E7, Developmental

Studies Hybridoma Bank), anti-HSP60(1:500 dilution, LK1, StressMarq Biosciences),

anti-Flag M2 (1:5,000 dilution, F3165, Sigma), anti-Histon H3 (1:5,000 dilution, ab1791,

abcam) monoclonal and anti-c-Myc (1:1,000 dilution, C3956, Sigma) polycronal

antibodies. After washing with T-PBS, the membranes were incubated with peroxidase-

conjugated anti-mouse IgG antibodies (1:5,000 dilution, 55558, Cappel) or anti-rabbit

IgG antibodies (1:5,000 dilution, 7074, Cell Signaling Technology). The membranes

were then reacted with Clarity Western ECL Substrate (Bio-Rad) and images were

collected using ChemiDoc XRS Plus System (Bio-Rad). The signal intensity was

quantified by Fiji (Ver. 2.0.0-rc-69/1.52p).[70]

Immunofluorescence

Immunofluorescence was performed as described previously.[16] OSCs were fixed with

4% formaldehyde in PBS. The cells were permeabilized with PBS containing 0.1% Triton

X-100, blocked with PBS containing 3% BSA, and incubated with the following primary

antibodies: anti-Armi (2F8A9, 1:200 dilution),[16] anti-Piwi (4D2, 1:200 dilution,[35] anti-Yb

(1G2H3, 1:200 dilution),[55] anti-Flag M2 (F3165, 1:1,000 dilution, Sigma) monoclonal and

anti-c-Myc (1:300 dilution, C3956, Sigma) polyclonal antibodies. Anti-Flag polyclonal

antibodies (F7425, 1:300 dilution, Sigma) were used to obtain the results shown in Fig.

2F. After washing with PBS, cells were incubated with the following secondary antibodies

(1:1,000 dilution): Alexa 488-conjugated anti-mouse IgG1, Alexa 488-conjugated anti-

mouse IgG2b, Alexa 555-conjugated anti-mouse IgG1, Alexa 555-conjugated anti-

mouse IgG2b, and Alexa 633-conjugated anti-rabbit IgG (all from Thermo Fisher

Scientific). Mitochondria were stained with MitoTracker Orange CMXRos (Thermo Fisher

Scientific) before fixation. Cells were mounted with VECTASHIELD Mounting Medium

with DAPI (Vector Laboratories). Images were obtained using LSM980 laser scanning

confocal microscope (Carl Zeiss).

Immunoprecipitation

Immunoprecipitation was performed as described previously.[16] Cells were suspended in

binding buffer A [30 mM HEPES-KOH (pH 7.3), 150 mM KOAc, 5 mM Mg(OAc)$_2$, 5 mM

DTT, 1% Triton X-100, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]

supplemented with 0.04 U/μL RNasin Plus RNase Inhibitor (Promega) and the cell

suspension was passed 10 times through a 30-gauge needle and centrifuged. The

obtained supernatants (cell lysates) were incubated at 4°C with Dynabeads Protein G

(Thermo Fisher Scientific) pre-bound with antibodies. After 1 h of incubation, the beads

were washed with binding buffer and the materials bound on the beads were analyzed.

Piwi-piRISC isolation from OSCs

Cells were lysed by sonication in binding buffer B [50 mM HEPES-KOH (pH 7.3), 200

mM KCl, 1 mM EDTA, 1 mM DTT, 1% Triton X-100, 0.1% Na deoxycholate, 2 μg/mL

leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin] supplemented with 0.1 U/μL RNasin

Plus RNase Inhibitor. Cell lysates were then incubated with anti-Piwi monoclonal

antibodies (3G11) pre-bound to Dynabeads Protein G at 4°C for 2 h. The beads were

washed with IP-wash buffer [20 mM HEPES-KOH (pH 7.3), 300 mM NaCl, 1 mM DTT,

0.05% NP-40, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin] and with

high-salt IP-wash buffer [20 mM HEPES-KOH (pH 7.3), 500 mM NaCl, 1 mM DTT, 0.05%

NP-40, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]. RNAs were eluted

from the beads with phenol-chloroform and precipitated with ethanol. Eluted RNAs were

separated on denaturing polyacrylamide gels and detected by SYBR Gold Nucleic Acid

Gel Stain (Thermo Fisher Scientific). Signal intensity was calculated using Fiji, and

displayed using Python3 and seaborn[75] (Ver.0.12.2). For piRNA sequencing, cells were

lysed in binding buffer C [20 mM HEPES-KOH (pH 7.3), 150 mM NaCl, 1mM EDTA, 1

mM DTT, 0.5% NP-40, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, and 0.5% aprotinin]

supplemented with 0.04 U/μL RNasin Plus RNase Inhibitor. RNAs were separated on

denaturing polyacrylamide gels and 20–40-nt RNAs were eluted from the gels and

cloned as described previously.[8] QIAseq FastSelect-rRNA Fly Kit (QIAGEN) was used

to remove rRNA from the library.

Northern blotting

Total RNAs were isolated from OSCs using ISOGEN II (Fujifilm Wako Pure Chemical)

and treated with 0.04 U/μL TURBO DNase (Life Technologies). Northern blotting was

carried out as described previously.[81] Total RNAs were separated on denaturing

polyacrylamide gels, blotted on Hybond-N+ hybridization membranes (Cytiva), and UV

cross-linked. The membranes were incubated at 42°C for 30 min in hybridization buffer

[200 mM sodium phosphate buffer (pH 7.2), 7% SDS, and 1 mM EDTA] and probed with

^{32}P-labeled oligonucleotides at 42°C overnight. The membranes were washed with NB-

wash buffer [30 mM sodium citrate (pH 7.0), 300 mM NaCl, and 0.1% SDS] at 42°C and

scanned using Typhoon FLA 9500 (Cytiva). Sequences of oligodeoxynucleotides are

shown in Table S3.

RT-qPCR

Total RNAs from OSCs were reverse-transcribed using ReverTra Ace (Toyobo).

Resulting cDNAs were amplified with StepOnePlus (Thermo Fisher Scientific) using

PowerUp SYBR Green Master Mix (Thermo Fisher Scientific). Sequences of primers are

shown in Table S4.

Recombinant proteins

GST-Daed ΔTM-His was expressed in *E. coli* (Rosetta DE3) at 16°C overnight. *E. coli*

was harvested and lysed by sonication in lysis buffer [30 mM Tris-HCl (pH 8.0), 500 mM

NaCl, 1% Triton X-100, 10 mM imidazole, 1× cOmplete ULTRA Tablets (Roche), and 1

mM DTT]. The supernatants after centrifugation were bound to Ni-Sepharose 6 Fast

Flow (Cytiva) at 4°C for 1 h. After washing with wash buffer A [30 mM Tris-HCl (pH 8.0),

500 mM NaCl, 1% Triton X-100, 10 mM imidazole, and 1 mM DTT], proteins were eluted

from the beads with elution buffer A [30 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% Triton

X-100, 250 mM imidazole, and 1 mM DTT]. Eluted proteins were bound to Glutathione

Sepharose 4B (Cytiva) at 4°C for 1 h. After washing with wash buffer B [30 mM Tris-HCl

(pH 8.0), 500 mM NaCl, 1% Triton X-100, and 1 mM DTT], proteins were eluted from the

beads with elution buffer B [30 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% Triton X-100, 1

mM DTT, and 15 mM glutathione], dialyzed in dialysis buffer [30 mM Tris-HCl (pH 8.0),

150 mM NaCl, 1 mM DTT, 10% glycerol, and 8 μg/mL GST-HRV3C protease], and

passed through Glutathione Sepharose 4B to remove the protease and GST.

Pull-down assay

OSCs expressing Armi WT-Flag and its ΔN34 mutant were suspended with bait-

preparing buffer [30 mM HEPES-KOH (pH 7.3), 150 mM KOAc, 500 mM NaCl, 5 mM

Mg(OAc)$_2$, 5 mM DTT, 1% Triton X-100, 2 μg/mL leupeptin, 2 μg/mL pepstatin A, 0.5%

aprotinin, and 25 μg/mL RNase A (Roche)]. The cell suspension was passed five times

through a 25-gauge needle and another five times through a 30-gauge needle, and then

centrifuged. The supernatants were incubated at 4°C for 1 h with anti-Flag antibodies

pre-bound onto Dynabeads Protein G. Beads were washed with bait-preparing buffer

and then binding buffer D [30 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM DTT, and 1%

Triton X-100]. Daed ΔTM-His was added to the beads and incubated at 4°C for 1.5 h.

After washing with binding buffer, proteins were eluted from the beads, separated on an

SDS polyacrylamide gel, and stained with CBB.

β-elimination assay

β-elimination was performed as described previously with some modifications.[36] piRNAs

were isolated from Piwi immunoprecipitated from OSCs. miRNAs were isolated from

Ago1 immunoprecipitated from OSCs using monoclonal anti-Ago1 antibodies[56]. RNAs

were incubated in 10 mM NaIO$_4$ for 40 min at 0°C in the dark. After ethanol precipitation,

RNAs were dissolved in 1 M L(+)-lysine-HCl (pH 8.5) and incubated for 90 min at 45°C.

After ethanol precipitation, RNAs were separated on a denaturing polyacrylamide gel

and detected by SYBR Gold Nucleic Acid Gel Stain.

piRNA sequencing and bioinformatic analysis

piRNAs were sequenced using MiSeq (Illumina) with MiSeq Reagent Kit v3 (Illumina).

piRNA-seq analysis was performed as described previously[8] with some modifications.

Adapter-trimmed reads of ≥ 18 nt in length were mapped against the *Drosophila* genome

(BDGP Release 6 + ISO1 MT/dm6) with FlyBase gene model (Release 6.21) using

STAR[63] (Ver. 2.7.8a). Uniquely mapped reads with up to one mismatch were used for

the subsequent analysis. Read length of each biological triplicate was used in the

analysis in Figs. 4B and S4I. After confirming high correlations between samples (Figure

S6A, B), reads were merged and used for subsequent analysis. Reads mapped to exons

of coding genes (Figure S6A, B) were counted with HTSeq[64] (Ver. 0.9.1) using

annotation file obtained from FlyBase and normalized to RPM+1 reads. Spearman's rho

was calculated using R[61] (Ver. 4.3.1). For transposon annotation (Fig. S4B,C), the

annotation file of repeat sequences was obtained from RepeatMasker (dm6-Apr 2006-

RepeatMasker open-4.0.6-Dfam 2.0) and transposons were extracted without distinction

between int and LTR. Reads mapped antisense to transposons were counted using

HTSeq. For the analysis in Fig. S4C, reads were normalized to RPM+1 reads. For the

analysis in Fig. S4B, transposons which mapped > 5 RPM reads in the sum of WT and

ΔSAM were selected, and the mean length of mapped reads for each transposon were

calculated. For 1U bias and +1U bias analyses (Figs. 4C, I and S4J), the 5′ ends and

their downstream sequences of piRNAs (1U bias) or *Drosophila* genome sequences

around the 3′ ends (+1U bias) were obtained using bedtools[65] (Ver. 2.30.0) and SAMtools

(Ver. 1.9).[66] Sequence logos were generated using ggseqlogo.[68] For the phasing

analysis (Fig. 4D), 3′ end positions around piRNA 5′ end position were normalized to

RPM and showed in heatmaps using RSeQC[69] (Ver. 2.6.4), deepTools[67] (Ver. 3.3.2),

SAMtools[66] (Ver. 1.9) and "wigToBigWig" from UCSC. Binary histograms were

generated using the matrix of heatmaps. Values were transformed into binary format

(counts = 0 → 0, counts > 0 → 1) in accordance with a previous study.[18] For the

comparison of 3′ end position in WT and mutant libraries of Daed or Armi (Figs. 4H and

S4K), 50,000 piRNA 5′ end positions, common to the WT and mutant libraries and most

abundant in the WT library, were extracted. The percentage of 3′ end positions

corresponding to each 5′ end position was calculated and shown in heatmaps. 5′ end

positions were ordered by the 3′ end position of WT library data and the same order was

maintained in the mutant library data. To obtain the results shown in Fig. S4G, 5′-end

positions that dominantly produce 26- or 27-nt piRNAs in the WT library were extracted,

and after combining, these piRNAs were classified as "Unchanged" (51,833 positions)

and "Elongated" (53,258 positions) according to 3′-end elongation caused by Daed SAM

loss and used for further analysis.

piRISC production simulation

The *flam* sequence was downloaded from FlyBase (Release FB2023_05) (21,629,891

to 21,792,731 on the X chromosome) and subjected to the digestion into 163 piRNA

precursors (starting at U and around 1,000 nt). The precursors were cleaved to 18–40-

nt piRNAs immediately before U. The cleavage position was determined based on the

distribution of the actual piRNA length, which was obtained by fitting a normal distribution

to the piRNA sequencing data. The simulation was performed using Python3[57] (Ver.

3.8.8) with add-on libraries [Numpy[58] (Ver. 1.20.1), Scipy[59] (Ver. 1.6.2), and Pandas[60]

(Ver. 1.2.4)].

Structure prediction

A structure model of the AmTEC (Fig. 1A) and Armi–AmTEC complex (Fig. 3A) was built

by structure prediction with ColabFold[82] (Ver. 1.5.5) and AlphaFold[31] (Ver. 2.3) using

protein amino acid sequences from FlyBase (Release FB2023_04). Five complex

models were built, and structural predictions and optimization were performed five times

for each complex model. The structure model with the highest pLDDT value underwent

relax steps. The N-terminal disorder region and TM domain orientations of Daed and

Gasz were appropriately changed at the linker region using ISOLDE[72] (Ver. 1.6). The

structure was visualized using UCSF ChimeraX[71] (Ver. 1.6.1).

Ortholog search and protein sequence alignment

Orthologs for Armi/MOV10L1, Gasz/ASZ1, PNLDC1, Papi/TDRKH, Nbr were searched

using OrthoDB v11. Orthologs for Daed were searched using OrthoDB v11[38] and NCBI

PSI-Blast. Species taxonomy was according to NCBI and phylogenetic tree (Figure 5A)

were generated following previous studies.[84,85] Multiple sequence alignment (Figure

S5B) was conducted using CLUSTAL Omega multiple alignment[73] and visualized using

ESPript (Ver. 3.0).[76] The phylogenetic tree (Figure 5SA) was generated using Neighbor-

Joining method[83] in MEGA11[77]. Multiple sequence alignment (Figure S5C) was

conducted using CLUSTAL Omega multiple alignment[73] and visualized using iTOL (Ver.

6.9).[74] The following protein sequences were acquired from NCBI: *D. melanogaster* Armi

(NP_001014556), *H. sapiens* Moloney leukemia virus 10-like protein 1 (MOV10L1)

(NP_061868.1), *M. musculus* MOV10L1 (XP_006521619.1), *G. gallus* MOV10L1

(XP_040514825.1), *X. tropicalis* MOV10L1 (NP_001072624.1), *D. rerio* MOV10L1

(NP_001070795.1), *Z. nevadensis* Armi (XP_021937793.1), *T. palmi* Armi

(XP_034251738.1), *A. echinatior* Armi (XP_011062616.1), *B. terrestris* Armi

(XP_012166280.1), *D. ponderosae* Armi (XP_048525683.1), *T. castaneum* Armi

(XP_008199251.1), *B. anynana* Armi (XP_023946786.2), *B. mori* Armi

(XP_037867032.1), *A. gambiae* Armi (XP_061506580.1), *A. aegypti* Armi

(XP_021698921.1), *B. dorsalis* Armi (XP_011214121.1), *C. capitata* Armi

(XP_004537921.1), *M. domestica* Armi (XP_058975516.1) and *S. calcitrans* Armi

(XP_013110641.2).

Quantification and statistical analysis

All experiments were performed over two independent experiments (biological

replicates). Sample sizes were predetermined following previously performed

experiments with similar setup in the field. Statistical methods, *p* value and number of

replicates are indicated in each figure and its legend. Statistical analyses were performed

in the R software environment (version 4.3.1). Shapiro-Wilk test was used for testing

normality when statistical methods based on the normal distribution was used.

Experiments were not blinded as consistent with the previous studies in the field.

Supplemental figure titles

Figure S1. Related to Figure1; Coiled-coil prediction and specificity of anti-Daed

antibodies

Figure S2. Related to Figure2; Functions and behaviors of Daed and Gasz in OSCs

Figure S3. Related to Figure3; Effects of Armi mutants on piRNA biogenesis and Piwi-

piRISC function

Figure S4. Related to Figure4; Effects of Daed and Armi mutants on piRNA biogenesis

and Piwi-piRISC function

Figure S5. Related to Figure5; Sequence alignment of Daed and Armi UPF1-like domain

in species

Figure S6. Related to STAR Method "piRNA sequencing and bioinformatic analysis";

Reproducibility of piRNA-seq

Supplemental table titles

Table S1. Related to STAR Method "Plasmid construction"; Sequences of primers used

for plasmid construction

Table S2. Related to STAR Method "RNAi and transfection"; Sequences of siRNAs

Table S3. Related to STAR Method "Northern blotting"; Sequences of oligos used for

northern blotting

Table S4. Related to STAR Method "RT-qPCR"; Sequences of primers used for RT-

qPCR

References

1.   Czech, B., Munafò, M., Ciabrelli, F., Eastwood, E.L., Fabry, M.H., Kneuss, E., and
Hannon, G.J. (2018). piRNA-Guided Genome Defense: From Biogenesis to
Silencing. Annual Review of Genetics *52*, 131–157.
https://doi.org/10.1146/annurev-genet-120417-031441.
2.   Huang, X., Fejes Tóth, K., and Aravin, A.A. (2017). piRNA Biogenesis in
Drosophila melanogaster. Trends in Genetics *33*, 882–894.
https://doi.org/10.1016/j.tig.2017.09.002.
3.   Ozata, D.M., Gainetdinov, I., Zoch, A., O'Carroll, D., and Zamore, P.D. (2019).

832   PIWI-interacting RNAs: small RNAs with big functions. Nat Rev Genet *20*, 89–108.

833   https://doi.org/10.1038/s41576-018-0073-3.

834  4. Yamashiro, H., and Siomi, M.C. (2018). PIWI-Interacting RNA in Drosophila:

835   Biogenesis, Transposon Regulation, and Beyond. Chem Rev *118*, 4404–4421.

836   https://doi.org/10.1021/acs.chemrev.7b00393.

837  5. Brennecke, J., Aravin, A.A., Stark, A., Dus, M., Kellis, M., Sachidanandam, R., and

838   Hannon, G.J. (2007). Discrete Small RNA-Generating Loci as Master Regulators

839   of Transposon Activity in Drosophila. Cell *128*, 1089–1103.

840   https://doi.org/10.1016/j.cell.2007.01.043.

841  6. Malone, C.D., Brennecke, J., Dus, M., Stark, A., McCombie, W.R.,

842   Sachidanandam, R., and Hannon, G.J. (2009). Specialized piRNA Pathways Act in

843   Germline and Somatic Tissues of the Drosophila Ovary. Cell *137*, 522–535.

844   https://doi.org/10.1016/j.cell.2009.03.040.

845  7. Goriaux, C., Desset, S., Renaud, Y., Vaury, C., and Brasset, E. (2014).

846   Transcriptional properties and splicing of the flamenco piRNA cluster. EMBO

847   reports *15*, 411–418. https://doi.org/10.1002/embr.201337898.

848  8. Hirakata, S., Ishizu, H., Fujita, A., Tomoe, Y., and Siomi, M.C. (2019).

849   Requirements for multivalent Yb body assembly in transposon silencing in

850   Drosophila. EMBO Rep *20*, e47708. https://doi.org/10.15252/embr.201947708.

851  9. Ishizu, H., Iwasaki, Y.W., Hirakata, S., Ozaki, H., Iwasaki, W., Siomi, H., and

852   Siomi, M.C. (2015). Somatic Primary piRNA Biogenesis Driven by cis-Acting RNA

853   Elements and trans-Acting Yb. Cell Reports *12*, 429–440.

854   https://doi.org/10.1016/j.celrep.2015.06.035.

855  10. Cora, E., Pandey, R.R., Xiol, J., Taylor, J., Sachidanandam, R., McCarthy, A.A.,

856   and Pillai, R.S. (2014). The MID-PIWI module of Piwi proteins specifies nucleotide-

857   and strand-biases of piRNAs. RNA *20*, 773–781.

858   https://doi.org/10.1261/rna.044701.114.

859  11. Stein, C.B., Genzor, P., Mitra, S., Elchert, A.R., Ipsaro, J.J., Benner, L., Sobti, S.,

860   Su, Y., Hammell, M., Joshua-Tor, L., et al. (2019). Decoding the 5′ nucleotide bias

861   of PIWI-interacting RNAs. Nat Commun *10*, 828. https://doi.org/10.1038/s41467-

862   019-08803-z.

863  12. Yamashiro, H., Negishi, M., Kinoshita, T., Ishizu, H., Ohtani, H., and Siomi, M.C.

864   (2020). Armitage determines Piwi–piRISC processing from precursor formation

865   and quality control to inter-organelle translocation. EMBO reports *21*, e48769.

866   https://doi.org/10.15252/embr.201948769.

867  13. Choi, S.-Y., Huang, P., Jenkins, G.M., Chan, D.C., Schiller, J., and Frohman, M.A.

868  (2006). A common lipid links Mfn-mediated mitochondrial fusion and SNARE-

869  regulated exocytosis. Nat Cell Biol *8*, 1255–1262. https://doi.org/10.1038/ncb1487.

870 14. Ipsaro, J.J., Haase, A.D., Knott, S.R., Joshua-Tor, L., and Hannon, G.J. (2012).

871  The structural biochemistry of Zucchini implicates it as a nuclease in piRNA

872  biogenesis. Nature *491*, 279–283. https://doi.org/10.1038/nature11502.

873 15. Nishimasu, H., Ishizu, H., Saito, K., Fukuhara, S., Kamatani, M.K., Bonnefond, L.,

874  Matsumoto, N., Nishizawa, T., Nakanaga, K., Aoki, J., et al. (2012). Structure and

875  function of Zucchini endoribonuclease in piRNA biogenesis. Nature *491*, 284–287.

876  https://doi.org/10.1038/nature11509.

877 16. Saito, K., Ishizu, H., Komai, M., Kotani, H., Kawamura, Y., Nishida, K.M., Siomi,

878  H., and Siomi, M.C. (2010). Roles for the Yb body components Armitage and Yb in

879  primary piRNA biogenesis in Drosophila. Genes Dev *24*, 2493–2498.

880  https://doi.org/10.1101/gad.1989510.

881 17. Han, B.W., Wang, W., Li, C., Weng, Z., and Zamore, P.D. (2015). Noncoding RNA.

882  piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA

883  production. Science *348*, 817–821. https://doi.org/10.1126/science.aaa1264.

884 18. Mohn, F., Handler, D., and Brennecke, J. (2015). piRNA-guided slicing specifies

885  transcripts for Zucchini-dependent, phased piRNA biogenesis. Science *348*, 812–

886  817. https://doi.org/10.1126/science.aaa1039.

887 19. Yashiro, R., Murota, Y., Nishida, K.M., Yamashiro, H., Fujii, K., Ogai, A.,

888  Yamanaka, S., Negishi, L., Siomi, H., and Siomi, M.C. (2018). Piwi Nuclear

889  Localization and Its Regulatory Mechanism in Drosophila Ovarian Somatic Cells.

890  Cell Reports *23*, 3647–3657. https://doi.org/10.1016/j.celrep.2018.05.051.

891 20. Dönertas, D., Sienski, G., and Brennecke, J. (2013). Drosophila Gtsf1 is an

892  essential component of the Piwi-mediated transcriptional silencing complex. Genes

893  Dev *27*, 1693–1705. https://doi.org/10.1101/gad.221150.113.

894 21. Fabry, M.H., Ciabrelli, F., Munafò, M., Eastwood, E.L., Kneuss, E., Falciatori, I.,

895  Falconio, F.A., Hannon, G.J., and Czech, B. (2019). piRNA-guided co-

896  transcriptional silencing coopts nuclear export factors. eLife *8*, e47999.

897  https://doi.org/10.7554/eLife.47999.

898 22. Murano, K., Iwasaki, Y.W., Ishizu, H., Mashiko, A., Shibuya, A., Kondo, S., Adachi,

899  S., Suzuki, S., Saito, K., Natsume, T., et al. (2019). Nuclear RNA export factor

900  variant initiates piRNA-guided co-transcriptional silencing. The EMBO Journal *38*,

901  e102870. https://doi.org/10.15252/embj.2019102870.

902 23. Ohtani, H., Iwasaki, Y.W., Shibuya, A., Siomi, H., Siomi, M.C., and Saito, K.

903  (2013). DmGTSF1 is necessary for Piwi–piRISC-mediated transcriptional

transposon silencing in the Drosophila ovary. Genes Dev *27*, 1656–1661.

https://doi.org/10.1101/gad.221515.113.

24. Onishi, R., Sato, K., Murano, K., Negishi, L., Siomi, H., and Siomi, M.C. (2020).

Piwi suppresses transcription of Brahma-dependent transposons via Maelstrom in

ovarian somatic cells. Science Advances *6*, eaaz7420.

https://doi.org/10.1126/sciadv.aaz7420.

25. Sienski, G., Dönertas, D., and Brennecke, J. (2012). Transcriptional silencing of

transposons by Piwi and maelstrom and its impact on chromatin state and gene

expression. Cell *151*, 964–980. https://doi.org/10.1016/j.cell.2012.10.040.

26. Sienski, G., Batki, J., Senti, K.-A., Dönertas, D., Tirian, L., Meixner, K., and

Brennecke, J. (2015). Silencio/CG9754 connects the Piwi–piRNA complex to the

cellular heterochromatin machinery. Genes Dev *29*, 2258–2271.

https://doi.org/10.1101/gad.271908.115.

27. Yu, Y., Gu, J., Jin, Y., Luo, Y., Preall, J.B., Ma, J., Czech, B., and Hannon, G.J.

(2015). Panoramix enforces piRNA-dependent cotranscriptional silencing. Science

350, 339–342. https://doi.org/10.1126/science.aab0700.

28. Czech, B., Preall, J.B., McGinn, J., and Hannon, G.J. (2013). A transcriptome-wide

RNAi screen in the Drosophila ovary reveals factors of the germline piRNA

pathway. Mol Cell *50*, 749–761. https://doi.org/10.1016/j.molcel.2013.04.007.

29. Munafò, M., Manelli, V., Falconio, F.A., Sawle, A., Kneuss, E., Eastwood, E.L.,

Seah, J.W.E., Czech, B., and Hannon, G.J. (2019). Daedalus and Gasz recruit

Armitage to mitochondria, bringing piRNA precursors to the biogenesis machinery.

Genes Dev. *33*, 844–856. https://doi.org/10.1101/gad.325662.119.

30. Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A.,

Bates, R., Blackwell, S., Yim, J., et al. (2022). Protein complex prediction with

AlphaFold-Multimer. Preprint at bioRxiv,

https://doi.org/10.1101/2021.10.04.463034

https://doi.org/10.1101/2021.10.04.463034.

31. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly

accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.

https://doi.org/10.1038/s41586-021-03819-2.

32. Madeo, G., Savojardo, C., Manfredi, M., Martelli, P.L., and Casadio, R. (2023).

CoCoNat: a novel method based on deep learning for coiled-coil prediction.

Bioinformatics *39*, btad495. https://doi.org/10.1093/bioinformatics/btad495.

33. Saito, K., Inagaki, S., Mituyama, T., Kawamura, Y., Ono, Y., Sakota, E., Kotani, H.,

940  Asai, K., Siomi, H., and Siomi, M.C. (2009). A regulatory circuit for piwi by the large
941  Maf gene traffic jam in Drosophila. Nature *461*, 1296–1299.
942  https://doi.org/10.1038/nature08501.

943 34. Olivieri, D., Sykora, M.M., Sachidanandam, R., Mechtler, K., and Brennecke, J.
944  (2010). An in vivo RNAi assay identifies major genetic and cellular requirements
945  for primary piRNA biogenesis in Drosophila. EMBO J *29*, 3301–3317.
946  https://doi.org/10.1038/emboj.2010.212.

947 35. Saito, K., Nishida, K.M., Mori, T., Kawamura, Y., Miyoshi, K., Nagami, T., Siomi,
948  H., and Siomi, M.C. (2006). Specific association of Piwi with rasiRNAs derived
949  from retrotransposon and heterochromatic regions in the Drosophila genome.
950  Genes Dev *20*, 2214–2222. https://doi.org/10.1101/gad.1454806.

951 36. Gunawardane, L.S., Saito, K., Nishida, K.M., Miyoshi, K., Kawamura, Y., Nagami,
952  T., Siomi, H., and Siomi, M.C. (2007). A Slicer-Mediated Mechanism for Repeat-
953  Associated siRNA 5' End Formation in Drosophila. Science *315*, 1587–1590.
954  https://doi.org/10.1126/science.1140494.

955 37. Horwich, M.D., Li, C., Matranga, C., Vagin, V., Farley, G., Wang, P., and Zamore,
956  P.D. (2007). The Drosophila RNA Methyltransferase, DmHen1, Modifies Germline
957  piRNAs and Single-Stranded siRNAs in RISC. Current Biology *17*, 1265–1272.
958  https://doi.org/10.1016/j.cub.2007.06.030.

959 38. Kuznetsov, D., Tegenfeldt, F., Manni, M., Seppey, M., Berkeley, M., Kriventseva,
960  E.V., and Zdobnov, E.M. (2023). OrthoDB v11: annotation of orthologs in the
961  widest sampling of organismal diversity. Nucleic Acids Research *51*, D445–D451.
962  https://doi.org/10.1093/nar/gkac998.

963 39. Vourekas, A., Zheng, K., Fu, Q., Maragkakis, M., Alexiou, P., Ma, J., Pillai, R.S.,
964  Mourelatos, Z., and Wang, P.J. (2015). The RNA helicase MOV10L1 binds piRNA
965  precursors to initiate piRNA processing. Genes Dev *29*, 617–629.
966  https://doi.org/10.1101/gad.254631.114.

967 40. Zheng, K., Xiol, J., Reuter, M., Eckardt, S., Leu, N.A., McLaughlin, K.J., Stark, A.,
968  Sachidanandam, R., Pillai, R.S., and Wang, P.J. (2010). Mouse MOV10L1
969  associates with Piwi proteins and is an essential component of the Piwi-interacting
970  RNA (piRNA) pathway. Proceedings of the National Academy of Sciences *107*,
971  11841–11846. https://doi.org/10.1073/pnas.1003953107.

972 41. Ikeda, S., Tanaka, K., Ohtani, R., Kanda, A., Sotomaru, Y., Kono, T., and Obata,
973  Y. (2022). Disruption of piRNA machinery by deletion of ASZ1/GASZ results in the
974  expression of aberrant chimeric transcripts in gonocytes. Journal of Reproduction
975  and Development *68*, 125–136. https://doi.org/10.1262/jrd.2021-146.

42. Ding, D., Liu, J., Dong, K., Midic, U., Hess, R.A., Xie, H., Demireva, E.Y., and
Chen, C. (2017). PNLDC1 is essential for piRNA 3′ end trimming and transposon
silencing during spermatogenesis in mice. Nat Commun *8*, 819.
https://doi.org/10.1038/s41467-017-00854-4.

43. Nishimura, T., Nagamori, I., Nakatani, T., Izumi, N., Tomari, Y., Kuramochi-
Miyagawa, S., and Nakano, T. (2018). PNLDC1, mouse pre-piRNA Trimmer, is
required for meiotic and post-meiotic male germ cell development. EMBO reports
19, e44957. https://doi.org/10.15252/embr.201744957.

44. Hayashi, R., Schnabl, J., Handler, D., Mohn, F., Ameres, S.L., and Brennecke, J.
(2016). Genetic and mechanistic diversity of piRNA 3' end formation. Nature *539*,
588–592. https://doi.org/10.1038/nature20162.

45. Honda, S., Kirino, Y., Maragkakis, M., Alexiou, P., Ohtaki, A., Murali, R.,
Mourelatos, Z., and Kirino, Y. (2013). Mitochondrial protein BmPAPI modulates the
length of mature piRNAs. RNA *19*, 1405–1418.
https://doi.org/10.1261/rna.040428.113.

46. Saxe, J.P., Chen, M., Zhao, H., and Lin, H. (2013). Tdrkh is essential for
spermatogenesis and participates in primary piRNA biogenesis in the germline.
The EMBO Journal *32*, 1869–1885. https://doi.org/10.1038/emboj.2013.121.

47. Feltzin, V.L., Khaladkar, M., Abe, M., Parisi, M., Hendriks, G.-J., Kim, J., and
Bonini, N.M. (2015). The exonuclease Nibbler regulates age-associated traits and
modulates piRNA length in Drosophila. Aging Cell *14*, 443–452.
https://doi.org/10.1111/acel.12323.

48. Joosten, J., Overheul, G.J., Van Rij, R.P., and Miesen, P. (2021). Endogenous
piRNA-guided slicing triggers responder and trailer piRNA production from viral
RNA in Aedes aegypti mosquitoes. Nucleic Acids Research *49*, 8886–8899.
https://doi.org/10.1093/nar/gkab640.

49. Wang, H., Ma, Z., Niu, K., Xiao, Y., Wu, X., Pan, C., Zhao, Y., Wang, K., Zhang,
Y., and Liu, N. (2016). Antagonistic roles of Nibbler and Hen1 in modulating piRNA
3′ ends in Drosophila. Development *143*, 530–539.
https://doi.org/10.1242/dev.128116.

50. Ishizu, H., Kinoshita, T., Hirakata, S., Komatsuzaki, C., and Siomi, M.C. (2019).
Distinct and Collaborative Functions of Yb and Armitage in Transposon-Targeting
piRNA Biogenesis. Cell Reports *27*, 1822-1835.e8.
https://doi.org/10.1016/j.celrep.2019.04.029.

51. Lee, Y.C.G., and Langley, C.H. (2012). Long-Term and Short-Term Evolutionary
Impacts of Transposable Elements on Drosophila. Genetics *192*, 1411–1432.

https://doi.org/10.1534/genetics.112.145714.

52. Obbard, D.J., Gordon, K.H.J., Buck, A.H., and Jiggins, F.M. (2008). The evolution

of RNAi as a defence against viruses and transposable elements. Philosophical

Transactions of the Royal Society B: Biological Sciences *364*, 99–115.

https://doi.org/10.1098/rstb.2008.0168.

53. Parhad, S.S., and Theurkauf, W.E. (2019). Rapid evolution and conserved function

of the piRNA pathway. Open Biology *9*, 180181.

https://doi.org/10.1098/rsob.180181.

54. Gamez, S., Srivastav, S., Akbari, O.S., and Lau, N.C. (2020). Diverse Defenses: A

Perspective Comparing Dipteran Piwi-piRNA Pathways. Cells *9*, 2180.

https://doi.org/10.3390/cells9102180.

55. Murota, Y., Ishizu, H., Nakagawa, S., Iwasaki, Y.W., Shibata, S., Kamatani, M.K.,

Saito, K., Okano, H., Siomi, H., and Siomi, M.C. (2014). Yb integrates piRNA

intermediates and processing factors into perinuclear bodies to enhance piRISC

assembly. Cell Rep *8*, 103–113. https://doi.org/10.1016/j.celrep.2014.05.043.

56. Okamura, K., Ishizuka, A., Siomi, H., and Siomi, M.C. (2004). Distinct roles for

Argonaute proteins in small RNA-directed RNA cleavage pathways. Genes Dev

18, 1655–1666. https://doi.org/10.1101/gad.1210204.

57. Van Rossum, G., and Drake, F.L. (2009). Python 3 Reference Manual

(CreateSpace).

58. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P.,

Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array

programming with NumPy. Nature *585*, 357–362. https://doi.org/10.1038/s41586-

020-2649-2.

59. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau,

D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0:

fundamental algorithms for scientific computing in Python. Nat Methods *17*, 261–

272. https://doi.org/10.1038/s41592-019-0686-2.

60. McKinney, W. (2010). Data Structures for Statistical Computing in Python.

Proceedings of the 9th Python in Science Conference, 56–61.

https://doi.org/10.25080/Majora-92bf1922-00a.

61. R Development Core Team (2023). R : A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

62. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

sequencing reads. EMBnet.journal *17*, 10–12. https://doi.org/10.14806/ej.17.1.200.

63. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,

Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq
aligner. Bioinformatics *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

64. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work
with high-throughput sequencing data. Bioinformatics *31*, 166–169.
https://doi.org/10.1093/bioinformatics/btu638.

65. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for
comparing genomic features. Bioinformatics *26*, 841–842.
https://doi.org/10.1093/bioinformatics/btq033.

66. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup
(2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*,
2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

67. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S.,
Heyne, S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web
server for deep-sequencing data analysis. Nucleic Acids Res *44*, W160-165.
https://doi.org/10.1093/nar/gkw257.

68. Wagih, O. (2017). ggseqlogo: a versatile R package for drawing sequence logos.
Bioinformatics *33*, 3645–3647. https://doi.org/10.1093/bioinformatics/btx469.

69. Wang, L., Wang, S., and Li, W. (2012). RSeQC: quality control of RNA-seq
experiments. Bioinformatics *28*, 2184–2185.
https://doi.org/10.1093/bioinformatics/bts356.

70. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-
source platform for biological-image analysis. Nat Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

71. Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I.,
Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for
researchers, educators, and developers. Protein Science *30*, 70–82.
https://doi.org/10.1002/pro.3943.

72. Croll, T.I. (2018). ISOLDE: a physically realistic environment for model building into
low-resolution electron-density maps. Acta Crystallogr D Struct Biol *74*, 519–530.
https://doi.org/10.1107/S2059798318002425.

73. Madeira, F., Pearce, M., Tivey, A.R.N., Basutkar, P., Lee, J., Edbali, O.,
Madhusoodanan, N., Kolesnikov, A., and Lopez, R. (2022). Search and sequence
analysis tools services from EMBL-EBI in 2022. Nucleic Acids Res *50*, W276–
W279. https://doi.org/10.1093/nar/gkac240.

74. Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates
to the phylogenetic tree display and annotation tool. Nucleic Acids Research,
gkae268. https://doi.org/10.1093/nar/gkae268.
75. Waskom, M.L. (2021). seaborn: statistical data visualization. Journal of Open
Source Software *6*, 3021. https://doi.org/10.21105/joss.03021.
76. Robert, X., and Gouet, P. (2014). Deciphering key features in protein structures
with the new ENDscript server. Nucleic Acids Research *42*, W320–W324.
https://doi.org/10.1093/nar/gku316.
77. Tamura, K., Stecher, G., and Kumar, S. (2021). MEGA11: Molecular Evolutionary
Genetics Analysis Version 11. Molecular Biology and Evolution *38*, 3022–3027.
https://doi.org/10.1093/molbev/msab120.
78. Nye, J., Buster, D.W., and Rogers, G.C. (2014). The use of cultured Drosophila
cells for studying the microtubule cytoskeleton. Methods Mol Biol *1136*, 81–101.
https://doi.org/10.1007/978-1-4939-0329-0_6.
79. Nishida, K.M., Iwasaki, Y.W., Murota, Y., Nagao, A., Mannen, T., Kato, Y., Siomi,
H., and Siomi, M.C. (2015). Respective functions of two distinct Siwi complexes
assembled during PIWI-interacting RNA biogenesis in Bombyx germ cells. Cell
Rep *10*, 193–203. https://doi.org/10.1016/j.celrep.2014.12.013.
80. Miyoshi, K., Tsukumo, H., Nagami, T., Siomi, H., and Siomi, M.C. (2005). Slicer
function of Drosophila Argonautes and its involvement in RISC formation. Genes
Dev *19*, 2837–2848. https://doi.org/10.1101/gad.1370605.
81. Saito, K., Ishizuka, A., Siomi, H., and Siomi, M.C. (2005). Processing of pre-
microRNAs by the Dicer-1-Loquacious complex in Drosophila cells. PLoS Biol *3*,
e235. https://doi.org/10.1371/journal.pbio.0030235.
82. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M.
(2022). ColabFold: making protein folding accessible to all. Nat Methods *19*, 679–
682. https://doi.org/10.1038/s41592-022-01488-1.
83. Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for
reconstructing phylogenetic trees. Mol Biol Evol *4*, 406–425.
https://doi.org/10.1093/oxfordjournals.molbev.a040454.
84. Lambkin, C.L., Sinclair, B.J., Pape, T., Courtney, G.W., Skevington, J.H., Meier,
R., Yeates, D.K., Blagoderov, V., and Wiegmann, B.M. (2013). The phylogenetic
relationships among infraorders and superfamilies of Diptera based on
morphological evidence. Systematic Entomology *38*, 164–179.
https://doi.org/10.1111/j.1365-3113.2012.00652.x.
85. Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen,

P.B., Ware, J., Flouri, T., Beutel, R.G., et al. (2014). Phylogenomics resolves the

timing and pattern of insect evolution. Science *346*, 763–767.

https://doi.org/10.1126/science.1257570.

1123

Supplemental figure legends

Figure S1. Coiled-coil prediction and specificity of anti-Daed antibodies

(**A**) Sequence-based CC prediction of Daed (upper) and Gasz (lower) by CoCoNat.[32]

Red line indicates the CC region predicted by AlphaFold2. (**B**) Western blotting shows

that anti-Daed monoclonal antibodies produced in this study are specific to Daed. Daed

KD: Daed knockdown by RNAi. control: EGFP KD. β-Tub: loading control.

1130

Figure S2. Functions and behaviors of Daed and Gasz in OSCs

(**A**) Northern blotting using probes that detect *idefix* piRNA and *tj* piRNA. miR-310:

loading control. KD: knockdown by RNAi. control: EGFP KD. pre-piRNA: piRNA

precursors. (**B**) Western blotting shows the levels of Flag-Gasz WT and Flag-Gasz Δα-

H2 where the same amounts of plasmids were used for transfection. β-Tub: loading

control. The statistics of western blotting signals (n=3; left) and the mRNA expression

levels (right). Signal intensities (left) or values (right) relative to F-Gasz WT were

normalized with those of β-Tub or *rp49*, respectively, and presented as mean values±SE.

***: p <0.005; n.s.: p >0.05 [*t*-test (unpaired, two-sided)]. (**C**) Western blotting shows

HSP60 (a mitochondrial fraction marker) and β-Tub (a cytoplasmic fraction marker) in

total lysates, and mitochondrial and cytoplasmic fractions in Figure 2G. (**D**) Summary of

Armi subcellular localization in OSCs under each condition.

1143

Figure S3. Effects of Armi mutants on piRNA biogenesis and Piwi-piRISC function

(**A**) Peptide sequence and domain structure of Armi. The ATP binding residues and

nucleotide binding residues are indicated in red and white, respectively. Asp1015,

Arg1026, and Arg1037 are highlighted in blue. (**B**) Quantification of piRNA levels (n=3)

(Figure 3H). Values relative to the control (EGFP KD) were normalized to Piwi level and

presented as mean values ± SE. ***: $p < 0.005$; n.s.: $p > 0.05$ [t-test (unpaired, two-sided)].

(**C**) This model summarizes the deleterious effects of Armi ∆N34 and KEE mutants on

Armi–AmTEC interaction, piRNA biogenesis, and Piwi-piRISC function in OSCs. Blue

arrowhead (WT): Zuc cleavage. Red arrowhead (KEE): obscured Zuc cleavage. White

arrowhead (KEE): Zuc cleavage corresponding to WT.

1154

Figure S4. Effects of Daed and Armi mutants on piRNA biogenesis and Piwi-piRISC

function

(**A**) Quantification of piRNA levels (n=3) (Figure 4A). Values relative to the control (EGFP

KD) were normalized to Piwi level and presented as mean values ± SE. ***: $p < 0.005$; *:

$p < 0.05$; [t-test (unpaired, two-sided)]. (**B**, **C**) Scatter plots showing the length distribution

of piRNAs upon subdivision by target transposons (**B**) and the abundance of reads

mapped antisense to transposons (**C**; Spearman's rho=0.98) in Daed WT and ∆SAM

libraries. (**D**) Left: Piwi-bound piRNAs obtained from cytoplasmic fraction (Cyto) and

nuclear extract fraction (Nuc). Immunoprecipitated Piwi was detected by western blotting.

Right: Western blotting showing β-Tub (a cytoplasmic marker) and Histon H3 (a nucleus

marker) in each fraction. (**E**) The expression levels of transposons (*gypsy*, *blood*, *gypsy5*,

quasimodo, *17.6*, and *mdg3*) in control OSCs (dark gray), Daed-lacking OSCs (Daed

KD; light gray), Flag-Daed WT rescued (WT; green) or Flag-Daed ∆SAM rescued cells

(∆SAM; red) (n=5). Values relative to the control (EGFP KD) were normalized to *rp49*

and presented as mean values ± SE **: $p < 0.01$; n.s.: $p > 0.05$ [Wilcoxon rank sum exact

test (two-sided)]. (**F**) Venn diagram showing the variety of positions of piRNA 5′ ends in

the WT and ΔSAM libraries. (**G**) Heatmaps showing 5′ end counts of piRNAs mapped

around the 10,000 most abundant piRNA 3′ ends (middle) and its binary histogram (top).

In sequence logos (bottom), piRNAs were aligned according to the 3′ end of the piRNAs

(black arrowhead) and its immediately downstream nucleotide is highlighted in green.

Left: Daed ΔSAM Unchanged. Right: Daed ΔSAM Elongated. (**H**) piRNA production

simulation. The U depletion in 29–34-nt piRNAs in WT and equivalent regions in ΔSAM

are highlighted in yellow. (**I**) Length distribution of piRNAs (18–40 nt) in Armi WT (green)

and KEE (red) libraries (n=3), presented as mean values ± SE. (**J**) Sequence logos

representing 11 nt from the 5′ end of piRNAs in the Armi WT and KEE libraries. (**K**)

Heatmaps showing the frequency of the 3′ end positions (red) corresponding to each 5′

end of piRNAs in Armi WT (upper) and KEE (lower) libraries. The 50,000 piRNA 5′ end

positions common to both libraries and most abundant in the WT library were used for

analysis.

1184

1185 **Figure S5. Sequence alignment of Daed and Armi UPF1-like domain in species**

1186 (**A**, **B**) Phyrogenic tree (A) and multiple sequence alignment (B) of Daed orthologs

1187 constructed using Clustal Omega and MEGA 11. (**C**) Multiple sequence alignment of *D.*

1188 *melanogaster* Armi and its orthologs. Red arrowhead: Asp1015, Arg1026, and Arg1037

1189 (D-R-R) of *D. melanogaster* Armi.

1190

1191 **Figure S6. Reproducibility of piRNA-seq**

1192 (**A,B**) Scatter plots showing reproducibility of piRNA-seq of Daed WT, ΔSAM mutant (**A**),

1193   Armi WT and KEE mutant libraries (**B**). Gray dots indicate the number of normalized

1194   piRNA reads mapped to coding genes.

A

AR

SAM   SAM

Daed

CC   α-H2

Gasz

TM

TM

B

Daed eCC

Gasz eCC   Gasz eCC

90°

CC   α-H2

Daed   Gasz

α-H2

Gasz

CC

Daed

Daed eCC

C

Daed eCC

Daed   SAM   CC TM   278 aa

140   150   160   170   180

ICNHPEAENYNNQILGNAGNHFMSLRASLAATNYKLQVSTPEDVVVGDKR

eCC

190   200   210   220   230

YASCFAQETLKSVNQITEEIAKDLAKIEANAQNARSQKKVQGNESHKKKW

CC

AR   Gasz eCC

Gasz   1 2 3 4 5 6   SAM   TM   461 aa

α-H2

340   350   360   370   380

DEMNNFYEILLITASHLQHLVIIQASLRFILKNQEHNKMGQPSEMQVAAM

eCC   α-**H2**

390   400   410

KSNLQGYRDVLNQITKTVKYLGSFSPSQNP

━━ α-helix

D

Gasz KD

input   Flag IP

(kDa)

70 —

55 —   Flag-Luc

45 —   *

                                                          Flag-Gasz

35 —

30 —   Daed

AR   SAM α-H2 TM

Gasz WT   1 2 3 4 5 6   SAM   461 aa

ΔAR1-3   4 5 6   SAM   360 aa

ΔAR4-6   1 2 3   SAM   363 aa

ΔSAM   1 2 3 4 5 6   384 aa

Δα-H2   1 2 3 4 5 6   SAM   432 aa

A

pre-piRNA

(nt)

idefix piRNA

tj piRNA

miR-310

C

HSP60

β-Tub

B

Flag-Gasz

WT
Δα-H2

β-Tub

D

	Armi	Daed	Gasz
Control	Yb body	+	+
[I]　Zuc KD	mitochondria	+	+
[II]　ZucKD + Daed KD	Yb body	–	–
[III] ZucKD + Daed KD, Flag-Daed	mitochondria	+	+
[IV] ZucKD + Daed KD, Flag-Gasz	Yb body	–	+
[V]　ZucKD + Gasz KD	Yb body	+	–
[VI] ZucKD + Gasz KD, Flag-Gasz	mitochondria	+	+

A

```
MFTYVSKFFT NPDRNREDIL ESLDRENSFL DQKLMEEKMD QQLKANPNEI   50
NGVLSNKIAE LTHGLSEMDV SKESSCTARK GVITSLDGDR GVIDKDVLFE  100
TKVAEDIILD LHVGCVVEYL TFTTGEAMRV VKVKSILEHS WEDTSQKEIE  150
KAVDNLKNEK PTFFNTETRS VLGLISQRLA SSIDVETEYG QLTVELDNIE  200
MNFIPTNGDR VRLECNIQLD DGFVDKQGEI LEVTKLFPTR IQEGEKCIVE  250
RVYVHMVVLG PETYILKTDL PTGTDLHLGD IVLADLIECQ YSKFTRRAIK  300
ITPLEKNFGA TKLTQQSSMG SSGSGKAVTV TGVNRFITAE LWQKESVSLK  350
LTNNLNRTLR LESITVCNDS ESQLSVVSPL ESKEISSGSE ITVTFEIHTQ  400
FLGEAIEKYV LNFDLLKVRR VFTVIVCKTK EEVAEAEKRM IAAEALMAPG  450
RNSQERSRFY ANQVWCNKVD VIPGQQIVTK RRFVALRLGC FEVPKELRQI  500
CLTSERRQEM IKAIEQHYSF LTEPLSIKTY MHRFRLLLHL EEIECFVNFR  550
NYDRDRAHFL RDGEFLTLQI ENLAERRPSL VIGDTLRVIN PWSDPDSQTT  600
KSYEGIIHKV LFDRILLKFH SSFQEKYNGE DYRLEFYFSR YSFRKQHHAI  650
SKIVGVMGED FLFPSKVTKR ENPQLDVYMK DDDMYLYDSK LEWYNQSLNS  700
IQKRAVFNIL RGEAENIPYV LFGPPGSGKT MTLIETLLQL VRNLPGARIL  750
VGTPSNSSAD LVTKRLIDSK ALLQGDFIRL VSYNQVEKDL IPPEIMSYCA  800
TSDVGAVGSC EDKMMVTESG LKLRCQAKFI GTHRITISTC TTLGNFLQLG  850
FPAGHFTHVL FDEAGQCTEP ETMVPIVMLT KKRSQVVLSG DPRQLQSIVT  900
SRIASKMGFS ISFLERLLER SPYRKDLQRF PESSGYNPLV LTKLLYNYRA  950
LPSIMSIYSR LFYDDELIPV LSEKDSRESR LLSKLRCVFE SEKDIPQAHG 1000
TFFYGIIGEN RQNNDSPSWF NPQEVREVFL MTIALYRANV TADQIGIITP 1050
YQKQVKMLRS MFIGTDVVMP KIGSVEEFQG QERDIILIST VRSSEEILRM 1100
DARFSLGFVR CSKRLNVAVS RARAMMIIFG NPHLLAVDEC WRQLILFCVK 1150
NNAYFGCDLP QMVINQKDEV PVCLETFVPS LNTTDDLN           1188
```

B

C

c

Figure S5 c. Protein sequence alignment of Armi Upf1-like region across species (Insecta: Hymenoptera, Coleoptera, Lepidoptera, Diptera). Positions D1015, R1026, R1037 in *D. melanogaster* are indicated.

Table S1. Primer sequences for plasmid construction

name	primer sequence	plasmid
Daed-clonig-insert_F	CTTGGTACCGAATTCATGGCAACGAAGACGCACAC	pAc-Flag-Daed
Daed-clonig-insert_R	GACTCGAGCGGCCGCTTAAGTCTGAACGGAGATCC	pAc-Flag-Daed
Daed-clonig-vector_F	TAAGCGGCCGCTCGAGTCTA	pAc-Flag-Daed
Daed-clonig-vector_R	CATGAATTCGGTACCAAGCT	pAc-Flag-Daed
Flag-Luc_insert_F	ATGGACTACAAAGACCATGACG	pAc-Flag-Luc
Flag-Luc_insert_R	CTTGTCATCGTCATCCTTGT	pAc-Flag-Luc
Flag-Luc_vector_F	GTCTTTGTAGTCCATGGTGGCTAGCCCGAT	pAc-Flag-Luc
Flag-Luc_vector_R	GGATGACGATGACAAGATGGAAGACGCCAAAAACATAAAGAAAG	pAc-Flag-Luc
Flag-Daed (RNAi resistant)_F	GATAAGCTCATTGCCGAGAACATT	pAc-Flag-Daed (RNAi resistant)
Flag-Daed (RNAi resistant)_R	AACGCATGTACTGGCGTAGTCC	pAc-Flag-Daed (RNAi resistant)
Flag-Daed ΔSAM_F	ATTCCCAACCAGAATCCCAGTTACG	pAc-Flag-Daed ΔSAM
Flag-Daed ΔSAM_R	TGTACTGGCGTAGTCCGTTTCCACG	pAc-Flag-Daed ΔSAM
Flag-Daed ΔCC_F	CACAAGAAGAAGAAGTGGAGCCTGG	pAc-Flag-Daed ΔCC
Flag-Daed ΔCC_R	GATCTGATTCACGCTTTTTAGAGTT	pAc-Flag-Daed ΔCC
GST-Daed_full_length_insert_F	GGGCCCCTGGGATCCATGGCAACGAAGACGCACAC	pGEX-GST-Daed
GST-Daed_full_length_insert_R	ACCCGGGAATTCCGGTTAAGTCTGAACGGAGATCC	pGEX-GST-Daed
GST-Daed_full_length_vector_F	CCGGAATTCCCGGGTCGACTCGAGC	pGEX-GST-Daed
GST-Daed_full_length_vector_R	GGATCCCAGGGGCCCCTGGAACAGA	pGEX-GST-Daed
GST-Daed ΔTM_F	TGGACCAAGTTCCGCAGCGCTCCCC	pGEX-GST-Daed ΔTM
GST-Daed ΔTM_R	CTTCTTCTTCTTGTGACTCTCGTTG	pGEX-GST-Daed ΔTM
GST-Daed ΔTM-His_F	CATCATCACTAACCGGAATTCCCGG	pGEX-GST-Daed ΔTM-His
GST-Daed ΔTM-His_R	ATGATGATGAGTCTGAACGGAGATC	pGEX-GST-Daed ΔTM-His
GST-Daed (N1-197)_insert_F	GGGCCCCTGGGATCCATGGCAACGAAGACGCA	pGEX-GST-Daed (N1-197)
GST-Daed (N1-197)_insert_R	ACCCGGGAATTCCGGTTATTCCTGGGCAAAGCAG	pGEX-GST-Daed (N1-197)
GST-Daed (N1-197)_vector_F	CCGGAATTCCCGGGT	pGEX-GST-Daed (N1-197)
GST-Daed (N1-197)_vector_R	GGATCCCAGGGGCC	pGEX-GST-Daed (N1-197)
Flag-Gasz ΔAR1-3_F	CACGGAGCCGTGATCAATGTTTCGG	pAc-Flag-Gasz ΔAR1-3
Flag-Gasz ΔAR1-3_R	CTTGGAGCGTTTCACCTCATCCGAG	pAc-Flag-Gasz ΔAR1-3
Flag-Gasz ΔAR4-6_F	CCAGCCTCCACTTACCTAGTGCCGA	pAc-Flag-Gasz ΔAR4-6
Flag-Gasz ΔAR4-6_R	CCGCAAGAGCAGGCCCACAATTCTC	pAc-Flag-Gasz ΔAR4-6
Flag-Gasz ΔSAM_F	TTCTATGAGATCCTACTGATCACCG	pAc-Flag-Gasz ΔSAM
Flag-Gasz ΔSAM_R	TTGAAAATATTCCGGGCAATCGCTC	pAc-Flag-Gasz ΔSAM
Flag-Gasz ΔCC_F	TTTAGCCCCTCTCAGAATCCCTTGT	pAc-Flag-Gasz ΔCC
Flag-Gasz ΔCC_R	CTCCGAAGGTTGCCCCATTTTATTG	pAc-Flag-Gasz ΔCC
Armi D1015K-Flag_F	ATAACAAATCaCCTTCCTGGTTCAA	pAc-Armi KEE-Flag
Armi D1015K-Flag_R	TTTGCCTGTTTTCGCCAATGATTCC	pAc-Armi KEE-Flag
Armi R1026E-Flag_F	AATCCCCAAGAGGTCGAGGAGGTGT	pAc-Armi KEE-Flag
Armi R1026E-Flag_R	GAACCAGGAAGGTGAATCGTTATTT	pAc-Armi KEE-Flag
Armi R1037E-Flag_F	ACGAGGCCAATGTGACTGCCGATCA	pAc-Armi KEE-Flag
Armi R1037E-Flag_R	ACAGGGCGATGGTCATTAGGAACAC	pAc-Armi KEE-Flag
MOV10L1-clonig-insert_F	ATCGGGCTAGCCACCATGCTGCGCCTTGCTGCCAA	pAc-MOV10L1-Flag
MOV10L1-clonig-insert_R	TTTGTAGTCCTCGAGCTTTTGGAGAGCCTGGAGTT	pAc-MOV10L1-Flag
MOV10L1-clonig-vector_F	CTCGAGGACTACAAAGACCATGACG	pAc-MOV10L1-Flag
MOV10L1-clonig-vector_R	CATGGTGGCTAGCCCGATCGGGGT	pAc-MOV10L1-Flag
ASZ1-clonig-insert_F	GGTACCGAATTCATGGCAGCGGGCACACTGAGGGG	pAc-Myc-ASZ1
ASZ1-clonig-insert_R	TCGAGCGGCCGCTTATTTTCTCTGCAAAGTTAGCT	pAc-Myc-ASZ1
ASZ1-clonig-vector_F	TAAGCGGCCGCTCGAGTCTAGAGGG	pAc-Myc-ASZ1
ASZ1-clonig-vector_R	CATGAATTCGGTACCAAGCTTGTTC	pAc-Myc-ASZ1

Table S2. siRNA sequences

name	sense	antisense
siEGFP	*GGCAAGCUGACCCUGAAGU*TT	*ACUUCAGGGUCAGCUUGCC*TT
siDaed	*CCUGUGUGGACAAACUUAUUG*	*AUAAGUUUGUCCACACAGGUG*
siGasz	*CGCUCGAAACCAGCUCCUU*TT	*AAGGAGCUGGUUUCGAGCG*TT
siArmi	*CAGCUUAAGGCGAAUCCAA*TT	*UUGGAUUCGCCUUAAGCUG*TT
siZuc	*GCAUUGCCGUCAGCACUGU*TT	*ACAGUGCUGACGGCAAUGC*TT
siPapi	*GAUUACGAGACAAAGACAG*TT	*CUGUCUUUGUCUCGUAAUC*TT
siNbr	*CCTAATGAGCAACCTGAAG*TT	*CTTCAGGTTGCTCATTAGG*TT

*: RNA is shown in itallics

Table S3. Probe sequences for northern blotting

name	probe sequence
idefix-piRNA	AAACTACTGGCAATCGTTTGGGAA
tj-piRNA	GGTAATGGGAATGCACTTCTCTTGAA
miR-310	AAAGGCCGGGAAGTGTGCAATA

Table S4. Primer sequences for qPCR

name	forward primer (5′ > 3′)	reverse primer (5′ > 3′)
rp49	CCGCTTCAAGGGACAGTATCTG	ATCTCGCCGCAGTAAACGC
Daed	GCAACGAAGACGCACACAAT	TCGGTGTGCGAGTGAAAGTT
Gasz	ACTTCCTGGGCTACAACACG	GGCCTCCAGTATGCCGTTAA
Flag-Gasz	ATGGACTACAAAGACCATGACGGTG	GAGCAGGCCCACAATTCTCTCCGCC
Nbr	GCCTTGGCAAGAAACTGAAC	ACCCCGCCTGAGAAAGTTAT
mdg1	AACAGAAACGCCAGCAACAGC	CGTTCCCATGTCCGTTGTGAT
tabor	ACAACAGAAGCCGCAAACTG	CAGCTATTGTGAGTGGGCGT
412	ATGCCACTGGTCTAAGCACC	ATAGTGGTGCAAGTCCCTGC
stalker2	CACACAGAAGAACAAGCGGC	TTAACCGCAACGCTTCCTCT
gypsy	CCAGGTCGGGCTGTTATAGG	GAACCGGTGTACTCAAGAGC
blood	AAGACTGCCCTCGACCATTG	ATTCGCTGTTGTCATGTGCG
quasimodo	CCCGACAACAGACAAGAGGC	AGCCCAAGTGTTGCACTGTT
gypsy5	GCCCAGAGACAACGACAGAA	CTGTCTTTGCTGTCCCGGAT
17.6	AATTGGCGCAGTCGATGTGA	TGCCCGTTATCGTGGTACTT
mdg3	ATGCCATACGATGCGCAAAC	CGCTCTTCCTTGTGGGACAT