UBS93 enables delineating intratumor subtype heterogeneity in breast cancer

Jing Li[1, #], Yuan Gao[2, #], Shouhui Guo[1], Shisong Zhang[3, 4, *], Weixing Yu[5, *], Ke Liu[1,*]

Affiliations:

1. Department of Medical Dataology, School of Public Health, Cheeloo College of Medicine, Shandong University, Jinan, China

2. Precision Medicine Laboratory for Chronic Non-communicable Diseases of Shandong Province, Institute of Precision Medicine, Jining Medical University, Jining, Shandong Province, China.

3. Department of Thoracic and Oncological Surgery, Children's Hospital Affiliated to Shandong University, Jinan, China

4. Department of Thoracic and Oncological Surgery, Jinan Children's Hospital, Jinan, China

5. Department of Biochemistry and Molecular Biology, School of Basic Medicine, Jining Medical University, Jining, China

# These authors contributed equally

* To whom correspondence should be addressed

zsli323@163.com

weixingy@mail.jnmc.edu.cn

keliu.iluke@email.sdu.edu.cn

Abstract

Breast cancer is a heterogeneous disease that comprises multiple molecular subtypes. Although several methods have been developed to subtype bulk tumor samples, their capacity to delineate intratumor heterogeneity at the single-cell level remains limited. In this study, we present UBS93, a novel classifier specifically designed for breast cancer subtyping in both bulk tumor samples and individual cancer cells. We rigorously validate UBS93 and demonstrate its superior performance compared to existing methods, particularly in the identification of Claudin-low samples. Application of UBS93 to single-cell RNA sequencing (scRNA-seq) data from triple-negative breast cancer patients uncovered a previously unrecognized "Basal-like/Claudin-low mixed" subtype, suggesting a Basal-like origin for Claudin-low cancer cells. Subsequent computational analyses and experimental validations reveal that downregulation of transcription factor *ELF3* induces a partial Claudin-low phenotype in Basal-like breast cancer cells. Collectively, our findings provide a powerful tool for elucidating intratumor subtype heterogeneity in breast cancer, with potential applicability to other cancer types.

Introduction

Breast cancer is a heterogeneous disease characterized by diverse genomic profiles, transcriptomic alterations, and clinical outcomes. The heterogeneity presents challenges in clinically relevant tumor classification[1].

Parker *et al.* utilized microarray technology to develop an intrinsic breast cancer classifier PAM50, which stratifies the disease into five subtypes: LuminalA, LuminalB, HER2-enriched, Basal-like, and Normal-like[2]. This classification system provides superior prognostic and predictive capabilities compared to traditional methods, improving upon pathological staging, histological grading, and standard clinical biomarkers[2]. In addition to PAM50, Herschkowitz *et al.* identified another subtype known as Claudin-low which is characterized by low expression levels of cell adhesion molecules (such as *CLDN3*, *CLDN4*, *CLDN7*, and *CDH1*) and marked by high stemness and mesenchymal traits[3, 4, 5, 6]. Fougner *et al.* conducted integrative multi-omics analyses on breast cancer samples from multiple cohorts and further confirmed Claudin-low as a distinct subtype[6].

The use of single-cell omics technology has profoundly enhanced our understanding of the complexity of breast cancer. Several recent studies have uncovered previously overlooked intratumor subtype heterogeneity, challenging the traditional classification strategy[7, 8, 9, 10, 11,12, 13, 14, 15, 16]. For instance, Wu *et al*. computationally identified Basal-like cancer cells within ER+ breast cancer patients and further confirmed the finding experimentally[16]. Similarly, the emergence of such "mixed subtype" phenomenon has been observed in mouse models through the activation of endogenous mutant *KRAS* or

exogenous *KRAS* expression[5]. These discoveries underscore the limitations of

traditional bulk omics-based subtyping and highlight the importance of achieving

single-cell resolution for more precise classification.

Despite advancements in computational methods for breast cancer subtyping, two

significant challenges remain. First, most of the existing methods are designed for

bulk omics data, which restricts their utility in single-cell analyses. Second, these

approaches often focus on PAM50 subtypes while overlooking the Claudin-low

subtype, a critical oversight that can lead to misclassification. Addressing these issues

is essential for refining breast cancer subtyping and improving diagnostic and

therapeutic strategies.

In this study, we developed a novel gene panel specifically tailored for breast cancer

subtyping and designed a classifier based on this panel. We rigorously evaluated its

performance using both bulk and single-cell RNA sequencing (scRNA-seq) data in

multiple species. Notably, we identified the presence of Claudin-low cancer cells

within Basal-like breast cancer cell lines and patient samples (a phenomenon we

define as the "Basal-like/Claudin-low mixed" subtype), demonstrating the utility of

our method in delignating intratumor subtype heterogeneity. Further computational

analysis and experimental assays revealed that *ELF3* knock-down induced a partial

Claudin-low phenotype in Basal-like breast cancer, suggesting a potential Basal-

like/Claudin-low trans-differentiation. Our work provides a powerful tool for

delineating intratumor subtype heterogeneity in breast cancer and should be

applicable to other cancers.

Results

Development of a uniform gene panel for breast cancer subtyping

Solid tumor tissue is a heterogeneous mixture of various cell types[16]. When quantifying the transcriptome of a tumor sample with bulk RNA sequencing technology, the expression of a gene could be considered as the combination of its expression in cancer cells and non-cancer cells. Given a gene g, suppose its average expression in cancer cells and non-cancer cells are C_g and M_g, respectively. If the tumor purity is α, then the expression value of g can be modeled as $\alpha * C_g + (1-\alpha) * M_g$. Given several random genes, their rankings (according to the measured expression in a bulk tumor sample) should significantly differ from those derived from pure cancer cells due to the contribution of microenvironment (**Figure 1a, left panel**); however, if these genes exhibit high cancer cell specificity (i.e., the value M for each gene is approximately zero), then the two ranking lists should be highly correlated (**Figure 1a, right panel**). Therefore, the rankings of cancer cell-specific genes might be informative for molecular subtyping.

To identify these genes, we analyzed the scRNA-seq dataset from Wu *et al.*'s study and calculated a value, termed the "C-M ratio", for each gene (see Materials and Methods). Among the top 100 genes with the highest C-M ratios, seven lacked ENSEMBL IDs and were subsequently removed, resulting in a uniform gene panel for breast cancer subtyping (UBS93) (**Supplementary Table 1**). As expected, the expression of UBS93 genes was significantly higher in cancer cells when compared to non-cancer cells (**Figure 1b**).

The Cancer Cell Line Encyclopedia (CCLE) project classifies breast cancer into five

subtypes: basal_A, basal_B, luminal, HER2-amp, and luminal-HER2-amp[17]. We

applied t-distributed Stochastic Neighbor Embedding (t-SNE) to the CCLE breast

cancer bulk RNA-seq data and observed that cell lines clustered according to their

subtypes, thereby validating the CCLE subtyping criteria (**Figure 1c**). As a result, we

primarily used CCLE subtypes in our study. However, to align with existing studies,

we made the following adjustments to subtype names. First, we renamed basal_A and

basal_B to Basal-like and Claudin-low, respectively. Second, we excluded the

luminal-HER2-amp subtype (see Materials and Methods). Ultimately, we classified

breast cancer into four molecular subtypes: Basal-like, Claudin-low, Luminal, and

HER2-amp.

We repeated the t-SNE analysis on the CCLE breast cancer bulk RNA-seq data with

only UBS93 genes (**Supplementary Figure 1**) and obtained results similar to

previous t-SNE analysis, confirming the effectiveness of the UBS93 panel in breast

cancer subtyping. Building on this, we developed our subtype classifier **(Figure 1d)**.

We first constructed a 93 $\times$ 3 subtype centroid matrix, where each column

represented the average expression vector of UBS93 genes across the CCLE cell lines

corresponding to Basal-like, Claudin-low, Luminal or HER2-amp, respectively. Given

the gene expression profile of a patient sample (or a single cancer cell), we computed

its Spearman rank correlation with each column of the subtype centroid matrix and

assigned it to the subtype with the highest correlation coefficient. If the sample is

subtyped as Luminal or HER2-amp, we further calculated an enrichment score of

HER2-amp signature genes and utilized this score to determine the sample's final

subtype (see Materials and Methods).

Evaluation of UBS93 classifier in human and mice

We first evaluated the performance of the UBS93 classifier using an assembled

dataset which contains the bulk RNA-seq profile of 47 breast cancer cell lines

extracted from the Gene Expression Omnibus (GEO) database[18, 19, 20]**(Supplementary**

Table 2). We found 93.6% of the samples were distributed on the diagonal of the

confusion matrix, suggesting the high overall accuracy of our method (**Figure 2a**).

Notably, our classifier also demonstrated optimal subtype-specific performance and

the F1 score for the four subtypes were 0.97 (Basal-like), 0.80 (Claudin-low), 1.00

(HER2-amp), and 0.92 (Luminal), respectively. In addition to cell lines, we also

evaluated our classifier on the breast cancer samples from The Cancer Genome Atlas

(TCGA) and derived ideal overall accuracy (81.9%) and subtype-specific

performance[21] (**Figure 2b**).

Next, we validated our classifier using an assembled scRNA-seq dataset of human

breast cancer cell lines (**Supplementary Table 3**)[22, 23]. For each cell line, we

computed the accuracy as the proportion of cells that were correctly subtyped. Our

classifier achieved an accuracy higher than 80% in 19 (63.3%) of the 30 cell lines and

the median accuracy was 85.7%. The subtype-specific median accuracy values were

99.0% (Basal-like), 98.3% (Claudin-low), 78.0% (HER2-amp), and 75.7% (Luminal)

(**Figure 2c**).

We assembled a dataset of 27 bulk RNA-seq samples to validate the performance of

our classifier in murine-derived datasets. In the dataset, 17 samples were from

Hollern's study[24] and experimentally confirmed as Claudin-low; the other 10 samples

were from Rädler's study[5] but lacked subtype labels. According to the results reported

in the literature, five of them were Basal-like and the other five were Claudin-low. We

performed hierarchical clustering based on the expression of five "Claudin-low

indicative" genes (*Cldn3, Cldn13, Cldn7, Vim, and Epcam*) to impute the missing

subtype labels (**Figure 2d**) and then applied our classifier to the 27 samples. We

found the subtype assigned by our classifier aligned with the true subtype in all

samples (**Supplementary Table 4**).

170
Comparison to existing methods

We systematically compared our classifier with genefu, a widely utilized R package

for performing PAM50 subtyping of breast cancer[25]. Notably, only six genes are

presented in both panels (**Supplementary Figure 2a**). In addition, UBS93 genes

exhibited significantly higher C-M-ratio values and cancer cell expression than

PAM50 genes (**Supplementary Figure 2b**). We compared the performance of the

two classifiers using the bulk RNA-seq data of TCGA breast cancer samples. The

genefu achieved slightly better overall accuracy (83.7%); however, the UBS93

classifier achieves better performance in Basal-like, Claudin-low, and HER2-amp

subtypes (**Figure 3a**). In murine datasets, the genefu package misclassified all Basal-

like samples as Claudin-low while our classifier accurately identified them as Basal-

like (**Supplementary Table 5**).

In addition to genefu, we also compared our classifier with SCSubtype (a method

developed for subtyping a single breast cancer cell) [16] on the assembled scRNA-seq

dataset that is used to evaluate our classifier. Since the Claudin-low subtype is not

included in SCSubtype, we therefore only made the comparisons on the three other

subtypes and found UBS93 classifier exhibited significantly higher accuracy than

SCSubtype in the Basal-like subtype (**Figure 3b**).

Discovery of "Basal-like/Claudin-low mixed" subtype

We observed that although our classifier generally performed well on scRNA-seq data,

the cell line HDQP1, which is believed to be a Basal-like cell line, appeared to be an

exception. Surprisingly, only 56.4% of HDQP1 single cells were subtyped as Basal-

like with the remainder subtyped as Claudin-low (**Figure 2c**). Interestingly, when

performing t-SNE visualization of the HDQP1 scRNA-seq data, the cells formed two

distinct clusters (**Figure 4a**). One cluster consisted of only Basal-like cells and 84.2%

of the other cluster were Claudin-low cells. We performed differential gene expression

analysis between UBS93-identified Claudin-low and Basal-like cells and found the

cells of Claudin-low group exhibited established characteristics of Claudin-low tumors[4,

5, 6, 26, 27]. First, there was significantly lower expression of cell-cell adhesion genes such

as *CLDN4*, *CLDN7*, *KRT19*, and *EPCAM* (**Figure 4b, Supplementary Figure 3,**

Supplementary Table 6). Second, there were higher levels of epithelial-to-

mesenchymal transition (EMT)-associated genes, as illustrated by up-regulation of *VIM*

and the enrichment of EMT signature genes in the up-regulated differentially expressed

(DE) genes (**Figure 4c**). Third, higher stemness was identified as indicated by an

elevated *CD44* (a marker of cancer stem cell) level and *CD44*/*CD24* ratio[4, 27, 28](**Figure**

4d). We next assembled a pseudo-bulk sample using the cells subtyped as Claudin-low

and computed its transcriptomic correlation with CCLE breast cancer cell lines[17]. The

cell line with the highest correlation coefficient was CAL120, a Claudin-low cell line

(**Figure 4e**). Moreover, four of the five cell lines with the highest correlation were

Claudin-low. In contrast, when we performed the same analysis using cells subtyped as

Basal-like, HDQP1 had the highest correlation coefficient (**Figure 4f**), which is

expected. Our computational analysis suggested that HDQP1 cell line was a mixture of

Basal-like and Claudin-low cells, therefore it is a cell line of "Basal-like/Claudin-low

mixed" subtype. To double confirm our finding, we replicated our analysis on the

HDQP1 scRNA-seq data from another dataset (GSE202771) and derived the same

conclusion (**Supplementary Figure 4, 5 and Supplementary Table 7**).

We next investigated whether the "Basal-like/Claudin-low mixed" subtype

phenomenon occurs in patients. We applied the UBS93 classifier to the scRNA-seq

data of 50 triple-negative breast cancer (TNBC) samples and ranked them according to

the median correlation coefficient (with the Claudin-low centroid) of UBS93-identified

Claudin-low cancer cells (**Figure 4g, Supplementary Table 8**). The metaplastic breast

cancer patient CID4513 (from dataset GSE176078) had the highest ranking, which may

be not surprising since previous studies have confirmed that metaplastic breast cancer

cells exhibit characteristics of Claudin-low subtype[29, 30]. Interestingly, the patient 0554

(from dataset GSE161529), who was reported as a Basal-like breast cancer patient in

the literature, had the second highest ranking[15]. Upon further exploration, we found

most (91.1%) of the cancer cells were subtyped as Basal-like while 7.9% were subtyped

as Claudin-low. Similar to HDQP1, the Claudin-low cells tended to form a distinct

cluster on the t-SNE plot and meanwhile exhibited the three established characteristics

of Claudin-low tumor (**Figure 5a-5d, Supplementary Figure 6, and Supplementary**

Table 9). In addition, the transcriptomic correlation analysis using pseudo-bulk

samples validated the results (**Figure 5e, 5f**). Therefore, we concluded that the Claudin-

low cancer cells identified in patient 0554 were highly convincible and this patient was

an instance of "Basal-like/Claudin-low mixed" subtype.

Together, these results highlight the existence of a Basal-like/Claudin-low mixed

subtype in both TNBC cell lines and patient samples, underscoring the complexity

within TNBC subtypes and the utility of UBS93 in delineating intratumor subtype

heterogeneity.

***ELF3* knock-down induces a partial Claudin-low phenotype in Basal-like cancer**

To date, the cellular origin of Claudin-low cancer cells remains unclear. While

Creighton *et al.* proposed that these cells originate from breast stem cells[31], evidence

from mouse models suggests they may instead arise from normal luminal epithelial

cells[5]. Pommier *et al.* performed large scale analysis on bulk cancer genomics data

and proposed the three potential origins of Claudin-low cancer cells: malignant

conversion from normal human mammary stem cells, transformation from luminal (or

basal-like) cancer cells[32]. However, no robust single cell-level evidence from patient

samples was provided in their study. The case of patient 0554, who exhibits the

"Basal-like/Claudin-low mixed" subtype, offers a unique opportunity to investigate

the cellular origin of Claudin-low cancer cells in greater depth. Hierarchical clustering

of various cell types based on their CNV (copy number variation) profiles revealed

that Claudin-low cancer cells share the highest similarity with Basal-like cancer cells

(**Figure 6a**), supporting the Basal-like origin of Claudin-low cancer cells in human

TNBC patients.

To investigate the genes which may play a pivotal role in the Basal-like/Claudin-low

transformation, we intersected the DE results performed in HDQP1 cell line and patient

0554, identifying 18 commonly upregulated and 37 commonly downregulated genes

(**Supplementary Table 10**). Among these, the transcription factor *ELF3* emerged as a

key player for two reasons **(Supplementary Figure 7). First,** overexpression of *ELF3*

in highly invasive Claudin-low cell lines (MDA-MB-231 and BT549) inhibits the EMT

process[33]; second, *ELF3* enhances the expression of *CLDN4* in ovarian cancer and non-

small cell lung cancer[34, 35]. Therefore, we questioned whether depletion of this gene in

Basal-like breast cancer would trigger the Basal-like/Claudin-low transformation.

Since HCC70 was previously identified as the cell line most closely resembling the

transcriptome of Basal-like breast cancer, we selected it for further studies[36]. First, we

performed scRNA-seq on HCC70 cells and applied the UBS93 classifier to the data.

As expected, 98.7% of the cells were classified as Basal-like, with no cells classified as

Claudin-low, confirming the high quality of the cell line. We then conducted scRNA-

seq on *ELF3* knockdown HCC70 cells to determine whether *ELF3* depletion could

convert Basal-like cells into Claudin-low cells, however UBS93 analysis indicated that

the majority of the cells (98.1%) remained Basal-like.

Despite failing to induce complete Basal-like/Claudin-low transformation, we explored

whether *ELF3* knockdown could partially induce Claudin-low characteristics in

HCC70 cells. To account for variable knockdown efficiency, we selected the 100 cells

with the lowest *ELF3* expression (sh-*ELF3* group) and the 100 cells with the highest

ELF3 expression (control group). Interestingly, the sh-*ELF3* group exhibited a

significantly lower correlation with the Basal-like centroid compared to the control

group (**Supplementary Figure 8a**). Differential gene expression analysis identified

210 upregulated and 79 downregulated genes (**Figure 6b, Supplementary Table 11**).

As expected, *CLDN4* was significantly down-regulated (p-value = 3.57e-14); in

addition, we also observed the down-regulation of the epithelial cell marker *KRT19*.

Interestingly, we found several of the down-regulated DE genes (such as *CLDN4*)

showed Claudin-low specific depletion (**Figure 6c**). We curated a set of such genes

(which we termed as the CLSD genes) and GSEA analysis revealed significant

enrichment of CLSD genes in the down-regulation part of DE gene list

(**Supplementary Figure 8b**).

We next examined how *ELF3* knock-down influenced the expression of *CLDN3*,

CLDN4, *CLDN7*, *EPCAM*, and *VIM* with RT-PCR (**Figure 6d**). Consistent with

scRNA-seq DE analysis, *CLDN4* was significantly down-regulated upon *ELF3* knock-

down (p-value = 0.002254). Interestingly, we also observed significant down-

regulation of *CLDN7* although it was not in the DE gene list (p-value = 0.018557); no

significant changes were detected for *CLDN3* and *EPCAM*. Unexpectedly, *VIM* was

significantly down-regulated (p-value = 0.003883), which is inconsistent with other

multiple studies reporting the suppression role of *ELF3* in EMT process. We conclude

there may be unknown mechanisms underlying the regulation of EMT by *ELF3* and

further investigation is needed. To double confirm our findings, we performed

replicated experiments and got same conclusion (**Supplementary Figure 9**). In

summary, our results suggest that *ELF3* knock-down induces a partial Claudin-low

phenotype in Basal-like cancer cell line, as evidenced by the down-regulation of CLSD

genes such as *CLDN4* and *CLDN7*.

Discussion

In this study, we introduced a novel classifier, UBS93, for the molecular subtyping of

breast cancer. We validated its performance using both bulk and single-cell RNA

sequencing data from human and mouse models. Notably, our classifier demonstrates

superior performance compared to classical methods, particularly in the identification

of Claudin-low subtypes. Unlike genefu, the most widely used method for breast

cancer subtyping, our classifier utilizes a single, uniform gene signature.

A key advantage of this approach is the simplification of the probability assignment

process. In UBS93, all correlation values (with subtype centroids) are calculated using

UBS93 genes, ensuring that the values are directly comparable. The probability of a

sample being classified as a specific subtype S_i is given by the formula:

314
$$Pr(Si) = cor(S_i) \, / \sum cor(S_j)$$

315 Where cor(Si) represents the correlation with the i-th centroid (values below zero are

316 assigned a value of zero). In contrast, when correlation values are computed across

317 different gene signatures (e.g., in genefu, the correlation with the Claudin-low

318 centroid is computed using non-PAM50 genes), additional adjustments are required to

319 make the correlations comparable.

320 Beyond computational analysis, our study lays the groundwork for a novel breast

321 cancer subtyping assay. We demonstrate that quantifying the expression of UBS93

322 genes is sufficient for molecular subtyping and can be achieved using cost-effective

323 technologies such as qRT-PCR. Future experiments will be conducted to validate our

324 concept.

325 Using UBS93, we identified human breast cancer cell lines and patients with the

326 "Basal-like/Claudin-low mixed" subtype. While Wu *et al*. previously reported "mixed

327 subtype" phenomenon in ER+ breast cancer patients[16], our finding was overlooked in

328 their study due to the exclusion of the Claudin-low subtype in the computational tool

329 SC Subtype. Our research complements and extends Wu *et al*.'s work, contributing to

330 a deeper understanding of intratumor heterogeneity especially in triple-negative breast

331 cancer.

332 Prior to our study, Pommier *et al*., through integrative analysis of large-scale cancer

333 genomics data, proposed three potential origins for Claudin-low cancer cells:

334 malignant conversion from normal human mammary stem cells, transformation from

335 luminal (or basal-like) cancer cells[32]. The discovery of the "Basal-like/Claudin-low

336 mixed" subtype, alongside our experimental findings, provides compelling evidence

supporting the Basal-like origin of Claudin-low cancer cells. In the future, we plan to

utilize UBS93 to obtain evidence for the other two hypotheses.

Our research also raises several interesting scientific questions for further

investigation. First, besides *ELF3* knock-down, which additional genetic perturbations

are needed to fully covert Basal-like cancer cells to Claudin-low cancer cells? High

EMT level is one of the three hallmark characteristics of Claudin-low cancer cells;

however, we did not detect significant up-regulation of mesenchymal marker genes

(such as *VIM* and *FN1*) upon *ELF3* knock-down. We noticed that *ZEB1,* a key EMT

regulator*,* was consistently up-regulated in the DE analysis between Claudin-low and

Basal-like cancer cells. Therefore, we intend to explore whether double genetic

perturbations (*ELF3*-knockdown and *ZEB1*-overexpression) would induce complete

Claudin-low phenotype.

Second, is *KRAS* mutation necessary for generation of Claudin-low cancer cells?

Rädler *et al.* reported that in mouse models activation of endogenous *KRAS* mutation

(or expression of exogenous *KRAS*) drove normal luminal epithelial cells to Claudin-

low and Basal-like cancer cells[5]; however, these findings have not been thoroughly

validated in human. In fact, *KRAS* mutation is rare in human breast cancer. According

to cBioPortal, we only identified 6 TCGA breast cancer patients carrying *KRAS*

mutation; in addition, the cell line HDQP1 also does not carry *KRAS* mutation.

Therefore, there might be *KRAS*-mutation-independent mechanisms which may

promote the generation of Claudin-low cancer cells.

Third, how does the presence of "Basal-like/Claudin-low mixed" subtype influence

the clinical outcomes? Distant metastasis remains the leading cause of cancer-related

deaths[37], and Claudin-low breast cancer has been shown to be highly aggressive

compared to other subtypes[27]. Therefore, it is reasonable to hypothesize that the

"Basal-like/Claudin-low mixed" subtype is an unfavorable prognostic marker in

TNBC. We plan to integrate scRNA-seq data with survival information to explore this

hypothesis further.

Last but not least, Fougner *et al.* reported that Claudin-low breast cancer exhibited high

lymphocyte infiltration and increased expression of PD-L1[6]. Therefore, we wonder

whether the presence of "Basal-like/Claudin-low mixed" subtype re-shapes the tumor

microenvironment. Briefly, we plan to apply the UBS93 classifier to spatial

transcriptomics data from patients with this "Basal-like/Claudin-low mixed" subtype,

identifying areas of Claudin-low cancer cells and comparing their surrounding

microenvironments to those of Basal-like cancer cells. This analysis could provide

valuable insights into potential immunotherapy opportunities.

In conclusion, our work presents a powerful tool for delineating intratumor subtype

heterogeneity in breast cancer. As high-quality tumor scRNA-seq data become

increasingly available, we anticipate that our strategy will be applicable to other

cancers and will help uncover novel biological mechanisms underlying tumor

progression.

Materials and Methods

Development of UBS93 gene panel

We obtained a breast cancer scRNA-seq dataset consisting of 8 samples from Wu *et al.*'s research[16]. Given a gene g, its C-M-ratio value in sample S is calculated with the following formula:

$$C\text{-}M\text{-ratio} = (C_g + 0.1) / (M_g + 0.1)$$

Here, C_g and M_g represent the average expression of G in cancer cells and non-cancer cells (of sample S), respectively.

Next, we divided all the samples into two groups: the TNBC group (2 samples) and the non-TNBC group (6 samples). We computed the group-wise C-M-ratio of gene g by taking the mean C-M-ratio within each group. Consequently, we obtained two group-wise C-M-ratio values for each gene g.

Finally, we ranked all genes according to their average group-wise C-M-ratio. We performed a quantile regression analysis between average group-wise C-M-ratio (Y) and rank (X) and noticed that there appears to be a change point of the slope. We picked out the top 400 genes to re-perform the analysis and found 300 appeared to be a reasonable estimate of the change point. Therefore, we finally kept the top 100 genes (**Supplementary Figure 10**). After excluding 7 genes that are lacked ENSEMBL gene IDs, we established the UBS93 gene panel.

Determination of breast cancer subtypes

The CCLE categorized breast cancer into five subtypes: basal_A, basal_B, Luminal, HER2-amp, and luminal-HER2-amp. In the t-SNE plot (**Figure 1c**), basal_A and basal_B form their own distinct cluster while the other three subtypes form the third cluster. The basal_A cluster is significantly enriched of cell lines whose PAM50

subtype is Basal-like; therefore, it is reasonable to rename it as Basal-like; similarly,

basal_B was renamed as Claudin-low. The luminal-HER2-amp subtype is not used in

our classifier since the two cell lines of this subtype do not appear to form a distinct

cluster.

Differentiating Luminal and HER2-amp subtypes

We conducted differential gene expression analysis between CCLE HER2-amp and

Luminal cell lines and identified 860 significantly up-regulated genes. We picked out

the 20 genes which are present in the UBS93 panel and derived the HER2-amp

signature.

In our classifier, ssGSEA analysis (UBS93 genes as background, and HER2-amp

signature as a geneset) is utilized to differentiate Luminal and HER2-amp subtypes.

We validated such strategy on a pooled bulk RNA-seq dataset (GSE212143 and

GSE48213) and obtained an AUC (area under the curve) value of 0.964

(**Supplementary Figure 11**). In the final classifier, we determined the cutoff value to

be 1.662 based on Youden's index calculated in the validation dataset.

Breast cancer subtyping with genefu

In genefu package, the gene sets used for Claudin-low and PAM50 subtyping are

different. When analyzing a breast cancer sample, we first employ the function

claudinLow to determine whether its subtype is Claudin-low; if not, we subsequently

conduct PAM50 subtyping. In PAM50 subtyping, we removed the Normal-like

subtype and merged LuminalA and LuminalB as Luminal.

Gene set enrichment analysis

The EMT signature was downloaded from MSigDB[38]. To derive the Claudin-low-

specific depletion (CLSD) gene set, z-score normalization was applied to gene

expression values (across all CCLE breast cancer cell lines) and the median

expression value for each gene was then calculated specifically for Claudin-low cell

lines. Genes with median expression values below -1 were selected. In total, resulting

in a total of 960 CLSD genes. The clusterProfiler R package was used to compute

enrichment score and p-value[39].

Processing of bulk RNA-seq data

We mapped raw bulk RNA-seq reads to transcriptome index with STAR[40] and then

utilized RSEM[41] to quantify each gene's TPM (Transcripts Per Million) value. The

genome versions used are hg38 and mm10, respectively. The genome annotation files

were downloaded from GENCODE[42].

The t-SNE analysis

Given two single cells C1 and C2, their distance is defined as 1- cor(C1, C2) where

cor(C1, C2) is the Spearman rank correlation value across used genes. After the

computation of the distance matrix, Rtsne function is used to generate the final

embedding.

Cell lines and cell culture conditions

HCC70 cells and HEK293T cells were obtained from Pricella Biotechnology Co., Ltd

and American Type Culture Collection. HCC70 cells maintained in RPMI 1640

(catalog no.22400089, Gibco). HEK293T cells maintained in DMEM (catalog no.

SH30285.01, Cytiva Life Sciences). All medium supplemented with 10% FBS (catalog

no. 164210-50, Pricella Biotechnology).

Lentivirus packaging and infection

HEK293T cells (2×10^6) were seeded in a 100-mm cell culture dish and transfected

with plasmids using PolyJet in vitro DNA transfection reagent PEI (C0537-20mg,

Beyotime Biotechnology). The transfection experiments were operated according to the

manufacturer's instructions. For knock-down target proteins, we co-transfected the

cells with the plasmids of pMDL, pVSVG, pREV and pLKO.1 (sh-*ELF3*, scrambled)

to generate lentivirus. For lentivirus infection, 500 μl of virus-containing medium was

added to the 5×10^4 cells with 1% polybrene.

Real-time PCR

Total RNA was extracted from cells using an RNAsimple Total RNA Kit

(TIANGEN, DP419). Complementary DNA was prepared using the High-Capacity

cDNA Reverse Transcription Kit purchased from Vazyme Biotech (catalog no. R333–

01). Real-time PCR analysis was performed using 2 × Universal SYBR Green Fast

qPCR Mix (catalog no. RK21203, Abclonal Technology) with the following

conditions: 3 min at 95 °C, and then 40 cycles at 95 °C for 5 s, 60 °C for 25 s, using a

Real-Time System (Roche LightCycler® 480). Data were normalized according to

expression of a control gene (ACTB) in each experiment. The data are presented as

the mean ± s.d. from three independent experiments. The p-values were computed

with unpaired t-test in GraphPad Prism 10 software.

Single-cell RNA-sequencing of HCC70

The single-cell RNA-sequencing of HCC70 cell line was performed by a commercial

company Novogene. Briefly, RNA from the barcoded cells was subsequently reverse-

transcribed and sequencing libraries were constructed with reagents from a Chromium

Single Cell 3' v2 reagent kit (10X Genomics) according to the manufacturer's

instructions. Sequencing was performed with Illumina sequencing machines

according to the manufacturer's instructions. After filtering low-quality read with

fastp[43], raw reads were demultiplexed and mapped to the human reference genome by

cellranger 7.2.0[44].

Software tools and statistical methods

All the analysis was conducted in R. Seurat[45, 46, 47, 48] was used for processing scRNA-

seq data. In DE analysis, genes with adjusted p-value < 0.05 and abs(log2FC) > 0.5)

were considered as differentially expressed genes. The ggplot2[49] package was used

for data visualization, pheatmap and ComplexHeatmap packages were used to

generate heat maps[50, 51], GSVA was used for ssGSEA analysis[52, 53]. If not specified,

the two-sided Wilcoxon rank-sum test was used to compute the p-value in hypothesis

testing.

Data Availability

The authors declare that all data supporting the findings of this study are available

within the article and its supplementary information files or from the corresponding

author upon reasonable request.

Code Availability

The code is available at github (https://github.com/bioklab/UBS93).

Acknowledgments

We thank Dr. Xiaohui Lin for helping us revise the manuscript. The research is

supported by National Natural Science Foundation of China (Fund 32370715),

Science Fund for Distinguished Oversea Young Scholars of Shandong Province

(2023HWYQ-015), Taishan Young Scholar Program of Shandong Province

(tsqn202312020), and Cheeloo Young Scholar Program of Shandong University. The

content is solely the responsibility of the authors and does not necessarily represent

the official views of sponsors.

Authors' contribution

J.L. and K.L. conceived the study. J.L. performed the majority of computational

analysis, Y.G. and W.X.Y. performed all the experimental assays, and K.L.

supervised the study. All authors contributed to writing, reviewing, and editing the

manuscript and approved the manuscript.

Competing Interests

The researchers declare no competing interests.

Reference

1. Nolan E, Lindeman GJ, Visvader JE. Deciphering breast cancer: from biology to the clinic. *Cell* **186**, 1708-1728.

2. Parker JS*, et al.* Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol* **27**, 1160-1167.

530   3.   Herschkowitz JI, *et al.* Identification of conserved gene expression features between murine ma
531   mmary carcinoma models and human breast tumors. *Genome Biol* **8**, R76.
532

533   4.   Prat A, *et al.* Phenotypic and molecular characterization of the claudin-low intrinsic subtype of
534   breast cancer. *Breast Cancer Res* **12**, R68.
535

536   5.   Rädler PD, *et al.* Highly metastatic claudin-low mammary cancers can originate from lumin al
537   epithelial cells. *Nat Commun* **12**, 3742.
538

539   6.   Fougner C, Bergholtz H, Norum JH, Sørlie T. Re-definition of claudin-low as a breast cancer
540   phenotype. *Nat Commun* **11**, 1787.
541

542   7.   Jordan NV, *et al.* HER2 expression identifies dynamic functional states within circulatin g
543   breast cancer cells. *Nature* **537**, 102-106.
544

545   8.   Chung W, *et al.* Single-cell RNA-seq enables comprehensive tumour and immune cell profi ling
546   in primary breast cancer. *Nat Commun* **8**, 15081.
547

548   9.   Azizi E, *et al.* Single-Cell Map of Diverse Immune Phenotypes in the Breast Tumor Micro
549   environment. *Cell* **174**, 1293-1308.e1236.
550

551   10.   Karaayvaz M, *et al.* Unravelling subclonal heterogeneity and aggressive disease states in T NBC
552   through single-cell RNA-seq. *Nat Commun* **9**, 3588.
553

554   11.   Kim C, *et al.* Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated by
555   Single-Cell Sequencing. *Cell* **173**, 879-893.e813.
556

557   12.   Savas P, *et al.* Single-cell profiling of breast cancer T cells reveals a tissue-reside nt memory
558   subset associated with improved prognosis. *Nat Med* **24**, 986-993.
559

560   13.   Gulati GS, *et al.* Single-cell transcriptional diversity is a hallmark of developmental p otential.
561   *Science* **367**, 405-411.
562

563   14.   Bassez A, *et al.* A single-cell map of intratumoral changes during anti-PD1 treatment of patients
564   with breast cancer. *Nat Med* **27**, 820-832.
565

566   15.   Pal B, *et al.* A single-cell RNA expression atlas of normal, preneoplastic and tumori genic states
567   in the human breast. *EMBO J* **40**, e107333.
568

569   16.   Wu SZ, *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nat Genet* **53**,
570   1334-1347.
571

17.   Barretina J, *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of anti

cancer drug sensitivity. *Nature* **483**, 603-607.

18.   Greer YE*, et al.* Mitochondrial Matrix Protease ClpP Agonists Inhibit Cancer Stem Cell F
unction in Breast Cancer Cells by Disrupting Mitochondrial Homeostasis. *Cancer Res Commun*
2, 1144-1161.

19.   Daemen A*, et al.* Modeling precision treatment of breast cancer. *Genome Biol* **14**, R110.

20.   Marcotte R*, et al.* Functional Genomic Landscape of Human Breast Cancer Drivers, Vulnerabi
lities, and Resistance. *Cell* **164**, 293-309.

21.   Cancer Genome Atlas Research N*, et al.* The Cancer Genome Atlas Pan-Cancer analysis project.
Nat Genet **45**, 1113-1120.

22.   Gambardella G, Viscido G, Tumaini B, Isacchi A, Bosotti R, di Bernardo D. A single-cell
analysis of breast cancer cell lines to study tumour het erogeneity and drug response. *Nat*
Commun **13**, 1714.

23.   Jovanović B*, et al.* Heterogeneity and transcriptional drivers of triple-negative breast ca ncer.
Cell Rep **42**, 113564.

24.   Hollern DP*, et al.* B Cells and T Follicular Helper Cells Mediate Response to Checkpoint I
nhibitors in High Mutation Burden Mouse Models of Breast Cancer. *Cell* **179**, 1191-1206.e1121.

25.   Gendoo DMA*, et al.* Genefu: an R/Bioconductor package for computation of gene expression-
b ased signatures in breast cancer. *Bioinformatics* **32**, 1097-1099.

26.   Pan C*, et al.* Research progress of Claudin-low breast cancer. *Front Oncol* **13**, 1226118.

27.   Dias K*, et al.* Claudin-Low Breast Cancer; Clinical & Pathological Characteristics. *PLoS One*
12, e0168669.

28.   Li W, Ma H, Zhang J, Zhu L, Wang C, Yang Y. Unraveling the roles of CD44/CD24 and ALDH1
as cancer stem cell marker s in tumorigenesis and metastasis. *Scientific reports* **7**, 13856.

29.   Reddy TP, Rosato RR, Li X, Moulder S, Piwnica-Worms H, Chang JC. A comprehensive
overview of metaplastic breast cancer: clinical featur es and molecular aberrations. *Breast*
Cancer Res **22**, 121.

30.   Weigelt B*, et al.* Metaplastic breast carcinomas display genomic and transcriptomic heter
ogeneity [corrected]. *Mod Pathol* **28**, 340-351.

31.   Creighton CJ*, et al.* Residual breast cancers after conventional therapy display mesenchymal as
well as tumor-initiating features. *Proc Natl Acad Sci U S A* **106**, 13820-13825.

32. Pommier RM*, et al.* Comprehensive characterization of claudin-low breast tumors reflects t he
impact of the cell-of-origin on cancer evolution. *Nat Commun* **11**, 3431.
620

621 33. Park S, Kwon Y. The role of ELF3 in metastasis of triple negative breast cancer. *Annals of*
622   *Oncology* **28**, x34 (2017).
623

624 34. Kuang L, Li La. E74-like factor 3 suppresses microRNA-485-5p transcription to trigger growth
625   and metastasis of ovarian cancer cells with the involvement of CLDN4/Wnt/β-catenin axis.
626   *Saudi J Biol Sci* **28**, 4137-4146.
627

628 35. Zhang X, Wang X, Wen Y, Chen S, Zhou C, Wu F. Single-cell transcriptomics reveal metastatic
629   CLDN4+ cancer cells unde rlying the recurrence of malignant pleural effusion in patients with
630   a dvanced non-small-cell lung cancer. *Clin Transl Med* **14**, e1649.
631

632 36. Liu K*, et al.* Evaluating cell lines as models for metastatic breast cancer through i ntegrative
633   analysis of genomic data. *Nat Commun* **10**, 2138.
634

635 37. Park M, Kim D, Ko S, Kim A, Mo K, Yoon H. Breast Cancer Metastasis: Mechanisms and
636   Therapeutic Implications. *Int J Mol Sci* **23**, 6806.
637

638 38. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP. Molecular
639   signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-1740.
640

641 39. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological themes
642   among ge ne clusters. *OMICS* **16**, 284-287.
643

644 40. Dobin A*, et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21.
645

646 41. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or wit
647   hout a reference genome. *BMC Bioinformatics* **12**, 323.
648

649 42. Frankish A*, et al.* GENCODE: reference annotation for the human and mouse genomes in 2023.
650   *Nucleic Acids Res* **51**, D942-D949.
651

652 43. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
653   *Bioinformatics* **34**, i884-i890.
654

655 44. Zheng GXY*, et al.* Massively parallel digital transcriptional profiling of single cells. *Nat*
656   *Commun* **8**, 14049.
657

658 45. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene
659   expression data. *Nat Biotechnol* **33**, 495-502.
660

661   46.   Stuart T, *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e1821.
662

663   47.   Hao Y, *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e3529.
664

665   48.   Hao Y, *et al.* Dictionary learning for integrative, multimodal and scalable single-ce ll analysis.
666   *Nat Biotechnol* **42**, 293-304.
667

668   49.   The ggplot2 package.https://cran.r-project.org/web/packages/ggplot2/index.html.).
669

670   50.   The pheatmap package.https://github.com/raivokolde/pheatmap.).
671

672   51.   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
673   multidimensional genomic data. *Bioinformatics* **32**, 2847-2849.
674

675   52.   Subramanian A, *et al.* Gene set enrichment analysis: a knowledge-based approach for interpret
676   ing genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550.
677

678   53.   Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and
679   RNA-seq data. *BMC Bioinformatics* **14**, 7.
680