1 **Pandemic *Vibrio parahaemolyticus* wave succession by gene loss**
2

3 Chao Yang[1,#*], Hongling Qiu[2,#], Sarah L. Svensson[1#], Chengpei Ni[3,#], Song Gao[3], Zhizhou Jia[1],
4 Huiqi Wen[1], Li Xie[1], Wenxuan Xu[1], Yujiao Qin[1], Shuzhu Lin[1], Jiancheng Wang[1], Yiquan
5 Zhang[4], Yinghui Li[5], Min Jiang[5], Xiaolu Shi[5], Qinghua Hu[5], Zhemin Zhou[6], Yanjie Chao[1],
6 Ruifu Yang[7], Yujun Cui[7], Jaime Martinez-Urtaza[8,*], Hui Wang[2,*], Daniel Falush[1,*]
7

[1] The Center for Microbes, Development and Health, CAS Key Laboratory of Molecular

Virology and Immunology, Shanghai Institute of Immunity and Infection, Chinese Academy

of Sciences, Shanghai, China;

[2] School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai,

China;

[3] Wuxi Center for Disease Control and Prevention, Wuxi, China;

[4] Department of Clinical Laboratory, Nantong Third People's Hospital, Affiliated Nantong

Hospital 3 of Nantong University, Nantong, Jiangsu, China;

[5] Shenzhen Center for Disease Control and Prevention, Shenzhen, China;

[6] Key Laboratory of Alkene-Carbon Fibers-Based Technology & Application for Detection of

Major Infectious Diseases, Cancer Institute, Suzhou Medical College, Soochow University,

Suzhou, China;

[7] State Key Laboratory of Pathogen and Biosecurity, Academy of Military Medical Sciences,

Beijing, China;

[8] Department of Genetics and Microbiology, Facultat de Biociències, Universitat Autònoma

de Barcelona, Barcelona, Spain

# These authors contributed equally to the article

* Corresponding authors:

C.Y.: cyang@siii.cas.cn; J.M.-U.: Jaime.Martinez.Urtaza@uab.cat;

H.W.: huiwang@shsmu.edu.cn; D.F.: daniel.falush@siii.cas.cn

29
30

Abstract

While the importance of spontaneous mutation and gene acquisition in the emergence and adaptive evolution of microbial pathogens is well recognized, the role of gene loss remains underexplored. Here, by large-scale phylogenomic analysis of 8,684 global isolates, we reconstructed the origin and evolutionary dynamics of the pandemic clone (PC) of *Vibrio parahaemolyticus*, the leading cause of seafood-associated infections worldwide. As has also been observed for the pandemic lineage of *V. cholerae*, the emergence of the clone involved the acquisition of multiple marker genes and genomic islands within a short timeframe (~60 years), followed by divergence into distinct, successive sublineages mediating independent waves of cross-country transmission. In *V. parahaemolyticus,* a recent wave succession event within the last two decades has been driven by gene-loss in the putrescine utilization (Puu) pathway, which confers phenotypic advantages related to environmental adaptation (enhanced biofilms) and human transmission (increased cell adhesion and intestinal colonization, reduced virulence in an animal model and milder clinical symptoms in humans), aligning with the trade-off hypothesis of pathogen virulence evolution. We identify similar lineage-specific Puu-gene-loss across multiple bacterial genera, with effects on biofilm and adhesion replicated in *V. cholerae* and *Escherichia coli*, suggesting convergent evolution and universal phenotypic effects across bacteria. Our findings highlight the indispensable role of gene loss in the adaptive evolution of bacterial pathogens.

Introduction

Microbial pathogens have recurrently emerged from environments or animals, with a few leading to disease epidemics or pandemics that pose notable public health challenges[1,2]. Understanding the genetic basis of pathogen emergence and adaptive evolution is critical for disease control and prevention, especially for pandemic pathogens. *Vibrio*, Gram-negative, halophilic bacteria that are ubiquitous in marine and aquatic habitats, are ideal models for investigating pathogen emergence and evolution. Among >100 *Vibrio* species, approximately 12 cause human infections[3]. Two species, *V. cholerae* and *V. parahaemolyticus*, are associated with disease pandemics, with the ongoing seventh cholera pandemic attributed to El Tor (7PET) isolates. *V. cholerae* 7PET was inferred to originate from a pre-pandemic strain that subsequently underwent explosive diversification - gaining multiple genetic elements within a short timeframe (~15 years), including the El Tor type cholera toxin prophage and *Vibrio* seventh pandemic islands (VSP-I and VSP-II)[4,5]. During the seventh pandemic, three successive sub-lineages of 7PET were identified, corresponding to the three independent but overlapping waves of global transmission[5,6], which may be driven by antibiotic resistance[7].

V. parahaemolyticus is the leading cause of seafood associated infections worldwide, with an increase in incidence over the last two decades. Its geographical range has also been expanding, possibly due to global warming[3,8,9]. The emergence of pandemic clone (PC), characterized by serotype O3:K6, substantially changed the *V. parahaemolyticus* epidemiology[10–12]. The O3:K6 PC strain was first isolated in 1995 from a traveler returning to Japan from Indonesia, and in 1996, it caused a disease outbreak in Calcutta, India[10]. Subsequently, it rapidly spread worldwide and came to dominate infections in many regions, along with rapid antigenic divergence (49 serotypes reported)[10–12]. Molecular and

epidemiological surveillances has provided evidence that the PC may have originated from a
serotype O3:K6 non-pathogenic strain, after acquisition of multiple marker genes (e.g.
virulence regulatory gene sequence *toxRS/new*[13], phage f237 and its unique open reading
frame *orf8*[14]) and marker genomic islands (VPaI-1 to VPaI-7), along with several
virulence-associated loci such as those encoding Type 3 or 6 secretion systems (T3SS2 or
T6SS1)[15,16]. However, how these events happened, the driving force and mechanism
underlying these events remain unclear. Currently, there is a lack of comprehensive genomic
view on the origin and evolution of the PC.
83
Leveraging a large-scale, global genome dataset of *V. parahaemolyticus*, which includes
strains from six continents and 34 countries, we comprehensively reconstructed the
evolutionary history of the PC. Intriguingly, we identified several similarities between PCs of
V. parahaemolyticus and *V. cholerae*. However, distinct from *V. cholerae*, we demonstrated
that gene-loss in the putrescine utilization pathway has driven the recent global wave
succession event by conferring phenotypic advantages, and these phenotypic effects are likely
universal among bacteria.
91

Results

From pre-pandemic (pre-PC) to PC

We assembled a total of 8,684 high-quality (completeness >90% and contamination <5%)
genomes of global *V. parahaemolyticus* isolates, including 333 newly sequenced genomes
(**Supplementary Table 1**). To characterize population structure and identify PC isolates, we
constructed a maximum-likelihood (ML) phylogenetic tree based on core-genome
single-nucleotide polymorphisms (SNPs). Based on the ML tree and pairwise SNP-distance,
2,815 were assigned as PC isolates (mean SNP-distance: 437), 39 as pre-PC (close relatives of
PC, mean SNP-distance to PC: 15,463), with the remaining being non-PC isolates (mean
SNP-distance to PC: 55,554, **Fig. 1a**).
102
Pre-PC isolates were further split into four sub-groups, designated pre-PC1 to pre-PC4,
respectively (**Fig. 1b,c**). Pre-PC1 to pre-PC3 were exclusively identified in Asia, with
pre-PC1 and pre-PC2 being clinical isolates, while pre-PC3 were mainly environmental
strains. The closest relative of the PC, pre-PC4, was represented mainly (13/16) by
environmental isolates from Asia (Japan and China) and the US. The serotypes of most
pre-PC isolates (33/39) were predicted to be O3:K6, and sequence types (STs) of pre-PC4b (a
subset of pre-PC4) isolates were ST3 except one strain (**Fig. 1c** and **Supplementary Table 2**).
110
Among Pre-PC and PC isolates, we identified a total of 88,754 core-genome SNPs, with 94%
attributed to homologous recombination, suggesting that recombination was the primary
driver of divergence from pre-PC to PC. We further assessed the temporal signal using the ML
tree based on recombination-stripped SNPs and strain isolation dates. A significant signal
($p<0.01$) was detected among strains from pre-PC3b to PC, allowing inference of the dated
phylogeny (**Fig. 1b,2b**). The most recent common ancestor (MRCA) of the PC was estimated
at 1983 (95% confidence interval [CI]: 1979-1985), 12 years preceding its first discovery in
1995. The strains are expected to be highly homogeneous at the point of their origin, but early

PC strains from 1996 already showed some diversity (an average of 38 non-recombined
SNPs), consistent with the long time-interval between their origination and first discovery
(**Fig. 2b**).
122
We investigated the distribution of known PC marker genes/islands and inferred the
evolutionary history from pre-PC to PC (**Fig. 1b,c** and **Extended Data Fig. 1**). The PC likely
originated from a serotype O3:K6 pathogenic strain encoding T3SS1, T6SS2, phage f237 and
VPaI-3, and subsequently acquired T6SS1 and the *toxRS/new* sequence. Between 1920
(MRCA of pre-PC4b and PC) and 1983 (MRCA-PC), MRCA-PC gained multiple known
marker genes/islands distributed across distinct genomic loci, including *orf8* and six out of the
seven VPaIs, as well as the gene encoding the well-known virulence factor thermostable
direct hemolysin, *tdh.* Additionally, 86 point mutations (34 synonymous; 38 non-synonymous)
and 80 homologous recombination events (524 kb, 10% of the reference genome) were
inferred along this branch. Moreover, *trh* (*tdh*-related hemolysin), VPaI-1 and VPaI-2 were
independently gained or lost by pre-PC strains.
134
We further investigated the source of these marker genes/islands by searching against non-PC
genomes. All of them were identified in non-PC isolates, although at low frequency (**Fig. 1c**,
Extended Data Fig. 2), indicating that these elements are more likely the result of
intra-species transfer from diverse non-PC *V. parahaemolyticus* strains than of cross-species
transfer from other *Vibrio* or bacterial species. Moreover, the differing distribution patterns of
these marker genes/islands among pre-PC and PC strains suggest that they were unlikely to
have been acquired simultaneously in a single burst. Apart from marker genes/islands, six
PC-specific genes were identified, three of which were within the superintegron
(**Supplementary Table 3**).
144

Fine-scale population structure of PC
To reduce the effect of repeat sampling (strains from disease outbreaks) and exclude outlier
strains, we generated a representative subset of 942 PC isolates for further analysis (see
methods), spanning 20 countries across five continents isolated between 1996 and 2022
(**Extended Data Fig. 3**). Among the representative isolates, we identified 26,377
core-genome SNPs, with 66% attributed to homologous recombination. Two recombination
hotspots were identified, including the known O:K antigen region[17] and a novel hotspot
(VP1883-1891) encoding a type II toxin-antitoxin system and a cold shock protein. There
were 281 homoplastic non-recombined SNPs, of which 26 were convergently identified 6-18
times in 16 genes associated with quorum sensing, environmental response and virulence
(**Extended Data Fig. 4** and **Supplementary Table 4**).
156

We constructed an ML tree and dated phylogeny for PC representative strains (**Fig. 2a,b**). By
integration with hierarchical clustering (**Extended Data Fig. 5**), we split the PC into four
sub-lineages corresponding to independent waves of cross-country transmission (**Fig. 2a-d**),
which was also observed in the *V. cholerae* 7PET[5]. Following the nomenclature of *V. cholerae*,
we designated these sub-lineages as wave-1 to 4, respectively. The estimated MRCA times for
wave-1 to wave-3 were close (1984-1987), while wave-4 emerged more recently in 1998.

We identified 11 serotypes and three STs in representative strains (**Fig. 2b**). While O3:K6 was dominant (79%), serotype conversions were identified, particularly in wave-2 and wave-3. ST3 (96%) remained the predominant ST. Several marker genes/islands (VPaI-1, 4, 5 and *orf8*) were lost in 2%-8% of representative isolates, with the others being conserved (**Extended Data Fig. 6a**). Notably, the genomic island VPaI-7, which encodes *tdh* and T3SS2, showed high differentiation between different waves (**Extended Data Fig. 6b**). Additionally, we observed a conserved antimicrobial resistance (AMR) gene profile (**Extended Data Fig. 6a**). Apart from intrinsic ampicillin resistance-related genes, no recently acquired gene was identified in >1% of representative isolates.

Wave succession and global transmission

Remarkably, we identified a clear 'wave succession' pattern: wave-1 to wave-4 succeeded each other temporally as the major wave in China, where sampling is sufficient. Similar trends were also seen in other countries with ≥10 representative strains, including the US, UK, Canada and Bangladesh, suggesting a possible global pattern. In contrast, in two South American countries, Peru and Chile, wave-1 has remained dominant for over a decade (**Fig. 2c,d**).

Wave-4 frequency has been increasing since its emergence, ultimately becoming predominant from 2012 onward. Since its origin, the effective population size (N_e) of the PC increased exponentially until 1998 (**Fig. 2e**). Subsequently, N_e fluctuated and showed a decreasing trend after 2010. Population dynamics of waves was consistent with temporal dynamics, with the N_e peaking alternatively for different waves. Post-2010, the N_e of wave-4 exceeded others and remained stable, while N_e of other waves declined rapidly.

We inferred the geographical transmission history of the PC by phylogeographical analysis, focusing on cross-continent events. A total of 67 transmission events were identified, mostly transient and not leading to local colonization (**Extended Data Fig. 7**). Only six 'TC' events (Transmissions resulting in local Colonization that persisted for ≥3 years) were identified (**Fig. 2b** and **Extended Data Fig. 7**). Wave-1 in South America likely sourced from a single TC event from Asia to Peru in 1990 (TC1), much earlier than its first discovery in South America (1997). Subsequently, wave-1 spread from Peru to Chile and backward transmissions were also identified (TC2-4). Additionally, two TC events from Asia to North America were identified (TC5-6) in other waves.

Puu-gene loss is associated with wave-4 replacement

To investigate the genetic basis of the wave-4 replacement of other waves, we comprehensively identified wave-4-specific variations and examined their functions. A total of 40 variations were identified (**Fig. 3a** and **Extended Data Table 1**), including 29 SNPs (located in 26 genes and 3 intergenic regions), two small indels (located in 2 genes) and most notably, the absence of nine adjacent genes (VP1775-VP1786). Of the nine genes, five are annotated as part of the putrescine utilization (Puu, VP1775-1781) pathway, one encodes a Na^+/H^+ antiporter (VP1782), and the others encode hypothetical or DUF3291

domain-containing proteins (VP1783-1786).

The fixation of gene loss is expected to be adaptive and selection-driven in free-living species,

which are generally believed to have large N_e[18]. Moreover, *Vibrio* species, which are abundant

in marine environments, have been broadly estimated to exhibit large N_e[19]. Therefore, we

hypothesize that the fixation of Puu-pathway gene loss in *V. parahaemolyticus* confers

phenotypic advantages.

The enzymes encoded by Puu-pathway genes degrade the polyamine putrescine into

4-aminobutanoate (GABA) or spermidine (**Fig. 3b**). Specifically, the *puuC* gene is involved

in three sub-pathways in the conversion of putrescine to GABA, and its deletion is expected

to completely block this conversion, mimicking the effect of a complete deletion of Puu-genes

(**Fig. 3b**). Therefore, we generated a *puuC* deletion mutant in the wildtype strain

RIMD2210633 (wave-3, hereafter RIMD-W3) to explore the function and consequences of

loss of Puu-genes in *V. parahaemolyticus*.

Phenotypic effects of Puu-gene loss

Biofilm formation

We first tested the phenotypic effects of Puu-gene loss on biofilm formation, as the

Puu-pathway has been reported to be involved biofilm formation in *Pseudomonas*

aeruginosa[20]. As reported in *P. aeruginosa,* Δ*puuC* showed significantly ($p<0.01$) enhanced

biofilm formation under M9 minimal media after 48 hours of culture (**Fig. 3c**), while

complementation of the deletion strains *in trans* (C-Δ*puuC*) significantly decreased biofilm

formation.

We next performed transcriptomic sequencing to identify differentially expressed genes

(DEGs) that may underlie the increased biofilm of Δ*puuC*, as well as to explore other

Puu-gene-loss related phenotypes. In line with increased biofilm formation of Δ*puuC*, we

identified multiple DEGs between wildtype and Δ*puuC* associated with biofilm or

biofilm-related pathways, including c-di-GMP and quorum sensing (**Fig. 3d, Supplementary**

Table 5). We next quantified intracellular c-di-GMP levels in wildtype and Δ*puuC* during

growth by introducing dual-fluorescence reporter plasmids. Consistent with differential

expression of several c-di-GMP-related enzymes and higher biofilm formation, c-di-GMP

levels were significantly higher in Δ*puuC* compared to wildtype in stationary phase (**Fig. 3e**).

Cell adhesion and intestinal colonization

The DEGs analysis give clues on the effects of Puu-gene-loss on other phenotypes in addition

to biofiom. Specifically, the most significant category of DEGs (Δ*puuC vs.* RIMD-W3) were

genes encoding CupE-fimbriae and Flp-pili (**Fig. 3d**), which are generally related to cell

adhesion for pathogens[21]. Motivated by these observations, we performed Caco-2 cell

adhesion assays and found significantly enhanced adhesion for Δ*puuC* compared to wildtype

(**Fig. 3f**). We also tested whether increased cell adhesion might transform into enhanced

intestinal colonization. We constructed bioluminescent wildtype and Δ*puuC* strains and traced

their colonization of mice after oral gavage using *in vivo* imaging. At 5 hours post-infection,

mice infected with Δ*puuC* exhibited stronger luminescence compared to wildtype (**Fig. 3g**).
Bacteria were then rapidly cleared, with no detectable luminescence observed for both
wildtype and mutant strains by 10 hours post-infection.
254
Virulence
An unexpected observation was the downregulation of the key virulence gene, *tdh,* in Δ*puuC*
(2.6-fold, *p*-adj=0.02, **Fig. 3d**), suggesting that Puu-gene-loss might reduce virulence.
Accordingly, we compared the lethality of wildtype and Δ*puuC* mutant strains in mice
following intraperitoneal infection. Consistent with *tdh* downregulation, Δ*puuC* exhibited
reduced lethality (**Fig. 3h**). These findings together revealed the potential impact of
Puu-gene-loss on phenotypes related to human infection, including cell adhesion, intestinal
colonization and virulence.
263
Phenotypes of wave-4 strains
The phenotypic experiments above provide clues about the selective advantages of wave-4
strains. To link the phenotypes related to Puu-gene-loss observed in a Δ*puuC* wave-3 strain
with the behavior of natural wave-4 strains, we next compared 12 natural strains, including
six wave-1 to wave-3 (2 strains each) and six wave-4 strains, for the same phenotypes as
described above. Consistent with the above phenotypic differences between wildtype and
mutant strains established above, the wave-4 strains exhibited enhanced biofilms and cell
adhesion, compared with wave-1 to wave-3 strains (**Fig. 4a**). Moreover, we complemented a
wave-4 strain (VPCZ58) with the nine genes (VP1775-VP1786) that are absent in wave-4
strains and identified significantly reduced biofilm formation and adhesion compared to the
wildtype strain (**Fig. 4b**), providing further evidence that the Puu-genes are responsible for
the observed phenotypic differences.
276
277 Additionally, we analyzed the clinical symptom data of 1,966 PC-related patients reported in
278 our previous study[22], and found that wave-4 was associated with milder symptoms, including
279 significantly lower rates of fever, vomiting and abdominal pain (**Fig. 4c**), consistent with
280 reduced virulence observed in our mouse infection experiment (**Fig. 3h**).
281
We hypothesized that these phenotypic differences mediated by Puu-gene-loss might confer
competitive advantages of wave-4 strains in the environment. Therefore, we performed
competition experiments for natural strains in seawater with chitin as a nutrition source,
mimicking a possible natural environment. We mixed wave-4 strains with those from other
waves in equal (1:1) or different (1:5) ratios. By targeted PCR amplification of a gene with a
wave-4-specific SNP allele (VPA0427 A919G), we quantified the relative abundance of
wave-4 based on the frequency of the wave-4-specific allele. After 72 hours of competition,
we measured a significant increase in the frequency of wave-4-specific alleles for both mixing
ratios (**Fig. 4d**), demonstrating the competitive advantage of wave-4 under the tested
condition.
292
Puu-gene-loss in bacteria
The convergent loss of Puu-genes and/or its fixation in specific bacterial lineages would

provide evidence supporting selection as the driving force. We analyzed the prevalence of
Puu-genes among 108,526 genomes from 36 bacterial species, including *Vibrio*, common
pathogens, and probiotics. Puu-genes were generally not conserved among bacteria, with part
of the pathway lost in some species (**Fig. 5** and **Extended Data Table 2**). Notably, the
absence/pseudogenization of Puu-genes in a subset of strains within species was convergently
identified across multiple species (**Fig. 5b** and **Extended Data Fig. 8**). Furthermore, this
absence/pseudogenization was fixed or nearly fixed in a specific lineage for species such as
Escherichia coli, *Shigella boydii*, *S. sonnei* and *P. aeruginosa*, or convergently occurred in
two or more lineages among species such as *Campylobacter coli, Yersinia pestis,*
Bifidobacterium longum and *Lactobacillus helveticus*. Additionally, Puu-gene-loss was not
only identified among *V. parahaemolyticus* PC strains, but also identified in the *V. cholerae*
7PET (**Fig. 5b**).
307
To test if Puu-gene-loss has a similar phenotypic effect in other bacteria, we generated *puuC*
deletion strains in *V. cholerae* (E7946 background) and *E. coli* (MG1655 background) and
tested their biofilm formation and cell adhesion abilities. Consistent with the results in *V.*
parahaemolyticus, the Δ*puuC* mutant strains in both species exhibited significantly enhanced
biofilms and adhesion compared to their parental wildtype strains (**Fig. 5b**). Together, these
results indicate that Puu-gene-loss is under selection in some species, and that the associated
phenotypic consequences are potentially universal across bacterial species.
315
Discussion
Leveraging a large-scale genome dataset of global isolates*,* we present the most
comprehensive genomic view to date on the origin and evolution of the *V. parahaemolyticus*
PC. Our findings suggest that the PC likely originated from a serotype O3:K6 pathogenic
strain through gain of multiple genes/islands within a short timeframe (~60 years) from *V.*
parahaemolyticus, rather than from non-pathogenic strains or from other *Vibrio* species[11].
Notably, several characteristics previously considered to be specific to the PC[10–12], including
ST3, *toxRS*/new, and VPaI-1 to VPaI-3, were also present in pre-PC strains, while the *orf8*
gene remained unique to the PC, thus remaining as a reliable marker. Moreover, we described
the fine-scale population structure and spatiotemporal dynamics of the PC, and identified
'wave succession' with potential underlying molecular mechanisms. Together, our findings
substantially enhance the current understanding of how epidemic lineages of *V.*
parahaemolyticus emerge and evolve.
329
During the evolution of the *V. parahaemolyticus* PC, we observed intriguing parallels with the
V. cholerae 7PET strains and other pathogens. First, we observed gain of genes/islands,
including key virulence genes (*tdh* and *ctxAB*) and marker genomic islands (six VPaIs and
two VSPs) for the *V. parahaemolyticus* PC, as has been reported for the *V. cholerae* 7PET[4,5].
This phenomenon has also been observed during human colonization by other pathogens[23]
and in the evolution of metabolic pathways across prokaryotic lineages[24]. Frequent
human-mediated cross-regional transmission and adaptation to new environments are
considered potential drivers for the *V. cholerae* 7PET[4], which might also apply to the *V.*
parahaemolyticus PC. Second, for both PCs from pandemic *Vibrio* species, the inferred times

of origin (1983 and 1955[4]) and introduction to specific geographical regions (e.g. South
America: 1990 and 1989[25]) were much earlier than their first epidemiological observations
(1995 and 1961; South America: 1997 and 1991). Three possible explanations for this include
inadequate surveillance, variability in the molecular clock rate[26] with higher early mutation
rates[27,28], and the possibility of early cryptic disease epidemics, with specific events such as El
Niño or warming in coastal conditions triggering a sudden surge in cases and thereby drawing
epidemiological attention. While the third explanation does not apply to *V. cholerae* due to the
high risk and strict surveillance of cholera, all explanations are applicable to *V.*
parahaemolyticus. Third, divergence into successive sublineages mediating waves of
infections, identified in both pandemic *Vibrio* and other pathogens such as SARS-CoV-2[29].
While antimicrobial resistance likely drove the wave succession of *V. cholerae* 7PET[7], *V.*
parahaemolyticus exhibits low resistance rates to first-line antibiotics, indicating a different
driving factor.
352
The importance of biofilm formation in bacterial environmental adaptation and survival under
adverse conditions (e.g. antibiotic exposure) is well recognized[30]. Therefore, the enhanced
biofilms mediated by Puu-gene-loss may improve the environmental adaptability of wave-4
strains, as suggested by our competition experiment results. Moreover, the ancestral strains of
the PC, as well as most non-PC strains, carry the complete Puu-pathway genes, indicating that
the loss of Puu-genes occurred recently, possibly in response to environmental changes.
Putrescine serves multiple physiological functions in diverse organisms, including roles in cell
growth and proliferation[31], and has been particularly reported to promote acid stress tolerance[32,33].
The loss of Puu-genes might have been selected for by recent increases in ocean or coastal
acidification.
363
Human activities are pivotal in pathogen transmission. Increased cell adhesion and intestinal
colonization could directly facilitate human-mediated transmission, while reduced virulence,
associated with milder clinical symptoms, may lower the likelihood of seeking medical
attention, thereby promoting cryptic transmission. Notably, increased adhesion and
colonization are not contradictory to reduced virulence, as different infection routes in the
experiments (gavage *vs.* intraperitoneal injection) and their underlying mechanisms are
known to differ[34].
371
The effect of the Puu-pathway on biofilms is likely a broader phenomenon among bacteria
than previously recognized. We have demonstrated that Puu-gene-loss enhances biofilms in
three different species. Furthermore, Puu-gene-loss in *P. aeruginosa* also leads to enhanced
biofilms[20], and *E. coli* phylogroup B2, characterized by absence of *puuA-D* (**Fig. 5b**) exhibits
a stronger biofilm-formation ability than other phylogroups[35,36], providing further evidence
for a more universal effect.
378
For the first time, we have established links between Puu-gene-loss, cell adhesion and
intestinal colonization, which are also likely universal across bacteria. We demonstrated that
Puu-gene-loss enhances cell adhesion in three different species and promotes intestinal
colonization in *V. parahaemolyticus*. Furthermore, *E. coli* phylogroup B2 strains are often

associated with urinary tract infections[37] and show increased persistence in the intestines of
human infants[38], both of which is consistent with the hypothesis of increased adhesion and
colonization mediated by Puu-gene-loss. Interestingly, the probiotic *E. coli* strain Nissle 1917,
known for its strong capability of human gut colonization[39], belongs to phylogroup B2 and
lacks Puu-genes (**Fig. 5b**).
388

The convergent loss or pseudogenization of Puu-gene within species (e.g., *C. coli*) and across
different bacterial species, along with its lineage-specific fixation (e.g., in *E. coli*), indicate
that these events are more likely selection-driven in these species and may confer fitness
advantages. Puu-gene loss/pseudogenization was uniquely fixed in major pathogenic clonal
complexes (CC) of *S. boydii* (CC145) and *S. sonnei* (CC152)[40], and recently fixed in *V.*
parahaemolyticus wave-4, suggesting potential fitness advantages for human infection or
transmission. However, this appears to not always be the case, as Puu-gene
loss/pseudogenization has not occurred in major pathogenic lineages of *C. coli* and *P.*
aeruginosa. Additionally, Puu-gene loss/pseudogenization appears to be associated with
lineage divergence in *E. coli* and *C. coli*, and Puu-gene gain is potentially linked to human
adaptation in the probiotic *B. longum,* with Puu-genes predominantly identified in adult
human strains and absent in infant or non-human strains. These observations together indicate
that while Puu-gene loss/pseudogenization is likely selection-driven in some species, the
underlying driving forces vary across different species.
403

While the significance of spontaneous mutation and gene acquisition in pathogen evolution is
well recognized[41–43], the role of gene loss remains underexplored. Gene loss is pervasive
across all life kingdoms[44,45]. In bacteria, it is commonly observed in symbionts and
host-dependent pathogens, where is generally considered neutral and driven by genetic drift
associated with transmission bottleneck[46,47], while loss of several specific genes has been
experimentally validated to be associated with pathogen virulence[48–50]. Notably, gene loss
affecting polyamine metabolism has also been reported in the pathogen *Shigella*, where the
polyamine cadaverine was found to inhibit enterotoxin activity[48]. This aligns with our
findings in *V. parahaemolyticus*, in which elevated levels of a different polyamine, putrescine,
are associated with reduced virulence. Together, these findings suggest a potentially broader
role for polyamines (e.g., cadaverine and putrescine) in modulating bacterial pathogenesis.
415

Free-living bacteria with a large N_e serve as better models for investigating adaptive evolution
driven by gene loss, as such fixation is more likely to be adaptative and selection-driven.
Some free-living bacteria, such as the abundant open-ocean bacterioplankton
Prochlorococcus, exhibit extensive gene loss, which has been attributed to accelerated
evolution and fitness advantages conferred by genomic and metabolic "streamlining"[51,52].
Moreover, previous studies of gene loss-mediated bacterial adaptation, whether for symbionts,
host-dependent pathogens, or free-living species, generally come with the cost of transition
from generalists capable of surviving in diverse environments to specialists adapted to
specific ecological niches[49–52]. The exact niche of the *V. parahaemolyticus* PC remains unclear
but it is more likely a generalist due to widespread distribution in marine and aquatic habitats.
If we consider different geographical regions as the macro-niches, the wave-4 strains

characterized by Puu-gene-loss have spread globally, without clear evidence supporting its
transition from a generalist to specialist strategy. Therefore, our study presents a novel pattern
of gene loss-mediated adaptation in free-living species.

In summary, our study provides the most comprehensive genomic view on the origin and
evolution of the *V. parahaemolyticus* PC. We identified several interesting common features
between the evolution of pandemic clones in two *Vibrio* species, but the driving forces
underlying wave succession events appear distinct. While spontaneous mutation and gene
gain-mediated antibiotic resistance may drive the *V. cholerae* 7PET succession[7], we
demonstrated that Puu-gene loss, which confers phenotypic advantages related to
environmental adaptation and human transmission, has likely driven the recent *V.*
parahaemolyticus wave succession. An interesting finding is that *V. parahaemolyticus* wave
succession has come along with reduced virulence, which aligns with the trade-off hypothesis
of pathogen virulence evolution[53]. We also found that Puu-gene-loss is general and frequent
among bacteria, selection-driven in multiple species, with its phenotypic impacts likely
universal across species. Furthermore, Puu-gene-loss may play key roles in bacterial
evolution, including lineage divergence (e.g. *E. coli, C. coli* and *B. longum*), human
adaptation (e.g. *B. longum*) and the epidemic expansion of specific pathogenic lineages (e.g. *V.*
parahaemolyticus, S. boydii and *S. sonnei*). Taken together, our study underscores the
indispensable role of gene loss in the adaptative evolution of bacterial pathogens.

Methods
***V. parahaemolyticus* genome dataset**
A total of 8,684 high-quality genomes were included in this study, including 333 newly
sequenced and 8,351 publicly available genomes (**Supplementary Table 1**). Public genomes
(up to November 2022) were downloaded from National Center for Biotechnology
Information GenBank or Sequence Read Archive (SRA) database, with metadata extracted
from corresponding BioSample information or the literature. New genomes were obtained
from archived strains isolated from multiple countries, including China, Japan, Senegal, Peru
and Chile.

Whole genome sequencing of new strains was performed using Illumina NovaSeq or Miseq
platforms. New and public sequencing reads were trimmed using Trimmomatic v0.39[54] and
were *de novo* assembled using shovill v1.1.0 pipeline (https://github.com/tseemann/shovill).
The quality assessment of genomes was performed using CheckM v1.1.3[55]. Whole genome
average nucleotide identity (ANI) against the reference genome of *V. parahaemolyticus*
(RIMD2210633, accession numbers: NC_004603.1 and NC_004605.1) was calculated using
fastANI v1.1[56]. Only high-quality genomes (completeness >90% or contamination <5%) with
ANI >95% were included in further analysis.

Phylogenetic and phylogeographical analysis
Genome sequences were aligned against the reference genome using MUMmer v3.23[57] to
generate whole-genome alignments. Core-genome (regions present in >99% strains)
single-nucleotide polymorphisms (SNPs) were identified using SNP-sites v2.5.1[58]. SNPs

located in repetitive regions identified by TRF v4.07b[59] or BLASTN self-search were
excluded. Non-repetitive SNPs were used to construct maximum-likelihood (ML)
phylogenetic trees using FastTree v2.1.10[60] for all the isolates.

For both pre-PC and PC strains, recombination regions and associated length were detected
using Gubbins v3.1.3[61]. Homoplastic SNPs and nutations (non-recombined SNPs) along the
phylogenetic branch were identified using SNPPar v1.0[62] based on ancestral state
reconstruction of these variations. Non-recombined and non-homoplastic SNPs were used for
the construction of ML tree using RAxML-NG v1.0.1[63] with the GTR+Γ model. Phylogenetic
trees were visualized using ggtree[64]. Hierarchical clustering analysis were performed using
fastBAPS v1.0.6[65] to assign strains into subgroups. The transitions in the isolation countries
of strains in the phylogenetic tree are considered cross-country events. These were identified
manually, and all events are shown in detail in Extended Data Fig. 7. TC was defined as
Transmissions resulting in local Colonization (persisted for $\geq$3 years).

Temporal analysis and population size dynamics
Temporal signals among pre-PC and PC isolates were assessed based on the
recombination-stripped ML tree using TempEst v1.5.3[66], by calculating the correlation
between root-to-tip distances and strain isolation dates (years). A significant temporal signal
was detected among strains from pre-PC3b to PC (p<0.01, R=0.62), whereas inclusion of
pre-PC3a and earlier pre-PC strains eliminated the significant signal. A significant temporal
signal was also detected within PC strains alone (p<0.01, R=0.57). For strains showing a
significant temporal signal, we applied BactDating v1.1.0[67] to infer the dated phylogenetic
trees with the additive relaxed clock (ARC) model. The effective sample sizes (ESS) of the
estimated parameters were all above 200. Effective population size dynamics was inferred
using skygrowth v0.3.1[68] based on dated trees with default parameters.

Serotype and MLST typing
In silico O- and K-antigen genotyping was performed using the *V. parahaemolyticus* Kaptive
database[69]. Multi-locus sequence typing (MLST) was performed using mlst
(https://github.com/tseemann/mlst) to scan genome sequences against the PubMLST
databases.

Representative genome dataset of PC
We generated a representative subset of 942 PC isolates. First, to reduce the effect of repeat
sampling such as strains from disease outbreaks, we selected the genome with maximum N50
from a cluster of strains with $\leq$6 SNPs from same country and within a year, representing
putative outbreaks[22]. Second, we calculated root-to-tip distances of phylogenetic trees based
on non-recombined SNPs, and removed putative hypermutators with extreme root-to-tip
distances and branch lengths (**Extended Data Fig. 3**).

Core-pangenome
All PC representative genomes were re-annotated using Prokka v1.13[70], and the annotation
outputs (GFF3 files) were used in Panaroo v1.2.8[71] to identify the core-pangenome and gene

presence/absence.

Wave-4 specific variations

We analyzed three types of variations, including SNPs, gene presence/absence and unitigs.
The SNPs and gene presence/absence data were extracted from the preceding analysis. The
unitig   matrix   was   determined   using   unitig-caller   v1.2.1
(https://github.com/bacpop/unitig-caller). Unitig sequences were aligned against assembled
genomes using BLASTN to identify corresponding genomic positions and related genes.
Wave-4 specific variations were defined as alleles, genes or unitigs present/absent in >95% of
wave-4 strains and <5% of non-wave-4 strains.

***puuC* gene in the reference genome**

The *puuC* gene (VP_RS08540/VP1776-7) in the reference genome RIMD2210633 was
annotated as a pseudogene due to a frameshift caused by insertion of one nucleotide within a
homopolymeric region (three consecutive adenines). However, we performed both Sanger
sequencing and Illumina re-sequencing of this region in our RIMD2210633 strain. Both
sequencing methods confirmed absence of this nucleotide and an intact *puuC* ORF in our
strain, with no frameshift detected. Given the low sequencing depth of the original reference
genome (average 8.9×) and the elevated error rate associated with homopolymeric regions, we
hypothesize that the frameshift reported in the reference genome is likely caused by a
sequencing error.

Puu-genes in bacteria

We investigated the prevalence of Puu-genes in 36 bacterial species, including *Vibrio*,
common pathogens, and probiotics. We included a total of 108,526 genomes (62-24,205
genomes per species, **Supplementary Table 6**) downloaded from Genome Taxonomy
Database (GTDB, release207)[72]. Puu-related protein sequences in *E. coli*, *P. aeruginosa* and *V.*
parahaemolyticus were used as initial queries, which were searched against all genomes of
the selected 36 species using TBLASTN to identify the reference genes of each species
(**Extended Data Table 2**). Nucleotide and protein sequences of the reference genes for each
species were then searched against all genomes of the corresponding species using BLASTN
and TBLASTN. We combined BLASTN and TBLASTN results to identify the presence,
absence or pseudogenization of genes, with identify >80% and coverage >70% as cutoffs. A
gene was classified as present if a BLASTN or TBLASTN match was identified. If only a
BLASTN match was identified, the gene was classified as a pseudogene; otherwise, it was
considered absent.

Bacterial strains and growth conditions

A total of 12 *V. parahaemolyticus* representative strains (pairwise SNP distances: >6) were
used for laboratory experiments, including reference strain RIMD2210633 (wave-3) and 11
previously or newly sequenced PC strains. Additionally, *V. cholerae* strain E7946 (El Tor,
Ogawa) and *E. coli* strain MG1655 (K12) were included. We chose *E. coli* and *V. cholerae* for
two main reasons: our detection of natural strains with Puu-gene loss, and the availability of
strains in our laboratory. In addition, *V. cholerae* is another *Vibrio* species associated with

disease pandemics, and *E. coli* is a widely used model organism. *V. parahaemolyticus* strains were grown routinely in rich broth (LB with 3% NaCl) with shaking or on agar plates at 30°C aerobically. *E. coli* and *V. cholerae* strains were cultured in LB at 37°C.

Mutant construction

Deletion mutants of *V. parahaemolyticus* were constructed by homologous recombination and counterselection with the suicide plasmid pDM4 as previously described[73], with 1kb of homology upstream and downstream of the desired mutation. For *V. cholerae*, a similar approach with pDM4 was used as for *V. parahaemolyticus*, except *E. coli* MFD*pir* was used as donor strain for conjugation. *E. coli* deletion mutants were constructed using λ red recombination as previously described[74]. *V. parahaemolyticus* deletion strains were complemented with the corresponding gene *in trans* on plasmid pMMB207 using the Ptac promoter. All deletion and complement clones were verified by colony PCR. All constructed strains and plasmids are listed in **Supplementary Table 7**.

Biofilm assays

Log-phase cultures were inoculated into M9 medium with 1% NaCl in 96-well plates (Kingmorn biotechnology) and incubated under static conditions at 37°C for 48 hours. Following incubation, culture medium was discarded, and 96-well plates were washed with 0.9% NaCl twice, dried at 60 °C for 30 min, and stained with 100 µl 0.1% (w/v) crystal violet solution for 10 min. After staining, the plates were washed with 0.9% NaCl three times. The bound crystal violet was resolubilized with 100 µl 95% ethanol for 10 min and quantified via OD_{595} with BioTek Synergy H1 (Agilent Technologies). Each strain was tested with 30 technical replicates.

RNA extraction and transcriptome analysis

Strains were grown to stationary phase in M9 medium with 1% NaCl and harvested after treatment with stop mix (5% phenol, 95% ethanol). Total RNA was extracted from cells using the hot-phenol method as described previously[75]. RNA was DNase I-digested and rRNA depleted. cDNA libraries were constructed at Novogene Co. Ltd. (Beijing, China) and sequenced on an Illumina NovaSeq platform.

Sequencing reads were trimmed using Trimmomatic v0.39[54] and then mapped to the reference genome using BWA v0.7.17[76]. The number of reads assigned to genes were counted using featureCounts v2.0.6[77]. Differentially expressed genes and pathway enrichment analysis were performed using R packages DESeq2 v1.34.0[78] and ClusterProfiler v4.2.2[79].

c-di-GMP assay

To measure intracellular c-di-GMP, we used a previously published riboswitch-based reporter system[80]. Log-phase cultures of strains carrying the reporter were inoculated into M9 medium with 1% NaCl and 50 µg/ml spectinomycin in 96-well plates (six replicate wells per strain). Additionally, six wells of blank control with 200 µl uninoculated medium were included. The plate was incubated in a BioTek Synergy H1 plate reader (Agilent Technologies) at 37°C for 48 hours. OD_{600} and fluorescence (455 nm excitation/489 nm emission for AmCyan and 540

nm excitation/574 nm emission for TurboRFP) were measured every 15 min. The average values of the control wells were subtracted from each measurement (OD$_{600}$, AmCyan, and TurboRFP) of the strains. The mean values were used to plot growth curves and the whisker length represents standard deviation.

Cell adhesion

Log-phage strain cultures were washed with PBS and then resuspended in cell culture medium without antibiotics. Caco-2 cells grown in 24-well plates were infected with the bacterial suspension at an MOI of 10:1. After 45 min, cells were washed three times with PBS to remove the non-attached bacteria and then lysed with 0.1% saponin for 10 min. The number of cell-associated bacteria was determined by plating serial dilutions of the lysate on LB agar plates. The percentage of cell-associated bacteria was calculated as (recovered CFU/inoculated CFU) × 100%, and the results were presented as adhesion relative to wild-type strain (RIMD-W3).

Mouse colonization

Female BALB/c mice aged 6-8 weeks were used to test strain colonization. Mice were randomly divided into three groups, including a control group and two groups inoculated with different strains (three mice each treated group). All mice were fasted for 8 hours prior to gavage.

We introduced plasmids with *lux* reporters (pWHQ205, encoding luciferase and substrate, constructed in this study) into two strains (RIMD-W3 and Δ*puuC*), respectively. Strains carrying *lux* reporters were cultured in LB medium with 3% NaCl supplemented with chloramphenicol (25 µg/ml) to logarithmic phase (OD$_{600}$=0.6~0.8) at 30°C. Strain concentration was then adjusted to 3×10^{10} CFU/ml with PBS, and each mouse was gavaged with 0.3 ml of the bacterial suspension. At 5 and 10 hours post-infection, the mice were anesthetized with 3% isoflurane for 5 minutes and exposed for 1 minute using the PerkinElmer IVIS Spectrum imaging system (E-Heng Import and Export Co. Ltd).

Mouse lethality

Female BALB/c mice aged 4-5 weeks were used for lethality experiment, which were randomly divided into three groups treated by different strains, with each consisting of 10 mice. Log-phage strain cultures were washed three times with PBS and then adjusted to a concentration of 5×10^7 CFU/ml with PBS. Mice were intraperitoneally injected with 100 µl of strain suspension, and the number of mice that died at a specific time point was record. The experimental protocols of colonization and lethality of mice were approved by the animal ethics committee of Shanghai Institute of Immunity and Infection (No. A2023011).

Growth competition assay

Log-phage cultures of 12 natural strains, including six wave-1/2/3 and six wave-4 strains (**Supplementary Table 7**), were individually diluted to an OD$_{600}$ of 0.08. Subsequently, the diluted cultures of wave-4 strains were mixed with those of wave-1 to wave-3 strains in equal (1:1) or different (1:5) ratios. Given the ecological relevance of chitin as a major nutrient

source for *Vibrio* spp. in marine environments, and its role as a sole carbon source in aquatic habitats[81], competitions were conducted in natural seawater (collected from Jinshan, Shanghai) supplemented with 0.5% chitin (Sigma). Mixed cultures were incubated at 30 °C under static conditions for 72 h.

Genomic DNA of harvested cultures was extracted using the TIANamp Bacteria DNA kit (Tiangen Biotech). PCR primers VPA0427SNPU (AGCGACGACGCCAGAGAAAG) and VPA0427SNPD (GGTGTCTAAAGTACGGTGCAG) were used to amplify the gene VPA0427, which contains a wave-4 specific SNP allele (A919G). The PCR amplification was performed for 30 cycles, with the parameters: 1) 3 min. at 95°C, 2) 30 sec. at 95°C, 3) 30 sec. at 58°C, 4) 45 sec. at 72°C, 5) repeat steps 2-4 for 30 cycles, 6) 5 min. at 72°C.

The PCR amplification products were used for library construction at Novogene Co. Ltd. (Beijing, China) and sequenced on an Illumina NovaSeq platform. Sequencing reads are trimmed using Trimmomatic v0.39[54] and were then mapped to *V. parahaemolyticus* reference genome using BWA v0.7.17[76]. The frequency of the wave-4 specific allele was calculated to represent the relative abundance of wave-4 strains.

Statistical analysis
Datasets were compared using the Mann-Whitney U test for continuous data and Chi-squared tests for categorical data. Log-rank (Mantel-Cox) tests were used for survival curve comparison. p-values less than 0.05 were considered statistically significant.

Data availability
The sequencing data have been deposited in the NCBI SRA or GenBank database under accession number PRJNA1117214 and PRJNA1062747. Background information of sequenced isolate is listed in Supplementary Table 1-2. Source data are provided with this paper.

References

1.   Bonneaud, C. & Longdon, B. Emerging pathogen evolution: Using evolutionary theory to understand the fate of novel infectious pathogens. *EMBO Rep* **21**, (2020).
2.   Balasubramanian, D., López-Pérez, M., Grant, T. A., Ogbunugafor, C. B. & Almagro-Moreno, S. Molecular mechanisms and drivers of pathogen emergence. *Trends in Microbiology* vol. 30 Preprint at https://doi.org/10.1016/j.tim.2022.02.003 (2022).
3.   Baker-Austin, C. *et al.* Vibrio spp. infections. *Nat Rev Dis Primers* **4**, 8 (2018).
4.   Hu, D. *et al.* Origins of the current seventh cholera pandemic. *Proc Natl Acad Sci U S A* **113**, E7730–E7739 (2016).
5.   Mutreja, A. *et al.* Evidence for several waves of global transmission in the seventh cholera pandemic. *Nature* **477**, 462–465 (2011).
6.   Didelot, X. *et al.* The Role of China in the Global Spread of the Current Cholera Pandemic. *PLoS Genet* **11**, (2015).
7.   Weill, F.-X. *et al.* Genomic History of the Seventh Pandemic of Cholera in Africa. Science, , 358: 785-789 (2017).

8. Baker-Austin, C., Stockley, L., Rangdale, R. & Martinez-Urtaza, J. Environmental
occurrence and clinical impact of Vibrio vulnificus and Vibrio parahaemolyticus: A
European perspective. *Environ Microbiol Rep* **2**, 7–18 (2010).

9. Yang, C. *et al.* Recent mixing of Vibrio parahaemolyticus populations. *ISME J* **13**,
2578–2588 (2019).

10. Nair, G. B. *et al.* Global dissemination of Vibrio parahaemolyticus serotype O3:K6 and its
serovariants. *Clin Microbiol Rev* **20**, 39–48 (2007).

11. Espejo, R. T., García, K. & Plaza, N. Insight into the origin and evolution of the Vibrio
parahaemolyticus pandemic strain. *Front Microbiol* **8**, 1–6 (2017).

12. Han, D., Yu, F., Chen, X., Zhang, R. & Li, J. Challenges in Vibrio parahaemolyticus
infections caused by the pandemic clone. *Future Microbiol* **14**, 437–450 (2019).

13. Matsumoto, C. *et al.* Pandemic spread of an O3:K6 clone of Vibrio parahaemolyticus and
emergence of related strains evidenced by arbitrarily primed PCR and toxRS sequence
analyses. *J Clin Microbiol* **38**, (2000).

14. Nasu, H. *et al.* A filamentous phage associated with recent pandemic Vibrio
parahaemolyticus O3:K6 strains. *J Clin Microbiol* **38**, (2000).

15. Hurley, C. C., Quirke, A. M., Reen, F. J. & Boyd, E. F. Four genomic islands that mark
post-1995 pandemic Vibrio parahaemolyticus isolates. *BMC Genomics* **7**, 1–19 (2006).

16. Boyd, E. F. *et al.* Molecular analysis of the emergence of pandemic Vibrio
parahaemolyticus. *BMC Microbiol* **8**, 1–14 (2008).

17. Cui, Y. *et al.* Epidemic clones, oceanic gene pools, and eco-LD in the free living marine
pathogen vibrio parahaemolyticus. *Mol Biol Evol* **32**, 1396–1410 (2015).

18. Sela, I., Wolf, Y. I. & Koonin, E. V. Theory of prokaryotic genome evolution. *Proc Natl
Acad Sci U S A* **113**, 11399–11407 (2016).

19. Luo, H. How Big Is Big? The Effective Population Size of Marine Bacteria. *Ann Rev Mar
Sci* **17**, 537–560 (2025).

20. Liu, Z., Hossain, S. S., Moreira, Z. M. & Haney, C. H. Putrescine and Its Metabolic
Precursor Arginine Promote Biofilm and c-di-GMP Synthesis in Pseudomonas aeruginosa.
J Bacteriol **204**, (2022).

21. Kline, K. A., Fälker, S., Dahlberg, S., Normark, S. & Henriques-Normark, B. Bacterial
Adhesins in Host-Microbe Interactions. *Cell Host and Microbe* vol. 5 Preprint at
https://doi.org/10.1016/j.chom.2009.05.011 (2009).

22. Yang, C. *et al.* Outbreak dynamics of foodborne pathogen Vibrio parahaemolyticus over a
seventeen year period implies hidden reservoirs. *Nat Microbiol* **7**, 1221–1229 (2022).

23. Datta, M. S. *et al.* Rapid methicillin resistance diversification in Staphylococcus
epidermidis colonizing human neonates. *Nat Commun* **12**, (2021).

24. Iwasaki, W. & Takagi, T. Rapid pathway evolution facilitated by horizontal gene transfers
across prokaryotic lineages. *PLoS Genet* **5**, (2009).

25. Domman, D. *et al.* Integrated View of Vibrio Cholerae in the Americas. Science **358**,
789-793 (2017).

26. Cui, Y. *et al.* Historical variations in mutation rate in an epidemic pathogen, Yersinia pestis.
Proceedings of the National Academy of Sciences **110**, 577–582 (2013).

27. Rabosky, D. L. & Lovette, I. J. Explosive evolutionary radiations: Decreasing speciation or
increasing extinction through time? *Evolution (N Y)* **62**, (2008).

28. Morlon, H., Kemps, B. D., Plotkin, J. B. & Brisson, D. Explosive radiation of a bacterial
species group. *Evolution (N Y)* **66**, (2012).

29. Markov, P. V. *et al.* The evolution of SARS-CoV-2. *Nature Reviews Microbiology* **21**
(2023).

30. Flemming, H. C. *et al.* Biofilms: An emergent form of bacterial life. *Nature Reviews*
Microbiology **14** (2016).

31. Wunderlichová, L., Buňková, L., Koutný, M., Jančová, P. & Buňka, F. Formation,
Degradation, and Detoxification of Putrescine by Foodborne Bacteria: A Review. *Compr*
Rev Food Sci Food Saf **13**, (2014).

32. Krammer, E. M. & Prévost, M. Function and Regulation of Acid Resistance Antiporters.
Journal of Membrane Biology **252**, (2019).

33. Romano, A., Ladero, V., Alvarez, M. A. & Lucas, P. M. Putrescine production via the
ornithine decarboxylation pathway improves the acid stress survival of Lactobacillus
brevis and is part of a horizontally transferred acid resistance locus. *Int J Food Microbiol*
175, (2014).

34. Hohmann, A. W., Schmidt, G. & Rowley, D. Intestinal colonization and virulence of
Salmonella in mice. *Infect Immun* **22**, (1978).

35. Wang, Y. *et al.* Isolation, phylogenetic group, drug resistance, biofilm formation, and
adherence genes of Escherichia coli from poultry in central China. *Poult Sci* **95**, (2016).

36. Flament-Simon, S. C. *et al.* Association between kinetics of early biofilm formation and
clonal lineage in Escherichia coli. *Front Microbiol* **10**, (2019).

37. Hogins, J., Xuan, Z., Zimmern, P. E. & Reitzer, L. The distinct transcriptome of
virulence-associated phylogenetic group B2 Escherichia coli . *Microbiol Spectr* **11**, (2023).

38. Nowrouzian, F. L., Wold, A. E. & Adlerberth, I. Escherichia coli strains belonging to
phylogenetic group B2 have superior capacity to persist in the intestinal microflora of
infants. *Journal of Infectious Diseases* **191**, (2005).

39. Grozdanov, L. *et al.* Analysis of the genome structure of the nonpathogenic probiotic
Escherichia coli strain Nissle 1917. *J Bacteriol* **186**, (2004).

40. Chattaway, M. A., Schaefer, U., Tewolde, R., Dallman, T. J. & Jenkins, C. Identification of
Escherichia coli and shigella species from whole-genome sequences. *J Clin Microbiol* **55**,
(2017).

41. Viana, D. *et al.* A single natural nucleotide mutation alters bacterial pathogen host tropism.
Nat Genet **47**, (2015).

42. Arnold, B. J., Huang, I. T. & Hanage, W. P. Horizontal gene transfer and adaptive
evolution in bacteria. *Nature Reviews Microbiology* **20** 206–218 (2022).

43. Rodríguez-Beltrán, J., DelaFuente, J., León-Sampedro, R., MacLean, R. C. & San Millán,
Á. Beyond horizontal gene transfer: the role of plasmids in bacterial evolution. *Nature*
Reviews Microbiology **19** 347–359 (2021).

44. Albalat, R. & Cañestro, C. Evolution by gene loss. *Nature Reviews Genetics* **17** 379–391
(2016).

45. Wolf, Y. I. & Koonin, E. V. Genome reduction as the dominant mode of evolution.
BioEssays **35**, (2013).

46. Pallen, M. J. & Wren, B. W. Bacterial pathogenomics. *Nature* **449**, 835–842 (2007).

47. McCutcheon, J. P. & Moran, N. A. Extreme genome reduction in symbiotic bacteria.

Nature Reviews Microbiology **10** 13–26 (2012).

48.   Maurelli, A. T., Fernández, R. E., Bloch, C. A., Rode, C. K. & Fasano, A. "Black holes"
and bacterial pathogenicity: A large genomic deletion that enhances the virulence of
Shigella spp. and enteroinvasive *Escherichia coli*. *Proc Natl Acad Sci U S A* **95**,
3943–3948 (1998).

49.   Maurelli, A. T. Black holes, antivirulence genes, and gene inactivation in the evolution of
bacterial pathogens. *FEMS Microbiology Letters* **267** (2007).

50.   Bliven, K. A. & Maurelli, A. T. Antivirulence genes: Insights into pathogen evolution
through gene loss. *Infection and Immunity* **80** 4061–4070 (2012).

51.   Dufresne, A., Garczarek, L. & Partensky, F. Accelerated evolution associated with genome
reduction in a free-living prokaryote. *Genome Biol* **6**, (2005).

52.   Morris, J. J., Lenski, R. E. & Zinser, E. R. The black queen hypothesis: Evolution of
dependencies through adaptive gene loss. *mBio* **3**, (2012).

53.   Alizon, S., Hurford, A., Mideo, N. & Van Baalen, M. Virulence evolution and the trade-off
hypothesis: History, current state of affairs and the future. *Journal of Evolutionary Biology*
22 (2009).

54.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

55.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
Assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043–1055 (2015).

56.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.
Nat Commun **9**, 1–8 (2018).

57.   Delcher, A. L., Salzberg, S. L. & Phillippy, A. M. Using MUMmer to Identify Similar
Regions in Large Sequence Sets . *Curr Protoc Bioinformatics* **00**, (2003).

58.   Page, A. J. *et al.* SNP-sites: rapid efficient extraction of SNPs from multi-FASTA
alignments. *Microb Genom* **2**, e000056 (2016).

59.   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*
Res **27**, 573–580 (1999).

60.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately
Maximum-Likelihood Trees for Large Alignments. *PLoS One* **5**, e9490 (2010).

61.   Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant
bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* **43**, e15 (2015).

62.   Edwards, D., Duchêne, S., Pope, B. & Holt, K. SNPPar: identifying convergent evolution
and other homoplasies from microbial whole-genome alignments. **264017**, 1–33 (2020).

63.   Kozlov, A. M. *et al.* RAxML-NG: A fast, scalable and user-friendly tool for maximum
likelihood phylogenetic inference. *Bioinformatics* (2019).

64.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. Y. ggtree: an r package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol Evol* **8**, (2017).

65.   Tonkin-Hill, G., Lees, J. A., Bentley, S. D., Frost, S. D. W. & Corander, J. Fast hierarchical
Bayesian analysis of population structure. *Nucleic Acids Res* **47**, 5539–5549 (2019).

66.   Rambaut, A., Lam, T. T., Carvalho, L. M. & Pybus, O. G. Exploring the temporal structure

of heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol* **2**, (2016).

67. Didelot, X., Croucher, N. J., Bentley, S. D., Harris, S. R. & Wilson, D. J. Bayesian
inference of ancestral dates on bacterial phylogenetic trees. *Nucleic Acids Res* **46**, (2018).

68. Volz, E. M. & Didelot, X. Modeling the growth and decline of pathogen effective
population size provides insight into epidemic dynamics and drivers of antimicrobial
resistance. *Syst Biol* **67**, 719–728 (2018).

69. van der Graaf-Van Bloois, L., Chen, H., Wagenaar, J. A. & Zomer, A. L. Development of
Kaptive databases for Vibrio parahaemolyticus O-and K-antigen genotyping. *Microb*
Genom **9**, (2023).

70. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069
(2014).

71. Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo
pipeline. *Genome Biol* **21**, 180 (2020).

72. Parks, D. H. *et al.* GTDB: An ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy.
Nucleic Acids Res **50**, D785–D794 (2022).

73. Chimalapati, S., Lafrance, A. E., Chen, L. & Orth, K. Vibrio parahaemolyticus: Basic
Techniques for Growth, Genetic Manipulation, and Analysis of Virulence Factors. *Curr*
Protoc Microbiol **59**, (2020).

74. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, (2000).

75. Sharma, C. M., Darfeuille, F., Plantinga, T. H. & Vogel, J. A small RNA regulates multiple
ABC transporter mRNAs by targeting C/A-rich elements inside and upstream of
ribosome-binding sites. *Genes Dev* **21**, (2007).

76. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

77. Liao, Y., Smyth, G. K. & Shi, W. FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, (2014).

78. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, (2014).

79. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
The Innovation **2**, 100141 (2021).

80. Zhou, H. *et al.* Characterization of a natural triple-tandem c-di-GMP riboswitch and
application of the riboswitch-based dual-fluorescence reporter. *Sci Rep* **6**, (2016).

81. Meibom, K. L. *et al.* The Vibrio cholerae chitin utilization program. *Proc Natl Acad Sci U*
S A **101**, (2004).

Acknowledgements
This work was supported by National Key Research and Development Program of China (No.
2022YFC2304700 to F.D.), National Natural Science Foundation of China (No. 32270003
and No. 32000008 to C.Y., No. 32170640 and No.32211550014 to D.F., No. 32250610209 to
S.S.), Youth Innovation Promotion Association, Chinese Academy of Sciences (No. 2022278
to C.Y.), Ministerio de Ciencia e Innovación (Spain) Grant PID2021-127107NB-I00 (J.M.U.),
Generalitat de Catalunya Grant 2021 SGR 00526 (J.M.U.), Shanghai Rising-Star Program
(No. 23QA1410500 to C.Y.), Shanghai Public Health System Construction Three-Year Action
Plan (No. GWVI-11.1-43 to H.Q.) and Wuxi Science and Technology Development Fund's

'Light of Taihu Lake' Science and Technology Research Program (Basic Research) (No.
K20231033 to C.N., No. K20231045 to S.G.). We are grateful to the labs of Prof. Qiyao Wang,
Kim Orth, and Jin He for providing strains and plasmids. We thank Prof. Lei Pan for valuable
comments.
873

Author contributions
C.Y., J.M.-U. and D.F. designed, initiated and coordinated the study. J.M.-U., L.X, Y.L., M.J.,
X.S. and Q.H. contributed to data collection. C.Y., Z.J., J.W., and W.X. performed
bioinformatics analysis. H.Q, S.S., C.N., S.G., Z.J., H.W., Y.Q. and S.L. performed
experiments. All authors contributed to interpretation of the data. C.Y. wrote the first draft of
the paper and H.Q., S.S., Y.Z., Z.Z., Y.C., J.M.-U., H.W., R.Y., Y.C. and D.F. reviewed and
revised the paper. All authors read and approved the final manuscript.
881

882 **Competing interests**
883 The authors declare no competing interests.
884

      Figure legends

Figure 1. From pre-pandemic (pre-PC) to pandemic clone (PC). a) Maximum likelihood

tree of 8,684 global isolates. Pre-PC and PC isolates are colored in blue and red, respectively.

b) Inferred evolutionary history from pre-PC to PC. The status of known PC marker

genes/islands in the ancestral strain is indicated by blue ellipses, while subsequent gains or

losses are represented by orange and black triangles. The numbers next to tree nodes indicate

inferred divergence times and 95% confidence intervals. Dashed lines represent lineages with

unknown origination times due to a lack of significant temporal signal. c) Distribution of

virulence-associated loci and known PC marker genes/islands among pre-PC, PC and non-PC

strains. Double slashes indicate artificially shortened phylogenetic branches. Heatmap fill

colors represent the frequency of presence for each locus across groups.

Figure 2. Fine-scale population structure, spatiotemporal and population dynamics of

PC. a) Maximum likelihood tree of 942 representative PC strains. The colors represent

different waves. Double slashes indicate artificially truncated branches. b) Dated phylogeny

of PC (left) and the geographical, subtyping and distribution of genes in the putrescine

utilization (Puu) pathway of strains (right). The colors of tree branches indicate geographic

areas, and numbers next to tree nodes indicate inferred divergence times and 95% confidence

intervals. TCs indicate transmissions resulting in local colonization. (c) Temporal dynamics of

four waves. Bar colors indicate the frequency of different waves. The line represents the

number of representative strains. (d) Temporal distribution of representative strains in

different countries and regions. The size of circles scales with the number of representative

strains. (e) Effective population size dynamics of different waves of the PC.

Figure 3. Effect of gene-loss in putrescine utilization (Puu) pathway. a) Positions of

wave-4 specific variations on the reference genome (top) and gene structure and coverage of

Puu-genes (colored in blue) in different waves (bottom). b) Summary of Puu-genes and

metabolites based on KEGG annotations. The *puuC* gene, which appears in three sub-pathway,

was highlighted in bold. c) Biofilm levels in M9 minimal media with 1% NaCl measured by

optical density at 595 nm (OD_{595}) of wildtype (RIMD-W3), mutant ($\Delta puuC$) and complement

strains (C-$\Delta puuC$). d) Transcriptomic volcano plots showing differentially expressed genes

(DEGs). The colors represent genes of specific category as in the legend. Red and blue

backgrounds represent up- and down-regulated genes in $\Delta puuC$ compared to wildtype strain. e)

c-di-GMP dynamics of wildtype (RIMD-W3) and mutant ($\Delta puuC$) strains. f) The adhesion

ability of strains to Caco-2 cells relative to wildtype (RIMD-W3). g) *In vivo* bioluminescent

imaging of mice at 5 hours post-infection by gavage. h) Survival curves of mice by

intraperitoneal infection. Mann-Whitney U test was used for pairwise comparisons except for

panel h, where log-rank (Mantel-Cox) tests were applied. *: $p<0.05$, **: $p<0.01$.

Figure 4. Phenotypes and clinical symptoms of wave1-3 and wave-4 strains. a) Biofilm

level and adhesion ability between wave1-3 and wave-4 strains. b) Biofilm level and adhesion

ability between the wave-4 wildtype strain (VPCZ58, wave-4 WT) and its complemented

strain (wave-4 WT C-(VP1775-1786)) carrying the nine puu-related genes (VP1775-1786). c)

Clinical symptoms of wave-4 and other wave-associated patients. d) Wave-4 specific allele

frequency after 72 hours of competition under seawater with chitin conditions. The light blue
and dark blue bars represent the mixture of natural strains before and after competition,
respectively. The Mann-Whitney U test was used for pairwise comparisons, except for panel c,
where Chi-squared tests were applied. *: $p<0.05$, **: $p<0.01$.
933
934 **Figure 5. Puu-genes in 36 bacterial species.** a) Prevalence of Puu-genes in 36 species.
935 Genes and sub-pathways (SP) on the left are as in Fig. 3b. The colors (right) indicate the
936 frequency of genes. b) Phylogenetic distribution of Puu-genes in 10 species and the biofilm
937 and adhesion of wildtype (E7946 and MG1655) and Δ*puuC* mutant strains in *V. cholerae* and
938 *E. coli*. Mann-Whitney U test was used for pairwise comparisons. *: $p<0.05$, **: $p<0.01$.

a

Pre-PC

Pandemic clone (PC)

0.01

b

1920 (1901-1934)

orf8, VPaI-1/2/4/5/6, VPaI-7 (*tdh*/T3SS2)

PC

1734 (1692-1769)

1919 (1901-1933)

%"**JA,K4$L5

Pre-PC4b

Pre-PC4a

toxRS/new

T6SS1

VPaI-2

Pre-PC3b

Pre-PC3a

trh

VPaI-3

Pre-PC2

O3:K6, T3SS1+, T6SS2+, f237+, VPaI-3+

Pre-PC1

1740   1770   1800   1830   1860   1890   1920   1950   1980   2010

c

	Virulence-associated loci							PC marker gene/island									
	Clinical	O3:K6	ST3	*tdh*	*trh*	T3SS1	T6SS1	T6SS2	*toxRS*/new	*orf8*	f237	VPaI-1	VPaI-2	VPaI-3	VPaI-4	VPaI-5	VPaI-6	VPaI-7 (T3SS2)
PC (n=2815)																		
Pre-PC4b (n=18) 1980-2014, Asia, US																		
Pre-PC4a (n=3) 2006/2018, US																		
Pre-PC3b (n=1) 2014, China																		
Pre-PC3a (n=3) 2015/2022, Asia																		
Pre-PC2 (n=18) 1985-2008, Asia																		
Pre-PC1 (n=1) 1980, Bangladesh																		
Non-PC (n=5830)																		

Presence frequency (%)

a

Chromosome 1 (NC_004603.1)   Chromosome 2 (NC_004605.1)

— SNP
— Accessory gene
— Indel

Putrescine utilization (Puu) pathway

Coverage (%)

— Wave 1
— Wave 2
— Wave 3
— Wave 4

Super-integron

Genome position (Mb)

b

Putrescine (Put)

puuA (6.3.1.11)
puuB (1.4.3.-)
puuC (1.2.1.99)
puuD (3.5.1.94)
SP1

speG (2.3.1.57)
1.4.3.4
puuC (1.2.1.3)
SP2

patA (2.6.1.82) *spuC* (2.6.1.113)
puuC (1.2.1.3)
puuC (1.2.1.54) *patD/puuC* (1.2.1.19)
SP3

1.5.1.43
4.1.1.96
spdH (1.5.99.6)
SP6

speE (2.5.1.16)
SP5

Spermidine

SP4

γ-aminobutyric acid (GABA)

c

Biofilm level (OD$_{595}$)

RIMD-W3   Δ*puuC*   C-Δ*puuC*

d

Δ*puuC vs.* RIMD-W3

-log$_{10}$ (*p*)

- Biofilm pathway
- c-di-GMP
- Quorum sensing
- Flp pili
- CupE fimbriae

log$_2$(FC)

VP2422
flp
VP1979
VP2421
rcpC
VPA0518
cusA
cusB
cylA *cusC*
VPA0609
opaR *cusD*
toxB2
epsD *epsC*
epsE VPA0869
cusE
VPA1511

ompU
VP2446
zur
toxB1
VP0699 *crp*
VP2888 *tdh*

f

Relative adhesion (Caco-2)

RIMD-W3   Δ*puuC*   C-Δ*puuC*

g

RIMD-W3

Δ*puuC*

PBS

1(×10^7)
Luminescence

e

Relative c-di-GMP level

— RIMD-W3
— Δ*puuC*

Time (h)

h

Probability of survival (%)

— RIMD-W3
— Δ*puuC*
— C-Δ*puuC*

Time (h)