Trait lability as a predictor of diversification dynamics in flowering plants

Abstract

Rates of diversification differ between angiosperm lineages. To date, attempts to explain this

heterogeneity have focused on the potential correlation between speciation and extinction rates

and particular key traits. However, an often-overlooked explanation is that evolutionary lability,

here defined as the rates of trait change, may be a better predictor of speciation and extinction

rate heterogeneity than the observed traits themselves. Here, we show how this can be tested by

using hidden Markov models (HMMs), which allow for several rate classes associated with

speciation, extinction, and transition between trait states across a phylogeny. Using a

phylogenetic dataset of 13 angiosperm clades including 10,474 species, we show that higher

rates of change between open and closed-canopy biomes is consistently associated with higher

lineage turnover rates (speciation + extinction rates) across clades. We demonstrate how HMMs

can be leveraged in ways that go beyond their conventional use as null models in diversification

analyses, and that comparing different rate classes can unveil novel patterns of biological

interest. These patterns result in a shift in focus from static traits to dynamic evolutionary

processes and may provide a more comprehensive understanding into how biodiversity is

generated and maintained, in angiosperms and other organisms.

Introduction

It is widely accepted that rates of speciation and extinction differ between clades. Nonetheless, the cause of this heterogeneity remains poorly understood despite the consequences being immediately apparent. Angiosperms, or flowering plants, are by far the most diverse clade of green plants alive today with c. 300,000 described species [1]. But less appreciated is that the same degree of disparity in species richness between angiosperms and other green plant clades is repeatedly observed among clades within angiosperms. For example, there are ten times more species of daisies (Asteraceae) than those of primroses (Primulaceae s.s.) and nine times more species of grasses (Poaceae) than sedges (Cyperaceae) [2]. This disparity can be observed at all taxonomic levels and between clades which have had the same amount of time for species accumulation [3].

Differences in clade richness are mainly explained by differences in rates of speciation and extinction. To date, most explanations of clade specific diversification patterns have relied on discovering innovative traits, novel landscapes, or both ("key opportunities") which correlate with diversification rates [4–7]. Such events are theorized to correlate with increased or decreased chances of speciation or extinction due to the opportunity for population isolation, differentiation, or resilience against extinction [4–7]. Though this search has been fruitful, there has been a lack of consistency in which traits are affecting diversification dynamics in general. In fact, despite hundreds of studies describing a relationship between traits, biogeography, and diversification (reviewed in [8]), studies which have utilized modern, more appropriate, null models (e.g. [9]) have often found little support for character dependent diversification. Even in cases where there is evidence for character dependent diversification, the characters are not consistent drivers of diversification across multiple clades [10].

This lack of generalities suggests that alternative explanations for observed heterogeneity in rates of diversification should be sought. One potential venue that has been suggested is that rates of speciation and extinction correlate with rates of trait change – that is, trait lability [6,8,11]. The botanist G. Ledyard Stebbins was perhaps one of the first to explicitly suggest this possibility in flowering plants [12]. He proposed that the high capacity of angiosperms to adapt to environmental change was the key to their rich diversity. His mechanism was that long term climatic instability led to cyclical changes in the location of climatic zones [13]. For lineages in these unstable regions, the constant fluctuation in the climatic setting would be responsible for

continuous allopatry and adaptive pressure for phenotypic change, as well as leading to many

ephemeral species and subsequent extinction [12,14]. Furthermore, it is expected that, although

observed patterns at the macroevolutionary scale suggest that plant lineages tend to track their

preferred environments over time [15,16], when change is rapid, tracking is likely to be

imperfect, and phenotypic change or extinction will follow [12,14]. In other words, the

continuous movement of lineages between biome types would increase probabilities of both

speciation and extinction over time, i.e. evolutionary turnover (sensu [17]).

Another possible reason for the lack of consensus on the drivers of diversification is

methodological. For instance, early work quantified diversification rates across angiosperm

clades revealing the uneven distribution of species richness without testing potential explanatory

factors [3]. As the field advanced, more complex models emerged, with State-dependent

Speciation and Extinction (SSE) models becoming particularly popular for testing hypotheses

about the link between specific traits and diversification rates [18]. However, this initial

enthusiasm was tempered by the growing recognition that standard SSE models were prone to

high rates of false positives [19]. This led to the development of new "null" SSE models which

incorporate "hidden" variation by applying a hidden Markov model (HMM) framework states

and account for rate heterogeneity that is not associated with the observed trait of interest [9].

Despite these methodological improvements, studies have often found a lack of support for trait-

dependent diversification when these more robust models are applied [10]. This suggests that

while hidden Markov models are useful for avoiding spurious correlations, they also highlight

the complexity of diversification dynamics.

Here, we test the hypothesis that rates of transition between open-canopy biomes and

closed canopy biomes positively correlate with turnover and net-diversification rates across

many angiosperm clades. To this end, we use a modeling framework that combines properties of

HMMs [20] and state-speciation and extinction (SSE) models [9,18] to allow for both jointly

estimating transition and diversification dynamics in a clade and accounting for heterogeneity in

those rates at different parts of the phylogenetic tree, i.e. different rate classes. For our dataset,

we sample 13 flowering plant clades, which combined span about 10,474 species where 5,257

species are found in closed-canopy, 3,305 in open-canopy and 1,912 are widespread across both

biome types. For each clade, we build hidden Markov SSE models and interpret parameter

estimates using non-parametric sign tests. In this way, we are able to explicitly test whether rate

classes with faster transition rates between biomes also tend to be those with faster rates of

speciation and extinction.

Methods

Dataset assembly

The criterion to select the 13 clades used in this study was to pick clades that had a

minimum of 250 tips, with at least 0.4 of sampling fraction for the ingroup, i.e. inclusion of at

least 40% of the species diversity assigned to the corresponding clades. Although this is a

somewhat arbitrary threshold, it has been demonstrated that very low sampling fraction often

lead to inaccurate parameter estimates in trait dependent diversification analyses [21,22]; and

trees with less than 300 tips are unlikely to have the necessary information to properly fit

complex SSE models [23] and to present rate class heterogeneity (see details about rate classes

below). We kept our thresholds slightly more inclusive than these (i.e., including clades with as

low as 0.4 sampling fraction in Solanaceae and 267 tips in Onagraceae) to retain a minimum

diversity of clades across the angiosperm tree of life. We also focused on trees produced by

taxonomists of each group, since they are more careful about voucher identification and choice

of molecular markers used in reconstructions (e.g. [24]). We avoided using large trees inferred

by genbank scraping (e.g. [25]) because of issues with uneven sampling across the tree. In that

way, we combined strengths of the large sample approach in which to observe generalities from

with the care for accuracy that comes from small trees (see debate between [26] and [27]).

Finally, we also excluded trees that were not reciprocally monophyletic, keeping only the ones

with larger sampling and/or more tips when the same clade appears in more than one tree (e.g.

we excluded the *Acacia* phylogeny of [28] and kept the phylogeny of Mimosoid legumes of [29],

which includes *Acacia*). Tree root ages ranged from 9.84 (NeoAstragalus; [30]) to 105.71

(Fagales; [31]) million years to the root and from 267 (Onagraceae; [32]) to 2,202 (Arecaceae,

[33]) tips. To score the main biome type for each species, we first standardized all tip names

according to the GBIF taxonomic backbone using the R package taxize [34] and removed

outgroups following the original publications of each phylogeny. We also pruned trees to remove

tips of species with multiple entries in some phylogenies, leaving only one tip per species.

We then downloaded all the 41,496,903 occurrence points for the originally 11,593

species sampled in the 13 trees from GBIF [35–47], using POWO [48] taxonomy and

geographical information to filter non-native distributions and distribution inaccuracies. Because

GBIF is prone to sampling bias, we also thinned occurrence points to five representative

occurrence points per 1x1 grid cell for all species before further analyses (see [49] for detailed

methodology). Our filtering decreased the number of species with trustworthy information to

10,474. The remaining 2,264,540 points after filtering were overlaid on the World Wildlife Fund

(WWF) map of terrestrial biomes [50]. Biomes are generally defined as a combination of

climatic factors that drive convergent and/or parallel evolution in certain plant traits, leading to

characteristic vegetation types (or physiognomies) under specific climatic conditions [51].

125   Following Stebbins' hypothesis, we binarized the original 13 terrestrial biomes into

closed or open-canopy, dividing them along a precipitation gradient. Occurrence points placed

on biomes where precipitation is higher ("Tropical & Subtropical Moist Broadleaf Forests",

"Tropical & Subtropical Dry Broadleaf Forests", "Tropical & Subtropical Coniferous Forests",

"Temperate Broadleaf & Mixed Forests", "Temperate Conifer Forests", and "Boreal

Forests/Taiga") were scored as closed-canopy biomes. Those occurring in areas of lower

precipitation ("Tropical & Subtropical Grasslands, Savannas & Shrubland", "Temperate

Grasslands, Savannas & Shrublands", "Flooded Grasslands & Savannas", "Montane Grasslands

& Shrublands", "Tundra","Deserts & Xeric Shrublands", "Mediterranean Forests, and

Woodlands & Scrub") as well as "Mangroves" were scored as open-canopy biomes. Of course,

that does not mean that closed-canopy biomes only have closed-canopy habitats, as open-canopy

vegetations are often found within closed-canopy biomes and vice versa (e.g. prairies in

temperate rainforests, riverine forests in tropical savannas). It means closed canopy biomes are

dominated by closed-canopy vegetations and open-canopy biomes by open-canopy vegetations.

Species were scored as occurring in closed or open-canopy biomes when at least 25% of their

occurrence points were found in that biome type. If at least 25% of their occurrence points were

found in both biome types, then the species was scored as widespread. All datasets and R scripts

used in the curation process are available at [10.5281/zenodo.17401294].

Diversification and hidden state modeling

145   For each of the 13 angiosperm clades we use the R-package 'hisse' [52] to test for

correlations between diversification dynamics and rate of biome shift. We fit a set of 36 models:

4 without hidden states (one rate class) and 32 with hidden states (two rate classes). The basic

discrete character model of evolution was an all-rates-different model, in which transitions

between open and closed-canopy biome could only occur through an intermediate widespread

state (Figure 1A). Models with two rate classes were categorized into two types of discrete

character models. The first type allows for discrete character rate heterogeneity by permitting

transition rates to differ between rate classes. The second type has transition rates that are fixed

to be equal between rate classes. For the MuHiSSE model, in addition to trait evolution, we must

define the structure of the diversification dynamics in terms of extinction fraction

(extinction/speciation) and turnover (speciation+extinction).

For single rate class models, parameters are fixed to be equal for all observed states (no

trait-dependent diversification) or all are set to be different (biome dependent diversification).

For two rate class models, we explore 4 different parameterizations: (1) no observed trait-

dependent diversification and no hidden trait-dependent diversification, (2) observed trait-

dependent diversification, but no hidden trait-dependent diversification, (3) no observed trait-

dependent diversification, but hidden trait-dependent diversification, and (4) both observed trait-

dependent diversification and hidden trait-dependent diversification. We examined all possible

combinations of the parameterizations resulting in 36 model structures, and the best fitting model

for each clade was used to estimate the transition, speciation and extinction parameters for that

clade (Tables S1). Estimates of sampling fraction, a fixed parameter in the model, were gathered

from the literature of each clade (Table S2).

Parameter interpretation and analyses

Following model selection, we analyzed the parameter estimates from the best-fit model

for each clade to test for a consistent association between the rate of biome shifts and

evolutionary turnover. For each clade whose best-fit model included two distinct rate classes

(e.g., rate class 'A' and rate class 'B'), we extracted the mean biome transition rate, the mean

turnover rate (speciation + extinction), and mean net diversification rate (speciation - extinction)

separately for both rate classes (Figure 1B). A positive relationship was recorded if the rate class

with higher biome transition rates also had higher turnover (or net diversification), while a

negative relationship was recorded if it had lower rates (Figure 1C). To evaluate if a positive

association occurred more frequently than expected by chance across all clades, we conducted

separate non-parametric sign tests for turnover and net diversification.

A non-parametric approach is particularly appropriate for this analysis. It treats each

clade as an independent evolutionary experiment, testing for the consistency of the relationship's

direction rather than its numerical magnitude. This is crucial because the absolute values of

speciation, extinction, and transition rates may not be directly comparable across clades with

vastly different ages, sizes, and unique evolutionary histories [53]. Furthermore, this method is

robust to the "label switching" issue inherent in hidden Markov models, where "rate class A" in

one clade is not necessarily equivalent to "rate class A" in another. Because our test only

considers the internal correlation, a positive relationship is recorded regardless of whether it is

rate class A or B that has the faster rates. Under the null hypothesis of no consistent association,

we would expect a random mix of positive and negative relationships across the clades, yielding

a median of zero.

Results and Discussion

Biome shift rates are positively correlated with turnover rates

The colonization of new biomes is frequently hypothesized to be a major driver of

diversification, with some studies showing that geographic movement into new areas can be

more tightly linked to diversification than morphological innovation [53,54]. However, this

relationship is known to be complex and context-dependent, as not all dispersal events trigger

diversification bursts [55–57]. These contingent results suggest that the unique history of each

clade plays a significant role in shaping its evolutionary trajectory, potentially obscuring more

general patterns. A promising way to move beyond these clade-specific effects is to examine the

dynamics within major clades, treating each as an independent evolutionary experiment. This

approach allows for a more focused test of whether the intrinsic lability of a lineage (its

propensity to transition between biomes) is generally associated with a faster tempo of

diversification, rather than focusing on the impact of singular biome shifts.

Our analysis reveals a consistent, positive relationship between the rate of biome shifts

and the rate of evolutionary turnover across angiosperm clades, but no corresponding association

with net diversification. Using a non-parametric sign test, we found that the rate class with

higher biome transition rates was significantly more likely to also have higher rates of turnover

(speciation + extinction). The median of the slopes describing this relationship was positive

(0.23), and the 95% confidence interval for the median [0.00, 4.41] excludes negative values,

providing statistical support for a consistent positive association (p = 0.016). In contrast, we

found no consistent directional relationship between biome shift rates and net diversification

rates (speciation - extinction). For this comparison, the median slope was zero, with a wide 95%

confidence interval [0.00, 6.48], indicating a lack of any consistent directional trend (p = 0.125).

These patterns are evident in the parameter estimates from the best-fit model for each of

the 13 clades analyzed (Table 1; Table S3, S4). All of the clades showed support for hidden rate

classes and the majority (7 of 13) showed a positive association between transition rates and

turnover rates (Figure 2). This pattern was particularly strong in clades like Fagales, Onagraceae,

and Arecaceae, where the rate class with more frequent biome shifts also exhibited substantially

higher turnover. The case of Onagraceae is particularly interesting because it possessed a strong

positive association between biome shifts and turnover, but a strongly negative association with

net diversification (mTurn = 8.637, mNetDiv = -501.167). While the extreme magnitude of the

net diversification slope should be interpreted with caution (the potential for high variance in

parameter estimates is why a non-parametric test was used here), the opposing direction of the

two relationships is interesting. The Onagraceae result suggests a scenario where higher lability

is associated with higher turnover (more macroevolutionary events), but this increased turnover

is associated with lower net diversification (less lineages surviving to the present). In other

words, for this clade, the state characterized by frequent biome shifts accumulates species at a

slower pace than the more stable state and the increase in speciation rates associated with lability

is less than the corresponding increase in extinction.

Finally, for six of the thirteen clades (e.g., Mimosoids, Solanaceae, NeoAstragalus), the

best-fit models constrained biome transition rates to be equal across the hidden rate classes

(Figure 2; Table 1). In these instances, our analysis by definition could not detect a relationship

between lability and diversification, resulting in a slope of zero. However, the lack of any

negative associations across the remaining clades is informative. In every case where a

relationship could be assessed, higher rates of biome shifts were associated with higher—never

lower—rates of turnover. This suggests that while the strength of this coupling may vary, or may

not be detectable in all lineages, we found no evidence that increased lability ever acts to slow

the pace of evolution. Taken together, our results support the hypothesis that lability in biome

preference has red-queen type dynamics [58], elevating the pace of both speciation and

extinction, but it is not a consistent predictor of long-term species accumulation.

An alternative perspective on diversification analyses

State dependent diversification analyses (SSE models) began as a way to correct biases in

transition rates due to the possibility that observed states may be correlated with speciation and

extinction rates [18]. It was later discovered that this class of models was biased towards finding

associations between a focal character and diversification rates [59]. This was because SSE

models included heterogeneity in diversification parameters that would outperform alternative

models which did not, regardless of whether there was a true association between diversification

rates and the focal character [60]. Hidden Markov models, which were introduced as a more

biologically plausible model of evolution by allowing for rate heterogeneity independent of the

focal character (e.g. HiSSE, [9]) were able to be utilized as better null hypotheses and thus

overcame challenges of previous SSE models (see also [61]). This body of work set the stage for

the typical diversification analysis in which biologists test for an association between the focal

character and diversification parameters with the inclusion of a hidden Markov model as a null

hypothesis. More recently, it has also been suggested that these models are also relatively robust

to noted identifiability issues of birth-death models [60].

Although treating hidden Markov SSE models as null hypotheses is common practice,

there is a particularly powerful way in which these models have been underutilized. Instead of

treating hidden states merely as a nuisance variable to be accounted for, the different rate classes

inferred by these models can themselves be treated as the "trait" of interest. This approach allows

for a direct test of the hypothesis that the lability of diversification rates—the propensity for a

lineage to shift between different rates of speciation and extinction—is itself an important factor

in explaining broad-scale patterns of diversity. Hidden Markov models, therefore, are not just

null models against which to test trait-dependent hypotheses; they also offer a framework for

exploring the role of evolutionary lability in shaping diversification dynamics.

HMMs are explicitly parameterized by the dynamics of the observed characters, as this is

where the information for estimating rate classes is derived from [20]. This means that evidence

for hidden Markov models can come from both differences in diversification dynamics and

differences in observed character transition rates, and the potential relationships between these

processes has been generally under-explored. This is not to say that biologists have ignored the

potential for trait lability to drive patterns of diversification. Several studies have attempted to

correlate these processes using independent modeling results (e.g. [8,62]), but not in a

framework where the diversification and trait evolution parameters were estimated jointly.

Furthermore, unlike many studies which utilize SSE models, we are not primarily interested in

whether a particular character influences diversification. The hidden Markov SSE models allow

us to frame our hypothesis in such a way that the rate classes themselves are of primary interest.

This opens up potential examinations of other rate class dynamics as the underlying character

evolution model can be structured to match different biological hypotheses. In our case, we are

testing whether the rates of biome shifts are positively correlated with rates of net-diversification

and turnover in a joint framework, but the tempo of the models is just one aspect of discrete

character evolution.

Finally, this approach offers the potential to better account for the clade-specific

differences. By modeling rate heterogeneity within each major clade, it is possible to test for a

consistent relationship between state transitions (e.g., biome shifts) and diversification rates

without the unique, contingent histories of those clades acting as confounding factors. In this

way, each clade can be treated as its own evolutionary experiment, providing independent tests

of the hypothesis that higher rates of trait transition are correlated with faster rates of

diversification. The framework outlined here can be implemented using existing tools as it does

not require any new methodological developments. Instead, it relies on evolutionary biologists to

refocus their hypotheses in an examination of how different processes influence diversification

rather than particular character traits.

Conclusion

The significant relationship between rates of biome shifts and turnover highlights the role

that trait lability can play in shaping the diversification dynamics across many clades. Though

lability is not a given for all clades, when it is present it may be a signature of important

biological shifts which are reflected in both diversification and trait evolution. This fact

highlights one of the main methodological differences of our study. Rather than using SSE

models and hidden Markov SSE models to test whether a particular phenotype influences

diversification, we examine the possibility that diversification dynamics are influenced by the

processes underlying phenotypic evolution, the tempo of biome change in this case. This

possibility was recognized by G. Ledyard Stebbins 50 years ago and proposed by several others

[e.g., 8,11] to be one of the primary reasons for the high species diversity in angiosperms.

However, this conclusion does not entirely bear out in our results because net-diversification, the

parameter related to species accumulation, is not significantly tied to the tempo of biome shifts.

Instead, biome lability may be a double-edged sword increasing the propensity of lineage

speciation, but equally increasing extinction risk.

References

1. Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A. 2021 The World Checklist of Vascular Plants, a continuously updated resource for exploring global plant diversity. *Scientific data* **8**, 215.

2. Group AP *et al.* 2016 An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical journal of the Linnean Society* **181**, 1–20.

3. Magallon S, Sanderson MJ. 2001 Absolute diversification rates in angiosperm clades. *Evolution* **55**, 1762–1780.

4. Anderson B *et al.* 2023 Opposing effects of plant traits on diversification. *Iscience* **26**.

5. Givnish TJ. 2015 Adaptive radiation versus 'radiation' and 'explosive diversification': why conceptual distinctions are fundamental to understanding evolution. *New Phytologist* **207**, 297–303.

6. Donoghue MJ, Sanderson MJ. 2015 Confluence, synnovation, and depauperons in plant diversification. *New Phytologist* **207**, 260–274.

7. Givnish TJ. 2010 Ecology of plant speciation. *Taxon* **59**, 1326–1366.

8. Onstein RE. 2020 Darwin's second 'abominable mystery': trait flexibility as the innovation leading to angiosperm diversity. *New Phytologist* **228**, 1741–1747.

9. Beaulieu JM, O'Meara BC. 2016 Detecting hidden diversification shifts in models of trait-dependent speciation and extinction. *Systematic biology* **65**, 583–601.

10. Helmstetter AJ *et al.* 2023 Trait-dependent diversification in angiosperms: Patterns, models and data. *Ecology Letters* **26**, 640–657.

11. Ricklefs RE, Renner SS. 1994 Species richness within families of flowering plants. *Evolution* **48**, 1619–1636.

12. Stebbins GL. 1974 *Flowering plants: evolution above the species level*. Harvard University Press.

13. Jansson R, Dynesius M. 2002 The fate of clades in a world of recurrent climatic change: Milankovitch oscillations and evolution. *Annual review of ecology and systematics* **33**, 741–777.

14. Vasconcelos T, O'Meara BC, Beaulieu JM. 2022 Retiring "cradles" and "museums" of biodiversity. *The American Naturalist* **199**, 194–205.

15. Donoghue MJ. 2008 A phylogenetic perspective on the distribution of plant diversity. *Proceedings of the National Academy of Sciences* **105**, 11549–11555.

16. Wiens JJ, Donoghue MJ. 2004 Historical biogeography, ecology and species richness. *Trends in ecology & evolution* **19**, 639–644.

17. Vrba ES. 1985 Environment and evolution: alternative causes of the temporal distribution of evolutionary events. *South African Journal of Science* **81**, 229–236.

18. Maddison WP, Midford PE, Otto SP. 2007 Estimating a binary character's effect on speciation and extinction. *Systematic biology* **56**, 701–710.

19. Rabosky DL, Goldberg EE. 2015 Model Inadequacy and Mistaken Inferences of Trait-Dependent
Speciation. *Systematic Biology* **64**, 340–355. (doi:10.1093/sysbio/syu131)

20. Boyko JD, Beaulieu JM. 2021 Generalized hidden Markov models for phylogenetic comparative
datasets. *Methods in Ecology and Evolution* **12**, 468–478.

21. Mynard P *et al.* 2023 Impact of phylogenetic tree completeness and mis-specification of sampling
fractions on trait dependent diversification models. *Systematic Biology* **72**, 106–119.

22. Chang J, Rabosky DL, Alfaro ME. 2020 Estimating diversification rates on incompletely sampled
phylogenies: theoretical concerns and practical solutions. *Systematic Biology* **69**, 602–611.

23. Davis MP, Midford PE, Maddison W. 2013 Exploring power and parameter estimation of the BiSSE
method for analyzing species diversification. *BMC evolutionary biology* **13**, 38.

24. Group NMW *et al.* 2024 Towards a species-level phylogeny for Neotropical Myrtaceae: Notes on
topology and resources for future studies. *American Journal of Botany* **111**, e16330.

25. Smith SA, Brown JW. 2018 Constructing a broadly inclusive seed plant phylogeny. *American
journal of botany* **105**, 302–314.

26. Beaulieu JM, O'Meara BC. 2019 Diversity and skepticism are vital for comparative biology.
American Journal of Botany **106**, 613–617.

27. Donoghue MJ, Edwards EJ. 2019 Model clades are vital for comparative biology, and ascertainment
bias is not a problem in practice. *American Journal of Botany* **106**, 327–330.

28. Renner MA, Foster CS, Miller JT, Murphy DJ. 2020 Increased diversification rates are coupled with
higher rates of climate space exploration in Australian Acacia (Caesalpinioideae). *New Phytologist*
226, 609–622.

29. Ringelberg JJ *et al.* 2023 Precipitation is the main axis of tropical plant phylogenetic turnover across
space and time. *Science Advances* **9**, eade4954.

30. Folk RA, Charboneau JL, Belitz M, Singh T, Kates HR, Soltis DE, Soltis PS, Guralnick RP,
Siniscalchi CM. 2024 Anatomy of a mega-radiation: Biogeography and niche evolution in
Astragalus. *American Journal of Botany* **111**, e16299.

31. Xing Y, Onstein RE, Carter RJ, Stadler T, Peter Linder H. 2014 Fossils and a large molecular
phylogeny show that the evolution of species richness, generic diversity, and turnover rates are
disconnected. *Evolution* **68**, 2821–2832.

32. Leal ES, Vasconcelos TN, Tuberquia D, Soto Gomez M, Michelangeli FA, Forzza RC, de Mello-
Silva R. 2022 Phylogeny and historical biogeography of the Panama-hat family (Cyclanthaceae,
Pandanales). *Taxon* **71**, 963–980.

33. Onstein RE, Baker WJ, Couvreur TL, Faurby S, Herrera-Alsina L, Svenning J-C, Kissling WD. 2018
To adapt or go extinct? The fate of megafaunal palm fruits under past global change. *Proceedings of
the Royal Society B* **285**, 20180882.

34. Chamberlain SA, Szöcs E. 2013 taxize: taxonomic search and retrieval in R. *F1000Research* **2**, 191.

35. GBIF.org. 2025 Solanaceae GBIF Occurrence Download. (doi:10.15468/dl.5nxt2y)

36. GBIF.org. 2025 Salvia GBIF Occurrence Download. (doi:10.15468/dl.evpkmg)

37. GBIF.org. 2025 Restionaceae GBIF Occurrence Download. (doi:10.15468/dl.5dtgjd)

38. GBIF.org. 2025 Onagraceae GBIF Occurrence Download. (doi:10.15468/dl.qy8x4h)

39. GBIF.org. 2025 NeoAstragalus GBIF Occurrence Download. (doi:10.15468/dl.d3j4yb)

40. GBIF.org. 2025 Mimosoids GBIF Occurrence Download. (doi:10.15468/dl.6ec5vq)

41. GBIF.org. 2025 Fagales GBIF Occurrence Download. (doi:10.15468/dl.shpqzu)

42. GBIF.org. 2025 Eucalypts GBIF Occurrence Download. (doi:10.15468/dl.2khevd)

43. GBIF.org. 2025 Arecaceae GBIF Occurrence Download. (doi:10.15468/dl.u4msqv)

44. GBIF.org. 2025 Carex GBIF Occurrence Download. (doi:10.15468/dl.6mtpue)

45. GBIF.org. 2025 Arabideae GBIF Occurrence Download. (doi:10.15468/dl.d72qng)

46. GBIF.org. 2025 Apioideae GBIF Occurrence Download. (doi:10.15468/dl.4n2dtt)

47. GBIF.org. 2025 Antirrhineae GBIF Occurrence Download. (doi:10.15468/dl.fpcpzf)

48. POWO. 2025 Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew.

49. Boyko JD, Hagen ER, Beaulieu JM, Vasconcelos T. 2023 The evolutionary responses of life-history
strategies to climatic variability in flowering plants. *New Phytologist* **240**, 1587–1600.

50. Olson DM *et al.* 2001 Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new
global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity.
BioScience **51**, 933–938.

51. Mucina L. 2019 Biome: evolution of a crucial ecological and biogeographical concept. *New
Phytologist* **222**, 97–114.

52. Beaulieu J, O'Meara B, Caetano D, Boyko J, Vasconcelos T. 2017 Package 'hisse'.

53. Nürk NM *et al.* 2020 Diversification in evolutionary arenas—Assessment and synthesis. *Ecology and
Evolution* **10**, 6163–6182.

54. Moore BR, Donoghue MJ. 2007 Correlates of Diversification in the Plant Clade Dipsacales:
Geographic Movement and Evolutionary Innovations. *The American Naturalist* **170**, S28–S55.
(doi:10.1086/519460)

55. Stephens PR, Farrell MJ, Davies TJ, Gittleman JL, Meiri S, Moreira MO, Roll U, Wiens JJ. 2025
Global Diversity Patterns are Explained by Diversification Rates and Dispersal at Ancient, Not
Shallow, Timescales. *Syst Biol* , syaf018. (doi:10.1093/sysbio/syaf018)

56. Stull GW *et al.* 2021 Gene duplications and phylogenomic conflict underlie major pulses of
409   phenotypic evolution in gymnosperms. *Nat. Plants* **7**, 1015–1025. (doi:10.1038/s41477-021-00964-4)

410   57. Givnish TJ *et al.* 2014 Adaptive radiation, correlated and contingent evolution, and net species
411   diversification in Bromeliaceae. *Molecular Phylogenetics and Evolution* **71**, 55–78.
412   (doi:10.1016/j.ympev.2013.10.010)

413   58. Van Valen L. 1973 A new evolutionary law. *Evolutionary Theory* **1**, 1–30.

414   59. Rabosky DL, Goldberg EE. 2015 Model inadequacy and mistaken inferences of trait-dependent
415   speciation. *Systematic biology* **64**, 340–355.

416   60. O'Meara B, Beaulieu J. 2021 Potential survival of some, but not all, diversification methods.

417   61. Boyko JD, Beaulieu JM. 2023 Reducing the biases in false correlations between discrete characters.
418   *Systematic Biology* **72**, 476–488.

419   62. Rabosky DL, Adams DC. 2012 Rates of morphological evolution are correlated with species richness
420   in salamanders. *Evolution* **66**, 1807–1818.

Tables

Table 1 – Parameter estimates from the best-fit hidden Markov models. For each of the 13 clades, the table shows the Akaike weight (AICwt) of the best-fit model and the estimated mean rates for biome transitions (meanTrans), turnover (meanTurn), and net diversification (meanNetDiv) for the two hidden rate classes (A and B). The slope columns (slopeTurn, slopeNetDiv) represent the relationship between the rate of biome shifts and diversification rates. A slope of zero indicates that the best-fit model constrained transition rates to be equal across the hidden states.

clade	nTip	AICwt	meanTransA	meanTurnA	meanNetDivA	meanTransB	meanTurnB	meanNetDivB	slopeTurn	slopeNetDiv
Fagales	526	0.898	0.076	0.322	0.093	28.834	1.382	0.4	27.132	93.844
Onagraceae	267	0.768	39.369	4.802	0.026	0.007	0.245	0.105	8.637	-501.167
Arecaceae	2202	0.992	1.76	0.367	0.367	0.169	0.104	0.104	6.032	6.032
Arabideae	320	0.571	0.163	0.752	0.051	9.783	5.789	0.393	1.909	28.156
Carex	1221	0.798	0.336	0.281	0.281	0.596	0.582	0.582	0.861	0.861
Salvia	484	0.816	0.073	0.132	0.105	0.167	0.515	0.409	0.246	0.31
Eucalypts	691	1	2.081	9.035	0.302	0.129	0.39	0.013	0.226	6.765
Antirrhineae	276	0.194	0.453	0.167	0.167	0.453	0.458	0.458	0	0
Apioideae	1033	0.938	0.934	1.187	0.378	0.934	0.55	0.163	0	0
Mimosoids	1758	1	0.367	0.161	0.133	0.367	0.416	0.35	0	0
NeoAstragalus	364	0.997	34.923	0.725	0.725	34.923	0.162	0.162	0	0
Restionaceae	332	0.402	0.081	0.027	0.027	0.081	0.122	0.122	0	0
Solanaceae	1000	1	0.457	0.432	0.207	0.457	1.105	0.529	0	0

432 **Figures**

A. Model fit

B. Mean parameter values from best model

C. Slope of rate class parameter means

433
Figure 1 – Modeling framework for testing the association between biome shifting and
diversification dynamics. (A) A visual representation of the MuHiSSE model, which
incorporates three observed biome states (O = Open-canopy, W = Widespread, C = Closed-
canopy) and two unobserved "hidden" rate classes (A and B). Within each state and class, the
model estimates rates of speciation (λ), extinction (μ), and biome transition (q). (B) Illustration
of the parameter extraction from the best-fit model for a single clade. For each of the two hidden
rate classes, we extracted the mean biome transition rate and the mean evolutionary turnover rate
(speciation + extinction). (C) Each clade is treated as an independent data point testing the
association between biome transition rate and turnover. A "positive relationship" (blue lines) is
recorded if the hidden rate class with higher transition rates also has higher turnover. A "negative
relationship" (red lines) is recorded if it has lower turnover.
445

446
Figure 2 – Phylogenetic distribution of biome affiliation and rate-class heterogeneity across 13
angiosperm clades. Each circular phylogeny represents a single focal clade, with tips color-coded
according to biome association: closed-canopy (dark green), open-canopy (orange), or
widespread (light green). Phylogenetic trees are scaled by evolutionary time, and internal
branching patterns illustrate inferred diversification dynamics from the best-fitting hidden-state
speciation and extinction (HiSSE) models. The bottom panel shows the global distribution of
binarized biome types (closed- versus open-canopy) used to score species occurrences based on
GBIF records and WWF biome classifications. Photographs depict representative closed- (top)
and open-canopy (bottom) habitats.
456
457

Table S1 – This table outlines the structure of the 36 unique models tested for each angiosperm
clade. Each row represents a distinct model defined by the combination of parameter settings for
four primary components: extinction fraction (ef), turnover (turn), the transition rate matrix (q),
and the rate class structure (rc). The number associated with each component (e.g., ef1, ef2)
indicates a specific parameterization that dictates how rates are allowed to vary. For instance, ef1
corresponds to a model with no variation in the extinction fraction across states. The 36 models
represent the full set of combinations of these different configurations for extinction, speciation,
and biome transition dynamics.

ef	turn	q	rc
1	1	1	1R
2	1	1	1R
1	2	1	1R
2	2	1	1R
1	1	1	2R
2	1	1	2R
3	1	1	2R
4	1	1	2R
1	2	1	2R
2	2	1	2R
3	2	1	2R
4	2	1	2R
1	3	1	2R
2	3	1	2R
3	3	1	2R
4	3	1	2R
1	4	1	2R
2	4	1	2R
3	4	1	2R
4	4	1	2R
1	1	2	2R
2	1	2	2R
3	1	2	2R
4	1	2	2R
1	2	2	2R
2	2	2	2R
3	2	2	2R
4	2	2	2R
1	3	2	2R
2	3	2	2R
3	3	2	2R
4	3	2	2R
1	4	2	2R
2	4	2	2R
3	4	2	2R
4	4	2	2R

model class	paramers to vary (ef)	paramers to vary (turn)
CID	ef1 = 0,1,1,1,0,1,1,1	turn1 = 0,1,1,1,0,1,1,1
CD	ef2 = 0,1,2,3,0,1,2,3	turn2 = 0,1,2,3,0,1,2,3
CID	ef3 = 0,1,1,1,0,2,2,2	turn3 = 0,1,1,1,0,2,2,2
CD	ef4 = 0,1,2,3,0,4,5,6	turn4 = 0,1,2,3,0,4,5,6

parameters to vary (q)	parameters to vary (rc)
q1 = no difference between rate classes	1R = one rate class
q2 = difference between rate classes	2R = tworate class

example	description
ef1,turn1,q1,1R	dull null. no state dependent differences.
ef2,turn2,q1,1R	bisse. classic state dependent diversification.
ef3,turn3,q1,1R	CID2. null model with hidden states
ef3,turn3,q2,1R	null model w.r.t. diversification, but allowing different trans rates

467
468
469
470

471 Table S2 – Sampling fractions for each clade.

label	group	author	year	doi	family	major_group	n_tips_ingroup	sf_ingroup
Antirrhineae-Gorospe_et_al-2020	Antirrhineae	Gorospe_et_al	2020	10.1111/jbi.13939	Plantaginaceae	asterids	336	1
Apioideae-Banasiak_et_al-2013	Apioideae	Banasiak_et_al	2013	10.1111/jbi.12071	Apiaceae	asterids	1194	0.41
Arabideae-Huang_et_al-2019	Arabideae	Huang_et_al	2019	10.1093/aob/mcz123	Brassicaceae	rosids	347	0.6
Arecaceae-Onstein_et_al-2018	Arecaceae	Onstein_et_al	2018	10.1098/rspb.2018.0882	Arecaceae	monocots	2048	1
Carex-MartinBravo_et_al-2019	Carex	MartinBravo_et_al	2019	10.1111/jse.12549	Cyperaceae	monocots	1386	0.69
Eucalypts-Thornhill_et_al-2019	Eucalypts	Thornhill_et_al	2019	10.1071/SB18015	Myrtaceae	rosids	716	0.9
Fagales-Xing_et_al-2014	Fagales	Xing_et_al	2014	10.1111/evo.12489	NA	rosids	579	0.42
Mimosoids_Ringelberg_et_al_2023	Mimosoids	Ringelberg_et_al	2023	10.1126/sciadv.ade4954	Fabaceae		2078	0.59
NeoAstragalus-Folk_et_al-2024	NeoAstragalus	Folk_et_al	2024	10.1002/ajb2.16299	Fabaceae		373	0.45
Onagraceae-Freyman_&_Hohna-2018	Onagraceae	Freyman_&_H√∂hna	2018	10.1093/sysbio/syy078	Onagraceae	rosids	292	0.45
Restionaceae-Linder-2020	Restionaceae	Linder	2020	10.1093/aob/mcaa129	Restionaceae	monocots	351	0.7
Salvia-Kriebel_et_al-2020	Salvia	Kriebel_et_al	2020	10.1111/evo.14030	Lamiaceae	asterids	519	0.52
Solanaceae-Sarkinen_et_al-2013	Solanaceae	Sarkinen_et_al	2013	10.1186/1471-2148-13-214	Solanaceae	asterids	1075	0.4

Table S3 – AIC weights for all 36 model fits for each clade. See Table S1 for the meaning of each model (M1 to M36).

Clade	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11	M12	M13	M14	M15	M16	M17	M18	M19	M20	M21	M22	M23	M24	M25	M26	M27	M28	M29	M30	M31	M32	M33	M34	M35	M36
ANTIRRHINEAE-GOROSPE_ET_AL-2020	0	0	0	0	0	0	0.028	0.012	0	0	0.005	0.006	0.097	0.024	0.083	0.034	0.194	0.026	0.014	0.004	0	0	0.01	0.022	0	0	0.084	0.001	0.016	0	0.026	0.008	0.194	0.095	0.011	0.005
APIOIDEAE-BANASIAK_ET_AL-2013	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.938	0	0	0	0	0	0	0	0	0	0	0	0	0	0.062	0	0	0	0	0	0	0
ARABIDEAE-HUANG_ET_AL-2019	0	0	0	0	0	0	0	0	0	0	0	0	0	0.001	0.004	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.571	0	0	0	0.423	0	0
ARECACEAE-ONSTEIN_ET_AL-2018	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.992	0	0	0.008
CAREX-MARTINBRAVO_ET_AL-2019	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.798	0.13	0.053	0.019
EUCALYPTS-THORNHILL_ET_AL-2019	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
FAGALES-XING_ET_AL-2014	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.898	0.001	0	0	0.055	0	0.046
MIMOSOIDS_RINGELBERG_ET_AL_2023	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NEOASTRAGALUS-FOLK_ET_AL-2024	0	0	0	0	0	0	0	0	0	0	0	0.003	0	0	0	0	0	0.997	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ONAGRACEAE-FREYMAN_&_HOHNA-2018	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.178	0.054	0	0	0	0.768
RESTIONACEAE-LINDER-2020	0	0	0.023	0	0.006	0	0.008	0.005	0	0	0.002	0	0	0.402	0.003	0.158	0.079	0	0.107	0.002	0	0	0	0	0	0	0	0	0.159	0.037	0	0.009	0	0	0	0
SALVIA-KRIEBEL_ET_AL-2020	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.002	0	0.003	0.005	0.012	0.022	0.001	0	0.816	0	0.076	0.034	0.025	0.001	0.002
SOLANACEAE-SARKINEN_ET_AL-2013	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

476 Table S4 – Parameter estimates for each of the best fitting models.

clade	model	loglik	AICc	AICwt	turnover01A	turnover10A	turnover11A	eps01A	eps10A	eps11A	q01A_11A	q10A_11A	q11A_01A	q11A_10A	q01A_01B	q10A_10B
Antirrhineae-Gorospe_et_al-2020	M17	-881.263	1787.713	0.194	0.108	0.068	0.324	0	0	0	0.635	0.137	0.529	0.51	0.016	0.016
Apioideae-Banasiak_et_al-2013	M15	-3636.94	7292.056	0.938	1.187	1.187	1.187	0.517	0.517	0.517	0.687	0.844	1.009	1.196	0.009	0.009
Arabideae-Huang_et_al-2019	M30	-672.328	1374.033	0.571	0.752	0.752	0.752	1.433	0.335	1.264	0.197	0.168	0.257	0.031	0.315	0.315
Arecaceae-Onstein_et_al-2018	M33	-7347.001	14726.252	0.992	0.154	0.481	0.467	0	0	0	0.24	0.683	5.905	0.212	0.005	0.005
Carex-MartinBravo_et_al-2019	M33	-3960.829	7954.109	0.798	0.638	0.185	0.02	0	0	0	0.836	0.197	0.133	0.18	0.042	0.042
Eucalypts-Thornhill_et_al-2019	M33	-1593.268	3219.342	1	13.122	0.258	13.724	0.935	0.935	0.935	1.007	0.815	0.165	6.337	0.039	0.039
Fagales-Xing_et_al-2014	M30	-1921.178	3871.178	0.898	0.322	0.322	0.322	0.959	0.113	0.904	0.019	0.129	0.156	0	0.052	0.052
Mimosoids_Ringelberg_et_al_2023	M16	-7063.029	14152.266	1	0.161	0.161	0.161	0	0.17	0.132	0.146	0.206	0.331	0.784	0.004	0.004
NeoAstragalus-Folk_et_al-2024	M18	-834.901	1699.004	0.997	1.87	0	0.305	0	0	0	1.528	38.974	0.385	98.807	0.502	0.502
Onagraceae-Freyman_&_Hohna-2018	M36	-974.584	1994.94	0.768	0	0	14.407	0.129	0.792	0.989	5.035	100	3.422	49.021	0.06	0.06
Restionaceae-Linder-2020	M13	-1169.841	2356.129	0.402	0.027	0.027	0.027	0	0	0	0.012	0.003	0.046	0.262	0.023	0.023
Salvia-Kriebel_et_al-2020	M30	-1582.358	3193.611	0.816	0.132	0.132	0.132	0.448	0	0	0.07	0.073	0.018	0.13	0.043	0.043
Solanaceae-Sarkinen_et_al-2013	M14	-3002.94	6026.103	1	0.432	0.432	0.432	0.586	0	0.702	0.249	0.545	0.783	0.253	0.025	0.025

q11A_11B	turnover01B	turnover10B	turnover11B	eps01B	eps10B	eps11B	q01B_11B	q10B_11B	q11B_01B	q11B_10B	q01B_01A	q10B_10A	q11B_11A
0.016	0.204	0.54	0.63	0	0	0	0.635	0.137	0.529	0.51	0.016	0.016	0.016
0.009	0.55	0.55	0.55	0.543	0.543	0.543	0.687	0.844	1.009	1.196	0.009	0.009	0.009
0.315	5.789	5.789	5.789	1.433	0.335	1.264	16.314	2.538	20.279	0	0.315	0.315	0.315
0.005	0.039	0.204	0.068	0	0	0	0.051	0.256	0.288	0.081	0.005	0.005	0.005
0.042	0.138	0.423	1.185	0	0	0	0.034	0.889	1.239	0.22	0.042	0.042	0.042
0.039	1.17	0	0	0.935	0.935	0.935	0.081	0.079	0.244	0.111	0.039	0.039	0.039
0.052	1.382	1.382	1.382	0.959	0.113	0.904	13.833	1.427	100	0.078	0.052	0.052	0.052
0.004	0.416	0.416	0.416	0.01	0.123	0.139	0.146	0.206	0.331	0.784	0.004	0.004	0.004
0.502	0.487	0	0	0	0	0	1.528	38.974	0.385	98.807	0.502	0.502	0.502
0.06	0.396	0.192	0.147	0.621	0.151	0.297	0	0.012	0.016	0	0.06	0.06	0.06
0.023	0.122	0.122	0.122	0	0	0	0.012	0.003	0.046	0.262	0.023	0.023	0.023
0.043	0.515	0.515	0.515	0.448	0	0	0.032	0.197	0.431	0.007	0.043	0.043	0.043
0.025	1.105	1.105	1.105	0.586	0	0.702	0.249	0.545	0.783	0.253	0.025	0.025	0.025

477