Genome-wide association mapping for growth rate at fluctuating and extreme temperatures

Emmi Räsänen[1, *], Martta Liukkonen[1], Pauliina A. M. Summanen[1], Tarmo Ketola[2] and Ilkka Kronholm[1]

[1]Department of Biological and Environmental Science, PO Box 35, FI-40014 University of Jyväskylä, Finland
[2] Department of Forest Science, PO Box 27, FI-00014, University of Helsinki, Finland
*Corresponding author's e-mail: emmi.j.rasanen@jyu.fi

Keywords: climate change, genetic architecture, growth rate, *Neurospora crassa*, quantitative trait, temperature fluctuation, thermal tolerance

ABSTRACT

Constant temperatures and fluctuations of varying frequency affect fitness differently, which
has led to suggestions of distinct genetic architectures and adaptation strategies between
constant and fluctuating thermal environments. However, very little is known about the
possible fitness trade-offs and genetic constraints underlying thermal adaptation and how
they affect species' ability to confront climatic changes. We addressed this gap in knowledge
by integrating quantitative genetics and genome-wide association mapping in the
filamentous fungus *Neurospora crassa*. Growth rates were measured for 434 strains under fast
and slow frequency fluctuations, at high and low temperature range with respect to the
species' tolerance, and at constant mean and extreme temperatures of these ranges. We found
strong genetic correlations between fast and slow frequency fluctuations, and between
fluctuations and their mean temperatures, but not with the highest extreme temperature.
Positive correlations were supported by high heritability values, pointing that in *N. crassa*
there are no significant trade-offs or genetic constraints in adaptation when variance in
temperature increases. Altogether, our results indicated clearly polygenic basis of thermal

tolerance, with most of the variation in overall performance (83 %), and clearly less in hot-cold trade-off (8 %), or heat stress tolerance (4 %). Interestingly, GWAS discovered many SNPs associated with growth rate only at constant temperatures or at fast and slow fluctuations at high and low thermal range. However, the cellular functions of the associated genes were overlapping, and no opposite allelic effects were found between treatments. Hence, large-effect loci indicated no trade-offs, but a shared physiology across temperatures, probably owing to the general stress response or individual's overall fitness.

1 INTRODUCTION

The global increases in mean and variance of temperature force species to tolerate more frequent climatic fluctuations such as severe heat waves (Meehl et al. 2000; IPCC 2018). Based on species performance projections and empirical data, adaptation to temperature fluctuations should be harder than tolerating just the rise in mean temperature (Vasseur et al. 2014; Colinet et al. 2015). Consequently, a better understanding of how species can cope with increased thermal variation is crucial for predicting their survival under climate change (Schulte et al. 2011; Kristensen et al. 2020). According to traditional hypotheses, the greater variability in temperature should lead to evolution of generalists with enhanced performance across a wider range of temperatures (Levins 1968; Lynch and Gabriel 1987; Gilchrist 1995). However, the ability to spend more time closer to critical thermal limits should evolve at the expense of overall fitness, also known as a specialist-generalist trade-off (Angilletta 2009). This hypothesis suggests that the only way to change the breadth of the thermal performance curves (TPCs) is by lowered elevation at optimal conditions, meaning that populations cannot be concurrently well-adapted to both mean and variance of temperature (Lynch and Gabriel 1987; Gilchrist 1995).

More recently, studies have acknowledged that adaptations to temperature fluctuations might not be predictable from TPCs measured across constant temperatures and shown evidence that distinct genetic architectures regulate adaptation in fluctuating versus constant conditions (Schulte et al. 2011; Ketola and Saarinen 2015; Sinclair et al. 2016; Sørensen et al. 2016; Ketola and Kristensen 2017). In addition, theoretical studies suggest that there are

evolutionary constraints between the types of fluctuations, for example, that particular adaptive mechanisms are operating in environments that fluctuate at different time scales (DeWitt and Langerhans 2004; Botero et al. 2015). If these hypotheses hold true, distinct genetic architectures and constraints are associated with adaptation to constant and fluctuating environments of varying frequency (Ketola and Kristensen 2017), which could limit evolutionary adaptation and heavily affect species survival when variation in temperature increases (Sgrò and Hoffmann 2004; Björklund et al. 2009).

The potential trade-offs in thermal adaptation are often investigated by the principal components of genetic variation in TPCs and quantitative genetic analysis of covariation (Kingsolver et al. 2001; Izem and Kingsolver 2005). Specifically, evolutionary constraints are defined as directions where selection cannot increase fitness due to low amounts of heritable genetic variation or negative genetic correlations across alternative environmental states (Gaydos et al. 2013). The fluctuations of varying frequency affect fitness differently and are expected to have their characteristic changes in TPCs (Schulte et al. 2011; Colinet et al. 2015; Schaum et al. 2022). Generally, three modes of variation are observed in the shape and position of TPCs; the elevation in overall performance, horizontal shift in optimum between hot and cold temperatures, and in breadth of the TPC (Izem and Kingsolver 2005; Angilletta 2009). Under fast fluctuations, selection should favour elevated performance and more robust tolerance across temperatures, whereas slower fluctuations should select for more fixed strategies, reflected as a narrower TPCs, or shift in optimum (DeWitt and Langerhans 2004; Ketola et al. 2014; Schaum et al. 2022). However, fast fluctuations could also select for reversible phenotypic plasticity aided by the fast production of heat shock proteins or the maintenance of regulatory enzymes, which could trade-off, due to their energetic requirements with tolerance to constant environments (Kristensen et al. 2000; Sørensen et al. 2003).

The traditional quantitative genetics studies have long focused on estimating statistical averages over a multitude of loci with small phenotypic effects (Fisher 1919). More recently, the next generation sequencing methods have made it possible to use whole genome sequence data and investigate the genetic architecture of polygenic traits on the level of individual loci (Fu et al. 2013). For example, genome-wide association studies (GWAS) link genotypes to phenotypes by mapping mutational markers such as single nucleotide

polymorphisms (SNPs). Using quantitative genetics and the latest molecular genetics methods together can give us more complete picture of the genetic variation in thermal tolerance and its evolutionary potential (Cortés et al. 2020; Buckley and Kingsolver 2021), but so far only few studies have adopted this approach (Gerken et al. 2015; Latimer et al. 2015; Duun Rohde et al. 2016; Rolandi et al. 2018; Lecheta et al. 2020). Furthermore, most of the quantitative genetics studies and studies using genome-wide sequence data have focused on evaluating the effects of constant or increasing heat (Riehle et al. 2001; Knies et al. 2006; Mitchell and Hoffmann 2010; Khan et al. 2022). Knowledge from differently fluctuating temperatures is sparse and studies have mostly used experimental evolution and resequencing (Tobler et al. 2014; Deatherage et al. 2017; Schaum et al. 2018; Lambros et al. 2021), whereas only one study has been made on natural genetic variation (Sørensen et al. 2016). As climate change keeps accelerating, there is a pressing need for experiments evaluating the predicted fitness trade-offs and constraints in thermal adaptation under various fluctuations (Chevin and Hoffmann 2017; Buckley and Kingsolver 2021).

To address this gap in knowledge, we investigated if distinct genetic architectures are found at constant and fluctuating temperatures of varying frequency, and if some genetic constraints might limit thermal adaptation. We used quantitative genetics and GWAS in the filamentous fungus *Neurospora crassa* to quantify how different components of genetic variation contribute to the total variation. The thermal sensitivity to growth rate was measured for a previously developed mapping population (Moghadam et al. 2020; Räsänen et al. 2024) under fast and slow frequency fluctuations, at high and low temperature range, and at constant mean and extreme temperatures of these ranges. In our analyses we used a multivariate approach in which growth rates at each temperature are treated as discrete and potentially genetically correlated traits (Kingsolver et al. 2004). We examined the following questions: (1) How much there is genetic variation in growth rate at constant and fluctuating temperatures and how strongly is this variation correlated between contrasting thermal environments? (2) To what extent the phenotypic variation in growth rate can be related to broader characteristics of the TPCs? (3) Are there genes associated with growth rate at constant and fluctuating temperatures and are these genes same between contrasting thermal environments? Our study is among the first ones to investigate the genetic basis of thermal tolerance at temperatures that fluctuate with different frequencies.

2 MATERIALS AND METHODS

2.1 *Neurospora crassa* mapping population

N. crassa is a clonally and sexually reproducing species that is well-suited for experiments studying genetic differences across environments (Fisher and Lang 2016). In our experiment we used 118 natural strains obtained from the Fungal Genetics Stock Center (Manhattan, Kansas, USA) (McCluskey et al. 2010), originally sampled from natural populations in southeastern USA (Louisiana), Central America and the Caribbean Basin (Ellison et al. 2011; Palma-Guerrero et al. 2013). Previously, some of the natural strains were crossed to form 316 additional offspring strains (Moghadam et al. 2020). In total our nested association mapping population had 434 strains genotyped for 1 473 869 single nucleotide polymorphisms (SNPs) (Räsänen et al. 2024).

2.2 Phenotyping by growth rate

We measured the mycelial growth rate as a proxy for thermal tolerance (Pringle and Taylor 2002). Before starting experiments, *N. crassa* cultures were maintained on Vogel's medium N (Metzenberg 2003) in slants at room temperature. The linear growth rates were measured for each strain by standard tube method using agar filled serological pipet tips (Ryan et al. 1943; White and Woodward 1995). A more detailed description of this method can be found from Moghadam et al. (2020) and Kronholm et al. (2016). We replicated each strain 3 times and put them to grow in 10 thermal treatments (MTM-313 Plant Growth Chamber, HiPoint Corp., Taiwan). We had two fluctuating regimes, low 25–35 °C and high 32–42 °C, and fast and slow fluctuations with frequencies of 120 min and 480 min respectively. Thus, fast fluctuations were 4 times faster, having a step duration of 15 min at each temperature compared to 60 min steps at slow fluctuations. At lower range the steps were 25–27.5–30–32.5–35–32.5–30–27.5–25 °C, and at higher range 32–34.5–37–39.5–42–39.5–37–34.5–32 °C. The fast fluctuations were estimated to span within the generation time of *N. crassa*, whereas the slow fluctuations

occurred between generations (Kronholm and Ketola 2018). The growth rates were measured also at constant mean and extreme temperatures of these ranges. The measurements at constant lower range temperatures 25 °C, 30 °C, and optimal 35 °C were collated from previously published data (Moghadam et al. 2020). At higher range the growth rates were measured at constant 32 °C, 37 °C, and extreme 42 °C which is known to be very stressful for *N.* crassa (Mohsenzadeh et al. 1998). Altogether we had 12 976 growth assays to use in our analyses.

2.3 Quantitative genetics

The statistical analysis for quantitative genetics parameters was conducted with Bayesian multilevel model using Hamiltonian Monte Carlo algorithm. We used R version 4.3.1 (R Core Team 2022) and R package 'brms' version 2.19.0 (Bürkner 2018) that fits Bayesian models by 'Stan' language. The phenotypic data including all growth assays was used in a multivariate model

$$y_i \sim MVN(\mu_i, E)$$
$$\mu_i = \alpha + \alpha_{g[i]}$$
$$\alpha_{g[i]} \sim MVN(0, G)$$
$$G = S_G R_G S_G$$
$$E = S_E R_E S_E$$

where the growth at each temperature treatment was explained by the strain genotype as a random variable and modeled as potentially correlated with growth at other temperatures. In this model α was the vector of intercepts, $a_{g[i]}$ was the vector of genotypic effects, S_G and S_E were 10 × 10 diagonal matrices with genetic or environmental standard deviations on the diagonal, and R_G and R_E were matrices for genetic and environmental correlations, respectively. We had weakly informative priors as we used the half location-scale version of Student's t-distribution with 3 degrees of freedom and 2.5 as the scale parameter. The prior for the intercept effects was

$$\alpha \sim \begin{pmatrix} hT\ (3, 3.1, 2.5) \\ hT\ (3, 3.9, 2.5) \\ hT\ (3, 4.2, 2.5) \\ hT\ (3, 4.3, 2.5) \\ hT\ (3, 3.8, 2.5) \\ hT\ (3, 0.7, 2.5) \\ hT\ (3, 3.6, 2.5) \\ hT\ (3, 3.8, 2.5) \\ hT\ (3, 2.8, 2.5) \\ hT\ (3, 3.2, 2.5) \end{pmatrix}$$

for growth rates from 25 °C to 42 °C, and for fluctuating temperatures in order 25–35 °C fast, 25–35 °C slow, 32–42 °C fast, and 32–42 °C slow. The prior for each standard deviation in the model was $\sigma \sim hT(3, 0, 2.5)$, and we used an lkj prior (McElreath 2015) for the correlation matrices: R_E, $R_G \sim LKJ(1)$. For MCMC estimation 2 chains were run with a warmup period of 1000 iterations, followed by 6000 iterations of sampling, with thinning set to 2. MCMC convergence was monitored by trace plots and $\hat{R}$ values.

From the model we obtained a G-matrix with estimates of genetic variance and covariance components for growth rate. The environmental variances were estimated as the variances between the clonal replicates of the same strain. We considered the posterior mean estimates to be statistically significant if their 95 % highest posterior density intervals (HPDI) did not overlap with 0. We extracted the posterior samples from the model and used these to calculate broad-sense heritabilities, genetic covariances and genetic correlations for growth rate in different thermal environments. We also quantified the coefficients of variation for both, the genetic and environmental components by standardizing the scales by the mean phenotype. The R package 'ggplot2' (Wickham 2016) was used for plotting.

2.4 Principal components analysis

We used a Principal Components Analysis (PCA) to characterize the main modes of variation in TPCs, and the amount of phenotypic variation that was attributed to each principal component (PC) (Izem and Kingsolver 2005). We implemented PCA with built-in R function 'princomp' for the growth rates averaged per temperature for each strain. The predictive values of the first 3 principal components (from here on PC1, PC2 and PC3) were saved and

used in GWAS so that the results could be compared with the associations of specific

temperature treatments. We counted also the Bayesian mean estimates and HPDIs for the

PCs. To form the posterior distributions, we sampled the intercepts extracted from the

multivariate model for each genotype and at each temperature 5000 times. Then we used the

PCA to calculate the proportions of variance explained by the first 6 PCs and the loadings of

PC1, PC2 and PC3 at each temperature.

2.5 Genome-wide association mapping

GWAS was used to study the genotype–phenotype associations and to answer whether the

same or different genes contribute to the thermal tolerance at fluctuating and constant

temperatures. We investigated also which SNPs are associated with PC1, PC2 and PC3

explaining the variation in average growth rate. We used nested association mapping that

combines information from recent and past recombination (Yu et al. 2008). We conducted the

GWAS by the genome association and prediction integrated tool (GAPIT) Version 3 R-

package (Wang and Zhang 2021). As an input data we had each strains SNP sequence in

HapMap format, and the strains average growth rate per temperature treatment. In GWAS,

GAPIT 3 identifies the quantitative trait loci via statistical association between the

phenotypes and single-nucleotide markers. It also corrects for the population structure and

kinship in these analyses and so sets control for the false positives. The basic mixed model

was

$$y = X\beta + Zu + e$$

where y is a vector of phenotypic observations, β is a vector of fixed effects including effects

of the genetic marker and population structure, u is a vector of random genetic effects for

each individual, X and Z are design matrices, and e is a vector of residual effects. The variance

matrix for u is σ_a^2 K where σ_a^2 is the additive variance and K is the kinship matrix. The

variance matrix for e is σ_e^2 I, where σ_e^2 is the residual variance and I is the identity matrix. For

the population structure correction, a principal component analysis was first done on the

SNPs, and the components were incorporated as fixed factors in the model (Price et al. 2006).

We chose the first 4 components for the analysis since they explained most of the variance in SNPs.

We run GWAS with Bayesian-information and linkage-disequilibrium iteratively nested keyway method (BLINK) (Huang et al. 2019). BLINK is a multi-locus test (Segura et al. 2012) and the most statistically powerful method in GAPIT 3 (Wang and Zhang 2021). In BLINK, the SNP markers are removed iteratively if they are in linkage disequilibrium with the most significantly associated reference SNP. The procedure of selecting a reference SNP among the remaining markers is repeated until no markers can be removed. We run the BLINK separately for the average growth rates in all temperature treatments and for the phenotypic variation attributed to PC1, PC2 and PC3 describing the TPC. The significance threshold for associations was set to 0.01 after a Bonferroni correction, which corresponded to a P-value of 6.78×10^{-9}. The signs of the allelic effects were with respect to the minor allele, i.e., a positive allelic effect indicated that the individuals carrying the minor allele had faster growth rate.

We searched the statistically significant SNPs from Ensemble Fungi release 57 (Yates et al. 2022) and identified the candidate genes affecting the growth rate. We used *Neurospora crassa* (NC12) whole genome alignment and annotations (Galagan et al. 2003). We also processed the SNP variants by the Variant Effect Predictor web tool (McLaren et al. 2016) to determine the possible effects on transcript and protein sequences. In the case of intergenic variants, we reported annotations for a gene that had the shortest distance from the marker to the transcript. We further searched for the functional protein annotations of the candidate genes from the UniProt (The UniProt Consortium 2023) and the FungiDB databases (Basenko et al. 2018).

3 RESULTS

3.1 Growth rates at different temperatures

We observed large variation in growth rates between temperature treatments (0.01–6.05mm/h) and there were clear differences in how well strains grow in fluctuating

temperatures (Fig. 1A). The genotype had a statistically significant effect on growth rate at each temperature treatment (the model estimates are reported as a mean with the 95 % HPD interval given in parentheses). On average, the growth rates were faster at lower temperature range and at slow frequency fluctuations (Fig. 1B). At constant temperatures, growth rates increased from 25 °C (3.04 mm/h, 3.00–3.08) to 30 °C (3.83 mm/h, 3.77–3.88), and at 32 °C (4.07 mm/h, 4.01–4.12). The growth rates were fastest at the optimal temperature 35 °C (4.16 mm/h, 4.10–4.23) and slowed down at hotter 37 °C (3.73 mm/h, 3.66–3.79) and at 42 °C (0.68 mm/h, 0.65–0.72). At lower 25–35 °C range, the mean growth rates were very similar, 3.53 mm/h (3.48–3.58) at fast fluctuations and 3.65 mm/h (3.61–3.70) at slow fluctuations. At higher 32–42 °C range, the mean growth rate was faster at slow frequency fluctuations 3.10 mm/h (3.05–3.15) than at fast fluctuations 2.76 mm/h (2.72–2.81).

Figure 1. (A) The mean growth rates for each *N. crassa* genotype (N = 434) and (B) the posterior mean estimates of growth rate when the growth in fluctuating environments was compared at high and low temperature range and at fast and slow frequency. In Fig. B the estimates for the constant temperatures within a temperature range are indicated by colors (red = high extreme temperature, green = mean temperature, and blue = low extreme temperature, reflecting 25 °C, 30 °C, and 35 °C at lower range, and 32 °C, 37 °C, and 42 °C at higher range). The 95 % HPD intervals are marked with dotted lines for constant temperatures. In fluctuating temperatures, the points are hiding the small 95 % HPD interval bars.

3.2 Quantitative genetics

The multivariate model was used to estimate the genetic variance and covariance
components for growth rate at different temperatures (Table 1). In all thermal treatments,
there was genetic variation for growth rate (posterior means 0.14–0.46) (Table 1), and this
was observable also from high heritability values (posterior means 0.64–0.92) (Fig. 2A). In
general, we found strong positive genetic correlations (posterior means 0.54–0.99) and
genetic covariances (posterior means 0.10–0.35) between treatments (Table 1). However, the
correlations with constant 42 °C were markedly lower (posterior means 0.14–0.39), which
was also true for the covariances (posterior means 0.02–0.07). There were most genetic and
environmental variance at constant 42 °C when scaled as the coefficients of variation (Fig.
2B).

Table 1. Genetic variances, covariances, correlations, and environmental variances for growth rate at constant and fluctuating temperatures estimated

from the multivariate model. In table: upper triangle genetic correlations, lower triangle genetic covariances, diagonal genetic variances (in bold), and

last column environmental variances. Estimates are posterior means with 95 % HPD intervals shown in parentheses.

(°C)	25	30	32	35	37	42	25–35 Fast	25–35 Slow	32–42 Fast	32–42 Slow	VarE
25	**0.15 (0.14–0.18)**	0.95 (0.93–0.96)	0.79 (0.75–0.83)	0.80 (0.76–0.84)	0.54 (0.46–0.61)	0.14 (0.04–0.23)	0.84 (0.81–0.87)	0.84 (0.81–0.88)	0.57 (0.49–0.64)	0.63 (0.57–0.69)	0.02 (0.02–0.02)
30	0.20 (0.18–0.23)	**0.29 (0.26–0.33)**	0.90 (0.87–0.92)	0.89 (0.86–0.91)	0.63 (0.56–0.68)	0.19 (0.09–0.28)	0.92 (0.89–0.94)	0.92 (0.89–0.94)	0.65 (0.59–0.71)	0.70 (0.65–0.75)	0.04 (0.03–0.04)
32	0.18 (0.16–0.21)	0.28 (0.24–0.32)	**0.34 (0.29–0.38)**	0.87 (0.84–0.90)	0.77 (0.72–0.81)	0.22 (0.13–0.31)	0.98 (0.97–0.99)	0.98 (0.97–0.99)	0.79 (0.74–0.83)	0.83 (0.79–0.86)	0.08 (0.07–0.09)
35	0.21 (0.18–0.25)	0.32 (0.28–0.37)	0.34 (0.30–0.39)	**0.46 (0.40–0.52)**	0.82 (0.78–0.85)	0.27 (0.17–0.36)	0.89 (0.86–0.92)	0.88 (0.85–0.91)	0.82 (0.78–0.86)	0.87 (0.84–0.90)	0.06 (0.06–0.07)
37	0.14 (0.11–0.16)	0.22 (0.18–0.26)	0.29 (0.24–0.33)	0.35 (0.31–0.41)	**0.41 (0.36–0.47)**	0.29 (0.20–0.38)	0.76 (0.71–0.80)	0.78 (0.74–0.82)	0.86 (0.83–0.90)	0.91 (0.89–0.93)	0.04 (0.03–0.04)
42	0.02 (0.01–0.03)	0.04 (0.02–0.06)	0.05 (0.03–0.07)	0.07 (0.04–0.09)	0.07 (0.05–0.09)	**0.14 (0.12–0.16)**	0.20 (0.10–0.29)	0.20 (0.11–0.29)	0.36 (0.27–0.45)	0.39 (0.30–0.48)	0.06 (0.06–0.07)
25–35 Fast	0.16 (0.14–0.18)	0.24 (0.20–0.27)	0.27 (0.23–0.31)	0.28 (0.25–0.33)	0.23 (0.20–0.27)	0.03 (0.02–0.05)	**0.22 (0.19–0.25)**	0.99 (0.98–1.00)	0.77 (0.72–0.82)	0.82 (0.78–0.85)	0.06 (0.05–0.07)
25–35 Slow	0.16 (0.14–0.18)	0.24 (0.21–0.27)	0.27 (0.24–0.31)	0.29 (0.25–0.33)	0.24 (0.21–0.28)	0.04 (0.02–0.05)	0.22 (0.19–0.25)	**0.23 (0.20–0.26)**	0.76 (0.71–0.81)	0.82 (0.78–0.85)	0.06 (0.05–0.07)
32–42 Fast	0.10 (0.08–0.12)	0.16 (0.13–0.18)	0.20 (0.17–0.23)	0.24 (0.21–0.28)	0.24 (0.21–0.28)	0.06 (0.04–0.08)	0.16 (0.14–0.19)	0.16 (0.14–0.19)	**0.19 (0.17–0.22)**	0.96 (0.94–0.97)	0.11 (0.10–0.12)
32–42 Slow	0.13 (0.11–0.15)	0.19 (0.16–0.23)	0.24 (0.21–0.28)	0.30 (0.26–0.34)	0.30 (0.26–0.34)	0.07 (0.05–0.09)	0.20 (0.17–0.23)	0.20 (0.17–0.23)	0.21 (0.19–0.25)	**0.26 (0.22–0.29)**	0.05 (0.05–0.06)

Figure 2. (A) Heritabilities of growth rate and (B) coefficients of genetic and environmental variation at each temperature treatment. Estimates are posterior means with 95 % HPD intervals (some points are larger than the error bars).

3.3 Principal component analysis

PCA found 3 principal components that explained 95 % of the total genetic variance in average growth rate (Fig. 3A). The loadings of these 3 principal components at different thermal treatments were easily interpreted as TPC attributes (Fig. 3B). PC1 explained 83 % of the total variation and had the largest loading (0.44) at the optimal temperature 35 °C. As all loadings were positive, PC1 was interpreted as the individual's overall fitness in growth i.e. the elevation of the TPC (Fig. 3C). PC2 explained 8 % of the variation and was interpreted as the horizontal shift in TPC optimum towards colder temperatures (Fig. 3D). Interestingly, the constant 37 °C and 42 °C, and 32–42 °C fluctuations all had negative loadings in PC2. The proportion of variance for PC3 was only 4 % and the loadings were interpreted as the change in the shape of the TPC (Fig. 3E). There was a strong positive loading (0.90) for PC3 at constant 42 °C, indicating increased growth rate at the high thermal extreme.

Figure 3. (A) Proportions of variance explained by the different principal components and (B) the loadings of PC1, PC2 and PC3 for each temperature treatment. Shown are the posterior means and 95 % HPD intervals as error bars. At lower line are the schematic pictures for the changes in TPC

317   attributes: (C) the elevation, (D) the hot and cold shift in optimum, and (E) the tolerance of high

318   thermal extreme.

3.4 Genome-wide association mapping

3.4.1 Associations and allelic effects

GWAS found 25 associations with growth rate at different temperatures, which were related

to 21 SNPs, and located in 19 genes or intergenic loci (Table 2). At constant temperatures

optimal or lower, significant SNPs were found from chromosomes 3, 5 and 6 (Fig. 4, Fig. 5).

At constant temperatures above the optimal, the associated SNPs were in chromosomes 1

and 6 (Fig. 5). At fluctuations, associations were in chromosomes 1, 3 and 6 (Fig. 4, Fig. 5).

The strongest associations were found at 42 °C in chromosome 1: 7 807 362 (P = 4.66 × 10^{-32}),

at slow 25–35 °C fluctuations in 1: 7 378 135 (P = 2.47 × 10^{-21}), and at fast 25–35 °C fluctuations

in 3: 2 881 603 (P = 2.77 × 10^{-19}) and in 6: 3 600 909 (P = 1.19 × 10^{-19}). From significantly

associated SNPs, 8 were detected only at fluctuating temperatures, 12 only at constant

temperatures, and 2 at both, fluctuating and constant temperatures. Altogether there were 18

unique SNPs associated only at one temperature treatment and 3 shared SNPs that were

associated at more than one temperature treatment (Table 2). Only unique SNPs were found

at the lowest extreme temperature 25 °C, above optimum at 37 °C and 42 °C, and at higher

32–42 °C range. The shared SNPs 3: 2 300 888, 3: 2 881 603, and 6: 3 600 716 were found at

favorable 30 °C, 32 °C, 35 °C, and fluctuating 25–35 °C temperatures. Interestingly, there were

no shared SNPs between the fast and slow fluctuations.

Most of the SNPs had a negative effect on growth rate when examined with respect to

the minor allele (Table 2). This was also true for the shared SNPs, for which the direction of

the allelic effects on growth rate were negative regardless of the temperature. The only

positive effect on growth rate (0.18 mm/h) was found at 37 °C for the SNP 1: 7 182 169. In

general, the minor alleles in our natural populations of *N. crassa* are known to be mildly

deleterious and occur at low frequencies (Räsänen et al. 2024). The magnitude of the allelic

effects was, on average, similar for the unique and the shared SNPs. The greatest allelic

effects were found at 37 °C (–0.69 mm/h), at 35 °C (–0.48 mm/h), and at fast 25–35 °C

fluctuations (–0.42 mm/h).

3.4.2 Allelic variants

Most of the allelic variants were synonymous or in non-coding area up- or downstream from

the gene (Table 2). GWAS found also 4 missense variants changing the amino acid and

affecting the protein structure. These were in chromosome 6: 3 669 489 at 25 °C changing

serine to proline, in 3: 2 179 382 at 35 °C changing glycine to serine, in 6: 430 408 at 42 °C

changing serine to phenylalanine, and in 3: 1 402 371 at fast 32–42 °C fluctuations, changing

serine to glycine. All missense variants were unique associations, whereas shared

associations were synonymous or in non-coding area. However, there was no obvious

connection between the types of allelic variants and the associations being found at constant

or fluctuating temperatures.

3.4.3 Gene annotations and protein functions

The associations found at constant lower range temperatures 25 °C, 30 °C, and at optimal 35

°C have been previously annotated in Räsänen et al. (2024). At constant 25 °C were associated

the *fpo-1* (NCU00236) gene which encodes flavoprotein oxygenase, a gene (NCU01043)

coding an uncharacterized protein, and a missense mutation in the gene (NCU17138)

encoding a haloacid dehalogenase (HAD) like protein (Table 2). At constant 30 °C and 32 °C,

and at fast fluctuations of both higher and lower range, 2 SNPs were associated in the

regulatory area of the gene (NCU00170) encoding DNAJ domain-containing protein, which

belongs to the HSP40 heat shock proteins. At constant 30 °C and 35 °C, and at fast 25–35 °C

fluctuations there were 2 synonymous mutations in the *ncw-1* (NCU05137) gene encoding a

highly conserved non-anchored cell wall protein 1. At constant 32 °C were associated a gene

(NCU05084) encoding an uncharacterized protein, and a gene (NCU03622) encoding a

UBA/TS-N domain-containing protein. This protein is known to function in endocytosis and

to have temperature-dependent regulation of expression in fungus *Aspergillus flavus*

(Georgianna et al. 2008). At constant 32 °C and at slow 25–35 °C fluctuations was associated

a gene (NCU00017) encoding a Zn(2)-C6 fungal-type domain-containing protein, that is a

transcription regulator. At optimal 35 °C was found a unique missense SNP in the *rbg-37*

(NCU06278) gene producing a variant in elongation factor 2.

At constant 37 °C was found a synonymous association in the *gpr-1* (NCU00786) gene

encoding a G-protein coupled receptor. The G-protein receptors work on the cell surface and

affect the gene regulation through signaling pathways. The *grp-1* gene has been found to be

important in the sexual reproduction of *N. crassa* (Krystofova and Borkovich 2006), and the

deletion mutants of another G-protein coupled receptor gene (*grp-4*), have been shown to

grow well at high temperatures up to 42 °C (Li and Borkovich 2006). At constant 37 °C was

also associated the *stk-32* (NCU04755) gene encoding a protein kinase domain-containing

protein ppk32 (large allelic effect of –0.69 mm/h). The protein kinases are universally

important coenzymes and enzyme regulators that bind to ATP in phosphorylation and work

in signal transduction. In the previous GWAS, the association with the *stk-32* gene was found

for the maximum growth rate at the optimal temperature (Räsänen et al. 2024).

At extreme 42 °C we found an association with the gene (NCU00621) encoding a TORC1

growth control complex subunit Kog1, and a missense mutation in the hypothetical protein

(NCU04701) that had a high minor allele frequency in the mapping population (0.47) (Table

2). The target of rapamycin complex 1 (TORC1) is a highly conserved serine/threonine kinase

that regulates fungal mycelial growth and protein synthesis in response to nutrient and

energy signaling pathways (Wang et al. 2023). In yeast cells, TORC1 has been shown to

regulate growth also by sensing environmental stressors such as high temperature (Urban et

al. 2007). In *N. crassa*, several studies have linked the TOR signaling pathway to cell cycle and

circadian rhythmicity, which are also affected by the changes in temperature (Lakin-Thomas

2019).

At fast 25–35 °C fluctuations was found an unique association in the gene (NCU06484)

encoding a AAA+ ATPase domain-containing protein (Table 2), that positively regulates

transcription. At slow 25–35 °C, the found SNPs were in the gene (NCU00735) encoding a

Ribosome biogenesis protein SLX9, and upstream from the gene (NCU08213) encoding a

hypothetical protein. At fast 32–42 °C fluctuations was associated a missense mutation in the

gene (NCU06434) encoding a hypothetical protein. At slow 32–42 °C fluctuations, the first

association was in the gene (NCU00669) producing an oligosaccharyl transferase (OST)

subunit. OST is a membrane protein complex, which catalyzes a central step in the
glycosylation pathway. Glycosylation affects the folding and thermodynamic stability of the
proteins. The N-glycosylation pathway has been found to be highly conserved across species
of filamentous fungi (Deshpande et al. 2008). The second association was upstream from the
fkr-3 (NCU04371) gene which produces a peptidyl-prolyl cis-trans isomerase. The *fkr-3* gene
accelerates the protein folding by catalyzing the cis-trans isomerization activity. In *N. crassa*,
the *fkr-3* deletion mutants have a temperature-sensitive phenotype (Pinto et al. 2008), and
this gene is upregulated during heat stress and recovery also in other fungi (Zou et al. 2018).

Table 2. SNPs that were significantly associated with growth rate at constant and fluctuating temperatures, their positions in genome (bp), allelic effects

(mm/h), variant types and annotated proteins. The signs of the allelic effects are with respect to the minor allele and annotations for the intergenic

variants are reported for the closest gene.

Temperature (°C)	Chromosome	Position (bp)	P-value	Allelic effect (mm/h)	Gene	Symbol	Annotation	Variant
25	3	3 119 653	1.31×10^{-10}	−0.20	NCU00236	*fpo-1*	flavoprotein oxygenase	synonymous
25	5	2 747 660	2.22×10^{-10}	−0.13	NCU01043		hypothetical protein	3'UTR
25	6	3 669 489	3.27×10^{-12}	−0.22	NCU17138		haloacid dehalogenase like protein	missense
30	3	2 881 603	2.69×10^{-11}	−0.26	NCU00170		DNAJ domain-containing protein	upstream
30	6	3 600 716	1.18×10^{-9}	−0.23	NCU05137	*ncw-1*	non-anchored cell wall protein 1	synonymous
32	3	2 300 888	7.10×10^{-13}	−0.32	NCU00017		Zn(2)-C6 fungal-type domain-containing protein	upstream
32	3	2 881 603	6.09×10^{-10}	−0.27	NCU00170		DNAJ domain-containing protein	upstream
32	5	1 883 866	5.55×10^{-15}	−0.22	NCU03622		UBA/TS-N domain-containing protein	3'UTR
32	6	3 797 277	1.99×10^{-10}	−0.29	NCU05084		hypothetical protein	3'UTR
35	3	2 179 382	9.39×10^{-17}	−0.48	NCU06278	*rbg-37*	elongation factor 2, variant 2	missense
35	6	3 600 716	1.32×10^{-10}	−0.26	NCU05137	*ncw-1*	non-anchored cell wall protein 1	synonymous
37	1	7 182 169	5.84×10^{-11}	0.18	NCU00786	*gpr-1*	G-protein coupled receptor	synonymous
37	6	686 146	1.25×10^{-13}	−0.69	NCU04755	*stk-32*	protein kinase domain-containing protein ppk32	5'UTR
42	1	7 807 362	4.66×10^{-32}	−0.38	NCU00621		TORC1 growth control complex subunit Kog1	5'UTR
42	6	430 408	6.38×10^{-13}	−0.11	NCU04701		hypothetical protein	missense
25–35 Fast	3	1 573 941	6.65×10^{-12}	−0.22	NCU06484		AAA+ ATPase domain-containing protein	3'UTR
25–35 Fast	3	2 881 603	2.77×10^{-19}	−0.40	NCU00170		DNAJ domain-containing protein	upstream
25–35 Fast	6	3 600 909	1.19×10^{-19}	−0.27	NCU05137	*ncw-1*	non-anchored cell wall protein 1	synonymous
25–35 Slow	1	7 378 135	2.47×10^{-21}	−0.24	NCU00735		Ribosome biogenesis protein SLX9	synonymous
25–35 Slow	3	2 300 888	2.26×10^{-16}	−0.29	NCU00017		Zn(2)-C6 fungal-type domain-containing protein	upstream
25–35 Slow	3	4 656 366	3.20×10^{-12}	−0.21	NCU08213		hypothetical protein	upstream
32–42 Fast	3	1 402 371	3.39×10^{-9}	−0.42	NCU06434		hypothetical protein	missense

32–42 Fast	3	2 881 312	5.95×10^{-13}	−0.22	NCU00170		DNAJ domain-containing protein	upstream
32–42 Slow	1	7 625 265	2.56×10^{-13}	−0.21	NCU00669		oligosaccharyl transferase subunit	3'UTR
32–42 Slow	6	3 599 935	3.46×10^{-12}	−0.23	NCU04371	*fkr-3*	peptidyl-prolyl cis-trans isomerase fkr-3	upstream

Figure 4. Manhattan plots for the growth rates at different temperatures. The solid horizontal lines denote a P-value threshold of 0.01 after a Bonferroni correction. The dotted vertical lines highlight the candidate SNPs that were significant in at least one of the temperature treatments within the lower range 25–35°C.

Figure 5. Manhattan plots for the growth rates at different temperatures. The solid horizontal lines denote a P-value threshold of 0.01 after a Bonferroni correction. The dotted vertical lines highlight the candidate SNPs that were significant in at least one of the temperature treatments within the higher range 32–42°C.

3.4.4 GWAS for the principal components

GWAS found 8 unique genes associated with the PCs characterizing the main modes of variation in TPCs (Table 3). No significant SNPs were found for the PC1 interpreted as a TPC elevation, even though it explained most of the phenotypic variation (Fig. 3A). However, the PC1 had a nearly significant association (P = 2.12×10^{-8}) with the gene (NCU00170) encoding a DNAJ domain-containing protein, which is likely to be a true association based on its occurrence in the GWAS for growth rates at different temperatures (Table 2). The PC2 was interpreted as a shift in TPC optimum towards colder temperatures and it had 3 associations located in chromosomes 1 and 7 (Table 3, Fig. 6). The first association was upstream from the gene (NCU03028) encoding a deubiquitination-protection protein dph1, which works in protein degradation and localization to vacuoles, and has been found to affect mycelial growth under starvation stress (Zhou et al. 2016). The second association was in the gene (NCU06948) encoding an EF-hand domain-containing protein and had a very strong allelic effect (–0.99 mm/h). The EF-hand proteins affect enzymatic activity and cell division through microtubule organization and have been found to be involved in vesicular trafficking and fungal hyphal growth (Kundu et al. 2022). The third association was a missense variant having a large allelic effect (–0.85 mm/h) and changing alanine to serine in the *pks-4* (NCU08399) gene. The *pks-4* encodes a polyketide synthase 4 variant which functions in the formation of fatty acids. The polyketide synthases have been linked to the sexual reproduction and synthesis of secondary metabolites against predation, which both are important to the survival of *N. crassa* (Zhao et al. 2017).

The PC3 was interpreted as the shape change towards higher temperatures and had 5 associations in chromosomes 1, 2, 3, and 5 (Table 3, Fig. 6). The strongest association (P = 6.30×10^{-20}) was in the gene (NCU00621) encoding a TORC1 growth control complex subunit Kog1, that was also associated at 42 °C, but with the PC3 there was a moderate positive allelic effect (Table 2, Table 3). There was also another association with positive allelic effect upstream from the gene (NCU03448) with unknown function (Table 3). In addition, 3 genes having negative allelic effects and encoding hypothetical proteins were associated with the PC3. One of them (NCU08496) was a missense variant changing alanine to serine and had a high minor allele frequency (0.45) in the mapping population. Other mutations were

synonymous (NCU10002) and intergenic upstream (NCU06730) variants. In the previous

GWAS for the growth rate of *N. crassa*, different genes were found to be important at optimal

temperature, and at hot and cold extremes of the TPC (Räsänen et al. 2024).

Table 3. SNPs that were significantly associated with the predictive values of the first 3 principal components, their positions in genome (bp), allelic

effects on phenotypic variance, allelic variant types and annotated proteins. The signs of the allelic effects are with respect to the minor allele and the

annotations for intergenic variants are presented for the closest gene.

PC	Chromosome	Position (bp)	P-value	Allelic effect	Gene	Symbol	Annotation	Variant
2	1	8 395 566	5.16×10^{-9}	−0.14	NCU03028		deubiquitination-protection protein dph1	upstream
2	7	871 838	4.21×10^{-11}	−0.99	NCU06948		EF-hand domain-containing protein	3'UTR
2	7	1 091 058	5.03×10^{-9}	−0.85	NCU08399	*pks-4*	polyketide synthase 4, variant	missense
3	1	7 807 362	6.30×10^{-20}	0.28	NCU00621		TORC1 growth control complex subunit Kog1	5'UTR
3	2	683 209	2.70×10^{-10}	0.12	NCU03448		hypothetical protein	upstream
3	2	3 468 329	7.39×10^{-12}	−0.09	NCU08496		hypothetical protein	missense
3	3	337 373	2.91×10^{-9}	−0.14	NCU10002		hypothetical protein	synonymous
3	5	4 421 459	4.56×10^{-16}	−0.21	NCU06730		hypothetical protein	upstream

Figure 6. Manhattan plots for the variance in growth rate explained by the PC1, PC2 and PC3. The solid horizontal lines denote a P-value threshold of 0.01 after a Bonferroni correction. The dotted vertical lines highlight the candidate SNPs that were significant at least for one of the PCs.

4 DISCUSSION

Climate change obscures the increases in the mean and variance of temperature, yet it is not well understood how different thermal environments are selecting individuals, genotypes and genes (Kingsolver et al. 2013; Sinclair et al. 2016; Kristensen et al. 2020). A more thorough understanding of the genetic basis of thermal tolerance can be achieved with studies integrating quantitative and molecular genetics methods (Cortés et al. 2020; Buckley and Kingsolver 2021). To establish whether distinct genetic architectures are regulating the growth rate of *N. crassa* at constant and fluctuating temperatures, we estimated the amounts of genetic variation and covariation, and interpreted the directions of most evolutionary

potential in TPCs. Interestingly, GWAS found many unique loci associated with growth rate

either at constant or fluctuating temperatures, or with the PCs reflecting the modes of the

TPCs. However, the general view of our results indicated a high level of shared genetic

variance across temperatures which was due to the elevated overall performance.

We found that the constant extreme temperature 42 °C and fast 32–42 °C fluctuations

had the most negative effects on the growth rate of *N. crassa*. Previous studies have suggested

that the fast frequency of fluctuations might be more important to the organism's fitness than

the duration or the magnitude of the exposure to extreme temperatures (Kearney et al. 2012;

Marshall and Sinclair 2015). In *N. crassa*, the upregulation of the heat shock protein (HSP)

production is activated in minutes at temperatures around 42 °C (Mohsenzadeh et al. 1998;

Kapoor and Roy 2014), but the HSP synthesis requires energy, which generally leads to

extensive fitness costs after repeated exposures to stressfully high temperatures (Sinclair et

al. 2016). In a cell, the allocation of energy to stress-related functions are known to mutually

repress growth-related pathways such as protein synthesis and ribosome biogenesis (López-

Maury et al. 2008). This mechanism is called a 'general stress response', which redirects the

resources from growth to stress resistance as a response to multiple stresses, meaning that

the most stress-resistant cells are non-growing (Kültz 2005). Under faster fluctuations, there

is also less time to acclimate. In *N. crassa*, there is a peak in HSP production in around 60 min

(at fast fluctuations, one step lasts 15 min), after which the recovery period with declining

mRNA levels and HSP degradation starts (Kapoor and Roy 2014). It has been found that

when temperature fluctuations became faster than the recovery time of *N. crassa*, the growth

rate slows down also at more optimal temperatures, as the recovery starts to take longer than

the acclimation to heat exposure (Kronholm and Ketola 2018).

The quantitative genetics analyses gave us estimates of the genetic variation in several

small-effect loci. We found high heritability values across temperatures, meaning that most

of the variation in growth was due to genetic variation among the strains. We found also

strong genetic correlations between the fast and slow frequency fluctuations, and between

fluctuations and their mean temperatures, predicting correlated responses to the selection

(Czesak et al. 2006). This indicated that in *N. crassa*, there should not be strong trade-offs or

constrains between adaptation to constant and fluctuating temperatures. However, the

genetic correlations and covariances with the high extreme temperature 42 °C were markedly lower, pointing that the growth at 42 °C could evolve more independently from the selection on fitness at other temperatures. Similar quantitative genetic results have been found before for the growth rate of *N. crassa* at constant 40 °C (Moghadam et al. 2020), suggesting that different genes or amounts of gene expression might affect acclimatization at the higher thermal limit.

The high heritability values and strong positive genetic correlations between environments might be found because some of the strains are superior in growth rate, so called 'supergeneralists' or 'master of all temperatures' (Huey and Hertz 1984; Kristensen et al. 2020). We verified this suggestion by PCA and found that most of the variation (PC1, 83 %) was in the individual's overall fitness in growth rate regardless of the temperature, meaning that some strains grew better, and some slower across all the tested temperatures. Previously, the TPC elevation has been found to predominate the growth rate in *N. crassa* (Moghadam et al. 2020) and performance in other species (Yamahira et al. 2007; Shama et al. 2011; Ketola et al. 2014; Latimer et al. 2015; Bartheld et al. 2017; Schaum et al. 2022). These observations are on the contrary to the traditional hypothesis of specialist-generalist trade-offs preventing the evolution of superior generalists (Angilletta 2009), and some studies that have found costs between performances at intermediate and extreme temperatures (reviewed in Logan and Cox 2020). According to a hypothesis of condition-dependent traits, the fact that many studies have failed to find evidence for the costs of plasticity (Berger et al. 2014; Latimer et al. 2015; Murren et al. 2015; Schaum et al. 2022), can be explained by a larger amount of genetic variation in individual's overall fitness traits compared to stress tolerance (Kristensen et al. 2020; Rowe and Houle 1996; Van Noordwijk and De Jong 1986). Based on this, the superiority in fitness could arise due to more efficient resource acquisition, metabolic assimilation, and allocation on dependent traits across temperatures. Indeed, it has been found that the traits associated with individual's overall fitness are usually those related to cells resource uptake and energy efficiency (Kristensen et al. 2005; Pedersen et al. 2008). In addition to allocation of energy in growth across temperatures, allocation trade-offs can exist between correlated traits sharing the same functional or developmental basis (Angilletta et al. 2003; Berger et al. 2014).

We found much less variation explained by the subsequent PCs, indicating little

evolutionary potential in TPCs for the optimum shift towards colder temperatures (PC2, 8

%) and for the shape change to better performance at high temperatures (PC3, 4 %). The

loadings of the PC2 were interesting since the tolerance to constant temperatures seemed to

also affect the tolerance to fluctuations within the same temperature range, indicating that in

our mapping population, there were, to some extent, cool and warm adapted genotypes

(Angilletta 2009; Berger et al. 2014). The PC3 showed little genetic variance in tolerance to

high extreme temperatures, which most likely means the induction or the magnitude of the

heat-shock response (Moghadam et al. 2020). One possibility is that our nature-derived

strains have undergone selection during their thermal history and are already well-adapted

to heat, decreasing the amount of variation in this tolerance trait. The ability to tolerate

prolonged heat is known to be a trait that is highly evolutionary conserved as temperature

affects all biochemical reactions (Araújo et al. 2013; Hochachka and Somero 2002). Many

experiments have found little variation in genomic regions important to heat stress,

suggesting evolutionary constraints on heat tolerance (Mitchell and Hoffmann 2010;

Kellermann et al. 2012; Kristensen et al. 2015). However, some studies have found evidence

that species can expand their heat tolerance if there is a strong selection for better

performance at hot temperatures, which is often observed in evolution and selection

experiments using constantly high temperatures (Bennett et al. 1990; Holder and Bull 2001;

Bubliy and Loeschcke 2005; Hangartner and Hoffmann 2016).

By GWAS we discovered unique large-effect loci associated with growth rate only at

constant temperatures or fluctuations of both ranges and frequencies, but also genes that had

shared associations at constant and fluctuating temperatures. Interestingly, at temperatures

close to extremes only unique SNPs were associated, whereas shared SNPs were associated

at more optimal temperatures, pointing that adaptation to extremes could be more specific.

Also, there were no shared SNPs between the fast and slow fluctuations. Most of the allelic

variants did not affect the protein structure or were at the regulatory area of the genes, being

potential regulatory variants. However, the types of allelic variants were not associated

clearly with growth at constant or fluctuating temperatures, even though regulatory area

mutations are often expected to dictate the dynamic responses under thermal fluctuations

(López-Maury et al. 2008). Almost all minor alleles slowed down the growth rate and no

opposite allelic effects on growth were found at different temperatures for the shared SNPs.

This supported our findings from quantitative genetics, which didn't show strong evidence

for the existence of trade-offs. The predominantly negative allelic effects and the absence of

any clear trade-offs have been found also in previous studies on thermal tolerance in *N. crassa*

(Moghadam et al. 2020; Räsänen et al. 2024).

There was not much overlap in SNPs associated with growth rate at different

temperatures and with the PCs, but the overlapping molecular functions of the associated

genes suggested a shared physiology, probably owing to the general stress response (López-

Maury et al. 2008; Gerken et al. 2015). Our genome-wide analysis found candidate genes that

encoded proteins essential for the basic functions in fungi, such as transcription, stress

resistance, cellular growth, and metabolism. Many of these proteins are known to be relevant

to the ecology and fitness of *N. crassa* and showed sensitivity to temperature or other

environmental stresses. It has been found that at lower temperatures, genes having

preparatory functions in thermal tolerance are activated, for example, preventing the damage

on cell structure through membrane composition and anticipatory production of stress

proteins (Lecheta et al. 2020). At higher temperatures more functions should be in dynamic

stress protein production, for example, in the upregulation of the HSP synthesis to repair the

unfolded proteins (Sørensen et al. 2003; Lecheta et al. 2020). Our results were in line with

these assumptions, as at temperatures up to optimum, proteins functioned in transcription,

ribosome biogenesis, nutrient uptake, and energy production. At temperatures of both lower

and higher range, and especially at fast fluctuations, we found a regulatory area mutation in

DNAJ (HSP40) protein, and there was also a nearly significant association with the TPC

elevation. At constant 37 °C the encoded proteins worked in signal transduction and there

was already heat related regulation for growth. At higher range 32–42 °C fluctuations we

found genes that affected the folding and thermodynamic stability of the proteins. It has been

suggested that at stressfully high temperatures, the adaptation can be driven also by other

genes than HSPs because of the costs of maintaining a dynamic stress response (Sørensen et

al. 2003). At constant 42 °C and with PC3 we found TORC1 that regulates mycelial growth

under heat stress (Urban et al. 2007; López-Maury et al. 2008), supported by the negligible

growth rates observed at high extreme temperature.

Overall, our results indicated that there is a shared polygenic basis of thermal tolerance

in *N. crassa* at constant and fluctuating temperatures. This tolerance seems to evolve mainly

via elevated TPCs, which could be due to individual's overall fitness or the general stress

response. Therefore, the increasing variance in temperature should not set bounds on the

evolutionary potential of *N. crassa*, but the ability to adapt to extreme heat is probably more

constrained. The identified candidate genes are also likely to be important for the thermal

tolerance in growth rate, which is an important fitness trait related to both pathogenicity and

mutualism in filamentous fungi (Pringle and Taylor 2002). As climate change keeps

escalating, more research is needed to scrutinize the genetic basis of thermal tolerance and

the species ability to adapt to various fluctuations. To this end, we argue that quantitative

genetics and molecular methods like GWAS should be used together, since solely these

methods do not provide a complete picture of the genetic architecture and might result in

opposite interpretations.

Acknowledgements

The authors wish to acknowledge CSC – IT Center for Science, Finland, for computational

resources and the Fungal Genetics Stock Center for the strains used in this study. The best

thanks also to Neda Moghadam and Karendeep Sidhu for their part in the data collection.

This study was funded by the University of Jyväskylä Doctoral Programme in Biological and

Environmental Science (ER), and grants from Emil Aaltonen foundation (ER), and the

Academy of Finland (TK: 278751, IK: 274769 and 321584).

Author contributions

IK, TK, and ER planned the questions and the experiment together. ER, ML, PAMS and IK

conducted the laboratory work. IK and ER run the GWAS and other statistical analyses with

advice from TK. ER was responsible for writing the manuscript, and all authors have read and approved the current version. Authors declare no conflicts of interests.

Data access

Genotypic data for the mapping population can be found at: XXXX.doi, and the phenotypic data and scripts at YYYY.doi.

REFERENCES

Angilletta, M. J., R. S. Wilson, C. A. Navas, and R. S. James. 2003. Tradeoffs and the evolution of thermal reaction norms. Trends Ecol. Evol. 18:234–240.

Angilletta, M. J. 2009. Thermal adaptation: a theoretical and empirical synthesis. Oxford Univ. Press, Oxford, U.K.

Araújo, M. B., F. Ferri-Yáñez, F. Bozinovic, P. A. Marquet, F. Valladares, and S. L. Chown. 2013. Heat freezes niche evolution. Ecol. Lett. 16:1206–1219.

Bartheld, J. L., P. Artacho, and L. Bacigalupe. 2017. Thermal performance curves under daily thermal fluctuation: A study in helmeted water toad tadpoles. J. Therm. Biol. 70:80–85.

Basenko, E. Y., J. A. Pulman, A. Shanmugasundram, O. S. Harb, K. Crouch, D. Starns, S. Warrenfeltz, C. Aurrecoechea, C. J. Stoeckert, J. C. Kissinger, D. S. Roos, and C. Hertz-Fowler. 2018. FungiDB: An Integrated Bioinformatic Resource for Fungi and *Oomycetes*. J. Fungi 4:39. Multidisciplinary Digital Publishing Institute.

Bennett, A. F., K. M. Dao, and R. E. Lenski. 1990. Rapid evolution in response to high-temperature selection. Nature 346:79–81.

Berger, D., R. J. Walters, and W. U. Blanckenhorn. 2014. Experimental evolution for generalists and specialists reveals multivariate genetic constraints on thermal reaction norms. J. Evol. Biol. 27:1975–1989.

Björklund, M., E. Ranta, V. Kaitala, L. A. Bach, P. Lundberg, and N. Chr. Stenseth. 2009. Quantitative Trait Evolution and Environmental Change. PLoS ONE 4:e4521.

Botero, C. A., F. J. Weissing, J. Wright, and D. R. Rubenstein. 2015. Evolutionary tipping

points in the capacity to adapt to environmental change. Proc. Natl. Acad. Sci. 112:184–

189.

Bubliy, O. A., and V. Loeschcke. 2005. Correlated responses to selection for stress resistance

and longevity in a laboratory population of *Drosophila melanogaster*. J. Evol. Biol. 18:789–

803.

Buckley, L. B., and J. G. Kingsolver. 2021. Evolution of Thermal Sensitivity in Changing and

Variable Climates. Annu. Rev. Ecol. Evol. Syst. 52:563–586.

Bürkner, P.-C. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. R J.

10:395.

Chevin, L.-M., and A. A. Hoffmann. 2017. Evolution of phenotypic plasticity in extreme

environments. Philos. Trans. R. Soc. B Biol. Sci. 372:20160138.

Colinet, H., B. J. Sinclair, P. Vernon, and D. Renault. 2015. Insects in Fluctuating Thermal

Environments. Annu. Rev. Entomol. 60:123–140.

Cortés, A. J., F. López-Hernández, and D. Osorio-Rodriguez. 2020. Predicting Thermal

Adaptation by Looking Into Populations' Genomic Past. Front. Genet. 11:564515.

Czesak, M. E., C. W. Fox, and J. B. Wolf. 2006. Experimental Evolution of Phenotypic

Plasticity: How Predictive Are Cross-Environment Genetic Correlations? Am. Nat.

168:323–335.

Deatherage, D. E., J. L. Kepner, A. F. Bennett, R. E. Lenski, and J. E. Barrick. 2017. Specificity

of genome evolution in experimental populations of *Escherichia coli* evolved at different

temperatures. Proc. Natl. Acad. Sci. 114:E1904–E1912.

Deshpande, N., M. R. Wilkins, N. Packer, and H. Nevalainen. 2008. Protein glycosylation

pathways in filamentous fungi. Glycobiology 18:626–637.

DeWitt, T.J. & R.B. Langerhans. 2004. Integrated solutions to environmental heterogeneity:

theory of multimoment reaction norms. In Phenotypic Plasticity. Functional and

Conceptual Approaches. T.J. DeWitt & S.M. Scheiner, Eds.: 98–111. New York: Oxford

University Press.

Duun Rohde, P., K. Krag, V. Loeschcke, J. Overgaard, P. Sørensen, and T. Nygaard

Kristensen. 2016. A Quantitative Genomic Approach for Analysis of Fitness and Stress

Related Traits in a *Drosophila melanogaster* Model Population. Int. J. Genomics 2016:1–11.

Ellison, C. E., C. Hall, D. Kowbel, J. Welch, R. B. Brem, N. L. Glass, and J. W. Taylor. 2011. Population genomics and local adaptation in wild isolates of a model microbial eukaryote. Proc. Natl. Acad. Sci. 108:2831–2836.

Fisher, R. A. 1919. XV.—The correlation between relatives on the supposition of Mendelian inheritance. Earth Env. Sci. T. R. So. 52:399–433.

Fisher, K. J., and G. I. Lang. 2016. Experimental evolution in fungi: An untapped resource. Fungal Genet. Biol. 94:88–94.

Fu, W., T. D. O'Connor, and J. M. Akey. 2013. Genetic architecture of quantitative traits and complex diseases. Curr. Opin. Genet. Dev. 23:678–683.

Galagan, J. E., S. E. Calvo, K. A. Borkovich, E. U. Selker, N. D. Read, D. Jaffe, W. FitzHugh, L.-J. Ma, S. Smirnov, S. Purcell, B. Rehman, T. Elkins, R. Engels, S. Wang, C. B. Nielsen, J. Butler, M. Endrizzi, D. Qui, P. Ianakiev, D. Bell-Pedersen, M. A. Nelson, M. Werner-Washburne, C. P. Selitrennikoff, J. A. Kinsey, E. L. Braun, A. Zelter, U. Schulte, G. O. Kothe, G. Jedd, W. Mewes, C. Staben, E. Marcotte, D. Greenberg, A. Roy, K. Foley, J. Naylor, N. Stange-Thomann, R. Barrett, S. Gnerre, M. Kamal, M. Kamvysselis, E. Mauceli, C. Bielke, S. Rudd, D. Frishman, S. Krystofova, C. Rasmussen, R. L. Metzenberg, D. D. Perkins, S. Kroken, C. Cogoni, G. Macino, D. Catcheside, W. Li, R. J. Pratt, S. A. Osmani, C. P. C. DeSouza, L. Glass, M. J. Orbach, J. A. Berglund, R. Voelker, O. Yarden, M. Plamann, S. Seiler, J. Dunlap, A. Radford, R. Aramayo, D. O. Natvig, L. A. Alex, G. Mannhaupt, D. J. Ebbole, M. Freitag, I. Paulsen, M. S. Sachs, E. S. Lander, C. Nusbaum, and B. Birren. 2003. The genome sequence of the filamentous fungus *Neurospora crassa*. Nature 422:859–868.

Gaydos, T. L., N. E. Heckman, M. Kirkpatrick, J. R. Stinchcombe, J. Schmitt, J. Kingsolver, and J. S. Marron. 2013. Visualizing genetic constraints. Ann. Appl. Stat. 7:860–882.

Georgianna, D. R., A. M. Hawkridge, D. C. Muddiman, and G. A. Payne. 2008. Temperature-Dependent Regulation of Proteins in *Aspergillus flavus*: Whole Organism Stable Isotope Labeling by Amino Acids. J. Proteome Res. 7:2973–2979.

Gerken, A. R., O. C. Eller, D. A. Hahn, and T. J. Morgan. 2015. Constraints, independence, and evolution of thermal plasticity: Probing genetic architecture of long- and short-term thermal acclimation. Proc. Natl. Acad. Sci. 112:4399–4404.

Gilchrist, G. W. 1995. Specialists and Generalists in Changing Environments. I. Fitness Landscapes of Thermal Sensitivity. Am. Nat. 146:252–270.

Hangartner, S., and A. A. Hoffmann. 2016. Evolutionary potential of multiple measures of upper thermal tolerance in *Drosophila melanogaster*. Funct. Ecol. 30:442–452.

Hochachka, P. W., and Somero, G. N. 2002. Biochemical Adaptation: Mechanism and Process in Physiological Evolution. New York: Oxford University Press.

Holder, K. K., and J. J. Bull. 2001. Profiles of Adaptation in Two Similar Viruses. Genetics 159:1393–1404.

Huang, M., X. Liu, Y. Zhou, R. M. Summers, and Z. Zhang. 2019. BLINK: a package for the next level of genome-wide association studies with both individuals and markers in the millions. GigaScience 8.

Huey, R. B., and P. E. Hertz. 1984. Is a jack-of-all-temperatures a master of none? Evolution 38:441–444.

IPCC 2018: Summary for policymakers. — An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty, World Meteorological Organization, Geneva, Switzerland.

Izem, R., and J. G. Kingsolver. 2005. Variation in Continuous Reaction Norms: Quantifying Directions of Biological Interest. Am. Nat. 166:277–289.

Kapoor, M., and S. S. Roy. 2014. A bird's eye-view of molecular chaperones and some facets of the heat shock response of *Neurospora crassa*. Annual Review of Plant Pathology 6:1.

Kearney, M. R., A. Matzelle, and B. Helmuth. 2012. Biomechanics meets the ecological niche: the importance of temporal data resolution. J. Exp. Biol. 215:1422–1424.

Kellermann, V., J. Overgaard, A. A. Hoffmann, C. Fløjgaard, J.-C. Svenning, and V. Loeschcke. 2012. Upper thermal limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. Proc. Natl. Acad. Sci. 109:16228–16233.

Ketola, T., V. M. Kellermann, V. Loeschcke, A. López-Sepulcre, and T. N. Kristensen. 2014. Does environmental robustness play a role in fluctuating environments? Evolution 68:587–594.

Ketola, T., and T. N. Kristensen. 2017. Experimental Approaches for Testing if Tolerance Curves Are Useful for Predicting Fitness in Fluctuating Environments. Front. Ecol. Evol. 5:129.

Ketola, T., and K. Saarinen. 2015. Experimental evolution in fluctuating environments: tolerance measurements at constant temperatures incorrectly predict the ability to tolerate fluctuating temperatures. J. Evol. Biol. 28:800–806.

Khan, M. I., Z. Kainat, S. Maqbool, A. Mehwish, S. Ahmad, H. M. Suleman, Z. Mahmood, M. Ali, A. Aziz, A. Rasheed, and H. Li. 2022. Genome-wide association for heat tolerance at seedling stage in historical spring wheat cultivars. Front. Plant Sci. 13:972481.

Kingsolver, J. G., S. E. Diamond, and L. B. Buckley. 2013. Heat stress and the fitness consequences of climate change for terrestrial ectotherms. Funct. Ecol. 27:1415–1423.

Kingsolver, J. G., R. Gomulkiewicz, and P. A. Carter. 2001. Variation, selection and evolution of function-valued traits. Genetica 112:87–104.

Kingsolver, J. G., G. J. Ragland, and J. G. Shlichta. 2004. Quantitative genetics of continuous reaction norms: thermal sensitivity of caterpillar growth rates. Evolution 58:1521–1529.

Knies, J. L., R. Izem, K. L. Supler, J. G. Kingsolver, and C. L. Burch. 2006. The Genetic Basis of Thermal Reaction Norm Evolution in Lab and Natural Phage Populations. PLoS Biol. 4:e201.

Kristensen, T. N., T. Ketola, and I. Kronholm. 2020. Adaptation to environmental stress at different timescales. Ann. N. Y. Acad. Sci. 1476:5–22.

Kristensen, T. N., J. Overgaard, J. Lassen, A. A. Hoffmann, and C. Sgrò. 2015. Low evolutionary potential for egg-to-adult viability in *Drosophila melanogaster* at high temperatures. Evolution 69:803–814.

Kristensen, T. N., P. Sørensen, M. Kruhøffer, K. S. Pedersen, and V. Loeschcke. 2005. Genome-Wide Analysis on Inbreeding Effects on Gene Expression in *Drosophila melanogaster*. Genetics 171:157–167.

786 Kronholm, I., H. Johannesson, and T. Ketola. 2016. Epigenetic Control of Phenotypic
787  Plasticity in the Filamentous Fungus *Neurospora crassa*. G3-Genes Genom. Genet.
788  6:4009–4022.

789 Kronholm, I., and T. Ketola. 2018. Effects of acclimation time and epigenetic mechanisms on
790  growth of *Neurospora* in fluctuating environments. Heredity 121:327–341.

791 Krystofova, S., and K. A. Borkovich. 2006. The Predicted G-Protein-Coupled Receptor GPR-
792  1 Is Required for Female Sexual Development in the Multicellular Fungus *Neurospora*
793  *crassa*. Eukaryot. Cell 5:1503–1516.

794 Kültz, D. 2005. Molecular and evolutionary basis of the cellular stress response. Annu. Rev.
795  Physiol. 67:225–257.

796 Kundu, P., A. Nehra, R. Gill, N. Tuteja, and S. S. Gill. 2022. Genomic collation revealed the
797  significance of *Piriformospora indica* (*Serendipita indica*) PiEF-hand protein in vesicle
798  trafficking and fungal hyphal growth. South Afr. J. Bot. 150:548–564.

799 Lakin-Thomas, P. 2019. Circadian rhythms, metabolic oscillators, and the target of rapamycin
800  (TOR) pathway: the *Neurospora* connection. Curr. Genet. 65:339–349.

801 Lambros, M., X. Pechuan-Jorge, D. Biro, K. Ye, and A. Bergman. 2021. Emerging adaptive
802  strategies under temperature fluctuations in a laboratory evolution experiment of
803  *Escherichia coli*. Front. Microbiol. 12:724982.

804 Latimer, C. A. L., B. R. Foley, and S. F. Chenoweth. 2015. Connecting thermal performance
805  curve variation to the genotype: a multivariate QTL approach. J. Evol. Biol. 28:155–168.

806 Lecheta, M. C., D. N. Awde, T. S. O'Leary, L. N. Unfried, N. A. Jacobs, M. H. Whitlock, E.
807  McCabe, B. Powers, K. Bora, J. S. Waters, H. J. Axen, S. Frietze, B. L. Lockwood, N. M.
808  Teets, and S. H. Cahan. 2020. Integrating GWAS and Transcriptomics to Identify the
809  Molecular Underpinnings of Thermal Stress Responses in *Drosophila melanogaster*.
810  Front. Genet. 11:658.

811 Levins, R. 1968. Evolution in Changing Environments: Some Theoretical Explorations.
812  Princeton, NJ: Princeton University Press.

813 Li, L., and K. A. Borkovich. 2006. GPR-4 Is a Predicted G-Protein-Coupled Receptor Required
814  for Carbon Source-Dependent Asexual Growth and Development in *Neurospora crassa*.
815  Eukaryot. Cell 5:1287–1300.

Logan, M. L., and C. L. Cox. 2020. Genetic Constraints, Transcriptome Plasticity, and the
Evolutionary Response to Climate Change. Front. Genet. 11:538226.

López-Maury, L., S. Marguerat, and J. Bähler. 2008. Tuning gene expression to changing
environments: from rapid responses to evolutionary adaptation. Nat. Rev. Genet.
9:583–593.

Lynch, M., and W. Gabriel. 1987. Environmental tolerance. Am. Nat. 129:283–303.

Marshall, K. E., and B. J. Sinclair. 2015. The relative importance of number, duration and
intensity of cold stress events in determining survival and energetics of an
overwintering insect. Funct. Ecol. 29:357–366.

McCluskey, K., A. Wiest, and M. Plamann. 2010. The Fungal Genetics Stock Center: a
repository for 50 years of fungal genetics research. J. Biosci. 35:119–126.

McElreath, R. 2015. Statistical rethinking—a Bayesian course with examples in R and Stan.
CRC Press. New York.

McLaren, W., L. Gil, S. E. Hunt, H. S. Riat, G. R. S. Ritchie, A. Thormann, P. Flicek, and F.
Cunningham. 2016. The Ensembl Variant Effect Predictor. Genome Biol. 17:122.

Meehl, G. A., T. Karl, D. R. Easterling, S. Changnon, R. Pielke, D. Changnon, J. Evans, P. Y.
Groisman, T. R. Knutson, K. E. Kunkel, L. O. Mearns, C. Parmesan, R. Pulwarty, T. Root,
R. T. Sylves, P. Whetton, and F. Zwiers. 2000. An Introduction to Trends in Extreme
Weather and Climate Events: Observations, Socioeconomic Impacts, Terrestrial
Ecological Impacts, and Model Projections. Bull. Am. Meteorol. Soc. 81:413–416.
American Meteorological Society.

Metzenberg, R. L. 2003. Vogel's Medium N salts: avoiding the need for ammonium nitrate.
Fungal Genet. Rep. 50:14–14.

Mitchell, K. A., and A. A. Hoffmann. 2010. Thermal ramping rate influences evolutionary
potential and species differences for upper thermal limits in *Drosophila*. Funct. Ecol.
24:694–700.

Moghadam, N. N., K. Sidhu, P. A. M. Summanen, T. Ketola, and I. Kronholm. 2020.
Quantitative genetics of temperature performance curves of *Neurospora crassa*.
Evolution 74:1772–1787.

Mohsenzadeh, S., W. Saupe-Thies, G. Steier, T. Schroeder, F. Fracella, P. Ruoff, and L. Rensing. 1998. Temperature Adaptation of House Keeping and Heat Shock Gene Expression in *Neurospora crassa*. Fungal Genet. Biol. 25:31–43.

Murren, C. J., J. R. Auld, H. Callahan, C. K. Ghalambor, C. A. Handelsman, M. A. Heskel, J. G. Kingsolver, H. J. Maclean, J. Masel, H. Maughan, D. W. Pfennig, R. A. Relyea, S. Seiter, E. Snell-Rood, U. K. Steiner, and C. D. Schlichting. 2015. Constraints on the evolution of phenotypic plasticity: limits and costs of phenotype and plasticity. Heredity 115:293–301.

Palma-Guerrero, J., C. R. Hall, D. Kowbel, J. Welch, J. W. Taylor, R. B. Brem, and N. L. Glass. 2013. Genome Wide Association Identifies Novel Loci Involved in Fungal Communication. PLoS Genet. 9:e1003669.

Pedersen, K. S., T. N. Kristensen, V. Loeschcke, B. O. Petersen, J. Ø. Duus, N. C. Nielsen, and A. Malmendal. 2008. Metabolomic Signatures of Inbreeding at Benign and Stressful Temperatures in *Drosophila melanogaster*. Genetics 180:1233–1243.

Pinto, D., M. Duarte, S. Soares, M. Tropschug, and A. Videira. 2008. Identification of all FK506-binding proteins from *Neurospora crassa*. Fungal Genet. Biol. 45:1600–1607.

Price, A. L., N. J. Patterson, R. M. Plenge, M. E. Weinblatt, N. A. Shadick, and D. Reich. 2006. Principal components analysis corrects for stratification in genome-wide association studies. Nat. Genet. 38:904–909.

Pringle, A., and J. W. Taylor. 2002. The fitness of filamentous fungi. Trends Microbiol. 10:474–481.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at https://www.R-project.org/.

Riehle, M. M., A. F. Bennett, and A. D. Long. 2001. Genetic architecture of thermal adaptation in *Escherichia coli*. Proc. Nat. Acad. Sci. 98:525–530.

Rolandi, C., J. R. B. Lighton, G. J. De La Vega, P. E. Schilman, and J. Mensch. 2018. Genetic variation for tolerance to high temperatures in a population of *Drosophila melanogaster*. Ecol. Evol. 8:10374–10383.

Rowe, L. & D. Houle. 1996. The lek paradox and the capture of genetic variance by condition dependent traits. Proc. R. Soc. B Biol. Sci. 263:1415–1421.

Ryan, F. J., G. W. Beadle, and E. L. Tatum. 1943. The Tube Method of Measuring the Growth
Rate of *Neurospora*. Am. J. Bot. 784-799.

Räsänen, E., N. N. Moghadam, K. Sidhu, P. A. M. Summanen, H.-R. Littunen, T. Ketola, and
I. Kronholm. 2024. Genome-wide association for loci influencing thermal performance
curves in *Neurospora crassa*. bioRxiv.

Schaum, C.-E., A. Buckling, N. Smirnoff, D. J. Studholme, and G. Yvon-Durocher. 2018.
Environmental fluctuations accelerate molecular evolution of thermal tolerance in a
marine diatom. Nat. Commun. 9:1719.

Schaum, C.-E., A. Buckling, N. Smirnoff, and G. Yvon-Durocher. 2022. Evolution of thermal
tolerance and phenotypic plasticity under rapid and slow temperature fluctuations.
Proc. R. Soc. B Biol. Sci. 289:20220834.

Schulte, P. M., T. M. Healy, and N. A. Fangue. 2011. Thermal Performance Curves,
Phenotypic Plasticity, and the Time Scales of Temperature Exposure. Integr. Comp.
Biol. 51:691–702.

Segura, V., B. J. Vilhjálmsson, A. Platt, A. Korte, Ü. Seren, Q. Long, and M. Nordborg. 2012.
An efficient multi-locus mixed-model approach for genome-wide association studies in
structured populations. Nat. Genet. 44:825–830.

Sgrò, C. M., and A. A. Hoffmann. 2004. Genetic correlations, tradeoffs and environmental
variation. Heredity 93:241–248.

Shama, L. N. S., M. Campero-Paz, K. M. Wegner, M. De Block, and R. Stoks. 2011. Latitudinal
and voltinism compensation shape thermal reaction norms for growth rate. Mol. Ecol.
20:2929–2941.

Sinclair, B. J., K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, S. Slotsbo, Y. Dong,
C. D. G. Harley, D. J. Marshall, B. S. Helmuth, and R. B. Huey. 2016. Can we predict
ectotherm responses to climate change using thermal performance curves and body
temperatures? Ecol. Lett. 19:1372–1385.

Sørensen, J. G., T. N. Kristensen, and V. Loeschcke. 2003. The evolutionary and ecological
role of heat shock proteins. Ecol. Lett. 6:1025–1037.

Sørensen, J. G., M. F. Schou, T. N. Kristensen, and V. Loeschcke. 2016. Thermal fluctuations affect the transcriptome through mechanisms independent of average temperature. Sci. Rep. 6:30975.

Tobler, R., S. U. Franssen, R. Kofler, P. Orozco-terWengel, V. Nolte, J. Hermisson, and C. Schlötterer. 2014. Massive habitat-specific genomic response in *D. melanogaster* populations during experimental evolution in hot and cold environments. Mol. Biol. Evol. 31:364–375.

Urban, J., A. Soulard, A. Huber, S. Lippman, D. Mukhopadhyay, O. Deloche, V. Wanke, D. Anrather, G. Ammerer, H. Riezman, J. R. Broach, C. De Virgilio, M. N. Hall, and R. Loewith. 2007. Sch9 Is a Major Target of TORC1 in *Saccharomyces cerevisiae*. Mol. Cell 26:663–674.

Van Noordwijk, A. J., and G. De Jong. 1986. Acquisition and Allocation of Resources: Their Influence on Variation in Life History Tactics. Am. Nat. 128:137–142.

Vasseur, D. A., J. P. DeLong, B. Gilbert, H. S. Greig, C. D. G. Harley, K. S. McCann, V. Savage, T. D. Tunney, and M. I. O'Connor. 2014. Increased temperature variation poses a greater risk to species than climate warming. Proc. R. Soc. B Biol. Sci. 281:20132612.

Wang, J., and Z. Zhang. 2021. GAPIT Version 3: Boosting Power and Accuracy for Genomic Association and Prediction. Genomics Proteomics Bioinformatics 19:629–640.

Wang, Y., X. Zheng, G. Li, and X. Wang. 2023. TORC1 Signaling in Fungi: From Yeasts to Filamentous Fungi. Microorganisms 11:218.

White, B., and D. Woodward. 1995. A simple method for making disposable race tubes. Fungal Genet. Rep. 42:79.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. URL: http://ggplot2.org

Yamahira, K., M. Kawajiri, K. Takeshi, and T. Irie. 2007. Inter- and intrapopulation variation in thermal reaction norms for growth rate: evolution of latitudinal compensation in ectotherms with a genetic constraint. Evolution 61:1577–1589.

Yates, A. D., J. Allen, R. M. Amode, A. G. Azov, M. Barba, A. Becerra, J. Bhai, L. I. Campbell, M. Carbajo Martinez, M. Chakiachvili, K. Chougule, M. Christensen, B. Contreras-Moreira, A. Cuzick, L. Da Rin Fioretto, P. Davis, N. H. De Silva, S. Diamantakis, S. Dyer,

J. Elser, C. V. Filippi, A. Gall, D. Grigoriadis, C. Guijarro-Clarke, P. Gupta, K. E. Hammond-Kosack, K. L. Howe, P. Jaiswal, V. Kaikala, V. Kumar, S. Kumari, N. Langridge, T. Le, M. Luypaert, G. L. Maslen, T. Maurel, B. Moore, M. Muffato, A. Mushtaq, G. Naamati, S. Naithani, A. Olson, A. Parker, M. Paulini, H. Pedro, E. Perry, J. Preece, M. Quinton-Tulloch, F. Rodgers, M. Rosello, M. Ruffier, J. Seager, V. Sitnik, M. Szpak, J. Tate, M. K. Tello-Ruiz, S. J. Trevanion, M. Urban, D. Ware, S. Wei, G. Williams, A. Winterbottom, M. Zarowiecki, R. D. Finn, and P. Flicek. 2022. Ensembl Genomes 2022: an expanding genome resource for non-vertebrates. Nucleic Acids Res. 50: D996–D1003.

Yu, J., J. B. Holland, M. D. McMullen, and E. S. Buckler. 2008. Genetic Design and Statistical Power of Nested Association Mapping in Maize. Genetics 178:539–551.

Zhao, Y., J. Ding, W. Yuan, J. Huang, W. Huang, Y. Wang, and W. Zheng. 2017. Production of a fungal furocoumarin by a polyketide synthase gene cluster confers the chemo-resistance of *Neurospora crassa* to the predation by fungivorous arthropods. Environ. Microbiol. 19:3920–3929.

Zhou, X.-G., P. Yu, C. Dong, C.-X. Yao, Y.-M. Ding, N. Tao, and Z.-W. Zhao. 2016. Proteomic analysis of mycelial proteins from *Magnaporthe oryzae* under nitrogen starvation. Genet. Mol. Res. 15.

Zou, Y., M. Zhang, J. Qu, and J. Zhang. 2018. iTRAQ-Based Quantitative Proteomic Analysis Reveals Proteomic Changes in Mycelium of *Pleurotus ostreatus* in Response to Heat Stress and Subsequent Recovery. Front. Microbiol. 9:2368.