Genome-wide association mapping for growth rate at fluctuating and extreme temperatures

Emmi Räsänen[1, *], Martta Liukkonen[1], Pauliina A. M. Summanen[1], Tarmo Ketola[2] and Ilkka

Kronholm[1]

[1]Department of Biological and Environmental Science, PO Box 35, FI-40014 University of

Jyväskylä, Finland

[2] Department of Forest Science, PO Box 27, FI-00014, University of Helsinki, Finland

*Corresponding author's e-mail: emmi.j.rasanen@jyu.fi

Keywords: climate change, genetic architecture, growth rate, *Neurospora crassa*, quantitative trait,

temperature fluctuation, thermal tolerance

Abstract

Constant temperatures and fluctuations of varying speed affect fitness differently, which has led to

suggestions of distinct genetic architectures and adaptation strategies between constant and fluctuating

thermal environments. However, very little is known about the possible fitness trade-offs and genetic

constraints underlying thermal adaptation and how they affect species' ability to confront climatic

changes. We addressed this gap in knowledge by integrating quantitative genetics and genome-wide

association mapping in the filamentous fungus *Neurospora crassa*. Growth rates were measured for

434 strains under fast and slow frequency fluctuations, at high and low temperature range with respect

to the species' tolerance. We found strong genetic correlations between fast and slow frequency

fluctuations, and between fluctuations and their mean temperatures, but not with the highest extreme

temperature. Positive correlations were supported by high heritability values, pointing that in *N. crassa*

there are no significant trade-offs or genetic constraints in adaptation when variance in temperature

increases. Altogether, our results indicated clearly polygenic basis of thermal tolerance, with most of

the variation in overall performance (83 %), and clearly less in hot-cold trade-off (8 %), or heat stress

tolerance (4 %). Interestingly, GWAS discovered many SNPs associated with growth rate only at

constant temperatures or at fluctuations of both ranges and frequencies. However, the cellular functions

of the associated genes were overlapping, and no opposite allelic effects were found between

treatments. Hence, large-effect loci indicated no trade-offs, but a shared physiology across

temperatures, probably owing to the core stress response or individual's general condition.

1.   Introduction

The global increases in mean and variance of temperature force species to tolerate more frequent

climatic fluctuations such as severe heat waves (Meehl et al. 2000; IPCC 2018). Based on species

performance projections and empirical data, adaptation to temperature fluctuations should be harder

than to tolerating the rise in mean temperature (Vasseur et al. 2014; Colinet et al. 2015). Consequently,

a better understanding of how species can cope with increased thermal variation is crucial for

predicting their survival under climate change (Schulte et al. 2011; Kristensen et al. 2020). According

to traditional hypotheses, the greater variability in temperature should lead to evolution of generalists

with enhanced performance across a wider range of temperatures (Levins 1968; Lynch and Gabriel

1987; Gilchrist 1995). However, the ability to spend more time closer to critical thermal limits could

evolve at the expense of overall fitness, also known as a specialist-generalist trade-off (Angilletta

2009). This hypothesis suggests that the only way to change the breadth of the thermal performance

curves (TPCs) is by lowered elevation at optimal conditions, meaning that populations cannot be

concurrently well-adapted to both mean and variance of temperature (Lynch and Gabriel 1987;

Gilchrist 1995). More recently, studies have acknowledged that adaptations to temperature fluctuations

might not be predictable from TPCs measured across constant temperatures and shown evidence that

distinct genetic architectures regulate adaptation in fluctuating versus constant conditions (Schulte et

al. 2011; Ketola and Saarinen 2015; Sinclair et al. 2016; Sørensen et al. 2016; Ketola and Kristensen

2017). In addition, theoretical studies suggest that there are evolutionary constraints between the types

of fluctuations, for example, that particular adaptive mechanisms are operating in environments that

fluctuate at different speeds (DeWitt and Langerhans 2004; Botero et al. 2015). If these hypotheses

hold true, distinct genetic architectures and constraints are associated with adaptation to constant and

fluctuating environments of varying frequency (Ketola and Kristensen 2017), which could limit

evolutionary adaptation and heavily affect species survival when variation in temperature increases

(Sgrò and Hoffmann 2004; Björklund et al. 2009).

The potential trade-offs in thermal adaptation are often investigated by the principal components of

genetic variation in TPCs and quantitative genetic analysis of covariation (Kingsolver et al. 2001; Izem

and Kingsolver 2005). Specifically, evolutionary constraints are defined as directions where selection

cannot increase fitness due to low amounts of heritable genetic variation or negative genetic

correlations across alternative environmental states (Gaydos et al. 2013). The fluctuations of varying

speed affect fitness differently and are expected to have their characteristic changes in TPCs (Schulte

et al. 2011; Colinet et al. 2015; Schaum et al. 2022). Generally, three modes of variation are observed

in the shape and position of TPCs; the elevation in overall performance, horizontal shift in optimum

between hot and cold temperatures, and in breadth of the TPC (Izem and Kingsolver 2005; Angilletta

2009). Under fast fluctuations, selection should favour elevated performance and more robust

tolerance across temperatures, whereas slower fluctuations should select for more fixed strategies,

reflected as a narrower TPCs, or shift in optimum (DeWitt and Langerhans 2004; Ketola et al. 2014;

Schaum et al. 2022). However, fast fluctuations could also select for reversible phenotypic plasticity

aided by the fast production of heat shock proteins or the maintenance of regulatory enzymes, which

could trade-off, due to their energetic requirements with tolerance to constant environments

(Kristensen et al. 2000; Sørensen et al. 2003).

The traditional quantitative genetics studies have long focused on estimating statistical averages over

a multitude of loci with small phenotypic effects (Fisher 1919). More recently, the next generation

sequencing methods have made it possible to use whole genome sequence data and investigate the

genetic architecture of polygenic traits on the level of individual loci (Fu et al. 2013). For example,

genome-wide association studies (GWAS) link genotypes to phenotypes by mapping mutational

markers such as single nucleotide polymorphisms (SNPs). Using quantitative genetics and the latest

molecular genetics methods together can give us more complete picture of the genetic variation in

thermal tolerance and its evolutionary potential (Cortés et al. 2020; Buckley and Kingsolver 2021),

but so far only few studies have adopted this approach (Gerken et al. 2015; Latimer et al. 2015; Duun

Rohde et al. 2016; Rolandi et al. 2018; Lecheta et al. 2020). Furthermore, most of the quantitative

genetics studies and studies using genome-wide sequence data have focused on evaluating the effects

of constant or increasing heat (Riehle et al. 2001; Knies et al. 2006; Mitchell and Hoffmann 2010;

Khan et al. 2022). Knowledge from differently fluctuating temperatures is parse and studies have

mostly used experimental evolution and resequencing (Tobler et al. 2014; Deatherage et al. 2017;

Schaum et al. 2018; Lambros et al. 2021), whereas only one study has been made on natural genetic

variation (Sørensen et al. 2016). As climate change keeps accelerating, there is a pressing need for

experiments evaluating the predicted fitness trade-offs and constraints in thermal adaptation under

various fluctuations (Chevin and Hoffmann 2017; Buckley and Kingsolver 2021).

To address this gap in knowledge, we investigated if distinct genetic architectures are found at constant

and fluctuating temperatures of varying frequency, and if some genetic constraints might limit thermal

adaptation. We used quantitative genetics and GWAS in the filamentous fungus *Neurospora crassa* to

quantify how different components of genetic variation contribute to the total variation. The thermal

sensitivity to growth rate was measured for a previously developed mapping population (Moghadam et al. 2020; Räsänen et al. 2024) under fast and slow frequency fluctuations, at high and low temperature range, and at constant mean and extreme temperatures of these ranges. In our analyses we used a multivariate approach in which growth rates at each temperature are treated as discrete and potentially genetically correlated traits (Kingsolver et al. 2004). We examined the following questions: (1) How much there is genetic variation in the growth rate at constant and fluctuating temperatures and how strongly is this variation correlated between contrasting thermal environments? (2) To what extent the phenotypic variation in growth rate can be related to broader characteristics of the TPCs? (3) Are there genes associated with growth rate at constant and fluctuating temperatures and are these genes same between contrasting thermal environments? Our study is among the first ones to investigate the genetic basis of thermal tolerance at temperatures that fluctuate with different frequencies.

2. Material and methods

2.1. *Neurospora crassa* mapping population

N. crassa is a clonally and sexually reproducing species that is well-suited for experiments studying genetic differences across environments (Fisher and Lang 2016). In our experiment we used 118 natural strains obtained from the Fungal Genetics Stock Center (Manhattan, Kansas, USA) (McCluskey et al. 2010), originally sampled from natural populations in southeastern USA (Louisiana), Central America and the Caribbean Basin (Ellison et al. 2011; Palma-Guerrero et al. 2013). Previously, some of the natural strains were crossed to form 316 additional offspring strains (Moghadam et al. 2020). In total our nested association mapping population had 434 strains genotyped for 1 473 869 single nucleotide polymorphisms (SNPs) (Räsänen et al. 2024).

2.3. Phenotyping by growth rate

We measured the mycelial growth rate as a proxy for thermal tolerance (Pringle and Taylor 2002).

Before starting experiments, *N. crassa* cultures were maintained on Vogel's medium N (VM)

(Metzenberg 2003) in slants at room temperature. The linear growth rates were measured for each

strain by standard tube method using agar filled serological pipet tips (Ryan et al. 1943; White and

Woodward 1995). A more detailed description of this method can be found from Moghadam et al.

(2020) and Kronholm et al. (2016). We replicated each strain 3 times and put them to grow in 10

thermal treatments (MTM-313 Plant Growth Chamber, HiPoint Corp., Taiwan). We had two

fluctuating regimes, low 25–35 °C and high 32–42 °C, and fast and slow fluctuations with frequencies

of 120 min and 480 min respectively. Thus, fast fluctuations were 4 times faster, having a step duration

of 15 min at each temperature compared to 60 min steps at slow fluctuations. At lower range the steps

were 25–27.5–30–32.5–35–32.5–30–27.5–25°C, and at higher range 32–34.5–37–39.5–42–39.5–37–

34.5–32 °C. The growth rates were measured also at constant mean and extreme temperatures of these

ranges. The measurements at constant lower range temperatures 25 °C, 30 °C, and optimal 35 °C were

collated from previously published data (Moghadam et al. 2020). At higher range the growth rates

were measured at constant 32 °C, 37 °C, and extreme 42 °C which is known to be very stressful for *N.*

crassa (Mohsenzadeh et al. 1998). Altogether we had 12 976 growth assays to use in our analyses.

2.4. Quantitative genetics

The statistical analysis for quantitative genetics parameters was conducted with Bayesian multilevel

model using Hamiltonian Monte Carlo algorithm. We used R version 4.3.1 (R Core Team 2022) and

R package 'brms' version 2.19.0 (Bürkner 2018) that fits Bayesian models by 'Stan' language. The

phenotypic data including all growth assays was used in a multivariate model

$$y_i \sim MVN(\mu_i, E)$$

$$\mu_i = \alpha + \alpha_{g[i]}$$

150
$$\alpha_{g[i]} \sim MVN(0, G)$$

151
$$G = S_G R_G S_G$$

152
$$E = S_E R_E S_E$$

where the growth at each temperature treatment was explained by the strain genotype as a random

variable and modeled as potentially correlated with growth at other temperatures. In this model α was

the vector of intercepts, $\alpha_{g[i]}$ was the vector of genotypic effects, S_G and S_E were 10×10 diagonal

matrices with genetic or environmental standard deviations on the diagonal, and R_G and R_E were

matrices for genetic and environmental correlations, respectively. We had weakly informative priors

as we used the half location-scale version of Student's t-distribution with 3 degrees of freedom and 2.5

as the scale parameter. The prior for the intercept effects was

162
$$\alpha \sim \begin{pmatrix} hT\,(3, 3.1, 2.5) \\ hT\,(3, 3.9, 2.5) \\ hT\,(3, 4.2, 2.5) \\ hT\,(3, 4.3, 2.5) \\ hT\,(3, 3.8, 2.5) \\ hT\,(3, 0.7, 2.5) \\ hT\,(3, 3.6, 2.5) \\ hT\,(3, 3.8, 2.5) \\ hT\,(3, 2.8, 2.5) \\ hT\,(3, 3.2, 2.5) \end{pmatrix}$$

for growth rates from 25 °C to 42 °C, and for fluctuating temperatures in order 25–35 °C fast, 25–35

°C slow, 32–42 °C fast, and 32–42 °C slow. The prior for each standard deviation in the model was σ

$\sim hT(3, 0, 2.5)$, and we used an lkj prior (McElreath 2015) for the correlation matrices: R_E, $R_G \sim$

LKJ(1). For MCMC estimation 2 chains were run with a warmup period of 1000 iterations, followed

by 6000 iterations of sampling, with thinning set to 2. MCMC convergence was monitored by trace

plots and $\hat{R}$ values.

From the model we obtained a G-matrix with estimates of genetic variance and covariance components

for growth rate. The environmental variances were estimated as the variances between the clonal

replicates of the same strain. We considered the posterior mean estimates to be statistically significant

if their 95 % highest posterior density intervals (HPDI) did not overlap with 0. We extracted the

posterior samples from the model and used these to calculate broad-sense heritabilities, genetic

covariances and genetic correlations for growth rate in different thermal environments. We also

quantified the coefficients of variation for both, the genetic and environmental components by

standardizing the scales by the mean phenotype. The R package 'ggplot2' (Wickham 2016) was used

for plotting.

2.5. Principal components analysis

We used a Principal Components Analysis (PCA) to characterize the main patterns of variation in

TPCs, and the amount of phenotypic variation that was attributed to each principal component (PC)

(Izem and Kingsolver 2005). We implemented PCA with built-in R function 'princomp' for the growth

rates averaged per temperature for each strain. The predictive values of the first 3 principal components

(from here on PC1, PC2 and PC3) were saved and used in GWAS. We counted also the Bayesian mean

estimates and HPDIs for the PCs. To form the posterior distributions, we sampled the intercepts

extracted from the multivariate model for each genotype and at each temperature 5000 times. Then we

used the PCA to calculate the proportions of variance explained by the first 6 PCs and the loadings of

PC1, PC2 and PC3 at each temperature.

2.6. Genome-wide association mapping

GWAS was used to study the genotype–phenotype associations and to answer whether the same or

different genes contribute to the thermal tolerance at fluctuating and constant temperatures. We also

investigated which SNPs are associated with PC1, PC2 and PC3 explaining the variation in mean

growth rate. We used nested association mapping that combines information from recent and past

recombination (Yu et al. 2008). We conducted the GWAS by the genome association and prediction

integrated tool (GAPIT) Version 3 R-package (Wang and Zhang 2021). As an input data we had each

strains SNP sequence in HapMap format, and the strains mean growth rate per temperature treatment.

In GWAS, GAPIT 3 identifies the quantitative trait loci via statistical association between the

phenotypes and single-nucleotide markers. It also corrects for the population structure and kinship in

these analyses and so sets control for the false positives. The basic mixed model was

204
$$y = X\beta + Zu + e$$

where y is a vector of phenotypic observations, β is a vector of fixed effects including effects of the

genetic marker and population structure, u is a vector of random genetic effects for each individual, X

and Z are design matrices, and e is vector of residual effects. Variance matrix for u is σ_a^2 K where σ_a^2

is the additive variance and K is the kinship matrix. Variance matrix for e is σ_e^2 I, where σ_e^2 is the

residual variance and I is the identity matrix. For the population structure correction, a principal

component analysis was first done on the SNPs, and the components were incorporated as fixed factors

in the model (Price et al. 2006). We chose the first 4 components for the analysis since they explained

most of the variance in SNPs.

We run GWAS with Bayesian-information and linkage-disequilibrium iteratively nested keyway

method (BLINK) (Huang et al. 2019). BLINK is a multi-locus test (Segura et al. 2012) and the most

statistically powerful method in GAPIT 3 (Wang and Zhang 2021). In BLINK, the SNP markers are

removed iteratively if they are in linkage disequilibrium with the most significantly associated

reference SNP. The procedure of selecting a reference SNP among the remaining markers is repeated

until no markers can be removed. We run the BLINK separately for the mean growth rates at all

temperature treatments and for the phenotypic variation attributed to PC1, PC2 and PC3 describing

the TPC. The significance threshold for associations was set to 0.01 after Bonferroni correction, which

corresponded to a P-value of 6.78×10^{-9}. The signs of the allelic effects were with respect to the minor

allele, i.e., a positive allelic effect indicated that the individuals carrying minor allele had faster growth

rate.

We searched the statistically significant SNPs from Ensemble Fungi release 57 (Yates et al. 2022) and

identified the candidate genes affecting growth rate. We used *Neurospora crassa* (NC12) whole

genome alignment and annotations (Galagan et al. 2003). We also processed the SNP variants by the

Variant Effect Predictor web tool (McLaren et al. 2016) to determine the possible effects on transcript

and protein sequences. In the case of intergenic variants, we reported annotations for a gene that had

the shortest distance from the marker to the transcript. We further searched for the functional protein

annotations of the candidate genes from the UniProt (The UniProt Consortium 2023) and the FungiDB

databases (Basenko et al. 2018).

3. Results

3.1. Growth rates at different temperatures

We observed large variation in growth rates between all temperature treatments (0.01–6.05mm/h) and

there were clear differences in how well strains grow in fluctuating temperatures (Fig. 1A). Genotype

had statistically significant effect on growth rate at each temperature treatment (the model estimates

are reported as mean with 95 % HPD interval given in parenthesis). On average, the growth rates were

faster at more favorable lower temperature range and at slower frequency fluctuations (Fig. 1B). At

constant temperatures, growth rates increased from 25 °C (3.04 mm/h, 3.00–3.08) to 30 °C (3.83 mm/h,

3.77–3.88), and 32 °C (4.07 mm/h, 4.01–4.12). The growth rates were fastest at the optimal

temperature 35 °C (4.16 mm/h, 4.10–4.23) and slowed down at hotter 37 °C (3.73 mm/h, 3.66–3.79)

and 42 °C (0.68 mm/h, 0.65–0.72). At 25–35 °C range, the mean growth rates were very similar, 3.53

mm/h (3.48–3.58) at fast fluctuations and 3.65 mm/h (3.61–3.70) at slow fluctuations. At 32–42 °C

range, the mean growth rate was faster at slow frequency fluctuations 3.10 mm/h (3.05–3.15) than at

fast fluctuations 2.76 mm/h (2.72–2.81).

Figure 1. (A) The mean growth rates for each *N. crassa* genotype (N = 434) and (B) the posterior mean estimates

of growth rate when the growth in fluctuating environments was compared at high and low temperature range

and at fast and slow frequency. In Fig. B the estimates for the constant temperatures within a temperature range

are indicated by colors (red = high extreme temperature, green = mean temperature, and blue = low extreme

temperature, reflecting 25 °C, 30 °C, and 35 °C at lower range, and 32 °C, 37 °C, and 42 °C at higher range).

The 95 % HPD intervals are marked with dotted lines for constant temperatures. In fluctuating temperatures,

the points are hiding the small 95 % HPD interval bars.

3.2. Quantitative genetics

The multivariate model was used to estimate the genetic variance and covariance components for

growth rate at different temperatures (Table 1). In all thermal treatments, there was genetic variation

for growth rate (posterior means 0.14–0.46) (Table 1), and this was observable also from high

heritability values (posterior means 0.64–0.92) (Fig. 2A). In general, we found strong positive genetic

correlations (posterior means 0.54–0.99) and genetic covariances (posterior means 0.10–0.35) between

treatments (Table 1). However, the correlations with constant 42 °C were markedly lower (posterior

means 0.14–0.39), which was also true for the covariances (posterior means 0.02–0.07). At constant

42 °C there were the most genetic and environmental variance when scaled as the coefficients of

variation (Fig. 2B).

Table 1. Genetic variances, covariances, correlations, and environmental variances for growth rate at constant and fluctuating temperatures estimated from the multivariate

model. In table: upper triangle genetic correlations, lower triangle genetic covariances, diagonal genetic variances (in bold), and last column environmental variances. Estimates

are posterior means with 95 % HPD intervals shown in parentheses.

(°C)	25	30	32	35	37	42	25–35 Fast	25–35 Slow	32–42 Fast	32–42 Slow	VarE
25	**0.15 (0.14–0.18)**	0.95 (0.93–0.96)	0.79 (0.75–0.83)	0.80 (0.76–0.84)	0.54 (0.46–0.61)	0.14 (0.04–0.23)	0.84 (0.81–0.87)	0.84 (0.81–0.88)	0.57 (0.49–0.64)	0.63 (0.57–0.69)	0.02 (0.02–0.02)
30	0.20 (0.18–0.23)	**0.29 (0.26–0.33)**	0.90 (0.87–0.92)	0.89 (0.86–0.91)	0.63 (0.56–0.68)	0.19 (0.09–0.28)	0.92 (0.89–0.94)	0.92 (0.89–0.94)	0.65 (0.59–0.71)	0.70 (0.65–0.75)	0.04 (0.03–0.04)
32	0.18 (0.16–0.21)	0.28 (0.24–0.32)	**0.34 (0.29–0.38)**	0.87 (0.84–0.90)	0.77 (0.72–0.81)	0.22 (0.13–0.31)	0.98 (0.97–0.99)	0.98 (0.97–0.99)	0.79 (0.74–0.83)	0.83 (0.79–0.86)	0.08 (0.07–0.09)
35	0.21 (0.18–0.25)	0.32 (0.28–0.37)	0.34 (0.30–0.39)	**0.46 (0.40–0.52)**	0.82 (0.78–0.85)	0.27 (0.17–0.36)	0.89 (0.86–0.92)	0.88 (0.85–0.91)	0.82 (0.78–0.86)	0.87 (0.84–0.90)	0.06 (0.06–0.07)
37	0.14 (0.11–0.16)	0.22 (0.18–0.26)	0.29 (0.24–0.33)	0.35 (0.31–0.41)	**0.41 (0.36–0.47)**	0.29 (0.20–0.38)	0.76 (0.71–0.80)	0.78 (0.74–0.82)	0.86 (0.83–0.90)	0.91 (0.89–0.93)	0.04 (0.03–0.04)
42	0.02 (0.01–0.03)	0.04 (0.02–0.06)	0.05 (0.03–0.07)	0.07 (0.04–0.09)	0.07 (0.05–0.09)	**0.14 (0.12–0.16)**	0.20 (0.10–0.29)	0.20 (0.11–0.29)	0.36 (0.27–0.45)	0.39 (0.30–0.48)	0.06 (0.06–0.07)
25–35 Fast	0.16 (0.14–0.18)	0.24 (0.20–0.27)	0.27 (0.23–0.31)	0.28 (0.25–0.33)	0.23 (0.20–0.27)	0.03 (0.02–0.05)	**0.22 (0.19–0.25)**	0.99 (0.98–1.00)	0.77 (0.72–0.82)	0.82 (0.78–0.85)	0.06 (0.05–0.07)
25–35 Slow	0.16 (0.14–0.18)	0.24 (0.21–0.27)	0.27 (0.24–0.31)	0.29 (0.25–0.33)	0.24 (0.21–0.28)	0.04 (0.02–0.05)	0.22 (0.19–0.25)	**0.23 (0.20–0.26)**	0.76 (0.71–0.81)	0.82 (0.78–0.85)	0.06 (0.05–0.07)
32–42 Fast	0.10 (0.08–0.12)	0.16 (0.13–0.18)	0.20 (0.17–0.23)	0.24 (0.21–0.28)	0.24 (0.21–0.28)	0.06 (0.04–0.08)	0.16 (0.14–0.19)	0.16 (0.14–0.19)	**0.19 (0.17–0.22)**	0.96 (0.94–0.97)	0.11 (0.10–0.12)
32–42 Slow	0.13 (0.11–0.15)	0.19 (0.16–0.23)	0.24 (0.21–0.28)	0.30 (0.26–0.34)	0.30 (0.26–0.34)	0.07 (0.05–0.09)	0.20 (0.17–0.23)	0.20 (0.17–0.23)	0.21 (0.19–0.25)	**0.26 (0.22–0.29)**	0.05 (0.05–0.06)

Figure 2. (A) Heritabilities of growth rate and (B) coefficients of genetic and environmental variation at each temperature treatment. Estimates are posterior means with 95 %

HPD intervals (some points are larger than the error bars).

3.3. Principal component analysis

PCA was able to find 3 principal components, explaining 95 % of the total genetic variance in mean

growth rate (Fig. 3A). The loadings of these 3 principal components at different thermal treatments

were easily interpreted as TPC attributes (Fig. 3B). PC1 explained 83 % of the total variation and had

the largest loading (0.44) at the optimal temperature 35 °C. As all loadings were positive, PC1 was

interpreted as the general fitness in growth i.e. the elevation of the TPC (Fig. 3C). PC2 explained 8 %

of the variation and was interpreted as the horizontal shift in TPC optimum towards colder

temperatures (Fig. 3D). Interestingly, the constant 37 °C and 42 °C, and 32–42 °C fluctuations all had

negative loadings in PC2. The proportion of variance for PC3 was only 4 % and the loadings were

interpreted as the change in the shape of the TPC (Fig. 3E). There was a strong positive loading (0.90)

for PC3 at constant 42 °C, indicating increased growth rate at the high thermal extreme.

Figure 3. (A) Proportions of variance explained by the different principal components and (B) the loadings of

PC1, PC2 and PC3 for each temperature treatment. Shown are the posterior means and 95 % HPD intervals as

error bars. At lower line are the schematic pictures for the changes in TPC attributes: (C) the elevation, (D) the

hot and cold shift in optimum, and (E) the tolerance of high thermal extreme.

3.4. GWAS

3.4.1. Associations and allelic effects

GWAS found 25 associations with growth rate at different temperatures, which were related to 21

SNPs, and located in 19 genes or intergenic loci (Table 2). At constant temperatures optimal or lower,

significant SNPs were found from chromosomes 3, 5 and 6 (Fig. 4, Fig. 5). At constant temperatures

above the optimal, the associated SNPs were in chromosomes 1 and 6 (Fig. 5). At fluctuations,

associations were in chromosomes 1, 3 and 6 (Fig. 4, Fig. 5). The strongest associations were found

at 42 °C in chromosome 1: 7 807 362 ($P = 4.66 \times 10^{-32}$), at slow 25–35 °C fluctuations in 1: 7 378 135

($P = 2.47 \times 10^{-21}$), and at fast 25–35 °C fluctuations in 3: 2 881 603 ($P = 2.77 \times 10^{-19}$) and in 6: 3 600

909 ($P = 1.19 \times 10^{-19}$). From significantly associated SNPs, 8 were detected only at fluctuating

temperatures, 12 only at constant temperatures, and 2 at both, fluctuating and constant temperatures.

Altogether there were 18 unique SNPs associated only at one temperature treatment and 3 shared

SNPs that were associated at more than one temperature treatment (Table 2). Only unique SNPs were

found at the lowest extreme temperature 25 °C, above optimum at 37 °C and 42 °C, and at higher 32–

42 °C range. The shared SNPs (3: 2 300 888, 3: 2 881 603, and 6: 3 600 716) were found at more

favorable 30 °C, 32 °C, and 35 °C temperatures, and at lower 25–35 °C range.

Most of the SNPs had a negative effect on growth rate when examined with respect to the minor allele

(Table 2). This was also true for the shared SNPs, for which the direction of the allelic effects on

growth rate was negative regardless of the temperature. The only positive effect on growth rate (0.18

mm/h) was found at 37 °C for SNP 1: 7 182 169. In general, the minor alleles in our natural

populations of *N. crassa* are known to be mildly deleterious and occur at low frequencies (Räsänen

et al. 2024). The magnitude of the allelic effects was, on average, similar for the unique and the shared

SNPs. The greatest allelic effects were found at 37 °C (–0.69 mm/h), at 35 °C (–0.48 mm/h), and at

fast 25–35 °C fluctuations (–0.42 mm/h).

3.4.2. Allelic variants

Most of the allelic variants were synonymous or in non-coding area up- or downstream from the gene

(Table 2). GWAS found also 4 missense variants changing the amino acid and affecting the protein

structure. These were in chromosome 6: 3 669 489 at 25 °C changing serine to proline, in 3: 2 179 382

at 35 °C changing glycine to serine, in 6: 430 408 at 42 °C changing serine to phenylalanine, and in

3: 1 402 371 at fast 32–42 °C fluctuations, changing serine to glycine. All missense variants were

unique associations, whereas shared associations were synonymous or in non-coding area. However,

there was no obvious connection between the types of allelic variants and the associations being found

at constant or fluctuating temperatures.

3.4.3. Gene annotations and protein functions

The associations found at constant lower range temperatures 25 °C, 30 °C, and optimal 35 °C have

been previously annotated in Räsänen et al. (2024). At constant 25 °C were associated the *fpo-1*

(NCU00236) gene which encodes flavoprotein oxygenase, a gene (NCU01043) coding

uncharacterized protein, and a missense mutation in the gene (NCU17138) encoding haloacid

dehalogenase (HAD) like protein (Table 2). At constant 30 °C and 32 °C, and at fast fluctuations of

both higher and lower range, 2 SNPs were associated in the regulatory area of the gene (NCU00170)

encoding DNAJ domain-containing protein, which belongs to the HSP40 heat shock proteins. At

constant 30 °C and 35 °C, and at fast 25–35 °C fluctuations there were 2 synonymous mutations in

the *ncw-1* (NCU05137) gene encoding a highly conserved non-anchored cell wall protein 1. At

constant 32 °C were associated a gene (NCU05084) encoding an uncharacterized protein, and a gene

(NCU03622) encoding UBA/TS-N domain-containing protein. This protein is known to function in

endocytosis and to have temperature-dependent regulation of expression in fungus *Aspergillus flavus*

(Georgianna et al. 2008). At constant 32 °C and at slow 25–35 °C fluctuations was associated a gene

(NCU00017) encoding Zn(2)-C6 fungal-type domain-containing protein, that is a transcription

regulator. At optimal 35 °C was found a unique missense SNP in the *rbg-37* (NCU06278) gene

producing a variant in elongation factor 2.

At constant 37 °C was found a synonymous association in the *gpr-1* (NCU00786) gene encoding G-

protein coupled receptor. The G-protein receptors work on the cell surface and affect the gene

regulation through signal pathways. The g*rp-1* gene has been found to be important in the sexual

reproduction of *N. crassa* (Krystofova and Borkovich 2006), and the deletion mutants of another G-

protein coupled receptor gene (*grp-4*), have been shown to grow well at high temperatures up to 42 °C

(Li and Borkovich 2006). At constant 37 °C was also associated the *stk-32* (NCU04755) gene

encoding protein kinase domain-containing protein ppk32 (large allelic effect of –0.69 mm/h). The

protein kinases are universally important coenzymes and enzyme regulators that bind to ATP in

phosphorylation and work in signal transduction. In the previous GWAS, the association with the *stk-*

32 gene was found for the maximum growth rate at optimal temperature, when the shifts in TPCs

were investigated (Räsänen et al. 2024).

At extreme 42 °C we found association with the gene (NCU00621) encoding TORC1 growth control

complex subunit Kog1, and a missense mutation in the hypothetical protein (NCU04701) that had a

high minor allele frequency in the mapping population (0.47) (Table 2). The target of rapamycin

complex 1 (TORC1) is a highly conserved serine/threonine kinase that regulates fungal mycelial

growth and protein synthesis in response to nutrient and energy signaling pathways (Wang et al.

2023). In yeast cells, TORC1 has been shown to regulate growth also by sensing environmental

stressors such as high temperature (Urban et al. 2007). In *N. crassa*, several studies have linked TOR

signaling pathway to cell cycle and circadian rhythmicity, which are also affected by the changes in

temperature (Lakin-Thomas 2019).

At fast 25–35 °C fluctuations was an unique association in the gene (NCU06484) encoding AAA+

ATPase domain-containing protein (Table 2), that positively regulates transcription. At slow 25–

35 °C, the found SNPs were in the gene (NCU00735) encoding Ribosome biogenesis protein SLX9,

and upstream from the gene (NCU08213) encoding a hypothetical protein. At fast 32–42 °C

fluctuations was associated a missense mutation in the gene (NCU06434) encoding a hypothetical

protein. At slow 32–42 °C fluctuations, the first association was in the gene (NCU00669) producing

an oligosaccharyl transferase (OST) subunit. OST is a membrane protein complex, which catalyzes

a central step in the glycosylation pathway. Glycosylation affects the folding and thermodynamic

stability of the proteins. The N-glycosylation pathway has been found to be highly conserved across

species of filamentous fungi (Deshpande et al. 2008). The second association was upstream from the

fkr-3 (NCU04371) gene which produces a peptidyl-prolyl cis-trans isomerase. The *fkr-3* gene

accelerates the protein folding by catalyzing the cis-trans isomerization activity. In *N. crassa*, the *fkr-*

3 deletion mutants have temperature-sensitive phenotype (Pinto et al. 2008), and this gene is

upregulated during heat stress and recovery also in other fungi (Zou et al. 2018).

Table 2. SNPs that were significantly associated with growth rate at constant and fluctuating temperatures, their positions in genome (bp), allelic effects (mm/h), variant types

and annotated proteins. The sign of the allelic effect is with respect to the minor allele and the annotations for intergenic variants are for the closest gene.

Temperature (°C)	Chromosome	Position (bp)	P-value	Allelic effect (mm/h)	Gene	Symbol	Annotation	Variant
25	3	3 119 653	1.31×10^{-10}	−0.20	NCU00236	fpo-1	flavoprotein oxygenase	synonymous
25	5	2 747 660	2.22×10^{-10}	−0.13	NCU01043		hypothetical protein	3'UTR
25	6	3 669 489	3.27×10^{-12}	−0.22	NCU17138		haloacid dehalogenase like protein	missense
30	3	2 881 603	2.69×10^{-11}	−0.26	NCU00170		DNAJ domain-containing protein	upstream
30	6	3 600 716	1.18×10^{-9}	−0.23	NCU05137	ncw-1	non-anchored cell wall protein 1	synonymous
32	3	2 300 888	7.10×10^{-13}	−0.32	NCU00017		Zn(2)-C6 fungal-type domain-containing protein	ustream
32	3	2 881 603	6.09×10^{-10}	−0.27	NCU00170		DNAJ domain-containing protein	upstream
32	5	1 883 866	5.55×10^{-15}	−0.22	NCU03622		UBA/TS-N domain-containing protein	3'UTR
32	6	3 797 277	1.99×10^{-10}	−0.29	NCU05084		hypothetical protein	3'UTR
35	3	2 179 382	9.39×10^{-17}	−0.48	NCU06278	rbg-37	elongation factor 2, variant 2	missense
35	6	3 600 716	1.32×10^{-10}	−0.26	NCU05137	ncw-1	non-anchored cell wall protein 1	synonymous
37	1	7 182 169	5.84×10^{-11}	0.18	NCU00786	gpr-1	G-protein coupled receptor	synonymous
37	6	686 146	1.25×10^{-13}	−0.69	NCU04755	stk-32	protein kinase domain-containing protein ppk32	5'UTR
42	1	7 807 362	4.66×10^{-32}	−0.38	NCU00621		TORC1 growth control complex subunit Kog1	5'UTR
42	6	430 408	6.38×10^{-13}	−0.11	NCU04701		hypothetical protein	missense
25–35 Fast	3	1 573 941	6.65×10^{-12}	−0.22	NCU06484		AAA+ ATPase domain-containing protein	3'UTR
25–35 Fast	3	2 881 603	2.77×10^{-19}	−0.40	NCU00170		DNAJ domain-containing protein	upstream
25–35 Fast	6	3 600 909	1.19×10^{-19}	−0.27	NCU05137	ncw-1	non-anchored cell wall protein 1	synonymous
25–35 Slow	1	7 378 135	2.47×10^{-21}	−0.24	NCU00735		Ribosome biogenesis protein SLX9	synonymous
25–35 Slow	3	2 300 888	2.26×10^{-16}	−0.29	NCU00017		Zn(2)-C6 fungal-type domain-containing protein	upstream
25–35 Slow	3	4 656 366	3.20×10^{-12}	−0.21	NCU08213		hypothetical protein	upstream
32–42 Fast	3	1 402 371	3.39×10^{-9}	−0.42	NCU06434		hypothetical protein	missense
32–42 Fast	3	2 881 312	5.95×10^{-13}	−0.22	NCU00170		DNAJ domain-containing protein	upstream
32–42 Slow	1	7 625 265	2.56×10^{-13}	−0.21	NCU00669		oligosaccharyl transferase subunit	3'UTR
32–42 Slow	6	3 599 935	3.46×10^{-12}	−0.23	NCU04371	fkr-3	peptidyl-prolyl cis-trans isomerase fkr-3	upstream

Figure 4. Manhattan plots for the growth rate at different temperatures. Solid horizontal line denotes P-value threshold of 0.01 after Bonferroni correction. Dotted vertical lines highlight candidate SNPs that were significant in at least one of the temperature treatments within lower range 25–35°C.

Figure 5. Manhattan plots for the growth rate at different temperatures. Solid horizontal line denotes P-value threshold of 0.01 after Bonferroni correction. Dotted vertical lines highlight candidate SNPs that were significant in at least one of the temperature treatments within higher range 32–42°C.

3.3.5 GWAS for the principal components

GWAS found 8 unique genes associated with PCs characterizing the main patterns of variation in

TPCs (Table 3). No significant SNPs were found for PC1 interpreted as TPC elevation, even though

it explained most of the phenotypic variation (Fig. 3A). However, PC1 had close to significant

association ($P = 2.12 \times 10^{-8}$) with the gene (NCU00170) encoding DNAJ domain-containing protein,

which is likely to be a true association based on its occurrence in the GWAS for growth rates at

different temperatures (Table 2). PC2 was interpreted as the shift in TPC optimum towards colder

temperatures and it had 3 associations located in chromosomes 1 and 7 (Table 3, Fig. 6). The first

association was upstream from the gene (NCU03028) encoding a deubiquitination-protection protein

dph1, which works in protein degradation and localization to vacuoles, and has been found to affect

mycelial growth under starvation stress (Zhou et al. 2016). The second association was in the gene

(NCU06948) encoding an EF-hand domain-containing protein and had a very strong allelic effect (–

0.99 mm/h). The EF-hand proteins affect enzymatic activity and cell division through microtubule

organization and have been found to be involved in vesicular trafficking and fungal hyphal growth

(Kundu et al. 2022). The third association was a missense variant having large allelic effect (–0.85

mm/h) and changing alanine to serine in the *pks-4* (NCU08399) gene. The *pks-4* encodes polyketide

synthase 4 variant which functions in the formation of fatty acids. The polyketide synthases have

been linked to the sexual reproduction and synthesis of secondary metabolites against predation,

which both are important to the survival of *N. crassa* (Zhao et al. 2017).

PC3 was interpreted as the shape change towards high extreme and had 5 associations in

chromosomes 1, 2, 3, and 5 (Table 3, Fig. 6). The strongest association ($P = 6.30 \times 10^{-20}$) was in the

gene (NCU00621) encoding TORC1 growth control complex subunit Kog1, that was also associated

at 42 °C, but with PC3 it had moderate positive allelic effect (Table 2, Table 3). Another association

with positive allelic effect was upstream from the gene (NCU03448) with unknown function (Table

3). In addition, 3 genes having negative allelic effects and encoding hypothetical proteins were

associated with PC3. One of them (NCU08496) was a missense variant changing alanine to serine

and had a high minor allele frequency (0.45) in the mapping population. Other mutations were

synonymous (NCU10002) and intergenic upstream (NCU06730) variants. Compared to our GWAS

for the PCs, different genes were found to be important at TPC optimal temperature, and hot and cold

extremes in a previous GWAS for the growth of *N. crassa* (Räsänen et al. 2024).

Table 3. SNPs that were significantly associated with the predictive values of the first 3 principal components, their positions in genome (bp), allelic effects on

phenotypic variance, allelic variant types and annotated proteins. The sign of the allelic effect is with respect to the minor allele and the annotations for intergenic

variants are for the closest gene.

PC	Chromosome	Position (bp)	P-value	Allelic effect	Gene	Symbol	Annotation	Variant
2	1	8 395 566	5.16×10^{-9}	−0.14	NCU03028		deubiquitination-protection protein dph1	upstream
2	7	871 838	4.21×10^{-11}	−0.99	NCU06948		EF-hand domain-containing protein	3'UTR
2	7	1 091 058	5.03×10^{-9}	−0.85	NCU08399	pks-4	polyketide synthase 4, variant	missense
3	1	7 807 362	6.30×10^{-20}	0.28	NCU00621		TORC1 growth control complex subunit Kog1	5'UTR
3	2	683 209	2.70×10^{-10}	0.12	NCU03448		hypothetical protein	upstream
3	2	3 468 329	7.39×10^{-12}	−0.09	NCU08496		hypothetical protein	missense
3	3	337 373	2.91×10^{-9}	−0.14	NCU10002		hypothetical protein	synonymous
3	5	4 421 459	4.56×10^{-16}	−0.21	NCU06730		hypothetical protein	upstream

Figure 6. Manhattan plots for the variance in growth rate explained by PC1, PC2 and PC3. Solid horizontal line

denotes P-value threshold of 0.01 after Bonferroni correction. Dotted vertical lines highlight candidate SNPs

that were significant in at least one of the PCs.

4. Discussion

Climate change obscures the increases in the mean and variance of temperature, yet it is not well

understood how different thermal environments are selecting individuals, genotypes and genes

(Kingsolver et al. 2013; Sinclair et al. 2016; Kristensen et al. 2020). A more thorough understanding

of the genetic basis of thermal tolerance can be achieved with studies integrating quantitative and

molecular genetics methods (Cortés et al. 2020; Buckley and Kingsolver 2021). To establish whether

distinct genetic architectures are regulating the growth rate of *N. crassa* at constant and fluctuating

temperatures, we estimated the amounts of genetic variation and co-variation, and interpreted the

directions of most evolutionary potential in TPCs. Interestingly, GWAS found many unique loci

associated with growth rate either at constant or fluctuating temperatures of varying frequency, or with

the PCs reflecting the components of the TPCs. However, the general view of our results indicated a

high level of shared genetic variance across temperatures which was due to the elevated overall

performance.

We found that the constant extreme temperature 42 °C and rapid 32–42 °C fluctuations had the most

negative effects on the growth rate of *N. crassa*. Previous studies have suggested that the fast frequency

of fluctuations might be more important to the organism's fitness than the duration or the magnitude

of the exposure to extreme temperatures (Kearney et al. 2012; Marshall and Sinclair 2015). In *N.*

crassa, the upregulation of the heat shock protein (HSP) production is activated in minutes at

temperatures around 42 °C (Mohsenzadeh et al. 1998; Kapoor and Roy 2014), but HSP synthesis

requires energy, which generally leads to extensive fitness costs after repeated exposures to stressfully

high temperatures (Sinclair et al. 2016). In a cell, the allocation of energy to stress-related functions

are known to mutually repress growth-related pathways such as protein synthesis and ribosome

biogenesis (López-Maury et al. 2008). This mechanism is called a 'core stress response', which

redirects the resources from growth to stress resistance as a response to multiple stresses, meaning that

the most stress-resistant cells are non-growing (Kültz 2005). Under faster fluctuations, there is also

less time to acclimate. In *N. crassa*, there is a peak in HSP production in around 60 min (at fast

fluctuations, one step lasts 15 min), after which the recovery period with declining mRNA levels and

HSP degradation starts (Kapoor and Roy 2014). It has been found that when temperature fluctuations

became faster than the recovery time of *N. crassa*, the growth rate slows down also at more optimal

temperatures, as the recovery starts to take longer than the acclimation to heat exposure (Kronholm

and Ketola 2018).

The quantitative genetics analyses gave us estimates of the genetic variation in several small-effect

loci. We found high heritability values across temperatures, meaning that most of the variation in

growth was due to genetic variation among the strains. We found also strong genetic correlations

between the fast and slow frequency fluctuations, and between fluctuating environments and their

mean temperatures, predicting correlated responses to the selection (Czesak et al. 2006). This indicated

that in *N. crassa*, there should not be strong trade-offs or constrains between adaptation to constant

and fluctuating temperatures. However, the genetic correlations and covariances with the high extreme

temperature 42 °C were markedly lower, pointing that the growth at 42 °C could evolve more

independently from the selection on fitness at other temperatures. Similar quantitative genetic results

have been found before for the growth rate of *N. crassa* at constant 40 °C (Moghadam et al. 2020),

suggesting that different genes or amounts of gene expression might affect acclimatization at the higher

thermal limit.

The high heritability values and strong positive genetic correlations between environments might be

found because some of the strains are superior in growth rate, so called 'supergeneralists' or 'master

of all temperatures' (Huey and Hertz 1984; Kristensen et al. 2020). We verified this suggestion by

PCA and found that most of the variation (PC1, 83 %) was in the general fitness in growth rate

regardless of the temperature, meaning that some strains grew better, and some slower across all the

tested temperatures. Previously, the TPC elevation has been found to predominate growth rate in *N.*

crassa (Moghadam et al. 2020) and performance in other species (Yamahira et al. 2007; Shama et al.

2011; Ketola et al. 2014; Latimer et al. 2015; Bartheld et al. 2017; Schaum et al. 2022). These

observations are on the contrary to the traditional hypothesis of a specialist-generalist trade-offs

preventing the evolution of superior generalists (Angilletta 2009), and some studies that have found

costs between performance at intermediate and extreme temperatures (reviewed in Logan and Cox

2020). According to a hypothesis of condition-dependent traits, the fact that many studies have failed

to find evidence for the costs of plasticity (Berger et al. 2014; Latimer et al. 2015; Murren et al. 2015;

Schaum et al. 2022), can be explained by the larger amount of genetic variation in general fitness traits

compared to stress tolerance (Kristensen et al. 2020; Rowe and Houle 1996; Van Noordwijk and De

Jong 1986). Based on this, the superiority in fitness could arise due to more efficient resource

acquisition, metabolic assimilation, and allocation on dependent traits across temperatures. Indeed, it

has been found that the traits associated with general fitness are usually those related to cells resource

uptake and energy efficiency (Kristensen et al. 2005; Pedersen et al. 2008). In addition to allocation of

energy in growth across temperatures, allocation trade-offs can exist between correlated traits sharing

the same functional or developmental basis (Angilletta et al. 2003; Berger et al. 2014).

We found much less variation explained by the subsequent PCs, indicating little evolutionary potential

in TPCs for the optimum shift towards colder temperatures (PC2, 8 %) and for the shape change to

better performance at high temperatures (PC3, 4 %). The loadings of the PC2 were interesting since

the tolerance to constant temperatures seemed to also affect the tolerance to fluctuations within the

same temperature range, indicating that in our mapping population, there were, to some extent, cool

and warm adapted genotypes (Angilletta 2009; Berger et al. 2014). The PC3 showed little genetic

variance in tolerance to high extreme temperatures, which most likely means the induction or the

magnitude of the heat-shock response (Moghadam et al. 2020). One possibility is that our nature-

derived strains have undergone selection during their thermal history and are already well-adapted to

heat, decreasing the amount of variation in this tolerance trait. The ability to tolerate prolonged heat is

known to be a trait that is highly evolutionary conserved as temperature affects all biochemical

reactions (Araújo et al. 2013; Hochachka and Somero 2002). Many experiments have found little

variation in genomic regions important to heat stress, suggesting evolutionary constraints on heat

tolerance (Mitchell and Hoffmann 2010; Kellermann et al. 2012; Kristensen et al. 2015). However,

some studies have found evidence that species can expand their heat tolerance if there is strong

selection for better performance at hot temperatures, which is often observed in evolution and selection

experiments using constantly high temperatures (Bennett et al. 1990; Holder and Bull 2001; Bubliy

and Loeschcke 2005; Hangartner and Hoffmann 2016).

By GWAS we discovered unique large-effect loci associated with growth rate only at constant

temperatures or fluctuations of both ranges and frequencies, but also genes that had shared associations

at constant and fluctuating temperatures. Interestingly, at temperatures close to extremes only unique

SNPs were associated, whereas shared SNPs were associated at more optimal temperatures, pointing

that adaptation to extremes could be more specific. Most of the allelic variants did not affect the protein

structure or were at the regulatory area of the genes, being potential regulatory variants. However, the

types of allelic variants were not associated clearly with growth at constant or fluctuating temperatures,

even though regulatory area mutations are often expected to dictate the dynamic responses under

thermal fluctuations (López-Maury et al. 2008). Almost all minor alleles slowed down the growth rate

and no opposite allelic effects on growth were found at different temperatures for the shared SNPs.

This supported our findings from quantitative genetics, which didn't show strong evidence for the

existence of trade-offs. The predominantly negative allelic effects and the absence of any clear trade-

offs have been found also in previous studies on thermal tolerance in *N. crassa* (Moghadam et al. 2020;

Räsänen et al. 2024).

There was not much overlap in SNPs associated with growth rate at different temperatures and with

PCs, but the overlapping molecular functions of the associated genes suggested a shared physiology,

probably owing to the core stress response (López-Maury et al. 2008; Gerken et al. 2015). Our genome-

wide analysis found candidate genes that encoded proteins essential for the basic functions in fungi,

such as transcription, stress resistance, cellular growth, and metabolism. Many of these proteins are

known to be relevant to the ecology and fitness of *N. crassa* and showed sensitivity to temperature or

other environmental stresses. It has been found that at lower temperatures, genes having preparatory

functions in thermal tolerance are activated, for example, preventing the damage on cell structure

through membrane composition and anticipatory production of stress proteins (Lecheta et al. 2020).

At higher temperatures more functions should be in dynamic stress protein production, for example,

in the upregulation of the HSP synthesis to repair the unfolded proteins (Sørensen et al. 2003; Lecheta

et al. 2020). Our results were in line with these assumptions, as at temperatures up to optimum, proteins

functioned in transcription, ribosome biogenesis, nutrient uptake, and energy production. At

temperatures of both lower and higher range, and especially at fast fluctuations, we found a regulatory

area mutation in DNAJ (HSP40) protein, and there was also a nearly significant association with the

TPC elevation. At constant 37 °C the encoded proteins worked in signal transduction and there was

already heat related regulation for growth. At higher range 32–42 °C fluctuations we found genes that

affected the folding and thermodynamic stability of the proteins. It has been suggested that at

stressfully high temperatures, the adaptation can be driven also by other genes than HSPs because of

the costs of maintaining a dynamic stress response (Sørensen et al. 2003). At constant 42 °C and with

PC3 we found TORC1 that regulates mycelial growth under heat stress (Urban et al. 2007; López-

Maury et al. 2008), supported by the negligible growth rates observed at high extreme temperature.

Overall, our results indicated that there is a shared polygenic basis of thermal tolerance in *N. crassa* at

constant and fluctuating temperatures. This tolerance seems to evolve mainly via elevated overall

performance, which could be due to individual's general condition or the core stress response.

Therefore, the increasing variance in temperature should not set bounds on the evolutionary potential

of *N. crassa*, but the ability to adapt to extreme heat is probably more constrained. The identified

candidate genes are also likely to be important for the thermal tolerance in growth rate, which is an

important fitness trait related to both pathogenicity and mutualism in filamentous fungi (Pringle and

Taylor 2002). As climate change keeps escalating, more research is needed to scrutinize the genetic

basis of thermal tolerance and the species ability to adapt to various fluctuations. To this end, we argue

that quantitative genetics and molecular methods like GWAS should be used together, since solely

these methods do not provide a complete picture of the genetic architecture and might result in opposite

interpretations.

Data access

Genotypic data for the mapping population can be found at: XXXX.doi, and the phenotypic data and

scripts at YYYY.doi.

Acknowledgements

The authors wish to acknowledge CSC – IT Center for Science, Finland, for computational resources

and the Fungal Genetics Stock Center for the strains used in this study. The best thanks also to Neda

Moghadam and Karendeep Sidhu for their part in the data collection. This study was funded by the

University of Jyväskylä Doctoral Programme in Biological and Environmental Science (ER), and

grants from Emil Aaltonen foundation (ER), and the Academy of Finland (TK: 278751, IK: 274769

and 321584).

Author contributions

IK, TK, and ER planned the questions and the experiment together. ER, ML, PAMS and IK conducted

the laboratory work. IK and ER run the GWAS and other statistical analyses with advice from TK. ER

was responsible for writing the manuscript, and all authors have read and approved the current version.

Authors declare no conflicts of interests.

References

Angilletta, M. J., R. S. Wilson, C. A. Navas, and R. S. James. 2003. Tradeoffs and the evolution of thermal reaction norms.

Trends Ecol. Evol. 18:234–240.

Angilletta, M. J. 2009. Thermal adaptation: a theoretical and empirical synthesis. Oxford Univ. Press, Oxford, U.K.

Araújo, M. B., F. Ferri-Yáñez, F. Bozinovic, P. A. Marquet, F. Valladares, and S. L. Chown. 2013. Heat freezes niche

evolution. Ecol. Lett. 16:1206–1219.

Bartheld, J. L., P. Artacho, and L. Bacigalupe. 2017. Thermal performance curves under daily thermal fluctuation: A study

in helmeted water toad tadpoles. J. Therm. Biol. 70:80–85.

Basenko, E. Y., J. A. Pulman, A. Shanmugasundram, O. S. Harb, K. Crouch, D. Starns, S. Warrenfeltz, C. Aurrecoechea,

C. J. Stoeckert, J. C. Kissinger, D. S. Roos, and C. Hertz-Fowler. 2018. FungiDB: An Integrated Bioinformatic

Resource for Fungi and *Oomycetes*. J. Fungi 4:39. Multidisciplinary Digital Publishing Institute.

Bennett, A. F., K. M. Dao, and R. E. Lenski. 1990. Rapid evolution in response to high-temperature selection. Nature

346:79–81.

Berger, D., R. J. Walters, and W. U. Blanckenhorn. 2014. Experimental evolution for generalists and specialists reveals

multivariate genetic constraints on thermal reaction norms. J. Evol. Biol. 27:1975–1989.

Björklund, M., E. Ranta, V. Kaitala, L. A. Bach, P. Lundberg, and N. Chr. Stenseth. 2009. Quantitative Trait Evolution

and Environmental Change. PLoS ONE 4:e4521.

Botero, C. A., F. J. Weissing, J. Wright, and D. R. Rubenstein. 2015. Evolutionary tipping points in the capacity to adapt

to environmental change. Proc. Natl. Acad. Sci. 112:184–189.

Bubliy, O. A., and V. Loeschcke. 2005. Correlated responses to selection for stress resistance and longevity in a laboratory

population of *Drosophila melanogaster*. J. Evol. Biol. 18:789–803.

Buckley, L. B., and J. G. Kingsolver. 2021. Evolution of Thermal Sensitivity in Changing and Variable Climates. Annu.

Rev. Ecol. Evol. Syst. 52:563–586.

Bürkner, P.-C. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. R J. 10:395.

Chevin, L.-M., and A. A. Hoffmann. 2017. Evolution of phenotypic plasticity in extreme environments. Philos. Trans. R.

Soc. B Biol. Sci. 372:20160138.

Colinet, H., B. J. Sinclair, P. Vernon, and D. Renault. 2015. Insects in Fluctuating Thermal Environments. Annu. Rev.

Entomol. 60:123–140.

Cortés, A. J., F. López-Hernández, and D. Osorio-Rodriguez. 2020. Predicting Thermal Adaptation by Looking Into

Populations' Genomic Past. Front. Genet. 11:564515.

Czesak, M. E., C. W. Fox, and J. B. Wolf. 2006. Experimental Evolution of Phenotypic Plasticity: How Predictive Are

Cross-Environment Genetic Correlations? Am. Nat. 168:323–335.

Deatherage, D. E., J. L. Kepner, A. F. Bennett, R. E. Lenski, and J. E. Barrick. 2017. Specificity of genome evolution in

experimental populations of *Escherichia coli* evolved at different temperatures. Proc. Natl. Acad. Sci. 114:E1904–

E1912.

Deshpande, N., M. R. Wilkins, N. Packer, and H. Nevalainen. 2008. Protein glycosylation pathways in filamentous fungi.

Glycobiology 18:626–637.

DeWitt, T.J. & R.B. Langerhans. 2004. Integrated solutions to environmental heterogeneity: theory of multimoment

reaction norms. In Phenotypic Plasticity. Functional and Conceptual Approaches. T.J. DeWitt & S.M. Scheiner,

Eds.: 98–111. New York: Oxford University Press.

Duun Rohde, P., K. Krag, V. Loeschcke, J. Overgaard, P. Sørensen, and T. Nygaard Kristensen. 2016. A Quantitative

Genomic Approach for Analysis of Fitness and Stress Related Traits in a *Drosophila melanogaster* Model

Population. Int. J. Genomics 2016:1–11.

Ellison, C. E., C. Hall, D. Kowbel, J. Welch, R. B. Brem, N. L. Glass, and J. W. Taylor. 2011. Population genomics and

local adaptation in wild isolates of a model microbial eukaryote. Proc. Natl. Acad. Sci. 108:2831–2836.

Fisher, R. A. 1919. XV.—The correlation between relatives on the supposition of Mendelian inheritance. Earth Env. Sci.

T. R. So. 52:399–433.

Fisher, K. J., and G. I. Lang. 2016. Experimental evolution in fungi: An untapped resource. Fungal Genet. Biol. 94:88–94.

Fu, W., T. D. O'Connor, and J. M. Akey. 2013. Genetic architecture of quantitative traits and complex diseases. Curr.

Opin. Genet. Dev. 23:678–683.

Galagan, J. E., S. E. Calvo, K. A. Borkovich, E. U. Selker, N. D. Read, D. Jaffe, W. FitzHugh, L.-J. Ma, S. Smirnov, S.

Purcell, B. Rehman, T. Elkins, R. Engels, S. Wang, C. B. Nielsen, J. Butler, M. Endrizzi, D. Qui, P. Ianakiev, D.

Bell-Pedersen, M. A. Nelson, M. Werner-Washburne, C. P. Selitrennikoff, J. A. Kinsey, E. L. Braun, A. Zelter,

U. Schulte, G. O. Kothe, G. Jedd, W. Mewes, C. Staben, E. Marcotte, D. Greenberg, A. Roy, K. Foley, J. Naylor,

N. Stange-Thomann, R. Barrett, S. Gnerre, M. Kamal, M. Kamvysselis, E. Mauceli, C. Bielke, S. Rudd, D.

Frishman, S. Krystofova, C. Rasmussen, R. L. Metzenberg, D. D. Perkins, S. Kroken, C. Cogoni, G. Macino, D.

Catcheside, W. Li, R. J. Pratt, S. A. Osmani, C. P. C. DeSouza, L. Glass, M. J. Orbach, J. A. Berglund, R. Voelker,

O. Yarden, M. Plamann, S. Seiler, J. Dunlap, A. Radford, R. Aramayo, D. O. Natvig, L. A. Alex, G. Mannhaupt,

D. J. Ebbole, M. Freitag, I. Paulsen, M. S. Sachs, E. S. Lander, C. Nusbaum, and B. Birren. 2003. The genome

sequence of the filamentous fungus *Neurospora crassa*. Nature 422:859–868.

Gaydos, T. L., N. E. Heckman, M. Kirkpatrick, J. R. Stinchcombe, J. Schmitt, J. Kingsolver, and J. S. Marron. 2013.

Visualizing genetic constraints. Ann. Appl. Stat. 7:860–882.

Georgianna, D. R., A. M. Hawkridge, D. C. Muddiman, and G. A. Payne. 2008. Temperature-Dependent Regulation of

Proteins in *Aspergillus flavus*: Whole Organism Stable Isotope Labeling by Amino Acids. J. Proteome Res.

7:2973–2979.

Gerken, A. R., O. C. Eller, D. A. Hahn, and T. J. Morgan. 2015. Constraints, independence, and evolution of thermal

plasticity: Probing genetic architecture of long- and short-term thermal acclimation. Proc. Natl. Acad. Sci.

112:4399–4404.

Gilchrist, G. W. 1995. Specialists and Generalists in Changing Environments. I. Fitness Landscapes of Thermal Sensitivity.

Am. Nat. 146:252–270.

Hangartner, S., and A. A. Hoffmann. 2016. Evolutionary potential of multiple measures of upper thermal tolerance in

Drosophila melanogaster. Funct. Ecol. 30:442–452.

Hochachka, P. W., and Somero, G. N. 2002. Biochemical Adaptation: Mechanism and Process in Physiological Evolution.

New York: Oxford University Press.

Holder, K. K., and J. J. Bull. 2001. Profiles of Adaptation in Two Similar Viruses. Genetics 159:1393–1404.

Huang, M., X. Liu, Y. Zhou, R. M. Summers, and Z. Zhang. 2019. BLINK: a package for the next level of genome-wide

association studies with both individuals and markers in the millions. GigaScience 8.

Huey, R. B., and P. E. Hertz. 1984. Is a jack-of-all-temperatures a master of none? Evolution 38:441–444.

IPCC 2018: Summary for policymakers. — An IPCC Special Report on the impacts of global warming of 1.5 °C above

pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the

global response to the threat of climate change, sustainable development, and efforts to eradicate poverty, World

Meteorological Organization, Geneva, Switzerland.

Izem, R., and J. G. Kingsolver. 2005. Variation in Continuous Reaction Norms: Quantifying Directions of Biological

Interest. Am. Nat. 166:277–289.

Kapoor, M., and S. S. Roy. 2014. A bird's eye-view of molecular chaperones and some facets of the heat shock response

of *Neurospora crassa*. Annual Review of Plant Pathology 6:1.

Kearney, M. R., A. Matzelle, and B. Helmuth. 2012. Biomechanics meets the ecological niche: the importance of temporal
data resolution. J. Exp. Biol. 215:1422–1424.

Kellermann, V., J. Overgaard, A. A. Hoffmann, C. Fløjgaard, J.-C. Svenning, and V. Loeschcke. 2012. Upper thermal
limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. Proc. Natl.
Acad. Sci. 109:16228–16233.

Ketola, T., V. M. Kellermann, V. Loeschcke, A. López-Sepulcre, and T. N. Kristensen. 2014. Does environmental
robustness play a role in fluctuating environments? Evolution 68:587–594.

Ketola, T., and T. N. Kristensen. 2017. Experimental Approaches for Testing if Tolerance Curves Are Useful for Predicting
Fitness in Fluctuating Environments. Front. Ecol. Evol. 5:129.

Ketola, T., and K. Saarinen. 2015. Experimental evolution in fluctuating environments: tolerance measurements at constant
temperatures incorrectly predict the ability to tolerate fluctuating temperatures. J. Evol. Biol. 28:800–806.

Khan, M. I., Z. Kainat, S. Maqbool, A. Mehwish, S. Ahmad, H. M. Suleman, Z. Mahmood, M. Ali, A. Aziz, A. Rasheed,
and H. Li. 2022. Genome-wide association for heat tolerance at seedling stage in historical spring wheat cultivars.
Front. Plant Sci. 13:972481.

Kingsolver, J. G., S. E. Diamond, and L. B. Buckley. 2013. Heat stress and the fitness consequences of climate change for
terrestrial ectotherms. Funct. Ecol. 27:1415–1423.

Kingsolver, J. G., R. Gomulkiewicz, and P. A. Carter. 2001. Variation, selection and evolution of function-valued traits.
Genetica 112:87–104.

Kingsolver, J. G., G. J. Ragland, and J. G. Shlichta. 2004. Quantitative genetics of continuous reaction norms: thermal
sensitivity of caterpillar growth rates. Evolution 58:1521–1529.

Knies, J. L., R. Izem, K. L. Supler, J. G. Kingsolver, and C. L. Burch. 2006. The Genetic Basis of Thermal Reaction Norm
Evolution in Lab and Natural Phage Populations. PLoS Biol. 4:e201.

Kristensen, T. N., T. Ketola, and I. Kronholm. 2020. Adaptation to environmental stress at different timescales. Ann. N.
Y. Acad. Sci. 1476:5–22.

Kristensen, T. N., J. Overgaard, J. Lassen, A. A. Hoffmann, and C. Sgrò. 2015. Low evolutionary potential for egg-to-adult
viability in *Drosophila melanogaster* at high temperatures. Evolution 69:803–814.

Kristensen, T. N., P. Sørensen, M. Kruhøffer, K. S. Pedersen, and V. Loeschcke. 2005. Genome-Wide Analysis on
Inbreeding Effects on Gene Expression in *Drosophila melanogaster*. Genetics 171:157–167.

Kronholm, I., H. Johannesson, and T. Ketola. 2016. Epigenetic Control of Phenotypic Plasticity in the Filamentous Fungus
Neurospora crassa. G3-Genes Genom. Genet. 6:4009–4022.

Kronholm, I., and T. Ketola. 2018. Effects of acclimation time and epigenetic mechanisms on growth of *Neurospora* in
fluctuating environments. Heredity 121:327–341.

Krystofova, S., and K. A. Borkovich. 2006. The Predicted G-Protein-Coupled Receptor GPR-1 Is Required for Female
Sexual Development in the Multicellular Fungus *Neurospora crassa*. Eukaryot. Cell 5:1503–1516.

Kültz, D. 2005. Molecular and evolutionary basis of the cellular stress response. Annu. Rev. Physiol. 67:225–257.

Kundu, P., A. Nehra, R. Gill, N. Tuteja, and S. S. Gill. 2022. Genomic collation revealed the significance of *Piriformospora*
indica (*Serendipita indica*) PiEF-hand protein in vesicle trafficking and fungal hyphal growth. South Afr. J. Bot.
150:548–564.

Lakin-Thomas, P. 2019. Circadian rhythms, metabolic oscillators, and the target of rapamycin (TOR) pathway: the
Neurospora connection. Curr. Genet. 65:339–349.

Lambros, M., X. Pechuan-Jorge, D. Biro, K. Ye, and A. Bergman. 2021. Emerging adaptive strategies under temperature
fluctuations in a laboratory evolution experiment of *Escherichia coli*. Front. Microbiol. 12:724982.

Latimer, C. A. L., B. R. Foley, and S. F. Chenoweth. 2015. Connecting thermal performance curve variation to the
genotype: a multivariate QTL approach. J. Evol. Biol. 28:155–168.

Lecheta, M. C., D. N. Awde, T. S. O'Leary, L. N. Unfried, N. A. Jacobs, M. H. Whitlock, E. McCabe, B. Powers, K. Bora,
J. S. Waters, H. J. Axen, S. Frietze, B. L. Lockwood, N. M. Teets, and S. H. Cahan. 2020. Integrating GWAS and
Transcriptomics to Identify the Molecular Underpinnings of Thermal Stress Responses in *Drosophila*
melanogaster. Front. Genet. 11:658.

Levins, R. 1968. Evolution in Changing Environments: Some Theoretical Explorations. Princeton, NJ: Princeton
University Press.

Li, L., and K. A. Borkovich. 2006. GPR-4 Is a Predicted G-Protein-Coupled Receptor Required for Carbon Source-
Dependent Asexual Growth and Development in *Neurospora crassa*. Eukaryot. Cell 5:1287–1300.

Logan, M. L., and C. L. Cox. 2020. Genetic Constraints, Transcriptome Plasticity, and the Evolutionary Response to
Climate Change. Front. Genet. 11:538226.

López-Maury, L., S. Marguerat, and J. Bähler. 2008. Tuning gene expression to changing environments: from rapid
responses to evolutionary adaptation. Nat. Rev. Genet. 9:583–593.

Lynch, M., and W. Gabriel. 1987. Environmental tolerance. Am. Nat. 129:283–303.
743

Marshall, K. E., and B. J. Sinclair. 2015. The relative importance of number, duration and intensity of cold stress events in

determining survival and energetics of an overwintering insect. Funct. Ecol. 29:357–366.

McCluskey, K., A. Wiest, and M. Plamann. 2010. The Fungal Genetics Stock Center: a repository for 50 years of fungal

genetics research. J. Biosci. 35:119–126.

McElreath, R. 2015. Statistical rethinking—a Bayesian course with examples in R and Stan. CRC Press. New York.

McLaren, W., L. Gil, S. E. Hunt, H. S. Riat, G. R. S. Ritchie, A. Thormann, P. Flicek, and F. Cunningham. 2016. The

Ensembl Variant Effect Predictor. Genome Biol. 17:122.

Meehl, G. A., T. Karl, D. R. Easterling, S. Changnon, R. Pielke, D. Changnon, J. Evans, P. Y. Groisman, T. R. Knutson,

K. E. Kunkel, L. O. Mearns, C. Parmesan, R. Pulwarty, T. Root, R. T. Sylves, P. Whetton, and F. Zwiers. 2000.

An Introduction to Trends in Extreme Weather and Climate Events: Observations, Socioeconomic Impacts,

Terrestrial Ecological Impacts, and Model Projections. Bull. Am. Meteorol. Soc. 81:413–416. American

Meteorological Society.

Metzenberg, R. L. 2003. Vogel's Medium N salts: avoiding the need for ammonium nitrate. Fungal Genet. Rep. 50:14–14.

Mitchell, K. A., and A. A. Hoffmann. 2010. Thermal ramping rate influences evolutionary potential and species differences

for upper thermal limits in *Drosophila*. Funct. Ecol. 24:694–700.

Moghadam, N. N., K. Sidhu, P. A. M. Summanen, T. Ketola, and I. Kronholm. 2020. Quantitative genetics of temperature

performance curves of *Neurospora crassa*. Evolution 74:1772–1787.

Mohsenzadeh, S., W. Saupe-Thies, G. Steier, T. Schroeder, F. Fracella, P. Ruoff, and L. Rensing. 1998. Temperature

Adaptation of House Keeping and Heat Shock Gene Expression in *Neurospora crassa*. Fungal Genet. Biol. 25:31–

43.

Murren, C. J., J. R. Auld, H. Callahan, C. K. Ghalambor, C. A. Handelsman, M. A. Heskel, J. G. Kingsolver, H. J. Maclean,

J. Masel, H. Maughan, D. W. Pfennig, R. A. Relyea, S. Seiter, E. Snell-Rood, U. K. Steiner, and C. D. Schlichting.

2015. Constraints on the evolution of phenotypic plasticity: limits and costs of phenotype and plasticity. Heredity

115:293–301.

Palma-Guerrero, J., C. R. Hall, D. Kowbel, J. Welch, J. W. Taylor, R. B. Brem, and N. L. Glass. 2013. Genome Wide

Association Identifies Novel Loci Involved in Fungal Communication. PLoS Genet. 9:e1003669.

Pedersen, K. S., T. N. Kristensen, V. Loeschcke, B. O. Petersen, J. Ø. Duus, N. C. Nielsen, and A. Malmendal. 2008.

Metabolomic Signatures of Inbreeding at Benign and Stressful Temperatures in *Drosophila melanogaster*.

Genetics 180:1233–1243.

Pinto, D., M. Duarte, S. Soares, M. Tropschug, and A. Videira. 2008. Identification of all FK506-binding proteins from
Neurospora crassa. Fungal Genet. Biol. 45:1600–1607.

Price, A. L., N. J. Patterson, R. M. Plenge, M. E. Weinblatt, N. A. Shadick, and D. Reich. 2006. Principal components
analysis corrects for stratification in genome-wide association studies. Nat. Genet. 38:904–909.

Pringle, A., and J. W. Taylor. 2002. The fitness of filamentous fungi. Trends Microbiol. 10:474–481.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing,
Vienna, Austria. Available at https://www.R-project.org/.

Riehle, M. M., A. F. Bennett, and A. D. Long. 2001. Genetic architecture of thermal adaptation in *Escherichia coli*. Proc.
Nat. Acad. Sci. 98:525–530.

Rolandi, C., J. R. B. Lighton, G. J. De La Vega, P. E. Schilman, and J. Mensch. 2018. Genetic variation for tolerance to
high temperatures in a population of *Drosophila melanogaster*. Ecol. Evol. 8:10374–10383.

Rowe, L. & D. Houle. 1996. The lek paradox and the capture of genetic variance by condition dependent traits. Proc. R.
Soc. B Biol. Sci. 263:1415–1421.

Ryan, F. J., G. W. Beadle, and E. L. Tatum. 1943. The Tube Method of Measuring the Growth Rate of *Neurospora*. Am.
J. Bot. 784-799.

Räsänen, E., N. N. Moghadam, K. Sidhu, P. A. M. Summanen, H.-R. Littunen, T. Ketola, and I. Kronholm. 2024. Genome-
wide association for loci influencing thermal performance curves in *Neurospora crassa*. bioRxiv.

Schaum, C.-E., A. Buckling, N. Smirnoff, D. J. Studholme, and G. Yvon-Durocher. 2018. Environmental fluctuations
accelerate molecular evolution of thermal tolerance in a marine diatom. Nat. Commun. 9:1719.

Schaum, C.-E., A. Buckling, N. Smirnoff, and G. Yvon-Durocher. 2022. Evolution of thermal tolerance and phenotypic
plasticity under rapid and slow temperature fluctuations. Proc. R. Soc. B Biol. Sci. 289:20220834.

Schulte, P. M., T. M. Healy, and N. A. Fangue. 2011. Thermal Performance Curves, Phenotypic Plasticity, and the Time
Scales of Temperature Exposure. Integr. Comp. Biol. 51:691–702.

Segura, V., B. J. Vilhjálmsson, A. Platt, A. Korte, Ü. Seren, Q. Long, and M. Nordborg. 2012. An efficient multi-locus
mixed-model approach for genome-wide association studies in structured populations. Nat. Genet. 44:825–830.

Sgrò, C. M., and A. A. Hoffmann. 2004. Genetic correlations, tradeoffs and environmental variation. Heredity 93:241–
248.

Shama, L. N. S., M. Campero-Paz, K. M. Wegner, M. De Block, and R. Stoks. 2011. Latitudinal and voltinism
compensation shape thermal reaction norms for growth rate. Mol. Ecol. 20:2929–2941.

Sinclair, B. J., K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, S. Slotsbo, Y. Dong, C. D. G. Harley, D. J.

Marshall, B. S. Helmuth, and R. B. Huey. 2016. Can we predict ectotherm responses to climate change using

thermal performance curves and body temperatures? Ecol. Lett. 19:1372–1385.

Sørensen, J. G., T. N. Kristensen, and V. Loeschcke. 2003. The evolutionary and ecological role of heat shock proteins.

Ecol. Lett. 6:1025–1037.

Sørensen, J. G., M. F. Schou, T. N. Kristensen, and V. Loeschcke. 2016. Thermal fluctuations affect the transcriptome

through mechanisms independent of average temperature. Sci. Rep. 6:30975.

Tobler, R., S. U. Franssen, R. Kofler, P. Orozco-terWengel, V. Nolte, J. Hermisson, and C. Schlötterer. 2014. Massive

habitat-specific genomic response in *D. melanogaster* populations during experimental evolution in hot and cold

environments. Mol. Biol. Evol. 31:364–375.

Urban, J., A. Soulard, A. Huber, S. Lippman, D. Mukhopadhyay, O. Deloche, V. Wanke, D. Anrather, G. Ammerer, H.

Riezman, J. R. Broach, C. De Virgilio, M. N. Hall, and R. Loewith. 2007. Sch9 Is a Major Target of TORC1 in

Saccharomyces cerevisiae. Mol. Cell 26:663–674.

Van Noordwijk, A. J., and G. De Jong. 1986. Acquisition and Allocation of Resources: Their Influence on Variation in

Life History Tactics. Am. Nat. 128:137–142.

Vasseur, D. A., J. P. DeLong, B. Gilbert, H. S. Greig, C. D. G. Harley, K. S. McCann, V. Savage, T. D. Tunney, and M. I.

O'Connor. 2014. Increased temperature variation poses a greater risk to species than climate warming. Proc. R.

Soc. B Biol. Sci. 281:20132612.

Wang, J., and Z. Zhang. 2021. GAPIT Version 3: Boosting Power and Accuracy for Genomic Association and Prediction.

Genomics Proteomics Bioinformatics 19:629–640.

Wang, Y., X. Zheng, G. Li, and X. Wang. 2023. TORC1 Signaling in Fungi: From Yeasts to Filamentous Fungi.

Microorganisms 11:218.

White, B., and D. Woodward. 1995. A simple method for making disposable race tubes. Fungal Genet. Rep. 42:79.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. URL: http://ggplot2.org

Yamahira, K., M. Kawajiri, K. Takeshi, and T. Irie. 2007. Inter- and intrapopulation variation in thermal reaction norms

for growth rate: evolution of latitudinal compensation in ectotherms with a genetic constraint. Evolution 61:1577–

1589.

Yates, A. D., J. Allen, R. M. Amode, A. G. Azov, M. Barba, A. Becerra, J. Bhai, L. I. Campbell, M. Carbajo Martinez, M.

Chakiachvili, K. Chougule, M. Christensen, B. Contreras-Moreira, A. Cuzick, L. Da Rin Fioretto, P. Davis, N. H.

De Silva, S. Diamantakis, S. Dyer, J. Elser, C. V. Filippi, A. Gall, D. Grigoriadis, C. Guijarro-Clarke, P. Gupta,

K. E. Hammond-Kosack, K. L. Howe, P. Jaiswal, V. Kaikala, V. Kumar, S. Kumari, N. Langridge, T. Le, M.

Luypaert, G. L. Maslen, T. Maurel, B. Moore, M. Muffato, A. Mushtaq, G. Naamati, S. Naithani, A. Olson, A.

Parker, M. Paulini, H. Pedro, E. Perry, J. Preece, M. Quinton-Tulloch, F. Rodgers, M. Rosello, M. Ruffier, J.

Seager, V. Sitnik, M. Szpak, J. Tate, M. K. Tello-Ruiz, S. J. Trevanion, M. Urban, D. Ware, S. Wei, G. Williams,

A. Winterbottom, M. Zarowiecki, R. D. Finn, and P. Flicek. 2022. Ensembl Genomes 2022: an expanding genome

resource for non-vertebrates. Nucleic Acids Res. 50:D996–D1003.

Yu, J., J. B. Holland, M. D. McMullen, and E. S. Buckler. 2008. Genetic Design and Statistical Power of Nested Association

Mapping in Maize. Genetics 178:539–551.

Zhao, Y., J. Ding, W. Yuan, J. Huang, W. Huang, Y. Wang, and W. Zheng. 2017. Production of a fungal furocoumarin by

a polyketide synthase gene cluster confers the chemo-resistance of *Neurospora crassa* to the predation by

fungivorous arthropods. Environ. Microbiol. 19:3920–3929.

Zhou, X.-G., P. Yu, C. Dong, C.-X. Yao, Y.-M. Ding, N. Tao, and Z.-W. Zhao. 2016. Proteomic analysis of mycelial

proteins from *Magnaporthe oryzae* under nitrogen starvation. Genet. Mol. Res. 15.

Zou, Y., M. Zhang, J. Qu, and J. Zhang. 2018. iTRAQ-Based Quantitative Proteomic Analysis Reveals Proteomic Changes

in Mycelium of *Pleurotus ostreatus* in Response to Heat Stress and Subsequent Recovery. Front. Microbiol.

9:2368.