Discovery of Novel Long Non-Coding RNAs with potential role in zebrafish brain regeneration.

Surbhi Kohli[1], Dasari Abhilash[1], Hemlata[2], Priyanka P. Srivastava[2], Vishantan Kumar[1] , Shilpi Minocha[2*], Ishaan Gupta[1*]

* corresponding author

1. Department of Biochemical Engineering and Biotechnology, Indian Institute of Technology, Delhi
2. Kusuma School of Biological Sciences, Indian Institute of Technology, Delhi

Abstract

Understanding brain regeneration mechanisms is vital for treating neurological conditions. Zebrafish (*Danio rerio*) are an excellent model due to their genetic similarity to humans and strong regenerative abilities. In this study we identified novel long non-coding RNAs (lncRNAs) in zebrafish brain regeneration post-traumatic brain injury (TBI) samples. RNA sequencing data of the zebrafish telencephalon from the BioStudies database was analyzed for novel long non-coding RNA expression (lncRNA) at control, one day-post-lesion (early wound healing), three days post-lesion (cell proliferation), and 14 days post-lesion (differentiation). We identified 664 potential lncRNAs using HISAT2, StringTie, FEELnc and PhastCon analysis tools. Principal component analysis (PCA) of identified lncRNAs revealed a distinct expression profile at 1-day post-lesion, indicating significant roles in early wound healing.

Weighted Gene Co-expression Network Analysis (WGCNA) identified two modules (brown and turquoise) showing unique expression patterns critical to brain regeneration. Pathway enrichment analysis linked brown module lncRNAs to peptide biosynthesis, cellular amide metabolism, and ribosome biogenesis. In contrast, turquoise module lncRNAs were associated with ion transmembrane transport and cell adhesion pathways. qPCR validation confirmed co-expression patterns of selected lncRNAs and correlated genes, emphasizing their regulatory roles. This study demonstrates that lncRNAs play crucial roles in zebrafish brain regeneration by modulating lncRNA gene expression during the early wound healing stage. These insights offer potential therapeutic applications of lncRNAs in neuro regenerative medicine.

Introduction

In order to effectively treat a variety of neurological conditions and traumas, it is essential to understand the mechanisms behind brain development and injury. With its estimated 100 billion neurons, the human brain is incredibly complex, which presents huge challenges for researchers. For this reason, model organisms that are simple to study but have important structural and functional similarities to the human brain are invaluable.

Zebrafish have shown to be an effective model for researching brain injuries and development. With almost 70% sequence similarity to humans, they are great models for studying brain function, development, and related illnesses. In-depth research on brain morphogenesis can be made possible by the structural parallels between the developing zebrafish brain and the brains of other vertebrates, including humans (Lowery et al. 2009). Zebrafish models have been very helpful in studying neurogenesis from embryonic stages to adulthood. They have shown us how new neurons are created and integrated into existing neural circuits (Schmidt, Strähle, and Scholpp 2013). The spinal cord, forebrain, midbrain, and hindbrain are among the primary structural components that the zebrafish brain shares with the human brain, despite the zebrafish brain's lower level of complexity (Kozol et al. 2016). These characteristics make zebrafish an excellent model for neurodevelopmental research, especially when combined with the abundance of embryos and sophisticated genetic methods for gene editing (Vaz, Hofmeister, and Lindstrand 2019). Furthermore, the transparency of zebrafish embryos allows for real-time observation of developmental processes, organogenesis, and injury responses.

Zebrafish can efficiently repair brain injuries, restoring lost function without any evident scar formation, which contrasts with the limited regenerative capabilities observed in mammals. Compared to the human brain's with 100 billion neurons, the zebrafish brain's manageable size of approximately 100,000 neurons also makes it easier to do in-depth study on neural growth and function (Herculano-Houzel 2009). The ease of use of automated tracking systems in high-throughput trials to evaluate diverse behaviors is made possible by this simplicity, which offers significant insights into the neurological mechanisms that underlie numerous activities like social interactions, learning, and locomotion (Kalueff, Stewart, and Gerlai 2014). Additionally,

their ability to regenerate injured neural tissue, including both neurons and glia, provides valuable insights into the cellular and molecular mechanisms governing the process of regeneration. According to (X. Yu and Li 2011), zebrafish are useful in the research of ischemia brain damage and drug screening because they are easily driven into hypoxia, can have it restored, and can have medications provided directly to the water.

LncRNAs are a group of RNA molecules longer than 200 nucleotides that do not encode proteins. lncRNAs have been connected to a number of human diseases, such as cancer, neurological disorders, and metabolic disorders. They also play critical roles in regulating developmental processes, such as embryogenesis and tissue differentiation (Bhattacharyya et al. 2021; Aliperti, Skonieczna, and Cerase 2021). LncRNAs also play a critical role in neural cell activity during brain development and have been linked to a number of human brain illnesses (Andersen and Lim 2018).

LncRNAs are essential modulators of gene expression in the brain in neurodevelopmental and mental disorders and could be candidates for therapeutic intervention (Aliperti, Skonieczna, and Cerase 2021). Dysregulated long noncoding RNAs (lncRNAs) have been associated with disorders like autism, schizophrenia, and bipolar disorder, indicating their importance in maintaining normal brain development and physiology (Andersen and Lim 2018). Furthermore, lncRNAs are important in acute brain injuries such as ischemic stroke (IS). They are specifically involved in the regulation of gene expression and cellular processes that are essential for the brain's repair response in ischemia-reperfusion injury (IR injury) (Wolska et al. 2021). Specific lncRNAs, including BACE1-AS, NEAT1, MALAT1, and SNHG1, which are linked to Alzheimer's disease and may function as biomarkers or therapeutic targets, have also been linked to neurodegenerative illnesses (Asadi et al. 2021).

The purpose of this work is to discover novel lncRNAs that are expressed in the injured zebrafish brain and to understand their roles. We aim to investigate their roles in brain regeneration, inflammation, and cellular repair mechanisms by defining their expression patterns and regulatory pathways. Understanding these mechanisms could lead to novel therapeutic approaches in neuroregenerative medicine, which could have wider consequences for managing neurodegenerative illnesses and brain injuries in people.

Methods

Figure 1: Schematic of lncRNA discovery pipeline

Data acquisition and sample description

RNA sequencing data were downloaded from the BioStudies database (Sarkans et al. 2018) under the accession number E-MTAB-11163. The data were part of a study performed by (Demirci et al. 2022), which investigated the gene expression patterns in the telencephalon region of the zebrafish brain following Traumatic Brain Injury (TBI). The dataset comprises 18 samples, categorised into four groups: control (5 samples), 1-day post-lesion (5 samples): early wound healing stage, 3 days post-lesion (4 samples): cell proliferation stage, and 14 days post-lesion (4 samples): differentiation stage.

Transcriptome Assembly and Mapping

Fastq files were downloaded from the biostudies database with Accession ID E-MTAB-11163. Raw reads were cleaned using Trimmomatic (v0.38) (Bolger, Lohse, and Usadel 2014) to remove adapters and low-quality bases using default parameters. The cleaned reads were then mapped to the zebrafish reference genome (GRCz11) using HISAT2 (v2.2.1) (Kim, Langmead, and Salzberg 2015) by disabling template length adjustment for RNA-seq reads. StringTie (v2.2.1) (Pertea et al. 2015) was used for reference-guided transcript assembly with parameters like reverse stranded library and leveraging HISAT2's output to construct transcriptomes for each sample, followed by merging all samples.

Identification and Classification of lncRNAs

To annotate and classify lncRNAs from our RNA-seq data, we employed FEELnc (FlExible Extraction of LncRNAs) (v0.2.1) (Wucher et al. 2017), an alignment-free program that differentiates mRNAs from lncRNAs using a Random Forest model. This model is trained on features such as multi k-mer frequencies and relaxed open reading frames (ORFs). FEELnc has been rigorously benchmarked against leading tools and demonstrates high performance, enabling precise and reliable lncRNA identification and classification in our study.

Using Feelnc, we filtered the spurious transcripts and exons, including monoexonic regions, using FEELnc_filter.pl. After filtering, we removed the transcripts with coding potential using FEELnc_codpot.pl. Then, the lncRNA was classified based on its proximity to the annotated genes in the genome.

Conservation Analysis

The conservation of identified lncRNA candidates across multiple species was assessed using PhastCons (V1.6) (Siepel and Haussler 2004), which employs a phylogenetic hidden Markov model to compute conservation scores. Due to the unavailability of Multiple Sequence Alignment (MSA) data for the new genome, the transcript data was first converted to version 7 of the zebrafish genome using BedFile Liftover.

Weighted Gene Co-expression Network Analysis (WGCNA)

Weighted Gene Co-expression Network Analysis was performed using the WGCNA package (1.72-5) (Langfelder and Horvath 2008) to identify clusters of co-expressed genes and lncRNAs. Gene expression data was preprocessed and TPM normalized prior to network construction. A soft-thresholding power of 30 was selected based on the scale-free topology criterion, and a signed network was used to emphasize positive correlations. The network was constructed with a deep split level of 2, a minimum module size of 30 genes, and a maximum block size of 4000 genes. Modules were defined with a reassign threshold of 0 and a merge cut height of 0.25, allowing closely related modules to be merged. This analysis enabled the identification of key gene networks potentially involved in the biological processes of interest.

Gene Proximity Analysis

lncRNAs present in the brown and turquoise modules of WGCNA were further shortlisted based on their genomic location. The gene proximity analysis was performed in Python v3.11.0. Known gene coordinates were extracted from the GTF file (GRCz11.109.gtf) and lncRNA coordinates were extracted from the Stringtie results. For each lncRNA, we checked if there was any known gene lying within 2.5 kb upstream or downstream and transcripts having with coding within 2,5 kb region were removed from consideration

Co-expression Analysis

The modules identified by WGCNA were subjected to further co-expression analysis to validate and refine the associations between lncRNAs and coding genes. This involved using statistical correlation tests and correlation of greater than 0.9 to ensure these relationships' robustness and

biological significance, focusing on those lncRNAs that consistently appeared central within the significant modules.

Pathway Enrichment Analysis

Pathway enrichment analysis was conducted to explore the potential biological functions and pathways associated with the co-expressed lncRNA clusters using the cluster profiler R package (G. Yu et al. 2012). This analysis helped identify enriched biological themes and pathways, that may be crucially regulated by or associated with the identified lncRNAs, thus providing insights into their functional roles in zebrafish development and physiology. Gene ontology enrichment of biological processes and KEGG pathway enrichment are performed with a P-value less than 0.05, and all the expressed genes are chosen as background for enrichment analysis.

Zebrafish maintenance and induction of Stab wound injury

The zebrafish were maintained at 25–28°C with a set photoperiod of 14 hours of light and 10 hours of darkness. Five to eight-month-old fish were utilised for the experiment. To investigate brain regeneration in zebrafish, a telencephalic stab wound injury was induced, and the zebrafish were momentarily anesthetized with MS222 (ethyl 3-aminobenzoate methanesulfonic acid). A 30G needle was inserted perpendicularly to the skull on the right hemisphere of the telencephalon to cause a telencephalic stab-wound injury. Thereafter, they were returned to the system water to heal. When evaluating the regeneration activity, the contralateral left hemisphere serves as the control.

RNA isolation, cDNA synthesis, and qRT-PCR

The regenerating brains at 1dpl (day post-lesion) were harvested, and the injured telencephalon were collected (5 pooled per sample). The uninjured telencephalon (5 pooled per sample) from the same brain were collected as control. RNA isolation was done using the RNeasy Lipid Tissue Mini Kit (Cat: 1023539) according to the manufacturer's protocol. An equal amount of RNA was used to synthesise cDNA using Takara PrimeScript 1st strand cDNA Synthesis Kit (Cat: 6110A) according to the manufacturer's protocol. An equal amount of cDNA was used for qRT-PCR using gene-specific primers (*MSTR 23642.1, MSTR 39546.3, vegaa, nsfl1c, rac3,* and *mdm4*) and *gapdh* (F- GATACACGGAGCACCAGGTT, R-GCCATCAGGTCACATACACG) as housekeeping gene using iTaq Universal SYBR Green Supermix (Biorad, Cat: 1725121) on a

Thermo Cycler (BIO-RAD CFX96). Two biological replicates and two technical replicates of each sample were used, and the data was analysed using $\Delta\Delta$ Ct method, and the fold change was calculated using the $2^{-\Delta\Delta Ct}$ method.

Results

RNA sequencing retrieved from the BioStudies database consisted of a total 18 samples. These were categorized based on the timeline post-traumatic brain injury (TBI) in zebrafish, an approach inspired by (Demirci et al. 2022). The samples included control (5 samples), 1 day post-lesion (5 samples), 3 days post-lesion (4 samples), and 14 days post-lesion (4 samples) (Figure 2). This classification was instrumental in investigating the dynamic changes in gene expression associated with different stages of brain regeneration.

Figure 2: Schematic diagram illustrating the experimental design for assessing gene expression changes post-brain injury in zebrafish. The study included four groups: control, 1 day post-lesion (initial wound healing), 3 days post-lesion (cell proliferation), and 14 days post-lesion (differentiation). Each time point post-lesion represents a critical stage in the brain's regeneration process.

Transcriptome Assembly and Mapping

The initial quality assessment of our raw sequencing data, conducted using FastQC, demonstrated high-quality metrics across all samples, with an average Phred score of 35. After using Trimmomatic to remove adapters and low-quality bases, we aligned the cleaned data to the zebrafish reference genome (GRCz11) via HISAT2. This alignment achieved a high mapping efficiency, with an average of 88% of reads aligning successfully. Transcriptome assembly with StringTie integrated these alignments, identifying roughly 119,757 transcripts, including known and novel isoforms. To enhance data relevance, we excluded transcripts derived from unplaced

zebrafish genome scaffolds, focusing only on those from defined chromosomes. This refinement reduced the number of considered transcripts to 100,408 (Figure 1).

Identification of potential LncRNA candidates

We utilised the FEELnc tool to categorise and analyse the transcripts from our transcriptome assembly, classifying them into long non-coding RNAs (lncRNAs) and mRNAs. The FEELnc analysis successfully identified 11,281 transcripts as potential lncRNAs (Figure 1), which were characterized by their lack of coding potential, specific multi k-mer frequencies, and open reading frame (ORF) characteristics.

Supplementary Figure 1: **PhastCons Score distribution for various transcripts**

To assess the evolutionary significance of these lncRNAs, we utilized the PhastCons tool, which applies a phylogenetic hidden Markov model across various vertebrate genomes. Our analysis showed that while the majority of lncRNAs displayed low or zero conservation, mRNAs demonstrated higher conservation scores (Supplementary Fig 1). This aligns with the generally accepted understanding that lncRNA exhibits low conservation. Given the typically low

conservation of lncRNAs, we hypothesized that those with any degree of conservation might play significant biological roles. Consequently, we focused on lncRNAs that exhibited non-zero conservation scores, narrowing our field from 11,281 to 6,600 transcripts. From this subset, we further selected the top 10% based on conservation scores, identifying 664 lncRNAs (Figure 1) with notably high conservation (over 88%). This select group of lncRNAs, due to their substantial conservation, suggests potential evolutionary importance and functional relevance, making them prime candidates for further in-depth functional analyses.

We assessed the expression profiles of identified lncRNA candidates across various samples using principal component analysis (PCA) of normalized lncRNA counts. The PCA results showed that samples from 1 day post-lesion (1dpl) distinctly clustered apart from other stages, which were closer to control samples (Figure 3). This shows the novel lncRNAs are highly active immediately after injury and slowly diminished during the recovery process. This pattern was corroborated by a sample-to-sample distance heatmap (Figure 4), where 1dpl samples formed a separate cluster. These observations suggest that the expression of these lncRNAs is particularly altered during the initial stage of wound healing, possibly playing critical roles in the initial stage of the brain regeneration process.

We performed Weighted Gene Co-expression Network Analysis (WGCNA) to pinpoint clusters of highly coexpressed lncRNAs with distinct sample-specific expression patterns. This approach aimed to prioritize lncRNAs potentially playing regulatory roles in brain regeneration. Our analysis identified six modules (Supplementary Figure 2) with unique expression profiles across the brain injury recovery timeline (Figure 5), with particular attention to two modules—brown and turquoise—that demonstrated unique and consistent patterns at the 1-day post-lesion (1dpl) stage. The brown module showed an increase in expression, while the turquoise module displayed decreased expression at 1dpl (Figure 6). We identified 385 lncRNAs in the turquoise module and 39 in the brown module. To ensure the accuracy of our functional inference, we implemented stringent criteria, excluding any lncRNAs with coding genes within a 2.5 kb region to avoid potential interference from nearby coding genes. This filtration left us with 73 lncRNAs in the turquoise module and 9 in the brown module, isolating those with potentially independent functional roles.

Figure 3: sample to sample lncRNA correlation heatmap

Figure 4: **Heatmap displaying sample distances based on lncRNA expression profiles.** One day post lesion samples lncRNA expression profile is different from other samples.

Supplementary Figure 2: Dendrogram of WGCNA Clusters

Figure 5: **Heatmap showing the correlation of expression profiles among lncRNA clusters.**
Brown and turquoise modules show consistently opposite expression profiles.

Figure 6: **Expression profiles of the individual long non-coding RNAs in the brown and turquoise co-expressed cluster.** Y-axis indicates normalized gene expression, and X-axis indicates samples that are labelled according to the treatment.

Functional significance of identified lncRNAs

To further explore the biological significance of these identified lncRNAs, we extended our analysis by correlating these lncRNAs with expressed known genes. We filtered genes highly that showed a high correlation (cutoff > |0.9|) with the lncRNAs, and conducted gene ontology (GO) and KEGG pathway enrichment analyses, aiming to elucidate the biological processes and pathways potentially influenced by these lncRNAs during the initial wound healing stage of brain regeneration.

Metabolic processes related to RNA splicing and peptide biosynthetic processes are enriched (Figure 7), showing that novel lncRNA-related co-expressed genes are involved in these processes, and they could be involved in relation to these RNA and protein metabolic processes.

Figure 7: **Emap plot of brown module GO enrichment of biological processes with a positive 0.90 correlation**.

Peptide biosynthetic process, cellular amide metabolic process, and cellular macromolecule biosynthetic process. These pathways are interconnected, and are involved in producing the building blocks of life.

ncRNA processing, rRNA processing, and mRNA splicing via spliceosome, highlighting their roles in production of transcripts which eventually translated into proteins. ribonucleoprotein complex biogenesis and ribosome biogenesis are main pathways that ensure the construction and maintenance of ribosomes, which are the molecular machines that drive protein synthesis.

Figure 8: **Dot plot of KEGG pathway enrichment analysis for genes in the brown module, using genes with positive correlations (r > 0.90)**

Figure 9: **Emap plot of turquoise module GO enrichment of biological processes with a positive 0.90 correlation.**

Regulation of ion transmembrane transport, cation transmembrane transport, and potassium ion transmembrane transport are enriched. These processes are critical for maintaining cellular homeostasis and membrane potential, which are essential for neuron firing and muscle contraction.

The Ephrin receptor signaling pathway and the transmembrane receptor protein tyrosine kinase signaling pathway are the signaling mechanisms essential for cells to communicate and respond to their environment, coordinating growth, differentiation, and responses to external stimuli.

Pathways like cell-cell adhesion via plasma-membrane adhesion molecules and cell adhesion pathways ensure that cells stick together, forming tissues and maintaining the structural integrity of neurons in the brain.

Figure 10: **Dot plot of KEGG pathway enrichment analysis for genes in the turquoise module, using genes with positive correlations (r > 0.90)**

Module	Before Filtering	After Filtering
Turquoise	385	73
Brown	39	9

Table 1: Gene proximity analysis filtering table

Figure 11: **Relative lncRNA and gene expression level in the pooled five hemispheres of telencephalon during the early wound injury at 1 dpi. Error bars represent ± standard error of the mean (SEM, n = 2)**. (A) MSTR 23642.1 lncRNA expression in combination with vegaa, rac3a, and nsfl1c genes and (B) MSTR 39546.3 lncRNA expression in combination with vagaa and mdm4 genes.

We chose 2 lncRNAs, one from brown (MSTRG 395546.3) and another from turquoise (MSTRG 23642.1) modules and chose specific gene pairs to perform experimental validation of co-expression using quantitative PCR. Since brown and turquoise show opposite expression pattern in 1-dpl we chose gene pairs having positive correlation with lncRNA from brown module and negatively correlated genes with lncRNA of turquoise module. qPCR validation of lncRNA and gene pairs were performed. qPCR confirmed the in silico results that lncRNA in the MSTRG 23642 (brown module) positively correlate with genes vegfaa, rac3a and nsfl1c and genes vegfaa and mdm4 negatively correlate with lncRNA MSTRG 39546.3 (turquoise module). The vascular endothelial growth factor Aa (vegfaa) gene is central to the angiogenesis process (Marín-Juez et al. 2016). Angiogenesis, activated within a few days post-cerebral ischemia,

plays a crucial role in neuronal remodeling and functional recovery. It provides guidance to sprouting axons and enhances the proliferation and differentiation of neural stem cells (Wang et al. 2007; Kanazawa et al. 2017; Ruan et al. 2015; Hatakeyama, Ninomiya, and Kanazawa 2020). Demirci et al. also observed an upregulation of vegfaa specifically at 1 day post-lesion (1 dpl), suggesting that injury triggers rapid angiogenic sprouting during early brain regeneration (Demirci). The negative correlation between lncRNAs in the turquoise module and vegfaa indicates that these lncRNAs might repress or inhibit vegfaa expression. The downregulation of these lncRNAs at 1 dpl may relieve their inhibitory effect on vegfaa, allowing for the necessary upregulation of vegfaa to support angiogenesis and neural regeneration processes. Additionally, the gene rac3a, associated with heart regeneration pathways, is also negatively correlated with turquoise module lncRNAs MSTRG 23642.1.

On the other hand The positive correlation between lncRNAs in the brown module and vegfaa suggests that these lncRNAs might support the upregulation of vegfaa, promoting angiogenesis and facilitating the initial stages of brain regeneration. These lncRNAs could be acting as enhancers or stabilizers of vegfaa expression, thereby contributing to the rapid angiogenic response observed at 1 dpl. Mdm4 also shows positive correlation with brown lncRNA MSTRG 23642.1. The MDM4/MDM2-p53-IGF1 axis, which controls axonal regeneration, sprouting, and functional recovery after CNS injury (Joshi et al. 2015), shows a positive correlation with lncRNAs in the brown module, suggesting that these lncRNAs may play a role in these critical regenerative processes as well.

Thus our findings suggest a dual regulatory mechanism where lncRNAs in the brown module promote the expression of genes participating in brain regeneration processes, while lncRNAs in the turquoise module inhibit genes involved in regeneration and recovery of brain

Discussion

Building on a recent study by Demirci et al., 2022, on gene expression during brain regeneration post-injury in zebrafish, our research investigated the regulatory role of long non-coding RNAs(lncRNAs) in this process. Demirci et al.'s work provided crucial insights into the dynamic gene expression changes across three key stages of brain regeneration. They discovered that the most significant changes occurred in the initial wound healing stage (one-day post-lesion), with notable similarities between these expression patterns and those observed in brain cancer.

We utilised their published RNA-seq data to identify and characterise lncRNAs that might influence these expression changes. The study of lncRNAs is vital as these molecules play key roles in regulating gene expression, and their dysregulation has been implicated in various pathologies, including cancer and potentially in the mechanisms of neuronal repair and regeneration (Hezroni, Perry, and Ulitsky 2019; Jiang et al. 2019).

The raw RNA seq was first refined to ensure quality and accuracy, followed by alignment using HISAT2 and transcriptome assembly using StringTie. Out of the 100,408 transcripts assembled by StringTie, FEELnc identified 11,281 potential lncRNAs based on coding potential and other structural features. To further refine our selection, we applied PhastCons conservation scores, which reduced the number of potential lncRNAs to 664 transcripts. Given that lncRNAs typically exhibit low conservation (Johnsson et al. 2014) across species, we prioritized those with high PhastCons scores to ensure the selection of functionally relevant lncRNAs.

The PCA plot and heatmap analysis of the 664 lncRNA candidates distinctly demonstrated that samples from the 1-day post-lesion (1dpl) stage clustered separately from other stages. Interestingly, while Demirci et al. showed distinct gene expression profiles at both the 1-day (initial wound healing) and 3-day (cell proliferation) stages, with the 14-day profile mostly resembling controls while our study observed that lncRNA expression profiles exhibited a distinct pattern primarily at the 1-day post-injury stage. This suggests that the expression patterns of identified lncRNAs are markedly distinct during the initial stage of wound healing, potentially playing important roles in initial wound healing roles. The significant divergence in lncRNA

expression specifically at 1dpl underscores their possible involvement in early regulatory mechanisms critical to initiating and guiding the regenerative processes.

To ascertain the biological functions of these lncRNA candidates, we performed a WGCNA analysis. WGCNA revealed six modules with distinct expression profiles over the brain injury recovery timeline. Notably, two modules (brown and turquoise) exhibited unique expression patterns specific to the 1-day post-lesion (1dpl) stage, with the brown module showing increased expression and the turquoise module showing decreased expression at this early stage. This pattern aligns with our previous observations of significant lncRNA activity during the initial wound-healing phase. We extended our analysis by correlating these lncRNAs with known genes and conducted gene ontology (GO) and pathway enrichment analysis of genes highly correlated (cutoff > |0.9|) with these lncRNAs. The analysis revealed that the peptide biosynthetic process, cellular amide metabolic process, and cellular macromolecule biosynthetic process are interconnected pathways involved in producing the building blocks of life.

Further, the GO analysis of the brown module revealed that many correlated genes are involved in the production of transcripts, which are eventually translated into proteins, and the pathways that ensure the construction and maintenance of ribosomes, which are the molecular machines that drive protein synthesis.

The GO analysis of the turquoise module revealed the processes that are critical for maintaining cellular homeostasis and membrane potential, which are essential for neuron firing and muscle contraction, the signaling mechanisms essential for cells to communicate and respond to their environment, coordinating growth, differentiation, and responses to external stimuli and the pathways ensure that cells stick together, forming tissues and maintaining the structural integrity of neurons in the brain.

Moreover, we selected two lncRNA targets and their highly correlated gene pairs, showing both negative and positive correlations, for experimental validation using qPCR. The qPCR results confirmed the expression patterns observed in our RNA-seq analysis, providing further support for the regulatory role of these lncRNAs in brain regeneration. These experimentally validated lncRNAs and their correlated genes suggest specific regulatory mechanisms where lncRNAs

might act as key modulators of gene expression, either enhancing or suppressing their correlated targets during the early stages of wound healing.

Our findings suggest that specific lncRNAs are intricately involved in the early stages of brain regeneration in zebrafish, potentially regulating key genes and pathways that drive the regenerative process. These insights not only enhance our understanding of lncRNA function in neuronal repair but also provide a foundation for future research aimed at leveraging lncRNAs as therapeutic targets for brain.

Funding

IG was supported by by the funds from Department of biotechnology Government of Indiathrough Ramalingasami fellowship ST/HRD/35/02/200 and Intramural MFIRP grant by IIT Delhi MI02552G to Ishaan Gupta. This research was also supported by the Council of Scientific & Industrial Research, Ministry of science and technology, Government of India through CSIR-JRF fellowship 09/086(1458)/2020-EMR-I to Dasari Abhilash. We are grateful for their support and funding, which enabled us to conduct this study.

SM was supported by a Start-Up Research Grant from the Science and Engineering Board (SRG/2021/000341), a Ramalingaswami re-entry fellowship from the Department of Biotechnology (BT/RLF/Re-entry/70/217) and IFCPAR/CEFIPRA (Indo-French Centre for Promotion of Advanced Research/Centre Franco-Indien pour la Promotion de la Recherche Avancée) grant no. 6503-J.

References

Aliperti, Vincenza, Justyna Skonieczna, and Andrea Cerase. 2021. "Long Non-Coding RNA (lncRNA) Roles in Cell Biology, Neurodevelopment and Neurological Disorders." *Non-Coding RNA* 7 (2). https://doi.org/10.3390/ncrna7020036.

Andersen, Rebecca E., and Daniel A. Lim. 2018. "Forging Our Understanding of lncRNAs in the Brain." *Cell and Tissue Research* 371 (1): 55–71.

Asadi, Mohammad Reza, Mehdi Hassani, Shiva Kiani, Hani Sabaie, Marziyeh Sadat Moslehian, Mohammad Kazemi, Soudeh Ghafouri-Fard, Mohammad Taheri, and Maryam Rezazadeh. 2021. "The Perspective of Dysregulated LncRNAs in Alzheimer's Disease: A Systematic Scoping Review." *Frontiers in Aging Neuroscience* 13 (September): 709568.

Bhattacharyya, Nirjhar, Vedansh Pandey, Malini Bhattacharyya, and Abhijit Dey. 2021. "Regulatory Role of Long Non Coding RNAs (lncRNAs) in Neurological Disorders: From Novel Biomarkers to Promising Therapeutic Strategies." *Asian Journal of Pharmaceutical Sciences* 16 (5): 533–50.

Bolger, Anthony M., Marc Lohse, and Bjoern Usadel. 2014. "Trimmomatic: A Flexible Trimmer for Illumina Sequence Data." *Bioinformatics* 30 (15): 2114–20.

Demirci, Yeliz, Guillaume Heger, Esra Katkat, Irene Papatheodorou, Alvis Brazma, and Gunes Ozhan. 2022. "Brain Regeneration Resembles Brain Cancer at Its Early Wound Healing Stage and Diverges From Cancer Later at Its Proliferation and Differentiation Stages." *Frontiers in Cell and Developmental Biology* 10 (February): 813314.

Hatakeyama, Masahiro, Itaru Ninomiya, and Masato Kanazawa. 2020. "Angiogenesis and Neuronal Remodeling after Ischemic Stroke." *Neural Regeneration Research* 15 (1): 16–19.

Herculano-Houzel, Suzana. 2009. "The Human Brain in Numbers: A Linearly Scaled-up Primate Brain." *Frontiers in Human Neuroscience* 3 (November): 31.

Hezroni, Hadas, Rotem Ben Tov Perry, and Igor Ulitsky. 2019. "Long Noncoding RNAs in Development and Regeneration of the Neural Lineage." *Cold Spring Harbor Symposia on Quantitative Biology* 84: 165–77.

Jiang, Ming-Chun, Jiao-Jiao Ni, Wen-Yu Cui, Bo-Ya Wang, and Wei Zhuo. 2019. "Emerging Roles of lncRNA in Cancer and Therapeutic Opportunities." *American Journal of Cancer Research* 9 (7): 1354–66.

Johnsson, Per, Leonard Lipovich, Dan Grandér, and Kevin V. Morris. 2014. "Evolutionary Conservation of Long Non-Coding RNAs; Sequence, Structure, Function." *Biochimica et Biophysica Acta* 1840 (3): 1063–71.

Joshi, Yashashree, Marília Grando Sória, Giorgia Quadrato, Gizem Inak, Luming Zhou, Arnau Hervera, Khizr I. Rathore, et al. 2015. "The MDM4/MDM2-p53-IGF1 Axis Controls Axonal Regeneration, Sprouting and Functional Recovery after CNS Injury." *Brain: A Journal of Neurology* 138 (Pt 7): 1843–62.

Kalueff, Allan V., Adam Michael Stewart, and Robert Gerlai. 2014. "Zebrafish as an Emerging Model for Studying Complex Brain Disorders." *Trends in Pharmacological Sciences* 35 (2): 63–75.

Kanazawa, Masato, Minami Miura, Masafumi Toriyabe, Misaki Koyama, Masahiro Hatakeyama, Masanori Ishikawa, Takashi Nakajima, et al. 2017. "Microglia Preconditioned by Oxygen-Glucose Deprivation Promote Functional Recovery in Ischemic Rats." *Scientific Reports* 7 (February): 42582.

Kim, Daehwan, Ben Langmead, and Steven L. Salzberg. 2015. "HISAT: A Fast Spliced Aligner

with Low Memory Requirements." *Nature Methods* 12 (4): 357–60.

Kozol, Robert A., Alexander J. Abrams, David M. James, Elena Buglo, Qing Yan, and Julia E. Dallman. 2016. "Function Over Form: Modeling Groups of Inherited Neurological Conditions in Zebrafish." *Frontiers in Molecular Neuroscience* 9 (July): 55.

Langfelder, Peter, and Steve Horvath. 2008. "WGCNA: An R Package for Weighted Correlation Network Analysis." *BMC Bioinformatics* 9 (December): 559.

Lowery, Laura Anne, Gianluca De Rienzo, Jennifer H. Gutzman, and Hazel Sive. 2009. "Characterization and Classification of Zebrafish Brain Morphology Mutants." *Anatomical Record* 292 (1): 94–106.

Marín-Juez, Rubén, Michele Marass, Sebastien Gauvrit, Andrea Rossi, Shih-Lei Lai, Stefan C. Materna, Brian L. Black, and Didier Y. R. Stainier. 2016. "Fast Revascularization of the Injured Area Is Essential to Support Zebrafish Heart Regeneration." *Proceedings of the National Academy of Sciences of the United States of America* 113 (40): 11237–42.

Pertea, Mihaela, Geo M. Pertea, Corina M. Antonescu, Tsung-Cheng Chang, Joshua T. Mendell, and Steven L. Salzberg. 2015. "StringTie Enables Improved Reconstruction of a Transcriptome from RNA-Seq Reads." *Nature Biotechnology* 33 (3): 290–95.

Ruan, Linhui, Brian Wang, Qichuan ZhuGe, and Kunlin Jin. 2015. "Coupling of Neurogenesis and Angiogenesis after Ischemic Stroke." *Brain Research* 1623 (October): 166–73.

Sarkans, Ugis, Mikhail Gostev, Awais Athar, Ehsan Behrangi, Olga Melnichuk, Ahmed Ali, Jasmine Minguet, et al. 2018. "The BioStudies Database-One Stop Shop for All Data Supporting a Life Sciences Study." *Nucleic Acids Research* 46 (D1): D1266–70.

Schmidt, Rebecca, Uwe Strähle, and Steffen Scholpp. 2013. "Neurogenesis in Zebrafish - from Embryo to Adult." *Neural Development* 8 (February): 3.

Siepel, Adam, and David Haussler. 2004. "Computational Identification of Evolutionarily Conserved Exons." In *Proceedings of the Eighth Annual International Conference on Research in Computational Molecular Biology*, 177–86. RECOMB '04. New York, NY, USA: Association for Computing Machinery.

Vaz, Raquel, Wolfgang Hofmeister, and Anna Lindstrand. 2019. "Zebrafish Models of Neurodevelopmental Disorders: Limitations and Benefits of Current Tools and Techniques." *International Journal of Molecular Sciences* 20 (6). https://doi.org/10.3390/ijms20061296.

Wang, Yaoming, Kunlin Jin, Xiao Ou Mao, Lin Xie, Surita Banwait, Hugo H. Marti, and David A. Greenberg. 2007. "VEGF-Overexpressing Transgenic Mice Show Enhanced Post-Ischemic Neurogenesis and Neuromigration." *Journal of Neuroscience Research* 85 (4): 740–47.

Wolska, Marta, Joanna Jarosz-Popek, Eva Junger, Zofia Wicik, Tahmina Porshoor, Lucia Sharif, Pamela Czajka, et al. 2021. "Long Non-Coding RNAs as Promising Therapeutic Approach in Ischemic Stroke: A Comprehensive Review." *Molecular Neurobiology* 58 (4): 1664–82.

Wucher, Valentin, Fabrice Legeai, Benoît Hédan, Guillaume Rizk, Lætitia Lagoutte, Tosso Leeb, Vidhya Jagannathan, et al. 2017. "FEELnc: A Tool for Long Non-Coding RNA Annotation and Its Application to the Dog Transcriptome." *Nucleic Acids Research* 45 (8): e57.

Yu, Guangchuang, Li-Gen Wang, Yanyan Han, and Qing-Yu He. 2012. "clusterProfiler: An R Package for Comparing Biological Themes among Gene Clusters." *Omics: A Journal of Integrative Biology* 16 (5): 284–87.

Yu, Xinge, and Yang V. Li. 2011. "Zebrafish as an Alternative Model for Hypoxic-Ischemic Brain Damage." *International Journal of Physiology, Pathophysiology and Pharmacology* 3 (2): 88–96.