A map of canine sequence variation relative to a Greenland wolf outgroup

Anthony K. Nguyen[1,2,*], Peter Z. Schall[1,*], Jeffrey M. Kidd[1,2,†]

[1]Department of Human Genetics, University of Michigan, Ann Arbor, Michigan USA
[2]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, Michigan USA

*A.K.N and P.Z.S. contributed equally to this study

[†]Author For Correspondence: Jeffrey M. Kidd, Department of Human Genetics, University of Michigan, Ann Arbor, Michigan USA. 734-647-3149, jmkidd@umich.edu

Abstract

For over 15 years, canine genetics research relied on a reference assembly from a Boxer breed dog named Tasha (i.e., canFam3.1). Recent advances in long-read sequencing and genome assembly have led to the development of numerous high-quality assemblies from diverse canines. These assemblies represent notable improvements in completeness, contiguity, and the representation of gene promoters and gene models. Although genome graph and pan-genome approaches have promise, most genetic analyses in canines rely upon the mapping of Illumina sequencing reads to a single reference. The Dog10K consortium, and others, have generated deep catalogs of genetic variation through an alignment of Illumina sequencing reads to a reference genome obtained from a German Shepherd Dog named Mischka (i.e., canFam4, UU_Cfam_GSD_1.0). However, alignment to a breed-derived genome may introduce bias in genotype calling across samples. Since the use of an outgroup reference genome may remove this effect, we have reprocessed 1,929 samples analyzed by the Dog10K consortium using a Greenland wolf (mCanLor1.2) as the reference. We efficiently performed remapping and variant calling using a GPU-implementation of common analysis tools. The resulting call set removes the variability in genetic differences seen across samples while maintaining general patterns of breed relationships. Using this sequence data, we inferred the history of population sizes and found that village dog populations experienced a 9-13 fold reduction in historic effective population size relative to wolves.

Key words
Canine, wolf, reference genome

Introduction

For almost two decades the canine genome research community relied on a reference genome derived from a Boxer breed dog named Tasha (Lindblad-Toh et al. 2005). This genome was assembled using a Sanger capillary sequencing whole genome shotgun approach coupled with limited finishing of selected BAC clones. The Boxer-derived genome served as a reference for gene annotation, for the discovery of genetic variation across canines, and the development of canine SNP genotyping arrays (Vaysse et al. 2011). The assembly was further improved through targeted finishing and gene annotation efforts (Hoeppner et al. 2014). The resulting genome, known as canFam3.1 based on its description in the UCSC Genome Browser, became the standard reference for canine genomics (Nassar et al. 2023). With the advent of routine whole genome sequencing using Illumina short-read technology, an increasing number of canine genomes were sequenced and aligned to this Boxer-derived reference (Jagannathan et al. 2019; Plassais et al. 2019). This included ancient DNA data from canines, which involves sequencing of short, damaged DNA fragments, which has refined the understanding of dog origins (Bergstrom et al. 2020; Bergstrom et al. 2022; Botigue et al. 2017; Frantz et al. 2016; Ni Leathlobhair et al. 2018; Ramos-Madrigal et al. 2021; Skoglund et al. 2015).

Although bias due to the ascertainment of SNPs that are genotyped on arrays is well known (Clark et al. 2005; Lachance and Tishkoff 2013), it is not clear how reliance on a single reference may affect genetic variation studies based on sequencing. A systematic bias toward reporting the reference allele has been reported previously, particularly when analyzing short-read data associated with ancient DNA (Gunther and Nettelblad 2019; Martiniano et al. 2020; Orlando et al. 2013) or allele-specific RNA-sequencing analysis (Stevenson et al. 2013). In 2017 Gopalakrishnan and colleagues reported the assembly of the genome of a Swedish wolf using Illumina short-read sequencing data (Gopalakrishnan et al. 2017). Although the resulting assembly was highly fragmented, with a contig N_{50} of 94 kbp, it captured most of the unique sequence present in canines. Reanalysis of existing dog and wolf short-read data using this reference revealed patterns of variation highly similar to those found when using the Boxer-derived genome reference. However, the authors noted that dogs showed a larger estimated heterozygosity when aligned to the canFam3.1 assembly as opposed to the Swedish wolf assembly. The contribution of assembly quality to this metric is unclear.

Soon after the release of the fragmented Swedish wolf genome, the development of long-read sequencing technologies revolutionized genomics. This led to the release of multiple high-quality reference assemblies from canines, including several different breed dogs (Edwards et al. 2021; Field et al. 2020; Field et al. 2022; Halo et al. 2021; Jagannathan et al. 2021; Player et al. 2021; Schall et al. 2023; Sinding et al. 2021; Wang et al. 2021). These assemblies have high accuracy and continuity, revealing new gene features that were absent from the canFam3.1 assembly. In 2023, the Dog10K consortium released a systematic analysis of canine genome variation based on whole genome sequence data from 1,929 dogs and wolves aligned to the UU_Cfam_GSD_1.0/canFam4 assembly derived from a German Shepherd Dog (Meadows et al. 2023). This included 57 wolves from across Eurasia, 281 village dogs sampled from 26 different countries, 1,579 dogs from 321 breeds, and 12 dogs with a mixed origin or that are not recognized by any international registering body. The Dog10K project represents the largest-

scale genome analysis performed in dogs using a reference other than the Boxer-derived canFam3.1 assembly.

Among the new suite of long-read reference genomes is mCanLor1.2, which is derived from a Greenland wolf (Sinding et al. 2021). This assembly was constructed using Pacific Biosciences HiFi circular consensus reads, a sequencing technology that creates highly-accurate long reads yielding higher quality assemblies compared to previous methods (Wenger et al. 2019). The HiFi read data was further supplemented with Hi-C and 10X Genomics Read Cloud Sequence data to generate a chromosome-scale assembly following methods developed by the Vertebrate Genomes Project (Rhie et al. 2021) and The Darwin Tree of Life Project (Darwin Tree of Life Project 2022). The resulting chromosomes-scale scaffolds were assigned to named chromosomes based on synteny with an existing Labrador retriever genome assembly (GCF_014441545.1). The Greenland wolf belongs to the polar wolf group in North America and can serve as an outgroup to dogs, which have an origin in Eurasia (Sinding et al. 2018).

In this study, we take advantage of recent informatics advances that use the highly parallel processing capabilities of graphics processing units (GPUs) to reanalyze whole genome sequencing data from 1,929 dogs and wolves sequenced by the Dog10K consortium using the mCanLor1.2 Greenland wolf genome as a reference. We first compare the structure of the mCanLor1.2 and UU_Cfam_GSD_1.0 assemblies, then describe patterns of SNP variation found when comparing to each assembly. Finally, we use the wolf-aligned data set to infer the history of population size changes in populations of village dogs and wolves.

Results
Global comparison of mCanLor1.2 and UU_Cfam_GSD_1.0 assemblies
Our previous analysis confirmed the high contiguity and completeness of the mCanLor1.2 assembly relative to other recently published canine assemblies and confirmed that mCanLor1.2 as an outgroup relative to the variation present in assemblies from breed dogs and a Basenji (Nguyen et al. 2024; Schall et al. 2023). Alignment of mCanLor1.2 and UU_Cfam_GSD_1.0 confirmed the large-scale continuity between the assemblies but revealed that the mCanLor1.2 chromosomal orientation is essentially random, with 19/39 analyzed chromosomes represented in the opposite orientation relative to UU_Cfam_GSD_1.0 (Figure 1). Consistent with the inverted orientation of the mCanLor1.2 X chromosome, analysis of read-depth profiles from male and female samples identified the mCanLor1.2 pseudo autosomal region (PAR) as chrX:117984001-124665963 (Figure S1).

Fig. 1 Chromosomal continuity between UU_Cfam_GSD_1.0 and mCanLor1.2
Corresponding segments between the UU_Cfam_GSD_1.0 (x axis) and mCanLor1.2 (y axis)
assemblies were identified using minimap2. The position of aligned segments at least 5,000 bp
in length are plotted. Alignments in the forward orientation are shown in red, alignments in the
reverse orientation are shown in blue.

The mCanLor1.2 assembly was derived from a male sample and contains a partial assembly of
the wolf Y chromosome. The sequence labeled as chrY (HG994382.1) consists of 6.5 Mbp of
ampliconic sequence in a palindromic configuration (Figure S2). Three additional unlocalized
sequences associated with the Y chromosome (CAJNRB020000005.1, CAJNRB020000035.1, and
CAJNRB020000036.1) show homology to KP081776.1, a partial dog Y chromosome assembly (Li
et al. 2013). The unique portions of KP081776.1, which have been previously used as a

reference to study Y chromosome sequence variation (Oetjens et al. 2018; Smeds et al. 2019), largely correspond to portions of CAJNRB020000035.1 and CAJNRB020000036.1. Read-depth analysis of Illumina sequencing reads from the same sample used to generate mCanLor1.2 indicates that CAJNRB020000005.1, CAJNRB020000035.1, and CAJNRB020000036.1 are composed of mixtures of amplified and unique sequence, and that there are additional copies of Y-chromosome palindromes that remain unassembled (Figure S3).

We have previously compared the duplication content of existing canine genome assemblies, finding that the mCanLor1.2 assembly shows the highest concordance between read-depth and assembly self-alignment approaches (Nguyen et al. 2024). To further assess the mCanLor1.2 assembly, we aligned it to UU_Cfam_GSD_1.0 and identified 35,192 segments spanning 18,102,207 bp that are missing in UU_Cfam_GSD_1.0 but present in mCanLor1.2. This includes 9,003 insertions (1,836,394 bp) associated with SINEC sequences and 1,462 (613,486 bp) associated with LINE-1 sequences. Of the 35,192 segments that are missing in UU_Cfam_GSD_1.0, we identified 418 that overlap with an exon annotated in ENSEMBL rapid release mCanLor1.2 annotation (Table S1).

Identification of Canine Sequence Variation Relative to mCanLor1.2
Using the GPU-accelerated pipeline implemented in the NVIDIA Clara Parabricks software (Franke and Crowgey 2020), we aligned reads from 1,929 dogs and wolves to the mCanLor1.2 assembly and identified variants using the same procedures previously used by the Dog10K platform. We identified a total of 36,908,138 SNVs and 13,640,837 indels. This represents an ~6.7% increase over the 34,566,550 SNVs identified based on alignment to UU_Cfam_GSD_1.0. When using the UU_Cfam_GSD_1.0 reference, which was derived from a German Shepherd Dog, the total distance from the UU_Cfam_GSD_1.0 reference varies widely across sample groups (Figure 2). Breed dogs range from 1,470,039 to 2,908,571 differences (median 2,381,059), village dogs range from 1,689,172 to 2,899,436 differences (median 2,535,491) and wolves range from 3,471,282 to 3,582,520.5 differences. The range of divergence among breed and village dogs is drastically reduced when using the mCanLor1.2 Greenland wolf. In this call set, breed dogs range from 3,720,950.5 to 3,803,669 differences (median 3785952.5), village dogs range from 3,720,129 to 3,817,528 (median 3,784,434.5) and wolves range from 3,563,551.0 to 3,810,647.0 (median 3,706,174.5). Although the total number of variants differs greatly based on the references employed, the number of heterozygous SNV genotypes identified in each sample shows a strong correlation between the call sets (Spearmen r = 0.9999, Figure S4). Principal component analysis recapitulates the sample relationships found when using UU_Cfam_GSD_1.0 as the reference. For example, when analyzing a single sample from each of 314 breeds, both SNP sets separate German Shepherds and related breeds from other samples along Principal Component 3 (Figure 3). Thus, the observed sample projections do not reflect a bias induced by alignment to a reference derived from a German Shepherd Dog.

Fig 2. Distance between sequenced samples and the UU_Cfam_GSD1.0 and mCanLor1.2 references
The distance between each sample and the reference genome was calculated as half the number of heterozygous genotypes plus the number of homozygous alternative genotypes. Histograms of the distance between each sample and each reference are shown with breed dogs, village dogs, and wolves plotted separately.

Fig 3. Principal Component Analysis of Breed Dogs is Similar Using Both References
Principal component analysis was performed using one sample from each of 314 breeds based on the variants derived from alignment to the UU_Cfam_GSD_1.0 (left) and mCanLor1.2 (right) assemblies. In both cases, projections of samples onto the 1st vs 2nd and 3rd vs 4th components are shown. Breeds related to German Shepherd Dogs are colored as indicated and separate from the other breeds along principal component 3.

Estimates of Village Dog and Wolf Demographic Histories
Using the alignments to the mCanLor1.2 assembly, we estimated the trajectory of effective population size (N_E) through time for four village dog and four wolf populations (Figure 4). The sampled village dogs and wolf populations do not share a similar population size trajectory until approximately 100,000 years ago, an observation consistent with findings from ancient DNA that indicate that a portion of dog genetic ancestry remained partially differentiated from sampled ancient wolves since before 100 kya (Bergstrom et al. 2022). From 20 – 40 thousand years ago village dog populations experienced a 9-13 fold reduction in effective population size.

Fig 4. Inferred effective population size histories for village dog and wolf populations
The population history of four wolf populations (blue lines: Greece (n=12), Iran (n=6), Sweden (n=6), and Xinjaing, China (n=6)) and four village dog populations (purple lines: Congo (n=16), Iran (n=24), Kenya (n=18), and Liberia (n=17)) were inferred using SMC++. Values were plotted assuming a generation time of 3 years and a mutation rate of $4 * 10^{-9}$ per site per generation.

Discussion

The advent of efficient technologies for genome assembly have enabled graph-based pan genome analyses (Eizenga et al. 2020; Gong et al. 2023). Despite the promise of these approaches, alignment of short-read genome sequence data to a single reference genome remains the dominant paradigm for studies of genetic variation. This may introduce a bias in comparisons among samples since sequencing reads that are more similar to the genome reference sequence are more likely to align. Although careful consideration of alignment tools and parameters can ameliorate this effect (Oliva et al. 2021), the potential effects of the reference on downstream analyses are largely unexplored. Due to the complex history of inbreeding that underlies the formation of modern dog breeds (Parker et al. 2017), the reliance on a single reference genome derived from a particular breed may bias the analysis of genetic differences within and among canines.

This potential bias can be overcome using an appropriate outgroup species as the reference. Recent advances in long-read genome sequencing and assembly approaches have led to the publication of multiple new canine reference genomes. However, reanalyzing existing genome-wide data collections requires an onerous and computationally expensive process of remapping

and variant calling. In this study we combined new, GPU-based implementations of standard alignment and variant identification pipelines to efficiently reanalyze whole genome sequence data from 1,929 dogs and wolves using a high-quality Greenland wolf genome as the reference. As expected, use of the wolf genome normalizes the apparent divergence between dog samples and the reference. General patterns of genome diversity are unchanged when either the Greenland wolf or a German Shepherd Dog reference are employed, including PCA analysis that shows a separation between German Shepherd-related breeds and others. Given the high degree of allele sharing reported among German Shepherd-related breeds and other breed clades (Meadows et al. 2023; Parker et al. 2017), the Greenland wolf-aligned data set may be a useful resource for studies of the origins and evolution of modern dogs.

Methods based on the sequentially Markovian coalescent are powerful approaches for inferring past population sizes based on genome-wide sequencing data (Mather et al. 2020). Taking advantage of alignment to the highly contiguous mCanLor1.2 assembly, we used SMC++ to infer the history of village dog and wolf populations. Unlike other approaches, SMC++ uses multiple individuals and does not require variation data to be phased, resulting in more accurate estimates (Terhorst et al. 2017). Application of SMC++ separately to wolves (6- 12 individuals per population) and village dogs (16-24 individuals per population) indicates that the ancestors of village dogs experienced a 9-13 fold reduction in effective population size 20-40 thousand years ago. This reduction likely reflects domestication from an unsampled wolf population (Bergstrom et al. 2020) and is similar, though slightly smaller, than the ~16-fold reduction in effective population size previously estimated using a Dingo and a Basenji using the pairwise sequential Markovian coalescent (PSMC) inference method (Freedman et al. 2014; Li and Durbin 2011).

Material and Methods
Genome Assembly Comparisons
The mCanLor1.2 (GCA_905319855.2) Greenland wolf assembly (Sinding et al. 2021) was downloaded from NCBI. The UU_Cfam_GSD_1.0 German Shepherd Dog assembly (Wang et al. 2021) ('Mischka', GCA_011100685.1, canFam4) was downloaded from the UCSC Genome Browser (RRID:SCR_005780). To assess continuity, the UU_Cfam_GSD_1.0 genome was aligned to mCanLor1.2 using Minimap2 version 2.26 (Li 2018, 2021) with option -x asm10. The resulting alignments were parsed to retain aligned segments with a query and target sequence length of at least 5,000 bp.

The approximate location of the Pseudoautosomal Region (PAR) on the X chromosome was determined by examination of read depth profiles from 3 male (VILLCR000002, CLUPGR000011, and BOXR000005) and 1 female (CLUPSE000004) samples using mosdepth version 0.3.2 (RRID:SCR_018929) (Pedersen and Quinlan 2018) with a window size of 1 kbp.

To assess representation of the Y chromosome, four mCanLor1.2 segments were compared: chrY (HG994382.1) as well as three unlocalized sequences associated with the Y chromosome: CAJNRB020000005.1, CAJNRB020000035.1, and CAJNRB020000036.1. Self-similarity dotplots were generated used Gepard version 2.1 (Krumsiek et al. 2007) with a word size of 100 bp and

ampliconic arms defined using iterative alignment with Minimap2 version 2.26. The mCanLor1.2 Y-chromosome sequences were compared with KP081776.1, an assembly of a portion of the dog Y chromosome generated by pooled sequencing of BAC clones from a Doberman (Li et al. 2013) that has been previously used as a reference to study Y chromosome sequence variation among canids using short-read sequencing (Oetjens et al. 2018; Smeds et al. 2019). The positions of the Y-chromosome SNVs used in (Oetjens et al. 2018) were converted to mCanLor1.2 coordinates by aligning 101 bp long fragments centered on each variant using Minimap2 version 2.26. Read-depth profiles based on the Illumina data generated as part of the mCanLor1.2 assembly were calculated using fastCN as previously described (Nguyen et al. 2024; Pendleton et al. 2018).

Insertions in mCanLor1.2 relative to UU_Cfam_GSD_1.0 were defined using SVIM-asm version 1.0.3 (Heller and Vingron 2021) based on Minimap2 alignments. Analysis was limited to the 38 assembled autosomes and the X chromosome. Putative SINE insertions were identified as insertions 150-250 bp in size that contained at least 90% SINE sequence. Potential LINE-1 insertions were not filtered by length but were required to have at least 90% of the insertion sequence annotated as LINE. Intersecting exons were identified using the ENSEMBL rapid release of the mCanLor1.2 GTF file ([https://ftp.ensembl.org/pub/rapid-release/species/Canis_lupus/GCA_905319855.2/ensembl/geneset/2022_01/Canis_lupus-GCA_905319855.2-2022_01-genes.gtf.gz)](https://ftp.ensembl.org/pub/rapid-release/species/Canis_lupus/GCA_905319855.2/ensembl/geneset/2022_01/Canis_lupus-GCA_905319855.2-2022_01-genes.gtf.gz)

Sequence Alignment and Variant Identification Relative to mCanLor1.2
We identified single nucleotide (SNV) and short insertion-deletion (indel) variants based on alignment of Illumina data to the mCanLor1.2 assembly. Our procedure began with the aligned CRAM files from 1,929 samples that passed quality control filters described by the Dog10K consortium (Meadows et al. 2023). We note that the base qualities in the CRAM files were already recalibrated based on alignment to the UU_Cfam_GSD_1.0 assembly. First, reads from the CRAM files were converted to fastq format using the collate and bam2fq commands from SAMtools version 1.13 (RRID:SCR_002105). The resulting reads were then processed using the fq2bam and haplotypecaller commands from the GPU-accelerated NVIDIA Clara Parabricks tool kit version 4.0.1-1 (Franke and Crowgey 2020). This resulted in a genomic variant call format (GVCF) file for each sample.

Sequence variants were then identified from the 1,929 GVCF files following the same procedures previously used by the Dog10K Consortium (Meadows et al. 2023). First, candidate variants were identified using the *GenomicsDBImport* and *GenotypeGVCFs* commands in GATK version 4.2.0.0 (RRID:SCR_001876). SNV loci were separated from loci containing indels using the GATK *SelectVariants* command. High-quality SNVs were identified using the variant quality score recalibration (VQSR) procedure using 680,458 SNPs from the Illumina Canine HD and Axiom Canine HD SNP arrays as a truth set. Positions were converted from UU_Cfam_GSD_1.0 to mCanLor1.2 coordinates using the GATK *LiftoverVcf* utility with a genome chain file generated constructed from BLAT alignments (Hinrichs et al. 2006; Kent 2002). Due to differences in read depth profiles, variants on the autosomes and X-par region were analyzed separately from variants on the non-PAR portion of the X chromosome. In each case, cut offs

were used such that 99.0% of variants in the truth set were retained, resulting in a total of 36,908,138 PASS SNVs for downstream analysis. Due to the absence of suitable training data, candidate indels were identified using the GATK *VariantFiltration* utility with the following criteria: QD < 2.0, FS > 200.0, *ReadPosRankSum* < -2.0, SOR > 10.0.

SNV analysis

The distance between each sample and the reference genome was calculated as the number of homozygous alternative SNV genotypes plus half the number of heterozygous genotypes. Principal Component Analysis (PCA) was performed using *smartpca* from the EIGENSOFT package (version 8.0.0 RRID:SCR_004965) (Patterson et al. 2006). First, samples were filtered to retain only one sample from each of 314 breeds, keeping the sample with the highest coverage in the original Dog10K data set. Breeds with large amounts of recent wolf ancestry (Czechoslovakian Wolfdog, Saarloos Wolfdog, and Tamaskan) were omitted. Next, biallelic autosomal SNVs with a minor allele frequency of at least 0.05 and a genotyping missing rate less than 0.10 were selected using PLINK version 1.9 (RRID:SCR_000084) (Chang et al. 2015). The resulting variants were pruned for linkage disequilibrium using PLINK with the command --indep-pairwise 50 10 0.1.

Trajectories of effective population size (N_E) through time were constructed using SMC++ (Terhorst et al. 2017) version v1.15.4.dev18+gca077da.d20210316. Analysis was performed on eight groups including 12 wolves from Greece, 6 wolves from Iran, 6 wolves from Sweden, 6 wolves form Xinjaing, China, 16 village dogs from Congo, 24 village dogs from Iran, 18 village dogs from Kenya, and 17 village dogs from Liberia. Variation within regions of the mCanLor12 assembly identified as segmental duplications was removed using the *–mask* option in SMC++ (Nguyen et al. 2024). In each case, analysis was performed for each autosome and was repeated using five different distinguished individuals, based on the lexigraphical order of sample identifiers. Results were converted from coalescent units using a generation time of 3 years and a mutation rate of $4 * 10^{-9}$ per site per generation, which is compatible with estimates from ancient DNA (Skoglund et al. 2015) and wolf pedigrees (Koch et al. 2019).

Supplementary Information

The online version contains supplementary material including Supplementary Table 1 and Supplementary Figures 1- 4.

Acknowledgements

We thank all members of the Dog10K consortium for contributing their samples and data to the project, the University of Michigan Advanced Research Computing Group for assistance with efficient GPU utilization.

Data Availability

SNV and indel variant data is available from the Zenodo archive under DOI 10.5281/zenodo.11248171.

Competing Interests

The authors declare that no competing interests exist.

Supplementary Figures

Fig. S1. Identification of the mCanLor1.2 pseudo autosomal region
Read depth profiles are shown for 1 female (CLUPSE000004) and 3 male (VILLCR000002, CLUPGR000011, and BOXR000005) samples along the end of the X chromosome. The dashed red line represents the inferred PAR boundary, which extends from chrX:117984001-124665963.

Fig S2. Ampliconic structure of chrY (HG994382.1)
A self-similarity dotplot of the mCanLor1.2 Y chromosome sequence drawn with a window size of 100 bp is shown. The colored arrows correspond to ampliconic sequences identified in the assembly. The entire sequence represents a complex palindromic structure.

Fig S3. Copy-number profile across Y-chromosome sequences
Copy-number was inferred using fastCN applied to Illumina genome sequence data generated from the sample used to create the mCanLor1.2 assembly. Results are shown for the mCanLor1.2 Y-chromosome sequence (top) as well as three additional unplaced assembled contigs that are associated with the Y chromosome.

Fig S4. Number of heterozygous sites identified based on alignment to UU_Cfam_GSD_1.0 and mCanLor1.2
A scatter plot of the number of heterozygous sites identified from alignment to each reference genome is shown for breed dogs, village dogs, and wolves.

Supplementary Table 1 Sequence deleted in UU_Cfam_GSD_1.0 that overlaps with annotated exons (see excel file)

References

Bergstrom A, Frantz L, Schmidt R, Ersmark E, Lebrasseur O, Girdland-Flink L, Lin AT, Stora J, Sjogren KG, Anthony D, Antipina E, Amiri S, Bar-Oz G, Bazaliiskii VI, Bulatovic J, Brown D, Carmagnini A, Davy T, Fedorov S, Fiore I, Fulton D, Germonpre M, Haile J, Irving-Pease EK, Jamieson A, Janssens L, Kirillova I, Horwitz LK, Kuzmanovic-Cvetkovic J, Kuzmin Y, Losey RJ, Dizdar DL, Mashkour M, Novak M, Onar V, Orton D, Pasaric M, Radivojevic M, Rajkovic D, Roberts B, Ryan H, Sablin M, Shidlovskiy F, Stojanovic I, Tagliacozzo A, Trantalidou K, Ullen I, Villaluenga A, Wapnish P, Dobney K, Gotherstrom A, Linderholm A, Dalen L, Pinhasi R, Larson G, Skoglund P (2020) Origins and genetic legacy of prehistoric dogs. Science 370, 557-564

Bergstrom A, Stanton DWG, Taron UH, Frantz L, Sinding MS, Ersmark E, Pfrengle S, Cassatt-Johnstone M, Lebrasseur O, Girdland-Flink L, Fernandes DM, Ollivier M, Speidel L, Gopalakrishnan S, Westbury MV, Ramos-Madrigal J, Feuerborn TR, Reiter E, Gretzinger J, Munzel SC, Swali P, Conard NJ, Caroe C, Haile J, Linderholm A, Androsov S, Barnes I, Baumann C, Benecke N, Bocherens H, Brace S, Carden RF, Drucker DG, Fedorov S, Gasparik M, Germonpre M, Grigoriev S, Groves P, Hertwig ST, Ivanova VV, Janssens L, Jennings RP, Kasparov AK, Kirillova IV, Kurmaniyazov I, Kuzmin YV, Kosintsev PA, Laznickova-Galetova M, Leduc C, Nikolskiy P, Nussbaumer M, O'Drisceoil C, Orlando L, Outram A, Pavlova EY, Perri AR, Pilot M, Pitulko VV, Plotnikov VV, Protopopov AV, Rehazek A, Sablin M, Seguin-Orlando A, Stora J, Verjux C, Zaibert VF, Zazula G, Crombe P, Hansen AJ, Willerslev E, Leonard JA, Gotherstrom A, Pinhasi R, Schuenemann VJ, Hofreiter M, Gilbert MTP, Shapiro B, Larson G, Krause J, Dalen L, Skoglund P (2022) Grey wolf genomic history reveals a dual ancestry of dogs. Nature 607, 313-320

Botigue LR, Song S, Scheu A, Gopalan S, Pendleton AL, Oetjens M, Taravella AM, Seregely T, Zeeb-Lanz A, Arbogast RM, Bobo D, Daly K, Unterlander M, Burger J, Kidd JM, Veeramah KR (2017) Ancient European dog genomes reveal continuity since the Early Neolithic. Nat Commun 8, 16082

Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ (2015) Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience 4, 7

Clark AG, Hubisz MJ, Bustamante CD, Williamson SH, Nielsen R (2005) Ascertainment bias in studies of human genome-wide polymorphism. Genome Res 15, 1496-1502

Darwin Tree of Life Project C (2022) Sequence locally, think globally: The Darwin Tree of Life Project. Proc Natl Acad Sci U S A 119

Edwards RJ, Field MA, Ferguson JM, Dudchenko O, Keilwagen J, Rosen BD, Johnson GS, Rice ES, Hillier D, Hammond JM, Towarnicki SG, Omer A, Khan R, Skvortsova K, Bogdanovic O, Zammit RA, Aiden EL, Warren WC, Ballard JWO (2021) Chromosome-length genome assembly and structural variations of the primal Basenji dog (Canis lupus familiaris) genome. BMC Genomics 22, 188

Eizenga JM, Novak AM, Sibbesen JA, Heumos S, Ghaffaari A, Hickey G, Chang X, Seaman JD, Rounthwaite R, Ebler J, Rautiainen M, Garg S, Paten B, Marschall T, Siren J, Garrison E (2020) Pangenome Graphs. Annu Rev Genomics Hum Genet 21, 139-162

Field MA, Rosen BD, Dudchenko O, Chan EKF, Minoche AE, Edwards RJ, Barton K, Lyons RJ, Tuipulotu DE, Hayes VM, A DO, Colaric Z, Keilwagen J, Skvortsova K, Bogdanovic O, Smith MA, Aiden EL, Smith TPL, Zammit RA, Ballard JWO (2020) Canfam_GSD: De novo chromosome-length genome assembly of the German Shepherd Dog (Canis lupus familiaris) using a combination of long reads, optical mapping, and Hi-C. Gigascience 9

Field MA, Yadav S, Dudchenko O, Esvaran M, Rosen BD, Skvortsova K, Edwards RJ, Keilwagen J, Cochran BJ, Manandhar B, Bustamante S, Rasmussen JA, Melvin RG, Chernoff B, Omer A, Colaric Z, Chan EKF, Minoche AE, Smith TPL, Gilbert MTP, Bogdanovic O, Zammit RA, Thomas T, Aiden EL, Ballard JWO (2022) The Australian dingo is an early offshoot of modern breed dogs. Sci Adv 8, eabm5944

Franke KR, Crowgey EL (2020) Accelerating next generation sequencing data analysis: an evaluation of optimized best practices for Genome Analysis Toolkit algorithms. Genomics Inform 18, e10

Frantz LA, Mullin VE, Pionnier-Capitan M, Lebrasseur O, Ollivier M, Perri A, Linderholm A, Mattiangeli V, Teasdale MD, Dimopoulos EA, Tresset A, Duffraisse M, McCormick F, Bartosiewicz L, Gal E, Nyerges EA, Sablin MV, Brehard S, Mashkour M, Balasescu A, Gillet B, Hughes S, Chassaing O, Hitte C, Vigne JD, Dobney K, Hanni C, Bradley DG, Larson G (2016) Genomic and archaeological evidence suggest a dual origin of domestic dogs. Science 352, 1228-1231

Freedman AH, Gronau I, Schweizer RM, Ortega-Del Vecchyo D, Han E, Silva PM, Galaverni M, Fan Z, Marx P, Lorente-Galdos B, Beale H, Ramirez O, Hormozdiari F, Alkan C, Vila C, Squire K, Geffen E, Kusak J, Boyko AR, Parker HG, Lee C, Tadigotla V, Wilton A, Siepel A, Bustamante CD, Harkins TT, Nelson SF, Ostrander EA, Marques-Bonet T, Wayne RK, Novembre J (2014) Genome sequencing highlights the dynamic early history of dogs. PLoS Genet 10, e1004016

Gong Y, Li Y, Liu X, Ma Y, Jiang L (2023) A review of the pangenome: how it affects our understanding of genomic variation, selection and breeding in domestic animals? J Anim Sci Biotechnol 14, 73

Gopalakrishnan S, Samaniego Castruita JA, Sinding MS, Kuderna LFK, Raikkonen J, Petersen B, Sicheritz-Ponten T, Larson G, Orlando L, Marques-Bonet T, Hansen AJ, Dalen L, Gilbert MTP (2017) The wolf reference genome sequence (Canis lupus lupus) and its implications for Canis spp. population genomics. BMC Genomics 18, 495

Gunther T, Nettelblad C (2019) The presence and impact of reference bias on population genomic studies of prehistoric human populations. PLoS Genet 15, e1008302

Halo JV, Pendleton AL, Shen F, Doucet AJ, Derrien T, Hitte C, Kirby LE, Myers B, Sliwerska E, Emery S, Moran JV, Boyko AR, Kidd JM (2021) Long-read assembly of a Great Dane genome highlights the contribution of GC-rich sequence and mobile elements to canine genomes. Proc Natl Acad Sci U S A 118

Heller D, Vingron M (2021) SVIM-asm: structural variant detection from haploid and diploid genome assemblies. Bioinformatics 36, 5519-5521

Hinrichs AS, Karolchik D, Baertsch R, Barber GP, Bejerano G, Clawson H, Diekhans M, Furey TS, Harte RA, Hsu F, Hillman-Jackson J, Kuhn RM, Pedersen JS, Pohl A, Raney BJ, Rosenbloom KR, Siepel A, Smith KE, Sugnet CW, Sultan-Qurraie A, Thomas DJ, Trumbower H, Weber RJ, Weirauch M, Zweig AS, Haussler D, Kent WJ (2006) The UCSC Genome Browser Database: update 2006. Nucleic Acids Res 34, D590-598

Hoeppner MP, Lundquist A, Pirun M, Meadows JR, Zamani N, Johnson J, Sundstrom G, Cook A, FitzGerald MG, Swofford R, Mauceli E, Moghadam BT, Greka A, Alfoldi J, Abouelleil A, Aftuck L, Bessette D, Berlin A, Brown A, Gearin G, Lui A, Macdonald JP, Priest M, Shea T, Turner-Maier J, Zimmer A, Lander ES, di Palma F, Lindblad-Toh K, Grabherr MG (2014) An improved canine genome and a comprehensive catalogue of coding genes and non-coding transcripts. PLoS One 9, e91172

Jagannathan V, Drogemuller C, Leeb T, Dog Biomedical Variant Database C (2019) A comprehensive biomedical variant catalogue based on whole genome sequences of 582 dogs and eight wolves. Anim Genet 50, 695-704

Jagannathan V, Hitte C, Kidd JM, Masterson P, Murphy TD, Emery S, Davis B, Buckley RM, Liu YH, Zhang XQ, Leeb T, Zhang YP, Ostrander EA, Wang GD (2021) Dog10K_Boxer_Tasha_1.0: A Long-Read Assembly of the Dog Reference Genome. Genes (Basel) 12

Kent WJ (2002) BLAT--the BLAST-like alignment tool. Genome Res 12, 656-664

Koch EM, Schweizer RM, Schweizer TM, Stahler DR, Smith DW, Wayne RK, Novembre J (2019) De Novo Mutation Rate Estimation in Wolves of Known Pedigree. Mol Biol Evol 36, 2536-2547

Krumsiek J, Arnold R, Rattei T (2007) Gepard: a rapid and sensitive tool for creating dotplots on genome scale. Bioinformatics 23, 1026-1028

Lachance J, Tishkoff SA (2013) SNP ascertainment bias in population genetic analyses: why it is important, and how to correct it. Bioessays 35, 780-786

Li G, Davis BW, Raudsepp T, Pearks Wilkerson AJ, Mason VC, Ferguson-Smith M, O'Brien PC, Waters PD, Murphy WJ (2013) Comparative analysis of mammalian Y chromosomes illuminates ancestral structure and lineage-specific evolution. Genome Res 23, 1486-1495

Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34, 3094-3100

Li H (2021) New strategies to improve minimap2 alignment accuracy. Bioinformatics 37, 4572-4574

Li H, Durbin R (2011) Inference of human population history from individual whole-genome sequences. Nature 475, 493-496

Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, Clamp M, Chang JL, Kulbokas EJ, 3rd, Zody MC, Mauceli E, Xie X, Breen M, Wayne RK, Ostrander EA, Ponting CP, Galibert F, Smith DR, DeJong PJ, Kirkness E, Alvarez P, Biagi T, Brockman W, Butler J, Chin CW, Cook A, Cuff J, Daly MJ, DeCaprio D, Gnerre S, Grabherr M, Kellis M, Kleber M, Bardeleben C, Goodstadt L, Heger A, Hitte C, Kim L, Koepfli KP, Parker HG, Pollinger JP, Searle SM, Sutter NB, Thomas R, Webber C, Baldwin J, Abebe A, Abouelleil A, Aftuck L, Ait-Zahra M, Aldredge T, Allen

N, An P, Anderson S, Antoine C, Arachchi H, Aslam A, Ayotte L, Bachantsang P, Barry A, Bayul T, Benamara M, Berlin A, Bessette D, Blitshteyn B, Bloom T, Blye J, Boguslavskiy L, Bonnet C, Boukhgalter B, Brown A, Cahill P, Calixte N, Camarata J, Cheshatsang Y, Chu J, Citroen M, Collymore A, Cooke P, Dawoe T, Daza R, Decktor K, DeGray S, Dhargay N, Dooley K, Dooley K, Dorje P, Dorjee K, Dorris L, Duffey N, Dupes A, Egbiremolen O, Elong R, Falk J, Farina A, Faro S, Ferguson D, Ferreira P, Fisher S, FitzGerald M, Foley K, Foley C, Franke A, Friedrich D, Gage D, Garber M, Gearin G, Giannoukos G, Goode T, Goyette A, Graham J, Grandbois E, Gyaltsen K, Hafez N, Hagopian D, Hagos B, Hall J, Healy C, Hegarty R, Honan T, Horn A, Houde N, Hughes L, Hunnicutt L, Husby M, Jester B, Jones C, Kamat A, Kanga B, Kells C, Khazanovich D, Kieu AC, Kisner P, Kumar M, Lance K, Landers T, Lara M, Lee W, Leger JP, Lennon N, Leuper L, LeVine S, Liu J, Liu X, Lokyitsang Y, Lokyitsang T, Lui A, Macdonald J, Major J, Marabella R, Maru K, Matthews C, McDonough S, Mehta T, Meldrim J, Melnikov A, Meneus L, Mihalev A, Mihova T, Miller K, Mittelman R, Mlenga V, Mulrain L, Munson G, Navidi A, Naylor J, Nguyen T, Nguyen N, Nguyen C, Nguyen T, Nicol R, Norbu N, Norbu C, Novod N, Nyima T, Olandt P, O'Neill B, O'Neill K, Osman S, Oyono L, Patti C, Perrin D, Phunkhang P, Pierre F, Priest M, Rachupka A, Raghuraman S, Rameau R, Ray V, Raymond C, Rege F, Rise C, Rogers J, Rogov P, Sahalie J, Settipalli S, Sharpe T, Shea T, Sheehan M, Sherpa N, Shi J, Shih D, Sloan J, Smith C, Sparrow T, Stalker J, Stange-Thomann N, Stavropoulos S, Stone C, Stone S, Sykes S, Tchuinga P, Tenzing P, Tesfaye S, Thoulutsang D, Thoulutsang Y, Topham K, Topping I, Tsamla T, Vassiliev H, Venkataraman V, Vo A, Wangchuk T, Wangdi T, Weiand M, Wilkinson J, Wilson A, Yadav S, Yang S, Yang X, Young G, Yu Q, Zainoun J, Zembek L, Zimmer A, Lander ES (2005) Genome sequence, comparative analysis and haplotype structure of the domestic dog. Nature 438, 803-819

Martiniano R, Garrison E, Jones ER, Manica A, Durbin R (2020) Removing reference bias and improving indel calling in ancient DNA data analysis by mapping to a sequence variation graph. Genome Biol 21, 250

Mather N, Traves SM, Ho SYW (2020) A practical introduction to sequentially Markovian coalescent methods for estimating demographic history from genomic data. Ecol Evol 10, 579-589

Meadows JRS, Kidd JM, Wang GD, Parker HG, Schall PZ, Bianchi M, Christmas MJ, Bougiouri K, Buckley RM, Hitte C, Nguyen AK, Wang C, Jagannathan V, Niskanen JE, Frantz LAF, Arumilli M, Hundi S, Lindblad-Toh K, Ginja C, Agustina KK, Andre C, Boyko AR, Davis BW, Drogemuller M, Feng XY, Gkagkavouzis K, Iliopoulos G, Harris AC, Hytonen MK, Kalthoff DC, Liu YH, Lymberakis P, Poulakakis N, Pires AE, Racimo F, Ramos-Almodovar F, Savolainen P, Venetsani S, Tammen I, Triantafyllidis A, vonHoldt B, Wayne RK, Larson G, Nicholas FW, Lohi H, Leeb T, Zhang YP, Ostrander EA (2023) Genome sequencing of 2000 canids by the Dog10K consortium advances the understanding of demography, genome function and architecture. Genome Biol 24, 187

Nassar LR, Barber GP, Benet-Pages A, Casper J, Clawson H, Diekhans M, Fischer C, Gonzalez JN, Hinrichs AS, Lee BT, Lee CM, Muthuraman P, Nguy B, Pereira T, Nejad P, Perez G, Raney BJ, Schmelter D, Speir ML, Wick BD, Zweig AS, Haussler D, Kuhn RM, Haeussler M, Kent WJ (2023) The UCSC Genome Browser database: 2023 update. Nucleic Acids Res 51, D1188-D1195

Nguyen AK, Blacksmith MS, Kidd JM (2024) Duplications and retrogenes are numerous and widespread in modern canine genomic assemblies. bioRxiv 2023.10.31.564742;

Ni Leathlobhair M, Perri AR, Irving-Pease EK, Witt KE, Linderholm A, Haile J, Lebrasseur O, Ameen C, Blick J, Boyko AR, Brace S, Cortes YN, Crockford SJ, Devault A, Dimopoulos EA, Eldridge M, Enk J, Gopalakrishnan S, Gori K, Grimes V, Guiry E, Hansen AJ, Hulme-Beaman A, Johnson J, Kitchen A, Kasparov AK, Kwon YM, Nikolskiy PA, Lope CP, Manin A, Martin T, Meyer M, Myers KN, Omura M, Rouillard JM, Pavlova EY, Sciulli P, Sinding MS, Strakova A, Ivanova VV, Widga C, Willerslev E, Pitulko VV, Barnes I, Gilbert MTP, Dobney KM, Malhi RS, Murchison EP, Larson G, Frantz LAF (2018) The evolutionary history of dogs in the Americas. Science 361, 81-85

Oetjens MT, Martin A, Veeramah KR, Kidd JM (2018) Analysis of the canid Y-chromosome phylogeny using short-read sequencing data reveals the presence of distinct haplogroups among Neolithic European dogs. BMC Genomics 19, 350

Oliva A, Tobler R, Llamas B, Souilmi Y (2021) Additional evaluations show that specific BWA-aln settings still outperform BWA-mem for ancient DNA data alignment. Ecol Evol 11, 18743-18748

Orlando L, Ginolhac A, Zhang G, Froese D, Albrechtsen A, Stiller M, Schubert M, Cappellini E, Petersen B, Moltke I, Johnson PL, Fumagalli M, Vilstrup JT, Raghavan M, Korneliussen T, Malaspinas AS, Vogt J, Szklarczyk D, Kelstrup CD, Vinther J, Dolocan A, Stenderup J, Velazquez AM, Cahill J, Rasmussen M, Wang X, Min J, Zazula GD, Seguin-Orlando A, Mortensen C, Magnussen K, Thompson JF, Weinstock J, Gregersen K, Roed KH, Eisenmann V, Rubin CJ, Miller DC, Antczak DF, Bertelsen MF, Brunak S, Al-Rasheid KA, Ryder O, Andersson L, Mundy J, Krogh A, Gilbert MT, Kjaer K, Sicheritz-Ponten T, Jensen LJ, Olsen JV, Hofreiter M, Nielsen R, Shapiro B, Wang J, Willerslev E (2013) Recalibrating Equus evolution using the genome sequence of an early Middle Pleistocene horse. Nature 499, 74-78

Parker HG, Dreger DL, Rimbault M, Davis BW, Mullen AB, Carpintero-Ramirez G, Ostrander EA (2017) Genomic Analyses Reveal the Influence of Geographic Origin, Migration, and Hybridization on Modern Dog Breed Development. Cell Rep 19, 697-708

Patterson N, Price AL, Reich D (2006) Population structure and eigenanalysis. PLoS Genet 2, e190

Pedersen BS, Quinlan AR (2018) Mosdepth: quick coverage calculation for genomes and exomes. Bioinformatics 34, 867-868

Pendleton AL, Shen F, Taravella AM, Emery S, Veeramah KR, Boyko AR, Kidd JM (2018) Comparison of village dog and wolf genomes highlights the role of the neural crest in dog domestication. BMC Biol 16, 64

Plassais J, Kim J, Davis BW, Karyadi DM, Hogan AN, Harris AC, Decker B, Parker HG, Ostrander EA (2019) Whole genome sequencing of canids reveals genomic regions under selection and variants influencing morphology. Nat Commun 10, 1489

Player RA, Forsyth ER, Verratti KJ, Mohr DW, Scott AF, Bradburne CE (2021) A novel canis lupus familiaris reference genome improves variant resolution for use in breed-specific GWAS. Life Sci Alliance 4

Ramos-Madrigal J, Sinding MS, Caroe C, Mak SST, Niemann J, Samaniego Castruita JA, Fedorov S, Kandyba A, Germonpre M, Bocherens H, Feuerborn TR, Pitulko VV, Pavlova EY, Nikolskiy PA,

Kasparov AK, Ivanova VV, Larson G, Frantz LAF, Willerslev E, Meldgaard M, Petersen B, Sicheritz-Ponten T, Bachmann L, Wiig O, Hansen AJ, Gilbert MTP, Gopalakrishnan S (2021) Genomes of Pleistocene Siberian Wolves Uncover Multiple Extinct Wolf Lineages. Curr Biol 31, 198-206 e198

Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, Uliano-Silva M, Chow W, Fungtammasan A, Kim J, Lee C, Ko BJ, Chaisson M, Gedman GL, Cantin LJ, Thibaud-Nissen F, Haggerty L, Bista I, Smith M, Haase B, Mountcastle J, Winkler S, Paez S, Howard J, Vernes SC, Lama TM, Grutzner F, Warren WC, Balakrishnan CN, Burt D, George JM, Biegler MT, Iorns D, Digby A, Eason D, Robertson B, Edwards T, Wilkinson M, Turner G, Meyer A, Kautt AF, Franchini P, Detrich HW, 3rd, Svardal H, Wagner M, Naylor GJP, Pippel M, Malinsky M, Mooney M, Simbirsky M, Hannigan BT, Pesout T, Houck M, Misuraca A, Kingan SB, Hall R, Kronenberg Z, Sovic I, Dunn C, Ning Z, Hastie A, Lee J, Selvaraj S, Green RE, Putnam NH, Gut I, Ghurye J, Garrison E, Sims Y, Collins J, Pelan S, Torrance J, Tracey A, Wood J, Dagnew RE, Guan D, London SE, Clayton DF, Mello CV, Friedrich SR, Lovell PV, Osipova E, Al-Ajli FO, Secomandi S, Kim H, Theofanopoulou C, Hiller M, Zhou Y, Harris RS, Makova KD, Medvedev P, Hoffman J, Masterson P, Clark K, Martin F, Howe K, Flicek P, Walenz BP, Kwak W, Clawson H, Diekhans M, Nassar L, Paten B, Kraus RHS, Crawford AJ, Gilbert MTP, Zhang G, Venkatesh B, Murphy RW, Koepfli KP, Shapiro B, Johnson WE, Di Palma F, Marques-Bonet T, Teeling EC, Warnow T, Graves JM, Ryder OA, Haussler D, O'Brien SJ, Korlach J, Lewin HA, Howe K, Myers EW, Durbin R, Phillippy AM, Jarvis ED (2021) Towards complete and error-free genome assemblies of all vertebrate species. Nature 592, 737-746

Schall PZ, Winkler PA, Petersen-Jones SM, Yuzbasiyan-Gurkan V, Kidd JM (2023) Genome-wide methylation patterns from canine nanopore assemblies. G3 (Bethesda) 13

Sinding MS, Gopalakrishan S, Vieira FG, Samaniego Castruita JA, Raundrup K, Heide Jorgensen MP, Meldgaard M, Petersen B, Sicheritz-Ponten T, Mikkelsen JB, Marquard-Petersen U, Dietz R, Sonne C, Dalen L, Bachmann L, Wiig O, Hansen AJ, Gilbert MTP (2018) Population genomics of grey wolves and wolf-like canids in North America. PLoS Genet 14, e1007745

Sinding MS, Gopalakrishnan S, Raundrup K, Dalen L, Threlfall J, Darwin Tree of Life Barcoding c, Wellcome Sanger Institute Tree of Life p, Wellcome Sanger Institute Scientific Operations DNAPc, Tree of Life Core Informatics c, Darwin Tree of Life C, Gilbert T (2021) The genome sequence of the grey wolf, Canis lupus Linnaeus 1758. Wellcome Open Res 6, 310

Skoglund P, Ersmark E, Palkopoulou E, Dalen L (2015) Ancient wolf genome reveals an early divergence of domestic dog ancestors and admixture into high-latitude breeds. Curr Biol 25, 1515-1519

Smeds L, Kojola I, Ellegren H (2019) The evolutionary history of grey wolf Y chromosomes. Mol Ecol 28, 2173-2191

Stevenson KR, Coolon JD, Wittkopp PJ (2013) Sources of bias in measures of allele-specific expression derived from RNA-sequence data aligned to a single reference genome. BMC Genomics 14, 536

Terhorst J, Kamm JA, Song YS (2017) Robust and scalable inference of population history from hundreds of unphased whole genomes. Nat Genet 49, 303-309

Vaysse A, Ratnakumar A, Derrien T, Axelsson E, Rosengren Pielberg G, Sigurdsson S, Fall T, Seppala EH, Hansen MS, Lawley CT, Karlsson EK, Consortium L, Bannasch D, Vila C, Lohi H, Galibert F, Fredholm M, Haggstrom J, Hedhammar A, Andre C, Lindblad-Toh K, Hitte C, Webster MT (2011) Identification of genomic regions associated with phenotypic variation between dog breeds using selection mapping. PLoS Genet 7, e1002316

Wang C, Wallerman O, Arendt ML, Sundstrom E, Karlsson A, Nordin J, Makelainen S, Pielberg GR, Hanson J, Ohlsson A, Saellstrom S, Ronnberg H, Ljungvall I, Haggstrom J, Bergstrom TF, Hedhammar A, Meadows JRS, Lindblad-Toh K (2021) A novel canine reference genome resolves genomic architecture and uncovers transcript complexity. Commun Biol 4, 185

Wenger AM, Peluso P, Rowell WJ, Chang PC, Hall RJ, Concepcion GT, Ebler J, Fungtammasan A, Kolesnikov A, Olson ND, Topfer A, Alonge M, Mahmoud M, Qian Y, Chin CS, Phillippy AM, Schatz MC, Myers G, DePristo MA, Ruan J, Marschall T, Sedlazeck FJ, Zook JM, Li H, Koren S, Carroll A, Rank DR, Hunkapiller MW (2019) Accurate circular consensus long-read sequencing improves variant detection and assembly of a human genome. Nat Biotechnol 37, 1155-1162