Group identification drives brain integration for collective performance

Enhui Xie[1], Shuyi Zha[1], Yiyang Xu[1], Xianchun Li[#,1,2,3]

[1] Shanghai Key Laboratory of Mental Health and Psychological Crisis Intervention, Affiliated

Mental Health Center (ECNU), School of Psychology and Cognitive Science, East China

Normal University, Shanghai, 200062, China

[2] Shanghai Changning Mental Health Center, Shanghai, China, 200335

[3] Institute of Wisdom in China, East China Normal University, Shanghai, China, 200062

[#] Correspondence to:

Xianchun Li (xcli@psy.ecnu.edu.cn)

School of Psychology and Cognitive Science

East China Normal University

Shanghai, China

Abstract

Group identification may influence collective behaviors and result in variations in collective performance. However, the evidence for this hypothesis and the neural mechanisms involved remain elusive. To this end, we conducted a study using both single-brain activation and multi-brain synchronization analyses to investigate how group identification influences collective problem-solving in a murder mystery case. Our results showed that groups with high levels of identification performed better individually compared to those with low identification, as supported by single-brain activation in the dorsolateral prefrontal cortex (DLPFC). Furthermore, high-identification groups also showed enhanced collective performance, supported by within-group neural synchronization (GNS) in the orbitofrontal cortex (OFC). The DLPFC-OFC connectivity played a crucial role in linking individual and collective performance. Overall, our study provides a two-in-one neural model to explain how group identification affects collective decision-making processes, offering valuable insights into the dynamics of group interactions.

Keywords: group identification, collective performance, single-brain activation, within-group neural synchronization (GNS)

Introduction

While group decision-making is commonly utilized and can yield positive outcomes, not all groups function effectively, and their performance can vary (De Wilde et al., 2017; Kerr and Tindale, 2003; Stasser and Abele, 2020; Xie et al., 2023). Even when two teams are equally skilled and exchange information, one team often outperforms the other. Interestingly, the winning team typically shows stronger cohesion, while the losing team appears to be more disorganized. Recent studies have highlighted the importance of group identification in determining how verbal information exchange affects collective performance (Reinero et al., 2021; Van Bavel and Cunningham, 2012; Xie et al., 2023). It appears that differences in group identification may play a role in the varying performance of different groups. Here we ask: How does group identification lead to differences in collective performance? What is the underlying mechanism behind this effect?

Social identity theory suggests that group identification is crucial in shaping collective behavior (Bicchieri, 2002; Gundlach et al., 2006; Tajfel and Turner, 1979). This theory elucidates that group identification refers to a person's attitudes toward a specific group and the emotional meaning associated with belonging to it. Research has consistently shown that stronger group identification is associated with better collective behavior, such as improved decision-making accuracy (Pärnamets et al., 2020), effectiveness in team-based interventions (Reinero et al., 2021), and overall group cooperation (Van Bavel et al., 2012). This is likely due to the positive impact of increased communication, cooperation, and reduced conflict among group members. Additionally, individuals with a strong sense of group identification are more likely to align their goals with those of the group, leading to increased effort towards achieving objectives (Dikker et al., 2022; Prochazkova et al., 2022). Based on this, we hypothesize that higher group identification will lead to improved collective performance.

Social cognitive functions, associated with specific brain regions such as the orbitofrontal cortex (OFC) and dorsolateral prefrontal cortex (DLPFC), may play a role in this process. When individuals make decisions within a group, the DLPFC is believed to be involved in coordinating and aligning individual choices with the collective goals and norms (Goupil et al., 2021; Jankovic, 2014; Yang et al., 2020). This brain region is responsible for cognitive regulation and is linked to

individual decision-making processes like moral and economic decision-making (Hu et al., 2015;

Stallen et al., 2018). In a group setting, the OFC is important for evaluating the potential outcomes

and consequences of different decisions (Bechara 2004; Izquierdo, 2017; Wallis, 2007). It is

thought to regulate group decision-making processes by integrating information about social

rewards and values (Dixon et al., 2014; Wallis 2007). Based on these findings, we hypothesized

that group identification and its impact on collective performance may be tracked by the DLPFC

and OFC.

To better comprehend the influence of group identification on collective performance, it is

important to uncover the dynamic neural processes involved. Previous research has focused on

either single-brain or multi-brain findings, but social interactions are complex and cannot be fully

captured by either approach alone (Frith, and Frith, 2012; Jiang et al., 2015; Xie et al., 2020; Xie

et al., 2023). In the present study, we combined single- and multi-brain analyses to address this

limitation. While recent research has identified both decreased single-brain activation and

increased synchronization as drivers of social decision-making (Cheng et al., 2022), the

interaction between single-brain and multi-brain activities in driving social decision-making is still

not fully understood. Yang et al. (2020) found that an increased brain activation connectivity

facilitates the integration of social information into decision-making, while Xie et al. (2023)

suggested that this connectivity could link two processes (i.e., strategy switching and influence

management). Moreover, Ni et al. (2024) indicated that neural synchronization was linked to the

strength of social-emotional communication and connections between individuals. An increase in

neural synchronization has also been shown to predict the coordination and cooperation abilities

of group members (Lu et al., 2023). Therefore, we hypothesize that neural synchronization may be

related to group performance. Based on the above, our goal was to explore the regulation of

single-brain activation and multi-brain synchronization concerning brain activation connectivity,

as well as to incorporate the temporal component to fully represent the dynamic nature of brain

processes. By using this innovative approach, we can uncover the intricate interplay between

individual and collective neural processes, providing insight into how the brain dynamically

responds and adapts during social interactions.

Taken together, we examined the impact of group identification on collective performance in

a collaborative problem-solving task. Additionally, we investigated the dynamic neural

mechanisms underlying this process.

Materials and Methods

Participants

G* Power 3.1 (Faul, 2007) indicates that for *t*-tests with a medium-to-large effect size (*d*

=□0.70), an alpha level of 0.05, and a desired statistical power of 0.80 (Cohen, 1988), a sample

size of at least 52 triads is needed. In total, the data of 60 triads, consisting of 180 healthy college

students (96 females, aged 20.45 ± 2.28 years), were included. In addition, 3 participants (2

female, 20.15 ± 1.56 years) without knowledge of the experimental design details were recruited

to rate the collective performance, while another 3 participants (2 females, 21.68 ± 1.32 years)

were recruited to rate the interactive frequency. The study had full ethical approval by the

University Committee on Human Research Protection (HR2-0189-2022), East China Normal

University. Informed written consent was obtained from each participant before each experiment.

Experimental procedures and tasks

The study aimed to investigate how collective performance was affected when group

identification was either high or low. Participants came to the laboratory and were randomly

assigned to 60 groups, each consisting of 3 people of the same gender. Each triad consisted of

participants who were unfamiliar with one another and had not previously engaged in similar tasks.

They were randomly assigned to either the high group identification or low group identification

manipulation and completed the task while undergoing hyperscanning with fNIRS.

Group identification manipulation. To manipulate group identification (High Group

Identification and Low Group Identification conditions), we referred to previous studies (Efferson

et al., 2008; Yang et al., 2020). For each High Group Identification condition, the participants in

each triad were invited to chat with each other to introduce themselves and find

three-person-shared features for minutes. For each Low Group Identification condition, the

participants in each triad were asked to chat with each other about the main courses they had been

taking during this semester without being explicitly asked to find shared features (Figure 1A). The

participants rated the group identification in their group before and after the manipulation (i.e.,

group identification _1 and group identification _2) (Figure 1B). Both discussions were conducted

for the same duration of three minutes, ensuring that the number of exchanges between the two

groups remained comparable. The group identification was assessed using a 3-item questionnaire

with each item being rated on a 100-point scale ranging from 0=strongly disagree to 100=strongly

agree (Van Bavel et al., 2012). The questionnaire was presented on participants' phones. The

reliability of this scale was confirmed to be high ($\alpha = 0.87$). The three participants conducted

face-to-face offline interaction throughout the manipulation process. In addition to explaining the

next phase of the task and controlling the timer, experimenters would be isolated from

participants.

Task. We adopted a well-validated collaborative problem-solving task that incorporated the

hidden profile task to solve a murder mystery case (De Wilde et al., 2017; Stasser et al., 1992; Xie

et al., 2023). The case description contained 24 relevant arguments that were either incriminating

or exonerating for each suspect (Suspect A, B, and C). In total, each suspect had 6 incriminating

arguments presented against them. Additionally, Suspects B and C each had three exonerating

arguments to their defense, while Suspect A did not. Accordingly, when combining all 24 relevant

arguments, suspect A was the real guilty suspect, while suspects B and C could be ruled out

because of the exonerating clues. However, each group member was not privy to all of the

relevant information. To uncover the truth, individuals had to exchange and combine their

knowledge with the knowledge of their group members, as the common information incorrectly

indicated that Suspects B or C were guilty.

Procedure. To start the procedure, participants were given 3 minutes to rest (Figure 1D).

They sat around a table without partitions between each other. Building on prior group

decision-making research (Stasser et al., 1992; Xie et al., 2023), we refined all stages to enhance

controllability throughout the process (i.e., a. Reading information, b. Sharing private information,

c. Discussing information, d. Decision). Each group member was then given 18 common and 2

private pieces of information, which they read in 5 minutes (Figure 1D). Each participant would

get a piece of paper, which presented the information. Participants could read independently.

Subsequently, each triad was required to complete verbal information exchange, comprising both

group sharing and group discussion (Figure 1D) (Xie et al., 2023). During the group sharing,

participants entered Tencent Meeting via their mobile phones and were able to text their private

information in the chat box to their group members for 5 minutes. During the group discussion,

each group discussed the information that had been disclosed orally for 20 minutes. Participants

were sitting and communicating around a table. The distance between adjacent participants was

about 15 cm, and the distance between face-to-face participants was about 40 cm. In this process

of discussion, the participants were able to communicate face-to-face and verbally. After

discussion, all triads were given 5 minutes to answer the following questions (i) the probability of

three suspects, 0%-100% for each suspect; (ii) the motivation and tool of crime; and (iii) deduced

the entire process of crime. The three questions were presented on paper, allowing participants to

write their answers directly on the same sheet. Subsequently, three independent raters used these

paper questionnaires to record and calculate the scores for each group.

163
Figure 1. Experimental procedure. (**A**) Group identification manipulation. For each High Group Identification

condition, the participants in each triad were invited to chat with each other to introduce themselves and find

three-person-shared features for minutes. For each Low Group Identification condition, the participants in each

triad were asked to chat with each other about the main courses they had been taking during this semester without

being explicitly asked to find shared features. (**B**) The rate of group identification. The participants in each triad

rated the group identification in their group before and after the manipulation. (**C**) Group identification

manipulation check. We examined how the level of group identification changed when we manipulated it, for both

High and Low Group Identification conditions. Group identification_1, Group identification before manipulation;

Group identification _2, Group identification after manipulation. (**D**) The procedure of task. First, participants

completed a series of individual difference questionnaires before the task. Then, each group member received 18

common information and 2 private information. They read their information within 5 minutes. After that, each

triad was required to complete verbal information exchange, comprising both group sharing and group discussion

Each group member texted the private information to other members by Tencent Meeting during group sharing for

5 minutes, and they discussed the information currently disclosed with others orally during the group discussion

for 20 minutes. Ultimately, the groups had a period of 5 minutes to answer the questions.

fNIRS data acquisition

In this study, the brain activities of participants in each group were simultaneously recorded

with fNIRS using an ETG-7100 optical topography system (Hitachi Medical Corporation, Japan).

The absorption of near-infrared light (two wavelengths: 695 and 830 nm) was measured with a

sampling rate of 10 Hz. The oxyhemoglobin (HbO) and deoxyhemoglobin (HbR) were obtained

under the modified Beer-Lambert law. We focused our analyses on the HbO signal for the

following reasons: (i) HbO concentration is sensitive to changes in regional cerebral blood flow

(Hoshi, 2003); (ii) the HbO signal was reported to have a higher signal-to-noise ratio than the HbR

signal (Mahmoudzadeh et al., 2013); and (iii) an increasing number of studies have revealed

neural synchronization based on the HbO signal (Yang et al., 2020).

Two optode probe sets were used to cover each participant's prefrontal and left TPJ regions

(Figure S1). The DLPFC plays a crucial role in group decision-making processes, with findings

suggesting that individuals exhibiting reduced prefrontal activity were more prone to out-group

exclusion and demonstrated stronger in-group preferences (Goupil et al., 2021; Jankovic, 2014;

Yang et al., 2020). Similarly, the left TPJ has been previously reported to be associated with

decision-making and information exchange (Freitas et al., 2019; Tindale et al., 2019). For each

participant, one 3×5 optode probe set (8 emitters and 7 detectors forming 22 measurement points

with 3 cm optode separation, see Table S1 for detailed MNI coordinates) was placed over the

prefrontal cortex (reference optode is placed at Fpz, following the international 10-20 system for

positioning). The other 2×4 probe set (4 emitters and 4 detectors forming 10 measurement points

with 3 cm optode separation, see Table S2 for detailed MNI coordinates) was placed over the left

TPJ (reference optode is placed at T3, following the international 10-20 system for positioning).

The probe sets were examined and adjusted to ensure consistency of the positions across the

participants. After the completion of data collection, we utilized the Vpen positioning system to

accurately locate the detection light poles, ultimately obtaining the MNI positioning coordinates.

Behavioral analyses

Group identification. To examine and quantify the manipulation of group identification, the

scores of the participants in each triad were averaged to determine the group identification, and an

ANOVA with repeated measures was performed to examine group identification, with the levels of

group identification (High/Low) serving as a between-subjects variable, and the orders of rating

group identification (group identification_1/group identification_2) as a within-subjects variable

(Figure 1C). Moreover, the Pearson's correlation was used to examine the relationship between

group identification_2 and collective performance.

Individual performance. Individual performance was evaluated based on participants'

accuracy in solving the case, including their probability estimates for the three suspects (0%-100%

each, 2 points per suspect, totaling 6 points), identification of the crime's motivation (1 point) and

tool (1 point), and deduction of the full crime process (20 points). Three independent raters

assessed each participant's performance, demonstrating high consistency (Cronbach's $\alpha = 0.89$).

Collective performance. Collective performance was calculated by averaging the individual

scores assigned by the three raters for each group, representing the group's overall accuracy in the

case solution.

An independent *t*-test was conducted to examine collective performance, with group

identification levels being the independent variable and collective performance as the dependent

variable. Moreover, we employed a regression model to examine how varying levels of group

identification affect collective performance, using group identification scores as the independent

variable and collective performance as the dependent variable.

The similarity in individual-collective performance. After the z-score normalization of

each item for individual and collective performance, we calculated the Euclidean distance (Eq. 1)

between individual and collective performance. In Eq.1, *x* is the individual score, *y* is the

collective score (y is calculated from the three per capita scores), and *i* stands for the group

number. So, x_i means the individual score of participants in the *i* group, and y_i means the collective

score of the *i* group. *d (x, y)* represents the distance from the individual to the collective score. A

smaller distance indicated a higher similarity in individual-collective performance, while a larger

distance suggested a lower similarity in individual-collective performance.

$$d(x, y) = \sqrt{\sum (x_i - y_i)}$$ **Eq. 1**

To investigate whether there was a significant difference in the similarity in

individual-collective performance between conditions, an independent t-test was conducted with

group identification levels as the independent variable and the similarity in individual-collective

performances as the dependent variable.

fNIRS data analyses

Overview. We aimed to investigate the neural mechanisms underlying the impact of different

levels of group identification on collective performance. (1) To do this, we sought to examine

whether the individual differences in individual performance were reflected in single-brain

activations. We first identified task-related brain regions and compared the single-brain activation

of different levels of group identification. We then examined the correlation between single-brain

activation and individual performance, and tested whether the relationship between an individual's

perceived group identification and individual performance was mediated by single-brain activation.

(2) We sought to examine whether the group differences in collective performance were reflected

in within-group neural synchronization (GNS). We first identified task-related GNS and compared

GNS of different levels of group identification. We then examined the correlation between

collective performance and GNS, and tested whether the relationship between the group

identification score of each triad and collective performance was mediated by GNS. (3)

Examining brain activation connectivity, we sought to bridge single-brain activations and the

corresponding GNS, thus unifying individual decision-making and collective performance. We

first identified task-related brain activation connectivity and compared brain activation

connectivity of different levels of group identification. We then examined the correlation between

the similarity in individual-collective performance and brain activation connectivity, and tested

whether the relationship between the individual's single-brain activations and the corresponding

GNS was mediated by brain activation connectivity.

Pre-processing approach. We sought to explore the neural mechanisms that manipulated

group identification and its effect on collective performance. Data were preprocessed using the

Homer2 package in MATLAB 2020b (Mathworks Inc., Natick, MA, USA). First, motion artifacts

were detected and corrected using a discrete wavelet transformation filter procedure. After that,

the raw intensity data were converted to optical density (OD) changes. Then, kurtosis-based

wavelet filtering (Wav Kurt) was applied to remove motion artifacts with a kurtosis threshold of

3.3 (Chiarelli, Maclin, Fabiani, & Gratton, 2015). Based on a prior multi-brain study of social

interactions (Cheng et al., 2022), the output was bandpass filtered using a Butterworth filter with

order 5 and cut-offs at 0.01 and 0.5 Hz to remove longitudinal signal drift and instrument noise.

Finally, OD data were converted to HbO concentrations.

Single-brain activation. The detailed steps are as follows:(1)We analyzed the data using

SPM-based software (Ye et al., 2009), focusing on the time series from information reading to

decision-making, and extracted the oxyhemoglobin (HbO) signal for each participant and triad.

The onsets and durations of these time series were used to generate the stimulus design, which

was convolved with a canonical hemodynamic response function (HRF) using NIRS-SPM. (2) We

then employed a general linear model (GLM) to fit the predicted signals to the actual data,

obtaining beta estimates (regression coefficients) for each parameter in the single-subject design

matrices. Within the GLM framework, contrast maps were generated to compare activation

between the baseline (resting phase) and task sessions (encompassing information reading to

decision-making) for each subject, based on the HbO signal. Next, the regression factor was

convolved with the HRF, and regression analysis provided brain activation values representing

decision-making activity for each participant across all task stages and channels. For group-level

analysis, we conducted second-level, random-effects analyses (Friston et al., 2007) by aggregating

the β values across participants. A one-sample t-test was performed for each channel to determine

whether mean activation significantly exceeded zero (i.e., greater task-related activation than rest),

with p values corrected for false discovery rate (FDR; $p < 0.05$, Benjamini & Hochberg, 1995). (3)

Channels showing significant single-brain activations were identified as task-related and included

in further analyses, and the results were visualized on a standard MNI brain template using

summary statistics.

Subsequently, we conducted independent t-tests on single-brain activations in task-related

brain regions, with group identification levels as the independent variable. The p values were also

adjusted to control for FDR ($p < 0.05$; Benjamini & Hochberg, 1995). Then, we used Pearson's

correlation analyses to investigate the relationship between single-brain activation and individual

performance (i.e., evaluated based on participants' accuracy in solving the case). Finally, we

employed the PROCESS model 4 to construct a mediation model with 5000 bootstrap resamples

(Preacher, & Hayes, 2008) to examine the relationship between an individual's perceived group

identification and individual performance, which was mediated by single-brain activation.

Within-group neural synchronization (GNS). After pre-processing, GNS was used as the

neural index (i.e., interpersonal brain activities that co-vary along the time course). Concerning

GNS, and similar to previous studies (Pan, Cheng, Hu, 2021; Yang et al., 2020), the wavelet

transform coherence (WTC) (Eq. 2) was used to assess the cross-correlation between two oxy-Hb

time series of pairs of participants. Here, t denotes the time, s indicates the wavelet scale, $\langle \cdot \rangle$

represents a smoothing operation in time, and W is the continuous wavelet transform (Grinsted,

Moore, & Jevrejeva, 2004). Within each triad (taking one triad with subject IDs of 1, 2, and 3 as

an example), WTC was applied to generate the brain-to-brain coupling of each pair in each triad

(Coherence1&2, Coherence 1&3, and Coherence 2&3). Then, three coherence values from three

pairs were averaged as the GNS for each triad, that is, GNS = (Coherence 1&2 + Coherence 1&3

+ Coherence 2&3) / 3.

$$WTC(t, s) = \frac{\left| \langle s^{-1} w^{ij}(t,s) \rangle \right|^2}{\left| \langle s^{-1} w^{i}(t,s) \rangle \right|^2 \left| \langle s^{-1} w^{j}(t,s) \rangle \right|^{2}}$$ **Eq. 2**

Regarding the first step, we estimated whether GNS was enhanced during the task compared

to the baseline. Time-averaged GNS (also averaged across channels in each group) was compared

between the baseline session (i.e., the resting phase) and the task session (from reading

information to making decisions) using a series of one-sample t-tests. Here, p-values were

thresholded by controlling for FDR (p < 0.05; Benjamini & Hochberg, 1995). When determining

the frequency band of interest, the time-averaged GNS was also averaged across channels. After

that, we analyzed the time-averaged GNS of each channel. Then, channels showing significant

GNS were regarded as regions of interest and included in subsequent analyses. An independent

t-test was conducted on GNS, with group identification levels being the independent variable.

Here, *p* values were thresholded by controlling for FDR ($p < 0.05$; Benjamini & Hochberg, 1995).

After that, the nonparametric permutation test was conducted on the observed interaction effects

on GNS of the real group against the 1,000 permutation samples. By pseudo-randomizing the data

of all participants, a null distribution of 1000 pseudo-groups was generated (e.g., time series from

member 1 in group 1 were grouped with member 2 in group 2 & member 3 in group 3). The GNS

of 1,000 reshuffled pseudo-groups was computed, and the GNS of the real groups was assessed by

comparing it with the values generated by 1000 reshuffled pseudo-groups. To provide a complete

picture of the underlying neural features, we also analyzed the GNS based on the HbR signal (see

Supplementary Materials). Second, the Pearson's correlation between GNS and collective

performance (i.e., calculated by averaging the individual scores assigned by the three raters for

each group) was performed. It is plausible that neural synchronization is closely associated with

group identification, and collective performance, suggesting that it serves as a promising

mechanism to explore how group identification influences collective outcomes. Moreover,

previous research has established that neural synchronization facilitates the emergence of group

identification, and the degree to which neural synchronization occurs among group members may

shape how individuals identify with the group (Xie et al., 2023; Reinero et al., 2021). Ultimately,

PROCESS model 4 with 5000 bootstraps resamples was used to test how GNS mediated the

relationship between group identification and collective performance (Preacher et al., 2008).

The brain activation connectivity. Studies in neuroimaging have indicated that brain

activation connectivity could be useful in understanding brain functional integration (Lu et al.,

2010; Montero-Hernandez et al., 2019; Yang et al., 2020). Consequently, an exploratory analysis

was conducted to test the hypothesis that brain activation connectivity could support the

connection between an individual's single-brain activations and the corresponding group's GNS,

thereby linking individual decision-making and collective performance.

To explore this hypothesis, we first isolated HbO brain activity associated with individual and

collective performance. Following that, we analyzed Pearson's correlations between the original

HbO data in the region related to individual and collective performance, denoted as brain

activation connectivity (Lu et al., 2010). Subsequently, we carried out one-sample *t*-tests on brain

activation connectivity to ascertain if there was any connection to the task. Furthermore, independent *t*-tests were conducted on brain activation connectivity with the group identification levels as the independent variable, accounting for the FDR ($p < 0.05$; Benjamini & Hochberg, 1995). Finally, we employed correlation and mediation analyses to assess if brain activation connectivity could explain the connection between individuals' single-brain activation and the related group's GNS. We examined the connection between the similarity in individual-collective performance and the correlation of brain activation, as well as whether the impact of each individual's single-brain activation on the corresponding group's GNS was regulated by their brain activation connectivity. We utilized the PROCESS tool in SPSS to investigate the proposed moderation effect. Specifically, we applied Model 1 with 5000 bootstrap resamples to examine the interaction between the independent variable (i.e., single-brain activation) and the moderator (i.e., brain activation connectivity) in predicting the dependent variable (i.e., GNS). It is noteworthy that prior to analysis, all variables in the moderation model were mean-centered to reduce multicollinearity and improve the interpretability of interaction terms.

Dynamic analyses. Our goal was to gain a more thorough understanding of how group identification influences collective performance through neural processes, with a focus on dynamic perspectives and tracking. We used one-minute epochs to analyze the average single-brain activation, GNS, and brain activation connectivity during the task. We then plotted the time course of dynamic single-brain activation and conducted one-sample *t*-tests to identify significant single-brain activation, GNS, and the correlation of brain activation periods. To better understand the interactive behavior revealed by dynamic single-brain activation, we linked the brain activation time series with video recordings of interactive behavior.

Additional modal measures and analyses

Additionally, we aimed to gain a more comprehensive understanding of how various group identifications influence collective performance. To this end, we obtained more evidence through other methods. Previous studies have shown that the quality of information exchange, such as verbal interactions, eye contact, and smiling, is a reliable indicator of group behavior and is associated with collective performance (Dikker et al., 2022; Hirsch et al., 2018; Jiang et al., 2015;

Liu et al., 2021; Xie et al., 2023). Therefore, three independent raters were asked to rate the quality of the information exchange of each group, with a Cronbach's alpha of 0.85. The raters were guided to consider verbal interactive frequency (e.g., "I agree with you", "You're right", "I understand what you mean") and nonverbal interactive frequency (e.g., eye contact and smiling) (Jiang et al., 2015; Xie et al., 2023) (Figure 8A). The evaluation period consisted of group sharing and discussion, in which the raters evaluated the suggested items in one-minute increments. The scores of the 25 periods were then compiled. The final quality of the group's information exchange was determined by taking the average of the scores of the three raters for each group. A higher quality of the group's information exchange entails communicating more fully.

Initially, independent *t*-tests were performed to examine the quality of information exchange between the group identification levels, which was the independent variable. Subsequently, the Pearson's correlation between the quality of information exchange and collective performance was assessed.

Behavioral metrics (i.e., the quality of information exchange) offer direct indicators of the strategies and patterns individuals use during the decision-making process. Neural data (i.e., GNS) can uncover the neural activity linked to decision-making processes in the brain. For a more comprehensive insight into the processes involved in group decision-making, we performed the hierarchical multiple regression analysis with collective performance as the dependent variable to determine the weight of prediction of the quality of information exchange, GNS, and their interaction.

Results

Group identification leads to differences in collective performance

We examined the impact of manipulated group identification on increasing intergroup discrimination. First, a repeated-measures ANOVA on group identification revealed a significant interaction between the levels of group identification and the order of rating group identification ($F_{1, 58} = 6.83$, $p < 0.012$, $\eta^2_p = 0.11$; Figure 1C). Post-hoc paired *t*-tests revealed that following the manipulation, participants in the High Group Identification condition reported higher group

identification (group identification_2) than those in the Low Group Identification condition (group

identification_2) (t_{58} = 4.83, p < 0.001). This effect was diminished before manipulation (t_{58} = 1.73,

p = 0.090). These results confirmed the effectiveness and validity of our paradigm in inducing

different levels of group identification.

Next, we observed significant differences in collective performance between the High and

Low groups. Specifically, participants in the High Group Identification group demonstrated a

higher level of collective performance (t_{58} = 2.18, p = 0.034, Cohen's d = 0.58) (Figure 2A). The

results from the regression model highlighted a significant association between the degree of

group identification and collective performance (β = 0.45, t = 4.56, p = 0.019). Furthermore, the

results of the Pearson's correlation indicated that groups with higher group identification were

more likely to exhibit better collective performance (r = 0.38, p = 0.003) (Figure 2B). These

findings suggested that manipulated group identification led to variations in collective

performance.

417
Figure 2. Group identification leads to differences in collective performance. (**A**) Manipulated group identification

led to group differences in collective performance. (**B**) Group identification was positively correlated with better

collective performance. The Pearson's correlation and its associated analyses were based on the data from group

identification_2. $^{*}p$ < 0.05.

The differences in collective performance are correlated with single-brain activation

We sought to identify single-brain activations that supported individual performance. First,

by performing one-sample t-tests for single-brain activations, we observed significantly increased

single-brain activations in the DLPFC (CH4, t_{59} = 14.54, p < 0.001, FDR corrected) and the TPJ

(CH1, $t_{59} = 3.89$, $p = 0.022$, FDR corrected). Subsequently, we performed independent t-tests on

single-brain activations in channels that exhibited significant results, with group identification

levels as the independent variable. The results showed significant differences in single-brain

activations between the High and Low Group Identification groups in the DLPFC (CH4, $t_{58} = 2.71$,

$p = 0.015$, FDR corrected, Cohen's $d = 0.42$) (Figure 3A).

Furthermore, single-brain activations in the DLPFC (CH4) were associated with individual

performance ($r = 0.27$, $p < 0.001$, Figure 3B), demonstrating that greater single-brain activations

in the DLPFC (CH4) were associated with a better individual performance. The mediation model

demonstrated a satisfactory fit (CFI = 0.93, TLI = 0.93, RMSEA = 0.04) (CFI-Comparative Fit

Index; TLI-Tucker-Lewis index; RMSEA-Root-Mean-Square Error of Approximation), suggesting

that the perceived group identification of each individual affected the alterations in single-brain

activations in the DLPFC, consequently leading to variations in their performance ($\beta_a = 0.16$, $t =$

2.20, $p = 0.030$; $\beta_b = 0.26$, $t = 3.56$, $p < 0.001$; $\beta_c = 0.18$, $t = 2.34$, $p = 0.020$) (Figure 3C).

Based on the results mentioned above, the DLPFC (CH4) has been determined as the key

neural substrate responsible for individual performance.

Figure 3. The differences in collective performance are correlated with single-brain activation. (**A**) Significant

differences in single-brain activations between the High and Low Group Identification groups were observed in the

DLPFC (CH4). (**B**) Greater single-brain activation in the DLPFC (CH4) was associated with higher individual

performance. (**C**) A serial mediation model suggested that single-brain activation mediated the relationship

between group identification of each person and individual performance. $^{*}p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$.

The differences in collective performance are correlated with neural synchronization

We sought to identify neural synchronization that supported collective performance. First, by

performing one-sample t-tests for GNS, we observed a significantly increased GNS in the OFC

(CH20, $t_{59} = 1.98$, $p = 0.046$, FDR corrected; CH21, $t_{59} = 2.49$, $p = 0.024$, FDR corrected) and the

TPJ (CH1, $t_{59} = 2.16$, $p = 0.030$, FDR corrected). Subsequently, independent t-tests were

conducted on GNS in the pertinent channels, which revealed that GNS was significantly different

between the High and Low Group Identification groups in the OFC (CH21, $t_{58} = 2.21$, $p = 0.037$,

FDR corrected, Cohen's $d = 0.58$) (Figure 4A). A permutation test confirmed that the observed

interactive effects on GNS in real groups were outside the 95% CI of a null distribution

comprising 1000 pseudo groups (Figure S2). Therefore, the neural synchronization was only

found in the 'real' groups who were interacting in the task. The pattern of associated results was

similar to that of HbO when the analyses of HbR were conducted (See the Supplementary

Materials).

Results showed that greater GNS in the OFC (CH21) was associated with higher collective

performance ($r = 0.46$, $p = 0.001$; Figure 4B). The mediation model demonstrated a satisfactory fit

(CFI = 0.86, TLI = 0.86, RMSEA = 0.04), suggesting that group identification of each group

caused changes in GNS in the OFC, and ultimately impacted the corresponding collective

performance ($\beta_a = 0.29$, $SE = 0.01$, $t = 2.24$, $p = 0.026$; $\beta_b = 0.25$, $SE = 7.28$, $t = 1.88$, $p = 0.049$; β_c

= 0.37, $SE = 0.01$, $t = 2.24$, $p = 0.005$) (Figure 4C).

Through correlation and regression model analysis, we found that in group decision-making,

the increase in group identity would affect group performance by improving GNS in the OFC

brain region.

Figure 4. The differences in collective performance are correlated with neural synchronization. (**A**) A significant difference between the High and Low Group Identification groups in the OFC (CH21) was observed (*p*-value, FDR corrected). (**B**) Greater GNS in the OFC (CH21) was associated with higher collective performance. (**C**) A mediation model suggested that GNS mediated the relationship between group identification of each group and collective performance. $^{*}p < 0.05$, $^{**}p < 0.01$.

Brain activation connectivity links the single-brain activation and the corresponding GNS

Given the above association between individuals' single-brain activations and individual performance, as well as GNS and collective performance, we subsequently explored individuals' single-brain activations and the corresponding group's neural synchronizations were linked by brain activation connectivity.

We first extracted the HbO brain activities related to individual performance (e.g., DLPFC, CH4) and collective performance (e.g., OFC, CH21) of each group member and conducted a Pearson's correlation between the two. The outcome of DLPFC-OFC connectivity was then utilized as brain activation connectivity in the following analyses. By performing one-sample *t*-tests, the DLPFC-OFC (CH4-CH21) correlation showed a significant increase during the task ($t_{59} = 9.198$, $p < 0.001$, FDR corrected). Subsequently, independent *t*-tests were conducted on DLPFC-OFC connectivity, which revealed that DLPFC-OFC connectivity was significantly

different between the High and Low Group Identification groups (t_{58} = 4.01, p < 0.001, FDR

corrected, Cohen's d = 0.61) (Figure 5A). An independent t-test was also conducted on the

corresponding behavioral indicators (i.e., the similarity in individual-collective performance) and

revealed a significant difference between the High and Low Group Identification groups (t_{58} =

4.03, p < 0.001, Cohen's d = 0.62) (Figure 5B). Subsequently, Pearson's correlation was used to

test whether individual differences in the similarity in individual-collective performance were

reflected by DLPFC-OFC connectivity. It was observed that the stronger DLPFC-OFC

connectivity was linked to a decrease in the distance between individual and collective

performance (r = -0.17, p = 0.026), indicating that stronger DLPFC-OFC connectivity led to a

greater similarity in individual-collective performance (Figure 5C). Finally, the relationship

between individual single-brain activation and the corresponding GNS was regulated by

DLPFC-OFC connectivity (β_{int} = 0.20, SE = 1.37, t = 2.22, p = 0.028) (Figure 5D).

The findings implied that DLPFC-OFC connectivity could link the individual's single-brain

activations and the group's GNS, thus unifying individual performance and collective

performance.

Figure 5. Brain activation connectivity links the single-brain activation and the corresponding GNS. (**A**) A

significant difference between the High and Low Group Identification groups in the DLPFC-OFC (CH4-CH21)

correlation was observed (p-value, FDR corrected). (**B**) A significant difference between the High and Low Group

Identification groups in the similarity in individual-collective performance was observed. (**C**) A stronger

DLPFC-OFC (CH4-CH21) correlation was linked to a decrease in the distance between individual and collective

performance. (**D**) The analyses suggested that the relationship between individual single-brain activation and the

corresponding GNS was regulated by brain activation connectivity. $^{*}p < 0.05$, $^{***}p < 0.001$.

The dynamic neural process

To better understand how group identification influences collective performance at a neural

level, we examined the dynamic activation of single brains, the group neural synchronization

(GNS), and brain activation connectivity throughout the entire task. Our research findings

revealed a significant increase in single-brain activation at approximately 7 minutes into the task,

during the group sharing stage ($t = 9.88$, $p < 0.001$; green line in Figure 6). Furthermore, there was

a notable increase in brain activation connectivity around 12 minutes into the task, during the

group discussion stage ($t = 4.70$, $p = 0.013$; black line in Figure 6), and GNS at approximately 17

minutes into the task, also during the group discussion stage ($t = 3.01$, $p = 0.042$; orange line in

Figure 6).

Considering the entire dynamic process, there was a time delay in the transition from

individual to collective decisions, and brain activation connectivity was in the middle. In addition

to the regulation model provided above, it further supported the processing of information flow in

the single-brain activation, DLPFC-OFC connectivity, and ultimately the GNS.

Figure 6. The dynamic neural process. There was a time delay in the transition from individual to collective

decisions, and brain activation connectivity was in the middle. The processing of information flow in the

single-brain activation (there was a significant increase in single-brain activation at approximately 7 minutes into

the task), DLPFC-OFC connectivity (there was a significant increase in connectivity at approximately 12 minutes

into the task), and ultimately the GNS (there was a significant increase in GNS at approximately 17 minutes into

the task). The red star sign indicates that at this time point, the neural signal began to increase significantly.

The two-in-one neural model of group identification influences collective performance

Building on the above results, we have developed a two-in-one neural model that explains

how group identification influences collective performance. This descriptive model aims to

illustrate the potential interrelationships among these indicators and establish a conceptual

framework to inspire forthcoming research endeavors. In the first step, group identification

influences individual performance, which is associated with significant single-brain activation in

the DLPFC of each group member. In the second step, group identification influences collective

performance, which is linked to significant within-group neural synchronization (GNS) in the

OFC. These two steps are linked by the DLPFC-OFC connectivity, which modulates the

relationship between individual DLPFC activation and GNS in the OFC. In particular, people with

a strong sense of group identification tend to have better individual performance, which is

associated with increased single-brain activation in the DLPFC of each group member. This, in

turn, leads to improved collective performance, which is linked to higher levels of within-group

neural synchronization (GNS) in the OFC. On the other hand, groups with a low sense of group

identification tend to have poorer individual performance, which is associated with decreased

activity in the DLPFC. This also results in poorer collective performance, which is correlated with

lower levels of GNS in the OFC. The DLPFC-OFC connectivity modulates the relationship

between the single-brain activation in the DLPFC and the corresponding group's neural

synchronizations in the OFC. This influences the similarity in individual and collective

performance, thus connecting individual performance and collective performance.

Figure 7. The two-in-one neural model explains how group identification influences collective performance. In the first step, group identification influences individual performance, which is associated with significant single-brain activation in the DLPFC of each group member. In the second step, group identification influences collective performance, which is linked to significant within-group neural synchronization (GNS) in the OFC. These two steps are connected by the DLPFC-OFC connectivity, which modulates the relationship between individual DLPFC activation and GNS in the OFC.

The quality of information exchange is correlated to the effect of group identification

To gain a better understanding of how various group identifications led to differences in collective performance, we gathered further evidence by assessing the quality of information exchange from the video. Independent *t*-tests were conducted on the quality of information exchange, which revealed that the quality of information exchange was significantly different between the High and Low Group Identification groups ($t_{58} = 2.53$, $p = 0.014$, Cohen's $d = 0.67$) (Figure 8B). Pearson's correlation showed that the higher quality of information exchange, the better collective performance ($r = 0.36$, $p = 0.007$) (Figure 8C). The results of our study indicated that interactive frequency played a role in indicating the relationship between group identification and collective performance.

The hierarchical multiple regression analysis showed that the model including both GNS and the quality of information exchange was the most effective in predicting collective performance

(R^2 = 0.32, SE = 3.33). Additionally, the model with GNS alone (R^2 = 0.27, SE = 4.17)

outperformed the model with only the quality of information exchange (R^2 = 0.24, SE = 4.82) in

predicting collective performance. These results indicated that combining GNS and the

information exchange quality can accurately predict collective performance. Moreover, GNS

proved to be a more reliable predictor of collective performance than behavioral metrics.

Figure 8. The quality of information exchange is correlated to the effect of group identification. (**A**) A significant

difference between the High and Low Group Identification groups in the quality of information exchange was

observed. (**C**) Higher quality of information exchange was linked to better collective performance. [*]$p < 0.05$.

Discussion

Previous research and social identity theory has demonstrated that variations in group

identification play a significant role in shaping collective behavior and can impact collective

performance (e.g., P¨arnamets et al., 2020; Reinero et al., 2021; Számadó et al., 2021; Solansky,

2011). Building upon existing findings and theory, our study adds information about the neural

mechanisms involved, both at the level of individual group members and at the level of the group.

Specifically, we found that group identification influenced individual performance, supported by

neural activity in the DLPFC. Moreover, we also found that high group identification enhanced

collective performance, supported by neural synchronization in the OFC. We highlighted the link

between individual performance and collective performance, which was associated with an increased correlation between the DLPFC and the OFC. Based on these results, we have developed a two-in-one neural model to illustrate the dynamic neural processes through which group identification influences collective performance.

Our findings converge with previous work which suggested that group identification is crucial in shaping collective behavior (Reinero et al., 2021; Számadó et al., 2021; Solansky, 2011). Group identification plays a significant role in shaping collective decision-making by promoting cohesion and reducing conflicts within a group (Bicchieri, 2002; Gundlach et al., 2006; Tajfel and Turner, 1979). Members collaborate to address conflicts through positive interactions to uphold positive intergroup relations and achieve better collective behavior (Brewer and Kramer, 1986; De Cremer and Van Vugt, 1999; Dikker et al., 2022; Prochazkova et al., 2022). Utilizing tools from neuroscience and video analysis, this study demonstrated the positive impact of increased communication and interaction among group members in groups with high group identification, leading to improved collective performance. These findings support existing research on the influential role of group identification in shaping collective behavior. Understanding how group identification operates can provide valuable insights into group decision-making processes and contribute to the development of strategies for effective group dynamics and decision-making.

Extending previous neuroimaging approaches, this work combined single-brain and multi-brain approaches to uncover how group identification influenced collective performance: through single-brain activation and multi-brain synchronization. Previous studies have shown that activation in the DLPFC is associated with controlled decision-making, such as moral and economic decision-making (Fecteau et al., 2007; Gläscher et al., 2012; Yang et al., 2020). Our findings further support this by demonstrating the involvement of DLPFC activation in collective decision-making during social interactions. Additionally, previous research has suggested that increased neural synchronization in the prefrontal area, including the OFC, is linked to collective performance (Liu et al., 2021; Wang et al., 2019; Xie et al., 2023). Our study expands on this by showing that neural synchrony in the OFC was correlated with collective performance. Interestingly, we found that the level of neural synchrony was higher in the high group identification group compared to the low group, underscoring the significance of group identification in influencing collective performance.

Our research suggests a new perspective on the connectivity between the DLPFC and OFC in

the brain, showing its potential role in understanding brain functional integration and the

relationship between individual brain activation and collective brain synchronization. While we

did not directly determine the reason for this connectivity, it may indicate a form of 'social

alignment' between the execution system (individual decision-making) and observation system

(collective decision-making) in the human brain (Adhikari, Goshorn, Lamichhane, & Dhamala,

2013; Schenk & Colloca, 2020; Yang et al., 2020). Our study aims to uncover the neural

mechanisms behind the impact of group identification on collective performance. We have

developed a two-in-one neural model that proposes the connectivity between the DLPFC and OFC

as a key mechanism for integrating individual and collective decision-making processes in the

brain.

Our study also has demonstrated significant increases in single-brain activation,

DLPFC-OFC functional connectivity, and GNS at 7, 12, and 17 minutes, respectively, following

task initiation. The significant increase in these neural activities together constructs the two-in-one

neural model that explains how group identification influences the collective performance we

proposed. As previous studies have shown, the BOLD signal collected by fNIRS is slowly

increasing compared to neuronal activity, which means that it has hysteresis (Turner et al., 1998).

In social interactions such as group decision-making, the time of neural synchronization is delayed

because people need to spend time increasing the number of dialogues to improve collaboration

efficiency and form the same preference (Zhang et al., 2019). For example, participants would

exhibit significant neural alignment, but only after they had completed a period of dialogue

(Sievers et al., 2024). In the task of cooperation, with the improvement of cooperation efficiency

between two participants, the higher degree of neural synchronization (Cui et al., 2012). Therefore,

the generation of neural synchronization depends on the interaction over a period of time, which

can affect the estimation of collaboration time. Prior research has shown that when the teaching

task with prior knowledge began 50-55 seconds, significant neural synchronization could be

generated between teacher and students, which meant that students and teacher achieved the same

goal of learning knowledge (Liu et al., 2019). Moreover, a noteworthy increase in GNS was

observed approximately 6 minutes into the group discussion period for better discussing and

solving the problem (Xie et al., 2023). These findings are similar to ours. Therefore, the time

points we found could reflect the dynamic time change of the neural process of team collaboration.

Our study has raised several interesting issues for further investigation. First, our laboratory task

was specifically created to assess collective performance. While we did observe significant

differences in collective performance between groups with high and low levels of identification,

future research needs to utilize a trial-by-trial problem-solving task rather than relying on

subjective evaluations. This approach would allow for the examination of collective performance

as a continuous variable, rather than being reliant on ratings provided by external raters. Second,

to enhance the validity of our findings, we utilized fNIRS technology to track both individual and

group brain activity simultaneously. Even though our primary focus was on the left temporal

regions, specifically lTPJ, we acknowledged that other brain regions such as rTPJ, vmPFC, IFG,

and right temporoparietal regions also played a role in Theory of Mind (ToM) tasks (Jiang et al.,

2015; Liu et al., 2023; Stolk et al., 2016) due to the constraints of fNIRS channels. Future studies

should consider utilizing MEG technology, which offers better spatial and temporal resolutions, to

further investigate the neural mechanisms underlying social cognition and collective behavior.

Although our study has found a new perspective, the analysis method still refers to and uses the

traditional fNIR-based hyperscanning analyses (Liu et al., 2019; P̈arnamets et al., 2020; Reinero

et al., 2021; Számadó et al., 2021; Solansky, 2011), which is generally accepted by the majority of

fNIR-based hyperscanning researchers. For example, we would first identify significant channels

through a one-sample t-test and then conduct further analyses, such as ANOVA or independent

samples t-tests. Selective analysis is a powerful tool and is perfectly justified whenever the results

are statistically independent of the selection criterion under the null hypothesis (Kriegeskorte et al.,

2019). However, it may lead to double dipping and missing information. In this study, the absence

of statistically significant TPJ activation in the analyzed data led to the TPJ being ignored. In the

future, it should be made explicit in the analysis, and the reliability of the results should be

ensured by appropriate statistical methods (e.g., cross-validation, independent data sets, or

techniques to control for selective bias). Next, our experimental design intentionally separated task

phases (reading, sharing, discussing, and decision) to enhance procedural clarity, experimental

control and ecological validity (Xie et al., 2023), departing from the integrated approaches used in

prior work (De Wilde et al., 2017; Stasser et al., 1992). However, the fixed (non-counterbalanced)

phase sequence introduces an important limitation: observed effects could potentially reflect

temporal factors rather than genuine task-related neural activity. While our analytical approach

controlled for some of these temporal confounds, future studies would benefit from implementing

counterbalanced phase sequences, jittered trial timing, or additional control tasks to more precisely

isolate phase-specific neural correlates. Finally, future research should investigate additional

variables, including sex differences and measures of attractiveness or hierarchy among

participants, such as students versus teachers.

To summarize, the present study revealed and connected the individual and group brain

processes involved in group identification that influenced collective performance, and constructed

a descriptive model to illustrate this process. This research not only enhances our understanding of

the neurobiological aspects of group dynamics and collaboration, but also offers a comprehensive

framework that synthesizes previous research findings and provides a theoretical foundation for

future investigations into the complexities of social interactions.

Data and Code Availability Statement

702 Data of this project were obtained from the GitHub repository (available at

703 https://github.com/xehui/group-identification). The code is only to be made available via a request

704 to the Authors, which requires a formal data sharing agreement.

Acknowledgments

706 This work is supported by the National Natural Science Foundation of China (71942001), STI

707 2030—Major Projects （2021ZD0200500）, the National Natural Science Foundation of China

708 (32071082), Key Specialist Projects of Shanghai Municipal Commission of Health and Family

709 Planning (ZK2015B01), the Programs Foundation of Shanghai Municipal Commission of Health

710 and Family Planning (201540114).

Reference

712 Adhikari, B. M., Goshorn, E. S., Lamichhane, B., & Dhamala, M. (2013). Temporal-order judgment of

audiovisual events involves network activity between parietal and prefrontal cortices. *Brain connectivity, 3*(5), 536-545.

Bechara, A. (2004). The role of emotion in decision-making: Evidence from neurological patients with orbitofrontal damage. *Brain and cognition, 55*(1), 30-40.

Bicchieri, C. (2002). Covenants without swords: group identity, norms, andcommunication in social dilemmas. *Rationality and Society, 14*(2), 192–228.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B: Methodological, 57*(1), 289-300.

Brewer, M.B., Kramer, R.M. (1986). Choice behavior in social dilemmas: effects of social identity, group size, and deci-sion framing. *Journal of Personality and Social Psychology, 50*(3), 543–49.

Cohen, J. (1988). Statistical power ANALYSIS for the Behavioral sciences. *Journal of the American Statistical Association, 2nd*(334).

Cheng, X. J., Zhu, Y. J, Hu, Y. Y, Zhou, X. L., Pan, Y. F., Hu, Y. (2022). Integration of social status and trust through interpersonal brain synchronization. *Neuroimage, 246*:118777.

Cui, X., Bryant, D. M., & Reiss, A. L. (2012). NIRS-based hyperscanning reveals increased interpersonal coherence in superior frontal cortex during cooperation. *NeuroImage, 59*(3), 2430-2437.

Dixon, M. L., & Christoff, K. (2014). The lateral prefrontal cortex and complex value-based learning and decision making. *Neuroscience & Biobehavioral Reviews, 45*, 9-18.

Dikker, S., Mech, E. N., Gwilliams, L., West, T., Dumas, G., & Federmeier, K. D. (2022). Exploring age-related changes in inter-brain synchrony during verbal communication.

De Wilde, T. R., Ten Velden, F. S., & De Dreu, C. K. (2017). The neuropeptide oxytocin enhances information sharing and group decision making quality. *Scientific reports*, 7(1), 1-8.

De Freitas, J., Thomas, K., DeScioli, P., & Pinker, S. (2019). Common knowledge, coordination, and strategic mentalizing in human social life. *Proceedings of the National Academy of Sciences, 116*(28), 13751-13758.

De Cremer, D., Van Vugt, M. (1999). Social identification effectsin social dilemmas: A transformation of motives. *European Journal of Social Psychology, 29*(7), 871–93.

Friston, K.J., Ashburner, J.T., Kiebel, S.J., Nichols, T.E., Penny, W.D.. (2007). Statistical Parametric Mapping: The Analysis of Functional Brain Images. Elsevier/Academic Press, London.

Eferson, C., Lalive, R. & Fehr, E. (2008). The coevolution of cultural groups and in-group favoritism. *Science, 321,* 1844–1849.

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior research methods, 39*(2), 175-191.

Frith, C. D. & Frith, U. (2012). Mechanisms of social cognition. *Annual Review Psychology. 63*, 287–313.

Fecteau, Shirley, Knoch, Daria, Fregni, & Felipe, et al. (2007). Diminishing risk-taking behavior by modulating activity in the prefrontal cortex: a direct current stimulation study. *Journal of Neuroscience, 27*(46), 12500-12505.

Gläscher, J., Adolphs, R., Damasio, H., Bechara, A., & Tranel, D. . (2012). Lesion mapping of cognitive control and value-based decision making in the prefrontal cortex. *Proceedings of the*

National Academy of Sciences of the United States of America, 109(36), p.14681-14686.

Grinsted, A., Moore, J. C., & Jevrejeva, S. (2004). Application of the cross wavelet transform and wavelet coherence to geophysical time series. *Nonlin. Processes Geophys., 11*(5/6), 561-566.

Gläscher, J., Adolphs, R., Damasio, H., Bechara, A., & Tranel, D. . (2012). Lesion mapping of cognitive control and value-based decision making in the prefrontal cortex. *Proceedings of the National Academy of Sciences of the United States of America, 109*(36), p.14681-14686.

Goupil, L., Wolf, T., Saint□Germier, P., Aucouturier, J. J., & Canonne, C. (2021). Emergent shared intentions support coordination during collective musical improvisations. *Cognitive Science, 45*(1), e12932.

Gundlach, M., Zivnuska, S., & Stoner, J. (2006). Understanding the relationship between individualism–collectivism and team performance through an integration of social identity theory and the social relations model. *Human relations, 59*(12), 1603-1632.

Hoshi, Y.. (2003). Functional near-infrared optical imaging: utility and limitations in human brain mapping. *Psychophysiology, 40*, 511–520.

Hirsch, J., Adam Noah, J., Zhang, X., Dravida, S., & Ono, Y. (2018). A cross-brain neural mechanism for human-to-human verbal communication. *Social cognitive and affective neuroscience, 13*(9), 907-920.

Hu, Y. , Sabrina, S. , & Bernd, W. . (2015). Helping or punishing strangers: neural correlates of altruistic decisions as third-party and of its relation to empathic concern. *Frontiers in Behavioral Neuroscience, 9*(24).

Izquierdo, A. (2017). Functional heterogeneity within rat orbitofrontal cortex in reward learning and decision making. *Journal of Neuroscience, 37*(44), 10529-10540.

Jiang, J., Chen, C., Dai, B., Shi, G., Ding, G., Liu, L., & Lu, C. (2015). Leader emergence through interpersonal neural synchronization. *Proceedings of the National Academy of Sciences, 112*(14), 4274-4279.

Jankovic, M. (2014). Communication and shared information. *Philosophical Studies, 169*(3), 489-508.

Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S. F., & Baker, C. I. (2009). Circular analysis in systems neuroscience: the dangers of double dipping. *Nature Neuroscience, 12*(5), 535-540.

Kerr, N. L., & Tindale, R. S. (2003). Group performance and decision making. *Annual Review of Psychology, 55*, 2201-2232.

Liu, T., Duan, L., Dai, R., Pelowski, M., & Zhu, C. (2021). Team-work, team-brain: exploring synchrony and team interdependence in a nine-person drumming task via multiparticipant hyperscanning and inter-brain network topology with fNIRS. *NeuroImage, 237*, 118147.

Lu, C. M. , Zhang, Y. J. , Biswal, B. B. , Zang, Y. F. , Peng, D. L. , & Zhu, C. Z. . (2010). Use of fnirs to assess resting state functional connectivity. *Journal of Neuroscience Methods, 186*(2), 242-249.

Lu, H., Wang, X., Zhang, Y., Huang, P., Xing, C., Zhang, M., & Zhu, X. (2023). Increased interbrain synchronization and neural efficiency of the frontal cortex to enhance human coordinative behavior: A combined hyper-tES and fNIRS study. *NeuroImage, 282*, 120385-120385.

Liu J., Zhang R., Geng B., Zhang T., Yuan D., Otani S., &Li X. (2019) Interplay between prior knowledge and communication mode on teaching effectiveness: Interpersonal neural synchronization as a neural marker. *NeuroImage, 193*,93-102.

Liu, J., Zhang, R., Xie, E., Lin, Y., Chen, D., Liu, Y., ... & Li, X. (2023). Shared intentionality modulates interpersonal neural synchronization at the establishment of communication system.

Communications Biology, 6(1), 832.

Mahmoudzadeh, M., Dehaene-Lambertz, G., Fournier, M., Kongolo, G., Goudjil, S., Dubois, J., . . .

Wallois, F. (2013). Syllabic discrimination in premature human infants prior to complete

formation of cortical layers. *Proceedings of the National Academy of Sciences, 110*(12),

4846-4851.

Montero-Hernandez, S. , Orihuela-Espina, F. , Sucar, E. , Pinti, P. , Hamilton, A. , Burgess P. , &

Tachtsidis I. (2018). Estimating functional connectivity symmetry between oxy-and

deoxy-haemoglobin: implications for fnirs connectivity analysis. *Algorithms*, *11*.

Ni, J., Yang, J., & Ma, Y. (2024). Social bonding in groups of humans selectively increases inter-status

information exchange and prefrontal neural synchronization. *Plos Biology, 22*(3),

e3002545-e3002545.

Pan, Y. , Cheng, X. , & Hu, Y. . (2021). Three heads are better than one: cooperative learning brains

wire together when a consensus is reached. *Cerebral Cortex*.

Preacher, K.J., & Hayes, A.F.. (2008). Asymptotic and resampling strategies for assessing and

comparing indirect effects in multiple mediator models. *Behav Res Methods, 40*(3), 879-891.

Prochazkova, E., Sjak-Shie, E., Behrens, F., Lindh, D., & Kret, M. E. (2022). Physiological synchrony

is associated with attraction in a blind date setting. *Nature Human Behaviour, 6*(2), 269-278.

P̈arnamets, P., Shuster, A., Reinero, D.A., Van Bavel, J.J. (2020). A value-based framework for

understanding cooperation. *Current Directions in Psychological Science, 29*(3), 227–34.

Reinero, D. A., Dikker, S., & Van Bavel, J. J. (2021). Inter-brain synchrony in teams predicts collective

performance. *Social cognitive and affective neuroscience, 16*(1-2), 43-57.

Sievers, B., Welker, C., Hasson, U., Kleinbaum, A. M., & Wheatley, T. (2024). Consensus-building

conversation leads to neural alignment. *Nature Communications,15*(1), 3936-3936.

Solansky, S. T. (2011). Team identification: a determining factor of performance. *Journal of*

Managerial Psychology.

Stasser, G., & Abele, S. (2020). Collective choice, collaboration, and communication. *Annual Review*

of Psychology, 71, 589-612.

Számadó, S., Balliet, D., Giardini, F., Power, E. A., & Takács, K. (2021). The language of cooperation:

reputation and honest signalling. *Philosophical Transactions of the Royal Society B,*

376(1838), 20200286.

Schenk, L. A., & Colloca, L. (2020). The neural processes of acquiring placebo effects through

observation. *Neuroimage, 209*, 116510.

Stallen, M., Rossi, F., Heijne, A., Smidts, A., Dreu, C., & Sanfey, A.G.. (2018). Neurobiological

mechanisms of responding to injustice. *Journal of Neuroscience*, 2944.

Stasser, G., & Stewart, D. (1992). Discovery of hidden profiles by decision-making groups: Solving a

problem versus making a judgment. *Journal of personality and social psychology, 63*(3), 426.

Stolk, A., Verhagen, L., & Toni, I. (2016). Conceptual alignment: How brains achieve mutual

understanding. *Trends in Cognitive Sciences, 20*(3), 180–191.

Tajfel, H., & Turner, J. C. (1979). An integrative theory of inter-group conflict. In W. G. Austin & S.

Worchel (Eds.), *The social psychology of inter-group relations* (pp. 33–47). Monterey, CA:

Brooks/Cole.

Tindale, R. S., & Winget, J. R. (2019). Group Decision-Making. *In Oxford Research Encyclopedia of*

Psychology.

Turner, R., Howseman, A., Rees, G. E., Josephs, O., & Friston, K. (1998). Functional magnetic

resonance imaging of the human brain: data acquisition and analysis. *Experimental Brain*
Research, 123(1-2), 5-12.

Van Bavel, J.J., Cunningham, W.A. (2012). A social identity approach to person memory: group
membership, collective identification, and social role shape attention and memory. *Personality*
and Social Psychology Bulletin, 38(12),1566–78.

Wallis, J. D. (2007). Orbitofrontal cortex and its contribution to decision-making. *Annu. Rev. Neurosci.,*
30, 31-56.

Wang, C., Zhang, T., Shan, Z., Liu, J., Yuan, D., & Li, X. (2019). Dynamic interpersonal neural
synchronization underlying pain□induced cooperation in females. *Human Brain Mapping,*
40(11), 3222-3232.

Xie, E., Li, K., Gu, R., Zhang, D., & Li, X. (2023). Verbal information exchange enhances collective
performance through increasing group identification. *NeuroImage, 279*, 120339.

Xie, E., Liu, M., Li, K., Nastase, S. A., Gao, X., & Li, X. (2023). The single-and dual-brain
mechanisms underlying the adviser's confidence expression strategy switching during
influence management. *NeuroImage, 270*, 119957.

Xie, H., Karipidis, I. I., Howell, A., Schreier, M., Sheau, K. E., Manchanda, M. K., Ayub, R., Glover, G.
H., Jung, M., Reiss, A. L., & Saggar, M. (2020). Finding the neural correlates of collaboration
using a three-person fMRI hyperscanning paradigm. *Proceedings of the National Academy of*
Sciences, *117*(37), 23066-23072.

Ye, J. C., Tak, S., Jang, K. E., Jung, J., & Jang, J. (2009). NIRS-SPM: Statistical parametric mapping
for near-infrared spectroscopy. *NeuroImage, 44*(2), 428-447.

Yang, J., Zhang, H., Ni, J., De Dreu, C. K., & Ma, Y. (2020). Within-group synchronization in the
prefrontal cortex associates with intergroup conflict. *Nature neuroscience, 23*(6), 754-760.

Zhang, H. J., Dong, Y. C., Chiclana, F., & Yu, S. (2019). Consensus efficiency in group decision
making: A comprehensive comparative study and its optimal design. *European Journal Of*
Operational Research, 275(2), 580-598.