CTPS1 inhibition synergizes with replication stress signaling inhibition in *MYC*-amplified Group 3 medulloblastoma

Matthew R Hathaway[1], Katherine E Gadek[1], Hawa L Jagana[1], Isabella C Terrones[1], John M Hemenway[1], Aya Miyaki[1], Ashmitha Rajendran[1,2], Michael Meechan[1], Leonel Elena-Sanchez[1], Nicholas A Vitanza[1,3,4,8], Barbara S Slusher[5,6], Siobhan S Pattwell[1,3,7], Myron K Evans II[1,3,7,8]*

[1] Ben Towne Center for Childhood Cancer Research, Seattle Children's Research Institute, Seattle, WA, USA
[2] Biomedical Informatics and Medical Education Graduate Program, University of Washington, Seattle, WA, USA
[3] Department of Pediatrics, University of Washington School of Medicine, Seattle, WA, USA
[4] Clinical Research Division, Fred Hutchinson Cancer Center, Seattle, WA, USA
[5] Johns Hopkins Drug Discovery, Baltimore, MD, USA
[6] Department of Neurology, The Johns Hopkins University School of Medicine, Baltimore, MD, USA
[7] Human Biology Division, Fred Hutchinson Cancer Center, Seattle, WA, USA
[8] Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, WA, USA

* Address correspondence to:
Myron K Evans II, Ph.D.
Ben Towne Center for Childhood Cancer Research
1920 Terry Avenue
Seattle, WA 98101
Telephone: +1 206-884-4363
Email: myron.evansii@seattlechildrens.org

Keywords: CTPS1; medulloblastoma; MYC; glutamine; CHK1

Financial support: National Cancer Institute of the National Institutes of Health under Award Numbers K22CA262343 (MKE) and K22CA258953 (SSP), Kate Amato Foundation (MKE), Steven Higgins Brain Tumor Research Fund (MKE), Unravel Pediatric Cancer (MKE), the Pediatric Brain Tumor Research Fund of Seattle Children's (MKE),

Seattle Children's Guild Association, and UW/FHCC/SCH Cancer Center Support Grant (P30CA015704).

Conflict of Interest

Author B.S.S. is an inventor of patent applications filed by Johns Hopkins Technology Ventures covering novel glutamine antagonists, including JHU083 and JHU395, and their utility which has been licensed to Dracen Pharmaceuticals. B.S.S. is also a co-founder and prior consultant for Dracen Pharmaceuticals. This arrangement has been reviewed and approved by Johns Hopkins University following their respective conflicts of interest policies.

N.A.V. serves as the scientific advisory board chair for BrainChild Bio, Inc. and is an inventor on issued and pending patents related to CAR T cell therapies. This arrangement has been reviewed and approved by Seattle Children's Research Institute following their respective conflicts of interest policies.

Abstract

MYC-driven medulloblastomas (MBs) represent the most aggressive and deadly

subgroup of MB, the most common malignant pediatric brain tumor. Direct targeting of

MYC itself remains an unmet clinical need, therefore focusing on vulnerabilities driven

by MYC may be a viable option for novel therapeutic interventions. Using whole-

genome CRISPR screening, we identified the *de novo* pyrimidine synthesis enzyme

CTP synthase (*CTPS1*) as a strong dependency in *MYC*-driven MB. CTPS1 is the final

and rate-limiting step in the *de novo* pyrimidine synthesis pathway. Targeted inhibition of

CTPS1 leads to decreased tumor cell proliferation and markedly reduces MYC

expression in G3 MB models. Mechanistically, we demonstrate that single agent CTPS1

inhibition activates the replication stress signaling pathway mediated by ATM-CHK2 and

ATR-CHK1. Blockade of CHK1 kinase activity increases sensitivity to CTPS1 inhibition

and significantly impedes heterotopic MB tumor growth. CTPS1 enzymatic activity

requires the amino acid glutamine, therefore we inhibited CTPS1 using the glutamine

antagonists, JHU083 and JHU395. These compounds are prodrugs of 6-diazo-5-oxo-L-

norleucine (DON) which were developed to exhibit better tumor targeting and enhanced

blood-brain barrier penetrability. Combining JHU083 and CHK1 inhibition demonstrates

potent synergy against patient-derived MB xenografts *in vivo*. Our findings strongly

suggest that combining *de novo* pyrimidine synthesis and ATR-CHK1 inhibitors is a

promising treatment for *MYC*-driven MBs.

Key Points

- *CTPS1* is a unique vulnerability in MYC-driven medulloblastoma

- CTPS1 inhibition activates the ATR-CHK1 replication stress response pathway for cell survival

- Blockade of CTPS1 enzymatic activity synergizes with CHK1 inhibition *in vitro* and *in vivo*

Importance of the Study

MYC hyperactivation in tumors drives multiple anabolic processes which contribute to tumor proliferation and aggressiveness in patients. We show that targeting *de novo* pyrimidine synthesis (via CTPS1) limits tumor growth and targets MYC itself through a feedback mechanism. CTPS1 inhibition potently combines with CHK1 blockade and enhances disease control in both heterotopic and orthotopic models of medulloblastoma (MB). Our results support the clinical evaluation of combined CTPS1 and CHK1 inhibition in patients with *MYC*-driven MB.

Introduction

Despite being the second most common oncological diagnosis, central nervous system tumors are the leading cause of cancer-related mortality in children[1]. Medulloblastoma (MB), the most common malignant pediatric brain tumor, exhibits a complex genomic landscape with diverse molecular subgroups which significantly impact clinical outcomes[2]. Recent advances in genomic, epigenomic, and proteomic profiling have identified four subgroups of MB: WNT, sonic hedgehog (SHH), Group 3, and Group 4. Patients with Group 3 MB account for ~25-30% of all diagnosed cases and have the poorest overall survival of all groups[2].

Dysregulation of the proto-oncogene MYC is a hallmark of G3 MB, with patients exhibiting recurrent gene amplification and genomic fusions[3] or high MYC activity due to increased protein stability[4]. Despite our understanding of MYC as a critical oncogene[5], no direct targeted agents have been tested clinically in pediatric patients. This has motivated the study of synthetic lethality approaches, targeting vulnerabilities specifically driven by MYC hyperactivation[6].

MYC supports cancer cell proliferation through numerous mechanisms, including rewiring of cellular metabolism, by modulating anabolic processes to sustain the heightened demands of rapidly dividing cells. Oncogenic MYC drives the upregulation of key enzymes crucial to nucleotide biosynthesis, ensuring a robust supply of purine and pyrimidine precursors needed for rapid tumor cell proliferation[7]. Simultaneously, MYC promotes metabolic rewiring of glycolysis and the pentose phosphate pathway (PPP), channeling resources toward nucleotide synthesis[7]. This dynamic interplay is a characteristic feature of G3 MB[8]. Further, it underscores the significance of MYC in

tailoring cellular metabolism to meet the unique demands of oncogenesis, unveiling

potential vulnerabilities that could be harnessed for targeted therapeutic interventions.

Inhibition of nucleotide synthesis and metabolism effectively induces cell death and is

commonly exploited by various chemotherapeutics sometimes used in the treatment of

MB[9,10]. However, these treatments commonly yield unwanted side effects in patients,

suggesting that targeting tumor specific metabolism pathways could alleviate negative

sequelae. CTP synthase 1 (CTPS1), the rate limiting enzyme in the *de novo* synthesis of

the nucleotide CTP, is one such metabolic target[11,12]. CTPS1 has previously been shown

as a vulnerability in various adult solid and hematologic tumors[13-16] and a small molecule

inhibitor of CTPS1 is currently in clinical trials (NCT05463263). Further, a recent paper

linked MYC expression to CTPS1 inhibitor sensitivity, demonstrating synthetic lethality in

colorectal cancer[17]. Low on-target, off-tumor toxicity of this molecule (and similar

compounds) is most likely due to significant compensation by the homolog CTPS2 in most

tissues, excluding the lymphoid compartment post-infection[18,19].

In this study, using functional genomics and small molecule inhibition, we characterize

CTPS1 as a targetable vulnerability in G3 MB and identify MYC-amplified MB as

exquisitely dependent on *de novo* pyrimidine but not purine synthesis. Unlike in

hematological malignancies where CTPS1 inhibition alone significantly inhibits growth

and increases apoptosis[13,16], we observe limited cytotoxicity as a single agent in G3 MB.

We demonstrate that inhibition of CTPS1 in G3 MB activates the ATR-CHK1 replication

stress response, enhancing cell survival; dual antagonism of *de novo* pyrimidine

synthesis and CHK1 synergizes *in vitro* and *in vivo*. Finally, we demonstrate that a blood-

brain barrier (BBB) penetrant inhibitor of nucleotide biosynthesis combines with CHK1

134 inhibition increasing survival in an orthotopic model of MB. Our findings identify CTPS1

135 as a novel target and demonstrate that this combinatorial approach may be a valid option

136 to move forward clinically for patients with G3 MB.

Results

CTPS1 and the *de novo* pyrimidine synthesis pathway are dependencies in Group 3 medulloblastoma

Functional genomics screens allow for unbiased identification of potentially druggable vulnerabilities in tumors and have been utilized to guide pre-clinical work and clinical trials in both adult and pediatric oncology. To evaluate dependencies specific to Group 3 medulloblastoma (MB), we first explored the Pediatric Cancer Dependency Map (DepMap)[20]. Our analysis of 4 well-characterized G3 MB cell lines (D283, D341, D425, and D458) versus all other pediatric entities, revealed CTP synthase 1 (*CTPS1*) as the third-strongest dependency (defined by a negative CHRONOS score, **Figure 1A**). Comparing G3 MB to only pediatric (peds) central nervous system (CNS) tumors still reveals a strong dependency on *CTPS1* (**Figure S1A**), suggesting a specific function in *MYC*-amplified MB. CTPS1 is the rate limiting enzyme responsible for the glutamine-dependent conversion of uridine triphosphate (UTP) to cytidine triphosphate (CTP), an essential step of the *de novo* pyrimidine synthesis pathway[12] (**Figure 1B**). Further analysis revealed significant dependencies on other enzymes in the *de novo* pyrimidine synthesis pathway including carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase (*CAD*), uridine monophosphate synthase (UMPS), and dihydroorotate dehydrogenase (*DHODH*) (**Figure 1C,** G3 MB vs. all peds, **S1B** G3MB vs. peds CNS). UMP-CMP kinase (*CMPK1*) scores as a vulnerability however, no difference is seen when compared to other pediatric tumors. This is further validated by a recent publication which revealed DHODH as a druggable target by CRISPR screening in another G3 MB cell line (SUMB002)[21]. Mining of CRISPR screening data in two fetal

neural stem cell (NSC) lines[22] shows no dependency on *CTPS1* or other *de novo*

pyrimidine synthesis enzyme genes (**Figure S1C**). Interestingly, neither disruption of

CTPS2 (**Figure 1C, S1B**) or genes in the salvage pathway (**Figure S1D**) had any effect

on cell growth, suggesting a strong, non-redundant role for *CTPS1* in G3 MB.

Higher *CTPS1* expression is associated with disease advancement and poor

prognosis in MB

Using a batch-normalized bulk RNA-seq resource (GSE124814)[23] comprised of 1350 MB

samples and 291 normal cerebellum revealed increased *CTPS1* expression in tumors

compared to normal cerebella (**Figure 1D**). Subgroup analysis uncovered the highest

average expression of *CTPS1* in G3 tumors in two independent datasets (p<0.05

G3vs.G4 and G3vs.SHH, **Figure 1E-F**). Analysis of a third dataset (GSE85217)[24] showed

a strong correlation between *MYC* and *CTPS1* expression in G3 MB tumors, with the

highest expression in those tumors characterized as Group3γ (**Figure 1G-H)**. *CTPS1*

expression also correlates with metastatic status of patients at time of diagnosis (**Figure**

1I), further demonstrating association with aggressive characteristics. We validated these

RNA-seq findings in a complementary proteomic dataset[4]. In this data, G3 MB tumors

were subclassified as G3a and G3b, where G3a tumors are characterized by a MYC-

activated state either by gene amplification or post-translational alterations which

increase MYC protein stability. We demonstrate higher CTPS1 protein levels in G3a vs.

G3b tumors (**Figure S1E**), as well as higher expression of other components of the *de*

novo pyrimidine pathway (**Figure S1F**). Immunohistochemistry of G3 MB patient-derived

xenografts confirms expression of CTPS1 in tumor tissue with little expression in normal

mouse tissue (**Figure 1J**). Weaker or no correlations were seen with other genes in the

pyrimidine synthesis pathway (**Figure S1G**). A correlation between *CTPS1* and *CCNB1*,

a key regulator of cell cycle progression and a proposed risk stratification gene[25], was

also noted (**Figure S1H**). Consistent with this, higher *CTPS1* expression was associated

with worse overall survival in all MB patients (**Figure 1K**) and in G3 MB patients with high

MYC expression (**Figure 1L**). Of the other *de novo* pyrimidine synthesis pathway

enzymes, only *DHODH* and *UMPS* expression correlated with poor survival in G3 MB

patients (**Figure S1I**).

Targeting CTPS1 reduces MB cell proliferation *in vitro*

Based on the significant attenuation of growth using CRISPR-Cas9 targeting of *CTPS1*,

we targeted *CTPS1* using shRNA. After selection with puromycin, sphere formation was

assessed and cell proliferation measured by BrdU incorporation. In two different G3 MB

cell lines, cells exhibited decreased sphere forming capacity (**Figure 2A-B**), and

diminished proliferation concurrent with downregulation of *CTPS1* expression (**Figure

2C**). Despite significant growth inhibition *in vitro*, animals implanted with sh*CTPS1* cells

showed limited survival benefit compared to control shRNA implanted animals. HD-MB03

sh*CTPS1* tumors retained CTPS1 expression suggesting a negative selection for cells

with *CTPS1* knockdown *in vivo*. We next employed treatment with 3-Deazauridine (DAU)

which is a structural analog of uridine that competitively inhibits the synthesis of CTP by

both CTPS1 and CTPS2[26]. DAU treatment caused a dose-dependent decrease in cell

proliferation across a panel of G3 MB cell lines (**Figure 2D**). This inhibition could be

reversed by supplementation of either uridine or cytidine, but not adenosine or guanosine

(**Figure 2E**). Similar to previously published data[14,27], CTPS1 pharmacological inhibition

resulted in S-phase blockade with decreased cells in G_0-G_1, which was reversed by

addition of cytidine or uridine (**Figure 2F, S2A**). Furthermore, DAU treatment also

decreased MYC expression in tumor lines similar to previous reports with DHODH

inhibition in G3 MB[21] (**Figure 2G**).

G3 MB cells are dependent on *de novo* pyrimidine but not *de novo* purine synthesis

Previous work in other tumor types has revealed that MYC regulates both pyrimidine and

purine biosynthesis through direct regulation of multiple enzyme-encoding genes in both

pathways[7]. Analysis of previously published ChIP-seq data from two human G3 MB

tumors demonstrates significant MYC binding at the *CTPS1* promoter, with lower binding

at the *CTPS2* promoter correlating with gene expression[28] (**Figure S3A,B**). Analysis of

the same dataset from **Fig 1D**. revealed strong correlation between *MYC* and numerous

genes encoding enzymes in the *de novo* purine synthesis pathway (**Figure S3C**).

Exploration of DepMap data revealed that while some genes in the pathway scored as

dependencies, they were no different than pediatric CNS tumors (**Figure S3D**). The only

purine related enzyme that scores is thymidylate synthetase (TYMS), a catabolic one-

carbon metabolism enzyme that has an indirect effect on purine by driving thymidine

synthesis from dUMP salvage (**Figure S3E**). This correlates with activity of TYMS

inhibitors pemetrexed and 5-fluorouracil[29,30] and suggests the possibility of recently

developed novel inhibitors for future pre-clinical modeling[31,32].

To further clarify that inhibition of pyrimidine but not purine synthesis was antagonistic to

growth of G3 MB cells, we treated both Med-2112FHTC and Med-411FHTC with

lometrexol (LTX), an inhibitor of the purine biosynthetic enzyme glycinamide

ribonucleotide formyltransferase (GARFT). We observed no effect on cell fitness after

treatment with LTX, similar to previous reports in IDH-mutant glioma and pediatric diffuse

midline glioma[33] (**Figure S3F**).

CTPS1 inhibition alone is cytostatic but combines with CHK1 inhibition

Western blot analysis of cells treated with DAU revealed little change in PARP cleavage,

a classic marker of apoptosis, at doses up to 5µM suggesting a cytostatic mechanism

rather than cytotoxicity (**Figure 3A**). Diehl et.al. recently published that nucleotide

imbalance, caused by targeting pyrimidine and/or purine enzymes, activates replication

stress signaling sensed by the ATR and ATM kinases[27]. These kinases sense both single-

and double- stranded breaks and phosphorylate CHK1 and CHK2, respectively, which

mediate survival and recovery of cells. We observed increased phosphorylation of both

CHK1 and CHK2 after treatment with DAU (**Figure 3B**). We then combined targeting of

CTPS1 with the potent CHK1/2 inhibitor prexasertib, which previously showed efficacy in

G3 MB cells[34]. Combining sublethal doses of prexasertib (**Figure S4A**) led to a dose-

dependent decrease in the IC_{50} of DAU in both HD-MB03 and Med-411FHTC cells

(**Figure 3C and D**). **Table 1** lists synergy scores as calculated by multiple reference

models[35] which revealed potent synergy between CTPS1 and CHK1/2 inhibition in G3

MB. This combination also led to apoptosis (**Figure 3C, inset** – cleaved PARP) and

significant DNA damage (measured by pH2AX) (**Figure 3E-F**). Prexasertib treatment also

synergized with BRQ, suggesting that CHK1 inhibition may broadly combine with

antagonism of *de novo* pyrimidine synthesis in G3 MB (**Figure S4B**). Utilizing a

heterotopic xenograft model of G3 MB in athymic nude mice, we assessed the efficacy of

combining DAU and prexasertib on tumor growth (**Figure 3G**). While single agent

targeting of either CTPS1 or CHK1 led to delays in tumor growth, the combination of DAU

and prexasertib synergized and completely abrogated growth in this model (**Figure 3H**).

Immunohistochemical analysis revealed decreased Ki67 corroborating our doubling time

findings (**Figure 3I**). Phosphorylation of H2AX was also elevated in combination treated

animals. Western blots revealed significant cleavage of PARP in our combination

treatment and sustained phosphorylation of CHK1/CHK2 (S345/S516) in combination

treatments (**Figure 3J**). We also noted decreased phosphorylation of the CHK1 target

CDC2 (Y15) in some combination treatment mice as well.

Targeted glutamine antagonism combines with CHK1 inhibition in orthotopic

models

While our *in vitro* and *in vivo* data with DAU are striking, there is only one reference to

blood-brain barrier penetration of this compound in humans[36]. This was evaluated in a

handful of patients with intracerebral tumors which commonly have abnormally leaky

vasculature[37]. MBs, apart from the WNT subgroup, are characterized to have a fully intact

BBB which modulates response to BBB-impermeable drugs[38]. The combination of these

make DAU an unlikely candidate for G3 MB therapeutic development. CTPS enzymatic

activity requires the amino acid glutamine as well as ATP to produce CTP[39]. Glutamine is

also required at the first step of the *de novo* synthesis pathway catalyzed by CAD (**Figure**

1B). Interestingly, the glutamine antagonist 6-diazo-5-oxo-l-norleucine (DON) was

previously shown to be an irreversible inhibitor of multiple nucleotide synthesis

enzymes[40], including CTPS1 and CAD. Single agent DON administration leads to

significant phosphorylation of CHK1/2 (**Figure 4A**) and cell death in G3 MB tumor cells[41].

We utilized two prodrugs of DON – 1) isopropyl 6-diazo-5-oxo-2-(((phenyl (pivaloyloxy)

methoxy) - carbonyl) amino) hexanoate (JHU395) and 2) Ethyl 2-(2-Amino-4-

methylpentanamido)-DON (JHU083) – which exhibit enhanced lipophilicity, brain

penetration, and were previously shown to target G3 MB tumor cells[41,42] (**Figure S5A**).

We first tested JHU083 and JHU395 as single agents in our patient-derived cell lines

which revealed efficacy with inter-cell line variability (**Figure 4B**). Co-administration of

prexasertib synergized with both JHU083 (**Figure 4C and D**) and JHU395 (**Figure 4E**

and F), decreasing the IC$_{50}$ in multiple G3 MB tumor lines. Synergy analyses (**Table 1**)

revealed stronger synergism of prexasertib with JHU083 than with JHU395, so for the

remainder of our study we used the combination of JHU083+prexasertib. Combining

JHU083 and prexasertib led to significant accumulation of DNA damage (**Figure 4G-H**)

similar to DAU+prexasertib. Previous work demonstrated significant BBB penetration and

tumor accumulation of both JHU083[42] and prexasertib[43] in G3 MB tumor bearing mice,

therefore we tested single agents and combinations after orthotopic implantation of

luciferase-expressing G3 MB cells. After confirmation of tumor growth by IVIS imaging,

mice were randomized to one of four groups and treated with JHU083 alone or in

combination with prexasertib (**Figure 4I**). While single agent administration had limited

effect on tumor growth or animal survival, the combination of JHU083 and prexasertib

significantly extended the median survival in animals implanted with either HD-MB03

(**Figure 4J, S5B**) or Med-411FHTC (**Figure 4K, S5C**). Further, this combination as well

as our DAU+prexasertib combination had limited systemic toxicities as measured by

weight loss in animals (**Figure S5D,E**). Western immunoblot analysis of tumors revealed

increased phospho-CHK1/2 and phospho-CDC2 in JHU083 single agent treated tumors.

Combination treatment led to a blockade of CHK1/2 and CDC2 phosphorylation as well

as a decrease in proliferation as measured by PCNA (**Figure 4L**).

Discussion

Herein, we report a dependency on *CTPS1* in *MYC*-amplified MBs when compared to other pediatric malignancies. Interrogation of patient data revealed high expression of *CTPS1* in tumors compared to normal brain (cerebellum/upper rhombic lip) with *MYC*-amplified G3γ tumors displaying the highest expression. This was confirmed using recently published proteomics data which correlates CTPS1 with G3a tumors, characterized by increased MYC expression and activity[4]. Consistent with these findings, *CTPS1* expression correlates with poor survival in *MYC* high patients. CTPS1, along with its homolog CTPS2, catalyze the rate limiting step of the *de novo* pyrimidine biosynthesis pathway. Interestingly, *MYC*-amplified MB tumors do not show a dependency on CTPS2, which also shows no correlation with survival. This is consistent with previous reports that demonstrate differential activity of the two enzymes, despite ~70% protein sequence homology[44]. Further analysis revealed vulnerabilities to deletion of other members of the *de novo* pyrimidine biosynthesis pathway, confirming previous reports of altered pyrimidine metabolism in MYC-amplified MB[21], and more broadly across a number of other brain tumor types[33]. Furthermore, recent work in a chemotherapy-resistant model of MB revealed upregulation of pyrimidine metabolic processes in resistant tumors[45], suggesting utility of CTPS1 targeting in both upfront and recurrence settings.

Using shRNAs and a small molecule inhibitor of CTPS1 (3-deazuridine, DAU) we demonstrate that single agent targeting of CTPS1 leads to cell cycle arrest and decreased proliferation which can be rescued by addition of exogenous pyrimidine nucleosides *in vitro*. This targeting of CTPS1 also leads to a decrease in MYC expression which may further impact tumor growth. Interestingly, germline mutations in *CTPS1* lead to no overt

phenotypes outside of the hematopoietic system, suggesting that inhibition may deliver

potent on-target effects in tumors while limiting deleterious side effects in patients[18,19].

This is highlighted by single agent efficacy of DAU in our flank model with no significant

weight loss observed. Mechanistically, amplified or hyperactive MYC in G3 MB induces

expression of CTPS1 required to synthesize nucleosides to maintain replicative capacity

of tumors. Inhibition of CTPS1 limits the pool of CTP available for replication[9], leading to

activation of the replication stress response pathway mediated by ATR-CHK1/ATM-CHK2

and subsequent cell cycle and proliferation arrest. Combined targeting of CTPS1 and the

replication stress pathway forces cells to continue through the cell cycle, accumulating

significant DNA damage and undergoing apoptosis (**Figure 5**).

This phenomenon has been observed in other tumor and normal cell types wherein

combining nucleotide depletion with ATR/ATM inhibitors leads to cell death[14,17,27]. We

demonstrate significant synergy between the CHK1 inhibitor, prexasertib, and DAU both

in vitro and *in vivo*. Further, we also demonstrate that the DHODH inhibitor, brequinar,

also synergizes with prexasertib suggesting the utility of dual *de novo* pyrimidine

synthesis and replication stress signaling inhibition in treatment of *MYC*-amplified MB.

Identification of this combination may further enhance the clinical utility of a DHODH

inhibitor from Bayer which shows potent BBB penetration, but whose clinical development

was halted due to single agent inactivity (NCT03404726).

Excitingly, CTPS1 inhibitors have recently entered clinical development (NCT05463263)

for hematological malignancies; however, blood-brain barrier (BBB) penetrance has not

been assessed. This compound (STP938) mimics CTP and acts an allosteric inhibitor of

enzymatic activity; impeding CTPS1 binding to UTP. CTPS1 also has a glutamine

amidotransferase domain which hydrolyzes glutamine to ammonia which is required for

catalysis of 4-phosphoryl UTP to CTP. Previous work identified that the glutamine

antagonist 6-Diazo-5-oxo-L-norleucine (DON) binds to this domain, as well as other

glutamine utilizing enzymes, to inhibit covalent binding of glutamine[40]. We observed a

similar activation of CHK1/2 when cells were treated with DON as with DAU. While DON

does accumulate in mouse models of brain tumors, previous clinical trials were

unsuccessful due to dose-limiting toxicities[46]. We utilized two DON prodrugs, JHU083 and

JHU395, with enhanced BBB penetrability and tumor-targeting, leading to reduced toxic

side effects[47,48]. These inhibitors combine strongly with CHK1 inhibition *in vitro*, with the

combination of JHU083 and prexasertib showing significant synergy. We tested this

combination *in vivo* and noted increased survival, decreased proliferation, and increased

apoptosis, and limited weight loss despite prolonged combination treatment. While our

studies are done in immune-deficient models, recent studies have identified immune

modulatory activity of JHU083[49,50] which may bolster effects of this drug in immune-

competent mouse models and patients. Together, the data presented herein identify a

targeted, combinatorial approach for the treatment of *MYC*-amplified MB patients who

currently have the least favorable outcomes.

Methods

Cell culture

Patient derived medulloblastoma cell lines (Med-411FHTC and Med-2112FHTC, a gift from J. Olson) were grown in NeuroCult NS-A Basal medium (Stemcell Technologies, 05750) with Neurocult NS-A Proliferation Supplement (Stemcell Technologies, 05753), and 1% Penicillin/Streptomycin (Sigma-Aldrich, P4333). HD-MB03 (obtained from T. Milde) and MB004 (a gift from P. Bandopadhayay) cell lines were grown in Gibco Neurobasal medium (Gibco, 21-103-049) with 2% B27 Supplement (Gibco, 12587010), 1% N2 Supplement (Gibco, 21103049), 1% Glutamine (Gibco, 25030-81), 1% Penicillin/Streptomycin (Sigma-Aldrich, P4333), and 1µg/mL Heparin (Stemcell Technologies, 07980). All cell lines were grown as spheroids and media supplemented with 0.2µg/mL each Human Epidermal Growth Factor Protein (Sino Biological, 10605-HNAE-1) and Human Recombinant bFGF/FGF2 Protein (Sino Biological, 10014-HNAE-1). HEK293T were obtained from American Type Culture collection and maintained in DMEM (Gibco) with L-glutamine (Fisher Sci.), 10% fetal bovine serum (Fisher Sci.), 1% L-Glutamine, and 1% Penicillin/Streptomycin (Sigma-Aldrich, P4333). All cells were grown at 37°C and 5% CO_2.

Drugs and small molecules

3-Deazuridine (DAU) was obtained from Santa Cruz Biotechnology (394445) or Cayman Chemical Company (23125). JHU395 and JHU083 were a gift from Barbara Slusher (Johns Hopkins). Prexasertib mesylate (206787) and Prexasertib HCl (206989) were procured from MedKoo Biosciences. Brequinar was procured from SelleckChem (S6626) and Lometrexol from Cayman Chemical Company (18049). 6-diazo-5-oxo-L-nor-Leucine

(DON) was procured from Cayman Chemical Company (17580). Guanosine (S2439),

Adenosine (S1647), Cytidine (S2053), and Uridine (S2029) were procured from

SelleckChem. For *in vitro* use, drugs were reconstituted in DMSO (Millipore Sigma,

D2438).

Proliferation measurements

For Cell Titer Glo, cells were treated with serial concentrations of Prexasertib (0-20nM),

DAU (0-30nM), JHU083 (0-20µM), JHU395 (0-20µM), Lometrexol (0-180nM), and

Brequinar (0-180nM) or indicated combinations, then incubated for 72 hours. Proliferation

was assessed with CellTiter-Glo 2.0 (Promega, G9242) according to manufacturer

instructions and quantified using a Spectramax iD5. For BrdU assessment, cells

(untreated or treated as indicated) were incubated with 10µM BrdU (ThermoFisher,

B5002) for 1hr. After incubation, cells were rinsed with 1X PBS (Cytiva, BSS-PBS-1X6),

fixed in ice-cold 70% ethanol, and stored at -20°C. Fixed cells were washed with DPBS

and incubated in 2N HCl/0.5% Triton X-100 for 20 min. Cells were stained with anti-BrdU

FITC antibody (BioLegend, 364104) for 1 hour and analyzed on a BD Accuri C6 and

FlowJo (BD Biosciences).

Cell cycle measurement

For cell cycle measurement, cells were treated as indicated and fixed in ice-cold 70%

ethanol after PBS wash. FxCycle PI/RNase Staining Solution (ThermoFisher, F10797)

was added to cells as per manufacturer instruction. Analysis was performed on a BD

Accuri C6 and cell cycle analysis performed in FlowJo (BD Biosciences).

Lentiviral shRNA production

Lentiviral shRNA plasmids (U6 promoter) were designed and produced by Vectorbuilder.

shRNA sequences were:

shCTPS1: GCTCTCACATTACCTCCAGAA (TRCN0000045349)

shCTPS2: CGCCTCACCAAGGACAATAAT (TRCN0000369478)

HEK293T cells were plated on a 10cm dish (Corning, 877222) until 60-70% confluent.

Cells were transfected with 10µg transfer plasmid (VectorBuilder), 9µg psPAX2

(Addgene, 12260) and 3µg pMD2.G (Addgene, 12259) using Xfect Transfection Reagent

(Takara, 631318) according to manufacturer instructions. After transfection, virus was

collected in Neurobasal media, filtered through a 0.45µm syringe filter (Globe Scientific,

SF-PES-4530-S) and stored at -80°C. Viral particles were added to tumorspheres with

8µg/mL polybrene (Millipore Sigma, TR-1003-G) overnight. Forty-eight hours post-

infection, puromycin (Millipore Sigma, P4512-1MLX10) was added at 1µg/mL for 14 days

to select for infected cells.

Western immunoblotting

After indicated treatments, cells were lysed in RIPA buffer (150 mM NaCl; 50 mM Tris pH

8.0; 1.0% IGEPAL CA-630; 0.5% sodium deoxycholate; 0.1% SDS) (Thermo Scientific,

PI89900) with Halt protease and phosphatase inhibitor (Thermo Scientific, PI78445). For

tumor tissue, tissue was homogenized using an electric homogenizer fitted with a

disposable pestle in RIPA buffer. Protein was quantified using Pierce 660nm Protein

Assay Reagent (Thermo Scientific, 22660) and immunoblotted with standard procedures.

Antibodies and concentrations are listed Table S1.

Immunofluorescence staining

FFPE sections from flank tumors were deparaffinized and rehydrated with Histoclear

(HS2001, Fisher) and ethanol according to manufacturer instructions. Antigen retrieval

was performed by pressurized retrieval in pH 6.0 citrate buffer (C9999, Millipore Sigma)

for 5 min on high pressure in a pressure cooker (Cuisinart CPC-600 6 Quart 1000 Watt

Electric Pressure Cooker). Samples were blocked in 0.1% PBS+TX-100 5% donkey

serum (S30-M, Millipore Sigma) for 1 hr at RT prior to staining with antibodies overnight

at 4C. Secondary Alexa fluor antibodies (donkey anti-488, 555 or 647; Invitrogen) were

diluted at 1:500 in blocking buffer for 1 hour at RT. Samples were mounted in Prolong

Gold with NucBlue (P36981, Fisher Scientific). All samples were imaged on a Leica DMi8

microscope with Thunder deconvolution software (LAS X, Leica Microsystems) and

quantified in ImageJ.

Synergy analysis

Visualization and analysis of drug combinations were performed using the SynergyFinder

web application (version 3.0, https://synergyfinder.fimm.fi) using viability as readout.

Outlier detection was on with LL4 curve fitting. Bliss, ZIP, HSA, and Loewe scores and p-

values were calculated as described[35].

Mouse xenograft experiments

All animal procedures were approved by Institutional Animal Care and Use Committee

(IACUC) of Seattle Children's Research Institute (Protocol #00000675). Nude female

mice (Jackson Laboratories, NU:J- 002019 or 007850) were procured at 5 weeks of age

and allowed to acclimate for at least 1 week prior to cell implantation. Luciferase-

expressing cells were made by transduction of single cells with pLenti CMV mCherry-

T2A-Luc blast (a gift from N. Vitanza) followed by hand selection of positive tumorsphere

colonies.

Subcutaneous implantation

Patient-derived MB cells were dissociated with Accutase (AT-104, Innovative Cell

Technologies), resuspended in Gibco Neurobasal Medium (21103049, Fisher Scientific),

and injected into the flank in a 100 µL aliquot containing 1.25×10^6 cells. Mice were

randomly segregated into groups when tumors reached 100 mm^3 volume: vehicle,

Prexasertib (10mg/kg subcutaneous twice daily), 3-Deazuridine (75mg/kg

intraperitoneal), or combination until health-defined end points. Tumor size was measured

with calipers. For growth curve analysis and survival studies, mice in study cohorts were

sacrificed when tumors reached a max of 3000 mm^3 in total volume. The tumors were

excised from sacrificed mice and subjected to routine histopathological processing.

Orthotopic implantation

MB tumorspheres were treated with Accutase and approximately 100,000 mCherry-

positive cells injected per mouse in 2µL of sterile PBS (Cytiva, BSS-PBS-1X6). For

surgery, mice were given buprenorphine ER for analgesia and anesthetized with a

ketamine/xylazine mixture. A 0.9mm burr hole was made in the right cerebellum (-1.0mm

AP 1.0mm ML) from lambda and a stereotactic needle dispensed the cells in two 1µL

injections (2.5mm and 2.0mm DV) to deliver the cells. To monitor tumor growth, mice were

injected with 150 mg/kg of D-luciferin (LUCK-4G, Gold Biotechnologies), and imaged

using the IVIS Spectrum Imaging System 15 minutes post-injection.

***In vivo* drug administration**

JHU083 was prepared in sterile PBS without Ca and Mg (Cytiva, BSSPBS1X6) and

administered at a concentration of 20mg/kg via oral gavage twice weekly. Prexasertib

mesylate (MEDKoo Biosciences, 206787) or prexasertib HCl (MEDKoo Biosciences,

206989) were prepared in 10% SBE-β-CD (MedChem Express, Cat# HY-17031/CS-

0731) and dosed at a concentration of 10mg/kg subcutaneously twice daily. 3-

Deazuridine (Santa Cruz Biotechnology, 394445) was diluted in corn oil (Millipore Sigma,

C8267-500ML) and dosed at a concentration of 75mg/kg IP once daily.

Dependency Map Analysis

Dependency data (CERES score) from the Broad Institute's Pediatric Cancer

Dependency Map[20] was accessed from the 23Q4 data deposit (depmap.org).

Patient gene expression and proteomic analyses

The processed expression of indicated genes were downloaded from the *R2: Genomics*

Analysis and Visualization Platform (**http://r2.amc.nl**) for indicated data sets[3,23,24]. When

calculating expression differences between different groups they were considered

significantly different when $P < 0.05$, as calculated using the one-way analysis of variance

(ANOVA) test included in the R2 online software. Linear regression was used to define

R-values and p-values for correlation plots. For significance of differences between

primary and metastatic tumors, the p-value was calculated with an unpaired

Student *t* test. For survival analyses in Kaplan–Meier plots, we used a 5-year/60-month

time point as a cut-off for survival due to the potential risk of including secondary tumors.

The *P* value for survival curves is calculated using a log-rank test of reported patient

survival data as described in R2 (**https://r2.amc.nl**). Only *a priori* fixed cut-offs (median

or top 1/3 v bottom 2/3) were used to calculate P values in the Kaplan–Meyer survival analysis.

Proteomics data was downloaded from the Medulloblastoma Data Explorer (https://medullo.shinyapps.io/archer2018/). One-way ANOVA was used to determine significant differences.

Acknowledgements

We thank the University of Washington/Fred Hutchinson Cancer Center/Seattle Children's Hospital Cancer Consortium for shared resources including Flow Cytometry and Cell Sorting and Research Pathology supported by NCI P30CA015704. We thank Virginia Hoglund and Galen Stewart for histopathology support, Jim Olson for providing PDX tissue and cell lines, Beth Lawlor for reagent support, Karuna Patil and Rajesh Utamanthil for animal husbandry and veterinary support. Research reported in this manuscript was funded by the National Cancer Institute of the National Institutes of Health under Award Numbers K22CA262343 (MKE) and K22CA258953 (SSP), Kate Amato Foundation (MKE), Steven Higgins Brain Tumor Research Fund, Unravel Pediatric Cancer, the Pediatric Brain Tumor Research Fund of Seattle Children's, the Run of Hope Seattle, Seattle Children's Guild Association, and UW/FHCC/SCH Cancer Center Support Grant (P30CA015704).

Author contributions

Matthew R Hathaway: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing – original draft, Writing – review & editing. **Katherine E Gadek:** Data curation, Formal analysis, Investigation,

Methodology, Validation, Visualization, Software, Writing – original draft, Writing –

review & editing. **Hawa L Jagana:** Data curation, Writing – review & editing. **Isabella C**

Terrones: Data curation, Formal analysis, Writing – review & editing. **John M**

Hemenway: Data curation, Formal analysis, Writing – review & editing. **Aya Miyaki:**

Data curation, Writing – review & editing. **Ashmitha Rajendran:** Investigation, Writing –

review & editing. **Michael Meechan:** Data curation, Investigation, Writing – review &

editing. **Leonel Elena-Sanchez:** Data curation, Investigation, Writing – review & editing.

Nicholas A. Vitanza: Methodology, Resources, Writing – review & editing. **Barbara S**

Slusher: Methodology, Resources, Writing – review & editing. **Siobhan S Pattwell:**

Methodology, Resources, Writing – review & editing. **Myron K Evans II:**

Conceptualization, Funding acquisition, Methodology, Project administration, Software,

Supervision, Validation, Writing – original draft, Writing – review & editing.

 Figure Legends

Figure 1: *CTPS1* is overexpressed in medulloblastoma and is a dependency in G3

MB. a, Volcano plot comparing CERES scores of G3 MB cell lines (D283, D425, D458,

D341) versus other pediatric tumor lines (n=111) in the Dependency Map using two class

comparison. **b,** Diagram of *de novo* pyrimidine synthesis pathway, **c,** CERES gene

dependency scores for pyrimidine synthesis genes in G3 MB cells (n=4) compared to

other pediatric cell lines (n=111) in the DepMap, 23Q4 release. **d,** Quantile normalized

expression of CTPS1 in normal (n=291) or primary medulloblastoma (n=1350) samples.

e-f, Log$_2$ normalized expression of CTPS1 in GSEs (e-GSE124814, f-

EGAS00001001953). Samples are separated by MB subgroup. **g,** Linear regression of

log$_2$ normalized CTPS1 and MYC expression from GSE85217 separated by subtype. **h,**

Log$_2$ normalized expression of CTPS1 in G3 MB subtypes from same samples in **g**. **i,**

Log$_2$ normalized CTPS1 expression in primary tumor samples from patients with or

without metastases at diagnosis from GSE85217. **j,** Immunohistochemistry of G3 patient-

derived xenograft tissue after orthotopic implantation in nude mice (red-CTPS1, green-

human nuclear antigen (hNuc), blue-DNA). Scale bar=100 µM. **k,** Kaplan-Meier survival

of patients diagnosed with MB separated by median expression of *CTPS1*, GSE85217. **l,**

Kaplan-Meier survival of *Myc*-high G3 MB patients separated by top 1/3 versus bottom

2/3 expression of *CTPS1*. For **k** and **l**, number of patients are shown on inset. *, P < 0.05;

**, P < 0.01; ***, P < 0.001, Data are means ± SEM.

Figure 2: CTPS inhibition suppresses cell proliferation in G3 MB tumor cells. a,

Representative images of HD-MB03 tumorspheres transduced with indicated shRNAs

72h after plating. **b,** Quantification of area of tumorspheres formed in shScramble,

shCTPS1.1, and shCTPS1.2 transduced HD-MB03 cells. **c,** Quantification of FACS for

BrdU-positive HD-MB03 and MED-411FHTC cells transduced with indicated shRNAs

(n=3). *Inset*, Western immunoblot analysis of CTPS1 expression in transduced cells in **a-**

c Scr= scramble, 1.1 =shCTPS1.1, 1.2= shCTPS1.2. **d,** Cell proliferation analysis of HD-

MB03, MB004, Med-411FHTC, and Med-2112FHTC tumorspheres treated with indicated

concentrations of DAU for 72h. **e,** Cell proliferation analysis of HD-MB03 and Med-

411FHTC tumorspheres treated with vehicle or DAU (5µM) alone or in the presence of

50µM cytidine, uridine, adenosine, or guanine (n=3). **f,** Cell-cycle analysis of HD-MB03

cells treated with the indicated concentrations of DAU alone or in the presence of cytidine

(C) or uridine (U) for 48h (n=3). **g,** Western immunoblot analysis of MYC expression in

HD-MB03 cells treated with increasing doses of DAU. *, $P < 0.05$; **, $P < 0.01$; ***, $P <$

0.001, Data are means ± SEM.

Figure 3: CTPS1 inhibition synergizes with ATR-CHK1 blockade *in vitro* and *in vivo*.

Western immunoblot for **a,** PARP and **b,** phospho-CHK1/2 in Med-411FHTC or HD-MB03

cells treated with vehicle, 2.5µM, or 5µM DAU for 48h. **c,** Cell proliferation analysis of

Med-411FHTC (left) and HD-MB03 (right) tumorspheres with increasing concentrations

of DAU alone or in the presence of indicated prexasertib (CHK1 inhibitor) concentrations

(n=3). *Inset*, Western immunoblot analysis of cleaved PARP in cells treated with

prexasertib alone (1.25nM), DAU alone (2.5µM), or the combination. **d,** 2D contour plots

of synergy (Bliss calculation) between DAU and prexasertib in Med-411FHTC (left) and

HD-MB03 (right) tumorspheres calculated by Synergy Finder. **e,** Immunofluorescence for

phospho-H2AX in D283 G3 MB cells treated as indicated. Scale bar 25 µM. **f,**

Quantification of images in **e** (n=5 images per condition, experiment repeated twice). **g,**

Pre-clinical mouse treatment protocol for data shown in **h**. **h,** left – Tumor volume of HD-

MB03 flank xenografts treated as indicated n=5 (vehicle) or 6 (all others). Tumor volume

statistics (p-value) – veh-prex: 3.9E-6, veh-combo: 3.1E-11, prex-combo: 8.7E-3. TGI:

tumor growth inhibition DT: doubling time; right – Quantification of tumor weights in grams

at Day 20 post-treatment (veh, n=2, DAU, n=3, prex, n=3, combo, n=5). **i,** Representative

images of immunohistochemistry for Ki-67 and phospho-H2AX in tumors at endpoint. **j,**

Western immunoblot for total PARP (* denotes cleaved PARP) and phospho-CHK1/2 in

tumors from **h,**. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$, Data are means ± SEM.

Figure 4: CTPS1 inhibition via glutamine analogs synergizes with CHK1 inhibition

in orthotopic G3 MB models. a, Western immunoblot for phospho-CHK1/2 in Med-

411FHTC (left) and HD-MB03 (right) tumorspheres treated with 10µM DON. **b,** Cell

proliferation analysis of HD-MB03, MB004, Med-411FHTC, and Med-2112FHTC

tumorspheres treated with indicated concentrations of JHU083 (left) and JHU395 (right)

for 72h (n=3). Cell proliferation analysis and 2D contour plots of synergy (Bliss calculation)

for **c, d** JHU083 and **e, f** JHU395 alone or in the presence of indicated prexasertib

concentrations (n=3). **g,** Immunofluorescence for phospho-H2AX in D283 G3 MB cells

treated as indicated. Scale bar 25 µM. **h,** Quantification of images in **g** (n=5 images per

condition, experiment repeated twice). **i,** Pre-clinical mouse treatment protocol for data

shown in **j-k**. Kaplan-Meier survival of mice orthotopically implanted with **j,** HD-MB03 or

k, Med-411FHTC cells transduced with luciferase expressing construct and treated as

indicated. **l,** Western immunoblot for phospho-CHK1/2, phospho-CDC2, and PCNA in

tumors from **j-k**. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$, Data are means ± SEM.

Figure S1: Dependency, gene expression and survival data for additional de *novo* pyrimidine synthesis and salvage pathway genes in G3 MB. a, Volcano plot comparing CERES scores of G3 MB cell lines (D283, D425, D458, D341) versus other pediatric CNS tumor lines (n=17) in the Dependency Map using two class comparison. **b,** CERES gene dependency scores for pyrimidine synthesis genes in G3 MB cells (n=4) compared to other pediatric CNS tumor lines (n=17) in the DepMap, 23Q4 release. **c**, CERES scores for normal NSCs (CB660 and U5) for *de novo* pyrimidine synthesis genes as well as *TOP2A* and *RPA2* (controls). **d**, CERES gene dependency scores for pyrimidine salvage genes in G3 MB cells (n=4) compared to all pediatric (left, n=111) and peds CNS (right, n=17). **e**, Relative CTPS1 protein expression in G3a (n=8) and G3b (n=6) tumors from(ref). **f**, Heatmap of relative protein expression for other *de novo* pyrimidine synthesis proteins. **g**, Linear regression of $\log_2$ normalized *MYC* and *de novo* pyrimidine synthesis gene expression from GSE85217 separated by subtype. **h**, Linear regression of $\log_2$ normalized *CTPS1* and *CCNB1* expression separated by G3 subtype. **i**, Kaplan-Meier survival of *MYC*-high G3 MB patients separated by top 1/3 versus bottom 2/3 expression of indicated genes. Number of patients are shown on inset.

Figure S2: DAU treatment alters cell cycle in G3 MB. a, Cell-cycle analysis of Med-411FHTC cells treated with the indicated concentrations of DAU alone or in the presence of cytidine (C) or uridine (U) for 48h (n=3). *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$, Data are means ± SEM.

Figure S3: MYC drives expression of pyrimidine and purine synthesis genes, but G3 MB cells are insensitive to purine synthesis inhibition. a, Representative ChIP-sequencing tracks for MYC in human G3 MB cells at *CTPS1* and *CTPS2* promoters. **b,**

Expression (transcripts per million, TPM) of *CTPS1* and *CTPS2* in tumor samples from **a**. **c,** Linear regression of $\log_2$ normalized *MYC* and *de novo* purine synthesis gene expression in G3 tumors. **d**, CERES gene dependency scores for *de novo* purine synthesis genes in G3 MB cells (n=4) compared to peds CNS (n=17) cell lines. **e**, left - CERES scores for *TYMS* in G3 MB (n=4), all peds (n=111), and peds CNS (n=17); right - Linear regression of $\log_2$ normalized *MYC* and *TYMS* expression in G3 tumors. **f**, Cell proliferation analysis of Med-411FHTC and Med-2112FHTC tumorspheres treated with indicated concentrations of lometrexol (LTX) for 72h (n=3). *, P < 0.05; **, P < 0.01; ***, P < 0.001, Data are means ± SEM.

Figure S4: Prexasertib alone has limited effect but synergizes with DHODH inhibition. Cell proliferation analysis of HD-MB03 and Med-411FHTC tumorspheres treated with increasing concentrations of **a,** prexasertib alone or **b,** brequinar in the presence of indicated doses of prexasertib. Data are means ± SEM.

Figure S5: JHU083+prexasertib combination limits tumor growth in vivo. a, chemical structures of DON, JHU083, and JHU395. IVIS imaging of mice orthotopically implanted with **b,** HD-MB03 or **c,** Med-411FHTC -luciferase expressing tumor lines treated as indicated. Images are from mid-point of treatment. Monitored weights of animals from **d,** heterotopic model treated with DAU and/or prexasertib and **e,** orthotopic model treated with JHU083 and/or prexasertib.

1. Siegel DA, Richardson LC, Henley SJ, et al. Pediatric cancer mortality and survival in the United States, 2001-2016. *Cancer.* 2020; 126(19):4379-4389.
2. Juraschka K, Taylor MD. Medulloblastoma in the age of molecular subgroups: a review. *J Neurosurg Pediatr.* 2019; 24(4):353-363.
3. Northcott PA, Shih DJ, Peacock J, et al. Subgroup-specific structural variation across 1,000 medulloblastoma genomes. *Nature.* 2012; 488(7409):49-56.

4. Archer TC, Ehrenberger T, Mundt F, et al. Proteomics, Post-translational Modifications, and Integrative Analyses Reveal Molecular Heterogeneity within Medulloblastoma Subgroups. *Cancer Cell.* 2018; 34(3):396-410.e398.

5. Conacci-Sorrell M, McFerrin L, Eisenman RN. An overview of MYC and its interactome. *Cold Spring Harb Perspect Med.* 2014; 4(1):a014357.

6. Wang C, Zhang J, Yin J, et al. Alternative approaches to target Myc for cancer treatment. *Signal Transduction and Targeted Therapy.* 2021; 6(1):117.

7. Dong Y, Tu R, Liu H, Qing G. Regulation of cancer cell metabolism: oncogenic MYC in the driver's seat. *Signal Transduction and Targeted Therapy.* 2020; 5(1):124.

8. Pham K, Hanaford AR, Poore BA, et al. Comprehensive Metabolic Profiling of MYC-Amplified Medulloblastoma Tumors Reveals Key Dependencies on Amino Acid, Tricarboxylic Acid and Hexosamine Pathways. *Cancers (Basel).* 2022; 14(5).

9. Parker WB. Enzymology of purine and pyrimidine antimetabolites used in the treatment of cancer. *Chem Rev.* 2009; 109(7):2880-2893.

10. Shelton J, Lu X, Hollenbaugh JA, Cho JH, Amblard F, Schinazi RF. Metabolism, Biochemical Actions, and Chemical Synthesis of Anticancer Nucleosides, Nucleotides, and Base Analogs. *Chem Rev.* 2016; 116(23):14379-14455.

11. Huang M, Graves LM. De novo synthesis of pyrimidine nucleotides; emerging interfaces with signal transduction pathways. *Cell Mol Life Sci.* 2003; 60(2):321-336.

12. Levitzki A, Koshland DE, Jr. Cytidine triphosphate synthetase. Covalent intermediates and mechanisms of action. *Biochemistry.* 1971; 10(18):3365-3371.

13. Asnagli H, Minet N, Pfeiffer C, et al. CTP Synthase 1 Is a Novel Therapeutic Target in Lymphoma. *Hemasphere.* 2023; 7(4):e864.

14. Pfeiffer C, Grandits AM, Asnagli H, et al. CTPS1 is a novel therapeutic target in multiple myeloma which synergizes with inhibition of CHEK1, ATR or WEE1. *Leukemia.* 2024; 38(1):181-192.

15. Wu F, Mao Y, Ma T, et al. CTPS1 inhibition suppresses proliferation and migration in colorectal cancer cells. *Cell Cycle.* 2022; 21(24):2563-2574.

16. Durand R, Bellanger C, Kervoëlen C, et al. Selective pharmacologic targeting of CTPS1 shows single-agent activity and synergizes with BCL2 inhibition in aggressive mantle cell lymphoma. *Haematologica.* 2024.

17. Sun Z, Zhang Z, Wang Q-Q, Liu J-L. Combined Inactivation of CTPS1 and ATR Is Synthetically Lethal to MYC-Overexpressing Cancer Cells. *Cancer Research.* 2022; 82(6):1013-1024.

18. Martin E, Minet N, Boschat AC, et al. Impaired lymphocyte function and differentiation in CTPS1-deficient patients result from a hypomorphic homozygous mutation. *JCI Insight.* 2020; 5(5).

19. Martin E, Palmic N, Sanquer S, et al. CTP synthase 1 deficiency in humans reveals its central role in lymphocyte proliferation. *Nature.* 2014; 510(7504):288-292.

20. Dharia NV, Kugener G, Guenther LM, et al. A first-generation pediatric cancer dependency map. *Nat Genet.* 2021; 53(4):529-538.

21. Gwynne WD, Suk Y, Custers S, et al. Cancer-selective metabolic vulnerabilities in MYC-amplified medulloblastoma. *Cancer Cell.* 2022; 40(12):1488-1502.e1487.

22. O'Connor SA, Feldman HM, Arora S, et al. Neural G0: a quiescent-like state found in neuroepithelial-derived cells and glioma. *Mol Syst Biol.* 2021; 17(6):e9522.

23.   Weishaupt H, Johansson P, Sundström A, et al. Batch-normalization of cerebellar and
medulloblastoma gene expression datasets utilizing empirically defined negative control
genes. *Bioinformatics.* 2019; 35(18):3357-3364.

24.   Cavalli FMG, Remke M, Rampasek L, et al. Intertumoral Heterogeneity within
Medulloblastoma Subgroups. *Cancer Cell.* 2017; 31(6):737-754.e736.

25.   de Haas T, Hasselt N, Troost D, et al. Molecular risk stratification of medulloblastoma
patients based on immunohistochemical analysis of MYC, LDHB, and CCNB1 expression.
Clin Cancer Res. 2008; 14(13):4154-4160.

26.   McPartland RP, Wang MC, Bloch A, Weinfeld H. Cytidine 5'-triphosphate synthetase as a
target for inhibition by the antitumor agent 3-deazauridine. *Cancer Res.* 1974;
34(11):3107-3111.

27.   Diehl FF, Miettinen TP, Elbashir R, et al. Nucleotide imbalance decouples cell growth from
cell proliferation. *Nat Cell Biol.* 2022; 24(8):1252-1264.

28.   Ecker J, Thatikonda V, Sigismondo G, et al. Reduced chromatin binding of MYC is a key
effect of HDAC inhibition in MYC amplified medulloblastoma. *Neuro Oncol.* 2021;
23(2):226-239.

29.   Morfouace M, Shelat A, Jacus M, et al. Pemetrexed and Gemcitabine as Combination
Therapy for the Treatment of Group3 Medulloblastoma. *Cancer Cell.* 2014; 25(4):516-529.

30.   Morrison LC, McClelland R, Aiken C, et al. Deconstruction of medulloblastoma cellular
heterogeneity reveals differences between the most highly invasive and self-renewing
phenotypes. *Neoplasia.* 2013; 15(4):384-398.

31.   Costantino L, Ferrari S, Santucci M, et al. Destabilizers of the thymidylate synthase
homodimer accelerate its proteasomal degradation and inhibit cancer growth. *Elife.* 2022;
11.

32.   Guijarro MV, Kellish PC, Dib PE, et al. First-in-class multifunctional TYMS nonclassical
antifolate inhibitor with potent in vivo activity that prolongs survival. *JCI Insight.* 2023;
8(10).

33.   Schild T, Keshari KR. Revealing de novo pyrimidine synthesis as a key vulnerability in brain
tumors. *Cancer Cell.* 2022; 40(12):1457-1458.

34.   Endersby R, Whitehouse J, Pribnow A, et al. Small-molecule screen reveals synergy of cell
cycle checkpoint kinase inhibitors with DNA-damaging chemotherapies in
medulloblastoma. *Science Translational Medicine.* 2021; 13(577):eaba7401.

35.   Ianevski A, Giri AK, Aittokallio T. SynergyFinder 3.0: an interactive analysis and consensus
interpretation of multi-drug synergies across multiple samples. *Nucleic Acids Research.*
2022; 50(W1):W739-W743.

36.   Stewart DJ, Benvenuto JA, Leavens M, et al. Penetration of 3-Deazauridine into Human
Brain, Intracerebral Tumor, and Cerebrospinal Fluid1. *Cancer Research.* 1979;
39(10):4119-4122.

37.   Hirano A, Matsui T. Vascular structures in brain tumors. *Hum Pathol.* 1975; 6(5):611-621.

38.   Phoenix TN, Patmore DM, Boop S, et al. Medulloblastoma Genotype Dictates Blood Brain
Barrier Phenotype. *Cancer Cell.* 2016; 29(4):508-522.

39.   Weng ML, Zalkin H. Structural role for a conserved region in the CTP synthetase glutamine
amide transfer domain. *J Bacteriol.* 1987; 169(7):3023-3028.

40.   Ortlund E, Lacount MW, Lewinski K, Lebioda L. Reactions of Pseudomonas 7A glutaminase-
asparaginase with diazo analogues of glutamine and asparagine result in unexpected
covalent inhibitions and suggests an unusual catalytic triad Thr-Tyr-Glu. *Biochemistry.*
2000; 39(6):1199-1204.
41.   Pham K, Maxwell MJ, Sweeney H, et al. Novel Glutamine Antagonist JHU395 Suppresses
MYC-Driven Medulloblastoma Growth and Induces Apoptosis. *J Neuropathol Exp Neurol.*
2021; 80(4):336-344.
42.   Hanaford AR, Alt J, Rais R, et al. Orally bioavailable glutamine antagonist prodrug JHU-083
penetrates mouse brain and suppresses the growth of MYC-driven medulloblastoma.
Transl Oncol. 2019; 12(10):1314-1322.
43.   Campagne O, Davis A, Maharaj AR, et al. CNS penetration and pharmacodynamics of the
CHK1 inhibitor prexasertib in a mouse Group 3 medulloblastoma model. *Eur J Pharm Sci.*
2020; 142:105106.
44.   Minet N, Boschat AC, Lane R, et al. Differential roles of CTP synthetases CTPS1 and CTPS2
in cell proliferation. *Life Sci Alliance.* 2023; 6(9).
45.   Mariotto E, Rampazzo E, Bortolozzi R, et al. Molecular and functional profiling of
chemotolerant cells unveils nucleoside metabolism-dependent vulnerabilities in
medulloblastoma. *Acta Neuropathol Commun.* 2023; 11(1):183.
46.   Lemberg KM, Vornov JJ, Rais R, Slusher BS. We're Not "DON" Yet: Optimal Dosing and
Prodrug Delivery of 6-Diazo-5-oxo-L-norleucine. *Mol Cancer Ther.* 2018; 17(9):1824-1832.
47.   Nedelcovych MT, Tenora L, Kim BH, et al. N-(Pivaloyloxy)alkoxy-carbonyl Prodrugs of the
Glutamine Antagonist 6-Diazo-5-oxo-l-norleucine (DON) as a Potential Treatment for HIV
Associated Neurocognitive Disorders. *J Med Chem.* 2017; 60(16):7186-7198.
48.   Rais R, Jančařík A, Tenora L, et al. Discovery of 6-Diazo-5-oxo-l-norleucine (DON) Prodrugs
with Enhanced CSF Delivery in Monkeys: A Potential Treatment for Glioblastoma. *J Med*
Chem. 2016; 59(18):8621-8633.
49.   Lee K, Thompson EA, Gharaie S, et al. T cell metabolic reprogramming in acute kidney
injury and protection by glutamine blockade. *JCI Insight.* 2023; 8(12).
50.   Praharaj M, Shen F, Lee AJ, et al. Metabolic reprogramming of tumor-associated
macrophages using glutamine antagonist JHU083 drives tumor immunity in myeloid-rich
prostate and bladder cancer tumors. *Cancer Immunol Res.* 2024.
768

Hathaway et al. Figure 1

Hathaway et al. Figure 2

Hathaway et al. Figure 3

Hathaway et al. Figure 4

Hathaway et al. Figure 5

Table 1: Synergy scores for CTPS1 inhibitiors with prexasertib

	Drug 1	Drug 2	ZIP_synergy	ZIP_synergy_p_value	HSA_synergy	HSA_synergy_p_value	Loewe_synergy	Loewe_synergy_p_value	Bliss_synergy	Bliss_synergy_p_value
HD-MB03	prexasertib	DAU	12.60639662	5.33E-20	17.40683667	3E-117	10.83592936	0.000055	12.0364247	6.25E-15
	prexasertib	JHU395	5.489365734	0.000469	8.794607635	2.34E-08	-1.769786284	0.886	5.237058489	0.00347
	prexasertib	JHU083	6.122734406	0.0211	12.55551008	8.69E-21	9.364874489	0.000188	5.965281089	0.027
Med-411FHTC	prexasertib	DAU	11.03533493	4.03E-19	13.89670691	2.03E-92	8.049172603	0.101	10.78442356	2.35E-14
	prexasertib	JHU395	-1.533340467	0.655	-4.356167358	0.0353	-7.164625504	0.172	-3.25086577	0.396
	prexasertib	JHU083	8.805722263	0.0000897	18.33827545	6.16E-18	4.763895713	0.534	8.852162691	0.0000665