The Wound Environment Agent-based Model (WEABM): a digital twin platform for characterization and complex therapeutic discovery for volumetric muscle loss

Chase Cockrell[1], Yoram Vodovotz[2,3], Ruben Zamora[2], Gary An[1]*

1. Department of Surgery, University of Vermont Larner College of Medicine
2. Department of Surgery, University of Pittsburgh
3. McGowan Institute of Regenerative Medicine, University of Pittsburgh

To Whom correspondence should be addressed:

docgca@gmal.com or gan@med.uvm.edu

Keywords: Volumetric muscle loss, wound healing, agent-based model, deep reinforcement learning, artificial intelligence, machine learning, adaptive control

Abstract:

Volumetric Muscle Loss (VML) injuries are characterized by significant loss of muscle mass, usually due to trauma or surgical resection, often with a residual open wound in clinical settings and subsequent loss of limb function due to the replacement of the lost muscle mass with non-functional scar. Being able to regrow functional muscle in VML injuries is a complex control problem that needs to override robust, evolutionarily conserved healing processes aimed at rapidly closing the defect in lieu of restoration of function. We propose that discovering and implementing this complex control can be accomplished by the development of a Medical Digital Twin of VML. Digital Twins (DTs) are the subject of a recent report from the National Academies of Science, Engineering and Medicine (NASEM), which provides guidance as to the definition, capabilities and research challenges associated with the development and implementation of DTs. Specifically, DTs are defined as dynamic computational models that can be personalized to an individual real world "twin" and are connected to that twin via an ongoing data link. DTs can be used to provide control on the real-world twin that is, by the ongoing data connection, adaptive. We have developed an anatomic scale cell-level agent-based model of VML termed the Wound Environment Agent Based Model (WEABM) that can serve as the computational specification for a DT of VML. Simulations of the WEABM provided fundamental insights into the biology of VML, and we used the WEABM in our previously developed pipeline for simulation-based Deep Reinforcement Learning (DRL) to train an artificial intelligence (AI) to implement a robust generalizable control policy aimed at increasing the healing of VML with functional muscle. The insights into VML obtained include: 1) a competition between fibrosis and myogenesis due to spatial constraints on available edges of intact myofibrils to initiate the myoblast differentiation process, 2) the need to biologically "close" the wound from atmospheric/environmental exposure, which represents an ongoing inflammatory stimulus that promotes fibrosis and 3) that selective, multimodal and adaptive local mediator-level control can shift the trajectory of healing away from a highly evolutionarily beneficial imperative to close the wound via fibrosis. Control discovery with the WEABM identified the following design principles: 1) multimodal adaptive tissue-level mediator control to mitigate pro-inflammation as well as the pro-fibrotic aspects of compensatory anti-inflammation, 2) tissue-level mediator manipulation to promote myogenesis, 3) the use of an engineered extracellular matrix (ECM) to functionally close the wound and 4) the administration of an anti-fibrotic agent focused on the collagen-producing function of fibroblasts and myofibroblasts. The WEABM-trained DRL AI integrates these control modalities and provides design specifications for a potential device that can implement the required wound sensing and intervention delivery capabilities needed. The proposed cyber-physical system integrates the control AI with a physical sense-and-actuate device that meets the tenets of DTs put forth in the NASEM report and can serve as an example schema for the future development of Medical DTs.

1.0 Introduction to Volumetric Muscle Loss Injury

Volumetric muscle loss (VML) is loss of skeletal muscle following traumatic injury that results in a functional deficit due to the healing of the wound with non-functional fibrotic scar instead of functional muscle [1, 2]. Wound healing of a VML (or more generalized skeletal muscle) injury in mammals is considered to progress in three broad phases: 1) hemostasis, occurring immediately to hours after the injury, 2) inflammation, occurring from hours to days post-injury, and finally, 3) repair, occurring in days to months post-injury [3, 4]. In the hemostatic phase, chemokines, cytokines and other molecular mediators indicating tissue injury are released, which will cause the initial recruitment of leukocytes. Polymorphonuclear neutrophils are the first inflammatory cells to arrive and have been observed as early as one-hour post-injury. [3, 5-7] Endothelial cells release interleukin-8 (IL-8), platelet activation factor (PAF), and a host of other mediators which influence the neutrophils to attach to the endothelial wall. The neutrophils then extravasate through the blood vessel wall and enter the wound, where they eliminate foreign debris and kill bacteria, if present [8]. Subsequently, the neutrophils undergo apoptosis and their pro-inflammatory functionality is replaced by M1-polarized macrophages [3, 9], which also have antimicrobial and phagocytic functions [10].

After this initial pro-inflammatory phase, negative feedback processes lead to macrophage population polarity transitions to the M2 (anti-inflammatory) phenotype, accentuating the suppression of pro-inflammatory processes and causing increased cell proliferation and influx of fibroblasts, extracellular matrix (ECM) synthesis, and capillary budding [11-16]. Over time, depending on the size of the VML injury, fibroblasts differentiate into myofibroblasts, which accentuate wound contraction, the formation of granulation tissue and subsequent conversion to fibrotic scar, marked by the replacement of collagen-3 with collagen-1 [6, 7, 12, 17-19]. The result is a healed, but nonfunctional volume of fibrotic scar tissue existing in place of muscle.

Evolutionary imperatives to enhance the survival of an organism to injury have led to the prioritization of fibrosis and wound closure over the regrowth of functional muscle. A VML injury involves significant loss of tissue and results in a relatively large wound surface area that is exposed to the external environment. To maximize survivability, it is critical to minimize pathogens from the external environment from invading the wound, and thus, the mammalian system quickly fills the missing volume with granulation tissue (i.e., extracellular matrix (ECM), fibrin, fibronectin, etc.) which is ultimately remodeled into stable fibrotic scar tissue [3, 12, 20]. It is this highly evolutionarily conserved, generally beneficial process that must be overridden to facilitate the regrowth of functional muscle in these VML. We propose that this is a complex control problem that requires manipulation of the cellular-molecular biology of skeletal muscle's response to injury and is suited to the application of a Digital Twin (DT) paradigm.

1.1 Medical Digital Twin of VML: Personalization and Precision Medicine

The ultimate goal of biomedical research is to provide the right treatment for the right patient at the right time. We have formalized these principles into the Axioms of "True" Precision Medicine [21]:

- Axiom 1: Patient A is not the same as Patient B (Personalization)
- Axiom 2: Patient A at Time X is not the same as Patient A at Time Y (Precision)
- Axiom 3: The goal of medicine is to treat, prognosis is not enough (Treatment)
- Axiom 4: Precision medicine should find effective therapies for every patient and not only identify groups of patients that respond to a particular regimen (Inclusiveness)

We believe that these Axioms have considerable overlap with the concept of Digital Twins (DTs). The concept of DTs derives from manufacturing and operations fields, including NASA, originating in the 1990-2000s as a means of using computer analogs to real world objects or processes to aid in these products' lifecycle management, facilitating product monitoring and maintenance. The original formal definition of a DT includes [22]:

- A data structure for the real-world system
- Some process that links data together to form dynamics
- Some link to the real world that feeds back data into the data-propagation/generation process

DTs have found extensive use in industrial applications, both for manufacturing processes and specific engineered systems (e.g., aircraft engines). The key appeal of DTs is that they are "twinned" to a specific real-world object, where the "twinness" between the digital and real-world object is maintained by an ongoing data link between the twins; this allows simulations to be run on the DT to predict the behavior of the real-world twin, anticipating potential failures and proscribing an individualized maintenance plan. Given the rapid and unstructured expansion of the use of the description "Digital Twin," the United States National Academies of Science, Engineering and Medicine (NASEM) sought to bring some structure and provide guidance by generating an extensive report on this technological approach. NASEM released its report "Foundational Research Gaps and Future Directions for Digital Twins (2023)" after an in-depth evaluation of the potential application of DT to Atmospheric Science, Engineering and Biomedicine [23]. We consider this report (henceforth referred to as the "NASEM Report") to be an authoritative statement on what constitutes a DT, its unique capabilities and associated research challenges for the promise of the technology to be eventually met.

One of the most common questions regarding DTs is "what is the difference between a DT and a computational model"? The NASEM Report (page 2) addresses this with the following text:

> "*Finding 2-1: A digital twin is more than just simulation and modeling.*
> *Conclusion 2-1: The key elements that comprise a digital twin include (1) modeling and simulation to create a virtual representation of a physical counterpart, and (2) a bidirectional interaction between the virtual and the physical. This bidirectional interaction forms a feedback loop that comprises dynamic data-driven model updating (e.g., sensor fusion, inversion, data assimilation) and optimal decision-making (e.g., control, sensor steering).*"

This definition distinguishes a DT from a personalized predictive computational model, such as an -omics based profile model. which are predicated on a feature set regarding a patient inputted at time of the creation of the model and is not (and often cannot be) updated in an ongoing fashion. We emphasize selected findings from the NASEM Report that directly impact the implementation of a DT to increase the functional recovery from VML. These include:

- Importance of Fit for purpose in defining the requirements of a DT
- Importance of ongoing data-linkage between the real and virtual worlds
- Importance of Establishing Trust in the DT (Validation and Uncertainty Quantification)
- Importance of control as a purpose of the DT

We propose that engineering a solution to the problem of VML will require an adaptive control process that instantiates our Axioms of Precision Medicine using the DT paradigm. Herein we present the computational specification for such a DT, the Wound Environment Agent-Based Model (WEABM). The WEABM is a computational representation of the process of wound healing from a VML injury. The current version of the WEABM is informed by a clinically relevant canine model of VML [24]. In the process of constructing this model, we have integrated knowledge concerning (muscle) wound healing on the volumetric and non-volumetric (e.g., muscle damage/repair from exercise), and have used a calibrated model to suggest broadly defined therapeutic strategies to increase functionality/volume on functional muscle in wounds healing from VML injuries.

1.2 Fit-for-purpose and overall description of WEABM

We applied our previously developed schema for determining the design criteria for a VML DT and its computational specification the WEABM [25]. The key features identified are:

- Fit for Purpose: Goal is to identify means of increasing muscle regrowth in VML by manipulation of cellular behavior via molecular-level interventions.
- Primary Scale of Representation: Cellular behavior, aggregated into system-level phenotype (e.g., the VML wound).
- Data Linkages: Hypothesized, based on requirements for effective control.

These results led to the decision to construct the WEABM as an Agent-based Model (ABM). Given the clinically relevant anatomic scale of the entire system, the WEABM was implemented in C++ to facilitate execution on high-performance computing platforms. Agent-based modeling is an object-oriented, discrete-event, rule-based, spatially-explicit, stochastic modeling method in which agents interact with each other and with their environment according to a pre-defined set of rules. In this work, the rules are informed by both literature and experimental data. Literature review has been used to establish the existence of a rule, e.g., interleukin-13 (IL13) stimulates fibroblasts to produce Transforming Growth Factor β1 (TGFβ1) [18], and the experimental data is then used to quantify the contribution/effect of IL13 on fibroblast production of TGFβ1 in the model. We note that parameter values of this type are inherently unitless and not directly translatable to the real world (see 'Model Calibration' in Methods). In this sense, an ABM is a computational/dynamic instantiation of mechanistic knowledge that has been established through rigorous and reproducible biological experiments. Because ABM's synthesize knowledge outside the domain expertise of a single researcher, behaviors generated by the aggregate behavior of the agents in the model can often be non-trivial or unintuitive, rendering ABM an excellent technique to investigate biomedical systems. The utility of ABM's spans from general purpose anatomic/cell-for-cell representations of organ systems capable of reproducing multiple independent phenomena [26, 27] to platforms for drug development,[28, 29] and are frequently used to model non-linear dynamical systems such as the human immune system [30-33]. For this initial implementation, the WEABM is calibrated to a canine model of VML. This model was chosen because it: 1) can produce a large enough wound such that it is contextually clinically relevant (12 cm x 7 cm x 4 cm) in the thigh, 2) allows for sequential tissue sampling so that tissue mediator time series can be obtained to capture the dynamics of the VML response, and 3) allows for experiments to be potentially performed to evaluate potential control strategies aimed at increasing the amount of functional muscle in the healing of the VML injury.

Figure 1: Schematic of the WEABM Immune cells are orange boxes, muscle regeneration cells and neurovascular components are depicted in pink, scar-promoting cells in blue and extracellular matrix/collagen in green. Green arrows denote promoting interactions, red denote downregulation blue represents upregulation and black represents differentiation.

In the WEABM, the individual agents are cells (see Figure 1 for a wire-frame schematic of the cellular-molecular interactions implemented and see Appendix A for "Facts and References for the WEABM") and the environment is discretized and represented as a three-dimensional grid of cubic voxels measuring 100 µm per side, as this is comparable to the diameter of a type-1 skeletal muscle fiber in the quadricep of a female German Shepherd [34, 35]. Each tissue voxel has three primary data elements associated with it: 1) volume of various tissues: each voxel can be filled with some combination of healthy muscle (which requires adjacent healthy muscle), collagen-3 (broadly representing granulation tissue), and collagen-1 (broadly representing fibrotic scar); [7, 19, 36-39] 2) a Boolean variable indication vascularization, which requires vascularization of a contiguous voxel; and 3) a Boolean variable indicating innervation, which required innervation of a contiguous voxel. Muscle fiber orientation is represented as a restriction on the movement of satellite cells such that, in the undamaged tissue, muscle fibers are oriented along the y-axis of the simulation and satellite cells are myoblasts are required to migrate along a single myofiber. In damaged regions, satellite cells and myoblasts are allowed free movement [5, 6, 40].

An execution of the WEABM proceeds by first instantiating the wound, and then incrementing the simulation in time. Each time step of the simulation represents 15 minutes. At the start of each step, the different cell types are randomly shuffled, and each cell then performs its actions sequentially. After the cells perform their actions, the diffusion and degradation of mediators and cytokines are simulated. Cells that have died (from apoptosis or necrosis) are then removed from the simulation and any new cells generated from proliferation or division are added to the simulation world.

The WEABM can be run in two primary modes: 1) the full anatomic scale that corresponds to the 12 cm long x 7 cm wide x 4 cm deep experimental wound in the reference canine model and can represent up to 400 million cells or 2) a biopsy scale that represents a smaller volume (approximately 6 mm long x 6 mm wide x 6 mm deep) corresponding to the size of tissue biopsies in the reference canine model [24].
In the anatomic scale configuration initial conditions are generated through processing data obtained from a CT scan of the canine model immediately after the application of the surgical wound; this allows for personalization of the WEABM to a specific individual consistent with the digital twin paradigm. We first transform the surface to remove the curvature of the dog's leg, as described in [41], which essentially projects the wound into a Euclidean space with the surface of the wound at the top of a rectangular prism and the depth of the wound going through the z-axis (See Figure 2).

Figure 2: Heatmap of XYZ Axis representation of the WEABM with initial conditions mapped from the first post-surgical Computer Tomography (CT) scan of a canine wound. Color scale is related to the depth of the wound (blue deep and yellow superficial). Tissue is shaded based on distance from the ideal wound surface, shaded from yellow (superficial) to blue (deep). The panel on the left shows a top-down view of the wound (looking down the z-axis at the x-y plane) and the panel on the right shows a side view (looking along the x-axis at the y-z plane).

We consider all tissue that is exposed to the atmosphere (and all tissue below that along the z-axis) to be healthy muscle. Given the fact that one of the hallmarks of acute VML is that the sizes of the defects preclude closure of the wound with biological tissue, we recognize that the surface of the wound is exposed to the atmosphere/environment and either that atmospheric exposure and/or any dressing/wound device applied to that surface will generate an ongoing inflammatory stimulus that would initiate and propagate the injury response process. A rendering of a personalized initialization of the anatomic scale WEABM can be seen in Figure 3.

Figure 3: Rendering of a personalized initialization of the WEABM. This image adds realistic tissue textures to the underlying voxel structure of the WEABM. Rendering produced by 4dThieves™.

In the biopsy configuration, the WEABM simulates a volume of tissue which is comparable to the experimental biopsies, specifically, a 6-mm (60 voxels on each side) cubic section of tissue represented with a rectangular prism of tissue. The WEABM in the biopsy configuration can grow to a larger depth than 6mm, however only the top 6mm of tissue are used to compare protein concentrations with the experiment and the very top of the Z-axis represents the wound-atmosphere interface noted above. Further, there is an additional abstraction here regarding the location of satellite stem cell pools. In the anatomic configuration, each muscle fiber has at least one satellite stem cell pool randomly located in the healthy (uninjured) tissue at the boundary of the simulation; in the biopsy configuration, the only healthy tissue is located at the base of the tissue volume, and thus, we allow myoblasts to initiate new fibers at one edge of the simulation-space.

There has been extensive prior work using multiscale mechanism-based modeling to study skeletal muscle injury and recovery, both using finite element methods [42, 43], and ABMs [44-46], including work using an ABM specifically examining and proposed means of improving muscle regrowth in VML [47]. The WEABM has many similarities to the work of Westman [47], particularly in terms of the inclusion of the primary cell types involved and the molecular signaling pathways present in VML. Primary features that distinguish the WEABM from prior work include: 1) the fact that the WEABM is a 3-dimensional model, 2) the WEABM is capable of representing clinically relevant, anatomic scale injuries and 3) the representation of the WEABM incorporates a critical pathophysiological feature seen in clinical VML, namely that the wound is not covered with biological tissue and therefore the wound-environment interface represents an ongoing injury/inflammatory source in VML.

To demonstrate the potential utility of the WEABM as the basis of a VML DT, we performed a series of simulation experiments with the WEABM with the goal of governing a complex adaptive control process that would increase the recovery of functional muscle in VML. The simulation experiments include:

- Initial Evaluation of Behavior Space of the WEABM to provide fundamental insights into the healing biology of VML
- Initial Calibration to Tissue Protein/Mediator Data to fulfill the validation requirement noted in the NASEM Report.
- Perform control discovery using simulation-based Deep Reinforcement Learning (DRL)
  - Simulated Control Modalities:
    - Multimodal mediator administration
    - Application of Extracellular Matrix (ECM)
    - Combination multimodal therapy including a hypothetical anti-fibrosis agent

The results of these simulation experiments are provided below. The code for the WEABM and the DRL environment can be found at https://github.com/An-Cockrell/WEABM_DRL/tree/main.

2.0 Results:

2.1 Pre-Calibration Insight: The essential role of the competition between fibrosis and myogenesis

Initial simulation experiments explored model behavior space arising from the inherent model structure of the WEABM. These simulations provided fundamental insights into the dynamics that governed healing of VML, and therefore served to frame subsequent investigation of potential control strategies aimed at healing these wounds with potentially functional muscle. The first of these insights was the essential role of the competition between fibrosis and myogenesis. Figure 4 shows a set of early pre-calibration simulations that examine the behavioral range of the WEABM.

Figure 4. Parameter space exploration demonstrating competition between fibrosis and myogenesis. With adjustment of parameters governing the rate of fibrosis functions and myogenesis functions, a progression of tissue composition during healing can be seen in the progression of Panel A through F, where Panel A represents the "native" configuration where fibrosis predominates, Panels B-E depict a shift between fibrosis and myogenesis, towards Panel F, where effective muscle regrowth is generated. Though uncalibrated, these dynamics show a potential path towards the desirable outcome of increased functional healing.

Specifically, we focused on the processes associated with fibrosis and myogenesis. The model structure demonstrated that there is a fundamental tradeoff between these two processes, a competition for binding sites on the edge of the damaged health muscle where myoblasts would fuse to differentiate into functional myofibrils. We term this competition "the race between fibrosis and myogenesis," and this dynamic informs our subsequent calibration simulations and the process of control discovery.

2.2 Calibration and Validation:

This first step in calibration is determining if the overall healing dynamics of the canine VML can be reproduced. These results can be seen in Figure 5, which compares the wound closure pattern between experimental data and WEABM simulations.

Figure 5. Comparison of tissue growth. Panel A: Total new tissue growth in each dog calculated from CT data, as well as the best-fit line in red. We note that there are many sources of experimental error when obtaining this experimental measurement, including dog motion and tissue swelling, and as such, we posit that the best-fit line is representative of true tissue growth. Panel B: Total tissue in the volume simulated as the wound evolves over 40 days.

The visual depiction of this process is seen in Figure 6, which shows successive renderings of the VML wound from a specific experimental animal as it heals.

Figure 6. Selected images of the WEABM as it heals from a VML over 40 days simulated time (Days, 0, 7, 30 and 40). Each panel displays a representation of the wound surface only (not the bulk tissue) for rendering reasons. Tissue is shaded based on distance from the ideal wound surface, shaded from yellow (superficial) to blue (deep). The panel on the left shows a top-down view of the wound (looking down the z-axis at the x-y plane) and the panel on the right shows a side view (looking along the x-axis at the y-z plane). Note that since this represents baseline biology the majority of the wound is filled with scar.

Having demonstrated face validity by reproducing the overall healing dynamics of VML injury the next step in calibration is the fitting of the WEABM to experimental time series data of mediator levels in the wound. Our novel process to complex model calibration and validation is described in the Methods, and involves the identification of all model structure and parameter configurations of the WEABM that are sufficient to

reproduce the range of the experimental data using a mathematical object called the Model Rule Matrix (MRM) [48, 49] and a machine learning (ML) search pipeline that includes genetic algorithms (GA) and active learning (AL) that we term Nested Active Learning [50]. The result of this process is a set of model structure and parameter configurations that we term *bioplausible* insomuch they cannot be falsified by any available data. As previously noted, the calibration data are drawn from Ref [24]. The results of this process are seen in Figure 7.

Figure 7: Comparison of WEABM bioplausible output to experimental data from different zones of wound for representative mediators MCP-1 (Panel A), IFNγ (Panel B), IL-10 (Panel C), TNFα (Panel D) and active TGFβ (Panel E). Blue bars are experimental data based on location within wound; the different wound zones are shown for demonstration purposes, with the center zone as the left-most blue bar, the intermediate zone as the center blue bar, and the edge as the right-most blue bar for each day, as the data across these regions are aggregated for determination of bioplausible trajectories of the WEABM (since all regions demonstrate values reachable/bioplausible in the real-world system). Red bars are simulation output of the WEABM at corresponding time points; note that WEABM simulated data projects what values would have been at Day 5 and 6.

We further define bioplausibility as model behavior being within the bounds of the experimental data while maintaining as much behavioral heterogeneity as possible; this is intended to mimic the ability of a particular model configuration to reproduce the breadth of behavior seen in a population, thereby providing generalizability as a property of the model representation. We also note that the Nested Active Learning fits to all selected variables (e.g., multidimensional fitting) at the same time (in this case all the measure mediators in the WEABM, of which 5 are shown in Figure 7). Parameter determination currently equally weighs every mediator for which experimental data exist, therefore there is differential quality of calibration in the 5 mediators shown here. For the most part, the experimental data are contained within the range of

ensemble-simulation generated data, with the exception of the spike in TNFα at day 7. Deeper investigation into these measurements identified that they derived from a single animal that was experiencing systemic inflammation due to causes unrelated to the experimental VML and therefore likely had increased systemic levels of TNFα. This finding, rather than showing a limitation of calibration through Nested Active Learning, demonstrates the ability of the process to determine the bounded expressiveness of the underlying model structures as reflected by the set of bioplausible MRMs and can even suggest alternative explanations for outlier measurements. This set of bioplausible MRMs now is the superset of WEABM configurations that are used on in subsequent simulation experiments.

The step in validation is the ability of the WEABM to provide forecasting of individual trajectories. The range and heterogeneity in the data set is considerable, as would be expected from a population of non-genetically identical animals (or a clinical population) undergoing a complex surgical procedure. It is impossible, therefore, to attempt to pre-specify where in that entire possible trajectory space an individual animal's data will reside. However, the DT paradigm, with its ongoing data link between the real world and the digital object, inherently accommodates the concept of a *rolling forecasting cone*, wherein, given a particular point in system state space, a range of possible trajectories can be generated using any ensemble of MRM configurations able to encompass that particular set of multidimensional data points. This cone is then updated with additional data feeds from the real world and propagates forward in time. This is the same concept used in forecasting hurricane paths. The capability of the WEABM is demonstrated in Figure 8, which shows a refined forecasting cone generated by a defined ensemble of MRM configurations for an individual animal as a subset of the total possible trajectories.

Figure 8. Demonstration of prediction cone for an individual animal. Trajectory spaces are shaded areas, actual experimental measurements are shown as dots. Blue Dots: Experimental data points from all dogs (N = 16). Blue Shaded Area: Total possible trajectory space generated by WEABM (encompasses all Blue Dots). Red Dots: Data points from an individual animal. Red Shaded Area: Forecasting cone for the individual animal, with re-fitting based on updated data. Panel A = TNFα, Panel B = IL-10, Panel C = active TGFβ. The refinement of the forecasting cone over time is analogous to the rolling prediction seen in hurricane prediction, and consistent with the DT paradigm in terms of an updating data feed.

The results of the calibration and validation simulations provided us a baseline level of trust that the WEABM plausibly reproduced critical features of the real-world VML model in terms of overall wound healing dynamics, reproduction of the heterogeneity and trajectories of mediator levels and the ability to individualize/personalize an ensemble of model configurations to a specific individual. As noted in the defined fit-for-purpose of the WEABM-based VML DT, we now turn to exploration of potential control modalities that would be able to increase the amount of functional muscle in the VML wound.

2.3 Examining Control Modalities:

2.3.1 Insights into control components:

Simulation experiments with the WEABM during the model exploration and calibration phases provided insights into the key dynamic features that would be needed to be targeted with putative control strategies. These insights included:

1. Confirmation of importance of competition between fibrosis and myogenesis (see Figure 4). Once collagen deposition occurs, caps the ends of the myofibrils, no myoblast fusion can occur, therefore no muscle regeneration. We tested this hypothesis by arbitrarily enhancing myoblast function to a 4-fold increase in proliferation. The results of this simulation can be seen in Figure 9.

Figure 49: Simulation experiments of hypothesized role of ECM in enhancing myoblast proliferation. Solid Lines: Baseline non-intervention simulation. Dotted Lines: Simulation of ECM effect that increases rate of myoproliferation 5-fold. Red: Healthy muscle. Black: Dead/necrotic muscle. Blue: Collagen deposited in the wound. Vertical Dashed Line: Application of intervention (timed to be applied after the loss of at-risk muscle from initial injury). Baseline simulation demonstrates expected loss of at-risk muscle early, followed by "natural" time course of healing, which involves the deposition of collagen and formation of scar. Simulated intervention leads to initial rise of muscle and myoblasts (as they spatially outcompete collagen formation) but lack of appropriate fusion points results in the production of myoblasts that subsequently necrose and form additional pro-inflammatory stimuli that kills healthy muscle.

3. The resulting control paradox: The M2 phenotype was needed to curtail the pro-inflammatory phase but counter-productive to the therapeutic goal of muscle regrowth.

4. The open nature of the VML wound provides an ongoing pro-inflammatory stimulus to the surface of the wound through atmospheric/environmental exposure, essentially providing an ongoing perturbation that "resets" the injury clock, enhancing the amplitude and duration of the pro-inflammatory phase of wound healing (M1) and a subsequent augmentation of the pro-growth/fibrosis processed generated by the compensatory M2 response. The robustness of the fibrosis response arises from the evolutionary imperative to biologically close the wound.

All these features point to a complex control problem that can generally be categorized into three main phases:

- Early (up to ~ 3 days): emphasizing the efficacy and subsequent plateau of the initial pro-inflammatory/M1 response

- Transition to Growth (~7-14 days): key interval where the initial pro-inflammatory response is attenuated (if possible) and pro-fibrosis functions are initiated as a result.
- Late (after ~ 14-21 days): in this phase the initial collagen deposition has occurred, and subsequent remodeling would require degradation of existing collagen to re-expose myofibrils for fusion to new myoblasts or rests of differentiated muscle tissue.

We identify the key transition point as being in the 7-10-day period, and therefore posit that this period should be targeted by mediator manipulation to change the relative strengths and efficacy of fibrosis versus myogenesis pathways to shift the conditions of the race between these two modules. Given the known heterogeneity seen in the tissue and the dynamic nature of the processes we propose using the WEABM in a DT context to provide an adaptive sense/actuate control approach to the processes in this phase. More specifically, since this transition phase represents a balance between pro-inflammation (nominally M1), appropriate inflammation suppression via anti-inflammatory mediators (nominally M2), and pro-growth factors associated with the anti-inflammatory processes, we pose that the ability to categorize the wound requires: 1) the ability to sense the pro-inflammatory state of the wound and 2) the ability to sense the canonical anti-inflammatory/pro-growth mediator milieu. These sensing targets will be successively refined in the results of the deep reinforcement learning (DRL) control discovery below.

We recognize the critical and essential nature of the initial pro-inflammatory response, the issue with VML is the extent and duration of this process, and therefore measures should be applied to biologically "close the wound." Prior research has suggested a role for extracellular matrix (ECM) [1, 51-53], and therefore this will be incorporated into the multimodal control policy.

In terms of therapeutically addressing the late phase and attempting to modulate collagen degradation and redeposition, we recognize that the challenge here will be how to degrade Collagen III without re-activating pro-inflammatory pathways, as that is a known function of ECM-degrading enzymes such as matrix metalloproteinases (MMPs). Therefore, we deferred the investigation of this target at this phase of our work.

The following sections describe the results of the control discovery investigations, carried out in a modular fashion such that the necessity of each component can be demonstrated.

2.3.2 Initial Multimodal control using mediator manipulation

Our developed approach to discovering novel multi-modal and adaptive control policies to affect cellular-molecular processes towards a specified clinical goal involves simulation-based Deep Reinforcement Learning (DRL) [54]; the specifics of this approach are discussed in the Methods and are explained at length in Refs [55, 56]. The process of simulation-based DRL is also consistent with the NASEM Report guidance on DTs, and the requirements of this approach can be identified in our specified design process for a DT [25], and which was applied to the WEABM.

As with our prior work on simulation-based DRL to cure sepsis the initial proof-of-concept implementation of DRL training on the WEABM used the most extensive sense/observation and actuate/control spaces in terms of mediator state of the wound and potential mediators for manipulation: every mediator could be observed in the 15-minute step interval of the WEABM, and every mediator could be augmented or inhibited at this interval as well. This allows us to determine the full extent of controllability given an initial representation of the potential control space based on the representation level of the WEABM; the inability to reach a desired end-state would suggest the need to expand the potential control space. With the initial maximal observation and action space the DRL training returned the top 10 performing control policies seen in Figure 10.

Figure 10: Full Control and Observation Space Top 10 DRL Policy Performance for 1000 steps = 10.4 days. Panel A shows the performance of the top 10 policies in terms of regrowth of muscle. Note that muscle regrowth is not seen at all in VML in nature, and therefore any positive curve represents a qualitative change in wound healing dynamics. Panel B shows the concurrent production and deposition of Collagen 3. Note that the end state of the wound is still primarily fibrotic, as the upward inflection in Collagen 3 production/deposition leads to the plateau of muscle regrowth.

The resulting policies are shown in a heatmap of direction of action and strength of action in Figure 11.

Figure 11: Heatmap of Policies, Top 10 policies to the Right of the figure. As noted in the Legend, Blue represents inhibition of a particular mediator whereas Red represents the augmentation of the mediator, darker shades represent higher levels of manipulation whereas lighter shades represent smaller manipulations (white = neutral).

The top 10 policies are all qualitatively similar. They are relatively static/constant through the entire 10-day course of simulated time, with actions near the maximum amplitudes of either augmentation or inhibition. The augmented mediators are: IL6, MIP1β, TNFα, HGF, IL1, IL13, VEGF, IL4, Myf5, sIL1r, IGF, a damage-associated molecular pattern (likely HMBG-1) and MCP1. The Inhibited mediators are: PAF, PDGF, TGFβ1, GCSF, IL10, IL17A, IL6, IL1ra, sTNFr, IFNγ, IL12, MRF4 and FNE.

Next, we attempted to identify a minimally sufficient observation/sense and control/actuator space by selecting canonical mediators for distinctly identified functions. The selected control targets were:

- Pro-inflammatory Mediator: TNFα
- Pro-inflammatory product: Reactive Oxygen Species (ROS)
- Anti-inflammatory Mediator: IL10
- Anti-inflammatory-Profibrosis Mediator: TGFβ1
- Pro-muscle Growth Factor: Hepatocyte Growth Factor (HGF)

In terms of the observation space, we posited that the key tissue states of interest in this phase of wound healing would be the pro-inflammatory state of the wound, which we represented by summing all the pro-inflammatory mediators, and the pro-fibrotic state of the wound, for which we selected TGFβ1 as the sensing target. We acknowledge that there might be a confounding effect with TGFβ1 as both an actuate and sense target but believe that given the central role TGFβ1 plays as a key regulator of fibrosis our initial studies would evaluate whether this dual role in DRL training would impact the resulting control policies. DRL was then performed with sensing targets of aggregated pro-inflammatory mediators and TGFβ1 and

con... ...of TNFα, ROS, IL10, TGFβ1 and HGF. We also increased the maximal augmentation allowed to 10-fold increase over baseline tissue levels as opposed to doubling in the prior DRL training sequence. The results of this discovered policy can be seen in Figure 12.

Figure 12: Top 10 control policies with restricted observation and action space for 1000 Steps = 1... targets of aggregated pro-inflammatory mediators and TGFβ and control targets of TNFα, ROS, IL10... Maximal action augmentation 10x greater than employed in results seen in Figure 10. Panel A shows... greater than 10-fold more muscle being produced. Panel B, however, shows a similarly increased le... produced. The inflection point where muscle growth plateaus is similar to that seen in Figure 10, reinfo... that the rate-limiting feature is the available damaged ends of healthy myofibrils.

These results show that with increased action magnitude there can be an increase in the am... muscle; we subsequently performed a parameter sweep of increased amplification levels t... effect plateaued at a 10-fold increase over baseline tissue levels. However, the plateau of r... remains, with the significant remainder of the wound still closed with Collagen 3, which wou... be converted to non-functional scar tissue. With a smaller set of control targets, we are a... more granular analysis of the implemented control actions and depict the time courses of th... of the top 10 policies in Figure 13.

Figure 13: Time courses of control actions in the top 10 policies of DRL training with restricted observation and action space. The different colored lines represent different control policies arrived at via the DRL training. While there are variable levels of control from time point to time point, in general the policies implement suppression/inhibition of pro-inflammatory mediators TNFα and ROS, while augmenting IL10, TGFβ and HGF.

The control actions in the discovered control policies are relatively consistent, with inhibition of pro-inflammation and suppression of the M1 phenotype by reduction of TNFα and ROS and augmentation of IL10, TGFβ1 and HGF. Note that there is a discrepancy in the direction of control for certain mediators (specifically IL10) with the entire versus restricted control space; we interpret this as a result of restricting the control space such that it reduces the number of parallel pathways being affected. The variability seen

between the depicted control actions for each individual time course is more likely to be due to variation on the specific behavioral dynamics from run arising from the stochastic nature of the WEABM. The policy is interpretable, both in terms of the desired goal of suppressing the initial pro-inflammatory phase and the need to augment myogenesis (via HGF). However, the dual effects of IL10 and TGFβ as anti-inflammatory but pro-growth/fibrosis mediators, is reflected by the enhanced degree of Collagen 3 deposition seen. Our understanding of the required effects needed to enhance muscle regrowth in VML suggests that what is needed is a more targeted anti-fibrosis controller that does not include ambiguity in terms of its targeted functions. We will return to this potential control point after we investigate the two other control modalities introduced at the beginning of this Section.

2.3.2 Implementing the effect of ECM

We have noted that one of the clinical hallmarks of VML is the fact that it is an open wound that needs to be covered to avoid ongoing contamination, desiccation, and trauma. Even wound coverage with a non-bio-active dressing is itself local tissue trauma (by triggering a foreign body reaction), and therefore we consider the open nature of VML injury as inducing an ongoing source of inflammation at the tissue-atmospheric environmental interface. This ongoing inflammatory stimulus leads to a persistence of the pro-inflammatory phase of wound healing, with consequent enhancement of the pro-fibrosis processes. Therefore, we consider an essential component of an effective control strategy to enhance muscle regrowth in VML to be a means of *biologically closing the wound* if native tissue cannot be used to accomplish this. We therefore consider the application of an artificial manufactured ECM to serve this purpose. While we recognize that there is likely a plethora of signaling effects of the ECM, these remain insufficiently characterized to the degree of detail and specificity needed to inform a mechanistic model such as the WEABM. Because of this we make the abstracting modeling decision to treat the primary effect of ECM as being the dampening of the pro-inflammatory stimulus of tissue at the surface of the wound (75% Reduction of DAMP generation at the wound surface). We then performed simulations to determine if this hypothesized mechanism is sufficient to explain experimental data that employs ECM in the canine wound. The reference model is again that which is presented in Ref [24], but now including interval application of an ECM hydrogel. The results of the MRM calibration runs are seen in Figure 14.

Figure 14: Comparison of different MRM sets fit to the baseline native healing versus with ECM application. The Red Dots are experimental data points from non-ECM animals, the Red Shaded areas are trajectory spaces of MRM configurations arrived at using the Nested Active Learning Pipeline. The Blue Dots are experimental data points from animals with ECM application within the first 10 days, the Blue Shaded areas are trajectory spaces from MRM configurations defined by the Nested Active Learning Pipeline. The ability of these two trajectory spaces to encompass their respective data while implementing the effect of ECM as 75% reduction of DAMPs at the surface of the wound demonstrates the sufficiency of that modeling assumption. Also note that there are two outlier data points in the TNFα plot; this is a similar situation to the outliers noted in Figure 7D, where out-of-bound TNFα values could be attributed to specific animals that suffered a systemic illness.

To further demonstrate the importance of ECM implemented as biologically covering the wound, we examine the efficacy of the policy discovered in Figure 13 in circumstances where the control device is applied for a limited period and then removed and standard wound care is applied during the off-control interval. These simulations involved: 1) episodic application of the control, which is assumed to include an

ECM barrier to prevent atmospheric irritation of the wound surface and 2) representation of non-control intervals assumed to have standard non-biological dressings in contact with the wound surface, with consequent pro-inflammatory stimulation due to the environmental/foreign body interaction. The episode intervals were application of the control for 6 hours, followed by an off-control interval of 42 hours, for a full 48 hours of a complete episode of treatment. The questions being investigated with simulating these conditions were: 1) is a previously efficacious control policy generalizable to this different context of use, and 2) is it even possible to use DRL to train an AI control policy that would be efficacious with this context of use? The first set of simulations just tested the previously derived control policy seen in Figure 13 in a standard continuous application case, the second set of simulations tested the previously trained AI control policy in the episodic control context and the third involved implementing a DRL training program on the episodic application context. The results of these simulation experiments are seen in Figure 15.

Figure 15: Simulations of episodic therapy for 1400 steps = 14.6 days. Panel A: Performance of Agents 1, 2 and 3 in terms of regrowth of muscle. Panel B: Performance of Agents 1, 2 and 3 in terms of generated Collagen 3. Agent 1 = AI Controller trained and reported in Figure 13. Agent 2 = AI Controller trained on episodic application of control. The blue plot shows the performance of Agent 1 in a set of WEABM simulations similar to the continuous control conditions upon which Agent 1 was trained; the performance of this control policy is equivalent to that seen in Figure 13. The orange plot shows the performance of Agent 1 in the episodic control context; in this setting the previous control policy completely fails. The green plot shows the performance of Agent 2, the AI control policy trained in the episodic context. Here there is marginal increase in muscle that quickly plateaus and becomes ineffective after about the first 48-hr control cycle.

Figure 15 shows that the previously trained AI control policy remains effective in a test condition similar to its training conditions (as expected). However, this control policy fails when applied to the episodic control context. Moreover, evening attempting to train in the episodic context does not result in an effective control policy (Agent 2). The depiction of the time series control actions for all 3 conditions is seen in Figure 16.

Figure 16: Time courses of control actions taken during episodic control application where the control is applied for 6 hours followed by a 42-hour period off-control with standard wound care. Agent 1 = AI Controller trained and reported in Figure 13. Agent 2 = AI Controller trained on episodic application of control. The blue plot shows the performance of Agent 1 in a set of WEABM simulations similar to the continuous control conditions upon which Agent 1 was trained; actions performed with this control policy are similar to those control actions in Figure 13. The orange plot shows the performance of Agent 1 in the episodic control context; in this setting the previously trained control policy generates maximal actions during the control interval that suggests its sensing of the state of the wound is continually reset after each off-control interval. The green plot shows the performance of Agent 2, the AI control policy trained in the episodic context. This policy appears to result in actions that also reset after each off-control interval that are at a maximal amplitude of intervention.

The control actions taken during the episodic application of the control with intervals off-control with standard wound care suggest that from the standpoint of the AI control agent the biology of the wound resets after each off-control interval. This is consistent with our concept of the robustness of the baseline biological imperative to close the wound, and points to the need to incorporate some means of mitigating the pro-inflammatory stimulus from the wound-atmospheric/environmental interface. We pose that ECM can serve this purpose.

2.3.3 The Missing Piece: Limiting Fibroblast collagen deposition

The control modalities thus implemented and tested include:

- Manipulation of mediator milieu to attenuate pro-inflammation/M1 and foster anti-inflammation/M2 via a DRL-derived AI control policy
- Augmentation of myogenesis via HGF supplementation and electrical stimulation
- Mitigation of the pro-inflammatory stimulus from the wound-atmospheric/environmental interface from the open wound with ECM

While these control modalities can promote the regrowth of muscle in the wound (again, a phenomenon not seen in the natural case), the robustness of the fibrosis imperative still results in most of the wound filling with Collagen 3 and subsequently with non-functional scar (see Figures 10, 12 and 15). What is missing in the control space is the ability to directly mitigate the production of Collagen 3 from fibroblasts, or attenuate the differentiation of fibroblasts to myofibroblasts, which produce substantially more Collagen 3 and contract the wound. While there are known anti-fibrosis agents that have been tested in VML, such as losartan [57], the small molecule IDT1 [58] and nintedanib [59], we have elected to avoid potential off-target effects by proposing a hypothetical anti-fibrosis agent that would directly suppress the Collagen 3 production from fibroblasts as an additional control agent in a multimodal control strategy. This hypothetical compound is implemented and incorporated in the multimodal control regimen noted in the section below.

2.3.5 Multimodal Control: Putting it all together

Our interpretation of the results of the different control modalities evaluated with the WEABM and our reading of the existing literature led us to conclude that there are 4 primary control areas for enhancing the regrowth of functional muscle in VML. These 4 areas are:

1. Controlling the transition from the early pro-inflammatory M1 phase to the counter-inflammation/pro-growth M2 phase in the initial response to VML injury. This we believe can be controlled at the mediator level.
2. Mitigating the pro-fibrotic aspects of the M2 pro-growth phase of the VML response by targeted suppression of fibroblast-myofibroblast function. We believe this could be accomplished by the targeted administration of known antifibrosis compounds and/or the suppression of differentiation of fibroblasts into myofibroblasts.
3. Mitigating the ongoing inflammatory stimulus from the wound-atmospheric/environmental interface that drives continued M1 stimulation and the generation of the evolutionarily beneficial wound

fibrosis phenotype. We believe this can be accomplished by "closing the wound" using engineered ECM.

4. Augmenting existing myogenesis pathways, enhancing myoblast migration, fusion, and differentiation. We propose to use HGF as an endogenous compound to facilitate these mechanisms.

Underpinning these control areas are two fundamental insights:

1. There is a spatial constraint in the competition between fibrosis and myogenesis due to the limited edge/surface of damaged muscle to which any new muscle must attach.
2. The evolutionary imperative is to close the wound to limit the risk of infection. This process is fibrosis, and therefore fibrosis is a heavily favored process in the endogenous healing of VML.

Therefore, we implemented a multi-modal control strategy in the WEABM that included:

1. Mediator manipulation governed by a DRL-trained AI control policy that targeted TNFα, ROS, IL10, TGFβ1 and HGF. To demonstrate the need to include a pro-myogenesis mediator an additional control policy was generated with no HGF.
2. The addition of a hypothetical anti-fibrosis compound that selective reduces fibroblast production of collagen (and subsequent differentiation into myofibroblasts).
3. A surface covering with an ECM that dampens the pro-inflammatory stimulus from the open surface of the wound.

To evaluate the contribution of each of these components in the overall performance of a control strategy, combinations of the control modalities were examined for efficacy for muscle regrowth. The results of these simulation experiments are seen in Figure 18.

Figure 18: Comparison in muscle regrowth between different combinations of control strategies with 3 different levels of mediator manipulation implemented in the DRL training. In all cases the strongest level of manipulation (100-fold increase over baseline tissue mediator level) led to the greatest muscle growth.

All the treatment regimens included a DRL-trained AI control policy that governed mediator manipulation (either all 5 above noted mediators or a set of 4 minus HGF); each combination included DRL training at 3

levels of maximal mediator manipulation, 1x, 10x, or 100x over baseline tissue levels. In all cases, the 100-fold increase over baseline led to the greatest amount of muscle regrowth, demonstrating the resistance of the underlying fibrosis trajectory to alteration. These results also show the critical role of a pro-myogenesis compound (in this case HGF) in facilitating muscle regrowth. Finally, the greatest amount of muscle regrowth required the addition of an anti-fibrosis compound that directly targeted fibroblast function and/or differentiation to myofibroblasts.

3.0 Summary

The WEABM fits into the DT Paradigm as described in the NASEM Report in the follow ways:

- Personalization at Initialization: Uses CT topography to initialize computational model
- Incorporates an ongoing data linkage between the real- and digital-twin (sense/actuate capability)
- Personalization based on updatable forecasting
- Personalization in terms of adaptive control

Biological insights provided by the development and use of the WEABM include:

- Competition between fibrosis and myogenesis due to spatial constraints on available edges of intact myofibrils to initiate the myoblast differentiation process.
- Need to biologically "close" the wound from atmospheric/environmental exposure, which represents an ongoing inflammatory stimulus and "resets" the disease process into the early M1 phase, also dynamics seen with chronic non-healing wounds.
- Selective, multimodal and adaptive control needed, inherent to this is the DT concept of ongoing sensing/data link between the virtual and real world.

Though we can generally characterize the dynamic phases of wound healing in VML, a primarily pro-inflammatory response in the first 3 days, with a transition to a predominance of anti-inflammatory functions from that time to about 7-10 days, followed by an initial acceleration of collagen production and early fibrosis from approximately 7-14 days, followed by collagen remodeling and the eventual formation of fibrotic scar, we also recognize that there is considerable heterogeneity both between individuals and within an individual's wound. Because of this, the ability of the control AI to adapt its recommended multi-mediator intervention to the state of the wound is a key aspect of being able to override the biological imperative leading to fibrosis. Based on the simulation experiments and DRL-based control discovery with the WEABM we would propose the following design specifications for a device to be applied to the surface of the wound:

- Sensing Capabilities:
    - Characterization of Pro-inflammatory State: We would suggest IL6 as a sense target given its well-known role as an aggregated marker of inflammation.
    - Characterization of Anti-inflammatory and Pro-growth State: We would suggest TGFβ as a marker for the translation between beneficial anti-inflammatory conditions and potentially pro-fibrosis conditions.
- Control Capabilities:
    - Variable delivery of the following mediators (minimal set) at levels up to 100-fold increase over maximum endogenous tissue levels or near complete inhibition via direct antagonists (either chemical agents or monoclonal antibodies):
        - Pro-inflammatory mediator: TNFα
        - Pro-inflammatory byproduct: ROS
        - Early Anti-inflammatory mediator: IL10
        - Transitional Anti-inflammatory/pro-fibrosis mediator: TGFβ1
        - Pro-myogenesis mediator: HGF
        - Anti-fibroblast/myofibroblast Mediator: Hypothetical, though could use IDT1 as a small molecule
    - An ECM-like interface with the wound surface

We propose that a device with these specifications can be a first-generation device that would be refined through its use and testing. The currently listed set of target mediators are readily modifiable based on the current version of the WEABM, and an expanded WEABM could introduce additional control targets (i.e., a more detailed representation of fibroblast function and differentiation into myofibroblasts).

As with all computational models the WEABM includes selective abstractions and is inherently incomplete. We can address the latter issue via our Model Calibration method (see Methods); the former can be refined should more detailed information become available. We have also attempted to avoid off-target effects of control compounds given the dynamic and adaptive nature of the needed control policies, and therefore have focused on delivering specific mediators and their direct antagonists. We recognize that we have heavily abstracted the potential biological interactions of the ECM, but this is due to the general lack of direct mechanistic signaling information regarding its interaction with specific cell types and regarding specific pathways of interest. Our current abstraction where the ECM serves primarily as a "inflammation sink" to mitigate the open nature of clinical VMLs has been sufficient to explain all the data generated in the associated reference canine model. A significant future aspect of clinical VMLs that will have to be addressed is the impact of microbial interactions with the wound, with attendant issues regarding how to suppress microbial virulence and pathogenicity by additional manipulation of the same host response that is governing the process of healing.

We cannot emphasize enough that the current version of the WEABM is vastly under-constrained with respect to the set of non-falsifiable configurations, and as such the proposed development and refinement process of the VML DT cyberphysical system requires the application and testing of the device *in situ* to generate the necessary iterative refinement loop. We assert that solving the problem of VML will require such an approach and hope that the introduction of this VML DT can both spur device development and provide a beneficial example of the application of the Medical DT paradigm to solving a complex and significant clinical problem.

4.0 Methods:

4.1 Model Calibration:

The level of abstraction implemented in this model (and our previous work) is informed by the rationale above, specifically, that the model is a representation of existing biological knowledge (noting that this does not preclude the model from positing biological features that should be investigated experimentally). This type of knowledge is typically of the form, "*cytokine x* upregulates *cytokine y*" or "*cytokine z* is necessary for this cell to perform a specific function." This is distinct from approaches that claim to be *ab initio* or from "first principles," as these approaches are not tractable to model organ/tissue systems at the desired level of resolution, of which a primary focus is representing biological/genetic/epigenetic heterogeneity.

Variation in biomedical data arises from two primary sources: stochastic responses to identical perturbations and genetic heterogeneity. We attempt to capture both of these effects in our modeling through both the stochastic replication of simulations with identical parameterizations and through the discovery of the set of all parameterizations that, when instantiated into the knowledge/rule structure of the model, cannot be invalidated by the available data (or some other established knowledge).[50, 60] This gives us an ensemble of putative model parameterizations that fit existing data, recognizing (and potentially hoping) that future experiments may invalidate some members of the ensemble. The result is not a model that is complete (as representing every possible biological interaction is intractable), but a model that is sufficient to explain biomedical phenomena (i.e., how the system works) in the context that there exists unavoidable epistemic incompleteness/uncertainty.

We recognize that the NASEM Report placed considerable emphasis on the importance of VVUQ for DT technology; as noted above VVUQ is the focus of NASEM Report Conclusions 2-2 through 2-4 and Recommendation 2. Specifically, there is a call for new approaches towards what may constitute VVUQ for digital twins based on their specified "fit for purpose." We support this conclusion and have asserted that there are particular properties associated with attempting to computationally represent biomedical systems in a multi-scale fashion, specifically being able to cross cellular-molecular biology to whole patient/clinical

phenotypes. We identify two key facts about biological systems that distinguish them from industrial/engineered systems and present intractable challenges to attempts to translate approaches known to be effective in many physical sciences and engineered systems:

- Biological systems are faced with perpetual epistemic uncertainty.
- Biological data, particularly time series data needed to establish system trajectories, will be perpetually sparse with respect to the range of possible data configurations.

We have presented an extensive discussion regarding the implications of these two features of biological systems in Ref [49]; we present a condensed discussion of herein. We pose that the nature of the mapping between computational representations and real-world biology is a mapping between two systems with unquantifiable uncertainty. There is uncertainty (perpetual and unquantifiable) in both the specification of the computational model (perpetual epistemic uncertainty) and the probability distributions of the real-world data (data sparsity and biological heterogeneity). Traditional approaches to validation of computational models are fundamentally incapable of dealing with this type of uncertainty mapping. However, we recognize that with a goal of defining robust control one needs to avoid failure of the proposed control in as broad a context as possible. Therefore, in cases where there are perpetual uncertainties in both the model and the data it is therefore essential to maximize the conservation of information in the how these systems are related to each other, e.g., "communicate" with each other. This insight leads to casting the Validation and Uncertainty description issue for medical DTs in the context of Shannon Information (from Information Theory) and the Maximal Entropy Principle (MEP) from Statistical Mechanics. These are analogous principles where the goal is to account for the uncertainty inherent in systems and their relationship/communication to each other by maintaining the largest set of possible configurations that provide concordance between those systems; i.e., Information as per Information Theory and Entropy as per Statistical Mechanics. The maximizing entropy/information content provides the broadest non-falsifiable set of mechanistic configurations capturing biological and clinical heterogeneity and upon which control can be discovered. Since, however, any putative control policy requires identification of ostensible control points, and any mechanism-based model must have a minimal set of represented components and interactions, we recognize that the discovery of robust control must use a mechanism-based structure that, while acknowledged to be incomplete, needs to be able to be expanded upon to meet the MEP. The task, then, is to design an approach that maximizes the information content given an initial knowledge structure.

Towards this end we have developed a mathematical object called the Model Rule Matrix, or MRM [48]. The MRM is a matrix that consists of columns that list all the entities/molecules/mediators chosen to be included in a simulation model and row that list all the rules representing biological functions/interactions chosen to be in the simulation model. The numerical values of each matrix element denote the strength and direction (inhibition or augmentation) of the contribution of the entity (column) to the functional rule (row). See Figure 2 for a depiction of the components of an MRM. The lack of inclusion of a particular entity in a particular rule is represented by a "0" for the corresponding matrix element in the initial/base MRM; this represents the choices made by the modeler regarding what to include in the computational model, which in turn is informed by what data types are available to link the model to the real-world and provides a set of putative control points.

As a result, the base MRM is sparse with most of the possible interactions not explicitly represented in the computational model. However, the selection of what is represented in the model represents a bias with respect to the full biological complexity (though an unavoidable one) and that there are numerous other potential/inevitable contributions from biological components/molecules/pathways/genes that influence the behavior of the explicitly represented model components. Thus the "0" elements in the base MRM represents a representation of a "latent" space of uncharacterized and unspecified interactions that are nonetheless known to be present in some form at some degree. We assert that it is the gap between the explicitly represented structure of a computational model and the recognized additional potential interactions (by whatever degree of connectivity) that contributes to a model to capture the richness (manifested as behavioral heterogeneity) in the real biological system. It is in this fashion that the MRM utilizes the Maximal Entropy Principle: the potential information content of the model is enriched by the representation of connections that are unknown or electively omitted that become necessary to represent the full heterogeneity

of a data set. The enriched MRM is then capable of representing a genetically, epigenetically, and functionally diverse cohort of in silico patients able to represent a range of heterogeneous experimental or clinical data. The process of evolving enriched MRMs involves the application of a ML pipeline that employs both Genetic Algorithms (GAs) [61-64] and Active Learning (AL) [65-69]. to a simulation model constructed such that the coefficients of the rules in the simulation model can be considered strengths of interactions of their associated variables (and is therefore able to be represented by an MRM) and a data set that manifests a large degree of variability. The ML pipeline identifies the set of MRMs (e.g., set of possible additional connectivity configurations) that are able to encompass the range of variability present in the data set. A more detailed description of the ML pipeline can be found at Ref [48, 50].

The large degree of variability in the data is desirable because the goal is to represent the widest range of biological behavior as possible; having such a target data set expands the descriptive comprehensiveness of an ensemble of MRMs. As opposed to classical parameter fitting, which seeks to find a minimal set (or single) optimal parameter configuration(s), this process does the exact opposite by identifying a very large set of non-falsifiable configurations that has the capability of generalizing beyond the training data set and reducing the risk of brittle, over-fitted models. Time series data is particularly desirable, because the requirement of characterizing of behavioral trajectories further constrains the set of non-falsifiable MRMs (because the knowledge-based form of the rules limits their possible behaviors).

Note that this approach addresses a series of Findings and Conclusions in the NASEM Report concerning fidelity of models, multiscale modeling and model refinement. We consider all these issues related to and inherent in the establishment of novel validity and uncertainty quantification methods such as presented above.

4.2 Control Discovery of Complex Systems: Beyond classical control using Deep Reinforcement Learning

While there are numerous classical control approaches to complex dynamical systems, we have previously suggested that for complex biological systems that are represented by agent-based models, such control approaches are limited in their applicability [70]. Instead, we have suggested that Deep Reinforcement Learning (DRL), a form of training ANNs that utilizes time series/outcome data to identify combinations of actions that can alter/change the behavior/trajectories of a system [54], can be applied to simulation models to discover robust control policies to cure sepsis [55, 56]. Our proposed approach is analogous to the successful application of DRL to playing and winning games [71-73]. We term this group of applications of DRL as simulation-based DRL; this describes the use of simulation-generated training data without a pre-defined structural model for the DRL agent. We term this approach *simulation-based DRL*. A key feature of this approach is that since it relies on actions taken on a simulation model there is transparency in the mechanistic state transitions of the simulated system and this allows for a degree of interpretability in the putative control action that is not possible in "standard" data-centric ML and AI systems. This feature has potential significant impact when it comes time to evaluate such a system from a regulatory standpoint on a path to clinical implementation. The details of the process of training a DRL AI can be seen in Ref [56], and are summarized here.

We utilize a Deep Deterministic Policy Gradient (DDPG) [54] to train an AI to cure sepsis. DDPG is a reinforcement learning (RL) algorithm that uses off-policy data and the Bellman equation (Equation 1) to learn a value function, or Q-function, to determine the most valuable action to take given any state of the simulation (obtained by some means of observation, e.g. a sensor in the biomedical case).

$$Q^*(s, a) = \mathop{\mathrm{E}}_{s' \sim P} \left[r(s, a) + \gamma \max_{a'} Q^*(s', a') \right]$$

Equation 1: The Bellman Equation. Value Q is a function of the current state and action (s, a), and is equal to the reward r from the current state and chosen action (s, a) summed with the discounted value of the next state (discount factor = γ) and action (s', a') where the next state is sampled from a probability distribution (s' ~ P).

The Q-function is discovered through the execution of numerous simulations of the target system in which the AI agent uses trial and error to optimize the Q-function based on observed rewards from chosen actions.

The basis of this process is a discrete version of Q-learning [74], where the next action taken by the AI is chosen from a set of discrete actions with the best action for the current state is the one that returns the highest value from the Q-function. Q-learning is an off-policy algorithm, which means that in the training phase the AI agent is able to choose actions not chosen by the Q-function, allowing the AI agent to explore and potentially discover actions that can lead to a greater reward than an already discovered policy. Q-learning has been demonstrated to be very effective at solving complex control problems in discrete space [54].

The full observation and action space for the WEABM consists of every mediator represented in the WEABM. This was used for the initial DRL shown in Figure 10. However, in terms of practical feasibility the results of the initial control policy were examined, and a subsequent restricted observation and control space were defined consisting of TNFα, ROS, IL10, TGFβ and HGF and eventually a hypothetical anti-fibrotic agent. Each training episode included ~ 10,000 simulations of up to 14 days of simulated time (= 1400 steps) distributed over 500 processors; these episodes took ~ 30 hours to execute. Specific simulation experiments are described in the Results.

Acknowledgements:

This work was supported in part by the National Institutes of Health Award UO1EB025825. This research is sponsored by the Defense Advanced Research Projects Agency (DARPA) through Cooperative Agreement D20AC00002 awarded by the U.S. Department of the Interior (DOI), Interior Business Center. The content of the information does not necessarily reflect the position or the policy of the Government, and no official endorsement should be inferred.

Special Acknowledgements to Dr. Stephen F Badylak and Scott Johnson from the University of Pittsburgh McGowan Institute of Regenerative Medicine for their work on the experimental canine model reported in Ref [24] and which provided the calibrating data for the WEABM presented herein.

References:

1. Greising, S.M., et al., *Unwavering Pathobiology of Volumetric Muscle Loss Injury.* Scientific Reports, 2017. **7**(1): p. 13179.
2. Grogan, B.F., J.R. Hsu, and S.T.R. Consortium, *Volumetric muscle loss.* JAAOS-Journal of the American Academy of Orthopaedic Surgeons, 2011. **19**: p. S35-S37.
3. Stroncek, J.D. and W.M. Reichert, *Overview of wound healing in different tissue types.* Indwelling neural implants: strategies for contending with the in vivo environment, 2008. **1**: p. 3-41.
4. Anderson, J.M., *Biological responses to materials.* Annual review of materials research, 2001. **31**(1): p. 81-110.
5. Chargé, S.B. and M.A. Rudnicki, *Cellular and molecular regulation of muscle regeneration.* Physiological reviews, 2004. **84**(1): p. 209-238.
6. Karalaki, M., et al., *Muscle regeneration: cellular and molecular events.* In vivo, 2009. **23**(5): p. 779-796.
7. Garg, K., B.T. Corona, and T.J. Walters, *Therapeutic strategies for preventing skeletal muscle fibrosis after injury.* Frontiers in pharmacology, 2015. **6**: p. 87.
8. Martin, P. and S.J. Leibovich, *Inflammatory cells during wound repair: the good, the bad and the ugly.* Trends in cell biology, 2005. **15**(11): p. 599-607.
9. Shukaliak, J.A. and K. Dorovini-Zis, *Expression of the β-chemokines RANTES and MIP-1β by human brain microvessel endothelial cells in primary culture.* Journal of Neuropathology & Experimental Neurology, 2000. **59**(5): p. 339-352.
10. Leibovich, S. and R. Ross, *The role of the macrophage in wound repair. A study with hydrocortisone and antimacrophage serum.* The American journal of pathology, 1975. **78**(1): p. 71.
11. Witte, M.B. and A. Barbul, *General principles of wound healing.* Surgical Clinics of North America, 1997. **77**(3): p. 509-528.
12. Kharraz, Y., et al., *Macrophage plasticity and the role of inflammation in skeletal muscle repair.* Mediators of inflammation, 2013. **2013**.
13. Villalta, S.A., et al., *IFN-γ promotes muscle damage in the mdx mouse model of Duchenne muscular dystrophy by suppressing M2 macrophage activation and inhibiting muscle cell proliferation.* The Journal of Immunology, 2011. **187**(10): p. 5419-5428.
14. Tidball, J.G. and S.A. Villalta, *Regulatory interactions between muscle and the immune system during muscle regeneration.* American Journal of Physiology-Regulatory, Integrative and Comparative Physiology, 2010. **298**(5): p. R1173-R1187.
15. Bosurgi, L., A.A. Manfredi, and P. Rovere Querini, *Macrophages in injured skeletal muscle: a perpetuum mobile causing and limiting fibrosis, prompting or restricting resolution and regeneration.* Frontiers in Immunology, 2011. **2**: p. 62.
16. Mantovani, A., et al., *The chemokine system in diverse forms of macrophage activation and polarization.* Trends Immunol, 2004. **25**(12): p. 677-86.
17. Uehara, O., et al., *Upregulated expression of MMP-9 in gingival epithelial cells induced by prolonged stimulation with arecoline.* Oncology Letters, 2017. **14**(1): p. 1186-1192.
18. Kim, E.-S., M.-S. Kim, and A. Moon, *TGF-β-induced upregulation of MMP-2 and MMP-9 depends on p38 MAPK, but not ERK signaling in MCF10A human breast epithelial cells.* International journal of oncology, 2004. **25**(5): p. 1375-1382.
19. McKleroy, W., T.-H. Lee, and K. Atabai, *Always cleave up your mess: targeting collagen degradation to treat tissue fibrosis.* American Journal of Physiology-Lung Cellular and Molecular Physiology, 2013. **304**(11): p. L709-L721.
20. Mann, C.J., et al., *Aberrant repair and fibrosis development in skeletal muscle.* Skeletal muscle, 2011. **1**(1): p. 1-20.
21. An, G., C. Cockrell, and J. Day, *Therapeutics as Control: Model-Based Control Discovery for Sepsis*, in *Complex Systems and Computational Biology Approaches to Acute Inflammation*. 2021, Springer. p. 71-96.
22. Grieves, M.W., *Virtually intelligent product systems: digital and physical twins*. 2019.

23.   National Academies of Sciences, E.a.M., *Foundational Research Gaps and Future Directions for Digital Twins (2023)*. 2023, National Academies of Sciences, Engineering and Medicine: Washington, DC. p. 164.

24.   Crum, R.J., et al., *Transcriptomic, proteomic, and morphologic characterization of healing in volumetric muscle loss.* Tissue Engineering Part A, 2022. **28**(23-24): p. 941-957.

25.   An, G. and C. Cockrell, *Drug development digital twins for drug discovery, testing and repurposing: a schema for requirements and development.* Frontiers in systems biology, 2022. **2**: p. 928387.

26.   Cockrell, C., S. Christley, and G. An, *Investigation of inflammation and tissue patterning in the gut using a Spatially Explicit General-purpose Model of Enteric Tissue (SEGMEnT).* PLoS Comput Biol, 2014. **10**(3): p. e1003507.

27.   Cockrell, R.C., et al., *Towards anatomic scale agent-based modeling with a massively parallel spatially explicit general-purpose model of enteric tissue (SEGMEnT_HPC).* PLoS One, 2015. **10**(3): p. e0122192.

28.   An, G., J. Bartels, and Y. Vodovotz, *In Silico Augmentation of the Drug Development Pipeline: Examples from the study of Acute Inflammation.* Drug Dev Res, 2011. **72**(2): p. 187-200.

29.   Cockrell, C. and D. Axelrod, *Optimization of Dose Schedules for Chemotherapy of Early Colon Cancer Determined by High Performance Computer Simulations.* 2018.

30.   Baldazzi, V., F. Castiglione, and M. Bernaschi, *An enhanced agent based model of the immune system response.* Cell Immunol, 2006. **244**(2): p. 77-9.

31.   Bailey, A.M., B.C. Thorne, and S.M. Peirce, *Multi-cell agent-based simulation of the microvasculature to study the dynamics of circulating inflammatory cell trafficking.* Ann Biomed Eng, 2007. **35**(6): p. 916-36.

32.   Cockrell, C. and G. An, *Sepsis reconsidered: Identifying novel metrics for behavioral landscape characterization with a high-performance computing implementation of an agent-based model.* J Theor Biol, 2017. **430**: p. 157-168.

33.   Cockrell, R.C. and G. An, *Examining the controllability of sepsis using genetic algorithms on an agent-based model of systemic inflammation.* PLoS Comput Biol, 2018. **14**(2): p. e1005876.

34.   Braund, K., J. McGuire, and C. Lincoln, *Observations on normal skeletal muscle of mature dogs: a cytochemical, histochemical, and morphometric study.* Veterinary Pathology, 1982. **19**(6): p. 577-595.

35.   Kuzon Jr, W.M., et al., *A comparative histochemical and morphometric study of canine skeletal muscle.* Canadian Journal of Veterinary Research, 1989. **53**(2): p. 125.

36.   Volk, S.W., et al., *Diminished type III collagen promotes myofibroblast differentiation and increases scar deposition in cutaneous wound healing.* Cells Tissues Organs, 2011. **194**(1): p. 25-37.

37.   Darby, I.A., et al., *Fibroblasts and myofibroblasts in wound healing.* Clinical, cosmetic and investigational dermatology, 2014. **7**: p. 301.

38.   Chaffey, N., *Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K. and Walter, P. Molecular biology of the cell. 4th edn*. 2003, Oxford University Press.

39.   Kendall, R.T. and C.A. Feghali-Bostwick, *Fibroblasts in fibrosis: novel roles and mediators.* Frontiers in pharmacology, 2014. **5**: p. 123.

40.   Dhawan, J. and T.A. Rando, *Stem cells in postnatal myogenesis: molecular mechanisms of satellite cell quiescence, activation and replenishment.* Trends in cell biology, 2005. **15**(12): p. 666-673.

41.   Larie, D., et al., *Quantifying the size and healing of volumetric muscle wounds using 3D Slicer on CT scans in a canine model.* bioRxiv, 2021: p. 2021.07. 13.452185.

42.   Khuu, S., J.W. Fernandez, and G.G. Handsfield, *Delayed skeletal muscle repair following inflammatory damage in simulated agent-based models of muscle regeneration.* PLOS Computational Biology, 2023. **19**(4): p. e1011042.

43.   Khuu, S., J.W. Fernandez, and G.G. Handsfield, *A coupled Mechanobiological model of muscle regeneration in cerebral palsy.* Frontiers in Bioengineering and Biotechnology, 2021. **9**: p. 689714.

44.   Martin, K.S., et al., *Computational modeling of muscle regeneration and adaptation to advance muscle tissue regeneration strategies.* Cells Tissues Organs, 2016. **202**(3-4): p. 250-266.

45. Martin, K.S., et al., *In silico and in vivo experiments reveal M-CSF injections accelerate regeneration following muscle laceration.* Annals of biomedical engineering, 2017. **45**: p. 747-760.

46. Haase, M., et al., *Agent-based model demonstrates the impact of nonlinear, complex interactions between cytokines on muscle regeneration.* bioRxiv, 2023: p. 2023.08. 14.553247.

47. Westman, A.M., et al., *Agent-based model provides insight into the mechanisms behind failed regeneration following volumetric muscle loss injury.* PLoS computational biology, 2021. **17**(5): p. e1008937.

48. Cockrell, C. and G. An, *Utilizing the heterogeneity of clinical data for model refinement and rule discovery through the application of genetic algorithms to calibrate a high-dimensional agent-based model of systemic inflammation.* Frontiers in physiology, 2021. **12**: p. 662845.

49. An, G. and R.C. Cockrell, *Generating synthetic multi-dimensional molecular-mediator trajectories: Considerations.* Frontiers in Systems Biology, 2023. **3**: p. 1188009.

50. Cockrell, C., et al., *Nested active learning for efficient model contextualization and parameterization: pathway to generating simulated populations using multi-scale computational models.* Simulation, 2021. **97**(4): p. 287-296.

51. Brown, B.N. and S.F. Badylak, *Extracellular matrix as an inductive scaffold for functional tissue reconstruction.* Translational Research, 2014. **163**(4): p. 268-285.

52. Badylak, S.F., et al., *Mechanisms by which acellular biologic scaffolds promote functional skeletal muscle restoration.* Biomaterials, 2016. **103**: p. 128-136.

53. Wolf, M.T., et al., *Naturally derived and synthetic scaffolds for skeletal muscle reconstruction.* Advanced drug delivery reviews, 2015. **84**: p. 208-221.

54. Lillicrap, T.P., et al., *Continuous control with deep reinforcement learning.* arXiv preprint arXiv:1509.02971, 2015.

55. Petersen, B.K., et al., *Deep reinforcement learning and simulation as a path toward precision medicine.* Journal of Computational Biology, 2019. **26**(6): p. 597-604.

56. Cockrell, C., D. Larie, and G. An, *Preparing for the next Pandemic: Simulation-based Deep Reinforcement Learning to discover and test multimodal control of systemic inflammation using repurposed immunomodulatory agents.* Frontiers in Immunology, 2022. **13**: p. 995395.

57. Garg, K., B.T. Corona, and T.J. Walters, *Losartan administration reduces fibrosis but hinders functional recovery after volumetric muscle loss injury.* Journal of Applied Physiology, 2014. **117**(10): p. 1120-1131.

58. Larouche, J.A., et al., *Spatiotemporal mapping of immune and stem cell dysregulation after volumetric muscle loss.* JCI insight, 2023. **8**(7).

59. Corona, B.T., et al., *Pharmacological mitigation of fibrosis in a porcine model of volumetric muscle loss injury.* Tissue Engineering Part A, 2020. **26**(11-12): p. 636-646.

60. Cockrell, C. and G. An, *Using Genetic Algorithms to reproduce the heterogeneity of clinical data through model refinement and rule discovery in a high-dimensional agent-based model of systemic inflammation.* Frontiers in Physiology, 2021. **12**: p. 726.

61. Fonseca, C.M. and P.J. Fleming. *Genetic Algorithms for Multiobjective Optimization: FormulationDiscussion and Generalization*. in *Icga*. 1993. Citeseer.

62. Goldberg, D.E. and J.H. Holland, *Genetic algorithms and machine learning.* Machine learning, 1988. **3**(2): p. 95-99.

63. Haupt, R.L. and S.E. Haupt, *Practical genetic algorithms*. 2004: John Wiley & Sons.

64. Cockrell, R.C. and G. An, *Examining the controllability of sepsis using genetic algorithms on an agent-based model of systemic inflammation.* PLoS computational biology, 2018. **14**(2): p. e1005876.

65. Cohn, D.A., Z. Ghahramani, and M.I. Jordan, *Active learning with statistical models.* Journal of Artificial Intelligence Research, 1996. **4**: p. 129-145.

66. Brinker, K., *On active learning in multi-label classification*, in *From Data and Information Analysis to Knowledge Engineering*. 2006, Springer. p. 206-213.

67. Huang, S.-J., R. Jin, and Z.-H. Zhou. *Active learning by querying informative and representative examples*. in *Advances in neural information processing systems*. 2010.

68. Schein, A.I. and L.H. Ungar, *Active learning for logistic regression: an evaluation.* Machine Learning, 2007. **68**(3): p. 235-265.

69. Tsymbalov, E., M. Panov, and A. Shapeev. *Dropout-Based Active Learning for Regression*. in *International Conference on Analysis of Images, Social Networks and Texts*. 2018. Springer.

70. An, G., et al., *Optimization and control of agent-based models in biology: a perspective.* Bulletin of mathematical biology, 2017. **79**: p. 63-87.
71. Silver, D., et al., *Mastering the game of Go without human knowledge.* Nature, 2017. **550**(7676): p. 354-359.
72. Silver, D., et al., *A general reinforcement learning algorithm that masters chess, shogi, and Go through self-play.* Science, 2018. **362**(6419): p. 1140-1144.
73. Vinyals, O., et al., *Grandmaster level in StarCraft II using multi-agent reinforcement learning.* Nature, 2019. **575**(7782): p. 350-354.
74. Mnih, V., et al., *Playing atari with deep reinforcement learning.* arXiv preprint arXiv:1312.5602, 2013.

Appendix A: Facts and References for WEABM

Stages of Wound Healing

Fact	Reference
Hemostasis (seconds to hours)	
Earliest signals of tissue injury are release of molecules (e.g., ATP) and exposure of collagen on blood vessel wall	1
Inflammation (hours to days) Also called 'muscle degeneration'	
contains, neutralizes, or dilutes injury-causing agent or lesion	1,2
Initiated by release of signaling molecules during hemostasis	1,2
Vasodilation, vasopermeability increased, initiating leukocyte recruitment	1
Neutrophils are first inflammatory cells to arrive, and show up within 1-6 hrs	1,19,33,48
Neutrophils migration influenced by chemoattractants released by platelets and chemokines from endothelial surface	1,3
Leukocyte receptor PSGL1 binds to P-selectin on platelets and endothelial cells, causing neutrophils to roll and tether to endothelial calls	1,4
During the rolling, the Neutrophils activation is increased from IL8 and MIP1B released from endothelial cells, leading them to strongly attach to endothelial cells	1,5
Neutrophils then extravasate through the cell wall into the injury site, releasing enzymes to digest foreign debris and kill bacteria through secretion of Ros and phagocytosis	1,6
Neutrophils apoptose after 24-48 hrs if wound decontamination is complete	1,7,33
Neutrophils are replaced by monocytes, macrophages attracted by the continued release of MIP1a and MIP1b	1,7
Macrophages then release pro-inflammatory cytokines (e.g., IL1, TNFa, TGFb)	1,2
Macrophages remove foreign debris	1,2
Wounds can heal without neutrophils, cant heal without macrophages	1,6,8
Macrophages can activate myogenic cells	19
Macrophages are the primary inflammatory cell type ~48 hrs post injury	19
Prolonged inflammation promotes muscle degeneration and fibrosis development	24
Neutrophils quickly apoptose and are undetectable after 3-4 days	24
Neutrophils release ROS and proteases, causing secondary damage, and facilitate phagocytosis and recruitment of monocytes	24
Neutrophils populations peak between 1-3 days post injury and quickly return to normal levels	22
Repair (Days to Weeks)	

Macrophages and their associated anti-inflammatory cytokines transitions the wound from the inflammatory stage to the repair stage; this typically happens 1 week after injury — 1

Repair phase has increased cell proliferation, capillary budding, and synthesis of extracellular matrix (initially fibrogen and fibronectin) — 1,9

ECM Fibrogen, Fibronectin is replaced by proteoglycans — 1

ECM produced by fibroblasts in connective tissue, chondrocytes, osteoblasts — 1

Excess ECM production leads to the formation of fibrous scar tissue — 1

Muscle regeneration relies on muscle stem cells, which are normally quiescent — 10,11

Quiescent satellite (muscle stem) cells are activated by inflammation-associated signals and begin proliferation — 10,12

The majority of the newly-proliferated stem cells will differentiate into myocytes, however some will remain and divide as stem cells to replenish the pool of quiescent satellite cells — 10,12

Myofibers are formed over the new ECM — 10,13

Excessive ECM (scar tissue) prevents restoration of original myofiber structure — 10,14

The fibroblast is the primary ECM producer — 10,14

The interactions between satellite cells and fibroblasts is a critical determinant of the efficiency of the repair process — 10,15-17

Proangiogenic factors come into play in the later stages of repair, revascularizing the new myofibers — 10,18

Myogenic cells increase their population, and muscle fibers are formed in the same manner as in embryonic morphogenesis — 19

New muscle fibers often express embryonic/developmental forms of MHC — 19

Once the myogenic cells have fused together, the new myofibers increase their size — 19

Basic Facts

Differentiation into a muscle cell requires up-regulation of MyoD and Myf5 — 19

Proliferative myogenic cells are MyoD and/or Myf5 positive are "myoblasts" — 19

Myoblasts become terminally differentiated myocytes, which express Myogenin and MRF4 — 19

Mononucleated myocytes fuse together, resulting in multinucleated syncytium, which will mature to contracting muscle fibers — 19

Muscle fibers are individually surrounded by connective tissue and group into bundles, forming skeletal muscle — 19

Muscle satellite cells are undifferentiated, mononuclear, myogenic — 19

Pax7 is the gene that directs progenitor cells to become satellite cells — 19

"Granulation tissue is new connective tissue and microscopic blood vessels that form on the surfaces of a wound during the healing process. Granulation tissue typically grows from the base of a wound and is able to fill wounds of almost any size." — Wikipedia

There are approximately 1000 capillaries per square mm of tissue — 68

Growth Factors

HGF (Hepatocyte Growth Factor)

Key regulator of satellite cell activity in muscle regeneration	19
HGF protein levels are increased in proportion to the severity of injury in the early phase of muscle regeneration	19
In vitro, HGF increases mpc (myogenic precursor cell) proliferation and inhibits differentiation	19
Injecting HGF to injured muscle blocks the repair process and increases the number of mpc's by a factor of 3	19
HGF promotes satellite cell migration to the source of injury (through activation of Ras-Ral pathway)	19
HGF is bound to the ECM in muscle and is release in response to injury	33
HGF activates satellite cells (possibly through activating MAPK and PI3K pathways	33
HGF inhibits precursor differentiation through upregulation of Twist, which inhibits MyoD expression, and downregulation of p27	33
HGF is also released from myotubes	33
HGF promotes myofibroblast apoptosis	57
HGF upregulated by IL1, IL6, TNFa	https://pubmed.ncbi.nlm.nih.gov/8527900/
HGF is downregulated by hypoxia	https://www.ahajournals.org/doi/full/10.1161/circ.100.suppl_2.Ii-301
produced by fibroblasts	https://en.wikipedia.org/wiki/Stromal_cell
Downregulated by TGFb	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5618492/
Prevents transition to myofibroblast	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5618492/

FGF (Fibroblast Growth Factor)

FGF's activate mpc proliferation and inhibit mpc differentiation	19
FGF-6 is upregulated during muscle regeneration	19
FGF-6 deficient mice had a decrease in the number of MyoD or myogenin positive cells and an increase in collagen deposition	19
FGF 4 may be redundant for FGF6	19
FGF6 is highly released during inflammation	33
FGF inhibits satellite cell differentiation and alters adhesion	33
FGF1 inhibits TGFb in endothelial cells	60,61

TNFa, IL1b inhibit FGF signaling	61
FGF contributes to innervation	66

IGF (insulin-like growth factor)

IGF-I and IGF-II alter MRF expression and promote proliferation and differentiation/fusion of myoblasts	19,33
Increased levels of IGF-I lead to increased muscle mass, both from increased satellite cell proliferation and from hypertrophy	19,33
IGF-1 promotes cell survival through the PI3K-Akt pathway	19
Akt is also involved in myogenic differentiation	19
IGF promotes terminal differentiation by inducing myogenin expression	33
IGF-1 down-regulates inflammatory cytokines and reduced fibrosis	33
IGF2 comes later and seems to be more related to myotube formation	33
IGF contributes to innervation	67

TGFb (Transforming Growth Factor Beta)

When TGFb binds to cell receptors, it activate phosphorylation of SMAD proteins which then translocate to the nucleus and activate various genes (including collagen transcription) depending on the state of the cell	19, 48
TGFb inhibits myoblast proliferation and differentiation	19,48
Myostatin is a member of the TGFb family and inhibits muscle growth	19
MSTN maybe inhibits satellite cell proliferation (MSTN expression higher in fast-twitch muscles, which have a lower number of satellite cells)	19,33
MSTN shows high expression in necrotic fibers and when satellite cell activation is low (early in muscle regeneration); these high levels may imply that MSTN could be a chemoattractant for inflammatory cells	19
Later in the regeneration process, MSTN levels are low	19
TGFb is released in response to injury by platelets and then induces its own expression in other cells (like myogenic cells)	33
TGFb can inhibit or induce or have no effect on cell proliferation, depending on the presence/combination of PDGF, FGF, and IGF	33
TGFb is chemotactic for macrophages, leukocytes, and myocytes, and activates monocytes	33
TGFb2 regulates where/when myoblasts fuse into myotubes	33
TGFb1 promotes fibroblast differentiation	33
TGFb induces the production of PDGF	33

Excessive TGFb activity can leads to excessive ECM production and fibrosis	33
TGFb is responsible for scar	45
TGFb1 is the principal fibrogenic signal	48
TGF's fibrogenic activity through SMAD2,3,4 can be suppressed by SMAD7	48
TGFb can also act through the MAPK pathway, which also leads to synthesis of ECM; MAPK can also be activated without TGF	48
TGFb1 inhibits MMP production	48
TGFb1 promotes differentiation of myoblasts into myofibroblasts	62
TGFb promotes Akt, and is thus antiapoptotic	62
LPS induces macrophages to make TGFb, not PMNs though	https://onlinelibrary.wiley.com/doi/10.1002/jcp.1041400226

PDGF

PDGF enhances fibroblast proliferation and production of ECM	30,56
PDGF is released from macrophages and injured vessels	33
PDGF stimulates proliferation of satellite cells and inhibits their differentiation, and is also chemotactic	33

VEGF

VEGF is produced by macrophages and keratinocytes and is significant for successful neovascularization	46
VEGF has partially fibrogenic	48
VEGF secreted by fibroblasts assists in angiogenesis	56
VEGF contributes to innervation	67

Muscle Lineage and Fibroblasts

Satellite Cells

As part of the regeneration process, satellite cells are activated and begin to proliferate. After multiple proliferations, the majority of the cells then differentiate into myocytes, which fuse into myofibers	19,33
Damage at one end of a muscle fiber will activate satellite cells, causing proliferation and migration towards the site of injury	19

When satellite cells are activated and begin proliferation, they are referred to as myogenic precursor cells (MPC's	19
This activation is marked by the rapid upregulation of the MRFs Myf5 and MyoD; these signals are not observed in quiescent satellite cells	19
M-cadherin is essential for myoblasts to fuse together	19
M-cadherin expression in satellite cells is activated by muscle injury	19
In a rat model, 80% of the satellite cells provided myonuclei to muscle repair, 20% divided more slowly and replenished the pool of satellite cells.	19,24
Alternatively, when the satellite cells divide, they may divide asymmetrically, and 1 cell may go to muscle repair, and 1 to replenish the pool	19,33
Pax3 is required for migration f muscle precursors from the somite; Pax7 is required for satellite cell specification	33
Cells that express Myf5 and MyoD are called Myoblasts	33
Upregulation of secondary Myogenic Regulatory Factors (MRFs) like myogenin and MRF4 induces terminal differentiation of the myoblasts into myocytes; The myocytes begin to express Myosin Heavy Chain (MHC) and Muscle Creatine Kinase (MCK)	33
After injury, Myf5 and MyoD are the first MRF's to be expressed; myogenin is next, and after that MRF4	33,36,48
Myf 5 promotes satellite cell self-renewal	33
M-cadherin expression in satellite cells is activated by muscle injury, and is required for myoblasts to fuse into muscle fibers; M-Calpain is also required for successful reconstruction and induced by injury	33
MyoD seems to up-regulate MRF4	33
MyoD up-regulates Slug (Snai2), which has unknown function but is significantly up-regulated in the late stages of muscle regeneration	33
MNF (Myocyte Nuclear Factor) is necessary for later regeneration to function correctly; at the time of SSC activation, MNFb is down-regulated and MNFa is more abundant; this reverses as muscle regeneration proceeds	33
STAT3 is expressed by activated satellite cells; STAT3 signaling is not activated in differentiated myoblasts or myotubes; STAT3 potentially increases proliferation and prevents apoptosis; STAT3 signaling induces the regeneration of injured muscle; STAT3 is phosphorylated by a variety of cytokines (IFNg, IL6, IL5, EGF, HGF), and then translocates to the nucleus where it regulates other signaling	33
Numb is thought to be part of the asymmetric division of satellite cells; Numb down-regulates the Notch1 pathway, which promotes proliferation of the myogenic precursors; lower levels of Numb in one of the progeny cells allows for up-regulation of the Notch-1 pathway, leading to continued proliferation	33
As satellite cells proliferate, p27 levels increase until they reach a threshold; the cells then exit cell cycle, differentiate, and turn off Twist expression	33
TNFa and IL6 released my macros promote recruitment, activation, proliferation of muscle satellite cells	48
M2 macros release IL4, Il10, which promotes myoblast differentiation, fusion to myotubes; these signals also up-regulate Myf4 and MEF2	48
Skeletal muscle expresses MMP1,2,9,13	

SSC pools located on fibers

SSC's make up about 5% of the muscle mass in adults

SSC's secrete MMPs

IL6, TNFa promote recruitment, activation, proliferation of muscle satellite cells — 48

ECM

Correct remodeling of the ECM around individual fibers is critical to the repair process; excessive/persistent ECM deposition (also called fibrosis) leads to a defective outcome	24
Fibroblasts proliferate and migrate to the injury site immediately after the damage occurs	24
Fibroblasts and satellite cells work together to repair; deletion of fibroblasts leads to a lack or proliferation and premature differentiation of satellite cells	24
The basal lamina is a thin layer of nonfibrallar collagen that is in direct contact with the myofiber plasma membrane	24
ECM acts as a reservoir for growth factors, including TGFb	33
Cell migration involves a number of complex cell-ECM interactions including adhesion to ECM via integrins	33
ECM remodeling requires MMPs (metalloproteinases)	33
MMPs can activate growth factors like TGFb and FGF	33
MMPs are secreted in latent form and need to be converted to active form for proteolysis	33
uPa and Plasmin activate MMPs and can lead to ECM degradation	
MMPs can degrade ECM Components	48
MMP 2,9 are gelatinases and degrade type 4 collagen, fibronectin, proteoglycans, laminin	48

MMP1,13 degrade the 1,3 collagen	48
MMP9 is up-regulated by NfKB, MAPK, STAT3	51
IL1b, TGFb up-regulate MMP2,9	52,53
In the synthesis of ECM process, collagen type 3 (major component of granulation tissue) is replaced by collagen type 1	54
Type 1 collagen is associated with scar; normal tissue is a mix of type 1 and type 3, scar is mostly type 1	58
TNFa or IL1b induced MMP1 secretion is mediated by MEK/ERK, so a MEK/ERK inhibitor will significantly inhibit the MMP secretion when induced by pro-inflammatory signals	71

Fibroblasts

TGFb promotes fibroblast proliferation, ECM production	31,56
ECM deposition occurs faster than myogenesis; excess ECM leads to scar tissue	31,48
TGFb can prevent Fibroblast apoptosis (triggered by TNF)	32
Fibrogenesis occurs the same way, regardless of how injury occurred (toxic, trauma, etc)	33
Ablation of fibroblast population leads to premature satellite cell differentiation and depletion of the satellite cell pool, and ultimately, smaller myofibers (maybe through TCF4)	39
TGFb1 causes fibroblasts to transform into myofibroblasts, which express alpha-smooth-muscle-actin (aSMA)	48
Fibronectin and Tenascin-C are produced first, followed by Collagen type III and collagen type I	48
For injuries that have persistent inflammation (i.e., VML), myofibroblasts do not undergo apoptosis, and remain in granulation tissue; the sustained presence of immune cells in this tissue causes the release of fibrogenic cytokines like TGFb1; this causes myofibroblasts to continue proliferation and production of ECM, which ultimately leads to scar	48
Fibroblasts secrete type 1 and 3 collagen	55
Fibroblasts can produce both MMPs and their inhibitors (TIMPs)	56
The fibroblast to myofibroblast transition enables the contraction of the ECM to seal open wounds	56
Fibroblasts are stimulated to differentiate into myofibroblasts from pro-inflammatory signals	56
Fibroblasts secrete TGFb1, IL1b, assisting in the activation/migration of macrophages	56
Fibroblasts can contribute to chronic inflammation, promoting myoblast differentiation and thus, fibrosis	56
Fibroblasts secrete VEGF, which is important for angiogenesis	56
IL13 stimulates (in fibroblasts) TGFb production, collagen production, MMP expression, fibroblast proliferation, myofibroblast differentiation	56
TGFb induces ROS production in fibroblasts	69
IL1 and IFNg reduce aSMA formation (preventing differentiation)	69
When fibroblasts are exposed to media from muscle cells which are treated with SMAD3 and TGFb, they preferentially express collagen3 over collagen 1	70

IL1 prevents collagen production	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1379579/
Dexamethasone prevents collagen production	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1379579/
Collagen 1 can be produced without presence of collagen3	https://www.karger.com/Article/Pdf/322399
Fibrin fragment E is produced as part of the granulation tissue process and recruits myofibroblasts	https://www.sciencedirect.com/science/article/pii/S0898656820301388

Myofibroblasts

In granulation tissue, fibroblasts are activated and become myofibroblasts (defined by there expression of a-SM actin)	54
In a normal wound healing, myofibroblasts undergo a wave of apoptosis when the tissue integrity is restored	54
Inhibiting myofibroblast apoptosis (for example, by mechanical stress) leads to hypertrophic scarring	54
TGFb1 inhibits myofibroblast apoptosis by activating Akt	54
HGF promotes myofibroblast apoptosis	57
Myofibroblasts secrete TGFb, VEGF, CTGF, IL1,6,8	62
TGFb induces expression of CTGF, VEGF in myofibroblasts	62

Inflammatory Cells

T Lymphocytes

T Lymphocytes have two major subtypes, Th1 and Th2 which can govern muscle repair and fibrosis by generating distinct cytokine profiles	24
Th1 cells (promote cell mediated immunity) produce anti-fibrotic cytokines such as IFNg, TNFa, IL12, and IL2	24
Th2 cells (promote humoral immunity) produce profibrotic cytokines such as IL-4, IL-5, IL-6, I-10 and IL-13	24,35
Th1 cytokines inhibit the development of Th2 cytokines and vice-versa	24
T lymphocytes can skew repair towards anti or profibrotic paths, as well as maintain the cytokine milieus which govern macrophage polarization	24
T lymphocytes are a significant source of TGFb	24

Mast cells accumulate around 8 hrs post injury, and are linked to the development of fibrosis   24

Mast cells release pro-inflammatory cytos like TNFa, IL1, and IL6   24

Th1 cytokines, especially TNFa and IFNg activate macrophages to the pro-inflammatory M1 phenotype   36

TH2 cytokines activate M2 macrophages (IL4, Il10, IL13)   36

Th17 cells produce IL17, which increases satellite cell proliferation, decreases satellite cell differentiation to muscle precursor cells (MPCs) aka myoblasts; but inhibits their fusion to myoblasts, prolonging the early stages of muscle regeneration   65

TH2, TH17 cells significantly improve revascularization   65

IL17 upregulates GCSF and IL8   Wikipedia

TGFb, IL1, IL6 promote Th17 recruitment   https://www.hindawi.com/journals/iji/2012/819467/

Immune cell concentration can be 100k/mm^3 in damaged tissue   https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4454513/

IL6 promotes Th17 differentiation   https://academic.oup.com/jmcb/article/11/2/144/5054600

Macrophages

Acutely injured muscle recruits Ly6Chi inflammatory monocytes, not Ly6Clo patrolling monocytes. Ly6Chi use the CCR2 receptor   27,46

Macrophages have historically been classified into M1 (classically activated) and M2 (alternatively activated) classes   27

M1 macros are pro-inflammatory and activated by IFNg and LPS   27

M2 macrophages are anti-inflammatory, pro-regenerative, and possibly pro-fibrotic, and are activated by IL4 and IL13   27

In vivo, macrophages can display mixed m1/m2 phenotypes depending on the milieu; also, macros may not expand clonally to maintain the phenotype   27

In the early stages of muscle repair, macros are broadly activated and express pro-inflammatory (TNFa, IL1b, IL6, IL23A), anti-inflammatory (CD163, arginase 1, Ym1, IL10, fibronectin, TGFb), and pro-regenerative signals.   27,28

Over time, macrophages shift from pro to anti-inflammatory, regulated by IL10   27

Phenotype switch (M1 to M2) occurs subsequent to phagocytosis of muscle debris   28

If Anti-inflammatory signals come too early, or are blocked when they arrive later, muscle regeneration will be impeded   28

M1 macrophages stipulate MPC proliferation and inhibit fusion   28

M2 macrophages promote MPC commitment to terminal differentiation and promote myotube formation   28

In vivo depletion of intramuscular macrophages during late phase regeneration leads to a decrease in the size of new myofibers   28

IFNg promotes the M1 phenotype, suppresses M2 phenotype   23

Macros extravasate from the blood in response to signals released in tissue injury   24

M1 macrophages are recruited to the injury, and subsequently switch to the M2 phenotype   24

Pro-inflammatory M1 macrophages are recruited from exposure to Th1 cytokines IFNg and TNFa, as well as LPS or endotoxin   24

M2 macrophages have three subtypes: M2a macrophages are associated with advanced stages of tissue repair, and are induced by Th2 cytokines such as Il4 and IL13; M2b macrophages are anti-inflammatory and can release large amounts of IL-10, but they can also release pro-inflammatory cytokines IL-1B and TNFa, but not IL12; M2c macrophages are anti-inflammatory and are induced by IL-10   24,37

3 days post injury, macrophages could be split in to high-TNF and low-TNF populations, though both express the same levels of IL10 and IL1b   22

Th1 cells promote pro-inflammatory macrophage function; Th2 cells promote anti-inflammatory macrophage function   21

Macrophages activate myogenic cells   33

There can be 100k inflammatory cells per cubic millimeter of regenerating muscle tissue, the majority of which are myeloid cells which can be precursors for monocytes   36

The first set of macrophages to the wound site is phagocytic, the following wave is nonphagocytic; 1st wave peaks at 2 days, 2nd wave peaks at 4 days   36

M1 Macrophages cause tissue damage from releasing NO   36

M2a macrophages are activated by Il4 and IL13; these promote wound healing and tissue repair   36,37

M2b macrophages are activated via TLR and release Th2 anti-inflammatory cytokines   36,37

M2c macrophages are activated by IL10 and release cytokines that deactivate M1 phenotypes and promote proliferation in nonmyeloid cells	36,37
Macrophages forced into an M1 phenotype through application of IFNg showed reduced TNFa secretion and increase in TGFb secretion after phagocytosis of injured muscle cells; these expressions signal a shift to the M2 phenotype and indicate the importance of phagocytosis in making that switch	36
Phagocytosis of apoptotic neutrophils increased TGFb secretion and suppressed Il1, Il10, and TNF, while phagocytosis of lysed neutrophils increased TNFa secretion	36,45
The m1 phenotype typically produces a lot of IL12, and small amounts of IL10; the M2 phenotype produces large amounts of IL10, but small amounts of IL12; Human M1's also produces a lot o fIL23	37
M2 macrophages typically produce very low amounts of pro-inflammatory cytokines, with the exception of those exposed to IC or LPS; those macrophages still produce high levels of pro-inflammatory cytokines, but also high levels of IL10 and low levels of IL12	37
M1 polarization produces inflammatory CC chemokines (CCL5, CXCL9,10,11,16) which recruit Th1, Tc1, and NK cells	37
M2a polarization, sustained by Il4 and Il13, produces CCL 18,17,22,24, which recruits eosinophils, basophils, and Th2 cells	37
M2b polarization produces CCL1 which recruits Tregs	37
M2c polarization is driven by exposure to IL10, and produces CCL 16,18 as well as CXCL13 (when co-stimulated with IL10 and LPS) which recruits eosinophils and naive t cells	37
LPS activation of monocytes or macrophages leads to NfKB-dependent transcription of inflammatory cytokines such as CXCL1,2,3,5,8,9,10 and CCL 2,3,4,5,11,17,22; LPS and IFNg induce CXCL10, CCXCL9, and CCL5	37
Evidence suggests that signals that induce M2 polarization (IL10,4,13) down-regulate NfKB and STAT1 activities	37
Il4 and IL13 selectively induce CCL17,22,24 in M2a macros through inhibition of IFNg	37
CCL17 binds to CCR4 and IL10 inhibit CpG-mediated M1 activation	45

Inflammatory Macrophages are activated by LPS, IFNg, or GM-CSF, and they produce TNFa, IL6, IL23, IL1, IL12p40, IFNi, RNI, COI, CXCL9, CXCL10, CXCL11   45

Tissue Healing macrophages are activated by IL4, IL13, IL10 and M-CSF, and they produce IL10, IL4, TGFb, IGF1, PTX3, PGE2, MMP9, Fizz1, YM1/2, CCL17, CCL22   46

(mice) Ly-6Clo/CCR2lo/CXCR1hi monocytes come into the wound after the Ly6chi/CCR2hi/CX3CR1lo (approximately 3 days)   46

(mice) When CX3CR1 is ko'd, wounds exhibit fewer macrophages, myofibroblasts, decreased TGFb and VEGF   46

(possible) Ly6Chi cells induce apoptosis in wound neutrophils, Ly6Clo cells remove post-apoptotic debris   46

IL4 increases macrophage production of TGFb   26

M2's produce TGFb1, PDGF, FGF-2, and VEGF   48

IL6 promtes the M2 phenotype   https://academic.oup.com/jmcb/article/11/2/144/5054600

Neutrophils
Akt regulates neutrophil apoptosis; when Akt runs out, cell dies. PTEN negatively regulates Akt
Neutrophils release MPO which causes muscle membrane damage, which induced macrophages to lyse them
Neutrophils are attracted to muscle injury from DAMPS, such as HMGB1, ATP, IL1a, CXCL8, and others

https://pubmed.ncbi.nlm.nih.gov/21558983/#:~:text=Polymorphonuclear%20neutrophil%20and%20iMos%20subpopulations,whereas%20iMos%20made%20IL%2D6.

Neutrophils make IL10

Monocytes
Monocytes are activated by TGFb and secrete FGF, TNF, and IL1   https://www.pnas.org/content/103/40/14836
                                                                36,37

Eosinophils   https://link.springer.com/article/10.1007/s00441-017-2785-7
Eosinophils contribute to muscle regeneration
Eosinophls are myeloids that are activated by IL5
Eosinophils modulate regeneration by decreasing number of Cytotoxic T lymphocytes   33

Mast Cells 36

resident mast cells may be responsible for initial cytokine spike
from the injury 36

Inflammation Signals

IL6

LIF (leukemia inhibitory factor) stimulated myoblast growth without affecting myoblast
differentiation/fusion 19

LIF activates JAK2-Stat pathway 19

In vitro, LIF applied to the site of injury increases myoblast proliferation and increases
regenerated myofiber number and size 19

IL6 affects myoblasts like LIF in vitro, but does not affect muscle repair in vivo 19

NfkB induces IL6 expression 36

IL6 increases myoblast proliferation, not fusion

IL6 upregulates IL10 https://journals.physiology.org/doi/pdf/10.1152/physrev.90100.2007

IL6 downregulates Il1 and TNFa https://journals.physiology.org/doi/pdf/10.1152/physrev.90100.2007

JAK/STAT

Stat3 reduces the efficacy of PI3K to produce Akt by regulating the subunits (of PI3K) p55alpha
and p50alpha 34,35

Janus Kinase (JAK) up-regulates Stat3 35

JAK is activated by: IFN-{gamma, lambda}; IL{2,4,6,10,11,13,15,22,26,31} 35

Stat3 is activated when phosphorylated by JAK. 35

TNFa

TNFa promotes activation of NfkB in muscle cells 33

TNFa activates leukocytes 33

TNFa upregulates atrogin-1, leading to the catabolism of muscle proteins 33

Post-injury, TNFa is released by macrophages and injured muscle fibers 33

The absence of TNFa only moderately affects the muscle repair process, so it likely has several
redundancies 33

TNFa induces iNOS in myeloid cells — 36
TNFa is chemotactic for myoblasts and satellite cells — 36
TNFa induces MMP release (especially MMP9) — 58,59

IFNg
IFNg (alone or combined with microbial products like LPS or with cytokines like TNF) activates macrophages — 37
IFNg decreases satellite cell differentiation to myoblasts — 63
IFNg limits MPC proliferation — 63
IFNg inhibits IL10 — https://pubmed.ncbi.nlm.nih.gov/14607900/

LIF
LIF is regulated by Il1a, IL1b, TNF, and FGF2, as well as TGFb — 33
LIF increases the size and number of myogenic precursor cells — 33
LIF affects ECM remodeling — 33
LIF promotes neuron survival — 33
LIF does not have an affect on myotube formation — 38
Giving LIF via IV results in a 50% decrease in area occupied by connective tissue 5 days after crush injury

p38
Both TNFa and IL1 activate p38 kinase, which promotes differentiation; inhibition of p38 inhibits myocytes from fusing into myotubes and reduces expression of MEF2 and myogenin — 36
p38 increases activity of MyoD — 36
p38 can promote myogenesis by phosphorylating and increasing transcriptional activity of MEF2; p38 can inhibit myogenesis by phosphorylating to MRF4 — 36

General
CXCR3 and CCR5 are preferentially expressed on Th1 cells; CCR3, CCR4, and CCR8 are preferentially expressed on Th2 cells — 37
CCR3 ligands attracts eosinophils — 37
The TLR+IFNg dependent induction of CXCL10, CCL5, and CXCL9 is inhibited by IL4 and IL10 — 37
IL10's inhibition on LPS activated macrophages relies on STAT3-dependent mechanisms and inhibiting the NfKB pathway — 37
Inhibition of LPS-activated genes by IL4 occurs through the STAT6 pathway — 37
CCL22 is deactivated by receptor D6, a decoy receptor — 37
CCL18 is induced by IL4,10,13, and inhibited by IFNg — 37

NFkB

IL1b

Mediators

Signal	Up-regulates	Down-regulates	upregulated By	downregulated by	Produced by	Effect

Molecule			Activators	Inhibitors	Cell Types	Function
MIP1b (aka CCL4) Macrophage Inflammatory Protein 1 beta			Damage	PTEN	Monocytes, Fibroblasts, T Cells, Endothelial Cells, Epithelial Cells	Chemoattractant for monocytes, macrophages
HGF (Hepatocyte growth factor)	CCR3, CCR4	MyoD	IL1, IL6, TNFa	TGFb	fibroblasts	increases mpc proliferation and inhibits differentiation, activates satellite cells; promotes myofibroblast apoptosis
PDGF			TGFB		macrophages, injured muscle	enhances fibroblast proliferation and production of ECM, enhances satellite cell proliferation
P-Selectin					endothelial cells	Causes neutrophils to roll and tether to endothelial calls
PI3K				STAT3	neutrophils	Phosphrylates PIP2 into PIP3 which activates Akt
PIP2					neutrophils	Akt regulation
PIP3					neutrophils	Akt regulation
PTEN (Phosphorin and Tensin Homolog)				TGFb	neutrophils	Down-regulates Akt by turning Pip3 back into Pip2
AKT			PI3K	PTEN	neutrophils	controls apoptosis (when Akt runs out, apoptosis in initiated)
ROS					neutrophils, macrophages	cell damage
MPO (myeloperoxidase)					neutrophils	causes muscle cell damage and marks them for phagocytosis by monocytes
IL1b	NfkB	IkB	TLR signaling, LPS	IL6	macrophages	Promotes pro-inflammatory macrophage configuration; inhibits collagen production
IL4		IFNg, IL12			Th2, Macrophages	Promotes Th0 into Th2
IL5					Th2	Recruits Eosinophils
IL6	IL1ra, IL10	TNFa, IL1			Macrophages, Th2	promotes Th17 differentiation
IL8			NfkB	IL4, IFNg	Endothelial Cells	Chemotaxis for neutrophils

IL10	STAT3	TNFa, IL1b, IL12, IFNg	TLR, p38, NfkB, TNF, IL1, IL6	IFNg	Th2, Macrophages	
IL12	IFNg, TNFa	IL4	Th1 signals	TH2 signals	Macrophages	Promotes Th0 into Th1; blocks formation of new blood vessels through upregulation of IFNg
IL13	STAT6				Th2	activates anti-onflammatory macrophages
IL17					Th17	Limits satellite cell differentiation to muscle precursor cells; promotes angiogenesis
IL23					Macrophages	promotes/activates Th17 cells
IFNg					Th1	Promotes Th0 into TH1; promotes pro-inflammatory macro configuration; anti-vascularization
TNFa	NfkB, p38, MAPK, JNK, IL10 (in macrophages)	IkB	IL1, LPS	IL6	Th1, neutrophils, macrophages	Promotes pro-inflammatory macrophage configuration; attracts neutrophils; stimulates phagocytosis in macrophages
CXCL9	IFNg	IL4			macrophages	Promotes Th0 into TH1; promotes pro-inflammatory macro configuration; promote T cell adhesion to endothelial;
CXCL10	IFNg				macrophages, fibroblasts	promote angiogenesis; chemoattraction for macrophages, T cells
CXCL13					macrophages	recruits naive t cells, eosinophils
TGFb1		NfkB	MMP2,9, ROS		macrophages	promotes anto-inflammatory macrophage activity; promotes fibroblast to myofibroblast differentiation; promotes collagen deposition; inhibits satellite cell proliferation, differentiation
IGF1 (Insulin-like Growth Factor 1)	IkB, Myogenin				macrophages	increase satellite cell proliferation, increase myoblast proliferation, fusion to myofibers; innervation
PTX3			TNF		macrophages	
PGE2 (Prostaglandin E2)					macrophages	
MMP2	TGFb		TGFb, IL1b		macrophages	promotes angiogenesis; degrade type 4 collagen

MMP9	TGFb		NfkB, MAPK, STAT3, TGFb, IL1b		macrophages	promotes angiogenesis, neovascularization; degrade type 4 collagen
MMP1						degrade type 1,3 collagen
MMP13	TGFb					degrade type 2 collagen
CCL17					macrophages	induces chemotaxis in t cells
NfkB (nuclear factor kappa-light-chain-enhancer of activated B cells)	IL1,6, TNFa, IFNg		TNF, ROS, TGFb	IkB		inhibits myogenesis, degradation of MyoD protein; promotes Th1 differentiation
IkB		NfkB	IGF, NfkB	IL1		
FGF	Akt		inflammation		fiibroblasts	activate mpc proliferation and inhibit mpc differentiation; promotes reinnervation
M-Cadherin			Muscle damage		Muscle cells	necessary for myoblast fusion to myofibers
MyoD	MRF4, myogenin, MyoD		Activation of satellite cells		Muscle cells	Promotes differentiation to skeletal muscle
Myf5			Activation of satellite cells, SHH		Muscle cells	renewal of satellite cell pool; prevents terminal differentiation
MRF4 (also known as Myf6)					Muscle Cells	necessary for terminal differentiation
Myogenin (Myf4)						necessary for terminal differentiation
STAT3		PI3K	IFNg, IL4,6,10,13		Satellite Cells	increases proliferation, prevents apoptosis in satellite cells
TCF4	MyoD				Fibroblasts	necessary for satellite cell proliferation
VEGF					Fibroblasts	Innervation, angiogenesis

References for Appendix

Stroncek, J.D. and Reichert, W.M., 2008. Overview of wound healing in different tissue types. Indwelling neural implants: strategies for contending with the in vivo environment, pp.3-40.

Anderson J. Biological responses to materials. Annual Review Materials Research. 2001;31:81.

Yager D, Nwomeh B. The proteolytic environment of chronic wounds. Wound Repair Regeneration. 1999;7:433

Diacovo TG, Roth SJ, Buccola JM, Bainton DF, Springer TA. Neutrophil rolling, arrest, and transmigration across activated, surface-adherent platelets via sequential action of P-selectin and the beta 2-integrin CD11b/CD18. Blood. 1996;88(1):146.

Springer TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. Cell. 1994;76(2):301.

Martin P, Leibovich SJ. Inflammatory cells during wound repair: the good, the bad and the ugly. Trends in Cell Biology. 2005;15(11):599.

Shukaliak J, Dorovini-Zis K. Expression of the [beta]-chemokines RANTES and MIP-1[beta] by human brain microvessel endothelial cells in primary culture. Journal Neuropathology Experimental Neurology. 2000;59(5):339.

Leibovich SJ, Ross R. The role of the macrophage in wound repair. A study with hydrocortisone and antimacrophage serum. Americal Journal of Pathology. 1975;78(1):71.

Witte M, Abarbul A. General principles of wound healing. Surgical Clinics of North America. 1997;3:509.

Kharraz, Y., Guerra, J., Mann, C.J., Serrano, A.L. and Muñoz-Cánoves, P., 2013. Macrophage plasticity and the role of inflammation in skeletal muscle repair. Mediators of inflammation, 2013.

Mauro, A., 1961. Satellite cell of skeletal muscle fibers. The Journal of biophysical and biochemical cytology, 9(2), p.493.

Rocheteau, P., Gayraud-Morel, B., Siegl-Cachedenier, I., Blasco, M.A. and Tajbakhsh, S., 2012. A subpopulation of adult skeletal muscle stem cells retains all template DNA strands after cell division. Cell, 148(1-2), pp.112-125.

Cornelison, D.D.W., 2008. Context matters: in vivo and in vitro influences on muscle satellite cell activity. Journal of cellular biochemistry, 105(3), pp.663-669.

Mann, C.J., Perdiguero, E., Kharraz, Y., Aguilar, S., Pessina, P., Serrano, A.L. and Muñoz-Cánoves, P., 2011. Aberrant repair and fibrosis development in skeletal muscle. Skeletal muscle, 1(1), p.21.

Murphy, M.M., Lawson, J.A., Mathew, S.J., Hutcheson, D.A. and Kardon, G., 2011. Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle regeneration. Development, 138(17), pp.3625-3637.

Joe, A.W., Yi, L., Natarajan, A., Le Grand, F., So, L., Wang, J., Rudnicki, M.A. and Rossi, F.M., 2010. Muscle injury activates resident fibro/adipogenic progenitors that facilitate myogenesis. Nature cell biology, 12(2), p.153.

Uezumi, A., Fukada, S.I., Yamamoto, N., Takeda, S.I. and Tsuchida, K., 2010. Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. Nature cell biology, 12(2), p.143.

Mounier, R., Chrétien, F. and Chazaud, B., 2011. Blood vessels and the satellite cell niche. In Current topics in developmental biology (Vol. 96, pp. 121-138). Academic Press.

Charge, S.B. and Rudnicki, M.A., 2004. Cellular and molecular regulation of muscle regeneration. Physiological reviews, 84(1), pp.209-238.

Sciorati, C., E. Rigamonti, A. A. Manfredi, and P. Rovere-Querini. "Cell death, clearance and immunity in the skeletal muscle." Cell Death & Differentiation 23, no. 6 (2016): 927-937.

Rigamonti, Elena, Paola Zordan, Clara Sciorati, Patrizia Rovere-Querini, and Silvia Brunelli. "Macrophage plasticity in skeletal muscle repair." BioMed research international 2014 (2014).

Novak, Margaret L., Eileen M. Weinheimer-Haus, and Timothy J. Koh. "Macrophage activation and skeletal muscle healing following traumatic injury." The Journal of pathology 232, no. 3 (2014): 344-355.

Villalta, S. Armando, Bo Deng, Chiara Rinaldi, Michelle Wehling-Henricks, and James G. Tidball. "IFN-γ promotes muscle damage in the mdx mouse model of Duchenne muscular dystrophy by suppressing M2 macrophage activation and inhibiting muscle cell proliferation." The Journal of Immunology 187, no. 10 (2011): 5419-5428.

Kharraz, Yacine, Joana Guerra, Christopher J. Mann, Antonio L. Serrano, and Pura Muñoz-Cánoves. "Macrophage plasticity and the role of inflammation in skeletal muscle repair." Mediators of inflammation 2013 (2013).

Das, Amitava, Mithun Sinha, Soma Datta, Motaz Abas, Scott Chaffee, Chandan K. Sen, and Sashwati Roy. "Monocyte and macrophage plasticity in tissue repair and regeneration." The American journal of pathology 185, no. 10 (2015): 2596-2606.

Novak, Margaret L., and Timothy J. Koh. "Macrophage phenotypes during tissue repair." Journal of leukocyte biology 93, no. 6 (2013): 875-881.

Wang, Xingyu, Wanming Zhao, Richard M. Ransohoff, and Lan Zhou. "Infiltrating macrophages are broadly activated at the early stage to support acute skeletal muscle injury repair." Journal of neuroimmunology 317 (2018): 55-66.

Chazaud, Bénédicte. "Macrophages: supportive cells for tissue repair and regeneration." Immunobiology 219, no. 3 (2014): 172-178.

Juban, Gaëtan, and Bénédicte Chazaud. "Metabolic regulation of macrophages during tissue repair: insights from skeletal muscle regeneration." FEBS letters 591, no. 19 (2017): 3007-3021.

Werner - Regulation of Wound Healingby Growth Factors and Cytokines

Greising - Unwavering pathobiology of volumetric muscle loss injury

Nilotinib reduces muscle fibrosis in chronic muscle injury by promoting TNF-mediated apoptosis of fibro/adipogenic progenitors

Karalaki – Muscle Regeneration: Cellular and molecular events

Abell, K., A. Bilancio, R. W. Clarkson, P. G. Tiffen, A. I. Altaparmakov, T. G. Burdon, T. Asano, B. Vanhaesebroeck, and C. J. Watson. 2005. Stat3-induced apoptosis requires a molecular switch in PI(3)K subunit composition. Nat. Cell Biol. 7:392-398.

Koch, Stefan, and Asma Nusrat. "The life and death of epithelia during inflammation: lessons learned from the gut." Annual Review of Pathology: Mechanisms of Disease 7 (2012): 35-60.

Tidball – Regulatory interactions between muscle and the immune system during regeneration

Mantovani - The chemokine system in diverse formsof macrophage activation andpolarization

White - Leukaemia inhibitory factor increases myoblast replication and survival and affects extracellular matrix production: combined in vivo and in vitro studies in post-natal skeletal muscle

Murphy - Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle regeneration

Edelblum KL, Yan F, Yamaoka T, Polk DB. Regulation of apoptosis during homeostasis and disease in the intestinal epithelium. Inflamm Bowel Dis. 2006;12(5):413–424.

Ghosh, Sankar, Michael J. May, and Elizabeth B. Kopp. "NF-{kappa} B and Rel proteins: evolutionarily conserved mediators of immune responses." Science Signalling 16, no. 1 (1998): 225.

Li, Xinyan, Song Jiang, and Richard I. Tapping. "Toll-like receptor signaling in cell proliferation and survival." Cytokine 49, no. 1 (2010): 1-9.

Kretz-Remy, Carole, and André-Patrick Arrigo. "Selenium: A key element that controls NF-κB activation and IκBα half life." Biofactors 14, no. 1 (2001): 117-125.

Hohmann, H. P., R. Remy, C. Scheidereit, and A. P. Van Loon. "Maintenance of NF-kappa B activity is dependent on protein synthesis and the continuous presence of external stimuli." Molecular and cellular biology 11, no. 1 (1991): 259-266.

Borsugi – Macrophages in injured skeletal muscle: a perpetum mobile causing and limiting fibrosis

Brancato – Wound Macrophages as key regulators of repair

Corona – Pathophysiology of Volumetric Muscle Loss Injury

Garg – Therapeutic strategies for preventing skeletal muscle fibrosis after injury

Pons, Sebastian, and Ignacio Torres-Aleman. "Insulin-like growth factor-I stimulates dephosphorylation of IκB through the serine phosphatase calcineurin (protein phosphatase 2B)." Journal of Biological Chemistry 275, no. 49 (2000): 38620-38625.

Karin, Michael. "How NF-κB is activated: the role of the IκB kinase (IKK) complex." Oncogene 18, no. 49 (1999): 6867-6874.

Uehara, Osamu, Kousuke Takimoto, Tetsuro Morikawa, Fumiya Harada, Rie Takai, Bhoj Raj Adhikari, Ryoko Itatsu et al. "Upregulated expression of MMP-9 in gingival epithelial cells induced by prolonged stimulation with arecoline." Oncology letters 14, no. 1 (2017): 1186-1192.

Kim, Eun-Sook, Mi-Sung Kim, and Aree Moon. "TGF-β-induced upregulation of MMP-2 and MMP-9 depends on p38 MAPK, but not ERK signaling in MCF10A human breast epithelial cells." International journal of oncology 25, no. 5 (2004): 1375-1382.

Huang, Qiaojia, Fenghua Lan, Xiaoting Wang, Yinghao Yu, Xuenong Ouyang, Feng Zheng, Junyong Han et al. "IL-1β-induced activation of p38 promotes metastasis in gastric adenocarcinoma via upregulation of AP-1/c-fos, MMP2 and MMP9." Molecular cancer 13, no. 1 (2014): 18.

Darby - Fibroblasts and myofibroblasts in wound healing

Molecular Biology of the Cell 4th ed – Alberts, Johnson, Lewis

Kendall – Fibroblasts in Fibrosis: Novel Roles and Mediators

68   Regirer, S. A., N. S. Utushkina, and N. Kh Shadrina. "Blood flow in the capillary network of muscle tissue." Fluid Dynamics 21, no. 6 (1986): 911-919.

69   Hinz, Boris. "Formation and function of the myofibroblast during tissue repair." Journal of Investigative Dermatology 127, no. 3 (2007): 526-537.

70   Goel, Shakti A., Lian-Wang Guo, Xu-Dong Shi, Rishi Kundi, Gregory Sovinski, Stephen Seedial, Bo Liu, and K. Craig Kent. "Preferential secretion of collagen type 3 versus type 1 from adventitial fibroblasts stimulated by TGF-β/Smad3-treated medial smooth muscle cells." Cellular signalling 25, no. 4 (2013): 955-960.

71   Tasaki, Kazuhito, Yutaka Shintani, Takao Saotome, Akira Andoh, Yoshihide Fujiyama, Shigenari Hozawa, and Tadao Bamba. "Pro-inflammatory cytokine-induced matrix metalloproteinase-1 (MMP-1) secretion in human pancreatic periacinar myofibroblasts." Pancreatology 3, no. 5 (2003): 414-421.