Pathogenicity of Commensal Gut Biofilm In Prefrail Aging

Guillaume Le Cosquer[1,2], Melissa Pannier[1], Elodie Meunier[1], Julie Thevenin[1], Elise

Pyhourquet[1], Sophie Guyonnet[3,*], Bruno Vellas[3,*], Yohan Santin[4,*], Bruno Guiard[5,*], Angelo

Parini[4,*], Louis Buscail[2], Barbara Bournet[2], Damien Guillemet[6], Celine Deraison[1], Nathalie

Vergnolle[1,7,#,*], Jean-Paul Motta[1,#,*,✉]   *for the IHU HealthAge INSPIRE/Open*

Science study group

Affiliations:

[1]Institute of Digestive Health Research (IRSD), INSERM, Toulouse University, INRAe, ENVT,

University Toulouse III Paul Sabatier (UPS), 31300 Toulouse, France

[2]Department of Gastroenterology and Pancreatology, Toulouse University Hospital (CHU

Toulouse), Toulouse, France and Toulouse University, University Toulouse III Paul Sabatier

(UPS), 31059 Toulouse, France

[3] Institute of Aging, Toulouse University Hospital (CHU Toulouse), Toulouse, France and

CERPOP, Toulouse University, INSERM, University Toulouse III Paul Sabatier (UPS), 31300

Toulouse, France

[4] Institute of Metabolic and Cardiovascular Diseases (I2MC), INSERM, Toulouse University,

University Toulouse III Paul Sabatier (UPS), 31432, Toulouse, France

[5] Research Center on Animal Cognition (CRCA), Center of Integrative Biology (CBI),

Université de Toulouse, CNRS UMR-5169, Univ Toulouse III - Paul Sabatier (UPS), Toulouse,

France.

[6] Nexira, 129 Chemin de Croisset, 76000 Rouen, France

[7] Department of Physiology and Pharmacology, University of Calgary Cumming School of

Medicine, 3330 Hospital Drive, NW Calgary, AB, T2N4N1, Canada

* members of the IHU HealthAge INSPIRE/Open Science study group (full list in

supplementary material)

[#] Co-supervision

✉Correspondence to: jean-paul.motta@inserm.fr

Keywords: gut microbiota; aging; microb-aging; gut biofilm; prefrailty; geroscience

Main text – ≈ 6.700 words or ≈38.000 characters, excluding Title Page, Tables, Figures, and

References

Abstract – 149 words

Display items – 6 Figures + 1 Table

Running title: Pathogenicity of prefrail gut biofilms

Abstract. Pathophysiological mechanisms of unhealthy aging, particularly the transition

from robustness to frailty, remain poorly understood. Despite extensive microbiome research

on taxonomy, the behavior of early prefrail gut bacteria in their natural community-host

mucosal tissue context remains unexplored. Using fecal samples from the INSPIRE-T aging

human cohort, we characterized gut microbiota phenotype during prefrailty stages using a

polymicrobial biofilm model. Results revealed that prefrail-derived biofilms exhibited distinct

taxonomic and physical alterations, enhanced dispersal, and increased epithelial virulence

compared to robust counterparts. Multiparametric analyses linked biofilm characteristics to

clinical traits, suggesting their potential as aging status indicators. Polyphenol-rich grape

pomace extract partially reversed prefrail biofilm alterations and reduced proinflammatory

prefrail biofilm responses *in vitro*. Microbiota from prefrail aged mice induced colon damage

in antibiotic-treated recipients, establishing a prefrail microbiome-inflammation causality.

Overall, the findings identified novel prefrail microbiome characteristics, established causal

inflammatory links, and support microbiota-targeted geroprotective interventions for the

prefrail populations.

1. INTRODUCTION

By 2030, the World Health Organization (WHO) projects that one in six individuals worldwide will be aged 60 years or older [1]. This shift to older populations coincides with an increase in age-associated diseases and unhealthy aging. One of the first steps in the development of unhealthy aging is linked to the appearance of frailty syndrome, a clinical condition defined as decreased physiological reserve and function that heightens vulnerability and impairs the ability to withstand health stresses[2]. The absence of frailty (robust individuals) has been identified as a reliable predictor of prolonged longevity without age-associated diseases [3]. The field of geroscience, which has recently emerged, is dedicated to understanding the underlying mechanisms of aging, finding methods to extend the healthspan – rather than lifespan – and to delay or impede frailty syndrome and age-associated diseases [4]. The scientific community has intensely focused on describing fecal microbiome composition and changes in the aging population[5-8]. From these studies, it has become clear that chronological age is linked to alterations within the gut microbiome, a process coined as 'microb-aging' by Bosco and Noti[6]. A current challenge is to identify additional aspects of microb-aging, beyond taxonomy, that could be related to early signs of unhealthy aging (in prefrail individuals)[9]. The involvement of microbiota behavior and virulence in such differences has not yet been investigated. Although clinical scales allow geroclinicians to categorize patients into three categories according to their frailty status (robust, prefrail, and frail individuals), biological data on the relationship between these progressive clinical changes and microb-aging dynamics are scarce. From a clinical perspective, prefrail individuals represent a potentially reversible stage before a significant decline in health. A report on community-dwelling elders indicated a notable prefrailty-to-robust transition, with a 16% conversion rate observed over an 18-to 36-month follow-up

period, compared to 0% during the same period for the frailty-to-robust transition [10]. Defining

conditions or treatments that would favor this transition from prefrailty-to-robust is among

the most important challenge of geroscience[10].

The motivation of this study was thus to focus on the aging population exhibiting prefrail

characteristics, in the search for interventions seeking at prefrailty-to-robust transition. For

that, we capitalized on the French INSPIRE-T cohorts at Toulouse University Hospital, two

clinically well-characterized human and mouse aging cohorts in which various non-invasive

samples were collected, including fecal samples. These cohorts were designed to identify

novel markers of aging, age-related diseases and health evolution that could be

therapeutically targetable[11-13]. Considering its key importance on gut health and disease, we

aimed at focusing on microbiota communities living on the surface of the mucosa, as a

polymicrobial anaerobic biofilm[14]. To that end, we used a static model of polymicrobial

biofilms grown on mucin-coated polystyrene microplates (MBEC device model[15-17]), providing

a straightforward, mid-throughput system for parallel community analysis. Our underlying

hypothesis was that these cultured biofilms, while partially preserving the fecal taxonomy,

and could retain key structural and phenotypic alterations associated with both aging and

donor health status. Ultimately, our goal was to establish a proof-of-concept platform for

geroprotective experimental strategies aimed at addressing these abnormalities.

2.   METHODS

Human clinical data and samples

This study used human feces collected from the INSPIRE-T cohort in 2021 (ID

NCT04224038, Table 1 and Supplementary Table 1)[12]. Donors were categorized into groups

of 13 adults (7 men/6 women; 30-70 years) and 29 elderly subjects (+ 70 years). The last group

was divided into 14 prefrail (6 men/8 women) and 15 healthy or robust individuals (7 men/8

women). Frailty syndrome was evaluated using the Fried frailty index (Supplementary Table

2 for detailed Fried index calculation[2]). Donors were excluded if they had received antibiotics,

probiotics, or prebiotics within the past three months and if they had history of any of the

following conditions: inflammatory bowel disease, irritable bowel syndrome, colorectal

cancer, immunosuppression, severe liver disease. Feces were collected at the initial visit,

aliquoted (≈1 g), and stored at −80°C until analysis in the INSPIRE-T biobank at a central

laboratory (Toulouse Bio Ressources CRB, CHU Toulouse/IFB Purpan, Toulouse, France).

Clinical characteristics, including age, gender, frailty status, and household status, were

documented (Supplementary Table 1: lifestyle). For the present analysis, we used the version

1.0 of the INSPIRE-T database. We calculated the nutritional status of each individual using

the Mini Nutritional Assessment (MNA; Supplementary Tables 3 and 4[18]) and calculated the

Charlson Comorbidity Index based on their past medical history[19](Supplementary Tables 5,

S6, and S7). Full list of members of the IHU HealthAge INSPIRE/Open Science study group is

included in Supplementary Material.

Mouse functional data and samples

This study involved colon tissue and feces collected from the INSPIRE mice cohort (CREFRE

– US006 / Inserm – University of Toulouse, Paul Sabatier). The cohort comprised male and

female SWISS outbred mice[13]. Animals underwent health assessments to measure frailty,

using a rodent equivalent to the human Fried frailty index, the Valencia Score[13,20]

(Supplementary Table 8). Behavioral phenotyping was conducted by the Mouse Behavioral

Core platform (Integrative Biology Center, CBI, Toulouse), at the CREFRE facility. In this study,

we collected samples from young (6 months, n=8), aged robust (24 months, n=8), and aged

prefrail animals (24 months, n=8). Distal colon was fixed in methacarn solution (6/3/1 of

methanol/chloroform/acetic acid), and embedded in paraffin. Feces were collected and

stored at −80°C. Both paraffin blocks and frozen feces were maintained in the INSPIRE mouse

biobank at a central laboratory (Centre de Ressources Biologiques-Cancer - Institut

Universitaire du Cancer Toulouse – Oncopole, CHU Toulouse, Toulouse, France).

Culture of biofilms from human feces

Human fecal samples (10-20 mg) were resuspended in supplemented Brain Heart Infusion

(BHI; Sigma-Aldrich 53286, St. Quentin Fallavier, France), enriched with hemin solution (final

5 µg/ml; Sigma-Aldrich H9039), Vitamin K1 (final 50µg/ml; Sigma-Aldrich V3501), and L-

Cysteine (0.5 mg/ml; Sigma-Aldrich W326305). Human microbiota biofilms was grown in

mucin-coated polystyrene pegs of MBEC Biofilm plates (Innovotech, Edmonton, AB, Canada)

by incubating the lids in a solution of 10mg/ml gastric porcine mucin (Sigma-Aldrich M1778,

a crude porcine mucin type III preparation, bound with sialic acid 0.5-1.5%) overnight at room

temperature without agitation (based on methods first described in [21]). After mucin-coating,

the lids were place onto another 96-well plate containing BHI supplemented and dilution of

fecal microbiota (OD 600nm =0.1; Varioskan® Flash, Thermo Scientific Massachusetts, MA,

USA). Biofilms were then matured for 72 hours in a humidified incubator set at 37 °C, under

anaerobic condition and continuous rocking (Rocker25 from Labnet set at 50 RPM) as

described in[15-17]. After maturation, the lid was gently rinsed and basal BHI plate was refreshed

to collect naturally dispersed bacteria from biofilms under the same conditions for an

additional 24 hours. Biofilm biomass, viability, and matrix content were characterized using

the methods described below. Additionally, the basal plate medium containing biofilm-

dispersed bacteria were collected and was used for 16S rRNA V3-V4 gene sequencing, to

estimate dispersal rates, and to perform interaction assays with human intestinal epithelial

cells. For all microbiological procedures described herein, original sample handling and

incubations were carried out in an anaerobic chamber (Don Whitley A20, 0.1% oxygen; 4.9%

CO2; 95.0% N2, 45-65% humidity) or in hermetic jars with an anaerobic sachet (Anaeropack;

Thermo Fisher).

Staining assays on MBEC biofilm plates

Total biofilm biomass was evaluated using Gram's safranin-O solution (Sigma-Aldrich),

biofilm matrix-associated glycoproteins content using fluorescein wheat germ agglutinin

(WGA, 1/100 diluted, ThermoFisher), and viability using a resazurin assay (1 mg/ml; Sigma-

Aldrich)[15-17]. While WGA has specific affinity for sialic acid (SA) and N-acetylglucosamine

(NAG), we used it to assess the presence of matrix-associated glycoproteins containing these

sugar moieties (SA/NAG-glycoproteins). The dispersal rate was estimated at 24h by measuring

the ratio of viable biofilm-dispersed bacteria to total biofilm biomass. Briefly, MBEC were

transferred to 96 well plate containing stains and incubated at 37°C for 2 h (resazurin and

WGA) or 20' (safranin-O). Stains were recovered by sonication at 45 kHz (USC-500T, VWR) in

a plate containing acetic acid (30%, safranin-O) or 1.5M sodium chloride (WGA). Specific

absorbance (safranin-O) or fluorescence (WGA and resazurin) was measured using an EnSight

spectrophotometer (Perkin Elmer, Villepinte, France). An average of at least 12 biofilm

replicates were performed for each individual. A positive control was included for each

experiment (*E. Coli* LF82, a high biofilm producer).

Donor fecal microbiota preparation

Feces obtained from SWISS aged mice were divided into three groups of interest: young

(n=8 mice, 4 males and 4 females, 6 months old), aged robust (n=8 mice, 5 males and 3

females, 24 months old), and aged prefrail (n=8 mice, 5 males and 3 females, 24 months

old[13]). Feces were pooled in 1.5 mL phosphate buffered saline (PBS, Eurobio Scientific, Les

Ulis, France), homogenized, and filtered through a sterile nylon mesh strainer. The filtered

contents were centrifuged twice, and the pellet was resuspended in PBS/20% glycerol at

OD_{600nm}=1.5 ($\approx 10^9$ CFU/ml). All the procedures were performed under anaerobic conditions

at 37°C in an anaerobic chamber (Don Whitley A20).

Post-antibiotics fecal microbiota inoculation

C57BL/6 mice (Janvier, Le Genest Saint-Isle, France), aged 8 weeks, were housed under

specific pathogen-free conditions (CREFRE U006, CHU Purpan). Mice were divided into four

groups: 6M (gavaged with pooled 6-month-old microbiota, n=9), 24M robust (gavaged with

pooled 24-month-old microbiota, n=8), 24M prefrail (gavaged with pooled 24-month-old

microbiota, n=7), and a control group receiving PBS vehicle without microbiota (n=5). Animals

were housed in large cages (for rats) (maximum to 10 mice per cage) and acclimatized for 2

weeks. To control for coprophagy and potential cage effects, every two days cage bedding –

including cotton nests – was thoroughly mixed every two days by a single experimenter using

the same gloves throughout the procedure. After acclimatization phase, mice were

distributed across four cages (PBS vehicle, 6M, 24M-robust and 24M-prefrail). The existing

microbiota was perturbed by broad-spectrum antibiotics in drinking water at concentrations

of 1 g/l (ampicillin and metronidazole, Sigma-Aldrich) and 0.5 g/l (neomycin and vancomycin,

Sigma-Aldrich) for the first five days, and half-diluted for the remaining days. Given this

standardized mixing and group distribution, we collected fecal pellets from five randomly

selected animals during the acclimatization period and after antibiotic exposure. On day 10,

the mice underwent fecal microbiota inoculation via oral gavage with 200 µL of previously

described fecal slurry. Four days post-inoculation, the colon was harvested for evaluation of

macroscopic damage, including colon thickness edema (in mm) and blind Wallace scoring

endpoints[22], was assessed by a blinded and skilled experimenter. Animal procedures were

approved by the National Laboratory Animal Ethics Committee (APAFIS 2023060610225611,

French Ministry of Research).

16S rRNA V3/V4 gene sequencing

Total DNA was extracted from feces or biofilms using the EZNA isolation kit stool DNA

(VWR for biofilms) according to the manufacturer's protocol. DNA extraction and specific PCR

amplification of 16S bacterial rDNA V3-V4 regions were targeted by 341f-806R primers and

sequenced on the Illumina MiSeq 2 × 250 bp technology at Vaiomer (Toulouse, France) or

Biomnigene (Besancon, France). Demultiplexed raw read sequences were processed using

FROGS bioinformatics pipeline (FROGS v. 4.1.0[23], Galaxy Toulouse server, GenoToul). Adaptor

and primer sequences were trimmed, and amplicons were merged using Vsearch plugin

(v.2.17.0). Sequences were processed using swarm v2.0 for cluster filtering and denoising to

identify amplicon sequence variants (ASVs), with a 0.005% abundance threshold. The ASV

table was filtered using three criteria: minimum count of 4, 10% prevalence, and 10%

variance. No rarefaction, scaling, or transformation was applied to the data. Taxonomic

assignments were performed using the 16S SILVA 138.1 database.

16S Fluorescent In situ hybridization (FISH)

For fluorescence in situ hybridization (FISH), methacarn-fixed mouse colon tissues were

embedded in paraffin. Slides were hybridized with a universal bacterial 16S fluorescent rRNA

probe (EUB338-Cy3, 5'GCTGCCTCCCGTAGGAGT-Cy5, Eurofins Scientific, France) and DNA was

counterstained with 4',6-diamidino-2-phenylindole (DAPI, Invitrogen) and polysaccharide

content (wheat germ agglutinin labeled with fluorescein, ThermoFisher)[16,17]. The biofilm

damage score was blindly evaluated in at least three microscopic fields for each mouse in the

groups of interest, according to the criteria described in[17]. Briefly, the score was established

on five criteria: mucus layer invasion, bacterial-epithelial distance, biofilm density, bacterial

translocation, and cell presence in the lumen. Each parameter was scored 0-3, with higher

scores indicating greater mucosal biofilm disruption. Samples that were unsatisfactorily fixed

for scoring were excluded from analysis. Images were acquired using a Leica LSM 710 confocal

microscope (TRI Genotoul, Purpan), and FIJI freeware was used for the final image mounting

(v.1.51, Mac Os).

Human intestinal epithelial cells

Caco-2 (ATCC, Manassas, USA) and HT29-MTX E12 (ATCC) were grown separately in

Dulbecco's modified Eagle medium (DMEM, Eurobio Scientific, Les Ulis, France)

supplemented with 10% fetal bovine serum (FBS, Sigma-Aldrich), penicillin/streptomycin

(Invitrogen, Massy, France) and non-essential amino acid solution (Thermofisher, Paisley, UK)

at 37°C in a humidified incubator with 5% CO2 as previously described [24]. Coculture at 3:1

(Caco-2:HT29) was done on flat bottom 24-wells plates at an initial total density of 2.10^5 cell

per well (Corning, Thermo Scientific). Cells were screened for mycoplasma infection

(MycoAlertTM, Lonza, Bâle, Switzerland) and treated before coculture if necessary

(PlasmocureTM, Lonza). Cells were cocultured with dispersed bacteria from biofilms

(normalized optical density; OD_{600nm}=0.01) in OPTIMEM (Thermo Fisher). Bacterial adhesion

to the intestinal epithelium was assessed after 4 h of coculture on a blood agar plate[16,17].

Relative mRNA expression by qRT-PCR

Total RNA was extracted using TRIzol reagent (Euromedex, France) after 4 h of coculture

(Qiagen, Montreal, Canada). Reverse transcription of isolated RNA was performed using the

Maxima First Strand Kit (Invitrogen, Thermo Scientific) according to the manufacturer's

instructions. Quantitative PCR (RT²qPCR Primer, Qiagen) was performed on a LightCycler

(Roche, Basel, Switzerland) based on SYBR Green technology, using the Takyon Low Rox SYBR

kit (Eurogentec, Seraing, Belgium). Data were analyzed by the $2^{-\Delta\Delta CT}$ method using HPRT and

GAPDH (Qiagen) as control genes and expressed as fold change compared to control cells

without bacteria. PCR primers used in this study are listed in Supplementary Table 9.

Grape pomace intervention

After maturation, MBEC Assay® was exposed to grape pomace solution diluted in

microbiological BHI supplemented broth at several concentrations (0.3 to 30 mg/mL) for 24h.

The medium was refreshed, and 4h later dispersed bacteria were collected to collect biofilm-

dispersed bacteria for interaction assays with human intestinal epithelial cells. Staining assays

were performed in parallel to estimate the effect of grape pomace on biofilm biomass and

matrix-associated glycoproteins content. The industrial grape pomace used (VinOgrape™,

Nexira, France) was an extract containing purified polyphenols (>90% content, quantified by

spectrophotometry method) anthocyanins, catechin, epicatechin and oligomeric

proanthocyanidins.

Statistics and visualization

We performed statistical analyses and data visualization using R 4.3.0 – Posit (Rstudio IDE)

v2023.0.9.1 built 494 (Posit Team 2023, Boston, MA, USA) and GraphPad Prism (v10 for

MacOs, La Jolla, USA). Baseline clinical characteristics were compared between the groups

with chi-squared test for categorical variables and ANOVA (Fisher test) for continuous

variables with Tukey's method to correct P-values. To explore the bacterial taxonomy

variability among sample groups, alpha and beta diversity metrics were performed in R using

phyloseq (v.1.46.0). Principal coordinate analysis (PCoA) and Partial least squares-

discriminant analysis (PLS-DA) and principal component analysis (PCA) were performed in R

using phyloseq (v. 1.46.0), mixomics (v. 6.26) and compared using PERMANOVA

(pairwiseAdonis v.0.4.1) with 9999 permutations. To calculate microbiome uniqueness—a

novel metric introduced in the context of age-related microbiome alterations[25]—we averaged

the lowest beta diversity values for each individual across multiple distance metrics, including Bray-Curtis, Jaccard, unweighted UniFrac, and weighted UniFrac, compared to all other individuals within the same experimental group. This metric quantifies how distinct each individual's microbiome is from its nearest neighbor within the cohort, with higher values indicating greater compositional distinctness. Rarefaction curves (Supplementary Figure 1A) and taxa bar plots were generated in R using phyloseq (v. 1.46.0) and ggnested (v. 0.1.0). Statistical differences between microbial communities were assessed in R using LEfSe (lefser v.1.12[26]), Deseq2 (v.1.42[27]), EdgeR (v.4.0.3[28]), metagenomeSeq (v.1.43.0[29], and ANCOM[30]. Boxplots, barplots and scatter plots were generated in R using ggplot2 (v.3.4.4) and ggpubr (v.0.6.0). Heatmaps were created in R using complexheatmap (v.2.18.0). Grout's test was used to filter statistical outliers in the qPCR dataset (GraphPad Prism). Statistical significance was set at P < 0.05. For group comparisons, we used Dunnett's or Tukey's method to correct P-values for ANOVA and the Benjamini-Hochberg method for differential abundant taxa analyses. Unless otherwise indicated, values are presented as the mean ± standard error of the mean.

3.   RESULTS

Characterization of the human cohort

Samples were collected from 42 individuals, stratified into 13 healthy individuals aged 30-70 years and 29 individuals aged ≥ 70 years. Within the aged group, 15 were robust based on the Fried frailty index[2], while 14 were prefrail (score of 1 and 2) (Table 1 and Supplementary Table 1). All the donors had satisfactory nutritional scores (Supplementary Table 3). As expected, the Charlson comorbidity index [19] indicated more comorbidities in aged individuals than in younger donors (0.923±1.19 versus 4.73±1.87, P<0.001), but this index was not

discriminant between prefrail and robust donors (5.07±1.59 versus 4.73±1.87, p=0.836;

Supplementary Table 6). General characteristics of donors used in the study are summarized

in Table 1.

Taxonomic characterization of microb-aging on human biofilms

To explore links between microb-aging and prefrailty syndrome, we cultured the fecal

microbiota of aged individuals on a mucin-coated device, in an attempt to reproduce a

mucosal polymicrobial anaerobic biofilm[14,16,17]. Despite a decline in Bacteroidota (formerly

Bacteroidetes) compared to fecal sample, biofilm communities remained dominated by

Bacillota (formerly Firmicutes), specifically *Ruminococcaceae*, *Christensenellaceae*,

Bifidobacteriaceae, *Akkermansiaceae*, *Lachnospiraceae*, and *Oscillospiraceae* (Figure 1A,

Supplementary Figure 1). These taxa are core members typically found in human fecal

microbiota (Supplementary Figure 2). The *ex vivo* biofilm model did not show statistically

significant differences in α-diversity compared to 30-70 years individuals (Supplementary

Figure 1C). Regarding β-diversity on weighted Unifrance distance, the groups were not

statistically separated (PERMANOVA >0.05, Supplementary Figure 1D). However, we

observed that biofilms from aged robust donors exhibited greater individual microbiome

uniqueness compared to those from prefrail donors (Figure 1B). We ran various statistical

models in metagenomics to identify differentially abundant taxa (i.e., Deseq2,

metagenomeSeq, ANCOM, and edgeR, Supplementary Tables 10 and 11). We found

correspondence in age-related taxa abundance between the biofilm model and other reports

on human aging cohorts[5-8] (Supplementary Figures 1E-F). Specifically, aged individuals

presented an enrichment of taxa including *Selemonadaceae*, *Prevotellaceae*, and potential

pathobionts such as *Enterobacteriaceae*. Concomitantly, there was a reduction in the

abundance of important commensals within aged biofilms, including *Dialister sp.,*

Akkermansia sp. and *Bifidobacteriaceae* (Supplementary Figure 1E). Biofilms from aged

robust individuals displayed a higher abundance of *Prevotellaceae* compared to prefrail

individuals, as well as an increased representation of *Selenomonadaceae (Megamonas sp.)*

and *Ruminococcaceae* (Supplementary Figure 1F).

We then expanded the investigation between clinical characteristics of donors and taxa

abundance in the biofilm model. We identified 34 statistically significant correlation pairs with

biofilm taxonomic features and clinical characteristics, primarily at the Family and Genus

levels (Figure 1C). First, aging alone was linked to a diminished abundance of core taxa such

as *Slackia sp.*, *Eggerthellaceae*, and Bifidobacteriales. In addition, we found a positive

correlation between prefrailty and the relative abundance of *Monogobales, Oscillibacter sp.*

and *Negativibacillus sp* (Figure 1C). Regarding clinical history, the Charlson comorbidity index

was negatively correlated with abundance of *Slackia sp.*, and *Veillonellaceae (Dialister sp.)*.

Abundances of *Pasteurellaceae* and *Haemophilus sp.* were inversely correlated with the MNA

score, shedding light on the potential links between these specific taxa and nutritional status

(Figure 1C). Gender exhibited no statistically significant correlation with the abundance of

microbial taxa in cultured biofilms (Figure 1C). Finally, regarding the living conditions of

donors, we uncovered a positive correlation between conjugal lifestyle and the microbial

abundance of several taxa, including *Tannerellaceae*, *Parabacteroides sp.*, and *Butyricimonas*

sp. (Figure 1C). Altogether, these data indicate that several microbial signatures associated

with aging and prefrailty are preserved in this biofilm model.

Physical structure of human biofilms is altered in prefrail donors

To explore microb-aging beyond taxonomy, we first characterized the physical structure

of biofilms. We observed a negative correlation between donor age and total biofilm biomass

(Pearson R = -0.6, P<0.05, Figure 2A). However, this feature was not significantly different

between prefrail and robust biofilms (Figure 2B). Regarding the extracellular matrix of biofilms, we found a positive correlation (R=0.39, P<0.05) between age and proportion of SA/NAG-glycoproteins within this matrix (Figure 2D). Interestingly, biofilms from prefrail donors exhibited a heightened glycoproteins matrix composition compared to the other samples (Figure 2E). We then investigated the links between the clinical characteristics of human donors and biofilm physical structure (Kendall's correlation approach, Figure 2C). In addition to confirming the previously described Fried and age relationship, we found a negative correlation between biofilm biomass and the Charlson comorbidity index (Figure 2C). Neither the donor's sex, household, nor diet (MNAscore) demonstrated a significant association with the biofilm structure (Figure 2C). Focusing on biofilm taxonomy (Figure 2F), we identified that high biomass biofilm, a feature of 30-70 yrs biofilms, was positively correlated with a high abundance of *Eggerthellaceae*, *Oscillospira sp.*, *Enterorhabdus sp.*, and *Slackia sp.* High biofilm matrix- SA/NAG-glycoproteins, a feature of prefrail biofilms, was correlated with donors harboring low abundance of Coriobacteriales and *Eggerthellaceae* (Figure 2F). Overall, these observations suggest that the physical structure of biofilms generated from 30-70 and aged individuals differs and, more importantly, that these physical modifications may serve as discriminant features between aged individuals predisposed to prefrailty or robustness.

Prefrailty-associated human biofilms have an hypervirulent phenotype

In addition to the physical structure, we evaluated the pathogenic behavior of human biofilms between young and aged donors. First, we assessed the stability of biofilms, measured as the rate of bacterial dispersal over a 24-hour period. High dispersal is considered a virulent trait that increases the likelihood of exposing the subjacent host epithelium to unwanted bacterial contact [15,16]. Donor age was not statistically correlated with this feature

(P=0.058, Figure 3A). However, prefrail biofilms had a significantly higher dispersal rate than

biofilms from younger groups (P<0.05, Figure 3B). We then explored the ability of biofilms to

disperse bacteria that could attach to the underlying intestinal epithelium. To this end, we

exposed enterocyte/mucus-producing human epithelial cell lines to a normalized

concentration of the dispersed biofilm bacteria. We found that higher chronological age

correlated with this behavior (P<0.05, Figure 3D). More importantly, it was dispersed bacteria

from prefrail biofilms that exhibited heightened adhesiveness compared to other groups

(P<0.01, Figure 3E). Intriguingly, the adhesiveness of bacteria was correlated with individuals

having a high Charlson comorbidity index, suggesting a potential connection between

bacterial behavior and clinical history (Figure 3C). In this cohort, neither the donor's gender,

household, nor diet (MNA score) demonstrated a statistically significant correlation's index

with biofilm pathogenic behavior (Figure 3C). Regarding taxonomy, biofilm dispersal was

statistically correlated with a lower abundance of *Ruminococcaceae* and *Oscillospirales sp*. In

addition, we identified several taxa whose abundance was either positively (e.g.,

Adlercreutzia sp. and Coriobacteriales) or negatively (e.g., *Akkermansiaceae*) correlated with

biofilm bacterial adhesiveness (Figure 3F). In summary, our findings underscore that while

chronological age plays a limited role in influencing biofilm behaviors, prefrailty status

emerges as a key factor shaping the instability and virulence of gut biofilms.

Prefrailty-associated biofilms trigger host innate response in human intestinal

epithelium

We then focused on the relative expression of 10 genes encoding proinflammatory

cytokines, antimicrobials, and tight junction proteins to discern patterns indicative of the host

epithelial innate response to these biofilm-dispersed bacteria. The scaled expression

heatmap revealed not only altered expression patterns in biofilms from aged versus younger

individuals, but also striking disparities within biofilms from prefrail versus robust individuals

(Figure 4A). Proinflammatory genes, such as *IL6* and *TNF*, and the tight junction genes claudin-

2 (*CLDN2*) and occludin (*OCLN*) (Figure 4B) exhibited a notable trend towards increased

expression in prefrail versus robust individuals. While such trends failed to reach statistical

significance when considered alone – except in the case of *TNF* and *IL6* – the PCA ordination

plot demonstrated a clear shift in the transcriptome of epithelia exposed to bacteria from

prefrail individuals compared to younger individuals (Figure 4C, PERMANOVA P= 0.007). As

visualized in the biplot vectors, genes with the greatest contribution in the separation were

those coding for proinflammatory signals (*TNF, IL8 and IL6* along the PC1 axis) and those

related to tight junction proteins (*CLDN2* and *OCLN* in the PC2 axis). Expanding this analysis

with biofilm features, we discovered that high dispersal rate – indicative of a virulent prefrail

phenotype – was correlated with a higher expression of inflammatory genes (*IL6* and *TNF*,

Figure 4D). Interestingly, high adhesiveness was also positively correlated with inflammatory

genes (*COX2*) as well as genes encoding tight junction proteins (*CLDN2* and *OCLDN*; Figure

4C). Finally, we found that an increased quantity of SA/NAG-glycoproteins in the biofilm

matrix – a feature linked to aging and prefrailty – was positively correlated with a higher

expression of proinflammatory genes (*IL6, IL8,* and *TNF*). Overall, these findings indicate that

in addition to aging host immunity, microbiota biofilms from +70 years prefrail individuals,

but not in 30-70 years and +70 years robust individuals, contain commensals with

proinflammatory triggering potential.

Prefrailty aging in mice is linked to altered mucosal biofilm biogeography in colon.

Motivated by human biofilm model findings, we aimed to confirm whether early signs of

prefrailty could influence the spatial organization of the gut microbiota in the colon. For this,

we used tissues from an aging mouse cohort, equivalent to humans aged 30 to 70+ years (= 6

to 24 months old, INSPIRE mouse cohort[13]). Among the 15 selected animals, thorough frailty

syndrome assessments were conducted, categorizing them into groups mirroring healthy

aging (robust, n=8) and those displaying early signs of frailty (prefrail, n=7, Valencia Frailty

Score, Supplementary Table 8[20]). Upon sacrifice, the colon-associated microbiota was

visualized using 16S fluorescent *in situ* hybridization (FISH) staining (Figures 5 A-C).

Remarkably, young mice exhibited a mucosal microbiota organized as a dense biofilm

community lining a layer of sterile mucus (outlined by dashed lines in yellow, Figure 5A). In

contrast, aged animals displayed a different spatial organization, featuring isolated bacteria

within the mucus layer that escaped from the dense mucosal biofilm (arrows, Figures 5B and

5C). Bacterial translocation, which was almost absent in the young cohort, was observed in

the aged animals (Figures 5B and 5C). Notably, aged prefrail animals exhibited more signs of

tissue-associated biofilm alterations than aged robust animals, including signs of virulent

behavior in gut microbiota such as invasion of the mucus sterile layer, epithelial adhesion, or

translocation (biofilm damage score[17], Figure 5D). These findings confirmed *in vivo* that

prefrail status in aging is linked to altered biofilm organization and emergence of virulent

phenotypes.

Prefrailty-associated microbiota directly causes colon damage when inoculated into

young mice

To experimentally demonstrate a causal relationship between prefrailty-associated

microbiota and local gut inflammation, we orally inoculated fecal microbiota in antibiotic-

treated young mice. We used as inoculate a pool of feces from young (6 months, n=8), robust

(24 months, n=8), and prefrail mice (24 months, n=8). As expected, the antibiotic regimen

caused taxonomic dysbiosis in gut microbiota, with severe reduction of total load, alpha

diversity, and an increased abundance of *Enterobacteriaceae*, which was partially restored

after 4 days of antibiotic washout (Figure 5E, Supplementary Figures 3-5). Post-inoculation, animals were monitored daily for 4 days, and no animals reached a distress endpoint (Supplementary Figure 6A). Mice inoculated with prefrail microbiota exhibited a significant difference in beta-diversity compared to those receiving aged robust or young microbiota (Supplementary Figure 4D). Furthermore, we observed a higher abundance of Proteobacteria, and a reduced abundance of *Muribaculaceae*, *Christensenellaceae*, *Parabacteroides* sp., and *Akkermansia* sp., in mice inoculated with prefrail microbiota compared to the other groups, mirroring previous findings in the litterature[31] (Supplementary Figure 5). Regarding intestinal tissue damage after the inoculation procedure, we found that young fecal microbiota did not cause macroscopic harm in the colon (total damage score of 1.7±0.44, Figure 5F-G, Supplementary Figure 6B). When inoculated with aged prefrail microbiota, recipient animals had significantly higher colon damage such as increased incidence of erythema, mucus release in the lumen, edema, and (total score 5.4±0.98, ANOVA followed by Dunnett's test, P = 0.002, Figure 5G). Additional histological analysis on these mice using FISH staining showed reduced bacteria-epithelium distance in prefrail mice compared to young mice (but not compared to aged-robusts), indicating a potentially compromised colonic barrier integrity in the prefrail group (Supplementary Figure 6C). These results combined with previous *in vitro* assays on intestinal epithelial cell lines, suggest that microbiota from prefrail – but not from young and robust – may directly causes intestinal damage when inoculated into recipient young mice.

Grape pomace partially reverses prefrailty-associated biofilm structure and virulence

Considering the virulent phenotype of the prefrail microbiota and its associated damage to the intestinal mucosa, we investigated whether this phenotype can be reversed by microbiota-targeted strategies. Grape-derived polyphenols have demonstrated prebiotic

properties via their activities on gut microbes[32-35]. Indeed, the majority of edible grape

polyphenols are poorly bioavailable and reach the colonic compartment where they shape

microbiome profile[36]. Therefore, we sought to determine whether industrial grape pomace,

which is already marketed as a food supplement in France, can affect the structure and

virulence of mucosal biofilms associated with prefrail aging. We previously observed that

prefrail biofilms had reduced biomass compared to young biofilms, and exposure to grape

pomace significantly increased it (Figure 6A, P<0.05 at 30 mg/ml). Importantly, we found a

statistically significant reduction of 79.5% in the dispersal rate in prefrail biofilms (Figure 6B,

P<0.05 at 30 mg/ml). However, grape pomace had no significant effect on the increased

adhesiveness of dispersed bacteria from prefrail biofilms (Figure 6C). At a concentration of 30

mg/ml, biofilm exposure to grape pomace led to a significant reduction of dispersed bacteria

able to trigger the expression of *IL6* (55% reduction, P<0.05, Figure 6D), while having no

statistically significant impact on levels of *IL8* and *TNF* (P>0.05, Figure 6E-F). Overall, these

experiments using a biofilm model suggested that grape pomace could directly reduce the

hypervirulent phenotype associated with the prefrail aging microbiota.

4.   DISCUSSION

Aging is commonly associated with changes in the gut microbiota, including shifts in

taxonomy and metabolism, a phenomenon referred to as microb-aging [6]. Despite extensive

research on fecal microbiome from various populations worldwide (reviewed in[5-8]), significant

knowledge gaps remain on the ecological implications of microb-aging within commensal gut

communities. Another under-explored area is the causal relationship between microb-aging

and low-grade inflammation of aging [9,37]. Additionally, the role of microbiota in frailty aging

and the potential of microbiota-targeted therapies in transitioning from prefrailty to

robustness remain open questions. Addressing these gaps is crucial for understanding the

nature of microb-aging and identifying non-chronological markers of unhealthy aging [4,11].

Using non-invasive samples in a clinically well characterized human aging cohort, we thus

aimed at generating a model for microbiota forming biofilm, which is the microbial lifestyle

that resides in close proximity to the host tissue and epithelium [14]. In this study, we aimed at

detecting microbiome shifts in biofilms from aged individuals at an early stage of frailty,

including but not limited to taxonomy, specifically to uncover potential therapeutic

interventions that can reverse these changes and establish new targets that promote the

transition from prefrailty to robustness [10].

From a taxonomic perspective, the cultured biofilm maintained important microb-aging

features already described in the literature, such as increased microbiome uniqueness in

robust but not in prefrail aged individuals (Figure 1B)[25]. By comparing biofilm taxonomy

results with a recent grouping proposed by Ghosh et al.[5] — Groups I, II, and III—we observed

an interesting parallelism, where gut pathobionts, such as Proteobacteria from the

Enterobacteriaceae family, exhibited increased abundance in aged individuals. We also noted

a decline of beneficial taxa in aged biofilms, such as *Bifidobacteriaceae*. Interestingly,

Akkermansia sp., a genus containing important commensals involved in host/mucus

homeostasis[38], showed slightly reduced abundance in aged biofilms. Intriguingly, biofilms

from aged robust individuals displayed increased taxa traditionally associated with cohorts of

healthy centenarians such as *Christencellaneaceae* [39-41]. As a limitation of these taxonomy

findings relates to small sample size, further investigation within the entire INSPIRE cohort

could be engaged to confirm and refine the identified taxonomic features (+1000

volunteers[12]). In addition, the longitudinal design of this cohort will enable collection from

the same individuals over the next 10 years, enabling us to track the progression of prefrailty

in individuals and determine whether they experience a shift to frailty, provided that microb-

aging is not therapeutically managed (with diet, exercise or pre- and probiotics).

In addition to taxonomic alterations, this study revealed new microbiological features that

may help distinguish between healthy and unhealthy aging in humans. Specifically, we

identified a reduced biofilm-forming capacity in most fecal samples of aged individuals

compared to that in younger individuals (Figure 2). This may indicate a common biological

pattern in the aging microbiome that is potentially linked to the loss or gain of specific taxa,

as previously discussed. More compelling are those features that distinguish between aged-

matched robust and prefrail donors, such as the amount of sialic acid/N-acetylglucosamine--

rich glycoproteins associated with the biofilm matrix (Figure 2). Extracellular polysaccharides

are indeed fundamental for retaining water within the biofilm matrix, generating a highly

hydrated and cohesive environment that influences cell movement [42,43]. Whether the altered

glycoproteins composition of the aged prefrail biofilms used in the present study contributes

to taxonomic changes, increased dispersal, and fragmented biofilm physical structure *in vitro*

(Figure 3) and *in vivo* (Figure 5) remains to be experimentally elucidated. Another discriminant

feature of prefrailty-associated biofilms is their capacity to release bacteria with higher

adhesiveness and which triggers proinflammatory signals from the underlying epithelium

(Figure 4). Although this phenomenon has not been previously documented, metagenomic

studies of fecal microbiota have indicated an increased proportion of virulence gene

pathways in microbiota from aged frail individuals [44].

It is yet to be fully understood whether local gut microbiota alterations directly contribute

to adverse outcomes in aged frail populations. Complementing *in vitro* findings, an *in vivo*

microbiota inoculation approach reinforces the hypothesis that prefrailty associated

microbes could directly cause a sub-inflammatory phenotype in the gut. Although this aligns

with other studies of aged microbiota transfer [45], the novel message from the present study

lies in the separation between robustness and prefrailty-associated sampling. Here, we found

that macroscopic colon damage induced by aged microbiota may be more pronounced with

prefrail microbiota than with robust (Figure 5G). Undoubtedly, the interaction between the

gut microbiome and its host is bidirectional and more complex links are expected *in vivo*. A

road to be explored is the impact of a prefrailty microbiota within an aging host, where

heightened inflammatory stress and immunosenescence may compromise the host's ability

to mount an effective response to local insults (the inflammaging hypothesis [37]). In addition

to the virulent behavior of prefrail biofilm, local inflammation may significantly alter microbial

biotope making it less favorable to commensal gut bacteria, or even promoting the blooming

of pathogenic taxa (e.g., *Enterobacteriaceae* [46]). This could be attributed to compromised

epithelial secretory functions in high age, such as defective mucus and antimicrobial peptide

production[47], or deficiencies in intestinal barrier integrity [48]– although the latter phenomenon

remains a subject of debate [49]. Natural polyphenols, including those present in grape pomace,

have been shown to improve cognitive function, extend the lifespan, and protect against age-

related muscle loss in several mice models [32-35]. Studies in mice have shown that grape

pomace in the diet can improve metabolic diseases by preventing gut microbiota dysbiosis

and increasing the abundance of beneficial taxa [50-52]. In this study, we found that an industrial

grape pomace preparation favored the restoration of the physical and non-pathogenic

features of prefrail biofilms *in vitro*, potentially indicating the improved resilience of microb-

aging communities (Figure 6). Additionally, this compound reduced the release of

proinflammatory triggering bacteria from prefrail biofilms. Previous study suggested that

grape pomace did not cause a significant taxonomic shift in human gut microbiota [53].

Therefore, further research is needed to determine the precise impact of grape polyphenols

on microbial virulence functions and to elucidate specific microbial genetic responses. Finally,

going beyond this particular compound, the methodology presented here may support the

clinical relevance of gut microbiota biofilm cultures (using non-invasive fecal samples) to

assess and validate the effects of new classes of geroprotectors, such as prebiotics or

probiotics, specifically targeting signs of unhealthy microb-aging.

Beyond the well-known taxonomic changes in the aging gut microbiome, this study points

to novel functional and behavioral changes in human gut biofilm, that switched to a virulent

phenotype in prefrail individuals but not in robusts. Importantly, we found that this

phenotype was reversible by polyphenol-rich grape pomace treatment. Altogether, these

results further establish the intestinal microbiota as a target to improve healthy aging. These

discoveries might represent a unique opportunity for geroclinicians to develop screening

tools and identify new therapeutic targets against microb-aging, to prevent frailty

development in the expanding aging population.

Author Contributions: GLC. performed most experiments, analyzed data, and provided clinical input; MP., EM., JT., EP., CD., NV. and JPM. contributed to *in vivo* and *in vitro* data acquisition and analysis; SG., and BV. provided human samples, contributed to clinical characterization, and reviewed manuscript; YS., BG., and A.P provided mice samples, contributed to clinical characterization and reviewed manuscript; LB., BB. reviewed manuscript and provided clinical input; DG. provided scientific reagents and reviewed the manuscript; JPM. performed 16S bioinformatics and statistics; GLC., NV. and JPM., wrote the manuscript; N.V and JPM. supervised the work and obtained fundings. All authors have read and agreed the final version of the manuscript.

Data Availability: 16S rRNA gene sequencing from the experimental biofilms and mouse data were uploaded to the NCBI Sequence Read Archive (SRA Bioproject PRJNA1076354). Additional biological or clinical metadata from the INSPIRE-T cohort can be obtained by contacting the corresponding author, who will submit the request to the INSPIRE biobanking and research committee for final approval. Other information from this study can be obtained, upon reasonable requests, from the corresponding author.

Ethics: The INSPIRE cohort protocol ID NCT04224038 was approved by the French Ethical Committee located in Rennes (CPP Ouest V) in October 2019. Mice experimentation protocol has been approved by local ethic committee of Toulouse University and by the Ministry of Superior Education and Research under the agreement APAFIS2023060610225611.

Acknowledgments: Authors thank the Department of Gastroenterology and Pancreatology at the University Hospital of Toulouse for an MD/PhD scholarship leave to GLC. Authors give credit to Dr. Felipe Sierra for its initial assessment of the research project, under the umbrella of INSPIRE Call for Geroscience Research Grant. Authors thank the INSPIRE biobanking committee for evaluating our project and approving sample collection. Authors

thank the staff of the human and mice Centre de Recherche Biologique (CRB) of the Toulouse

Hospital for samples preparation. Authors thank the CREFFRE core facility (UMS 006) for

hosting animals used in the project and contributing to histopathology analyses (R. Balouzat

and S. Milia), and the TRI-Genotoul platform for cellular imaging facilities (U1043, S. Allart).

Authors are thankful to Dr. Racaud-Sultan for critically reading the manuscript. Fundings came

from INSPIRE-Grant AMIGO / ANR dysBIOFILM ANR-22-CE14-0041-01 (JPM), ANR-PARCURE

PRCE-CE18, 2020 (NV), and the national program "Microbiote" from INSERM (NV and JPM).

The INSPIRE-T study was supported by grants from the Region Occitanie/Pyrénées-

Méditerranée (Reference number: 1901175), the European Regional Development Fund

(ERDF) (Project number: MP0022856), and the Inspire Chairs of Excellence funded by:

Alzheimer Prevention in Occitania and Catalonia (APOC), EDENIS, KORIAN, Pfizer, Pierre-

Fabre. The IHU HealthAge Open Science initiative was supported by the French National

Research Agency (ANR) as part of the France 2030 program (ANR-23-IAHU-0011), and builds

on the work conducted in the Data Sharing Alzheimer project.

Competing Interests: Prebiotic product (grape pomace) was gratuitously furnished by the

NEXIRA company (Rouen, France) and covered by a collaborative research grant between DG.

and NV. The company acknowledges having no influence on the results obtained during

assays presented in this manuscript.

References

World Health, O. *Decade of healthy ageing: baseline report*. (World Health Organization, 2020).

Fried, L. P. *et al.* Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* **56**, M146-156 (2001). https://doi.org/10.1093/gerona/56.3.m146

Romero-Ortuno, R. & Kenny, R. A. The frailty index in Europeans: association with age and mortality. *Age Ageing* **41**, 684-689 (2012). https://doi.org/10.1093/ageing/afs051

Sierra, F. The Emergence of Geroscience as an Interdisciplinary Approach to the Enhancement of Health Span and Life Span. *Cold Spring Harb Perspect Med* **6**, a025163 (2016). https://doi.org/10.1101/cshperspect.a025163

Ghosh, T. S., Shanahan, F. & O'Toole, P. W. The gut microbiome as a modulator of healthy ageing. *Nat Rev Gastroenterol Hepatol* **19**, 565-584 (2022). https://doi.org/10.1038/s41575-022-00605-x

Bosco, N. & Noti, M. The aging gut microbiome and its impact on host immunity. *Genes Immun* **22**, 289-303 (2021). https://doi.org/10.1038/s41435-021-00126-8

DeJong, E. N., Surette, M. G. & Bowdish, D. M. E. The Gut Microbiota and Unhealthy Aging: Disentangling Cause from Consequence. *Cell Host Microbe* **28**, 180-189 (2020). https://doi.org/10.1016/j.chom.2020.07.013

O'Toole, P. W. & Jeffery, I. B. Gut microbiota and aging. *Science* **350**, 1214-1215 (2015). https://doi.org/10.1126/science.aac8469

Le Cosquer, G., Vergnolle, N. & Motta, J. P. Gut microb-aging and its relevance to frailty aging. *Microbes Infect* **26**, 105309 (2024). https://doi.org/10.1016/j.micinf.2024.105309

Gill, T. M., Gahbauer, E. A., Allore, H. G. & Han, L. Transitions between frailty states among community-living older persons. *Arch Intern Med* **166**, 418-423 (2006). https://doi.org/10.1001/archinte.166.4.418

Ader, I. *et al.* Healthy Aging Biomarkers: The INSPIRE's Contribution. *J Frailty Aging* **10**, 313-319 (2021). https://doi.org/10.14283/jfa.2021.15

de Souto Barreto, P. *et al.* The INSPIRE Research Initiative: A Program for GeroScience and Healthy Aging Research Going from Animal Models to Humans and the Healthcare System. *J Frailty Aging* **10**, 86-93 (2021). https://doi.org/10.14283/jfa.2020.18

13 Santin, Y. *et al.* Towards a Large-Scale Assessment of the Relationship between Biological

and Chronological Aging: The INSPIRE Mouse Cohort. *J Frailty Aging* **10**, 121-131 (2021).

https://doi.org/10.14283/jfa.2020.43

14 Motta, J. P., Wallace, J. L., Buret, A. G., Deraison, C. & Vergnolle, N. Gastrointestinal

biofilms in health and disease. *Nat Rev Gastroenterol Hepatol* **18**, 314-334 (2021).

https://doi.org/10.1038/s41575-020-00397-y

15 Motta, J. P. *et al.* Hydrogen sulfide protects from colitis and restores intestinal microbiota

biofilm and mucus production. *Inflamm Bowel Dis* **21**, 1006-1017 (2015).

https://doi.org/10.1097/MIB.0000000000000345

16 Motta, J. P. *et al.* Iron Sequestration in Microbiota Biofilms As A Novel Strategy for Treating

Inflammatory Bowel Disease. *Inflamm Bowel Dis* **24**, 1493-1502 (2018).

https://doi.org/10.1093/ibd/izy116

17 Motta, J. P. *et al.* Active thrombin produced by the intestinal epithelium controls mucosal

biofilms. *Nat Commun* **10**, 3224 (2019). https://doi.org/10.1038/s41467-019-11140-w

18 Rubenstein, L. Z., Harker, J. O., Salva, A., Guigoz, Y. & Vellas, B. Screening for undernutrition

in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J*

Gerontol A Biol Sci Med Sci **56**, M366-372 (2001). https://doi.org/10.1093/gerona/56.6.m366

19 Charlson, M. E., Pompei, P., Ales, K. L. & MacKenzie, C. R. A new method of classifying

prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* **40**,

373-383 (1987). https://doi.org/10.1016/0021-9681(87)90171-8

20 Gomez-Cabrera, M. C. *et al.* A New Frailty Score for Experimental Animals Based on the

Clinical Phenotype: Inactivity as a Model of Frailty. *J Gerontol A Biol Sci Med Sci* **72**, 885-891

(2017). https://doi.org/10.1093/gerona/glw337

21 Dague, E. *et al.* Probing in vitro interactions between Lactococcus lactis and mucins using

AFM. *Langmuir* **26**, 11010-11017 (2010). https://doi.org/10.1021/la101862n

22 Wallace, J. L., Le, T., Carter, L., Appleyard, C. B. & Beck, P. L. Hapten-induced chronic colitis

in the rat: alternatives to trinitrobenzene sulfonic acid. *J Pharmacol Toxicol Methods* **33**, 237-

239 (1995). https://doi.org/10.1016/1056-8719(95)00001-x

23 Bernard, M., Rue, O., Mariadassou, M. & Pascal, G. FROGS: a powerful tool to analyse the

diversity of fungi with special management of internal transcribed spacers. *Brief Bioinform* **22**

(2021). https://doi.org/10.1093/bib/bbab318

24 Rolland-Fourcade, C. *et al.* Epithelial expression and function of trypsin-3 in irritable bowel

syndrome. *Gut* **66**, 1767-1778 (2017). https://doi.org/10.1136/gutjnl-2016-312094

25 Wilmanski, T. *et al.* Gut microbiome pattern reflects healthy ageing and predicts survival

in humans. *Nat Metab* **3**, 274-286 (2021). https://doi.org/10.1038/s42255-021-00348-0

26 Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biol* **12**, R60

(2011). https://doi.org/10.1186/gb-2011-12-6-r60

27 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org/10.1186/s13059-

014-0550-8

28 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140

(2010). https://doi.org/10.1093/bioinformatics/btp616

29 Paulson, J. N., Stine, O. C., Bravo, H. C. & Pop, M. Differential abundance analysis for

microbial marker-gene surveys. *Nat Methods* **10**, 1200-1202 (2013).

https://doi.org/10.1038/nmeth.2658

30 Mandal, S. *et al.* Analysis of composition of microbiomes: a novel method for studying

microbial composition. *Microb Ecol Health Dis* **26**, 27663 (2015).

https://doi.org/10.3402/mehd.v26.27663

31 Langille, M. G. *et al.* Microbial shifts in the aging mouse gut. *Microbiome* **2**, 50 (2014).

https://doi.org/10.1186/s40168-014-0050-9

32 Ziegler, T., Tsiountsioura, M., Meixner-Goetz, L., Cvirn, G. & Lamprecht, M. Polyphenols'

Impact on Selected Biomarkers of Brain Aging in Healthy Middle-Aged and Elderly Subjects: A

Review of Clinical Trials. *Nutrients* **15** (2023). https://doi.org/10.3390/nu15173770

33 Luo, J., Si, H., Jia, Z. & Liu, D. Dietary Anti-Aging Polyphenols and Potential Mechanisms.

Antioxidants (Basel) **10** (2021). https://doi.org/10.3390/antiox10020283

34 Xu, Q. *et al.* The flavonoid procyanidin C1 has senotherapeutic activity and increases

lifespan in mice. *Nat Metab* **3**, 1706-1726 (2021). https://doi.org/10.1038/s42255-021-

00491-8

35 Davinelli, S. & Scapagnini, G. Interactions between dietary polyphenols and aging gut

microbiota: A review. *Biofactors* **48**, 274-284 (2022). https://doi.org/10.1002/biof.1785

36 Rodriguez-Lopez, P. *et al.* Grape and Grape-Based Product Polyphenols: A Systematic

Review of Health Properties, Bioavailability, and Gut Microbiota Interactions. *Horticulturae* **8**,

583 (2022).

37 Franceschi, C., Garagnani, P., Parini, P., Giuliani, C. & Santoro, A. Inflammaging: a new

immune-metabolic viewpoint for age-related diseases. *Nat Rev Endocrinol* **14**, 576-590

(2018). https://doi.org/10.1038/s41574-018-0059-4

38 Cani, P. D., Depommier, C., Derrien, M., Everard, A. & de Vos, W. M. Author Correction:

Akkermansia muciniphila: paradigm for next-generation beneficial microorganisms. *Nat Rev*

Gastroenterol Hepatol **19**, 682 (2022). https://doi.org/10.1038/s41575-022-00650-6

39 Biagi, E. *et al.* Gut Microbiota and Extreme Longevity. *Curr Biol* **26**, 1480-1485 (2016).

https://doi.org/10.1016/j.cub.2016.04.016

40 Kong, F. *et al.* Gut microbiota signatures of longevity. *Curr Biol* **26**, R832-R833 (2016).

https://doi.org/10.1016/j.cub.2016.08.015

41 Biagi, E. *et al.* Through ageing, and beyond: gut microbiota and inflammatory status in

seniors and centenarians. *PLoS ONE* **5**, e10667 (2010).

https://doi.org/10.1371/journal.pone.0010667

42 Karygianni, L., Ren, Z., Koo, H. & Thurnheer, T. Biofilm Matrixome: Extracellular

Components in Structured Microbial Communities. *Trends Microbiol* **28**, 668-681 (2020).

https://doi.org/10.1016/j.tim.2020.03.016

43 Flemming, H. C. & Wingender, J. The biofilm matrix. *Nat Rev Microbiol* **8**, 623-633 (2010).

https://doi.org/10.1038/nrmicro2415

44 Maffei, V. J. *et al.* Biological Aging and the Human Gut Microbiota. *J Gerontol A Biol Sci Med*

Sci **72**, 1474-1482 (2017). https://doi.org/10.1093/gerona/glx042

45 Parker, A. *et al.* Fecal microbiota transfer between young and aged mice reverses hallmarks

of the aging gut, eye, and brain. *Microbiome* **10**, 68 (2022). https://doi.org/10.1186/s40168-

022-01243-w

46 Winter, S. E. *et al.* Host-derived nitrate boosts growth of E. coli in the inflamed gut. *Science*

339, 708-711 (2013). https://doi.org/10.1126/science.1232467

47 Sovran, B. *et al.* Age-associated Impairment of the Mucus Barrier Function is Associated

with Profound Changes in Microbiota and Immunity. *Sci Rep* **9**, 1437 (2019).

https://doi.org/10.1038/s41598-018-35228-3

48 Man, A. L. *et al.* Age-associated modifications of intestinal permeability and innate

immunity in human small intestine. *Clin Sci (Lond)* **129**, 515-527 (2015).

https://doi.org/10.1042/CS20150046

49 Wilms, E. *et al.* Intestinal barrier function is maintained with aging - a comprehensive study

in healthy subjects and irritable bowel syndrome patients. *Sci Rep* **10**, 475 (2020).

https://doi.org/10.1038/s41598-019-57106-2

50 Mezhibovsky, E. *et al.* Grape Polyphenols May Prevent High-Fat Diet-Induced Dampening

of the Hypothalamic-Pituitary-Adrenal Axis in Male Mice. *J Endocr Soc* **7**, bvad095 (2023).

https://doi.org/10.1210/jendso/bvad095

51 Roopchand, D. E. *et al.* Dietary Polyphenols Promote Growth of the Gut Bacterium

Akkermansia muciniphila and Attenuate High-Fat Diet-Induced Metabolic Syndrome.

Diabetes **64**, 2847-2858 (2015). https://doi.org/10.2337/db14-1916

52 Chacar, S. *et al.* The Impact of Long-Term Intake of Phenolic Compounds-Rich Grape

Pomace on Rat Gut Microbiota. *J Food Sci* **83**, 246-251 (2018). https://doi.org/10.1111/1750-

3841.14006

53 Ramos-Romero, S. *et al.* Modifications of Gut Microbiota after Grape Pomace

Supplementation in Subjects at Cardiometabolic Risk: A Randomized Cross-Over Controlled

Clinical Trial. *Foods* **9** (2020). https://doi.org/10.3390/foods9091279

Figure 1.

Figure 2.

Figure 3.

755   **Figure 4.**

758   **Figure 5.**

762   **Figure 6.**

765   **Table 1.**

	Adults (30-70 yrs) (n=13)	>70yrs robusts (n=15)	>70yrs prefrails (n=14)	p
Gender				0.846
Male	7 (53.8)	6 (42.9)	7 (46.7)	
Female	6 (46.2)	8 (57.1)	8 (53.3)	
Mean age (years)	47.8 (±11.6)	77.8 (±5.21)[1]	76.2 (±5.07)	**< 0.001**
Household status				0.786
Solo	9 (69.2)	10 (71.4)	9 (60)	
With partner	4 (30.8)	4 (28.6)	6 (40)	
Mean Fried Frailty Index	0 (±0)	0 (±0)[2]	1.14 (±0.363)	**< 0.001**
Mean Charlson Comorbidity Index	0.923 (±1.19)	5.07 (±1.59)[3]	4.73 (±1.87)	**< 0.001**
Mean MNA score	27.7 (±1.58)	28.1 (±1.27)	27.5 (±2.30)	0.713

Figure and Table legends.

Figure 1: Taxonomic analyses of biofilms from human aging cohort. Fecal microbiota cultured *in vitro* on a polymicrobial anaerobic biofilm model underwent 16S rRNA V3-V4 gene sequencing. The study groups include individuals aged 30-70 years (n=13), individuals aged 70 years and above characterized as healthy aging (robust, n=15) and +70 years prefrail (n=14). **A.** The stacked bar plots illustrate the relative abundance of the 5 major Phyla and their top 5 most abundant Families. Each individual is depicted as a separate bar plot, highlighting taxonomic variations **B.** For each individual, the average of lowest values from four different beta-diversity matrices (Bray-Curtis, Jaccard, unifrac, and weighted unifrac) represented individual uniqueness. Statistical significance was determined by ANOVA followed by Tukey's for multiple comparisons, where P<0.05 was considered significant (* for P< 0.05, *** for P<0.0010). **C.** Heatmap represents Kendall's correlation coefficients between biofilm taxa counts clinical parameters from human donors. Clinical parameters include discrete and continuous clinical features: household (1: solo 2: with a partner), gender (1: male, 2: female), MNA score, Charlson index, age, and Fried Frailty Index. The color gradient from blue to red signifies negative to positive correlations, and * corresponds to Kendall's P value <0.05. For enhanced clarity, the heatmap was filtered to include only significantly positive or negative correlations (P<0.05).

Figure 2. Prefrailty associated microb-aging is linked with physical alterations of human biofilms. Feces from individuals aged 30-70 years, 70+ years robust, and 70+ years prefrail, were cultured *in vitro* as a polymicrobial anaerobic biofilm. **(A-B)** The total biofilm biomass and **(D-E)** and the total sialic acid/N-acetylglucosamine glycoproteins content of extracellular biofilm matrix were measured with WGA stain. **(A-D)** Pearson correlation relationship between biofilm measures (biomass in **A.**, WGA-matrix in **D.**) and donor's age are presented.

Shaded area around the regression line represents the 95% confidence interval. **(B-E)** Scatter

plots depict for each individual's its corresponding value of biofilm biomass (in **B.**) and biofilm

WGA-matrix (in **E.**) and were grouped for each three cohort of individuals. In panels B and E

each dot is the average value of an individual donor biofilm (at least 12 replicates). The

gender of each donor is indicated using open(male)/filled(female) circles. Statistical

significance was determined by ANOVA followed by Tukey's for multiple comparisons, where

$P<0.05$ was considered significant. **(C-F)** Heatmaps represent Kendall (in **F.**) or Pearson (in **C.**)

correlation coefficients between biofilm physical structure and clinical data (in **C.**) or biofilm

taxa counts (in **F.**). Clinical parameters include discrete and continuous clinical features:

household (1: solo 2: with a partner), gender (1: male, 2: female), MNA score, Charlson index,

age, and Fried Frailty Index. The color gradient ranges from blue (negative) to red (positive)

correlations, with statistically significant associations ($P<0.05$) marked with *. For enhanced

clarity, the heatmap in panel **F.** was filtered to include only taxa rank with significantly

positive or negative correlations ($P<0.05$).

Figure 3. Prefrailty associated microb-aging is linked with pathogenic traits of human

biofilms. Feces from individuals aged 30-70 years, 70+ years robust, and 70+ years prefrail,

were cultured *in vitro* as a polymicrobial anaerobic biofilm. **(A-B)** The rate of dispersal of

planktonic bacteria, biofilm-dispersed bacteria, during a 24-hour period from biofilms was

measured. Raw values were normalized to the average dispersal rate of the 30-70 years

group. **(D-E)** Normalized counts of biofilm-dispersed bacteria were cultured on the apical

surface of the human Caco2/HT29MTX monolayer. After a 4-hour coculture, the number of

bacteria adhered to the epithelium was quantified. **(A-D)** Pearson correlations between

biofilm rate of dispersal (in **A.**) or adhesiveness (in **D.**) with the age of donors is presented.

(B-E) Scatter plots depict for each individual's its corresponding value of biofilm dispersal

rate (in **B.**) or adhesiveness (in **E.**) and were grouped for each three cohort of individuals. In

panels B and E each dot is the average value of an individual donor biofilm (at least 12

replicates). The gender of each donor is indicated using open(male)/filled(female) circles.

Statistical significance was determined by ANOVA followed by Tukey's for multiple

comparisons, where P<0.05 was considered significant. **(C-F)** Heatmaps represent Kendall (in

F.) or Pearson (in **C.**) correlation coefficients between biofilm pathogenic traits and clinical

data (in **C.**) or biofilm taxa counts (in **F.**). Clinical parameters include discrete and continuous

clinical features: household (1: solo 2: with a partner), gender (1: male, 2: female), MNA

score, Charlson index, age, and Fried Frailty Index. The color gradient ranges from blue

(negative) to red (positive) correlations, with statistically significant associations (P<0.05)

marked with *. For enhanced clarity, the heatmap in panel **F.** was filtered to include only taxa

rank with significantly positive or negative correlations (P<0.05).

Figure 4. Increased host epithelial inflammatory response to prefrailty-associated human

biofilms. After coculture with a normalized amount of biofilm-dispersed bacteria, epithelial

cells were collected for RNA extraction. **A.** The heatmap represents scaled relative expression

of a targeted list of host response genes. Specifically, these genes are involved in innate

inflammatory signaling (*IL6*, *TNF*, *IL8*, *IL10*, *COX2*, and *BD2*), secretion (*BD2* and *MUC2*), and

tight junction function (*OCLN* and *CLDN2*). The color gradient from blue to red correspond to

scale relative mRNA expression. **B.** Scatter plot depicts relative mRNA expression of *IL6*, *TNF*,

CLDN2 and *OCLN* for each individual's and were grouped for each three cohort of individuals.

The gender of each donor is indicated using open(male)/filled(female) circles. Dotted line at

1 corresponds to basal mRNA expression in wells without bacteria. Statistical significance

was determined by ANOVA followed by Tukey's for multiple comparisons, where P<0.05 was

considered significant. **C.** Gene expression dataset was ordinated using Partial least squares-

discriminant analysis (PLS-DA) with its corresponding loading vectors. Pairwise PERMANOVA

revealed statistical differences between the group of 30-70 years and +70 years prefrail

(P=0.007). **D.** The heatmap corresponds to Pearson correlation coefficients between host

epithelial gene response and biofilm features. The color gradient from blue to red signifies

negative to positive correlations, and * corresponds to Pearson P value <0.05.

Figure 5. Prefrailty aging in mice is linked with altered gut biofilm and emergence of

pathogenic behavior when transferred into recipient mice. (A-C.) Using 16S fluorescent *in*

situ hybridization, we visualized tissue-associated microbiota in the distal colon of young (6

months, 6M, n=8, **A.**), aged (24 months, 24M) robust (n=7, **B.**), and aged prefrail (n=7, **C.**)

mice. Representative images are presented with blue is DAPI staining for host nuclei, green

is fluorescein-coupled wheat germ agglutinin for glycoproteins-rich content (e.g., mucus

layer), and red is the 16S-Cyanine3 probe for all bacteria. Dashed lines represent the limits

of mucosal tissue and the "inner mucus layer". Arrows highlight areas of interest. Scale bars

are 25 μm. **D.** Biofilm-damage scores were calculated for the three groups of mice, with dots

representing the mean score from at least 3 fields per animal. **(E-G.)** C57Bl/6 animals (8

weeks old) were previously exposed to 10-day antibiotics and underwent oral gavage with

either vehicle (PBS, n=5), or with feces from young (6M, n=9), aged robust (24M robust, n=8),

and aged prefrail (24M prefrail, n=7) mice. **E.** Four days after inoculation, feces were

collected for microbial taxonomy assessment using 16S rRNA V3-V4 gene sequencing. Colons

were harvested for (**F.**) parietal thickness measurement and (**G.**) macroscopic damage score.

(D, F, G) Bar plots corresponds to mean value per group ±SEM. Statistical significance was

determined by ANOVA followed by Dunnett's for multiple comparisons, where P<0.05 was

considered significant (* for P<0.05, ** for P<0.01).

 Figure 6 Positive impact of grape pomace on prefrailty-associated human biofilm structure

and virulence. Polymicrobial anaerobic biofilms were generated from human feces of

prefrails individuals aged >70yrs, n=4-5 per group) and exposed to grape pomace diluted in

microbial broth at concentrations of 3 to 30 mg/ml for 24 hours. **A.** Total biomass was

evaluated using safranin-O stain, and the data are presented as the fold change of mean

biomass from control prefrail biofilms (broth media). **B.** The rate of dispersal for each biofilm

was evaluated as a ratio between amount of biofilm-dispersed bacteria and biofilm biomass

previously assessed. **C.** Normalized counts of biofilm-dispersed bacteria were cultured on the

apical surface of the human Caco2/HT29MTX monolayer. After a 4-hour coculture, the

number of bacteria adhered to the epithelium was quantified by plating and was represented

as fold change of vehicle treated biofilms (broth media). **D-F**. Dispersed bacteria were

collected from biofilms previously exposed to grape pomace. Human intestinal epithelial cell

lines (Caco2 and HT29MTX, at a 3:1 ratio) were then exposed to these bacteria for 4 hours,

and mRNA expression of *IL6* (**D.**), *TNF* (**E.**) and *IL8* (**F.**) genes were assessed. **(A-C)** Bar plots

corresponds to average value ± SEM of 12 biofilm replicates from human donors

(represented as individual dots). **(D-F)** Bar plots corresponds to average value ± SEM of

relative mRNA expression of vehicle control group (n=4 human donors). Statistical

significance was determined by ANOVA followed by Dunnett's for multiple comparisons,

where P<0.05 was considered significant and * for P<0.05 and *** for P<0.001.

Table 1. Human cohort characteristics. Data are represented as mean (± standard deviation)

for continuous variables and n (%) for categorical variables. MNA: Mini nutritional

assessment. Statistical significance was determined by ANOVA (Fisher test) followed by

Tukey's for multiple comparisons for continuous variables and Chi squared test for

categorical variables, where P<0.05 was considered significant. [1] Prefrail vs robust (p= 0.846)

886   and Prefrail vs Adults (p< 0.001), [2] Prefrail vs robust (p< 0.001) and Prefrail vs Adults (p<

887   0.001), [3] Prefrail vs robust (p= 0.836) and Prefrail vs Adults (p< 0.001).