Pathogenicity of Commensal Gut Biofilm In Prefrail Aging

Guillaume Le Cosquer[1,2], Melissa Pannier[1], Elodie Meunier[1], Julie Thevenin[1], Sophie

Guyonnet[3,*], Bruno Vellas[3,*], Yohan Santin[4,*], Bruno Guiard[4,*], Angelo Parini[4,*], Louis Buscail[2],

Barbara Bournet[2], Damien Guillemet[5], Celine Deraison[1], Nathalie Vergnolle[1,6,#,*], Jean-Paul

Motta[1,#,*,✉]   *for the IHU HealthAge INSPIRE/Open Science study group*

Affiliations:

[1]Institute of Digestive Health Research (IRSD), INSERM, Toulouse University, INRAe, ENVT,

University Toulouse III Paul Sabatier (UPS), 31300 Toulouse, France

[2]Department of Gastroenterology and Pancreatology, Toulouse University Hospital (CHU

Toulouse), Toulouse, France and Toulouse University, University Toulouse III Paul Sabatier

(UPS), 31059 Toulouse, France

[3] Institute of Aging, Toulouse University Hospital (CHU Toulouse), Toulouse, France and

CERPOP, Toulouse University, INSERM, University Toulouse III Paul Sabatier (UPS), 31300

Toulouse, France

[4] Institute of Metabolic and Cardiovascular Diseases (I2MC), INSERM, Toulouse University,

University Toulouse III Paul Sabatier (UPS), 31432, Toulouse, France

[5] Nexira, 129 Chemin de Croisset, 76000 Rouen, France

[6] Department of Physiology and Pharmacology, University of Calgary Cumming School of

Medicine, 3330 Hospital Drive, NW Calgary, AB, T2N4N1, Canada

* members of the IHU HealthAge INSPIRE/Open Science study group (full list in

supplementary material)

[#] Co-supervision

[✉]Correspondence to: jean-paul.motta@inserm.fr

Keywords: gut microbiota; aging; microb-aging; gut biofilm; prefrailty; geroscience

Main text – ≈.6.700 words or ≈38.000 characters, excluding Title Page, Tables, Figures, and

References

Abstract – 217 words

Display items – 6 Figures + 1 Table

Running title: Pathogenicity of prefrail gut biofilms

30 **Abstract.** Achieving healthy aging is a global challenge requiring a deeper understanding

31 of aging mechanisms, spanning from robustness to frailty. Although age-related alteration of

32 the gut microbiome (microb-aging) is well studied, there is a lack of research investigating the

33 gut microbiota during the early stages of prefrailty and understanding its behavior within the

34 context of a natural community lifestyle, in interaction with host tissue. Therefore, we used

35 fecal samples from the human aging cohort INSPIRE-T, to characterize polymicrobial biofilms.

36 Intestinal tissues and feces from an aging mouse cohort were also used. Food supplement

37 was tested for potential geroprotective effects on biofilms. Findings pointed to taxonomic

38 and physical alterations, increased instability, and virulence with regards to intestinal

39 epithelium, for biofilms of prefrail individuals, compared to their age-matched robust

40 counterparts. Multiparametric analyses further linked biofilm characteristics to clinical traits,

41 suggesting that biofilm culture could be a proxy tool for evaluating a healthy or unhealthy

42 aging status. Exposure of biofilms to the grape pomace prebiotic partially reversed the prefail

43 biofilm phenotype. In aged mice with prefrailty, but not in robust aged mice, gut microbiota

44 biofilm structure was altered and caused colon damage when transplanted into antibiotic-

45 depleted mice. These findings suggest a causal link between microb-aging in prefrailty and

46 gut inflammation, advocating microbiota-targeted therapies to reverse prefrail phenotype

47 and thereby promote healthspan.

   ## *1.*   **INTRODUCTION**

By 2030, the World Health Organization (WHO) projects that one in six individuals worldwide will be aged 60 years or older (World Health, 2020). This shift to older populations coincides with an increase in age-associated diseases and unhealthy aging. One of the first steps in the development of unhealthy aging is linked to the appearance of frailty syndrome, a clinical condition defined as decreased physiological reserve and function that heightens vulnerability and impairs the ability to withstand health stresses (Fried et al., 2001). The absence of frailty (robust individuals) has been identified as a reliable predictor of prolonged longevity without age-associated diseases (Romero-Ortuno & Kenny, 2012). The field of geroscience, which has recently emerged, is dedicated to understanding the underlying mechanisms of aging, finding methods to extend the healthspan – rather than lifespan – and to delay or impede frailty syndrome and age-associated diseases (Sierra, 2016). The scientific community has intensely focused on describing fecal microbiome composition and changes in the aging population (Bosco & Noti, 2021; DeJong et al., 2020; Ghosh et al., 2022; O'Toole & Jeffery, 2015). From these studies, it has become clear that chronological age is linked to alterations within the gut microbiome, a process described as microb-aging (Bosco & Noti, 2021). A current challenge is to identify additional aspects of microb-aging, beyond taxonomy, that could be related to early signs of unhealthy aging (in prefrail individuals)(Le Cosquer et al., 2024). The involvement of microbiota behavior and virulence in such differences has not yet been investigated. Although clinical scales allow geroclinicians to categorize patients into three categories according to their frailty status (robust, prefrail, and frail individuals), biological data on the relationship between these progressive clinical changes and microb-aging dynamics are scarce. From a clinical perspective, prefrail individuals represent a potentially reversible stage before a significant decline in health. A

report on community-dwelling elders indicated a notable prefrailty-to-robust transition, with a 16% conversion rate observed over an 18-to 36-month follow-up period, compared to 0% during the same period for the frailty-to-robust transition (Gill et al., 2006). Defining conditions or treatments that would favor this transition from prefrailty-to-robust is among the most important challenge of geroscience(Gill et al., 2006).

The motivation of this study was thus to focus on the aging population exhibiting prefrail characteristics, in the search for interventions seeking at prefrailty-to-robust transition. For that, we capitalized on the French INSPIRE-T cohorts at Toulouse University Hospital, two clinically well-characterized human and mouse aging cohorts in which various non-invasive samples were collected, including fecal samples. These cohorts were designed to identify novel markers of aging, age-related diseases and health evolution that could be therapeutically targetable (Ader et al., 2021; de Souto Barreto et al., 2021; Santin et al., 2021). Considering its key importance on gut health and disease, we aimed at focusing on microbiota communities living on the surface of the mucosa, as a polymicrobial anaerobic biofilm (Motta et al., 2021). The underlying hypothesis was that microbial characteristics within biofilms might undergo alterations not only with age but also with the health status of human donors. Ultimately, our aim was to offer proof-of-concept geroprotective experimental approaches to rectify these abnormalities.

2. MATERIALS & METHODS

Human clinical data and samples

This study used human feces collected from the INSPIRE-T cohort in 2021 (ID NCT04224038) (de Souto Barreto et al., 2021). Donors were categorized into groups of 13 adults (7 men/6 women; 30-70 years) and 29 elderly subjects (+ 70 years). The last group was

divided into 14 prefrail (6 men/8 women) and 15 healthy or robust individuals (7 men/8

women). Frailty syndrome was evaluated using the Fried frailty index (see Supplementary

Table 2 for detailed Fried index calculation(Fried et al., 2001)). Donors were excluded if they

had received antibiotics, probiotics, or prebiotics within the past three months and if they

had history of any of the following conditions: inflammatory bowel disease, irritable bowel

syndrome, colorectal cancer, immunosuppression, severe liver disease. Feces were collected

at the initial visit, aliquoted (≈1 g), and stored at −80°C until analysis in the INSPIRE-T biobank

at a central laboratory (Toulouse Bio Ressources CRB , CHU Toulouse/IFB Purpan, Toulouse,

France). Clinical characteristics, including age, gender, frailty status, and household status,

were documented (Supplementary Table 1: lifestyle). For the present analysis, we used the

version 1.0 of the INSPIRE-T database. We calculated the nutritional status of each individual

using the Mini Nutritional Assessment (MNA; Supplementary Tables 3 and 4(Rubenstein et

al., 2001)) and calculated the Charlson Comorbidity Index based on their past medical

history(Charlson et al., 1987)(Supplementary Tables 5, 6, and 7). Full list of members of the

IHU HealthAge INSPIRE/Open Science study group is included in Supplementary Material.

Mouse functional data and samples

This study involved colon tissue and feces collected from the INSPIRE mouse cohort

(CREFRE – US006 / Inserm – University of Toulouse, Paul Sabatier). The cohort comprised male

and female SWISS outbred mice(Santin et al., 2021). Animals underwent health assessments

to measure frailty, using a rodent equivalent to the human Fried frailty index, the Valencia

Score(Gomez-Cabrera et al., 2017; Santin et al., 2021) (Supplementary Table 8). Behavioral

phenotyping was conducted by the Mouse Behavioral Core platform (Integrative Biology

Center, CBI, Toulouse), at the CREFRE facility. In this study, we collected samples from young

(6 months, n=8), aged robust (24 months, n=8), and aged prefrail animals (24 months, n=8).

Distal colon was fixed in methacarn solution (6/3/1 of methanol/chloroform/acetic acid), and

embedded in paraffin. Feces were collected and stored at −80°C. Both paraffin blocks and

frozen feces were maintained in the INSPIRE mouse biobank at a central laboratory (Centre

de Ressources Biologiques-Cancer - Institut Universitaire du Cancer Toulouse – Oncopole,

CHU Toulouse, Toulouse, France).

Culture of biofilms from human feces

Human fecal samples (10-20 mg) were resuspended in supplemented Brain Heart Infusion

(BHI, Sigma-Aldrich, St. Quentin Fallavier, France) as described in(Motta et al., 2018; Motta et

al., 2019; Motta et al., 2015). Biofilms were generated using an MBEC Assay® (Innovotech,

Edmonton, Canada) with identical initial densities for each donors (OD600nm= 0.1). MBEC

plates were pre-coated overnight at room temperature with gastric porcine mucin (10 mg/ml,

Sigma-Aldrich). After 72 h of maturation, the biofilm biomass, viability, and matrix content

were characterized as described below. Alternatively, the medium was refreshed for 24h to

collect biofilm-dispersed bacteria for 16S rDNA metabarcoding, to estimate the rate of

dispersal, and to perform interaction assays with human intestinal epithelial cells. Short-term

microbiological handling was performed in a hypoxic chamber (0.1% oxygen; Don Whitley

H35; Bingley, UK). Longer incubations were performed at 37°C in an anaerobic chamber (Don

Whitley A20) or hermetic jars with an anaerobic sachet (Anaeropack, Thermo Fisher).

Staining assays on MBEC biofilm plates

Total biofilm biomass was evaluated using Gram's safranin-O solution (Sigma-Aldrich),

biofilm matrix-associated sugar content using fluorescein wheat germ agglutinin (WGA, 1/100

diluted, ThermoFisher), and viability using a resazurin assay (1 mg/ml; Sigma-Aldrich)(Motta

et al., 2018; Motta et al., 2019; Motta et al., 2015). The dispersal rate was estimated at 24h

by measuring the ratio of viable biofilm-dispersed bacteria to total biofilm biomass. Briefly,

MBEC were transferred to 96 well plate containing stains and incubated at 37°C for 2 h

(resazurin and WGA) or 20' (safranin-O). Stains were recovered by sonication at 45 kHz (USC-

500T, VWR) in a plate containing acetic acid (30%, safranin-O) or 1.5M sodium chloride

(WGA). Specific absorbance (safranin-O) or fluorescence (WGA and resazurin) was measured

using an EnSight spectrophotometer (Perkin Elmer, Villepinte, France). An average of at least

12 biofilm replicates were performed for each individual. A positive control was included for

each experiment (*E. Coli* LF82, a high biofilm producer).

In vivo transplantation: Donor microbiota preparation

Feces obtained from SWISS aged mice were divided into three groups of interest: young

(n=8 mice, 4 males and 4 females, 6 months old), aged robust (n=8 mice, 5 males and 3

females, 24 months old), and aged prefrail (n=8 mice, 5 males and 3 females, 24 months

old(Santin et al., 2021)). Feces were pooled in 1.5 mL phosphate buffered saline (PBS, Eurobio

Scientific, Les Ulis, France), homogenized, and filtered through a sterile nylon mesh strainer.

The filtered contents were centrifuged twice, and the pellet was resuspended in PBS/20%

glycerol at $OD_{600nm}=1.5$ ($\approx 10^9$ CFU/ml). All the procedures were performed under anaerobic

conditions at 37°C in an anaerobic chamber (Don Whitley A20).

In vivo transplantation: Post-antibiotics fecal microbiota transfer (FMT)

C57BL/6 mice (Janvier, Le Genest Saint-Isle, France), aged 8 weeks, were housed under

specific pathogen-free conditions (CREFRE U006, CHU Purpan). Mice were assigned to three

experimental cages: FMT-6M (n=9), FMT-24M robust (n=8), and FMT-24M prefrail (n=7), and

acclimatized for 2 weeks before the experiment. The existing microbiota was depleted by

broad-spectrum antibiotics in drinking water at concentrations of 1 g/l (ampicillin and

metronidazole, Sigma-Aldrich) and 0.5 g/l (neomycin and vancomycin, Sigma-Aldrich) for the

first five days, and half-diluted for the remaining days. On day 10, the mice underwent FMT

via oral gavage with 200 µL of previously described fecal slurry. Four days post-

transplantation, the colon was harvested for evaluation of macroscopic damage, including

colon thickness edema (in mm) and blind Wallace scoring endpoints(Wallace et al., 1995),

was assessed by a blinded and skilled experimenter. Animal procedures were approved by

the National Laboratory Animal Ethics Committee (APAFIS 2023060610225611, French

Ministry of Research).

16S rDNA amplicon sequencing

Total DNA was extracted from feces or biofilms using the EZNA isolation kit stool DNA

(VWR for biofilms) according to the manufacturer's protocol. DNA extraction and specific PCR

amplification of 16S bacterial rDNA V3-V4 regions were targeted by 341f-806R primers and

sequenced on the Illumina MiSeq 2 × 250 bp technology at Vaiomer (Toulouse, France) or

Biomnigene (Besancon, France). Demultiplexed raw read sequences were processed using

FROGS bioinformatics pipeline (FROGS v. 4.1.0(Bernard et al., 2021), Galaxy Toulouse server,

GenoToul). Adaptor and primer sequences were trimmed, and amplicons were merged using

Vsearch plugin (v.2.17.0). Cluster filtering and denoising (swarm v2.0) plugins were then used

for quality control and identification of amplicon sequence variants (ASVs). Taxonomic

assignments were performed using the 16S SILVA 138.1 database.

16S rDNA Fluorescent In situ hybridization (FISH)

For fluorescence in situ hybridization (FISH), methacarn-fixed mouse colon tissues were

embedded in paraffin. Slides were hybridized with a universal bacterial 16S fluorescent rRNA

probe (EUB338-Cy3, 5'GCTGCCTCCCGTAGGAGT-Cy5, Eurofins Scientific, France) and DNA was

counterstained with 4',6-diamidino-2-phenylindole (DAPI, Invitrogen) and polysaccharide

content (wheat germ agglutinin labeled with fluorescein, ThermoFisher)(Motta et al., 2018;

Motta et al., 2019). The biofilm damage score was blindly evaluated in at least three

microscopic fields for each mouse in the groups of interest, according to the criteria described

in(Motta et al., 2019). Samples that were unsatisfactorily fixed for scoring were excluded from

analysis. Images were acquired using a Leica LSM 710 confocal microscope (TRI Genotoul,

Purpan), and FIJI freeware was used for the final image mounting (v.1.51, Mac Os).

Human intestinal epithelial cells

Caco-2 (ATCC, Manassas, USA) and HT29-MTX E12 (ATCC) were grown separately in

Dulbecco's modified Eagle medium (DMEM, Eurobio Scientific, Les Ulis, France)

supplemented with 10% fetal bovine serum (FBS, Sigma-Aldrich), penicillin/streptomycin

(Invitrogen, Massy, France) and non-essential amino acid solution (Thermofisher, Paisley, UK)

at 37°C in a humidified incubator with 5% CO2 as previously described (Rolland-Fourcade et

al., 2017). Coculture at 3:1 (Caco-2:HT29) was done on flat bottom 24-wells plates at an initial

total density of 2.10^5 cell per well (Corning, Thermo Scientific). Cells were screened for

mycoplasma infection (MycoAlertTM, Lonza, Bâle, Switzerland) and treated before coculture

if necessary (PlasmocureTM, Lonza). Cells were cocultured with dispersed bacteria from

biofilms (normalized optical density; OD_{600nm}=0.01) in OPTIMEM (Thermo Fisher). Bacterial

adhesion to the intestinal epithelium was assessed after 4 h of coculture on a blood agar

plate(Motta et al., 2018; Motta et al., 2019).

Relative mRNA expression by qRT-PCR

Total RNA was extracted using TRIzol reagent (Euromedex, France) after 4 h of coculture

(Qiagen, Montreal, Canada). Reverse transcription of isolated RNA was performed using the

Maxima First Strand Kit (Invitrogen, Thermo Scientific) according to the manufacturer's

instructions. Quantitative PCR (RT²qPCR Primer, Qiagen) was performed on a LightCycler

(Roche, Basel, Switzerland) based on SYBR Green technology, using the Takyon Low Rox SYBR

kit (Eurogentec, Seraing, Belgium). Data were analyzed by the $2^{-\Delta\Delta CT}$ method using HPRT and

GAPDH (Qiagen) as control genes and expressed as fold change compared to control cells

without bacteria. PCR primers used in this study are listed in Supplementary Table 9.

Grape pomace (prebiotic) intervention

After maturation, MBEC Assay® was exposed to grape pomace solution diluted in

microbiological BHI supplemented broth at several concentrations (0.3 to 30 mg/mL) for 24h.

The medium was refreshed, and 4h later dispersed bacteria were collected to collect biofilm-

dispersed bacteria for interaction assays with human intestinal epithelial cells. Staining assays

were performed in parallel to estimate the effect of grape pomace on biofilm biomass and

matrix-associated sugar content. The industrial grape pomace used (VinOgrape™, Nexira,

France) was an extract containing purified polyphenols (>90% content, quantified by

spectrophotometry method) anthocyanins, catechin, epicatechin and oligomeric

proanthocyanidins.

Statistics and visualization

We performed statistical analyses and data visualization using R 4.3.0 – Posit (Rstudio IDE)

v2023.0.9.1 built 494 (Posit Team 2023, Boston, MA, USA) and GraphPad Prism (v10 for

MacOs, La Jolla, USA). Baseline clinical characteristics were compared between the groups

with chi-squared test for categorical variables and ANOVA (Fisher test) for continuous

variables with Tukey's method to correct P-values.

To explore the bacterial taxonomy variability among sample groups, alpha and beta

diversity metrics were performed in R using phyloseq (v.1.46.0). Principal coordinate analysis

(PCoA) and principal component analysis (PCA) were performed in R using phyloseq (v. 1.46.0)

and compared using PERMANOVA (pairwiseAdonis v.0.4.1) with 9999 permutations. Taxa bar

plots were generated in R using phyloseq (v. 1.46.0) and ggnested (v. 0.1.0). Statistical

differences between microbial communities were assessed in R using LEfSe (lefser

v.1.12(Segata et al., 2011)), Deseq2 (v.1.42(Love et al., 2014)), EdgeR (v.4.0.3(Robinson et al.,

2010)), metagenomeSeq (v.1.43.0(Paulson et al., 2013), and ANCOM(Mandal et al., 2015).

Boxplots, barplots and scatter plots were generated in R using ggplot2 (v.3.4.4) and ggpubr

(v.0.6.0).

Heatmaps were created in R using complexheatmap (v.2.18.0). Grout's test was used to

filter statistical outliers in the qPCR dataset (GraphPad Prism). Statistical significance was set

at P < 0.05. For group comparisons, we used Dunnett's or Tukey's method to correct P-values

for ANOVA and the Benjamini-Hochberg method for differential abundant taxa analyses.

Unless otherwise indicated, values are presented as the mean ± standard error of the mean.

3.   RESULTS

Characterization of the human cohort

Samples were collected from 42 individuals, stratified into 13 healthy individuals aged 30-

70 years and 29 individuals aged ≥ 70 years. Within the aged group, 15 were robust based on

the Fried frailty index(Fried et al., 2001), while 14 were prefrail (score of 1 and 2) (Table 1 and

Supplementary Table 1). All the donors had satisfactory nutritional scores (Supplementary

Table 3). As expected, the Charlson comorbidity index (Charlson et al., 1987) indicated more

comorbidities in aged individuals than in younger donors (0.923±1.19 versus 4.73±1.87,

P<0.001), but this index was not discriminant between prefrail and robust donors (5.07±1.59

versus 4.73±1.87, p=0.836; Supplementary Table 6). General characteristics of donors used in

the study are summarized in Table 1.

Taxonomic characterization of microb-aging on human biofilms

To explore links between microb-aging and prefrailty syndrome, we cultured the fecal

microbiota of aged individuals on a mucin-coated device, in an attempt to reproduce a

mucosal polymicrobial anaerobic biofilm (Motta et al., 2021). Using 16S V3-V4 metabarcoding, we identified that microbial community was dominated by Bacillota (formerly Firmicutes), followed by Actinobacteria and Bacteroidota (formerly Bacteroidetes). At Family level, biofilm core taxa belong to *Ruminococcaceae*, *Christensenellaceae,* *Bifidobacteriaceae, Akkermansiaceae, Lachnospiraceae*, and *Oscillospiraceae* (Fig.1A, Supplementary Fig.1). Being a recognized hallmark of human microb-aging, we confirmed that biofilms from aged donors had reduced α-diversity compared to 30-70 years old individuals (Chao1, ACE, Simpson, Shannon, and Fisher indices, Fig.1B). Following with β-diversity analyses (Fig. 1C, Supplementary Fig. 1B), we found that biofilms from aged robust donors exhibited heightened individual microbiome uniqueness compared with those from prefrail. Regarding the relative abundance distribution at the Family level, no hierarchical clustering was evident among the donor groups or with the Fried frailty index (Supplementary Fig.1A). We then ran various statistical models in 16S metagenomics to identify differentially abundant taxa (i.e., Deseq2, metagenomeSeq, ANCOM, and edgeR, Supplementary Tables 10 and 11, Supplementary Fig.1). Importantly, we found correspondence in age-related taxa abundance between the biofilm model and other reports on human aging cohorts(Bosco & Noti, 2021; DeJong et al., 2020; Ghosh et al., 2022; O'Toole & Jeffery, 2015) (Supplementary Fig.1C, 1D). Specifically, aged individuals presented an enrichment of taxa including *Selemonadaceae*, *Prevotellaceae*, and potential pathobionts such as *Enterobacteriaceae*. Concomitantly, there was a reduction in the abundance of important commensals within aged biofilms, including *Dialister sp., Akkermansia sp.* and *Bifidobacteriaceae* (Supplementary Fig.1C). Biofilms from aged robust individuals displayed a higher abundance of *Prevotellaceae* compared to prefrail individuals, as well as an increased representation of *Selenomonadaceae (Megamonas sp.)* and *Ruminococcaceae* (Supplementary Fig.1D).

We then expanded the investigation between clinical characteristics of donors and taxa

abundance in the biofilm model. Employing Kendall's correlation approach and excluding

non-statistically significant scores, we identified 34 taxonomic biofilm features, primarily at

the Family and Genus levels, that were significantly associated with clinical characteristics

(Fig.1D). First, aging alone was linked to a diminished abundance of core taxa such as *Slackia*

sp., *Eggerthellaceae*, and Bifidobacteriales. In addition, we found a positive correlation

between prefrailty and the relative abundance of *Monogobales, Oscillibacter sp.* and

Negativibacillus sp. Regarding clinical history, the Charlson comorbidity index was negatively

correlated with abundance of *Slackia sp.*, and *Veillonellaceae (Dialister sp.)*. Abundances of

Pasteurellaceae and *Haemophilus sp.* were inversely correlated with the MNA score, shedding

light on the potential links between these specific taxa and nutritional status. Gender

exhibited no statistically significant correlation with the abundance of microbial taxa in

cultured biofilms. Finally, regarding the living conditions of donors, we uncovered a positive

correlation between conjugal lifestyle and the microbial abundance of several taxa, including

Tannerellaceae, *Parabacteroides sp.*, and *Butyricimonas sp.* In summary, these findings

indicate that biofilms derived from fecal samples harbor alterations indicative of microb-

aging, with a reduced abundance of core taxa members coupled with an increased abundance

of Proteobacteria. Importantly, *cultured* biofilm maintains specific composition related to the

prefrailty status of their donors. In addition to these taxonomic observations, this culture

approach allowed us to further elucidate the structural and behavioral features of microb-

aging.

Physical structure of human biofilms is altered in prefrail donors

To explore microb-aging beyond taxonomy, we first characterized the physical structure

of biofilms. We observed a negative correlation between donor age and total biofilm biomass

(Pearson R = -0.6, P<0.05, Fig.2A). However, this feature was not significantly different

between prefrail and robust biofilms (Fig.2B). Regarding the extracellular matrix of biofilms,

we found a positive correlation (R=0.39, P<0.05) between age and proportion of sugar within

this matrix (Fig.2D). Interestingly, biofilms from prefrail donors exhibited a heightened sugar

matrix composition compared to the other samples (Fig.2E). We then investigated the links

between the clinical characteristics of human donors and biofilm physical structure (Kendall's

correlation approach, Fig.2C). In addition to confirming the previously described Fried and

age relationship, we found a negative correlation between biofilm biomass and the Charlson

comorbidity index (Fig.2C). Neither the donor's sex, household, nor diet (MNAscore)

demonstrated a significant association with the biofilm structure (Fig.2C). Focusing on biofilm

taxonomy (Fig.2F), we identified that high biomass biofilm, a feature of 30-70 yrs biofilms,

was positively correlated with a high abundance of *Eggerthellaceae*, *Oscillospira sp.*,

Enterorhabdus sp., and *Slackia sp.* High biofilm matrix-sugar, a feature of prefrail biofilms,

was correlated with donors harboring low abundance of Coriobacteriales and

Eggerthellaceae (Fig.2F). Overall, these observations suggest that the physical structure of

biofilms generated from 30-70 and aged individuals differs and, more importantly, that these

physical modifications may serve as discriminant features between aged individuals

predisposed to prefrailty or robustness.

Prefrailty-associated human biofilms have an hypervirulent phenotype

In addition to the physical structure, we evaluated the pathogenic behavior of human

biofilms between young and aged donors. First, we assessed the stability of biofilms,

measured as the rate of bacterial dispersal over a 24-hour period. High dispersal is considered

a virulent trait that increases the likelihood of exposing the subjacent host epithelium to

unwanted bacterial contact (Motta et al., 2018; Motta et al., 2015). Donor age was not

statistically correlated with this feature (P=0.058, Fig.3A). However, prefrail biofilms had a

significantly higher dispersal rate than biofilms from younger groups (P<0.05, Fig.3B). We then

explored the ability of biofilms to disperse bacteria that could attach to the underlying

intestinal epithelium. To this end, we exposed enterocyte/mucus-producing human epithelial

cell lines to a normalized concentration of the dispersed biofilm bacteria. We found that

higher chronological age correlated with this behavior (P<0.05, Fig.3D). More importantly, it

was dispersed bacteria from prefrail biofilms that exhibited heightened adhesiveness

compared to other groups (P<0.01, Fig.3E). Intriguingly, the adhesiveness of bacteria was

correlated with individuals having a high Charlson comorbidity index, suggesting a potential

connection between bacterial behavior and clinical history (Fig.3C). In this cohort, neither the

donor's gender, household, nor diet (MNA score) demonstrated a statistically significant

correlation's index with biofilm pathogenic behavior (Fig.3C). Regarding taxonomy, biofilm

dispersal was statistically correlated with a lower abundance of *Ruminococcaceae* and

Oscillospirales sp. In addition, we identified several taxa whose abundance was either

positively (e.g., *Adlercreutzia sp.* and Coriobacteriales) or negatively (e.g., *Akkermansiaceae*)

correlated with biofilm bacterial adhesiveness (Fig.3F). In summary, our findings underscore

that while chronological age plays a limited role in influencing biofilm behaviors, prefrailty

status emerges as a key factor shaping the instability and virulence of gut biofilms.

Prefrailty-associated biofilms trigger host innate response in human intestinal

epithelium

We then focused on the relative expression of 10 genes encoding proinflammatory

cytokines, antimicrobials, and tight junction proteins to discern patterns indicative of the host

epithelial innate response to these biofilm-dispersed bacteria. The scaled expression

heatmap revealed not only altered expression patterns in biofilms from aged versus younger

individuals, but also striking disparities within biofilms from prefrail versus robust individuals

(Fig.4A). Proinflammatory genes, such as *IL6* and *TNF*, and the tight junction genes claudin-2

(*CLDN2*) and occludin (*OCLN*) (Fig.4B) exhibited a notable trend towards increased expression

in prefrail versus robust individuals. While such trends failed to reach statistical significance

when considered alone – except in the case of *TNF* and *IL6* – the PCA ordination plot

demonstrated a clear shift in the transcriptome of epithelia exposed to bacteria from prefrail

individuals compared to younger individuals (Fig.4C, PERMANOVA P= 0.007). As visualized in

the biplot vectors, genes with the greatest contribution in the separation were those coding

for proinflammatory signals (*TNF*, *IL8 and IL6* along the PC1 axis) and those related to tight

junction proteins (*CLDN2* and *OCLN* in the PC2 axis). Expanding this analysis with biofilm

features, we discovered that high dispersal rate – indicative of a virulent prefrail phenotype

– was correlated with a higher expression of inflammatory genes (*IL6* and *TNF*, Fig.4D).

Interestingly, high adhesiveness was also positively correlated with inflammatory genes

(*COX2*) as well as genes encoding tight junction proteins (*CLDN2* and *OCLDN*; Fig.4C). Finally,

we found that an increased quantity of sugar in the biofilm matrix – a feature linked to aging

and prefrailty – was positively correlated with a higher expression of proinflammatory genes

(*IL6*, *IL8*, and *TNF*). Overall, these findings indicate that in addition to aging host immunity,

microbiota biofilms from +70 years prefrail individuals, but not in 30-70 years and +70 years

robust individuals, contain commensals with proinflammatory triggering potential.

Prefrailty aging in mice is linked to altered mucosal biofilm biogeography in colon.

Motivated by *cultured* human biofilm models, we aimed to confirm whether early signs of

prefrailty could influence the spatial organization of the gut microbiota in the colon. For this,

we used tissues from an aging mouse cohort, equivalent to humans aged 30 to 70+ years (= 6

to 24 months old, INSPIRE mouse cohort(Santin et al., 2021)). Among the 15 selected animals,

thorough frailty syndrome assessments were conducted, categorizing them into groups

mirroring healthy aging (robust, n=8) and those displaying early signs of frailty (prefrail, n=7,

Valencia Frailty Score, Supplementary Table 8(Gomez-Cabrera et al., 2017)). Upon sacrifice,

the colon-associated microbiota was visualized using 16S rDNA fluorescent *in situ*

hybridization (FISH) staining (Fig.5A). Remarkably, young mice exhibited a mucosal microbiota

organized as a dense biofilm community lining a layer of sterile mucus (outlined by dashed

lines in yellow, Fig.5A). In contrast, aged animals displayed a different spatial organization,

featuring isolated bacteria within the mucus layer that escaped from the dense mucosal

biofilm (arrows, Fig.5B and 5C). Bacterial translocation, which was almost absent in the young

cohort, was observed in the aged animals (Fig.5B and 5C). Notably, aged prefrail animals

exhibited more signs of tissue-associated biofilm alterations than aged robust animals,

including signs of virulent behavior in gut microbiota such as invasion of the mucus sterile

layer, epithelial adhesion, or translocation (biofilm damage score(Motta et al., 2019), Fig.5D).

These findings confirmed *in vivo* that prefrail status in aging is linked to altered biofilm

organization and emergence of virulent phenotypes.

Prefrailty-associated microbiota directly causes colon damage when transplanted into

young mice

To experimentally demonstrate a causal relationship between prefrailty-associated

microbiota and local gut inflammation, we conducted fecal microbiota transplantation (FMT)

in antibiotic-treated young mice. We used as transplant a pool of feces from young (6 months,

n=8), robust (24 months, n=8), and prefrail mice (24 months, n=8). As expected, the antibiotic

regimen caused transient dysbiosis in microbiota, with an increased abundance of

Enterobacteriaceae, which was partially restored after 4 days of antibiotic washout (Fig.5E

and Supplementary Fig.2). In recipient mice, the fecal 16S composition of FMT-prefrail was

significantly different from that of mice that underwent FMT with young and aged robust

microbiota (Fig.5E, Supplementary Fig.3 and 4). Mirroring previous findings in aging mice

(Langille et al., 2014) – as well as our own findings in human biofilms – mice transplanted with

prefrail microbiota conserved a significantly higher abundance of harmful Proteobacteria

taxa, combined with a reduced abundance of core commensal taxa (*Muribaculaceae*,

Christensenellaceae, *Parabactoides sp.* and *Akkermansia sp.*, Supplementary Fig.4). Regarding

intestinal tissue damage after the FMT procedure, we found that young fecal microbiota did

not cause macroscopic harm in the colon (total damage score of 1.7±0.44, Fig.5F-G). However,

when exposed to aged microbiota FMT, recipient animals had significantly higher colon

damage such as increased incidence of erythema, mucus release in the lumen, edema, and

ulcers (1.7±0.44 vs. 3.6±0.73, P= 0.035, Fig.5F-G). Further stratifying aged animals into robust

and prefrail categories revealed a substantially higher damage score in the prefrail group

compared to robust animals (total score 5.4±0.98 vs. 2±0.98, ANOVA followed by Dunnett's

test, P = 0.002, Fig.5G). These results combined with previous *in vitro* assays, confirmed that

microbiota from prefrail – but not from young and robust – directly causes intestinal damage

when transplanted into recipient young mice.

Grape pomace partially reverses prefrailty-associated biofilm structure and virulence

Considering the virulent phenotype of the prefrail microbiota and its associated damage

to the intestinal mucosa, we investigated whether this phenotype can be reversed by

microbiota-targeted strategies. Grape-derived polyphenols have long demonstrated

prebiotic properties via their activities on gut microbes (Davinelli & Scapagnini, 2022; Luo et

al., 2021; Xu et al., 2021; Ziegler et al., 2023). Indeed, the majority of edible grape polyphenols

are poorly bioavailable and reach the colonic compartment where they shape microbiome

profile (Rodriguez-Lopez et al., 2022). Therefore, we sought to determine whether industrial

grape pomace, which is already marketed as a food supplement in France, can affect the

structure and virulence of mucosal biofilms associated with prefrail aging. Focusing on the

discriminant biofilm features described above in the aged human biofilms, we found that the

grape pomace extracts restored the biomass of prefrail biofilms, at a level similar to that

found in biofilms from healthy individuals (Fig.6A, P<0.001 at 30 mg/ml). Importantly, we

found a statistically significant reduction of 79.5% in the dispersal rate in prefrail biofilms,

returning to the level recorded in biofilms from robust individuals (Fig.6B, P<0.05 at 30

mg/ml). Grape pomace had no significant effect on the increased adhesiveness of dispersed

bacteria from prefrail biofilms (Fig.6C). At a concentration of 30 mg/ml, biofilm exposure to

grape pomace led to a significant reduction of dispersed bacteria able to trigger the

expression of *TNF* (55% reduction, P<0.05, Fig.6D) or *IL6* (43.9% reduction, P>0.05, Fig.6E),

while having no impact on *IL8* (P>0.05, Fig.6F). Overall, these experiments using a biofilm

model suggested that grape pomace could directly reduce the hypervirulent phenotype

associated with the prefrail aging microbiota.

4.   DISCUSSION

Aging is commonly associated with changes in the gut microbiota, including shifts in

taxonomy and metabolism, a phenomenon referred to as microb-aging (Bosco & Noti, 2021).

Despite extensive research on fecal microbiome from various populations worldwide

(reviewed in(Bosco & Noti, 2021; DeJong et al., 2020; Ghosh et al., 2022; O'Toole & Jeffery,

2015)), significant knowledge gaps remain on the ecological implications of microb-aging

within commensal gut communities. Another under-explored area is the causal relationship

between microb-aging and low-grade inflammation of aging (Franceschi et al., 2018; Le

Cosquer et al., 2024). Additionally, the role of microbiota in frailty aging and the potential of

microbiota-targeted therapies in transitioning from prefrailty to robustness remain open

questions. Addressing these gaps is crucial for understanding the nature of microb-aging and

identifying non-chronological markers of unhealthy aging (Ader et al., 2021; Sierra, 2016).

Using non-invasive samples in a clinically well characterized human aging cohort, we thus

aimed at generating a model for microbiota forming biofilm, which is the microbial lifestyle

that resides in close proximity to the host tissue and epithelium (Motta et al., 2021). In this

study, we aimed at detecting microbiome shifts in biofilms from aged individuals at an early

stage of frailty, specifically to uncover potential therapeutic interventions that can reverse

these changes and establish new targets that promote the transition from prefrailty to

robustness (Gill et al., 2006).

From a taxonomic and alpha-diversity perspective, the cultured biofilm maintained

important microb-aging features already described in the literature, such as reduced alpha-

diversity (Fig.1B)(Bosco & Noti, 2021; Ghosh et al., 2022; Jackson et al., 2016; Maffei et al.,

2017; O'Toole & Jeffery, 2015), as well as increased microbiome uniqueness in robust but not

in prefrail aged individuals (Fig.1D)(Wilmanski et al., 2021). By comparing biofilm taxonomy

results with a recent grouping proposed by Ghosh et al.(Ghosh et al., 2022) — Groups I, II,

and III—we observed an interesting parallelism, where gut pathobionts, such as

Proteobacteria from the *Enterobacteriaceae* family, exhibited increased abundance in aged

individuals. We also noted a decline of beneficial taxa in aged biofilms, such as

Bifidobacteriaceae. Interestingly, *Akkermansia sp.*, a genus containing important commensals

involved in host/mucus homeostasis(Cani et al., 2022), showed slightly reduced abundance

in aged biofilms. Intriguingly, biofilms from aged robust individuals displayed increased taxa

traditionally associated with cohorts of healthy centenarians such as *Christencellaneaceae*

(Biagi et al., 2016; Biagi et al., 2010; Kong et al., 2016). Other disparities between robust and

prefrail individuals were also observed (Fig.2B), with taxa producing short-chain fatty acids

(SCFAs) being more abundant in biofilms from robust individuals (e.g., *Prevotellaceae*).

Interestingly, biofilms from prefrail donors displayed a slightly higher abundance of disease-

associated taxa such as *Megamonas* species, compared to robust donors biofilm (Fig.2B). As

a limitation of these taxonomy findings relates to small sample size, further investigation

within the entire INSPIRE cohort could be engaged to confirm and refine the identified

taxonomic features (+1000 volunteers, (de Souto Barreto et al., 2021)). In addition, the

longitudinal design of this cohort will enable collection from the same individuals over the

next 10 years, enabling us to track the progression of prefrailty in individuals and determine

whether they experience a shift to frailty, provided that microb-aging is not therapeutically

managed (with diet, exercise or pre- and probiotics).

In addition to taxonomic alterations, this study revealed new microbiological features that

may help distinguish between healthy and unhealthy aging in humans. Specifically, we

identified a reduced biofilm-forming capacity in most fecal samples of aged individuals

compared to that in younger individuals (Fig.3). This may indicate a common biological

pattern in the aging microbiome that is potentially linked to the loss or gain of specific taxa,

as previously discussed. More compelling are those features that distinguish between aged-

matched robust and prefrail donors, such as the amount of sugar associated with the biofilm

matrix (Fig.3). Extracellular polysaccharides are fundamental for retaining water within the

biofilm matrix, generating a highly hydrated and cohesive environment that influences cell

movement (Flemming & Wingender, 2010; Karygianni et al., 2020). Whether the altered sugar

composition of the aged prefrail biofilms used in the present study contributes to taxonomic

changes, increased dispersal, and fragmented biofilm physical structure *in vitro* (Fig.3) and *in*

vivo (Fig.6) remains to be experimentally elucidated. Another discriminant feature of

prefrailty-associated biofilms is their capacity to release bacteria with higher adhesiveness

and which triggers proinflammatory signals from the underlying epithelium (Fig.4). Although

this phenomenon has not been previously documented, metagenomic studies of fecal

microbiota have indicated an increased proportion of virulence gene pathways in microbiota

from aged frail individuals (Maffei et al., 2017).

It is yet to be fully understood whether local gut microbiota alterations directly contribute

to adverse outcomes in aged frail populations. Complementing *in vitro* findings, an *in vivo*

FMT approach reinforces the hypothesis that prefrailty associated microbes could directly

cause a sub-inflammatory phenotype in the gut. Although this aligns with other studies of

aged microbiota transfer (Parker et al., 2022), the novel message from the present study lies

in the separation between robustness and prefrailty-associated sampling. Here, we found

that macroscopic colon damage induced by aged microbiota may be more pronounced with

prefrail microbiota than with robust (Fig.6C). Undoubtedly, the interaction between the gut

microbiome and its host is bidirectional and more complex links are expected *in vivo*. A road

to be explored is the impact of a prefrailty microbiota within an aging host, where heightened

inflammatory stress and immunosenescence may compromise the host's ability to mount an

effective response to local insults (the inflammaging hypothesis (Franceschi et al., 2018)). In

addition to the virulent behavior of prefrail biofilm, local inflammation may significantly alter

microbial biotope making it less favorable to commensal gut bacteria, or even promoting the

blooming of pathogenic taxa (e.g., *Enterobacteriaceae* (Winter et al., 2013)). This could be

attributed to compromised epithelial secretory functions in high age, such as defective mucus

and antimicrobial peptide production(Sovran et al., 2019), or deficiencies in intestinal barrier

integrity (Man et al., 2015)– although the latter phenomenon remains a subject of debate

(Wilms et al., 2020). Natural polyphenols, including those present in grape pomace, have been

shown to improve cognitive function, extend the lifespan, and protect against age-related muscle loss in several mice models (Davinelli & Scapagnini, 2022; Luo et al., 2021; Xu et al., 2021; Ziegler et al., 2023). Studies in mice have shown that grape pomace in the diet can improve metabolic diseases by preventing gut microbiota dysbiosis and increasing the abundance of beneficial taxa (Chacar et al., 2018; Mezhibovsky et al., 2023; Roopchand et al., 2015). In this study, we found that an industrial grape pomace preparation favored the restoration of the physical and non-pathogenic features of biofilms *in vitro*, potentially indicating the improved resilience of microb-aging communities (Fig.7). Additionally, this compound reduced the release of proinflammatory triggering bacteria from prefrail biofilms. Previous study suggested that grape pomace did not cause a significant taxonomic shift in human gut microbiota (Ramos-Romero et al., 2020). Therefore, further research is needed to determine the precise impact of grape polyphenols on microbial virulence functions and to elucidate specific microbial genetic responses. Finally, going beyond this particular compound, the methodology presented here may support the clinical relevance of gut microbiota biofilm cultures (using non-invasive fecal samples) to assess and validate the effects of new classes of geroprotectors, such as prebiotics or probiotics, specifically targeting signs of unhealthy microb-aging.

5. CONCLUSION

Beyond the well-known taxonomic changes in the aging gut microbiome, this study points to novel functional and behavioral changes in human gut biofilm, that switched to a virulent phenotype in prefrail individuals but not in robusts. Importantly, we found that this phenotype was reversible by prebiotic treatment. Altogether, these results further establish the intestinal microbiota as a target to improve healthy aging. These discoveries might

represent a unique opportunity for geroclinicians to develop screening tools and identify new

therapeutic targets against microb-aging, to prevent frailty development in the expanding

aging population.

Author Contributions: GLC. performed most experiments, analyzed data, and provided

clinical input; MP., EM., JT., CD., NV. and JPM. contributed to *in vivo* and *in vitro* data

acquisition and analysis; SG., and BV. provided human samples, contributed to clinical

characterization, and reviewed manuscript; YS., BG., and A.P provided mice samples,

contributed to clinical characterization and reviewed manuscript; LB., BB. reviewed

manuscript and provided clinical input; JPM. performed 16S bioinformatics and statistics;

GLC., NV. and JPM., wrote the manuscript; N.V and JPM. supervised the work and obtained

fundings. All authors have read and agreed the final version of the manuscript.

Funding: Fundings came from INSPIRE-Grant AMIGO / ANR dysBIOFILM ANR-22-CE14-

0041-01 (JPM), ANR-PARCURE PRCE-CE18, 2020 (NV), and the national program "Microbiote"

from INSERM (NV and JPM). The INSPIRE-T study was supported by grants from the Region

Occitanie/Pyrénées-Méditerranée (Reference number: 1901175), the European Regional

Development Fund (ERDF) (Project number: MP0022856), and the Inspire Chairs of Excellence

funded by: Alzheimer Prevention in Occitania and Catalonia (APOC), EDENIS, KORIAN, Pfizer,

Pierre-Fabre.

Data Availability Statement: 16S rDNA sequences from the experimental biofilms and

mouse data were uploaded to the NCBI Sequence Read Archive (SRA Bioproject

PRJNA1076354). Additional biological or clinical metadata from the INSPIRE-T cohort can be

obtained by contacting the corresponding author, who will submit the request to the INSPIRE

biobanking and research committee for final approval. Other information from this study can

be obtained, upon reasonable requests, from the corresponding author.

Ethics: The INSPIRE cohort protocol ID NCT04224038 was approved by the French Ethical

Committee located in Rennes (CPP Ouest V) in October 2019. Mice experimentation protocol

has been approved by local ethic committee of Toulouse University and by the Ministry of
Superior Education and Research under the agreement APAFIS2023060610225611.

Acknowledgments: Authors thank the Department of Gastroenterology and
Pancreatology at the University Hospital of Toulouse for an MD/PhD scholarship leave to GLC.
Authors give credit to Dr. Felipe Sierra for its initial assessment of the research project, under
the umbrella of INSPIRE Call for Geroscience Research Grant. Authors thank the INSPIRE
biobanking committee for evaluating our project and approving sample collection. Authors
thank the staff of the human and mice Centre de Recherche Biologique (CRB) of the Toulouse
Hospital for samples preparation. Authors thank the CREFFRE core facility (UMS 006) for
hosting animals used in the project and contributing to histopathology analyses (R. Balouzat
and S. Milia), and the TRI-Genotoul platform for cellular imaging facilities (U1043, S. Allart).
Authors are thankful to Dr. Racaud-Sultan for critically reading the manuscript.

Conflicts of Interest: Prebiotic product (grape pomace) was gratuitously furnished by the
NEXIRA company (Rouen, France) and covered by a collaborative research grant between DG.
and NV. The company acknowledges having no influence on the results obtained during
assays presented in this manuscript.

	Adults (30-70 yrs) (n=13)	>70yrs robusts (n=15)	>70yrs prefrails (n=14)	p
Gender				0.846
Male	7 (53.8)	6 (42.9)	7 (46.7)	
Female	6 (46.2)	8 (57.1)	8 (53.3)	
Mean age (years)	47.8 (±11.6)	77.8 (±5.21)[1]	76.2 (±5.07)	**< 0.001**
Household status				0.786
Solo	9 (69.2)	10 (71.4)	9 (60)	
With partner	4 (30.8)	4 (28.6)	6 (40)	
Mean Fried Frailty Index	0 (±0)	0 (±0)[2]	1.14 (±0.363)	**< 0.001**
Mean Charlson Comorbidity Index	0.923 (±1.19)	5.07 (±1.59)[3]	4.73 (±1.87)	**< 0.001**
Mean MNA score	27.7 (±1.58)	28.1 (±1.27)	27.5 (±2.30)	0.713

590   **Table 1. Human cohort characteristics.** Data are represented as mean (± standard deviation) for continuous variables and n (%) for categorical

591   variables. MNA: Mini nutritional assessment. Statistical significance was determined by ANOVA (Fisher test) followed by Tukey's for multiple

592   comparisons for continuous variables and Chi squared test for categorical varibales, where P<0.05 was considered significant. [1] Prefrail vs robust

593   (p= 0.846) and Prefrail vs Adults (p< 0.001), [2] Prefrail vs robust (p< 0.001) and Prefrail vs Adults (p< 0.001), [3] Prefrail vs robust (p= 0.836) and

594   Prefrail vs Adults (p< 0.001)

Page 29

Fig.1: Taxonomic analyses of biofilms from human aging cohort. Fecal microbiota cultured *in vitro* on a polymicrobial anaerobic biofilm model underwent 16S rDNA metabarcoding targeting V3-V4 regions. The study groups include individuals aged 30-70 years (n=13), individuals aged 70 years and above characterized as healthy aging (robust, n=15) and +70 years prefrail (n=14). **A.** The stacked bar plots illustrate the relative abundance of the 5 major Phyla and their top 5 most abundant Families. Each individual is depicted as a separate bar plot, highlighting taxonomic variations. **B.** The heatmap displays scaled mean values of five alpha-diversity indices calculated for each individual within the three study groups. The first row, representing combined total alpha diversity. **C.** For each individual, the average of lowest values from four different beta-diversity matrices (Bray-Curtis, Jaccard, unifrac, and weighted unifrac) represented individual uniqueness. Statistical significance was determined by ANOVA followed by Tukey's for multiple comparisons, where P<0.05 was considered significant (* for P< 0.05, *** for P<0.0010). **D.** Heatmap represents Kendall's correlation coefficients between biofilm taxa counts clinical parameters from human donors. Clinical parameters include discrete and continuous clinical features: household (1: solo 2: with a partner), gender (1: male, 2: female), MNA score, Charlson index, age, and Fried Frailty Index. The color gradient from blue to red signifies negative to positive correlations, and * corresponds to Kendall's P value <0.05. For enhanced clarity, the heatmap was filtered to include only significantly positive or negative correlations (P<0.05).

Fig.2. Prefrailty associated microb-aging is linked with physical alterations of human biofilms. Feces from individuals aged 30-70 years, 70+ years robust, and 70+ years prefrail, were cultured *in vitro* as a polymicrobial anaerobic biofilm. **(A-B)** The total biofilm biomass and **(D-E)** and the total sugar content of extracellular biofilm matrix were measured. **(A-D)** Pearson correlation relationship between biofilm measures (biomass in **A.**, sugar-matrix in **D.**) and donor's age are presented. Shaded area around the regression line represents the 95% confidence interval. **(B-E)** Scatter plots depict for each individual's its corresponding value of biofilm biomass (in **B.**) and biofilm sugar-matrix (in **E.**) and were grouped for each three cohort of individuals. In panels B and E each dot is the average value of an individual donor biofilm (at least 12 replicates). The gender of each donor is indicated using open(male)/filled(female) circles. Statistical significance was determined by ANOVA followed by Tukey's for multiple comparisons, where P<0.05 was considered significant. **(C-F)** Heatmaps represent Kendall (in **F.**) or Pearson (in **C.**) correlation coefficients between biofilm physical structure and clinical data (in **C.**) or biofilm taxa counts (in **F.**). Clinical parameters include discrete and continuous clinical features: household (1: solo 2: with a partner), gender (1: male, 2: female), MNA score, Charlson index, age, and Fried Frailty Index. The color gradient from blue to red signifies negative to positive correlations, and * corresponds to P value <0.05. For enhanced clarity, the heatmap in panel **F.** was filtered to include only significantly positive or negative correlations (P<0.05).

Fig.3. Prefrailty associated microb-aging is linked with pathogenic traits of human biofilms. Feces from individuals aged 30-70 years, 70+ years robust, and 70+ years prefrail, were cultured *in vitro* as a polymicrobial anaerobic biofilm. **(A-B)** The rate of dispersal of planktonic bacteria, biofilm-dispersed bacteria, during a 24-hour period from biofilms was measured. Raw values were normalized to the average dispersal rate of the 30-70 years group. **(D-E)** Normalized counts of biofilm-dispersed bacteria were cultured on the apical surface of the human Caco2/HT29MTX monolayer. After a 4-hour coculture, the number of bacteria adhered to the epithelium was quantified. **(A-D)** Pearson correlations between biofilm rate of dispersal (in **A.**) or adhesiveness (in **D.**) with the age of donors is presented. **(B-E)** Scatter plots depict for each individual's its corresponding value of biofilm dispersal rate (in **B.**) or adhesiveness (in **E.**) and were grouped for each three cohort of individuals. In panels B and E each dot is the average value of an individual donor biofilm (at least 12 replicates). The gender of each donor is indicated using open(male)/filled(female) circles. Statistical significance was determined by ANOVA followed by Tukey's for multiple comparisons, where $P<0.05$ was considered significant. **(C-F)** Heatmaps represent Kendall (in **F.**) or Pearson (in **C.**) correlation coefficients between biofilm pathogenic traits and clinical data (in **C.**) or biofilm taxa counts (in **F.**). Clinical parameters include discrete and continuous clinical features: household (1: solo 2: with a partner), gender (1: male, 2: female), MNA score, Charlson index, age, and Fried Frailty Index. The color gradient from blue to red signifies negative to positive correlations, and * corresponds P value <0.05. For enhanced clarity, the heatmap in panel **F.** was filtered to include only significantly positive or negative correlations ($P<0.05$).

A.

Human epithelium relative mRNA expression

B.

C.

D.

Fig.4. Increased host epithelial inflammatory response to prefrailty-associated human biofilms. After coculture with a normalized amount of biofilm-dispersed bacteria, epithelial cells were collected for RNA extraction. **A.** The heatmap represents scaled relative expression of a targeted list of host response genes. Specifically, these genes are involved in innate inflammatory signaling (*IL6*, *TNF*, *IL8*, *IL10*, *COX2*, and *BD2*), secretion (*BD2* and *MUC2*), and tight junction function (*OCLN* and *CLDN2*). The color gradient from blue to red correspond to scale relative mRNA expression. **B.** Scatter plot depicts relative mRNA expression of *IL6*, *TNF*, *CLDN2* and *OCLN* for each individual's and were grouped for each three cohort of individuals. The gender of each donor is indicated using open(male)/filled(female) circles. Dotted line at 1 corresponds to basal mRNA expression in wells without bacteria. Statistical significance was determined by ANOVA followed by Tukey's for multiple comparisons, where $P<0.05$ was considered significant. **C.** Gene expression dataset was ordinated using Principal Component Analysis (PCA) with its corresponding bi-plot vectors. Pairwise PERMANOVA revealed statistical differences between the group of 30-70 years and +70 years prefrail ($P=0.007$). **D.** The heatmap corresponds to Pearson correlation coefficients between host epithelial gene response and biofilm features. The color gradient from blue to red signifies negative to positive correlations, and * corresponds to Pearson P value <0.05.

Fig.5. Prefrailty aging in mice is linked with altered gut biofilm and emergence of pathogenic behavior when transferred into recipient mice.

(A-C.) Using 16S rDNA fluorescent *in situ* hybridization, we visualized tissue-associated microbiota in the distal colon of young (6 months, 6M, n=8, A.), aged (24 months, 24M) robust (n=7, B.), and aged prefrail (n=7, **C.**) mice. Representative images are presented with blue is DAPI staining for host nuclei, green is fluorescein-coupled wheat germ agglutinin for sugar-rich content (e.g., mucus layer), and red is the 16S-Cyanine3 probe for all bacteria. Dashed lines represent the limits of mucosal tissue and the "inner mucus layer". Arrows highlight areas of interest. Scale bars are 25 μm.**D.** Biofilm-damage scores were calculated for the three groups of mice, with dots representing the mean score from at least 3 fields per animal. **(E-G.)** C57Bl/6 animals (8 weeks old) were previously exposed to 10-day antibiotics and underwent oral gavage with feces (fecal microbiota transplantation, FMT) from young (FMT-6M, n=9), aged robust (FMT-24M robust, n=8), and aged prefrail (FMT-24M prefrail, n=7) mice. **E.** Four days after transplantation, feces were collected for microbial taxonomy assessment using 16S rDNA metabarcoding. Colons were harvested for (**F.**) parietal thickness measurement and (**G.**) macroscopic damage score. **(D, F, G)** Bar plots corresponds to mean value per group ±SEM. Statistical significance was determined by ANOVA followed by Dunnett's for multiple comparisons, where $P<0.05$ was considered significant (* for $P<0.05$, ** for $P<0.01$).

Fig.6 Positive impact of grape pomace on prefrailty-associated human biofilm structure and virulence. Polymicrobial anaerobic biofilms were generated from human donor feces (30-70yrs old, >70yrs robust; >70yrs prefrail, n=4 per group) and exposed to grape pomace diluted in microbial broth at concentrations of 3 to 30 mg/ml for 24 hours. **A.** Total biomass was evaluated using safranin-O stain, and the data are presented as the fold change of mean biomass from individuals aged 30-70 years. **B.** The rate of dispersal for each biofilm was evaluated as a ratio between amount of biofilm-dispersed bacteria and biofilm biomass previously assessed. **C.** Normalized counts of biofilm-dispersed bacteria were cultured on the apical surface of the human Caco2/HT29MTX monolayer. After a 4-hour coculture, the number of bacteria adhered to the epithelium was quantified. **D-F**. Dispersed bacteria were collected from biofilms previously exposed to grape pomace. Human intestinal epithelial cell lines (Caco2 and HT29MTX, at a 3:1 ratio) were then exposed to these bacteria for 4 hours, and mRNA expression of *TNF* (**D.**), *IL6* (**E.**) and *IL8* (**F.**) genes were assessed. **(A-C)** Bar plots corresponds to average value ± SEM of 12 biofilm replicates from human donors (n=4). **(D-F)** Bar plots corresponds to average value ± SEM of relative mRNA expression of vehicle control group (n=4 human donors). Statistical significance was determined by ANOVA followed by Dunnett's for multiple comparisons, where $P < 0.05$ was considered significant and * for $P < 0.05$ and *** for $P < 0.001$.

References

Ader, I., Penicaud, L., Andrieu, S., Beard, J. R., Davezac, N., Dray, C., . . . Casteilla, L. (2021).

Healthy Aging Biomarkers: The INSPIRE's Contribution. *The Journal of Frailty & Aging, 10*(4),

313-319. doi:10.14283/jfa.2021.15

Bernard, M., Rue, O., Mariadassou, M., & Pascal, G. (2021). FROGS: a powerful tool to analyse

the diversity of fungi with special management of internal transcribed spacers. *Brief*

Bioinform, 22(6). doi:10.1093/bib/bbab318

Biagi, E., Franceschi, C., Rampelli, S., Severgnini, M., Ostan, R., Turroni, S., . . . Candela, M.

(2016). Gut Microbiota and Extreme Longevity. *Current biology: CB, 26*(11), 1480-1485.

doi:10.1016/j.cub.2016.04.016

Biagi, E., Nylund, L., Candela, M., Ostan, R., Bucci, L., Pini, E., . . . De Vos, W. (2010). Through

ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians.

PLoS ONE, 5(5), e10667. doi:10.1371/journal.pone.0010667

Bosco, N., & Noti, M. (2021). The aging gut microbiome and its impact on host immunity.

Genes Immun, 22(5-6), 289-303. doi:10.1038/s41435-021-00126-8

Cani, P. D., Depommier, C., Derrien, M., Everard, A., & de Vos, W. M. (2022). Author

Correction: Akkermansia muciniphila: paradigm for next-generation beneficial

microorganisms. *Nat Rev Gastroenterol Hepatol, 19*(10), 682. doi:10.1038/s41575-022-

00650-6

Chacar, S., Itani, T., Hajal, J., Saliba, Y., Louka, N., Faivre, J. F., . . . Fares, N. (2018). The Impact

of Long-Term Intake of Phenolic Compounds-Rich Grape Pomace on Rat Gut Microbiota. *J*

Food Sci, 83(1), 246-251. doi:10.1111/1750-3841.14006

Charlson, M. E., Pompei, P., Ales, K. L., & MacKenzie, C. R. (1987). A new method of classifying

prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis,*

40(5), 373-383. doi:10.1016/0021-9681(87)90171-8

Davinelli, S., & Scapagnini, G. (2022). Interactions between dietary polyphenols and aging gut

microbiota: A review. *Biofactors, 48*(2), 274-284. doi:10.1002/biof.1785

de Souto Barreto, P., Guyonnet, S., Ader, I., Andrieu, S., Casteilla, L., Davezac, N., . . . Vellas,

B. (2021). The INSPIRE Research Initiative: A Program for GeroScience and Healthy Aging

Research Going from Animal Models to Humans and the Healthcare System. *The Journal of*

Frailty & Aging, 10(2), 86-93. doi:10.14283/jfa.2020.18

DeJong, E. N., Surette, M. G., & Bowdish, D. M. E. (2020). The Gut Microbiota and Unhealthy

Aging: Disentangling Cause from Consequence. *Cell Host Microbe, 28*(2), 180-189.

doi:10.1016/j.chom.2020.07.013

Flemming, H. C., & Wingender, J. (2010). The biofilm matrix. *Nature Reviews Microbiology,*

8(9), 623-633. doi:10.1038/nrmicro2415

Franceschi, C., Garagnani, P., Parini, P., Giuliani, C., & Santoro, A. (2018). Inflammaging: a new

immune-metabolic viewpoint for age-related diseases. *Nature Reviews. Endocrinology,*

14(10), 576-590. doi:10.1038/s41574-018-0059-4

Fried, L. P., Tangen, C. M., Walston, J., Newman, A. B., Hirsch, C., Gottdiener, J., . . .

Cardiovascular Health Study Collaborative Research, G. (2001). Frailty in older adults:

evidence for a phenotype. *The Journals of Gerontology. Series A, Biological Sciences and*

Medical Sciences, 56(3), M146-156. doi:10.1093/gerona/56.3.m146

Ghosh, T. S., Shanahan, F., & O'Toole, P. W. (2022). The gut microbiome as a modulator of

healthy ageing. *Nat Rev Gastroenterol Hepatol, 19*(9), 565-584. doi:10.1038/s41575-022-

00605-x

Gill, T. M., Gahbauer, E. A., Allore, H. G., & Han, L. (2006). Transitions between frailty states

among community-living older persons. *Arch Intern Med, 166*(4), 418-423.

doi:10.1001/archinte.166.4.418

Gomez-Cabrera, M. C., Garcia-Valles, R., Rodriguez-Manas, L., Garcia-Garcia, F. J., Olaso-
Gonzalez, G., Salvador-Pascual, A., . . . Vina, J. (2017). A New Frailty Score for Experimental
Animals Based on the Clinical Phenotype: Inactivity as a Model of Frailty. *The Journals of*
Gerontology. Series A, Biological Sciences and Medical Sciences, 72(7), 885-891.
doi:10.1093/gerona/glw337

Jackson, M. A., Jeffery, I. B., Beaumont, M., Bell, J. T., Clark, A. G., Ley, R. E., . . . Steves, C. J.
(2016). Signatures of early frailty in the gut microbiota. *Genome Medicine, 8*(1), 8.
doi:10.1186/s13073-016-0262-7

Karygianni, L., Ren, Z., Koo, H., & Thurnheer, T. (2020). Biofilm Matrixome: Extracellular
Components in Structured Microbial Communities. *Trends Microbiol, 28*(8), 668-681.
doi:10.1016/j.tim.2020.03.016

Kong, F., Hua, Y., Zeng, B., Ning, R., Li, Y., & Zhao, J. (2016). Gut microbiota signatures of
longevity. *Current biology: CB, 26*(18), R832-R833. doi:10.1016/j.cub.2016.08.015

Langille, M. G., Meehan, C. J., Koenig, J. E., Dhanani, A. S., Rose, R. A., Howlett, S. E., & Beiko,
R. G. (2014). Microbial shifts in the aging mouse gut. *Microbiome, 2*(1), 50.
doi:10.1186/s40168-014-0050-9

Le Cosquer, G., Vergnolle, N., & Motta, J. P. (2024). Gut microb-aging and its relevance to
frailty aging. *Microbes Infect, 26*(3), 105309. doi:10.1016/j.micinf.2024.105309

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-
014-0550-8

Luo, J., Si, H., Jia, Z., & Liu, D. (2021). Dietary Anti-Aging Polyphenols and Potential
Mechanisms. *Antioxidants (Basel), 10*(2). doi:10.3390/antiox10020283

Maffei, V. J., Kim, S., Blanchard, E. t., Luo, M., Jazwinski, S. M., Taylor, C. M., & Welsh, D. A.
(2017). Biological Aging and the Human Gut Microbiota. *The Journals of Gerontology. Series*
A, Biological Sciences and Medical Sciences, 72(11), 1474-1482. doi:10.1093/gerona/glx042

Man, A. L., Bertelli, E., Rentini, S., Regoli, M., Briars, G., Marini, M., . . . Nicoletti, C. (2015).
Age-associated modifications of intestinal permeability and innate immunity in human small
intestine. *Clin Sci (Lond), 129*(7), 515-527. doi:10.1042/CS20150046

Mandal, S., Van Treuren, W., White, R. A., Eggesbo, M., Knight, R., & Peddada, S. D. (2015).
Analysis of composition of microbiomes: a novel method for studying microbial composition.
Microb Ecol Health Dis, 26, 27663. doi:10.3402/mehd.v26.27663

Mezhibovsky, E., Tveter, K. M., Villa-Rodriguez, J. A., Bacalia, K., Kshatriya, D., Desai, N., . . .
Roopchand, D. E. (2023). Grape Polyphenols May Prevent High-Fat Diet-Induced Dampening
of the Hypothalamic-Pituitary-Adrenal Axis in Male Mice. *J Endocr Soc, 7*(9), bvad095.
doi:10.1210/jendso/bvad095

Motta, J. P., Allain, T., Green-Harrison, L. E., Groves, R. A., Feener, T., Ramay, H., . . . Buret, A.
G. (2018). Iron Sequestration in Microbiota Biofilms As A Novel Strategy for Treating
Inflammatory Bowel Disease. *Inflamm Bowel Dis, 24*(7), 1493-1502. doi:10.1093/ibd/izy116

Motta, J. P., Denadai-Souza, A., Sagnat, D., Guiraud, L., Edir, A., Bonnart, C., . . . Vergnolle, N.
(2019). Active thrombin produced by the intestinal epithelium controls mucosal biofilms. *Nat*
Commun, 10(1), 3224. doi:10.1038/s41467-019-11140-w

Motta, J. P., Flannigan, K. L., Agbor, T. A., Beatty, J. K., Blackler, R. W., Workentine, M. L., . . .
Wallace, J. L. (2015). Hydrogen sulfide protects from colitis and restores intestinal microbiota
biofilm and mucus production. *Inflamm Bowel Dis, 21*(5), 1006-1017.
doi:10.1097/MIB.0000000000000345

Motta, J. P., Wallace, J. L., Buret, A. G., Deraison, C., & Vergnolle, N. (2021). Gastrointestinal
biofilms in health and disease. *Nat Rev Gastroenterol Hepatol, 18*(5), 314-334.
doi:10.1038/s41575-020-00397-y

O'Toole, P. W., & Jeffery, I. B. (2015). Gut microbiota and aging. *Science (New York, N.Y.),*
350(6265), 1214-1215. doi:10.1126/science.aac8469

Parker, A., Romano, S., Ansorge, R., Aboelnour, A., Le Gall, G., Savva, G. M., . . . Carding, S. R.
(2022). Fecal microbiota transfer between young and aged mice reverses hallmarks of the
aging gut, eye, and brain. *Microbiome, 10*(1), 68. doi:10.1186/s40168-022-01243-w

Paulson, J. N., Stine, O. C., Bravo, H. C., & Pop, M. (2013). Differential abundance analysis for
microbial marker-gene surveys. *Nat Methods, 10*(12), 1200-1202. doi:10.1038/nmeth.2658

Ramos-Romero, S., Martinez-Maqueda, D., Hereu, M., Amezqueta, S., Torres, J. L., & Perez-
Jimenez, J. (2020). Modifications of Gut Microbiota after Grape Pomace Supplementation in
Subjects at Cardiometabolic Risk: A Randomized Cross-Over Controlled Clinical Trial. *Foods,*
9(9). doi:10.3390/foods9091279

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics, 26*(1), 139-
140. doi:10.1093/bioinformatics/btp616

Rodriguez-Lopez, P., Rueda-Robles, A., Borrás-Linares, I., Quirantes-Piné, R. M., Emanuelli, T.,
Segura-Carretero, A., & Lozano-Sánchez, J. (2022). Grape and Grape-Based Product
Polyphenols: A Systematic Review of Health Properties, Bioavailability, and Gut Microbiota
Interactions. *Horticulturae, 8*(7), 583.

Rolland-Fourcade, C., Denadai-Souza, A., Cirillo, C., Lopez, C., Jaramillo, J. O., Desormeaux, C.,
. . . Vergnolle, N. (2017). Epithelial expression and function of trypsin-3 in irritable bowel
syndrome. *Gut, 66*(10), 1767-1778. doi:10.1136/gutjnl-2016-312094

Romero-Ortuno, R., & Kenny, R. A. (2012). The frailty index in Europeans: association with age
and mortality. *Age Ageing, 41*(5), 684-689. doi:10.1093/ageing/afs051

Roopchand, D. E., Carmody, R. N., Kuhn, P., Moskal, K., Rojas-Silva, P., Turnbaugh, P. J., &
Raskin, I. (2015). Dietary Polyphenols Promote Growth of the Gut Bacterium Akkermansia

muciniphila and Attenuate High-Fat Diet-Induced Metabolic Syndrome. *Diabetes, 64*(8), 2847-
2858. doi:10.2337/db14-1916

Rubenstein, L. Z., Harker, J. O., Salva, A., Guigoz, Y., & Vellas, B. (2001). Screening for
undernutrition in geriatric practice: developing the short-form mini-nutritional assessment
(MNA-SF). *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences,*
56(6), M366-372. doi:10.1093/gerona/56.6.m366

Santin, Y., Lopez, S., Ader, I., Andrieu, S., Blanchard, N., Carriere, A., . . . Parini, A. (2021).
Towards a Large-Scale Assessment of the Relationship between Biological and Chronological
Aging: The INSPIRE Mouse Cohort. *The Journal of Frailty & Aging, 10*(2), 121-131.
doi:10.14283/jfa.2020.43

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C.
(2011). Metagenomic biomarker discovery and explanation. *Genome Biol, 12*(6), R60.
doi:10.1186/gb-2011-12-6-r60

Sierra, F. (2016). The Emergence of Geroscience as an Interdisciplinary Approach to the
Enhancement of Health Span and Life Span. *Cold Spring Harb Perspect Med, 6*(4), a025163.
doi:10.1101/cshperspect.a025163

Sovran, B., Hugenholtz, F., Elderman, M., Van Beek, A. A., Graversen, K., Huijskes, M., . . .
Wells, J. M. (2019). Age-associated Impairment of the Mucus Barrier Function is Associated
with Profound Changes in Microbiota and Immunity. *Sci Rep, 9*(1), 1437. doi:10.1038/s41598-
018-35228-3

Wallace, J. L., Le, T., Carter, L., Appleyard, C. B., & Beck, P. L. (1995). Hapten-induced chronic
colitis in the rat: alternatives to trinitrobenzene sulfonic acid. *J Pharmacol Toxicol Methods,*
33(4), 237-239. doi:10.1016/1056-8719(95)00001-x

Wilmanski, T., Diener, C., Rappaport, N., Patwardhan, S., Wiedrick, J., Lapidus, J., . . . Price, N.
D. (2021). Gut microbiome pattern reflects healthy ageing and predicts survival in humans.
Nature Metabolism, 3(2), 274-286. doi:10.1038/s42255-021-00348-0

Wilms, E., Troost, F. J., Elizalde, M., Winkens, B., de Vos, P., Mujagic, Z., . . . Masclee, A. A. M.

(2020). Intestinal barrier function is maintained with aging - a comprehensive study in healthy

subjects and irritable bowel syndrome patients. *Sci Rep, 10*(1), 475. doi:10.1038/s41598-019-

57106-2

Winter, S. E., Winter, M. G., Xavier, M. N., Thiennimitr, P., Poon, V., Keestra, A. M., . . .

Baumler, A. J. (2013). Host-derived nitrate boosts growth of E. coli in the inflamed gut. *Science*

(New York, N.Y.), 339(6120), 708-711. doi:10.1126/science.1232467

World Health, O. (2020). *Decade of healthy ageing: baseline report*: World Health

Organization.

Xu, Q., Fu, Q., Li, Z., Liu, H., Wang, Y., Lin, X., . . . Sun, Y. (2021). The flavonoid procyanidin C1

has senotherapeutic activity and increases lifespan in mice. *Nature Metabolism, 3*(12), 1706-

1726. doi:10.1038/s42255-021-00491-8

Ziegler, T., Tsiountsioura, M., Meixner-Goetz, L., Cvirn, G., & Lamprecht, M. (2023).

Polyphenols' Impact on Selected Biomarkers of Brain Aging in Healthy Middle-Aged and

Elderly Subjects: A Review of Clinical Trials. *Nutrients, 15*(17). doi:10.3390/nu15173770