Muscone-specific olfactory protein reveals the putative scent-marking pheromone in the Sunda pangolin (*Manis javanica*)

Zhongbo.Yu[1,2], Tao. Meng[3], Luyao.Yu[1,2], Yichen. Zhou[1,2], Tengcheng.Que[4,5],

Meihong.He[4], Haijing.Wang[3], Yingjiao.Li[4], Liling. Liu[4], Wenjian.Liu[5], Yinliang.Wang[1,2*],

Bingzhong.Ren[1,2]*

[1]Jilin Provincial Key Laboratory of Animal Resource Conservation and Utilization,

School of Life Science, Northeast Normal University, Changchun 130024, China;

[2]Key Laboratory of Vegetation Ecology, MOE, Northeast Normal University, Changchun

130024, China

[3]Guangxi Forestry Inventory and Planning Institute, Nanning, 530011, China.

[4]Guangxi Zhuang Autonomous Terrestrial Wildlife Rescue Research and Epidemic

 Diseases Monitoring Center, Nanning, 530003, China

[5]Faculty of Data Science City University of Macau, Macau, 999078, China

*Corresponding author: Dr. Yinliang. Wang and Dr. Bingzhong. Ren

Email: wangyl392@nenu.edu.cn ; bzren@nenu.edu.cn

Abstract

The Sunda pangolin (*Manis javanica*) is a burrowing and nocturnal animal, and they have poor vision, thus, intraspecies communication relies on olfaction, such as mating, warning, and scent-marking. However, the intraspecies pheromone in pangolins remains unknown. In this study, all the odorant-binding proteins in Sunda pangolins were functionally expressed, and they were screened against a panel of 32 volatiles that were derived from the pangolin's urine, feces, and anal gland secretions. Reverse chemical ecology identified that *M. javanica* odorant-binding protein 3 (MjavOBP3) possesses the highest binding affinity to muscone. A subsequent behavior-tracking assay showed that only males can sense muscone; thus, we hypothesize that muscone is a male-specific scent-marking pheromone. Meanwhile, the structural study showed that Tyr117 contributes the most to muscone's binding, which was further validated by site-directed mutagenesis. The findings clarify the scent-marking mechanism in pangolins, and muscone could potentially be used to support the monitoring and conservation of this endangered animal.

Author Summary

The Sunda pangolin is an endangered mammal that is native to Southeast Asia and is threatened due to its economic value. They are cave-living and nocturnal, poor vision; thus, their intraspecies communication is highly reliant on olfaction. Although they are generally solitary, they have been observed to have some social aspects in the wild, such as breeding and territorial behaviors, which are mediated by scents. However, no previous study has investigated the type of pheromones and how they are detected. Using the reverse chemical ecology approach, MjavOBP3 was found to bind to muscone with high affinity, and behavior-tracking assay was performed under well-controlled artificial rearing conditions, which showed only male pangolins can recognize muscone, suggesting its potential male-specific pheromone role.

Introduction

Scent-marking is one of the most conspicuous behaviors of mammals and other terrestrial vertebrates. It has long intrigued researchers from various disciplines [1]. Although the chemical composition, source, and behavioral effect of scent-marking are highly diverse among different taxa, it has been suggested to have four main functions: 1) attracting females to increase mating and reproductive success [2]; 2) advertising males' territory ownership and social dominance status [3]; 3) marking of food resources to increase foraging efficiency [4]; and 4) individual recognition among different species and populations [5]. In some cases, scent marking not only advertises the existence of the scent owners but also includes information about the individual's health, infections, age, sex, and mating state [5-7]. The source of the scent marking that is used by mammals is usually urine, feces, and glandular secretions [8]. For instance, Siberian wolves (*Canis lupus*) deposit feces at crossroads to increase the effectiveness of territory maintenance [9, 10], and female Asian elephants (*Elephas maximus*) use urine to signal males of their readiness to mate [11]. Anal gland secretions are also used by some Carnivora, such as ferrets (*Mustela putorius furo*), meerkats (*Suricata suricatta*), domestic cats (*Felis catus*), and giant panda (*Ailuropoda melanoleuca*), and the scent that is emitted from their anal gland provides important information for intraspecies individual recognition [12-15].

The Sunda pangolin belongs to Manidae, Pholidota, and is closely related to the family Carnivora. There are only eight extant species in the world [16]. Due to the economic value of their meat, scales (attributed medicinal properties), and skins, enormous numbers of pangolins have been seized and traded in the black market, especially in Southeast Asia [17]. Currently, *M. javanica* is classified as a critically endangered species and is listed in Appendix II of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) [18]. Malayan pangolins is nocturnal and has a highly specialized diet, foraging only on ants and

termites, with genomic evidence showing that their olfactory receptor gene families are

significantly expanded, supporting that they have a well-developed olfactory system [19].

Pangolins are generally solitary but they also have home ranges [20], which are more

distinguished in males. Male pangolins have been observed to display aggressive behavior toward

each other to defend their home ranges, suggesting that they are territorial [21]. Most of the social

interactions of *M. javanica* are believed to be scent-based and generally comprised of urine and

anal glands secretions [20]; however, well-controlled behavioral experiments on *M. javanica* are

still lacking, and our knowledge of the behaviors of *M. javanica* still relies on the observations

from zoos and nature reserves.

Consequently, the reverse chemical ecology approach may shed light on the communication

behaviors of Sunda pangolin, and it is a simple and effective tool for screening biologically active

scents and clarifying their molecular mechanisms [22]. Among the olfactory proteins, odorant

binding proteins (OBPs), which belong to the lipocalin family, might be specifically dedicated to

pheromonal communication [23]. When compared with insect OBPs, the function of vertebrate

OBPs remains less well understood; however, evidence suggests that these proteins might also be

involved in scent markings. For instance, three OBPs were identified in the anal sac gland of dogs

(*Canis lupus familiaris*) and are involved in the signal release and nasal mucosa reception of scent

secretions [24]. Similarly, rodent urine also contains a class of lipocalins that play a key role in

individual recognition [25]. Although the genome of *M. javanica* was sequenced many years ago,

at present, the functional expression of proteins of this kind has not been studied in this species.

Moreover, the behavioral and molecular mechanisms of scent marking in Sunda pangolin remain

elusive.

Consequently, in this study, we analyzed the *M. javanica* transcriptome from different

tissues, and identified, cloned, and analyzed the expression of all three *M. javanica* OBPs

(MjavOBPs). Based on the reverse chemical ecology approach, we screened these OBPs against a

panel of volatiles from their urine, feces, and anal gland secretions with the aim to explore the

scent-marking-related chemical signals. Additionally, the behavioral effect of the active ligands

was evaluated using a behavioral tracking assay. Lastly, the key binding sites of the OBP-ligand

complex were identified and verified by molecular modeling to clarify the structural-based

selectivity of the MjavOBPs.

Results

Candidate volatiles were associated with scent marking

Forty-nine volatiles were identified by a gas chromatography-mass spectrometry (GC-MS)

analysis of urine, feces, and anal gland secretions (Fig 1; S1 Table), namely: 21 fatty acids, seven

ketones, seven alcohols, four esters, three alkanes, three aromatic compounds, and four other

compounds. The top three abundant volatiles in the feces of the males and females were 4-cresol

($\male$: 40.01 ± 2.7%, $\female$: 44.67 ± 4.33%), isovaleric acid ($\male$: 10.58 ± 0.76%, $\female$: 9.01 ± 1.72%), and

butanoic acid ($\male$: 7.05 ± 0.75%, $\female$: 10.44 ± 1.24%; Fig 1A and B). Then, 3-methylindole, a

common volatile in feces, was detected ($\male$: 0.66 ± 0.51%, $\female$: 4.84 ± 1.55%), and, the ant volatiles

dimethyl trisulfide ($\male$: 2.67 ± 1.64%, $\female$: 4.24 ± 1.02%) and farnesene ($\male$: 0.24 ± 0.01%, $\female$: 0.09

± 0.04%) were also detected, which might be correlated with the pangolin's foraging habitats. The

top three volatiles in the urine of the males and females had minor differences, with hexadecanoic

acid (33.63 ± 1.20%), zibeton (22.37 ± 4.97%), and benzaldehyde (15.00 ± 4.03%) being the top

three volatiles in males, and those in females were hexadecanoic acid (31.73 ± 7.19%),

tetradecanoic acid (22.04 ± 3.05%), and zibeton (12.50 ± 2.29%; Fig 1C and D). Additionally,

phenol (14.82 ± 4.06%), 2-methylheptanoic acid (10.76 ± 0.21%), and 2-methyldecanoic acid

(10.22 ± 0.62%) were the most abundant volatiles in the anal gland secretions (Fig 1E).

126 Interestingly, muscone, which is the pheromone in musk deer, and 8-cyclohexadecen-1-one, a

127 volatile that has a similar chemical structure to muscone, were detected in the anal secretions.

Muscone binds to MjavOBP3 with high affinity

A transcriptome analysis identified 34 lipocalin proteins, which belong to 14 different

families, including three OBPs, 12 fatty acid-binding proteins, five retinoid-binding proteins, two

β-lactoglobulins, and 12 other lipocalin proteins (Table 1). MjavOBP1 was only expressed in the

liver, while MjavOBP2 and MjavOBP3 were specifically expressed in the nasal cavity,

suggesting their potential olfactory sensing roles (Fig 2A; S2 Table).

A homology analysis showed that pangolins are most closely related to the order Carnivora,

and they diverged during approximately the post-Cretaceous period (74 Mya). The number of

OBPs highly varied among different species, ranging from 1 to 27 (S1 Fig; S3 Table). The

number of OBPs in Chiroptera, Primates, and Carnivora was close to that of pangolins, and the

order Artiodactyla had up to 27 OBPs. In the Cetacea, all the OBPs appeared to be pseudogenes

that contained a premature termination codon, suggesting the degeneration of their olfactory-

sensing in underwater habits (Fig 2B).

We then analyzed the functional expression of all three MjavOBPs. After purification,

sodium dodecyl-sulfate polyacrylamide gel electrophoresis (SDS-PAGE) showed that the

MjavOBPs had a target band at approximately 20 kDa (S2 Fig), which was consistent with the

predicted length of the amino acid sequence of those proteins. A liquid chromatography-mass

spectrometry (LC-MS) analysis showed that the coverage rates of the measured amino acid

sequences ranged from 83% to 100% when compared with the reference sequences, validating the

MjavOBPs that were purified (Fig 3A-C).

Subsequently, three MjavOBPs were screened against a panel of 32 candidate scent-marking

and plant volatiles using N-phenyl-1-naphthylamine (1-NPN) as a fluorescent reporter (Table 2,

Fig 3D-F; S3 Fig). A competitive binding assay showed that both MjavOBP1 and MjavOBP2

bound to 2-undecanone with K_d values of $56.55 \pm 4.20 \, \mu M$ and $54.48 \pm 10.75 \, \mu M$, respectively.

Furthermore, MjavOBP1 had the best binding ability with the urinary volatile δ-dodecalactone

$(38.74 \pm 2.04 \, \mu M)$ and also bound weakly with other ligands, such as Citral and 1-octen-3-ol.

Then, MjavOBP2 had the highest binding affinity for Skatole $(40.83 \pm 8.93 \, \mu M)$, and MjavOBP3

showed strong binding $(IC50, 2.23 \pm 0.80 \, \mu M)$ with muscone, which is an anal gland secretion

volatile, and MjavOBP3 also binds to other ketones, sesquiterpenes, and aldehydes, such as 1-

octen-3-one, β-farnesene, and Hexanal, with the IC50 value being higher than $2.23 \, \mu M$ (S4

Table).

160 Behavioral responses of the pangolins to muscone

To examine the behavioral effect of muscone, a trace tracking assay was performed. The

results showed that both male and female pangolins visited and stayed in the f5 area more

frequently, where was the entrance of the resting place. After placing the control odor source

(propylene glycol), both the males and females visited and stayed on the left side of the room

more frequently, which was far away from the odor source (b8), suggesting that neither the males

nor females were able to recognize the control odor (Fig. 4A and C; S2 and 4 videos). After the

muscone was placed, the males could accurately recognize the muscone (area b8, Fig 4 B), and

obvious sniffing behavior was observed (S3 Video). In contrast, the females were mainly

distributed in f5-6, e5-6, and b5-6 (Fig 4 D), and no preference for muscone was detected (S5

Video).

Then, the average staying time of the pangolins in each area was based on three males and

three females. In the control group, the highest staying time of the males was in the order of f5

(45.09 ± 12.46 s), b2 (35.62 ± 15.04 s), and b1(31.11 ± 15.23 s), while that for the females was in

the order of f5 (38.26 ± 8.25 s), e5 (31.39 ± 15.37 s), and c2 (26.67 ± 10.49 s). The average

staying time for both sexes in b8 was 0 (Fig. 4 E and G), suggesting that neither the males nor

females could sense the control odor. In the muscone group, the highest staying time for the males

was for b8 (78.10 ± 20.04 s), f5 (30.29 ± 5.57 s), and a8 (29.48 ± 9.14 s), and the staying time in

the muscone area (b8) was significantly ($P < 0.01$) higher than that of any of the other areas (Fig.

4 F). In contrast, the highest staying time for the females occurred in f5 (35.72 ± 6.94 s), d2

(18.25 ± 8.85 s), and c2 (15.18 ± 5.78 s), and no staying or sniffing behavior was observed in the

b8 area (Fig 4 H; S5 Video). In summary, the behavioral tracking results showed that the male

pangolins showed a strong preference for muscone when compared with the control group, and

the females could not detect it.

Key binding sites for muscone on MjavOBP3

The 3D modeling and docking showed that muscone is wrapped in the center of a binding

pocket on MjavOBP3 (Fig. 5 A and B; S4 Fig). Additionally, Tyr117 can form a hydrogen bond

with muscone, with a molecular distance of 2.8 Å, and the binding of muscone is also attributed

to a Pi-alkyl interaction (Phe52), an alkyl interaction (Ile101), and Van der Waal forces (Glu68,

Tyr81, and Tyr87; Fig. 5 C). The 100 ns time-evolution root-mean-squared deviation (RMSD)

test showed that the MjavOBP3-muscone complex systems experienced a large fluctuation during

0–14 ns and reached equilibrium at 2.26 Å within 50 ns. The standard deviation of the RMSD

after 50 ns was 0.14 Å, indicating the stability and reliability of the MD trajectory (Fig. 5 D). In

the equilibrium phase, Tyr117 formed a hydrogen bond, and the other non-polar residues (Phe52,

Glu68, Tyr81, Tyr87, and Ile101) formed Van der Waal forces and a (pi)-alkyl bond, which also

contributed to the MjavOBP3-muscone interactions (Fig. 5 E). The binding free energy of the

MjavOBP3-muscone complex was predicted to be -39.42 kcal/mol (S5 Table). Subsequently, the residues with a total energy contribution (ΔG bind) exceeding −1.00 kcal/mol were chosen for further validation (Fig. 5 F), which included Tyr117, Phe52, Ile101, Glu68, Tyr81, Tyr87, and two negative control residues (Met97, inside the binding pocket; Lys130, far away from the binding pocket).

When compared with the wild type, the binding affinity of muscone in the four mutations (Y117F, F52A, Y87A, and Y81A) was significantly decreased (P < 0.01; Fig. 6 A-C; Fig. S5), which shows that Y117F does not bind to muscone with a K_d value of 116.78 ± 4.41 μ M. The fluorescence binding results were highly consistent with the modeling and docking predictions, indicating that the Tyr117, Phe52, Tyr87, and Tyr81 residues are crucial for the binding of muscone, and the Tyr117 residues contributed the most to the binding process.

The amino acids alignment on MjavOBP3 showed a conserved YxxxYxG motif (positions 77–83) and 16 conserved residues (indicated by *) among its orthologues (Fig. 6 D). The similarity of the homologous OBP3s among the different species ranged from 28% to 90%, and the highest similarity was with its close relative, *Manis pentadactyla*. The above four key residues (Tyr117, Phe52, Tyr87, and Tyr81) were highly conserved between *M. javanica* and *M. pentadactyla*, suggesting a similar olfactory sensing pathway for muscone. Among the key binding residues, Phe52 (Pi-alkyl) and Tyr81 (van der Waals force) were highly conserved among the mammals, indicating their crucial role in the OBP3 orthologues. In contrast, Tyr117 (hydrogen bond) was only conserved in *M. javanica* and *M. pentadactyla*, suggesting that it contributes the most to the muscone binding and has a crucial role in specific sensing of muscone in pangolins.

Discussion

The volatile organic compounds (VOCs) that were identified from the urine, feces, and anal gland secretions of the pangolins were mainly carboxylic acids, ketones, and alcohols, which were followed by lactones and aldehydes. Among them, carboxylic acids, aldehydes, phenols, and indoles are often detected in mammal excretions [26] and these VOCs may not convey intraspecies information. Notably, some of the VOCs, such as δ-dodecalactone, farnesene, and Muscone, are rare in nature or pangolin-specific, indicating their potential pheromone role. Specifically, δ-dodecalactone may act as a scent-marking signal in male Bengal tigers (*Panthera tigris*) [27], and Farnesene, which is derived from the urine of male mice, was shown to be a territory-marking pheromone [28]. Muscone is a rare VOC in mammal secretions, and only musk deer have been recorded to synthesize it using long-chain fatty acids [29]; they use it for territory marking and female attraction [30]. The food sources of pangolins (ants and termites) do not contain any muscone-relevant compounds [31, 32], indicating that there might be a specific biosynthesis pathway for muscone in pangolins, which would reflect their unique biological function.

The OBPs' main function is to increase the solubility of odorants and transport odorants to olfactory receptors in a hydrophilic environment [23]. The number of OBPs varies greatly among the different animal groups; for instance, the number of OBPs in insects mainly ranges from 40 to 60 [33], while the number of OBPs in mammals is comparatively small. Our study showed that the number of OBPs in the orders Pholidota, Carnivora, Perissodactyla, Chiroptera, and Primates were only 1-3, and the reduction of OBPs in these species was likely due to an insufficient olfactory transport ability to varied environmental volatiles, which limited their function to pheromone carrying [23]. Previous studies have shown that OBPs in mammals are not synthesized in the olfactory organ but are synthesized in the vomeronasal organ (VON) and nearby areas [34-37]. The VON is the main pheromone-sensing organ in mammals, and it

shortens the transportation distance of the pheromone. A recent study showed that AimelOBP3 in

giant pandas is bound with linear aldehydes, and they may be involved in pheromone recognition

[22]. Pangolins are fossorial, and their vision quality is low but their olfactory system is well

developed, and, thus, chemical signals play a key role in their intraspecies communications [20].

The fluorescence binding assay showed that MjavOBP3 had the highest binding affinity to

muscone, suggesting that muscone has a putative pheromone-sensing role.

To further explore the behavioral effect of muscone on pangolins, a well-controlled

behavioral assay was performed on this endangered species. The behavioral tracking assay

indicated that male pangolins could recognize muscone but females could not, which was

consistent with the fact that MjavOBP3 is only expressed in the olfactory epithelium of males.

Muscone is released by the male anal glands, and the anal gland is a common organ in mammals,

its main function is to synthesize anal gland secretions and lubricate the feces. However, in some

cases, the anal gland is also responsible for territory marking and intraspecies recognition (such as

in *Meles meles* and *Herpestes javanicus*) [38, 39]. Male pangolins prefer to live in solitary, and

they usually have a fixed living place with a home range of approximately 40 ha [21]. Field

observations have shown that pangolins are territorial animals, and males sometimes fight with

each other for territory. Considering that pangolins have degraded vision but a well-developed

olfaction, thus, territorial marking is believed to rely on chemical signals. Similar behavior was

also observed in other Carnivora mammals, and they are close relatives to Pholidota. For

example, *Crocuta crocuta* and *Canis lupus* use anal gland secretion for scent and territory

marking [40, 41]. The fluoresce binding assays showed that muscone is the best ligand for

MjavOBP3, and it is a macrocyclic ketone that is composed of 15 carbons with a slow release and

long-lasting aroma, which is consistent with the persistence characteristics of scent-marking

pheromones [42]. Moreover, muscone is a rare chemical in nature and has only been detected in a

few mammals, such as muskrats (*Ondatra zibethicus*) and musk deer, where it is used for scent

marking [*29*]. Therefore, we hypothesized that muscone is a male scent-marking pheromone in *M.*

javanica.

Reverse chemical ecology approaches were applied to identify the pheromone in pangolins.

The current conservation strategy would benefit from understanding how male pangolins

communicate with each other via chemical signals at a molecular level. The discovery of scent-

marking pheromones in pangolins could be used to monitor this endangered species [*43, 44*], and

a scent-based pheromone trap could be developed for the observation and relocation of this

endangered animal.

Materials and Methods

Ethics statement

280   All the animal procedures were approved by the ethics committee for animal experiments at

281 the Terrestrial Wildlife Rescue and Epidemic Diseases Surveillance Center of Guangxi, and the

282 recommended guidelines were followed. (S6 Fig)

Gas chromatography-mass spectrometry analysis

Feces, urine, and anal secretions from Sunda pangolins were collected in January 2021. The

individuals that were used in this study were three adult males and three adult females, and the

detailed information for each individual is listed in S6 Table. The volatile samples were collected

in a 10 mL bottle and adsorbed using a solid-phase microextraction syringe (Supelco 50/30 μm

DVB/CAR/PDMS, Manual Holder, 3pk, USA) for 30 min. The samples were analyzed using a

GC-MS chromatograph (Agilent 6890N/5973I, USA), and the temperature was maintained

isothermally at 80 °C for 3 min and increased to 350 °C at a rate of 10 °C/min for 27 min. The

retention time of each peak was used to identify the relevant chemicals based on the National

Institute of Standards and Technology database (http://webbook.nist.gov/chemistry), and the

relative peak areas were used to assess the abundance of each volatile.

Transcriptome analysis

Frozen samples of one male (already dead when intercepted by customs in November 2020) and one female (unable to eat due to intestinal disease and died in December 2020) were used to extract the total RNA. Nine tissues from the frozen samples mentioned above were used for RNA sequencing, including the nasal olfactory epithelium of the male and female, tongue of the male and female, ovary, testis, and sex-mixed samples of the brain, heart, kidney, liver, and stomach. The total RNA was extracted using TRIzol reagent (Invitrogen, USA), and the complementary DNA libraries were constructed using magnetic beads and Illumina's NEBNext Ultra RNA Library Prep Kit (Illumina, USA). The RNA sequencing was conducted on an Illumina platform, and the genome sequence of *M. javanica* (GCF_014570535.1) was used as a reference for annotations. Then, Fragments Per Kilobase of transcript per Million mapped reads (FPKM) was used to calculate the gene expression level.

The homology analysis was performed on an OBP family among 59 mammalian species. Mafft V7 (European Bioinformatics Institute, UK) was used to compare the amino acid sequences, and TBtools-II (Chengjie Chen) was used to construct the maximum likelihood tree with 1000 bootstrap runs and default settings. Meanwhile, the Time Tree of Life database (https://timetree.org/) was used to construct the species tree.

Gene cloning, expression, and purification

The open reading frame of MjavOBP1-3 was obtained by polymerase chain reaction (PCR) with specific primers that were designed by Primer Premier 5.0 (PREMIER Biosoft International, USA; Table S9) with 2 × TransTaq HiFi PCR SuperMix I (Transgen Bio, China), and they were subcloned to pET30a (Novagen, Germany) between EcoRI and XhoI (NEB, USA) using a pEASY-Basic Seamless Cloning and Assembly Kit (Transgen Bio, China). The plasmids containing the target OBPs were subsequently transformed into BL21(DE3) competent cells and induced by 1 mM isopropyl β-D-1-thiogalactopyranoside at 34 °C for 10 h. The protein

319 expression was checked using 15% SDS-PAGE and loaded onto a Ni-TED 6FF pre-packed

320 chromatography column (Sangon Bio, China) in a High-Performance Liquid Chromatography

321 system (ＡKTA pure, Sweden). The His tag was removed by Recombinant Bovine Enterokinase

322 (Sangon Bio, China).

Liquid chromatography-mass spectrometry analysis

The LC-MS was conducted according to Bortolussi et al. with a few modifications [45]. In

brief, the gel was destained, washed, alkylated, and trypsinized. The samples were then desalted

with a Zip-TipC18 column (Sigma Aldrich, USA) and eluted with a 67% acetonitrile solution

containing 2% formic acid. The protein sample was analyzed by a nano LC-ESI-Q Orbitrap

tandem mass spectrometry (MS/MS) platform that consisted of the Dionex Ultimate 3000 HPLC

RSLC nano system (Thermo Fisher Scientific, USA) and Q-Exactive Plus mass spectrometer

(Thermo Fisher Scientific) mounted to a Nanoflex ion source (Thermo Fisher Scientific). The

sample was first bound to a C18 Reversed Phase Column (Acclaim PepMap RSLC, 75 μM × 150

mm, 2 μM 100 Å; Thermo Fisher Scientific) and eluted by gradient acetonitrile (from 0% to 80%

over 80 min) containing 0.1% formic acid at 1 mL/min. The mass spectrometer was operated

under the data-dependent analysis mode (MS1 scan resolution: 70000; scanning range: 350–2000

m/z), which was followed by an MS/MS scan of the ten most abundant ions in high-energy

collisional dissociation mode (maximum ion injection time: 50 ms; collision energy: 28 eV;

dynamic exclusion: 25 s). Finally, the mass spectrum was converted into mgf format using

ProteoWizard 3.0 software [46] and imported into Mascot v. 2.6.1 (Matrix Science, UK) for

protein identification. The settings were as follows: database, uniprot; enzyme, trypsin; maximal

number of missed cleavages, 1; MS tolerance, 20 ppm; MS/MS tolerance, 0.05 Da; protein score,

$\geq 95\%$).

Fluorescence binding assays

The reaction was conducted in a 2 mL transparent quartz cell (Hellma, Germany). A mixture was constructed of 2 μM protein dissolved in 50 mM Tris·HCl (pH 7.4) and 2 μM probe 1-NPN dissolved in 50 mM Tris·HCl (pH 7.4), ligands were dissolved in methanol eight times to ensure the final content of the ligands was 2-16 μM under a Fluoromax-4 spectrofluorometer (HORIBA Jobin Yvon, USA), and they were added to this mixture. Then, 1-NPN was used as a fluorescent reporter. The excitation wavelength was set to 337 nm, and the emission spectrum was recorded between 380 and 450 nm under an emission slit of 5 nm. The dissociation constants of the competitors were calculated using the equation $K_d = IC_{50} / (1+ [1\text{-NPN}]/K_{1\text{-NPN}})$, where IC_{50} is the concentration of ligands halving the initial fluorescence value of 1-NPN, [1-NPN] is the free concentration of 1-NPN, and $K_{1\text{-NPN}}$ is the dissociation constant of the complex protein/1-NPN. GraphPad Prism 9.0 software (GraphPad by Dotmatics, USA) was used for the K_d calculation.

Behavioral tracking

The behavioral tracking assay was conducted on six individuals (S6 Table). The experiment was conducted during the activity peak of the pangolins (8 pm to 2 am) in a 1.6×1.2 m breeding room. The room was artificially divided into 48 small regions (20×20 cm) to record the visiting time of the different areas. Then, 450 μg of muscone dissolved in 40 μL propylene glycol was used as the odor stimulus, and pure propylene glycol was used as a negative control. The stimulus was added to an ethanol-cleaned plastic round box (2.4 cm in diameter, 1.2 cm in height) and placed at the corner of the room, and the behavior of the pangolins was recorded as soon as they left the resting area. A two-day interval was used between each experiment to avoid conditional reflex effects. The behaviors were recorded using an infrared camera (Cuddeback Expert, USA) under dark conditions, with at least three technical replicates for the six individuals mentioned above. After recording, the behavioral path of the pangolins was tracked by the position of their

368 noses with Tracker 6.1.5 software (https://physlets.org/tracker/trackerJS/; supplementary video 1).

369 The visiting frequency was calculated using the average staying time in each region.

Molecular modeling and docking of MjavOBP3

The three-dimensional structure of MjavOBP3 was predicted using the Alphafold2 platform

based on the amino acid sequence, and the highest score model was selected from the top five.

The topology of the protein model was optimized by Amber 20 [47], and the stereochemical

quality of the modeled protein structure was verified by the PROCHECK method. Next, Sybyl

v7.3 (Tripos, USA) was applied to calculate the interaction between the protein and ligand; the

muscone was energy minimized, then it was docked to MjavOBP3 with "auto" binding cavity

settings, and the total score was used to evaluate the binding affinity of the ligand to the target

protein. Finally, the modeling and docking results were visualized by PyMOL (DeLano Scientific

LLC).

Molecular dynamic simulation and site-directed mutagenesis

The topology fields of the ligand and protein were generated under the AMBER ff14SB

force field and general AMBER force field in Amber 20 software, respectively, and then the

complex was immersed in a cubic box with TIP3P water molecules. Then, Na^+ was added to

neutralize the entire system, and the energy of the complex was minimized by the steepest descent

and conjugate gradient methods. Next, the system was heated from 0 to 300 K and the MD

simulations were performed at constant 300 K and 1.0 Atmos for 100 ns using the PMEMD

module. The RMSD of MjavOBP3-muscone was calculated to assess the equilibrium of the MD

trajectories. The binding free energy between the residues and muscone was calculated using the

Molecular Mechanics Poisson-Boltzmann Surface Area method [48]. The hydrogen bonding sites

and the residues that contributed more than -1.00 kcal/mol to the binding free energy were

selected for the mutagenesis validation. Site-directed mutagenesis was achieved by using the Fast

Mutagenesis System Kit (TransGen, China), and the mutagenesis primers were designed by

Primer Premier 5.0 (S7 Table).

Acknowledgments

We thank all the reviewers for comments on the manuscript and Yuxin Zhou, Xiaoyan Zhu and Haoqin Ke for their help with molecular biology experiments. as well as MJEditor (www.mjeditor.com) for providing English editing services during the preparation of this manuscript.

References

1. Ralls, K. (1971). Mammalian Scent Marking: Mammals mark when dominant to and intolerant of others, not just when they possess a territory. Science 171, 443-449.

2. Thonhauser, K.E. (2013). Scent marking increases male reproductive success in wild house mice. Animal Behaviour 86, 1013-1021.

3. Gosling, L.M. (2001). Scent-marking by male mammals: cheat-proof signals to competitors and mates. In Advances in the Study of Behavior 30.169-217.

4. Thompson, C.L. (2018). Do common marmosets (*Callithrix jacchus*) use scent to communicate information about food resources? Folia Primatologica *89*, 305-315.

5. Berns, G.S. (2015). Scent of the familiar: An fMRI study of canine brain responses to familiar and unfamiliar human and dog odors. Behavioural Processes 110, 37-46.

6. Zala, S.M. (2004). Scent-marking displays provide honest signals of health and infection. Behavioral Ecology 15, 338-344.

7. Ferkin, M.H. (2010). Age affects over☐marking of opposite☐sex scent marks in meadow voles, Microtus pennsylvanicus 116, 24-31.

8. Wyatt, T.D. (2003). Pheromones and animal behaviour: communication by smell and taste, (Cambridge university press).

9. Peters, R. (1978). Scent-marking in wolves. In Wolf and Man.133-147

10. Barja, I. (2004). The importance of crossroads in faecal marking behaviour of the wolves (*Canis lupus*). Naturwissenschaften 91, 489-492.

11. Rasmussen, L.E.L. (1996). Insect pheromone in elephants. Nature 379, 684-684.

12. Clapperton, B.K. (1988). An olfactory recognition system in the ferret Mustela furo L. (Carnivora: Mustelidae). Animal Behaviour 36, 541-553.

13. Leclaire, S. (2013). Odour-based kin discrimination in the cooperatively breeding meerkat. Biology letters 9, 2012.1054.

14. White, A.M. (2003). Chemical communication in the giant panda (*Ailuropoda melanoleuca*): the role of age in the signaller and assessor. Journal of Zoology 259, 171-178.

15. Miyazaki, T. (2018). Olfactory discrimination of anal sac secretions in the domestic cat and the chemical profiles of the volatile compounds. Journal of Ethology 36, 99-105

16. Xue, Lv. (2021). Diverse phylogenomic datasets uncover a concordant scenario of laurasiatherian interordinal relationships. Molecular Phylogenetics and Evolution *157*, 107065.

17. Van Thai, N. (2009). Tapping into Local Knowledge to Help Conserve Pangolins in Viet Nam. In workshop on trade and conservation of Pangolins Native to Southeast Asia 6.163-168

18. Inskipp, T. (2005). Checklist of CITES species and annotated CITES appendices and reservations: a reference to the appendices to the Convention on International Trade in Endangered Species of Wild Fauna and Flora.

19. Choo, S.W. (2016). Pangolin genomes and the evolution of mammalian scales and

immunity. Genome Research 26, 1312-1322.

20. W.S.Challender, D. (2016). Pangolins science, society and conservation, Biodiversity of

the World: Conservation from Genes to Landscapes.

21. Lim, N.T.L. (2008). Home range, activity cycle and natal den usage of a female Sunda

pangolin *Manis javanica* (Mammalia: Pholidota) in Singapore. Endangered Species

Research 4, 233-240.

22. Zhu, J. (2017). Reverse chemical ecology: Olfactory proteins from the giant panda and

their interactions with putative pheromones and bamboo volatiles. Proceedings of the

National Academy of Sciences 114, E9802-E9810.

23. Pelosi, P. (2022). Odorant-binding proteins of mammals. Biological Reviews 97, 20-44.

24. Janssenswillen, S. (2021). Odorant-binding proteins in canine anal sac glands indicate an

evolutionarily conserved role in mammalian chemical communication. BMC Ecology and

Evolution 21, 182.

25. Hurst, J.L. (2001). Individual recognition in mice mediated by major urinary proteins.

Nature 414, 631-634.

26. Burger, B.V. "[Mammalian semiochemicals]" in The Chemistry of Pheromones and Other

Semiochemicals II. 240 231-278 (2005).

27. Burger, B. (2008). Chemical characterization of territorial marking fluid of male Bengal

tiger, *Panthera tigris*. Journal of chemical ecology 34, 659-671.

28. Harvey, S. (1989). Pattern of volatile compounds in dominant and subordinate male

mouse urine. Journal of Chemical Ecology 15, 2061-2072.

29. Ward, J., and van Dorp, D.J.E. (1981). The animal musks and a comment of their

biogenesis. Experientia 37, 917-922

30. Shirasu, M., Yoshikawa, K., Takai, Y., Nakashima, A., Takeuchi, H., Sakano, H., and
Touhara, K.J.N. (2014). Olfactory receptor and neural pathway responsible for highly
selective sensing of musk odors. Neuron 81, 165-178.

31. Brophy, J., Cavill, G., McDonald, J., Nelson, D., and Plant, W.J.I.b. (1982). Volatile
constituents of two species of Australian formicine ants of the genera Notoncus and
Polyrhachis. Insect Biochemistry 12, 215-219.

32. Mitaka, Y., Matsuyama, S., Mizumoto, N., Matsuura, K., and Akino, T. (2020). Chemical
identification of an aggregation pheromone in the termite Reticulitermes speratus.
Scientific Reports 10, 7424.

33. Gong, D.-P., Zhang, H.-J., Zhao, P., Xia, Q.-Y., and Xiang, Z.-H.J.B.g. (2009). The
odorant binding protein gene family from the genome of silkworm, *Bombyx mori*. BMC
Genomics 10, 1-14.

34. Lobel, D., Strotmann, J., Jacob, M., and Breer, H. (2001). Identification of a third rat
odorant-binding protein (OBP3). Chem Senses 26, 673-680.

35. Krishna, N.S., Getchell, M.L., Margolis, F.L., and Getchell, T.V. (1995). Differential
expression of vomeromodulin and odorant-binding protein, putative pheromone and
odorant transporters, in the developing rat nasal chemosensory mucosae. Neurosci Res 40,
54-71.

36. Ohno, K., Kawasaki, Y., Kubo, T., and Tohyama, M. (1996). Differential expression of
odorant-binding protein genes in rat nasal glands: implications for odorant-binding protein
II as a possible pheromone transporter. Neuroscience 71, 355-366.

37. Pes, D., Mameli, M., Andreini, I., Krieger, J., Weber, M., Breer, H., and Pelosi, P. (1998).
Cloning and expression of odorant-binding proteins Ia and Ib from mouse nasal tissue.
Gene 212, 49-55.

38. Roper, T., Shepherdson, D., and Davies, J.J.B. (1986). Scent marking with faeces and anal

secretion in the European badger (Meles meles): seasonal and spatial characteristics of

latrine use in relation to territoriality. Behaviour 97, 94-117.

39. Kusuda, S., Hoson, O., Nakaya, Y., Ogura, G., Oshiro, S., Takara, J., Matsuda, A., Doi,

O., Nagamine, T., and Murata, K.J.M.s. (2010). Induced estrus in female small Asian

mongooses (*Herpestes javanicus*) for the purpose of controlling invasive alien species in

Okinawa Island. Mammal Study *35*, 217-219.

40. Dorrigiv, I., Hadian, M., and Bahram, M. (2023). Comparison of volatile compounds of

anal sac secretions between the sexes of domestic dog (*Canis lupus familiaris*). Veterinary

Research Forum, 14, 169-176.

41. Drea, C.M., Vignieri, S.N., Kim, H.S., Weldele, M.L., and Glickman, S.E. (2002).

Responses to olfactory stimuli in spotted hyenas (Crocuta crocuts): II. Discrimination of

conspecific scent. Journal of Comparative Psychology 116, 342-349.

42. Apps, P., Mmualefe, L., and McNutt, J.W.J.J.o.C.E. (2012). Identification of volatiles

from the secretions and excretions of African wild dogs (Lycaon pictus). Journal of

Chemical Ecology 38, 1450-1461.

43. Kerley, L.L., and Salkina, G.P. (2010). Using Scent☐Matching Dogs to Identify

Individual Amur Tigers from Scats. The Journal of Wildlife Management 71, 1349-1356.

44. Campbell-Palmer, R., and Rosell, F. (2011). The importance of chemical communication

studies to mammalian conservation biology: A review. Biological Conservation 144,

1919-1930.

45. Bortolussi, G., Codarin, E., Antoniali, G., Vascotto, C., Vodret, S., Arena, S., Cesaratto,

L., Scaloni, A., Tell, G., Muro, A.J.C.d., et al. (2015). Impairment of enzymatic

antioxidant defenses is associated with bilirubin-induced neuronal cell death in the

cerebellum of Ugt1 KO mice. Cell Death & Disease 6, e1739.

46. Chambers, M.C., Maclean, B., Burke, R., Amodei, D., Ruderman, D.L., Neumann, S., Gatto, L., Fischer, B., Pratt, B., Egertson, J., et al. (2012). A cross-platform toolkit for mass spectrometry and proteomics. Nature Biotechnology 30, 918-920.

47. D.A. Case, K.B., I.Y. Ben-Shalom, S.R. Brozell, D.S. Cerutti, T.E. Cheatham, III, V.W.D. Cruzeiro, (2020). AMBER 2020 software.

48. Feig, M., Onufriev, A., Lee, M.S., Im, W., Case, D.A., and Brooks III, C.L.J.J.o.c.c. (2004). Performance comparison of generalized born and Poisson methods in the calculation of electrostatic solvation energies for protein structures. Journal of computational chemistry 25, 265-284.

Table

Table 1. Sunda pangolin lipocalins.

Protein Family	Gene Name	NR Annotation	Length (nt)	Length (aa)	Molecular Weight
Odorant binding protein (OBP)	MjavOBP1	allergen Bos d 2-like isoform X1	522	173	19.34 kD
	MjavOBP2	odorant-binding protein 2b-like	519	172	19.40kD
	MjavOBP3	odorant-binding protein-like	522	173	19.15 kD
Fatty-acid-binding protein (FABP)	MjavFABP1A1	fatty acid-binding protein, liver	384	127	14.09 kD
	MjavFABP1A2	fatty acid-binding protein, liver	402	133	14.68 kD
	MjavFABP2	fatty acid-binding protein, intestinal	399	132	15.29 kD
	MjavFABP3	fatty acid-binding protein, heart	402	133	14.92 kD
	MjavPMP2	myelin P2 protein	402	133	14.87 kD
	MjavFABP9	fatty acid-binding protein 9-like	390	129	14.79 kD
	MjavFABP4	fatty acid-binding protein, adipocyte	399	132	14.89 kD
	MjavFABP5A1	fatty acid-binding protein, epidermal	408	135	15.17 kD
	MjavFABP5A2	fatty acid-binding protein, epidermal	408	135	15.08 kD
	MjavFABP6	gastrotropin isoform X2	387	128	14.68 kD
	MjavFABP7	fatty acid-binding protein, brain	399	132	14.90 kD
	MjavFABP12	fatty acid-binding protein 12	462	153	17.17 kD
Retinol binding protein (RBP)	MjavRBP1	retinol-binding protein 1	408	135	15.80 kD
	MjavRBP2	retinol-binding protein 2	282	93	10.73 kD
	MjavRBP4	retinol-binding protein 4	594	197	22.54 kD
	MjavRBP5	retinol-binding protein 5	408	135	15.91 kD
	MjavRBP7	retinoid-binding protein 7 isoform X2	384	127	14.71 kD
Beta-lactoglobulin (BLG)	MjavBLG1	beta-lactoglobulin-1-like, partial	543	180	20.31 kD
	MjavBLG2	lactoglobulin-2	543	180	20.30 kD
Cellular retinoic acid-binding protein (CRABP)	MjavCRABP1	cellular retinoic acid-binding protein 1	414	137	15.57 kD
	MjavCRABP2	cellular retinoic acid-binding protein 2	417	138	15.72 kD
Epididymal-Specific Lipocalin	MjavLCN6	epididymal-specific lipocalin-6	498	165	17.88 kD
	MjavLCN9	epididymal-specific lipocalin-9 isoform X2	498	165	18.88 kD
Apolipoprotein D	MjavAPOD	apolipoprotein D	570	189	21.47 kD
Orosomucoid	MjavORM1	alpha-1-acid glycoprotein-like isoform X1	669	222	24.76 kD
Prostaglandin D2 synthase	MjavPTGDS	prostaglandin-H2 D-isomerase	576	191	21.18 kD
Von Ebner's gland protein	MjavLCN1	lipocalin-1-like	552	183	19.80 kD
Neutrophil gelatinase-associated lipocalin	MjavLCN2	Neutrophil gelatinase-associated lipocalin	600	199	22.91 kD
Salivary lipocalin protein	MjavSAL1	minor allergen Can f 2-like	555	184	20.41 kD
Human-specific lipocalin	MjavLCN15	lipocalin-15 isoform X3	492	163	17.85 kD
Complement component 8	MjavC8G	complement component C8 gamma chain isoform X2	831	276	30.92 kD

Table 2. Compounds used for functional validation of odorant binding proteins (OBPs)

in Sunda pangolins.

Compound	Source	Cas Number	Molecular Structure	Purity	Company
Muscone	Secretion	541-91-3		97.0%	Aladdin
Phenol	Secretion	108-95-2		99.0%	Aladdin
2-Undecanone	Secretion	112-12-9		99.0%	Aladdin
3-Undecanone	Secretion	2216-87-7		≥85%	Aladdin
Oleic acid	Secretion	112-80-1		98.0%	Aladdin
Indole	Secretion feces	120-72-9		99.0%	Aladdin
2-Piperazinone	Secretion feces	5625-67-2		98.0%	Aladdin
Tridecane	Feces	629-50-5		98.0%	Aladdin
β-farnesene	Feces	18794-84-8		90.0%	Aladdin
3-methylindole	Feces	83-34-1		98.0%	Aladdin
Dimethyl trisulfide	Feces	3658-80-8		98.0%	Aladdin
Palmitic acid	Feces urine	57-10-3		97.0%	Aladdin
δ-Dodecalactone	Feces urine	713-95-1		98.0%	Aladdin
Phenethyl alcohol	Feces urine	60-12-8		99.0%	Aladdin
γ-Undecanolactone	Urine	104-67-6		98.0%	Aladdin
Myristic acid	Urine	544-63-8		98.0%	Aladdin
Benzaldehyde	Urine	100-52-7		99.5%	Aladdin
2-Isobutyl-3-methoxypyrazine	Habitat Volatiles	24683-00-9		99.0%	Sigma
1-Octen-3-one	Habitat Volatiles	4312-99-6		95.0%	Aladdin
Hexanal	Habitat Volatiles	66-25-1		99.0%	Aladdin
β-Pinene	Habitat Volatiles	18172-67-3		98.0%	Rhawn
Geraniol	Habitat Volatiles	106-24-1		98.0%	Sigma

α-pinene	Habitat Volatiles	80-56-8		98.0%	Sigma
Myrcene	Habitat Volatiles	123-35-3		99.0%	Aladdin
Undecane	Habitat Volatiles	1120-21-4		97.0%	Aladdin
Acetophenone	Habitat Volatiles	98-86-2		99.5%	Aladdin
Citral	Habitat Volatiles	5392-40-5		97.0%	Aladdin
1-octen-3-ol	Habitat Volatiles	3391-86-4		98.0%	Aladdin
Nonanal	Habitat Volatiles	124-19-6		95.0%	Sigma
6-Methyl-5-hepten-2-one	Habitat Volatiles	110-93-0		99.0%	Sigma
3-Octanone	Habitat Volatiles	106-68-3		98.0%	Aladdin
Octanal	Habitat Volatiles	124-13-0		99.0%	Sigma
Eugenol	Habitat Volatiles	97-53-0		99.0%	Sigma
Pentadecane	Habitat Volatiles	629-62-9		98.0%	Aladdin
3-Carene	Habitat Volatiles	13466-78-9		≥90%	Aladdin
Linoleic acid	Habitat Volatiles	60-33-3		95.0%	Aladdin
β-Lonone	Habitat Volatiles	14901-07-6		96.0%	Sigma
Citronellal	Habitat Volatiles	106-23-0		96.0%	Heowns
3-Phenylpropionic acid	Habitat Volatiles	501-52-0		99.0%	Aladdin

555 **Figure 1**. GC-MS analysis. Total ion chromatogram of the feces, urine and anal gland secretion

556   of adult *M. javanica* on a polyethylene glycol column (Table S1).

Figure 2. Transcriptome and Homology analysis. (A) Expression profiles of lipocalins in

M.javanica. (FN: female nasal olfactory epithelium, MN: male nasal olfactory epithelium, FT:

female tongue, MT: male tongue, O: ovary, T: testis, H: mixed sample of heart; K: mixed

sample of Kidney; L: mixed sample of liver, S: mixed sample of stomach). The left side of the

figure is a phylogenetic tree constructed using neighbor-joining for pangolin lipocalins, and

the right side is the corresponding gene name (Table S3). (B) Homology analyses of odorant

binding proteins (OBPs) in 59 mammalian species. The gene tree constructed for odorant

binding proteins using the maximum likelihood method is divided into three clades,

corresponding to the OBP1-3 subfamily (Fig. S1, Table S4). The right side of the figure

shows the number and location number of different types of OBPs in different color located

on the chromosome.

Figure 3. Sequencing analysis and ligand binding ability determination of three MjavOBPs.

(A-C) Mass Spectrum of peptides from three MjavOBPs are shown, respectively. The

fragments are reported in different color depending on peptides present in *M. javanica* and

corresponding b and y ion series. (D-F) Binding curves of all MjavOBPs to different ligands.

The ligand names are shown on the right of the curves with different colors, and binding

affinity data are listed in Table S6.

Figure 4. Preferential behavioral responses of female and male pangolins to muscone.

(A-D) An individual test pangolin is introduced into the room, can freely enter and exit from the sleeping place (gray area), and muscone odor or control odor (propylene glycol) is placed in b8, and the behavioral trajectories of pangolins to control odors and muscone were recorded separately, ★ represents the entrance, control odor is in cyan, and muscone is in yellow, blue

502　dots represent males, red dots represent females; (E-H) The residence time of pangolins in

503　different areas after placing control odor or muscone. Blue histograms represent males and red

504　histograms represent females. Bars indicate the mean time that a pangolin spent investigating

505　each test sample within a 5-min period, ±S.E.M. Asterisks indicate significant differences

506　between two samples (paired Student's t-test, **$P < 0.01$, n=10).

Figure 5. Prediction of key sites in the MjavOBP3-Muccone complex. (A) The 3D structure

of MjavOBP3 presents a barrel-like structure, mainly composed of 8 β-sheets (β1-β8) and 1 α-

helix; (B) The predicted binding pocket is shown in green. The ligand Muscone (dark pink) is

tightly wrapped by the binding pocket; (C) 3D visual analysis of the binding mode of

MjavOBP3 and muscone. Important amino acid residues and muscone are shown as baseballs,

in which hydrogen bonds are represented by pink dotted lines, and molecular distances are

calculated; (D) The root-mean-square deviation (RMSD) curve of the MjavOBP3–Muscon

complex. (E) Protein-ligand interaction diagram of MjavOBP3 bound to muscone at steady

state; (F) Total energy contribution of each amino acid residue in the MjavOBP3–muscone

complexes. The residues with a total energy contribution of more than −1.00 kcal/mol are

marked.

Figure 6. Verification of key sites of MjavOBP3-muscone complex. (A) Target site in the

MjavOBP3 gene (wild type) and results obtained from the mutagenesis of MjavOBP3 through

site-directed mutagenesis, and bases A, T, C and G are represented by green, pink, blue and

black respectively, and the eight mutation positions are represented by different color; (B)

Binding curves of MjavOBP3 mutants to muscone. The lines (different colors) represent the

different mutants, and the line (black) represents the wild type (WT); (C) Comparison of the

binding affinities, which were indicated by K d value, of MjavOBP3 WT and mutants to

muscone, respectively. The significant differences were compared using one-way analysis of

variance (ANOVA) with Tukey's honestly test and indicated by different lowercase letters (p-

value < 0.01, n =3); (D) Comparison of the amino acid sequence similarity of MjavOBP3

homologous proteins in Laurasia. The sequence similarity is expressed by per ident. The stars

(*) show the same amino acids in all proteins. Conserved cysteine amino acid sites involved

in the formation of disulfide bridges are shown in light blue, while Key binding sites in

MjavOBP3 for Muscone are shown in light pink

Supporting information

S1 Fig. Mammalian OBP gene tree based on Maximal likelihood method

S2 Fig. SDS/PAGE of MjavOBP1-3 before and after purification

S3 Fig. Binding curve of 1-NPN to all MjavOBP1-3

S4 Fig. Ramachandran plot of MjavOBP3 model

S5 Fig. Binding curve of 1-NPN to MjavOBP3 WT and Mutants

S6 Fig. Ethics statement

S1 Table (Microsoft Excel format). Determination of VOCs in urine, feces and anal gland secretions in pangolins.

S2 Table. (Microsoft Excel format). The expression levels of lipocalins in Sunda pangolins from different tissues.

S3 Table. (Microsoft Excel format). List of identified mammalian OBP genes.

S4 Table. (Microsoft Excel format). [IC]50 and Kd values between three MjavOBP1-3 and candidate ligands.

S5 Table. (Microsoft Excel format). Binding free energy of MjavOBP3-muscone complex.

S6 Table. (Microsoft Excel format). Individual information used for experimental pangolins.

S7 Table. (Microsoft Excel format). Primers used for plasmid construction of MjavOBP1-3 and mutants.

S1 Video (separate files). Pangolin activity videos converted into tracks.

S2 Video (separate files). Video of male pangolin's behavioral response to control odor.

706   **S3 Video (separate files). Video of male pangolin's behavioral response to muscone.**

707   **S4 Video (separate files). Video of female pangolin's behavioral response to control odor.**

708   **S5 Video5 (separate files). Video of female pangolin's behavioral response to muscone.**