Diet predicts performance trade-offs along Atlantic salmon ontology

Tutku Aykanat[1*], Jan Arge Jacobsen[2], Kjetil Hindar[3]

1. Organismal and Evolutionary Biology Research Program, Faculty of Biological and

Environmental Sciences, PO Box 56, FI-00014 University of Helsinki, Finland.

tutku.aykanat@helsinki.fi

2. Faroe Marine Research Institute, Nóatún 1, FO-100 Tórshavn, Faroe Islands. Council

3. Norwegian Institute for Nature Research (NINA), NO-7485 Trondheim, Norway.

kjetil.hindar@gmail.com

[*] Corresponding author: tutku.aykanat@helsinki.fi

Running page head: Age dependent diet-performance trade-offs

Abstract

Understanding the mechanistic basis of life history divergence is fundamental for predictive ecology. Variation in resource use impacts life history divergence and provides a framework for studying life history trade-offs. Contrary to the theory, empirical evidence does not support that resource allocation alone mediates life history divergence in the wild, which subsequently suggests a pivotal role for resource acquisition, but this has not been empirically demonstrated in free-living populations. Here, we sampled wild Atlantic salmon (*Salmo salar*) from different age groups during their marine feeding stage and hypothesized that prey taxa (consumption of crustaceans *versus* fish) has an age-dependent effect on the condition factor. We found that salmon in their first year at sea had a higher condition factor when foraging on larger numbers of crustaceans and a lower condition factor when foraging on larger numbers of fish, and *vice versa* in older sea age groups. Provided that condition factor is a predictor of age-at-maturity, our results show that resource acquisition may be an important process shaping life history variation in Atlantic salmon via modulating age-dependent performance trade-offs. Our results also emphasize the importance of bottom-up trophic cascades for maturation age and help to better predict the maturation dynamics of wild populations in the marine environment.

Keywords: life history variation, resource acquisition, age at maturity, ontogenetic diet shift, trade-offs, Atlantic salmon.

1. INTRODUCTION

An important focus of evolutionary biology has been to identify and study the underlying mechanisms of life history trade-offs in wild populations (Reznick et al. 2000, Stearns 2000, Leroi 2001, Roff 2002, Braendle et al. 2011). By characterizing the physiological basis of the trade-offs between maturation timing and survival, the response to selection can be predicted more accurately (Roff 2002, Stearns 2011). Competitive resource allocation —the allocation of available energy to support various biological processes such as growth and reproduction—, has been suggested as the major contributor to the trade-offs dynamics in many life history models, such as in a trade-off between investing to current and future reproductive success (reviewed by Flatt et al. 2011). However, the expectation of negative phenotypic (or genetic) correlations between survival and other fitness traits lack confirmation in natural populations (Haave-Audet et al. 2022, Chang et al. 2024). Essentially, traditional trade-off models do not account for variation in resource acquisition, —the processes associated with obtaining energy from the environment to support biological processes—. Resource acquisition, however, could not only explain the observed positive phenotypic (or genetic) correlation between fitness components (Van Noordwijk & de Jong 1986, Houle 1991, Snell-Rood et al. 2015), but also provide a realistic framework to assess the mechanistic basis and the role of life history trade-offs in natural systems. For example, a recent meta-analysis in non-human animals found that differences in resource acquisition were more important than differences in resource allocation in mediating the relationship between behavior and fitness (Haave-Audet et al. 2022).

Atlantic salmon (*Salmo salar*) is a good model to study life history evolution and how life history variation is maintained (Schaffer 2003). Atlantic salmon is an anadromous species in which juveniles migrate to resourceful marine environment to enhance growth and subsequently increase reproductive success. The number of years that Atlantic salmon stay at sea prior to first reproduction (henceforth,

sea-age at maturity) and fecundity are coupled because fish size and, consequently, fecundity increases

with age. By extension reproductive success and mortality are inherently negatively correlated due to

deceasing overall survival probability of salmon by staying longer at sea before reproduction. The

maintenance of trait variation represents a classic case of fitness trade-off between reproductive success

and survival: longer times at sea increase fitness via a size-dependent increase in reproductive success

but at the expense of a higher risk of mortality prior to reproduction (Fleming & Einum 2010, Mobley

et al. 2021). Subsequently, the species also represents a wild model to study the potential contribution

of resource acquisition or allocation in mediating the optimal age structure, which also has substantial

practical value for predicting the dynamics of the demographic structure of the populations. Whether

resource acquisition is involved in life history trade-offs has yet to be empirically shown in wild

systems.

Sea age at maturity is a threshold trait (Thorpe et al. 1998) with a strong genetic component (Barson et

al. 2015, Sinclair-Waters et al. 2020), and environmental factors also play a central role in shaping the

trait variation (Jonsson & Jonsson 2004, Jonsson et al. 2012). In the marine environment, Atlantic

salmon achieve high growth rates and accumulate fat reserves to support energetically expensive

maturation and reproduction (Jonsson & Jonsson 2011). As such, variation in resource acquisition

between individuals may play a central role in shaping sea age at maturity via its effect on lipid

content, which is suggested as the underlying liability trait of maturation (Rowe et al. 1991, Thorpe et

al. 1998). The changes in lipid content of individuals may be mirrored by changes in condition factor,

hence condition factor can predict maturation (Herbinger 1987, Herbinger & Friars 199, Duston &

Saunders 1999, Peterson & Harmon 2005, Jonsson et al. 2012, Jonsson et al. 2013, Debes et al. 2021).

For example, Herbinger (1987) found by studying individually marked salmon, that maturation in the

autumn was associated with a high growth rate during the previous winter, and that this growth was

attained by weight rather than length increase, resulting in a higher condition factor. Concordantly, it

has been suggested that food web changes in the ocean may have a substantial impact on the

maturation age of salmon populations (Friedland et al. 2005, Vollset et al. 2022), and salmon

populations may show evolutionary responses to changes in food availability (Czorlich et al. 2022). At

sea, the diet of sub-adult Atlantic salmon (the marine stage prior to maturation) mainly consists of

crustaceans and fish, with large spatiotemporal variation (Jacobsen & Hansen 2001, Rikardsen &

Dempson 2010). These two major taxonomic groups have fundamentally different properties as

potential prey, such as size, behavior, and availability, which ultimately require different foraging

strategies to utilize them (Mittelbach & Persson 1998, Ferry-Graham et al. 2002, Sanchez-Hernandez et

al. 2019). Feeding also has an ontogenetic component: crustaceans-based diet in early marine phases

switches to a more piscivorous diet as individuals grow larger. However, there is a considerable

overlap in the distribution of prey groups between different sea ages (Jacobsen & Hansen 2001).

Therefore, age-dependent dynamics of diet intake and how it affects condition factor, a strong predictor

of maturation timing (Herbinger & Friars 1991, Jonsson et al. 2012), is unclear in wild Atlantic salmon.

The ontogenetic diet shift is an adaptive trait in which an optimal feeding strategy is attained to supply

the increasing energetic demands of a larger body, and is mostly linked to changes in habitat and life-

stage transitions (Villéger et al. 2017, Sanchez-Hernandez et al. 2019). However, few studies have

quantified inter-individual variation in ontogenetic diet dynamics (Mittelbach & Persson 1998,

Sanchez-Hernandez et al. 2019), with no assessment of potential performance trade-offs in free-living

populations. An ontogenetic diet shift in the marine environment implies that the net adaptive value of

energetic gain obtained from a particular feeding regime (e.g., diet type) changes as a function of

ontogeny. As such, acquiring optimal resource type increases the net energy gain, which may facilitate

lipid accumulation and condition factor (Jonsson et al. 2013), and subsequently probability of maturing

(Herbinger & Friars 1991, Rowe et al. 1991, Thorpe et al. 1998). Therefore, resource acquisition may

be suggested as a functional ecological process explaining the dynamics of age at maturity, by

demonstrating an age dependent relation between performance (condition factor) and diet composition

(Haave-Audet et al. 2022, Chang et al. 2024).

Fig. 1. Conceptual model that links performance (condition factor) to diet type (crustacean versus fish)
of sub-adult Atlantic salmon during the marine feeding stage.

Based on the above predictions, we hypothesized there is an ontogeny-dependent performance trade-off

for consuming different diet items (Fig. 1). Specifically, we hypothesize that eating smaller food items

early in the marine stage is an optimal feeding strategy for smaller-sized salmon, and increases their

probability of maturation in the following year by facilitating lipid depositions, which may be predicted

by the condition factor (Herbinger & Friars 1991, Todd et al. 2008). Lipids are essential, for example,

during vitellogenesis (egg production), which comprises a substantial amount of energy expenditure of

females during the reproduction period (Jonsson & Jonsson 2003), as well as for other energetically

expensive migration and reproductive activities (Jonsson & Jonsson 2011). Concordantly, lipid content

has been suggested as the underlying, liability trait that triggers the developmental switch to initiate the

maturation the following year (Thorpe et al. 1998). Likewise, we hypothesize that switching to larger

food items is more beneficial to older, larger-sized salmon, leading to higher net energy gain, also to

support future reproduction activities. Such age dependent trade-off between performance (condition

factor) and diet composition would provide an ecological explanation of how a life history trait (age at

maturity) is mediated by resource acquisition variation (marine diet composition). As such, we

quantified the diet of wild Atlantic salmon sub-adults sampled in the oceanic feeding ground,

quantified their stomach contents (consisting of crustaceans and fishes, Jacobsen and Hansen 2001),

and compared the age-dependent changes in condition factor in relation to diet.

2. MATERIALS & METHODS

2.1. Study system

Atlantic salmon used in this study were originally described by (Jacobsen & Hansen 2001), and the

final data were identical to Aykanat et al. (2024) except that we retained N=68 individuals that were

filtered out in the prior study due to missing genotype information in two focal life history loci. Briefly,

the dataset comprises 1604 individuals collected from the Northeast Atlantic Ocean in their ocean

feeding ground North of the Faroe Islands, of which 178, 995, and 431 are from 1SW, 2SW, and 3SW

sea-age groups, respectively (SW, "sea-winter" is the number of winters a fish has been in the marine

environment and is equivalent to the sea age). Samples were collected at 55 locations during fishing

periods in autumn (November and December) and winter (February and March) in 1992-93 and 1993-

94 (Fig. S1). The fish were genetically assigned to large-scale reporting groups (Ireland and the UK,

Southern Norway, and Northern Norway; the latter encompassing fish assigned to Barents Sea

drainages (O'Sullivan et al. 2022), which descibes broad phylogenetic groups associated with large

geographical regions. Compared to the original study (Jacobsen & Hansen 2001), which comprised of

N=1765 wild individuals from the 1992-93, and 1993-94 fishing periods, we excluded 161 individuals,

of which N=34 did not have any material to genotype (i.e. no scales), N=9 did not have sex

information, N=15 had sea age larger than 3SW, and 103 were not assigned to the reporting groups

indicated above (out of 103, N=88 was assigned to North American phylogenetic groups). Individual

salmon had their fork lengths measured in centimeters, and gutted, and gutted fish were weighed in

kilograms to two decimal places. The gut contents were identified to taxon level and enumerated. Fish

scales were used to determine sea age (via scale reading), and for DNA extraction and genotyping. The

fish sampled in this study were sub-adults with no phenotypic indications of maturation. An

independent, earlier study from the Faroese feeding grounds indicates that 80% of individuals are likely

to mature in the following autumn, but these estimates are based on limited number of tag recapture

studies, or based on assessing the maturation probability via hormonal expression using hatchery reared

salmon as the baseline (Youngson & McLay 1984), hence may not be reliable.

2.2. Data analysis

We employed a two-step multi-model inference approach using linear mixed models, using condition

factor as the response variable. Condition factor is a strong predictor of lipid content in Atlantic salmon

(Herbinger & Friars 1991, Todd et al. 2008) which is suggested as the liability trait of the maturation

timing (Thorpe et al. 1998). In this framework, we first identify the most parsimonious null model

structure by comparing the fit of the models parameterized with different combination of non-focal

variables, and their interactions. The global null model structure is as follow:

Global null model = CF ~ μ + fishing period + Season x Sea Age x Sex x Length$_{age.std}$ + ε$_{reporting\ group}$ +

ε$_{location}$ + ε$_{error}$

Whereby, μ is the intercept and ε$_{error}$ is the residual errors. The condition factor (CF) is Fulton's

condition factor K, (K=10^5 w / l^3), in which w is the gutted weight in kilograms divided by 0.9, to

account for gut weight, and l is fork length in cm (Fig. S2). Fulton's condition factor is a common

condition index that has been a sufficient standard in physiological studies in fishes (Stevenson &

Woods 2006). The condition factor is based on the weight obtained from gutted fish; therefore, it

reflects the condition factor associated with body weight excluding the gut weight and visceral fat and

gonadal tissue weights. This is pertinent since subcutaneous adipose tissue and musculature lipids have

been suggested as the main energy reserves during the spawning period (Jonsson et al. 1997).

The model included season, fishing period, sea age, sex, and age-standardized length (Length$_{age.std}$) as

fixed effects. Spatial variation in sampling effort was accounted for by including sampling location as a

random effect (ε$_{location}$, N=55). Likewise, variation due to geographical origin of individuals was

accounted for by including reporting groups (ε$_{reporting\ group}$, N=3) as random terms. Length$_{age.std}$ was

scaled to have a mean of zero and a standard deviation of one to improve convergence. All four-way

combinations of season, sex, sea age, and age-standardized length and their reduced models were fitted

and compared. Models were fitted by parameterizing sea-age both as a continuous or factorial variable

(Table 1). All models were then compared using corrected Akaike Information Criteria (AICc), which

corrects for bias for small sample sizes. The model with lowest AICc value is considered as the most

parsimonious null model and used in the following analyses.

Next, on top of the most parsimonious null model, we fitted models that also include combinations of

focal variables ($Diet_{crus}$ and $Diet_{fish}$) and their interaction with sea age, the fit of which was then

compared using AICc. These models reflected hypotheses explaining the diet type effect on condition

(Table 2). The global model structure is as follow:

Global model: CF ~ terms as in the most parsimonious null model + Sea Age x ($Diet_{crus}$ + $Diet_{fish}$)

Whereby, $Diet_{crus}$ and $Diet_{fish}$ are log-normalized number of diet items in the stomach. Both crustacean

and fish diets had highly left-skewed distributions with many zero entries (i.e., empty stomach, Fig.

S3). To avoid undefined zero values after log transformation, we added 1 to the number of diet

variables. As above, sea-age was modelled both as a continuous or categorical variables.

All models were run using the glmmTMB package (version 1.1.5, Brooks et al. 2017) in R version

4.1.0 (R Core Team 2019), using a Gaussian error structure. Null model selection and AICc

calculations was performed using MuMin package (version 1.48.4, Barton et al 2024). Marginal means

for genotype effects and their confidence intervals were calculated using the emmeans package, and

significance was assessed by t-test (version 1.8.5, Lenth, 2023). Model diagnostics were performed

using the DHARMa package in R (version 0.4.6, Hartig 2022).

2.3. Post-hoc analyses

We implemented a number of post-hoc analysis in which we revise original parameterization of terms

to further break down the structure of the diet effect on condition.

2.3.1. The effect of different taxonomic groups within crustaceans.

We evaluated whether any taxonomic groups within crustaceans drive the relationship associated with

the crustaceans diet, by partitioning crustaceans diet into lower taxonomic groups, provided that the

numbers within a specific taxon were sufficiently large. As such, we implemented the most

parsimonious model structure, but by replacing $Diet_{crus}$ term with either $Diet_{Hyperiid.amphipods}$ or

$Diet_{Euphausiids}$, which are the log-normalized number of hyperiid amphipods (*Themisto* spp.) and

euphausiids in the diet, respectively. These two taxa comprise the majority of the crustaceans in the

stomach (Table S1), and were substantially different in size, with the latter being larger than the former

(Table S1). We did not implement the same approach in fishes due to small number of prey items per

species (Table S1).

2.3.2.Partitioning the underlying resource acquisition processes

The relationship between the condition factor and number of prey items might be driven by different

ecological processes associated with foraging, such as active foraging frequency or foraging efficiency

(Aykanat et al. 2024). We assessed the contribution of these two possibilities by repeating the analysis

after restructuring diet variables to account for these factors. As such, the presence or absence of prey

was used as a proxy for active foraging frequency, using the model structure described above, but

replacing the diet variable with a Boolean parameter, for crustaceans and fish in two separate binomial

models. Using the data with which diet was present, log-transformed number of diet items was

modelled as a proxy for foraging efficiency (when actively foraging for prey), for crustaceans and fish

in two separate models.

2.3.3. Comparing the importance of weight vs number of diet in the stomach

219 The number and total weight of the diet items in the stomach were highly correlated (Fig. S4); thus, any

220 effect due to the number of diet items could be attributed to diet weight. To assess whether the number

221 or total mass of the diet better explain the change in the condition factor, we implemented the same

222 model as above but used total diet weight as a factor instead of the number of prey items acquired.

3. RESULTS

3.1. Null model structure and parametrization of sea age

The most parsimonious null model included fishing period, season, length$_{age.std}$, sex, and sea age as

parameters, and second order interactions between length$_{age.std}$, sex, and sea age, (Table S2). Notably, a

model that parameterizes sea age categorically was substantially better than a model with continuous

parametrization (ΔAICc = 67.18, Table S3). However, we modelled sea age continuously when

interacting with diet variables. Such a model was more parsimonious than a model that parameterizes

sea age continuously (ΔAICc = 87.78) or categorically (ΔAICc = 4.01) across all the interactions. In all

three models, main results regarding age-dependent diet association on condition factor was

qualitatively similar (data not shown).

Fig. 2. Marginal effect of the number of crustaceans (a) and fish (b) prey items (log scale) on condition
factor in Atlantic salmon sub-adults in the marine environment. Dashed line indicates a zero slope, no
relation between the number of diet items and condition factor. Error bars indicate 95% confidence
intervals.

3.2. Age dependent diet effect on condition factor

Our analysis supported the hypothesis that suggests an age dependent diet effect on condition factor

(Table 1). In this model, the number of diet items in the stomach had a significant age dependent

association with condition factor, but the direction of the age effect was in contrasting for crustaceans

and fish diets. In other words, the age dependent slope of the relation was negative for crustaceans diet

(for crustaceans diet by sea age interaction: t-test, t_{1585} = -3.208, p = 0.001, Fig. 1a), but it was positive

for fish diet (for fish diet by sea age interaction t_{1585} = 2.088, p = 0.037, Fig. 1b). Within 1SW age

group, the number of crustaceans in the diet was positively associated with condition factor (t_{1585} =

1.901, p = 0.058), but the relation was increasingly negative as the age of salmon progresses (for 2SW:

t_{1585} = -1.794, p = 0.073, for 3SW: t_{1585} = -3.572, p < 0.001. Fig. 2a). The number of fish in the diet

did not affect the condition factor in the 1SW age group (t_{1585} = -0.702, p = 0.48), but the relation was

increasingly positive as the age of Atlantic salmon progresses (for 2SW: t_{1585} = 2.177, p = 0.030, for

3SW: t_{1585} = 3.367, p < 0.001, Fig. 2b, Fig. S5). The scenario where crustaceans and fish diets were

modelled age dependently with continuous parameterization of sea age significantly outperformed any

other alternative hypotheses (Table 1).

3.3. Age dependent sex and length effects on condition factor

Sex and length also exhibited age-dependent patterns, which had an apparent non-linear dependency

with sea age (Fig. 3). Notably, there was a substantial reversal of sex-dependent changes in the

condition factor between the 1SW and 2SW age groups (Fig. 3a). Males had significantly higher

condition factors in the 1SW age group (t_{1585} = 2.693, p = 0.007) and significantly lower condition

factors in the 2SW age group (t_{1585} = -5.254, p < 0.001), and there was no difference between sexes in

the 3SW age group (t_{1585} = 0.026, p = 0.98, Fig. 3a). Length was positively associated with the

condition factor positively in the 2SW age group (t_{1585} = 5.040, p <0.001) but the association was not

significant in other age groups (Fig. 2b). As expected, winter was associated with a lower condition

factor (t_{1585} = -8.123, p < 0.001), and there was a marginal fishing-period effect ($_{1585}$ = -1.989, p <

0.047, see also Table S2).

Fig. 3. Marginal effects of sex (a) and length (b) on condition factor in Atlantic salmon sub-adults in
the marine environment. Dashed line in b indicates a zero slope, no relation between (age standardized)
length of salmon and condition factor. Error bars indicate 95% confidence intervals.

3.4. Taxon specificity of crustacean diet effect

The crustacean diet in the stomach consisted of a few major taxonomic groups, the majority of which

were hyperiid amphipods (~93% and 69% by number and weight, respectively, and mostly *Themisto*

species) and euphausiids (krill, ~ 6% and 15% by number and weight, respectively, Table S1). When

the number of diet specific to these taxa was separately parameterized in the model, instead of

crustaceans as a whole, hyperiid amphipods, but not krill numbers, explained the ontogenetic diet effect

on the condition factor (Fig. 4, Table S4), suggesting that the overall pattern was driven by hyperiid

amphipods acquisition. As such, a model that contained hyperiid amphipods explained the data

substantially better than the original model that had parameterized crustaceans as a whole (ΔAICc =

2.53, Table S4). Concordantly, the age-dependent slope of the association between the number of

hyperiid amphipods in the diet and condition factor was strongly significant (for hyperiid amphipods

diet by sea age interaction: t_{1583} = -3.621, p < 0.001). The condition factor of 1SW was significantly

higher when the diet contained more hyperiid amphipods (t_{1583} = 2.146, p = 0.032), but the relation was

directly opposite direction in 2SW (t_{1583} = -2.112, p = 0.035) and 3SW age groups (t_{1583} = -3.941, p <

0.001, Fig. 4a). A parallel analysis for the fish diet was not performed because the number of diet items

was prohibitively small when the fish diet data were partitioned into smaller taxonomic groups.

Fig. 4. Marginal effects of the number Hyperiid amphipods (a) and Euphausiids (b) prey items (log
scale) on condition factor in Atlantic salmon sub-adults in the marine environment. Dashed line
indicates a zero slope, no relation between the number of diet items and condition factor. Error bars
indicate 95% confidence intervals.

3.5. Contribution of diet weight vs. number of diet items on condition factor

The relationship between the condition factor and number of prey items might be driven by different ecological processes associated with foraging, such as foraging frequency or foraging efficiency (Aykanat et al. 2024). We assessed the contribution of these two processes after restructuring the response variable into two parts, first to quantify foraging frequency (approximated with a Boolean diet presence and absence), and second to quantify foraging efficiency (i.e. approximated with log-transformed number of diet items, when diet was present in the stomach), and by repeating the analysis above. The effects of diet on the condition factor were qualitatively similar to each other and to the original models for both crustaceans and dish diets (Fig. S7), ruling out the sole contribution of any of the two processes.

When the total diet weight was parameterized in the model, instead of the number of prey items acquired, it did not explain ontogenetic effects in the diet (Fig. S6), suggesting that number of diet items but not weight were the primary factor associated with the ontogenetic effect.

4. DISCUSSION

We demonstrated that resource acquisition predicts ontogenetic performance trade-offs via its effect on condition factor. Competitive resource allocation has been the major mechanism suggested to mediate life history trade-offs within species but lacks evidence in natural systems (Flatt & Heyland 2011, Haave-Audet et al. 2022, Chang et al. 2024). The theoretical premise of resource acquisition in shaping life history trade-offs dynamics has been acknowledged directly (Flatt & Heyland 2011), or indirectly across different approaches, as a potential outcome through the effect of signaling networks (Stearns 2011) or via the effect of changing environmental plasticity to couple-decouple the correlations between life history traits. However, the empirical studies involve only common garden designs under

controlled conditions (Messina & Fry 2003, Ernande et al. 2004, Lailvaux et al. 2005). Our results,

obtained in the wild, demonstrate a link between the numbers of different prey species in the diet and

condition factor, a strong predictor of the liability trait of age-at-maturity. Therefore, we suggest that

Atlantic salmon life history dynamics may be shaped by resource acquisition dynamics in the wild via

the availability and composition of different prey species.

Among crustacean taxa, hyperiid amphipods (mostly *Themisto* spp.), but not euphausiids (krill), were

associated with an age-dependent decrease in condition factor. Hyperiid amphipods are much more

abundant prey type in the marine feeding grounds of the Northeast Atlantic (Jacobsen & Hansen 2001)

but are smaller than krill or other crustaceans taxa observed in the diet (Table S1). Optimal foraging

theory predicts that organisms select prey that maximize net energy gain (Pyke et al. 1977). It is

plausible that acquiring hyperiid amphipods becomes increasingly energetically expensive as salmon

get older and larger, perhaps due to size dependent changes in morphology and swimming ability

(Sanchez-Hernandez et al. 2019). In contrast, the influence of fish diet on the condition factor increases

with sea age. Consuming fish increases the condition factor only in older sea age groups, suggesting

that some energetic constraints associated with smaller fish size exist, which may be associated with

gape size (Keeley & Grant 1997, Scharf et al. 2000), or size dependent attributes that makes younger

fish to foraging for fish less efficient (Webb et al. 1984, Sanchez-Hernandez et al. 2019, see also

below). Taken together, foraging predominantly on hyperiid amphipods at younger ages and shifting to

foraging fish prey at older ages may signify an age-dependent optimal foraging strategy for Atlantic

salmon, at least in the environmental settings that the sampling have taken place.

Combined with the fact that winter condition factor at sea is a strong predictor of maturation timing in

the following season (Herbinger 1987, Herbinger & Friars 1991), our results may have significant

implication for understanding the maturation dynamics of Atlantic salmon. For example, we predict

that a younger age structure at maturity will be associated with relatively higher crustacean abundances

(mainly hyperiid amphipods). Likewise, relatively higher fish prey abundances might negatively affect

condition factor at younger age groups, leading to delayed maturation and older age structure at

maturity. Vollset et al. (2022) suggested that broad-scale changes in oceanic conditions led to an

ecological regime shift that subsequently affected the age at maturity. In their 28-year time series data,

the decrease in planktonic biomass was highly correlated to an increase in the proportion of older sea

age groups (Vollset et al. 2022), which corroborates our prediction. In their study, there was a strong

covariation across several abiotic and biotic factors during the regime shift, making it challenging to

break down the drivers of this change further. Our study linked individual-based diet patterns to

condition factor across a heterogeneous spatial distribution of Atlantic salmon and its prey, and within

a narrow temporal window, thus providing an assessment that is independent from the effect of

temporally covarying factors, such as temperature. As such, we can suggest, among other factors

outlined in Vollset et al. (2022), that decreased crustacean abundances are likely the driver of observed

older age structure.

While the ontogenetic diet shift towards a more piscivorous diet has been demonstrated in this system

earlier (Jacobsen & Hansen 2001), its effect on condition factor was not known until this study. Based

on our results, one can suggest that the net benefit of a crustaceans oriented diet is diminishing with

age, while a fish-based diet becomes more rewarding. Since the ontogenetic relationship is driven by

prey numbers, but not mass (Fig. S6), we can postulate that the age-dependent diet benefits are caused

not by post-acquisition processes, such as differences in prey energy content or potential differences in

nutrient stoichiometry between diet types, but by differences in energy use prior to or during prey

acquisition. The result that this relationship is mostly driven by the smaller (and more numerous)

hyperiid amphipods (Fig. 4) further supports this notion. Therefore, our results are consistent with a

scenario that foraging for smaller crustaceans becomes increasingly energetically unsustainable for larger salmon. Similarly, we suggest that foraging for fish is only energetically advantageous for larger salmon, which may be due to size-dependent changes in morphology or behavior (Smith & Skúlason 1996). For example, gape size may restrict smaller salmon from capturing larger fish (Keeley & Grant 1997, Scharf et al. 2000), or they may not sustain cruising speeds that are required for foraging for fish prey as efficiently as larger salmon (Webb et al. 1984, Sanchez-Hernandez et al. 2019).

The ontogenetic effect of diet on condition was not solely explained by age-dependent changes in foraging strategy, as demonstrated by similar effects between when diet was parameterized as presence/absence (as an indication of frequency of active foraging), or total prey numbers in the diet (when diet items were non-zero, and indication of foraging outcome). This suggests that ontogenetic effects are independent of the foraging strategy. Interestingly, Aykanat et al. (2024) suggested that the *vgll3* locus, a major locus influencing age at maturity, was associated with age-dependent differences in crustaceans foraging frequency, but not foraging outcome. Fish with *vgll3* early maturing genotype appeared to gain similar amounts of food with less foraging activity, suggesting that an increase in acquisition efficiency may be one of the factors that allow salmon with the *vgll3* early genotype to mature earlier, that is, by allocating surplus resources towards reproductive investment (Aykanat et al. 2024). However, they did not find any evidence of an age-dependent association between condition factor and *vgll3* genetic variation. These results suggest that the association between diet and condition factor in the present study likely has an independent physiological pathway than how *vgll3* locus effects diet.

Similar to diet, our results show that sex has an ontogenetic effect on condition factor, in which 1SW salmon males have a higher condition factor, whereas this pattern reversed for 2SW age group (Fig. 2b). In Atlantic salmon, males mature more often as 1SW, and females more often delay maturation to

achieve a larger body size since size is a stronger determinant of reproduction success in females via its

effect on fecundity (Fleming & Einum 2010). Similarly, energy expenditure during spawning is higher

among younger males than females but increases more by size in females than in males (Jonsson &

Jonsson 2003, Persson & De Roos 2006). The reversal in sex-specific differences in condition factor

between 1SW and 2SW age groups (Fig. 4) is consistent with the indicated maturation trends above,

providing that condition factor predicts maturation via its effect on the lipid conetnt. This result is also

consistent with the proposed sex-specific variation in the liability distribution of of age at maturity,

which explains the emergence of sex-specific dominance reversal (Reid 2022), an adaptation to offset

sex differences in optimal age at maturity in Atlantic salmon (Jonsson & Jonsson 2003, Fleming &

Einum 2010, Barson et al. 2015).

Linking stomach content to general resource acquisition patterns has sometimes been criticized because

it is only a snapshot of the diet, a measure of a very recent feeding activity, as opposed to methods such

as stable isotopes that can assess long-term trends (e.g., Power et al. 2023). While integration of

multiple approaches (that includes analysis of stomach content, and stable isotope and fatty acid

profiling) may help to better understand complex trophic relationships (Nielsen et al. 2017), a careful

formulation of hypotheses and statistical methods could avoid limitations associated with sole use of

stomach content data (Nielsen et al. 2017, Amundsen & Sánchez-Hernández 2019). As such, our

results remarkably explain an ontogenetic diet shift in the species (Klemetsen et al. 2003), and provide

a well-supported and predictable relationship with condition factor, a body index that should not be

affected by only very recent feed intake. Hence, our results confirmed that stomach content analysis

reflects general feeding patterns.

The sampling site for this study was one of the major marine feeding grounds for European salmon

populations, especially for older, 2SW and 3SW sea age groups (Jacobsen & Hansen 2001, O'Sullivan

et al. 2022). However, many salmon populations from geographically distant localities, such as North

America and the Baltic Sea, utilize other marine feeding areas, and salmon of the youngest sea age

groups (1SW) may inhabit marine feeding areas that are closer to their native rivers (Rikardsen &

Dempson 2010, Gilbey et al. 2021, Rikardsen et al. 2021). Subsequently, prey compositions may be

substantially different in those habitats which potentially different factors effecting feeding and

maturation dynamics. Therefore, the age-dependent diet performance relation that we observed here,

and its potential effect on maturation may not readily be generalized across species range.

Data collected in the wild has high dimensionality. Our model contains several parameters that could

otherwise be controlled under common garden conditions. While important, testing all possible

interactions between covariates (which may covary with focal covariates) is not feasible owing to the

number of possible combinations, and unobserved variables are also likely important for diet variation.

Nonetheless, our model appears robust to the exclusion of observed variables from the model (see

Results). Furthermore, despite a strong statistical support, and multiple lines of indirect evidence

supporting the main conclusion, such as observing an ontogenetic diet shift that is common in

salmonids (Klemetsen et al. 2003) and functional morphological variation associated with size and diet

type, the age-dependent diet effect on condition factor is based on correlation of observational data.

Ideally, controlled experiments are required to confirm the causality, but this may be unfeasible for this

case, given impracticalities, at the very least, associated with rearing salmon long periods and using

forage species as food sources.

5. CONCLUSIONS

In this study, we provide a clear case of an ontogenetic performance trade-off associated with diet in a

freely roaming species across large geographical areas. The empirical evidence here supports the notion

that trade-offs mediating life history divergence could be explained by variation in resource acquisition

in the wild, and could explain the lack of genetic or phenotypic correlations observed between life

history traits. Our study provides a mechanistic basis to better predict Atlantic salmon maturation

dynamics and performance trajectories across large spatiotemporal scales. Climate-induced effects on

Atlantic salmon performance and demography have been suggested previously (Jonsson & Jonsson

2004, Todd et al. 2008, Friedland et al. 2009, Vollset et al. 2022). However, covariation between

critical biotic and abiotic physiological factors (such as temperature and forage composition and

abundances) in longitudinal data is insufficient to identify primary factors driving these patterns

(Friedland et al. 2009). Our results contribute to the knowledge by providing a spatial analysis in which

the effect of forage species (diet) on performance (condition factor) could be factored out from

covarying factors that intrinsically exist in longitudinal data structures. Our results indicate bottom-up

trophic dynamics as likely underlying factor that links large-scale ecological regime shifts to changes in

maturation patterns (Vollset et al. 2022). Our study also highlights the importance of exploiting and

repurposing old but gold datasets to address novel ecological questions.

Acknowledgements: This study was funded by the Research Council of Finland (grant nos. 328860,

353388, and 325964 to TA) and the Research Council of Norway (project number 280308 to KH). We

thank three anonymous reviewers whose comments greatly improved the quality of the manuscript.

Author contributions: TA conceptualized the paper. KH organized sample retrieval from scale

archives at NINA. JAJ coordinated the Faroese fieldwork and collected diet data. TA conducted the

statistical analyses. TA and KH drafted the MS and all coauthors contributed to subsequent drafts.

Conflict of Interest: The authors declare no competing interests.

LITERATURE CITED

Amundsen PA, Sanchez-Hernandez J (2019) Feeding studies take guts - critical review and recommendations of methods for stomach contents analysis in fish. Journal of Fish Biology 95(6) 1364-1373

Aykanat T, Jacobsen JA, Hindar K (2024) Ontogenetic variation in the marine foraging of Atlantic salmon functionally links genomic diversity with a major life history polymorphism. Molecular Ecology e17465

Barson NJ, Aykanat T, Hindar K, Baranski M and others (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. Nature 528 405-408

Bartoń K (2024) MuMIn: Multi-Model Inference. R package version 1.48.4, https://CRAN.R-project.org/package=MuMIn.

Braendle C, Heyland A, Flatt T (2011) Integrating mechanistic and evolutionary analysis of life history variation. In: Mechanisms of Life History Evolution: The Genetics and Physiology of Life History Traits and Trade-Offs (eds. Flatt T, Heyland A). Oxford University Press

Brooks ME, Kristensen K, van Benthem KJ, Magnusson A and others (2017) glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. R Journal, 9, 378-400

Chang CC, Moiron M, Sanchez-Tojar A, Niemela PT, Laskowski KL (2024) What is the meta-analytic evidence for life-history trade-offs at the genetic level? Ecology Letters 27 e14354

Czorlich Y, Aykanat T, Erkinaro J, Orell P, Primmer CR (2022) Rapid evolution in salmon life history induced by direct and indirect effects of fishing. Science 376 420-423

Debes PV, Piavchenko N, Ruokolainen A, Ovaskainen O and others (2021) Polygenic and major-locus contributions to sexual maturation timing in Atlantic salmon. Molecular Ecology 30 4505-4519

Duston J, Saunders RL (1999) Effect of winter food deprivation on growth and sexual maturity of

Atlantic salmon (*Salmo salar*) in seawater. Canadian Journal of Fisheries and Aquatic Sciences

56 201-207

Ernande B, Boudry P, Clobert J, Haure J (2004) Plasticity in resource allocation based life history traits

in the Pacific oyster, *Crassostrea gigas*. I. Spatial variation in food abundance. Journal of

Evolutionary Biology 17 342-356

Ferry-Graham LA, Bolnick DI, Wainwright PC (2002) Using functional morphology to examine the

ecology and evolution of specialization. Integrative and Comparative Biology 42 265-277

Flatt T, Heyland A (2011) Mechanisms of life history evolution : the genetics and physiology of life

history traits and trade-offs. Oxford University Press, Oxford; New York

Flatt T, Heyland A, Stearns SC (2011) What mechanistic insights can or cannot contribute to life

history evolution: An exchange between Stearns, Heyland, and Flatt. In: Mechanisms of Life

History Evolution: The Genetics and Physiology of Life History Traits and Trade-Offs (eds.

Flatt T, Heyland A). Oxford University Press

Fleming IA, Einum S (2010) Reproductive Ecology: A Tale of Two Sexes. In: Atlantic Salmon

Ecology (eds. Aas Ø, Einum S, Klemetsen A, Skurdal J). Wiley-Blackwell pp 33-65

Friedland KD, Chaput G, MacLean JC (2005) The emerging role of climate in post-smolt growth of

Atlantic salmon. ICES Journal of Marine Science 62 1338-1349

Friedland KD, MacLean JC, Hansen LP, Peyronnet AJ and others (2009) The recruitment of Atlantic

salmon in Europe. ICES Journal of Marine Science 66 289-304

Gilbey J, Utne KR, Wennevik V, Beck AC and others (2021) The early marine distribution of Atlantic

salmon in the North-east Atlantic: A genetically informed stock-specific synthesis. Fish and

Fisheries 22 1274-1306.

Haave-Audet E, Besson AA, Nakagawa S, Matho, KJ (2022) Differences in resource acquisition, not

allocation, mediate the relationship between behaviour and fitness: a systematic review and

meta-analysis. Biological Reviews of the Cambridge Philosophical Society 97 708-731

Hartig F (2022) DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression

Models. R package version 0.4.6

Herbinger CM (1987) A study of Atlantic salmon (*Salmo salar*) maturation using individually

identified fish. Dalhousie University Halifax, Canada

Herbinger CM, Friars GW (1991) Correlation between condition factor and total lipid content in

Atlantic salmon, *Salmo salar* L., parr. Aquaculture Research 22 527-529

Houle D (1991) Genetic covariance of fitness correlates - what genetic correlations are made of and

why it matters. Evolution 45 630-648

Jacobsen JA, Hansen LP (2001) Feeding habits of wild and escaped farmed Atlantic salmon, *Salmo*

salar L., in the Northeast Atlantic. ICES Journal of Marine Science 58 916-93

Jacobsen JA, Lund RA, Hansen LP, O'Maoileidigh N (2001) Seasonal differences in the origin of

Atlantic salmon (*Salmo salar* L.*)* in the Norwegian Sea based on estimates from age structures

and tag recaptures. Fisheries Research 52:169-177

Jonsson B, Jonsson N, Finstad AG (2013) Effects of temperature and food quality on age and size at

maturity in ectotherms: an experimental test with Atlantic salmon. Journal of Animal Ecology

82(1) 201-210

Jonsson N, Jonsson B, Hansen LP (1997) Changes in proximate composition and estimates of energetic

costs during upstream migration and spawning in Atlantic salmon *Salmo salar*. Journal of

Animal Ecology 66(3) 425-436

Jonsson B, Finstad AG, Jonsson N, Bradford M (2012) Winter temperature and food quality affect age

at maturity: an experimental test with Atlantic salmon (*Salmo salar*). Canadian Journal of

Fisheries and Aquatic Sciences 69 1817-1826

Jonsson B, Jonsson N (2011) Ecology of Atlantic salmon and brown trout: habitat as a template for life

histories. Springer, Dordrecht

Jonsson N, Jonsson B (2003) Energy allocation among developmental stages, age groups, and types of

Atlantic salmon (*Salmo salar*) spawners. Canadian Journal of Fisheries and Aquatic Sciences

60 506-516

Jonsson N, Jonsson B (2004) Size and age of maturity of Atlantic salmon correlate with the North

Atlantic Oscillation Index (NAOI). Journal of Fish Biology 64 241-247

Keeley ER, Grant JWA (1997) Allometry of diet selectivity in juvenile Atlantic salmon (*Salmo salar*).

Canadian Journal of Fisheries and Aquatic Sciences 54 1894-1902

Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, Mortensen E

(2003) Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus*

alpinus (L.): a review of aspects of their life histories. Ecology of Freshwater Fish 12 1–59

Lailvaux SP, Hathway J, Pomfret J, Knell RJ (2005) Horn size predicts physical performance in the

beetle *Euoniticellus intermedius* (Coleoptera: Scarabaeidae). Functional Ecology 19:632-639

Lenth R (2023) emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version

1.8.5

Leroi AM (2001) Molecular signals versus the Loi de Balancement. Trends in Ecology and Evolution

16 24-29

Messina FJ, Fry JD (2003) Environment-dependent reversal of a life history trade-off in the seed beetle

Callosobruchus maculatus. Journal of Evolutionary Biology 16 501-509

Mittelbach GG, Persson L (1998) The ontogeny of piscivory and its ecological consequences. Canadian
Journal of Fisheries and Aquatic Sciences 55 1454-1465

Mobley KB, Aykanat T, Czorlich Y, House A and others (2021) Maturation in Atlantic salmon (*Salmo*
salar, Salmonidae): a synthesis of ecological, genetic, and molecular processes. Reviews in Fish
Biology and Fisheries 31 523-571

Nielsen, JM Clare EL, Hayden B, Brett MT, Kratina P, Gilbert MTP (2018) Diet tracing in ecology:
Method comparison and selection. Methods in Ecology and Evolution 9(2) 278-291

Nowicki RC, Borgå K, Gabrielsen GW, Varpe Ø (2023) Energy content of krill and amphipods in the
Barents Sea from summer to winter: variation across species and size. Polar Biology 46 139-
150

O'Sullivan RJ, Ozerov M, Bolstad GH, Gilbey J and others (2022) Genetic stock identification reveals
greater use of an oceanic feeding ground around the Faroe Islands by multi-sea winter Atlantic
salmon, with variation in use across reporting groups. ICES Journal of Marine Science 79 2442-
2452

Pedersen J, Hislop JRG (2005) Seasonal variations in the energy density of fishes in the North Sea.
Journal of Fish Biology 59 380-389

Peterson RH, Harmon PR (2005) Changes in condition factor and gonadosomatic index in maturing
and non-maturing Atlantic salmon (*Salmo salar* L.) in Bay of Fundy sea cages, and the
effectiveness of photoperiod manipulation in reducing early maturation. Aquaculture Research
36(9): 882-889

Persson L, De Roos AM (2006) Food-dependent individual growth and population dynamics in fishes.
Journal of Fish Biology, 69, 1-20.

Power M, Thorstad EB, Forseth T, Fiske P (2023) Temporal shifts in the marine feeding of individual
Atlantic salmon inferred from scale isotope ratios. Ecology and Evolution 13 e10656

Pyke GH, Pulliam HR, Charnov EL (1977) Optimal foraging: a selective review of theory and tests.

The Quarterly Review of Biology 52:137-154

R Core Team (2019) R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria

Reid JM (2022) Intrinsic emergence and modulation of sex-specific dominance reversals in threshold

traits. Evolution 76 1924-1941

Reznick D, Nunney L, Tessier A (2000) Big houses, big cars, superfleas and the costs of reproduction.

Trends in Ecology and Evolution 15 421-425

Ricker WE (1975) Computation and inter pretation of biological statistics of fish populations. Bulletin

of the Fisheries Research Board of Canada 191:1-382

Rikardsen AH, Dempson JB (2010) Dietary Life-Support: The Food and Feeding of Atlantic Salmon at

Sea. In: Atlantic Salmon Ecology (eds. Aas Ø, Einum S, Klemetsen A, Skurdal J). Wiley-

Blackwell pp 115-143

Rikardsen AH, Righton D, Strom JF, Thorstad EB and others (2021) Redefining the oceanic

distribution of Atlantic salmon. Scientific Reports 11 12266

Roff DA (2002) Life history evolution. Sinauer Associates, Sunderland, Mass.

Rowe DK, Thorpe JE, Shanks AM (1991) Role of fat stores in the maturation of male Atlantic salmon

(*Salmo salar*) parr. Canadian Journal of Fisheries and Aquatic Sciences 48(3) 405-413

Sanchez-Hernandez J, Nunn AD, Adams CE, Amundsen PA (2019) Causes and consequences of

ontogenetic dietary shifts: a global synthesis using fish models. Biological Reviews of the

Cambridge Philosophical Society 94 539-554

Schaffer W (2003) Life histories, evolution, and salmonids. In: Evolution illuminated: Salmon and

Their Relatives (eds. Hendry AP, Stearns SC). Oxford University Press pp 20-51

Scharf FS, Juanes F, Rountree RA (2000) Predator size-prey size relationships of marine fish predators:

interspecific variation and effects of ontogeny and body size on trophic-niche breadth. Marine

Ecology Progress Series 208 229-248

Sinclair-Waters M, Odegard J, Korsvoll SA, Moen T and others (2020) Beyond large-effect loci: large-

scale GWAS reveals a mixed large-effect and polygenic architecture for age at maturity of

Atlantic salmon. Genetics Selection Evolution 52 9

Smith TB, Skúlason S (1996) Evolutionary Significance of Resource Polymorphisms in Fishes,

Amphibians, and Birds. Annual Review of Ecology and Systematics 27 111-133

Snell-Rood E, Cothran R, Espeset A, Jeyasingh P, Hobbie S, Morehouse NI (2015) Life-history

evolution in the anthropocene: effects of increasing nutrients on traits and trade-offs.

Evolutionary Applications 8 635-649

Stearns SC (1989) Trade-Offs in Life-History Evolution. Functional Ecology, 3.

Stearns SC (2000) Life history evolution: successes, limitations, and prospects. Naturwissenschaften 87

476-486

Stearns SC (2011) Does impressive progress on understanding mechanisms advance life history

theory? In: Mechanisms of Life History Evolution: The Genetics and Physiology of Life

History Traits and Trade-Offs (eds. Flatt, T, Heyland, A). Oxford University Press.

Stevenson RD, Woods WA Jr (2006) Condition indices for conservation: new uses for evolving tools.

Integrative and Comparative Biology 46(6) 1169-1190

Thorpe JE, Mangel M, Metcalfe NB, Huntingford FA (1998) Modelling the proximate basis of

salmonid life-history variation, with application to Atlantic salmon, *Salmo salar* L.

Evolutionary Ecology 12 581-599

Todd CD, Hughes SL, Marshall CT, MacLean JC, Lonergan ME, Biuw EM (2008) Detrimental effects

of recent ocean surface warming on growth condition of Atlantic salmon. Global Change

Biology 14 958-970

Van Noordwijk AJ, de Jong G (1986) Acquisition and allocation of resources - their influence on

variation in life-history tactics. American Naturalist 128 137-142

Villéger S, Brosse S, Mouchet M, Mouillot D, Vanni MJ (2017) Functional ecology of fish: current

approaches and future challenges. Aquatic Sciences 79 783-801

Vollset KW, Urdal K, Utne K, Thorstad EB and others (2022) Ecological regime shift in the Northeast

Atlantic Ocean revealed from the unprecedented reduction in marine growth of Atlantic salmon.

Science Advances 8 eabk2542

Youngson AF, McLay HA (1985) Changes in steroid hormone levels in blood serum as an early

predictor of approaching maturity in Atlantic salmon. ICES Journal of Marine Science 69(1-2)

145-157

Webb PW, Kostecki PT, Stevens ED (1984) The Effect of Size and Swimming Speed on Locomotor

Kinematics of Rainbow Trout. Journal of Experimental Biology 109 77-95

Table 1: Parsimony of models with different parametrizations. Null model is the most parsimonious structure containing non-focal terms, and
this structure included in hypothesis-specific models. LL indicates log likelihood of the model.

Model type	Hypothesis	Model structure	LL	df	AICc	dAICc
Null	No diet effect on condition factor. Other non-focal variables explains the diet variation.	Fishing period + season + SeaAge$_{factorial}$ x Length$_{age.std}$ + Length$_{age.std}$ x Sex + SeaAge$_{factorial}$ x Sex + $\varepsilon_{regional.group}$ + $\varepsilon_{location}$ + ε_{error}	1210.16	15	-2390.01	14.4
No sea age effect on diet	Both crustaceans and fish diet effects condition factor age independently.	Null model parameters + Diet$_{crus}$ + Diet$_{fish}$	1210.92	16	-2389.49	14.96
	Crustaceans diet effects condition factor age independently.	Null model parameters + Diet$_{crus}$	1213.76	16	-2395.17	9.28
	Fish diet effects condition factor age independently.	Null model parameters + Diet$_{fish}$	1215.17	17	-2395.95	8.50
Diet effect on condition factor changes non-linearly with sea age (categorical parametrization)	Crustaceans diet effects condition factor age dependently. Fish diet affects condition factor age independently.	Null model parameters + SeaAge$_{factorial}$ x Diet$_{crus}$ + Diet$_{fish}$	1219.30	19	-2400.12	4.33
	Fish diet affects condition factor age dependently. Crustaceans diet affects condition factor age independently.	Null model parameters + SeaAge$_{factorial}$ x Diet$_{fish}$ + Diet$_{crus}$	1216.34	19	-2394.2	10.25
	Both crustaceans and fish diet effects condition factor age dependently.	Null model parameters + SeaAge$_{factorial}$ x (Diet$_{crus}$ + Diet$_{fish}$)	1221.51	21	-2400.44	4.01
Diet effect on condition factor changes linearly with of sea age (continuous parametrization)	Crustaceans diet effects condition factor age dependently. Fish diet affects condition factor.	Null model parameters + SeaAge$_{continious}$ x Diet$_{crus}$ + Diet$_{fish}$	1219.29	18	-2402.14	2.31
	Fish diet effects condition factor age dependently. Crustaceans diet affects condition factor age independently.	Null model parameters + SeaAge$_{continious}$ x Diet$_{fish}$ + Diet$_{crus}$	1216.34	18	-2396.24	8.21
	Both crustaceans and fish diet effects condition factor age dependently.	Null model parameters + SeaAge$_{continious}$ x (Diet$_{crus}$ + Diet$_{fish}$)	1221.46	19	-2404.45	0

31