Mid-Cenozoic Rhinocerotid Dispersal via the North Atlantic

Danielle Fraser[1,2,3,4], Natalia Rybczynski[1,2,3], Marisa Gilbert[1], and Mary R. Dawson[5]

[1]Palaeobiology, Canadian Museum of Nature, PO Box 3443 Stn "D", Ottawa ON K1P 6P4

[2]Department of Biology, Carleton University, 1125 Colonel By Drive, Ottawa, Ontario, Canada

K1S 5B6

[3]Department of Earth Sciences, Carleton University, 1125 Colonel By Drive, Ottawa, Ontario,

Canada K1S 5B6

[4]Department of Paleobiology, Smithsonian National Museum of Natural History, Washington,

DC, 20560, USA

[5]Vertebrate Paleontology, Carnegie Museum of Natural History, 4400 Forbes Ave, Pittsburgh,

PA 15213, United States

Abstract

The North Atlantic Land Bridge (NALB), which connected Europe to North America, enabled

high latitude dispersal, particularly during globally warm periods such as the Paleocene-Eocene

Thermal Maximum, a period of dramatic faunal reorganization. It has been generally accepted by

paleontologists, based on faunal comparisons between Europe and North America, that terrestrial

vertebrates did not disperse via the NALB more recently than the early Eocene. Herein, we

describe a new Early Miocene rhinocerotid species from the Canadian High Arctic with

proximity to the NALB and present novel phylogenetic hypotheses for rhinocerotids. The new

species, a member of the genus *Epiaceratherium*, is differentiated from the four other members

of the genus by characteristics of the P3, M1-2, mandible, and lower premolars. Global-scale

biogeographic analyses reveal a high number of dispersals between Europe and North America,

in both directions; cumulatively, they near the number of dispersals within Eurasia. Notably,

multiple dispersals occurred in the Oligo-Miocene, suggesting that the NALB may have been

crossable for mammals for at least 20 Ma longer than previously considered, consistent with

emerging geological and palaeoclimatological models. We thus provide insight into the

importance of the NALB as persistent high latitude connector of geographically disparate

terrestrial faunas, underscoring the pivotal role of the Arctic in mammalian evolution.

Introduction

Cenozoic terrestrial mammalian faunas have been shaped and re-shaped globally through biotic interchange (i.e., the dispersal of organisms among regions and continents) across the land bridges of the Arctic and subarctic (e.g.,[1-7]). Driven by episodic geographical rearrangements and climate changes that reduced barriers to dispersal[8], biotic interchange enables clades to colonize new environments and diversify to fill new ecological niches (e.g.,[9,10]). It brings species into contact that otherwise would not be and can trigger profound changes to ecosystems through enhanced species richness, extinction of endemics, and alteration of the numbers and types of biotic interactions among organisms (e.g.,[11-18]). Biotic interchange has therefore structured species assemblages over large spatiotemporal scales, ultimately leading to the distributions of organisms as we know them today (e.g.,[12,14]).

Simpson [19] recognized Holarctic mammal dispersal based on faunal similarity but, at the time, considered only a single dispersal route, the Bering Land Bridge, which connected northern western North America and eastern Asia. Both fossil mammal and plant occurrences suggest that the Bering Land Bridge was intermittently crossable through much of the Cenozoic[20-22]. For example, time-calibrated molecular phylogenies for mammals and plants suggest that that dispersal via the Bering Land Bridge peaked during the Oligo-Miocene, coinciding with periods of climate warming[23]. Fossil assemblages from the High Arctic show that biotic interchange via the Bering Land Bridge also played a pivotal role in Late Neogene mammal evolution until regional subsidence led to the opening of the Bering Strait 5.5 – 5.4 Ma (mega-annum)[24-28]. Thereafter, biotic interchange was paced by the episodic formation of the Bering Land Bridge due to Pleistocene fluctuations in land ice volume and, thus, sea

level[29,30], shaping the population genetics of extant species such as wolves (*Canis lupus*),

lions (*Panthera* sp.), brown bears (*Ursus arctos*), and horses (*Equus* sp.)[31-33].

It is now clear that biotic interchange among the Northern Hemispheric continents

occurred over three possible routes, one via the Bering Land Bridge, and two via the North

Atlantic (i.e., North Atlantic Land Bridge; NALB), which connected eastern North

America and western Europe[34-37]. The NALB comprises the De Geer and Thulean

routes[35,36,38,39]. The De Geer route required crossing from Northern Europe over Svalbard

and Greenland to Ellesmere Island and is generally thought to have formed a continuous

land bridge only during the Paleocene[22,34-38,40-43]. The more southerly Thulean route

connected Britain to North America via the Greenland-Iceland-Faroe Ridge Complex and

was similarly considered traversable for terrestrial organisms only during the late

Paleocene through early Eocene[34-36,38,44-52]. During the Paleocene, the presence of these

high latitude land bridges in combination with warm climates drove the synchronous

appearance of new mammal taxa across multiple continents[53-55]. Much of the evidence for

biotic interchange via the NALB comes from comparison of mid-latitude North American

and European floral and faunal fossil assemblages of Paleocene and Early Eocene age[44,56].

Similarly, fossils from the Eocene-age Eureka Sound Formation of Ellesmere Island

strongly resemble the contemporary faunas of Western Europe, suggesting some degree of

faunal exchange[41,57,58]. While paleontologists recognize that the NALB played an

important role in biotic interchange during the early Cenozoic, dispersal via the NALB

more recently than the early Eocene has been assumed all but impossible (reviewed by

Brikiatis[34]). Recent geological and palaeontological data, however, suggest that the NALB

was not fully submerged until the Oligocene or even the Early Miocene, with some

portions submerged as late as the Quaternary (e.g.,[19,37,44,46,56]). Terrestrial dispersal via the

NALB may therefore have been possible beyond the early Eocene. High-latitude vertebrate

fossils with proximity to the NALB that are younger than the early Eocene are, however,

very rare. For most terrestrial groups, we also lack a thorough understanding of their

biogeographic history.

Herein, we describe a near-complete specimen (~75%) of a new rhinocerotid species

recovered from the Haughton Astrobleme (Devon Island, Nunavut, Canada), an

extraterrestrial impact crater located near the NALB[59]. The age of the impact has been

contentious. The impact event was originally ascribed a ~23 Ma cooling age based on

$^{40}Ar/^{39}Ar$ dating[60] and apatite fission track thermochronology[61]. It was revised to 39 Ma using

a spot-dating $^{40}Ar/^{39}Ar$ approach on glass within the gneiss[62], 23 Ma based on (U-Th)/He

ages from zircons[63], and 31 Ma based on $^{40}Ar/^{39}Ar$ ages from shocked feldspar clasts[64]. The

Haughton Formation is a crater-lake deposit preserving pollen, megaflora, and vertebrate

fossil remains[65] (see *Supplementary Text*). Biostratigraphy of the Haughton Formation is most

consistent with an Early Miocene age. Occurrences of ?*Desmatolagus schizopetrus*[66] and a

shrew belonging to the subfamily Heterosoricinae with resemblance to *Wilsonosorex*[67]

support a Hemingfordian (20.43 – 15.97 Ma) age for the Haughton Crater fauna. The pinniped

discovered in the crater, *Puijila darwini*,[68] is also a close relative of the well-known European

genus, *Potamotherium*, which occurs between ~24 – 19 Ma[69]. Furthermore, the presence of

Gramineae pollen in combination with *Acer* (maple) and *Tilia* (lindens and basswoods)

suggests an age between Miocene and Pliocene[70,71]. Thus, we consider the likely age of the

Haughton Formation to be ~23 Ma.

Rhinocerotids are an excellent group for investigation of high latitude dispersal

because, although extant forms comprise only five species and are limited to Southern Asia

and Africa[72-75], they achieved near global distribution during the Cenozoic, including North

America and Europe. Furthermore, to date, ~66 genera of extinct rhinocerotid have been

named[76]. Yet, the current lack of global-scale phylogenetic hypotheses for rhinocerotids limits

our understanding of their Holarctic dispersal. To this end, we present novel phylogenetic

hypotheses for rhinocerotids that includes the new species from the Haughton Formation of

the Canadian High Arctic. We compiled morphological character matrices from published

sources and scored morphological characters for additional North American species. We also

downloaded cytochrome B sequences for all five extant rhinocerotids (see *Methods*). Our final

dataset includes 57 species of rhinocerotid, representing most of the presently named

genera[76]. To simultaneously infer tree topology and divergence dates, we performed a

combined evidence analysis for stratigraphic range data using a fossilized birth-death (FBD)

model, a sampling model that describes how speciation, extinction, and sampling produce a

given phylogeny[77,78], implemented in RevBayes[79]. We generated the tree topology based on

estimated dates for the age of the Haughton Lake of ~23 Ma[60,61,63]. We also generated a set of

phylogenetic hypotheses using maximum parsimony applied only to the morphological data

partition as well as the character matrix of Lu *et al.*[80] implemented in TNT[81] (see *Methods*).

We then time scaled all the most parsimonious trees using the *cal3* approach in the paleotree

R package[82]. The *cal3* method enables the generation of posterior distributions of time scaled

trees that we used for downstream analyses (see *Methods*).

Finally, to characterize the timing and number of dispersals, we performed a global-

scale analyses of rhinocerotid historical biogeography using the maximum likelihood

approach available in the BioGeoBears R package[83-85]. To test for differences in dispersal

rates among regions, we used Biogeographical Stochastic Mapping[83-86] applied to our FBD

and maximum parsimony derived phylogenetic hypotheses as well as the phylogenetic

hypothesis from Lu *et al.*[80] (see *Methods*). The present study provides, to the best of our

knowledge, the most taxonomically inclusive and spatially extensive study of rhinocerotid

biogeographic history.

We find that the Haughton rhinocerotid represents a North Atlantic dispersal event no

earlier than the latest Eocene and that rhinocerotids underwent repeated dispersals via the

North Atlantic, including during the Oligo-Miocene. These dispersals included the genus

Epiaceratherium as well as ancestors to *Ronzotherium* and *Amphicaenopus*, *Prosantorhinus*

Aphelops, and *Peraceras*, taxa that vary widely in their likely palaeoecology. We therefore

present evidence of a persistent NALB supported by the terrestrial vertebrate fossil record.

What is emerging is a picture of a NALB that played a much larger role in mammal dispersal

than previously recognized, prompting a reevaluation of our understanding of Northern

Hemisphere mammalian biogeography and evolution.

Systematic palaeontology

Perissodactyla Owen, 1848

Rhinocerotoidea Owen, 1845

Rhinocerotidae Gray, 1821

Epiaceratherium Abel, 1910

Type Species

Epiaceratherium bolcense Abel, 1910

Included Species

Epiaceratherium bolcense Abel, 1910, *Epiaceratherium itjilik* sp. nov., *Epiaceratherium*

magnum Uhlig, 1999, *Epiaceratherium naduongense* Böhme et al., 2013, *Epiaceratherium*

delemontense Becker & Antoine, 2013[93].

Generic Diagnosis

Epiaceratherium is a rhinocerotid lacking i3 and a lower canine, with a wide postfossette on P2–

P4, a protoloph usually constricted on M1–M2, a straight posterior half of the ectoloph on M1–

M2, and a posterior valley usually closed on p2[94].

Stratigraphic Distribution

Late middle Eocene to early Oligocene (South Asia), Early to early late Oligocene (Europe), and

Early Miocene (Canada).

Geographical Distribution

Northern Vietnam, Pakistan, Czech Republic, Northern Italy, Germany, Switzerland, France, and

the High Arctic of Canada.

Epiaceratherium itjilik sp. nov.

Diagnosis

A species of *Epiaceratherium* characterized by an interrupted P3 protoloph, the presence of a

medifossette on the M2, low and interrupted posterior cingulum on the M1-2, vertically-oriented

mandibular ramus, labial cingulum on the lower premolars, enlargement of the fifth metacarpal,

and reduction of the third trochanter of the femur. *Epiaceratherium itjilik* sp. nov. differs from

the type species, *E. bolcense*, in lacking dental cement, a labial cingulum on the upper premolars

(sometimes present in *E. bolcense*), and a constricted protocone on the M3[90]. *E. itjilik* sp. nov.

differs from *E. magnum* in possessing comparatively reduced lingual cingula on the P3-4,

protocone constriction on M1-2, an upraised symphysis, and i2 tusks that do not diverge

rostrally, as well as the absence of cristae on the upper dentition and antecrochet on the M3[91]. *E.*

itjilik sp. nov. also differs from *E. delemontense* by lacking a forked occipital crest and by

possessing a metacone fold on the M2[93]. *E. itjilik* sp. nov. differs from *E. naduongense* in lacking

a long metastyle on the M3, lacking corrugated and wrinkled enamel, lacking a labial cingulum

on the upper molars (sometimes present in *E. naduongense*), and in possessing reduced lingual

cingula on P2-4[92].

Etymology

itjilik (Inuktitut): frost or frosty, in reference to the fact that the specimen was collected in the

High Arctic of Canada, which today is characterized by cold climates that support glaciers and

permafrost. The name was chosen in collaboration with Jarloo Kiguktak, an elder from Grise

Fiord, the most northerly Inuit community in Nunavut, Canada.

Holotype

CMNFV 59632: nearly complete skeleton of one individual

Type horizon and locality

Haughton Crater, Devon Island, Nunavut, Canada (Early Miocene), 10.8 m above the base of the

Haughton Formation.

Stratigraphical Distribution

~23 Ma, Arikareean North American Land Mammal Age

Geographical distribution

Devon Island, Nunavut, Canada.

Description

Here, we briefly describe the new taxon and provide a more extensive description with

comparison to other Eocene through Miocene rhinocerotids in the *Supplementary Text*. The

position of CMNFV 59632 within *Epiaceratherium* was confirmed by morphological

comparison and phylogenetic analysis. We performed phylogenetic analyses with the type

species for the genus and with the additional species of *Epiaceratherium* to confirm the

placement of *Epiaceratherium itjilik* sp. nov. We only present the results of the latter analysis

here. We exclude CMNFV 59632 from *Teletaceras*, *Trigonias*, and *Ronzotherium*, other early

rhinocerotids. We also exclude CMNFV 59632 from the other Oligo-Miocene North American

rhinocerotids, *Aphelops*, *Peraceras*, *Diceratherium*, *Floridaceras*, *Penetrigonias*, *Menoceras*,

Amphicaenopus, and *Subhyracodon* (see *Supplementary Text*).

  The nearly complete skeleton of one individual was found in an area measuring $5 - 7 \text{ m}^2$.

The bones occurred both as surface lag and within the active layer, above the permafrost. The

specimen preserves portions of the skull, the left and right mandibulae, and most of the

postcrania (Fig. 1-3; Fig. S1-20).

Remains of the skull are restricted to the posterio-dorsal portions of the brain case and

portions of the right and left zygomatic arch, partial skull roof, incomplete basicranium, and

braincase floor (Fig. 1; Fig. S1). The preserved right nasal bone lacks rugosity, suggesting the

absence of a horn (Fig. S1). *Epiaceratherium itjilik* sp. nov. possesses a high zygomatic arch, a

depression between the temporal and nuchal crests, and the occipital side of the skull is inclined

backward (Fig. 1; Fig. S1).

The cheek teeth are moderately worn, suggesting early to mid-adulthood (Fig. 2-3). The

maxillae are well preserved. The left maxilla preserves the P2-M3, though the M1 is heavily

damaged. The right maxilla similarly preserves the P2-M3, though the labial aspects of the P4

and M1 are damaged (Fig. 2). Though the P1 is not preserved, the anterior root is small and

rounded, while the posterior root is transversely wide. The P2 is submolariform with a protocone

and hypocone connected by a lingual bridge. The P3 is smaller than the P4. The M2 bears a

distinct metastyle. The dentition lacks cement and the upper premolars lack a labial cingulum,

though the P2-4 possesses reduced lingual cingula. There is an antecrochet on the P3, a low and

interrupted posterior lingual cingulum on the M1-2 (Fig. 2).

E. itjilik sp. nov. is definitively a rhinocerotid based on possession of a tusk-like i2 as

well as apparent absence of i3 and lower canine (Fig. 3). The left mandible bears crowns of p3,

p4, and m2 and the right mandible crowns of p3-m3, though the m1-m3 crowns are broken to

varying degrees. There is no lingual groove on the mandibular corpus and the mandibular ramus

is vertically oriented. The mandibular symphysis is upraised and transversely narrow, extending

posteriorly in line with the alveolus for the talonid of the p1 (Fig. 3). The alveoli i1 and i2 are

preserved. The i2 is a tusk with a nearly horizontal root. The right p1 and p2 are preserved as

isolated teeth and posseses two roots. The p2 posses a narrower trigonid than talonid. The other

lower teeth are characterized by long paralophids and the presence of labial cingula on the lower

molars (Fig. 3).

A large portion of the post-cranial skeleton is also preserved, including the cervical

vertebrae, humeri, radii, ulnae, scapulae, femora, tibiae, ribs, forefeet, hindfeet, many of the

thoracic, lumbar, caudal vertebrae, and an incomplete pelvis. All postcranial bones except some

of the manus and pes are broken, likely due to cryoturbation, before collection (Fig. S2-20).

The atlas facets on the axis are straight (Fig. S2), the scapula is very elongated (Fig. S6),

the radius and ulna are in contact, an anterior tubercle is present on the distal end of the ulna

(Fig. S7). Adulthood is corroborated by the closure of the long bone epiphyses (Fig. S7 & 14).

The manus and pes possess the nearly complete complement of carpals and tarsals (Fig. S

8-12 & 16-20). The carpals of *E. itjilik* sp. nov. are large and deep. The manus bares a fifth

metacarpal (Fig. S8-9). The anterior height of the scaphoid is less than the posterior height, the

anterior side of the semilunate is keeled, and the proximal facet of the Mc IV is trapezoid in

shape. The third trochanter of the femur is reduced to a small area of rugosity. The proximal

articulation of the femur is high, the medio-distal gutter of the tibia is shallow, and the posterior

apophysis of the tibia is low (Fig. S14). The depth to height ratio of the astragalus is less than

1.2, the fibula facet of the astragalus is subvertical, there is presence of an expansion of the first

calcaneus facet on the astragalus (Fig. S18), and insertion of the *m. interossei* on the lateral

metapodials is short.

Results

Phylogenetic analyses. — Our Fossilized Birth-Death (FBD) tree recovered Elasmotheriinae and

Elasmotheriini. We reconstruct the split between the Rhinocerotinae and Elasmotheriinae

between ~38.9 Ma and ~34.1 Ma. We also recovered the Aceratherini, Rhinocerotina, and

Teleoceratina (Fig. S21).

Our FBD analysis returned *Parelasmotherium schansiense* as a sampled ancestor to

Elasmotherium sibiricum and *Sinotherium lagrelii* with a moderate posterior probability (0.58;

only slightly more likely than a coin flip). We also reconstruct *Penetrigonias dakotensis* as a

sampled ancestor to *Floridaceras whitei* (posterior probability of 0.44; Fig. S21), which we do

not recover as a member of the Aceratheriinae, potentially owing to the large number of

characters for which we scored their state as unknown (see *Supplementary Text*).

Our FBD tree returns CMNFV 59632 as a member of the genus *Epiaceratherium* and sister

taxon to *E. bolcense*, a species found in Europe, with a divergence date of ~37.3 Ma to ~31.7 Ma

(Fig. S21).

Using maximum parsimony, we recover three most parsimonious trees. Few of the nodes are

well supported (Bremer support >= 3; Fig. S22). The 50% majority rule consensus tree broadly

recovers the same clades as our FBD tree but with some differences in clade membership (Fig.

S22; see *Supplementary Text*).

Using maximum parsimony, we recover CMNFV 59632 as a member of the genus

Epiaceratherium, sister to both *E. delemontense* and *E. magnum*, species found in Europe and

Europe plus the Middle East, respectively. Sister to this clade, is *E. bolcense*, a European

species. The nodes within the genus *Epiaceratherium* are relatively well supported (Bremer

supports > = 3; Fig. S22).

Our reanalysis of the Lu *et al.*[80] morphological character matrix recovered 12 most

parsimonious trees. The 50% majority rule consensus tree broadly recovers the same clades with

some differences in sister relationships (Fig. S23; see *Supplementary Text*).

Biogeographic analyses. — The best fit biogeographic model was the BAYAREALIKE model

with jump dispersal (Table 1). Employing biogeographic stochastic mapping under each model

suggests the occurrence of numerous dispersal events among all biogeographic regions excepting

Africa (Fig. 4-5). Under both unstratified and stratified models, the greatest number of dispersals

occurred between Europe and the Middle East (Fig. 5). Under an unstratified model, the number

of dispersals directly between Europe and North America (i.e., total in both directions) nears the

same magnitude as between Europe and Asia (Fig. 5A). Under a stratified model where the

North Atlantic Land Bridge (NALB) was crossable until 21 Ma, the number of dispersals

between Europe and North America is marginally greater than for Europe and Asia (Fig. 5B).

Notably, dispersals between Europe and North America are reconstructed as occurring as late as

the Early Miocene (Fig. 4). However, under a model where the NALB was only crossable until

35 Ma, the number of dispersals between Europe and North America is approximately half of

that for Europe and Asia (Fig. 5C). Under all model types, dispersals between Asia and North

America are low (Fig. 5;). The results are comparable whether based on our FBD tree or

posterior distribution of time scaled most parsimonious trees (Fig. S24-27), suggesting that our

results are not characteristic of only our FBD derived phylogenetic hypothesis nor differences

between our phylogenetic hypotheses and other published hypotheses.

Discussion

Biotic interchange via the North Atlantic Land Bridge (NALB) during the Paleocene and

early Eocene is well-established based on floral and faunal similarity between mid-latitude

Europe and North America[19,44,53-56]. Cessation of dispersal via the NALB during the early

Eocene was inferred in the past from faunal dissimilarity as well as geological models that

suggested simultaneous opening of waterways between the Arctic Ocean and North

Atlantic (reviewed by Brikiatis [34]). Recent geological evidence, however, suggests that the

NALB may have been crossable for terrestrial organisms longer than previously proposed

(e.g.,[19,37,44,46,56]). Owing to the lack of global-scale phylogenetic hypotheses for most

terrestrial taxa and rarity of terrestrial vertebrate fossils with proximity to the NALB, many

high-latitude dispersal events beyond the early Eocene have likely remained undetected.

Herein, we describe a new rhinocerotid, *Epiaceratherium itjilik* sp. nov., recovered from

the Haughton Formation of Devon Island, Nunavut, a site near the western end of the

NALB. We also present global-scale biogeographic analyses based on novel time scaled

phylogenetic hypotheses for rhinocerotids. We find that *E. itjilik* sp. nov. represents a

North Atlantic dispersal event no earlier than the latest Eocene and that rhinocerotids

underwent repeated dispersals via the North Atlantic, including during the Oligo-Miocene,

necessitating a reevaluation of the role of the NALB in terrestrial dispersal.

Our phylogenetic hypotheses are similar, in many ways, to other published hypotheses

(e.g.,[93-96]; see *Supplementary Text*). We recover both Elasmotheriinae and Rhinocerotinae

using both Fossilized Birth-Death model (FBD) and parsimony (Fig. S21 & 22). We

reconstruct the divergence date between the two using FBD as ~38.9 Ma – 34.1 Ma, which

is consistent with recently proposed divergence dates[97,98]. We also recover Aceratherini,

Rhinocerotina, and Teleoceratina, consistent with other, recent phylogenetic analyses

(e.g.,[93-96]). The primary differences include the phylogenetic positions for *Floridaceras*,

Aphelops, and *Peraceras*, the latter two of which we reconstruct as rhinocerotines. They

have, however, been previously recovered within[94,95,99] and outside[80] of Rhinocerotinae

(see *Supplementary Text*). There are also topological differences between our FBD- and

parsimony-derived phylogenetic hypotheses (see *Supplementary Text*). However, both

reconstruct CMNFV 59632 as a member of the genus *Epiaceratherium*, a genus of

rhinocerotids with four previously described species from the Eocene-Oligocene of

Eurasia[90-93].

The Haughton Crater rhinocerotid is diagnosable as *Epiaceratherium* but differentiated from

existing species by characteristics of the P3, M1, M2, mandible, and lower premolars (see

Differential Diagnosis), necessitating the erection of a new species, *Epiaceratherium itjilik* sp.

nov. The occurrence of *E. itjilik* sp. nov. in the Haughton Crater extends the duration of

Epiaceratherium by millions of years, given an estimated age of 23 Ma for the Haughton

Formation[60,61,63]. Moreover, its geographic range is expanded to North America, making it one of

the most widespread rhinocerotid genera. *E. itjilik* sp. nov. also represents the highest latitude

record of a rhinocerotid, to date, and the first occurrence in the Canadian Arctic Archipelago.

The presence of rhinocerotids in the Western Low Arctic (i.e., Yukon Territory) during the

Tertiary is apparent based on an enamel fragment that is not identifiable to genus or species[100];

the much higher latitude occurrence of *E. itjilik* sp. nov. reported herein is represented by a

nearly complete skeleton. The presence of a rhinocerotid in the High Arctic represents an

interesting palaeoecological puzzle. Though the flora of the Haughton Crater suggests a

temperate palaeoclimate[65,70], there would have been a near modern light regime with months of

winter darkness. Analysis of palaeodiet and migration patterns using tooth wear and isotopes

may, in the future, illuminate the ecological strategies of high latitude herbivores during the

period of deposition of the Haughton Formation.

Our FBD phylogenetic analysis reconstructs *E. itjilik* sp. nov. as sister to *E. bolcense*, a

European species, with an estimated divergence date of ~39 Ma – 34 Ma (Fig. S21), consistent

with estimates from the enamel proteome of CMNFV 59632[98]. The presence of *E. itjilik* sp. nov.

in the Canadian High Arctic may therefore be the result of a dispersal event from Europe to

North America via the NALB during the latest Eocene. In contrast, our parsimony analysis

reconstructs *E. itjilik* sp. nov. as sister to both *E. delemontense* and *E. magnum* (Fig. S22),

species found in Europe and Europe plus the Middle East[76] (defined here to include Pakistan),

respectively. Sister to this clade, is *E. bolcense*, a European species[76]. Our tree topology thus

implies a Europe to North America dispersal via the NALB among the ancestor to *E. itjilik* sp.

nov. (Fig. 4; Fig. S24-25). Among *E. itjilik*, *E. magnum*, and *E. delemontense* our tree topology

implies either that the ancestor of the latter two species underwent at least one North America to

Eurasia dispersal (it is unclear whether via the NALB, Bering Land Bridge or both) or that the

three species derived from different populations, North American and Eurasian, of a widespread

ancestor. Thus, our parsimony analysis may suggest multiple dispersal events between North

America and Eurasia within the genus *Epiaceratherium*, whereas our FBD tree implies one.

Our global-scale biogeographic analyses show that rhinocerotids underwent

repeated intercontinental dispersals (Fig. 4-5), illuminating a much more complex history

than previously known. The oldest rhinocerotids appear almost simultaneously in North

America (Utah and Wyoming) and Russia (northeastern), the result of dispersal via the

Bering Land Bridge during the Eocene[76,101]. Our models suggest a North American origin

for rhinocerotidae due to our choice of outgroup taxon (i.e., *Hyrachyus eximius*). Our

models, however, show repeated dispersals within Eurasia during the Oligo-Miocene (Fig.

4), consistent with their known biogeographic history[76,96]. For example, *Epiaceratherium*

dispersed from Asia into Europe during the late Eocene (Fig. 4), reflecting the opening of a

terrestrial corridor coinciding with the Eocene-Oligocene Transition[102]. We reconstruct

dispersal from North America to Eurasia among the ancestors to the Elasmotheriinae

during the late Oligocene (Fig. 4; Fig. S24 & 25), resulting from their closest relatives

being North American[80]. We also model frequent rhinocerotid dispersals within Eurasia,

associated with the diversification of the Elasmotheriinae as well as the Rhinocerotinae

(Fig. 4). Elasmotheres diversified in Asia and dispersed into Europe while rhinocerotines

first appeared in the Middle East, dispersing into both Asia and Europe[76]. The colonization

of Afro-Arabia by rhinocerotids then followed during the Miocene[103], and is reflected in

our models (Fig. 4; Fig. S24 & 25).

Our most novel finding is that direct dispersal between North America and Europe

occurred with surprisingly high frequency, second only to dispersals within Eurasia (Fig.

5). The oldest Europe to North America dispersal occurred no earlier than the divergence

of *Epiaceratherium itjilik* sp. nov. from other members of the genus during the latest

Eocene (Fig. 4). We model a North American origin for the ancestors of the

Elasmotheriinae and Rhinocerotinae (Fig. 4; Fig. S24 & 25) and, thus, a North American to

Europe dispersal via the NALB prior to their divergence. Lu *et al.*[80] also reconstruct the

sister taxa to the rhinocerotines and elasmotherines as North American, similarly implying

dispersal via the NALB prior to their divergence. We also model more recent dispersals

between North America and Europe among the ancestors of taxa including *Prosantorhinus*

as well as *Peraceras* and *Aphelops* (Fig. 4). Our reconstruction of a dispersal event

between Europe and North America within the Teleoceratini is consistent with other

phylogenetic hypotheses that reconstruct the sister taxon of *Teleoceras* as European

(*Prosantorhinus*) [80]. We, however, find a unique phylogenetic position for *Peraceras* and

Aphelops (see *Supplementary Text*) that is indicative of a dispersal from Eurasia (modelled

as likely from Europe [Fig. 4; Fig. S24 & 25]) into North America within the

Rhinocerotinae. The node placing *Aphelops* and *Peraceras* within the Rhinocerotinae is,

however, poorly supported (posterior probability of 0.28; Bremer Support < 3) (Fig. S21 &

22). The topology of Lu *et al.*[80] also implies that both *Aphelops* and *Peraceras* were

immigrant taxa, though they are recovered as members of the Aceratheriinae with Asian

sister taxa. Despite the curious position of taxa such as *Peraceras* and *Aphelops* in our

phylogenetic hypotheses, we reconstruct similarly high rates of dispersal between North

America and Europe when reanalyzing the morphological matrix of Lu *et al.*[80], suggesting

that our findings are not simply a result of unique tree topologies. Employing various

possible dates (i.e., 35 Ma and 21 Ma) for the opening of impassable North Atlantic

waterways does not change the overall pattern but, as expected, we reconstruct fewer

North Atlantic dispersals when assuming NALB submergence by 35 Ma (Fig. 5; Fig. S26

& 27).

Taxa descending from nodes where we reconstruct North Atlantic dispersals are

surprisingly variable in their likely palaeoecology, ranging from comparatively small-bodied

(e.g., *Epiaceratherium*) to large-bodied (e.g., *Prosantorhinus*), and brachydont (e.g.,

Ronzotherium) to mesodont (e.g., *Aphelops*). *Teleoceras* and *Prosantorhinus*, sister taxa in our

phylogenetic hypotheses (Fig. S21 & 22) were short-legged and barrel-chested, both having been

reconstructed as semi-aquatic, suggesting they and their most recent common ancestor would

have been highly dependent on water bodies. Furthermore, strontium isotope analyses suggest

Teleoceras major was not highly mobile[104]. The phylogenetic reconstruction of Sun, et al. [105] is

consistent with our reconstruction of a North America to Europe dispersal within the

Teleoceratini. We have therefore uncovered a palaeoecological puzzle that may be illuminated

via additional isotopic and morphological analyses.

Our phylogenetic and biogeographic analyses reveal dispersals that have been previously

considered unlikely by palaeontologists (e.g.,[19,37,44,46,56]); we present phylogenetic and model

evidence of mammalian dispersal via the NALB after the early Eocene. We suggest that the

NALB supported terrestrial dispersal for at least 20 Ma longer than previously suggested (i.e.,

intermittently crossable for at least half of the Cenozoic). Other lineages of mammals and plants

may also have undergone late dispersal via the North Atlantic. Dawson[106] recognized six

Miocene mammal genera found only in North America and Europe (i.e., *Potamotherium*,

Enhydritherium, *Amphictis*, *Cynelos*, *Ysengrinia*, and *Pseudocyon*) that hinted at late dispersal

over the NALB. Similarly, occurrences of nuts and leaves of two species of *Fagus* (i.e., beech)

on Iceland suggest that one dispersed from North America and the other from Europe during the

Late Miocene[107]. The seeds of *Fagus* are not distributed long distances (~95% disperse within 25

m of the parent plant), necessitating some degree of land continuity until the Miocene. Similarly,

the common ancestors of two genera of freshwater perch, *Perca* and *Sander*, may have been

continuously distributed across the NALB until the Middle Miocene, after which they

diverged[108-110]. A continuous distribution of freshwater fish across the NALB suggests the

presence of contiguous freshwater connections until the Miocene, consistent with our

biogeographic models for rhinocerotids.

Terrestrial crossing via the NALB to Greenland occurred via the De Geer and Thulean

routes[34-36,38,39]. Dispersal via the De Geer route (~75°N) required organisms to cross from

Sweden and Norway (i.e., Fennoscandia) via Svalbard to Greenland[35,36], a route that today, is

impeded, from east to west, by the Barents Sea and Fram Strait, [36]. Dispersal via the more

southerly Thulean route (~65°N) involved crossing the North Atlantic via the Greenland-Iceland-

Faroe Ridge Complex (collective name for thick volcanic crust between the Faroes, Iceland, and

Greenland), which formed a land connection between Scotland and Greenland[35,36]. Today,

terrestrial crossing via the Thulean route is currently impeded by the Faroe-Bank Channel,

Faroe-Shetland Channel, and the Denmark Strait[36]. Understanding of the timing of submergence

of the NALB east of Greenland, based on faunal, floral, palaeoclimate, and geological evidence,

has changed considerably since paleontologists first recognized it as a mammal dispersal route

during the Paleocene[34,36,111].

Paleontologists have generally accepted that the De Geer route was subaerial from the

latest Cretaceous through to the early Palaeocene based on geotectonics and faunal similarity

between Eurasia and North America during the Paleocene (e.g.,[1-7]; reviewed by Brikiatis[34]).

Understanding of the formation of the Barents Sea and Fram Strait have, however, been revised.

The modern Barents Sea plays key roles in coupling oceanic heat to the atmosphere and, thus,

modulation of Arctic climates[112]. Recent numerical modeling indicates the presence of a

persistent highland (i.e., the Paleo-Barents) connecting Svalbard to northernmost Europe from

the Paleocene-Eocene transition (~56 Ma) through to the Quaternary (~2.7 Ma)[113]. Today, the

Fram strait provides the only deep-water connection between the Atlantic and Arctic oceans and

is pivotal in the formation of North Atlantic Deep-Water, Atlantic Meridional Overturning

Circulation, and, thus, climate evolution during the late Cenozoic[114]. Dates for the formation of

deep water in the Fram Strait based on an array of data sources (e.g., sedimentology of ODP Site

909, Arctic Ocean sediment cores, aeromagnetic surveys [111,114-120]) now range from ~21 Ma – 14

Ma (i.e., the Miocene)[111]. During the Early Miocene, the Fram Strait was narrow (i.e., 2 km in

width), only reaching a width of 2.5 km – 2.8 km by the Middle Miocene[119]. The Fram Strait was

likely fully open by the Late Miocene [111]. Considering revisions to our understanding of the

formation of the Barents Sea and Fram Strait, it is now apparent that they provided little

geographical impedance to terrestrial dispersal via the De Geer Route until the Late Miocene.

Paleontologists have considered the De Geer and Thulean routes as non-

contemporaneous, reconstructing the Thulean route as subaerial during the latest Paleocene and

earliest Eocene based on seismic, stratigraphic, and vertebrate occurrence data (reviewed by

Brikiatis[34]). The presence of the Thulean route has been hypothesized to explain the enhanced

similarity of North American and European mammalian faunas during the Paleocene-Eocene

Thermal Maximum (~56 Ma; e.g.,[55]) and early Eocene[121]. Estimates for the timing of the

submergence of the Thulean route are, however, highly variable ranging from 35 Ma – 6 Ma

based on vertebrate occurrences, geophysical modelling, benthic foraminifera, and plant fossils,

among others[35,36,48,122-125]. Uncertainty may relate to variations in Iceland Mantle Plume activity,

which produced repeated periods of uplift and subsidence resulting in the opening and closing of

the channels of the Greenland-Scotland Ridge complex[111]. Vertebrate and geological evidence

suggest that the Faroe-Shetland Channel likely opened first, during the late Eocene or early

Oligocene[35,36,123,124] but, possibly, as late as the Miocene[125,126]. The Iceland-Faroe and

Greenland-Iceland Ridges were subsequently submerged between the late Oligocene and early to

Late Miocene[123-129]. Thus, it is apparent that the Thulean route was also likely crossable for

much longer than previously considered.

West of Greenland, there are two major barriers to dispersal between Europe and North

America. The Nares Strait separates Greenland from Ellesmere Island and the nearby, more

southerly Davis Strait separates Greenland from Baffin Island. During the Cretaceous, Greenland

and the Islands of the Canadian Arctic Archipelago, including Devon Island, formed a

contiguous landmass[130]. The Davis Strait opened prior to 57 Ma [131], while the Nares Strait was

formed during the Paleocene in two phases. Northeasterly movement of Greenland was followed

by northwesterly plate convergence, opening, and then narrowing the Nares Strait[132] Much later,

during the Quaternary, the channels of the Canadian Arctic Archipelago, were formed when

glacial activity incised the drainage patterns of the once contiguous land mass[130]. Thus, from the

late Eocene through Early Miocene, rhinocerotids dispersing between Europe to North America

via the NALB likely would have crossed at least two narrow waterways (i.e., the Fram Strait and

Nares Strait or Faroe-Shetland channel and Nares Strait). While it is possible that rhinocerotids

dispersed via swimming (e.g., the Asian rhinoceros is known to be a capable swimming[133]), we

hypothesize an additional mechanism facilitated dispersal between Europe and North America.

Biotic interchange is regulated by geography and climate, changes to which enhance or

reduce barriers to dispersal[8]. Today, cold conditions at high northern latitudes drive the

formation of sea ice, impacting animal behavior. Both caribou and muskox, for example, use sea

ice to move among Arctic islands[134,135] and, thus, connectivity and gene flow among populations

may be threatened by sea ice loss[136]. During the Paleocene and much of the Eocene, however,

the Northern Hemisphere was very likely ice-free year-round[137-140]. Estimates of shallow marine

temperatures between 11 and 22 °C are consistent with an ice-free Arctic Ocean[141]. The

transition from the globally warm hothouse (i.e., Paleocene through late Eocene) to the global

cool house (i.e., Oligocene through to Pliocene), driven by decreasing atmospheric CO_2, resulted

in the seasonal formation of sea ice in the Arctic Ocean and North Atlantic during the

Oligocene[137-139,142,143]. Sub-Arctic Ice-Rafted Debris (IRD) records (IODP site 909) suggest that

perennial Northern Hemisphere glaciation only commenced during the late Middle Miocene

(~14 Ma)[139,144] or Late Miocene (~10.8 Ma)[145]. IRD records from the ACEX (i.e., Arctic Coring

Expedition) sedimentary sequence, however, provide evidence of middle Eocene (47 – 46 Ma or

44 – 41 Ma) perennial (i.e., seasonal) ice in the Arctic Ocean[146-149], which is supported by

estimates of sea surface temperatures from micropalaeontology[150]. The ACEX sequence,

represents conditions in the central Arctic Ocean, and, so, may not be indicative of ice formation

at the lower latitudes of the NALB. ODP site 913, situated east of Greenland between Svalbard

and Iceland, however, also contains IRD, suggesting the presence of at least perennial sea ice by

the late Eocene and Oligocene[36,144,145]. Furthermore, source determination of Fe-oxide grains

from ODP site 913 tell a very similar story to the ACEX sequence. Source determination

suggests the presence of glacial and sea ice in Greenland and the Canadian Arctic, as well as the

Western, Northwestern, Northeastern, and Eastern Arctic by the late Eocene and Oligocene[143].

We therefore hypothesize that, from the late Eocene onward, winter sea ice may have

episodically provided seasonal dispersal routes for terrestrial mammals. Thus, the picture that is

emerging is one of a NALB that may have been crossable by terrestrial animals into the mid

Cenozoic and, possibly, the Neogene.

While ice may have provided a physical means of crossing the NALB after the early

Eocene (in addition to swimming), climate cooling from the mid Eocene onward may have

presented a significant barrier, particularly during the Eocene-Oligocene Transition (EOT; ~34.1

– 33.55 Ma) [137,138,151]; peaks in high-latitude biotic interchange are most often associated warm

intervals (e.g., [23]), presumably due to enhanced food availability and more favorable survival

conditions at high latitudes. *Epiaceratherium itjilik* sp. nov. represents a dispersal event between

Europe and North America that occurred prior to the EOT, when the Arctic was transitioning

from the comparatively wet rainforest "hot house" of the early and middle Eocene[152] to cooler

more temperate forests of the late Eocene "warm house"[151]. Thus, the population(s) dispersing

across the NALB during the latest Eocene that gave rise to *E. itjilik* sp. nov. would have

experienced relatively warm conditions while also dispersing among landmasses over winter sea

ice. While the EOT would have presented much harsher high latitude winters[137,138,151], during the

Oligocene global "cool house," the Arctic was anomalously warm and comparable to the late

Eocene [142]. Similarly, Middle Miocene high latitudes were comparatively warm, despite the

apparent presence of at least seasonal glaciation[139,144,153]. The Early Miocene floral assemblage

associated with *E. itjilik* sp. nov. (Haughton Formation, Devon Island) is suggestive of a cool

temperate climate with sub-freezing winter temperatures[70]. Extant ungulates (e.g., caribou and

Muskox), however, endure much harsher winters using their hooves and antlers to dig for food,

through ice and snow[154]. These strategies support large, non-migratory herds in the High Arctic

through long, cold, and dark winters. Some caribou populations, such as on Svalbard, are also

known to feed on washed up seaweed[155], though such diets are not long-term strategies. High

northern latitude populations of rhinocerotids could have adopted similar survival strategies to

survive cold, dark winters. Furthermore, high latitude ecosystems from the mid Eocene through

early Miocene supported forests that are now found at much lower latitudes[70]. Woody stems

from bushes and trees, which support ungulates such as moose (*Alces alces*)[156], could therefore

also have provided winter sustenance for dispersing rhinocerotids.

Presently, there are, to our knowledge, no records of post-middle Eocene terrestrial

rhinocerotids from the islands that comprised the NALB. The discovery of rhinocerotid fossils or

palaeomolecules in sedimentary rocks of Oligo-Miocene age on Iceland or Greenland would

support the findings of the present paper. However, such rocks are rare on both islands, and those

that are fossiliferous primarily preserve the remains of invertebrates such as marine bivalves,

insects, and plants[122,126,157,158]. The remains of a possible deerlet, however, have been recovered

from Iceland dating to the Pliocene[159,160]. Despite the current limitations of the northern

vertebrate fossil record to support or refute our biogeographic findings, we believe they are not

artefactual. For example, we consider it unlikely, based on current knowledge, that the Europe to

North America dispersals we model represent Europe to Asia to North America dispersals

obscured by an incomplete fossil record. The vertebrate-bearing sedimentary rocks of Oligo-

Miocene age, a period during which we model biotic interchange between Europe and North

America among rhinocerotids, are well represented across Asia[76,161]. Moreover, rhinocerotids are

large-bodied mammals with a comparatively high probability of entering the fossil record[162,163],

thus, their fossil record is expected to be more complete than for smaller vertebrates (e.g.,

rodents). However, the fossil record for the Eocene in east Asia remains a comparative "black

box"[76]. Discovery of a sister taxon to *Epiaceratherium itjilik* sp. nov. in Russia, for example,

would change our findings for the biogeographic history of the genus. Similarly, discovery of

otherwise European taxa in east Asia could change the results of our biogeographic models,

depending on their phylogenetic affinities. Equally, however, discovery of new European forms

with close affinities to North America taxa could strengthen them. While the fossil record for

rhinocerotids certainly offers more to discover, we hypothesize, given the results of our models

and emerging geological findings, that the NALB played a more important role in the dispersal

of terrestrial vertebrates than previously recognized.

Conclusions

Herein, we describe the highest latitude occurrence of a rhinocerotid that represents a

new species, *Epiaceratherium itjilik*, sp. nov. We reconstruct *E. itjilik* as sister to one or two

species of *Epiaceratherium* with geographic ranges encompassing Europe and the Middle East,

implying direct dispersal(s) between Europe and North America as early as the latest Eocene,

potentially enabled by high latitude warmth prior to the Eocene-Oligocene Transition.

Modelling of rhinocerotid biogeographic histories employing new, novel phylogenetic

hypotheses shows repeated dispersals between Europe and North America during mid

Cenozoic and, potentially, Neogene. We suggest that persistence of the NALB into the late

Oligocene or Early Miocene, the presence of periodic climate warm periods in combination

with the formation of seasonal sea ice, and, possibly, swimming facilitated stepwise

rhinocerotid dispersal across the NALB. Rhinocerotids represent evidence that mammalian

biotic interchange occurred via the NALB more recently than the early Eocene, possibly into

the Miocene.

Methods

Specimens. — The newly described specimen CMNFV 59632 from the Haughton Astrobleme

(Devon Island, Nunavut) is housed at the Canadian Museum of Nature, Ottawa, Canada. Other

specimens scored for phylogenetic analysis are housed at the American Museum of Natural

History (New York, New York) and the Museum of Comparative Zoology (Cambridge,

Massachusetts) (Table S3).

Terminology. — The morpho-anatomical characters described here follow the terminology of

Antoine, et al. [164] and [165]. The dental abbreviations are as follows: c/C, lower/upper canine; i/I,

lower/ upper incisor; m/M, lower/upper molar; p/P, lower/upper premolar. Measurements are

provided in mm.

Phylogenetic analysis. — To simultaneously infer tree topologies and divergence times, we

performed combined-evidence analyses for stratigraphic range data using a fossilized birth-death

(FBD) model implemented in RevBayes[166,167]. FBD is a sampling model that describes how

speciation, extinction, and sampling produce a given phylogeny[77]. FBD models diverge from

other methods for inferring tree topologies in that they incorporate fossil ages, either as a series

of point observations or temporal ranges, thus ascribing higher likelihoods to tree topologies

congruent with the fossil record[78]. FBD models thus eliminate the need for *ad hoc* calibration of

node ages and produce estimates that are robust to biased sampling of fossils and extant species.

FBD models are comprised of two processes, the birth-death and fossilization

processes[78,168]. The parameters of the birth-death process include rates of speciation (λ) and

extinction (μ), the origin time (ϕ) (i.e., the starting time of the stem lineage), and the sampling

probability (ρ) (i.e., the probability that an extant species is sampled)[168]. The fossilization

process provides a distribution for the sampling of fossil lineages modeled on a Poisson process

with rate parameter ψ[168,169]. We used the birth-death range process described by [170].

Unknown parameters, λ, μ, and ψ, are sampled via MCMC. In setting up the FBD

process, we placed exponential priors on the speciation (λ) and extinction (μ) rates. Each

parameter is drawn independently from an exponential distribution with rates equal to 10. The

FBD process is conditioned on the origin time ϕ, specified as a uniform distribution on the origin

age. All extant rhinocerotids are sampled, thus $\rho = 1.0$. FBD models also estimate the probability

of ancestor-descendent pairs, which is correlated with turnover rate μ / λ and fossil recovery rate

ψ and probably of sampling ρ.

Discrete morphological character changes are represented by the substitution model,

branch rate model, and site rate model. The substitution model describes how discrete characters

change over time, modeled using a single-rate Mk model, which uses a generalized Jukes-Cantor

matrix and assumes that state changes occur symmetrically (i.e., character state changes in any

direction are equally likely). This a conservative means of modelling discrete morphological

character evolution. We used a model where $k = 4$, given that the characters used in this study

have between two and four states. The branch rate model describes variations in the rate of

morphological evolution along different branches in the phylogeny. We assumed a strict clock

model (i.e., a constant rate of morphological character changed throughout the tree). The site-rate

model describes variation in the rates of evolution among morphological characters. We allowed

gamma-distributed rate heterogeneity among characters.

Nucleotide substitution was modeled using a general time-reversible model of nucleotide

evolution with rate heterogeneity across sites modeled on a gamma distribution (i.e., the site rate

model)[168]. For the branch rate model, we relaxed the assumption of a strict molecular clock[171].

Thus, for the branch rates model, we used an uncorrelated exponential relaxed clock where

substitution rates for each lineage were independent and distributed identically according to an

exponential distribution.

For the morphological data partition, we compiled morphological character matrices from

an array of sources (Table S4)[93,95,96,164] using the set of characters retained by Becker et al.[93],

which included 214 characters modified from Antoine[172]. We opted to use the set of characters

from Becker et al.[93] because it allowed us to include the broadest sample of rhinocerotid taxa

without introducing errors due to combining character states for single taxa as scored by multiple

authors (i.e., avoiding within taxon differences in character interpretation; we cannot avoid

similar errors among taxa). All characters were aligned where character numbers had changed

and character state numbers were edited where necessary (e.g., if a character state was scored 0

in one matrix but 1 in another, they were changed to align with the Becker et al.[93]matrix). We

also scored morphological character states for specimens of *Aphelops megalodus*, *Diceratherium*

armatum, *Peraceras profectum*, and *Floridaceras whitei* (Table S3) to include a wider range of

North American genera. Character states were also scored for the Haughton Crater rhinocerotid.

Our final dataset included 57 species, representing most of the presently named genera.

Stratigraphic range data (i.e., first and last appearance dates) were downloaded from the

Paleobiology Database or taken from the relevant literature. For the molecular data partition, we

downloaded cytochrome B sequences for all extant rhinocerotids from GenBank (Table S5) and

aligned them using ClustalW2 in Jalview. We chose Cytochrome B because: a) it displays clock-

like behavior and to produce accurate phylogenies for mammals[173], and c) to limit computational

demands.

We performed a separate analysis by maximum parsimony of the morphological data

partition using TNT v. 1.61.6[81] with 1000 replicates, collapsing branches with no possible

support (rule three), and 100 trees saved per replicate. We selected *Hyrachyus eximius* Leidy,

1871 as the outgroup. All characters were coded as ordered except characters 51, 70, 76, 103,

and 134 as per Becker et al.[93] and Antoine et al[164]. We generated a 50% majority rule tree and

calculated Bremer Supports[174] using TNT. We retained all most parsimonious trees for

downstream analysis.

We also repeated the analyses of Lu *et al.*[80] using maximum parsimony. We downloaded

the TNT file and kept the various character settings (i.e., 98 characters were ordered). We

performed a maximum parsimony analysis with 1000 replicates, tree bisection-reconnection

(TBR) branch swapping, and 100 trees saved per replicate, as per Lu *et al.*[80]. We generated a

50% majority rule tree and calculated Bremer Supports[174] using TNT v. 1.6.1.6[81]. We retained all

most parsimonious trees for downstream analysis.

Time scaling most parsimonious trees.— We scaled the branch lengths of the tree topologies

generated via parsimony to reflect time in millions of years using species first and last

appearance dates sourced from the Paleobiology Database (https://paleobiodb.org) and literature.

We binned first and last occurrence dates into stages based on the International

Chronostratigraphic Chart[175] because many of the dates are based on localities that are dated by

biostratigraphy rather than direct dating methods (i.e., the dates can be constrained only to dates

associated with the stage to which fossil localities can be referred). We used cal3 dating

method[176] available in the paleotree R package[82]. The cal3 method requires estimates of

instantaneous per-capita rates of speciation and extinction[176], which we calculated by estimating

the sampling probability per bin using a likelihood function (make_durationFreqDisc in

Paleotree) and converting to instantaneous or per capita rates (sProb2sRate in Paleotree) as per

the example provided in the Paleotree documentation[82]. The cal3 method then stochastically

timescales trees using a probability distribution (gamma distribution with a shape parameter of

two) of waiting times between speciation and first appearance in the fossil record[176].

We then generated posterior distributions of 5 trees (morphological matrix employed

herein; 15 trees total) and 3 trees (Lu *et al.*[80] morphological matrix; 36 trees total) per most

parsimonious tree because cal3 allows for the random resolution of polytomies and stochastic

estimation of branch lengths based on sampling and rate estimates. Due to computational

limitations, we did not generate larger posterior distributions of time scaled most parsimonious

trees. We therefore performed all further analyses using the generated posterior distributions of

time scaled tree topologies.

Biogeographic analyses. — Each species was scored as living in one or more of five

biogeographic regions: North America, Europe, Middle East (including Pakistan and Turkey),

Asia, and Africa. Geographic ranges were taken from searches for individual species in the

Paleobiology Database, the published literature, and the database of Old World Neogene and

Quaternary rhino occurrences[177] as summarized by Antoine[76]. To test for differences in dispersal

rates among regions, we used a modelling approach and Biogeographical Stochastic Mapping

(BSM) as implemented in the BioGeoBears R package[83-86]. BioGeoBears enables fitting of an

array of biogeographic models in a likelihood framework including Dispersal-Extinction

Cladogenesis (DEC), Dispersal Vicariance Analysis (DIVA), and Bayesian Inference of

Historical Biogeography for Discrete Areas (BayArea). Each of the included models allows for a

range of processes as summarized in Matzke [83], which are assigned probabilities and used to

calculate the likelihood of the observed geographic range tip data. All three models have two free

parameters, the rate of dispersal (d) and range contraction or extinction (e) as in Matzke [83]. All

allow for species to occupy one or more regions. All three models allow for anagenesis via range

expansion and contraction. The DEC model assumes that daughter species inherit the ancestral

range, live in a subset of the ancestral range, or undergo vicariance within one area (but not

splitting off multiple areas)[86,178,179]. In contrast, the DIVA model allows for multi-region

vicariance but not inheritance of a subset of the ancestral range by a daughter species (subset

speciation in)[179]. The BayArea model assumes no range evolution at cladogenesis but allows for

sympatry across multiple regions[180]. DEC, DIVA, and BayArea can all also be implemented

incorporating founder event speciation, the process whereby a daughter species "jumps" to a new

region, a process that may play a significant role in speciation in island systems.

Herein, we assumed three possible biogeographic scenarios, the first where the North

Atlantic Land Bridge (NALB) was passable for the entire Cenozoic (i.e., an unstratified

analysis), until 35 Ma, and until 21 Ma based. We assumed equal probability of dispersal among

the other biogeographic regions throughout rhinocerotid evolutionary history. For each scenario,

we then fit all three models with and without the founder event or j parameter and compared

model fit using the Akaike Information Criterion with a correction for small sample sizes

(AICc).

We then used the best fit model for Biogeographical Stochastic Mapping (BSM)[85,181].

BSM uses simulation based on likelihood models of geographic range evolution. It performs

Ancestral Character Estimation at nodes and simulates potential histories consistent with the

observed geographic range tip data. Summarizing the various simulated range evolution

histories, BSM can be used to calculate the numbers of different types of biogeographic

events[84,86]. We performed 50 simulations and analyzed the total number of dispersal events

among all five geographic regions.

Data Availability

Phylogenetic character data for *Aphelops megalodus*, *Diceratherium armatum*, *Peraceras*

profectum, and *Floridaceras whitei* were collected by the authors from the American Museum of

Natural History and Museum of Comparative Zoology. Specimen numbers are listed in Table S3.

Biogeographic data were collected from the Paleobiology Database (https://paleobiodb.org),

relevant literature, or the database of Old World Neogene and Quaternary rhino occurrences[177].

The data used in this study have been deposited in a Github repository

(https://doi.org/10.5281/zenodo.16966587).

Code Availability

The code used for the current study is available in a Github repository

(https://doi.org/10.5281/zenodo.16966587).

LITERATURE CITED

1   Tabuce, R., Clavel, J. & Antunes, M. T. A structural intermediate between triisodontids

and mesonychians (Mammalia, Acreodi) from the earliest Eocene of Portugal.

Naturwissenschaften **98**, 145-155 (2011).

2   Rose, K. D. *The beginning of the age of mammals*. (JHU Press, 2006).

3   Missiaen, P. An updated mammalian biochronology and biogeography for the early

Paleogene of Asia. *Vertebrata Palasiatica* **49**, 29 (2011).

4   Lofgren, D. L., Lillegraven, J.A., Clemens, W.A., Gingerich, P.D. & Williamson, T.A. in

Late Cretaceous and Cenozoic mammals of North America: biostratigraphy and

biochronology   (ed M.O. Woodburne) 43–105 (Columbia University Press, 2004).

5   McKenna, M. C. & Bell, S. K. *Classification of Mammals Above the Species Level*.

(Columbia University Press, 1997).

6   Lucas, S. G. in *Evolution of Tertiary mammals of North America*   (eds C. M. Janis, K.

M. Scott, & L. L. Jacobs and) 274-283 (1998).

7   Lüthje, C. J., Milàn, J. & Hurum, J. r. H. Paleocene tracks of the mammal pantodont

genus *Titanoides* in coal-bearing strata, Svalbard, Arctic Norway. *J. Vert. Paleontol.* **30**,

521-527 (2010).

8   Vermeij, G. J. When biotas meet: understanding biotic interchange. *Science* **253**, 1099-

1104 (1991).

9   Cantalapiedra, J. L., Prado, J. L., Fernández, M. H. & Alberdi, M. T. Decoupled

ecomorphological evolution and diversification in Neogene-Quaternary horses. *Science*

355, 627-630 (2017).

10   Pires, M. M., Silvestro, D. & Quental, T. B. Continental faunal exchange and the

asymmetrical radiation of carnivores. *Proceedings of the Royal Society B: Biological*

Sciences **282**, 20151952 (2015).

11   Pires, M. M., Silvestro, D. & Quental, T. B. Interactions within and between clades

shaped the diversification of terrestrial carnivores. *Evolution* **71**, 1855-1864 (2017).

12   Fraser, D. & Lyons, S. K. Biotic interchange has structured Western Hemisphere

mammal communities. *Global Ecol. Biogeogr.* **26**, 1408-1422, doi:10.1111/geb.12667

(2017).

13   Fraser, D. & Lyons, S. K. Mammal community structure through the Paleocene-Eocene

Thermal Maximum. *Am. Nat.* **196**, 1-20 (2020).

14   Faurby, S. & Svenning, J.-C. The asymmetry in the Great American Biotic Interchange in

mammals is consistent with differential susceptibility to mammalian predation. *Global*

Ecol. Biogeogr. **25**, 1443-1453, doi:10.1111/geb.12504 (2016).

15   Kempf, H. L., Castro, I. O., Dineen, A. A., Tyler, C. L. & Roopnarine, P. D.

Comparisons of Late Ordovician ecosystem dynamics before and after the Richmondian

invasion reveal consequences of invasive species in benthic marine paleocommunities.

Paleobiology **46**, 320-336 (2020).

16   McCreless, E. E. *et al.* Past and estimated future impact of invasive alien mammals on

insular threatened vertebrate populations. *Nat Commun* **7** (2016).

17   Sax, D. F. & Gaines, S. D. Species invasions and extinction: the future of native

biodiversity on islands. *Proceedings of the National Academy of Sciences* **105**, 11490-

11497 (2008).

18   Fraser, D. *et al.* Investigating Biotic Interactions in Deep Time. *Trends Ecol. Evol.* **36**,

61-75 (2020).

19   Simpson, G. G. Holarctic mammal faunas and continental relationships during the

Cenozoic. *Geological Society of America Bulletin* **58**, 613-688, doi:10.1130/0016-

7606(1947)58[613:hmfacr]2.0.co;2 (1947).

20   Manchester, S. R. Biogeographical relationships of North American tertiary floras.

Annals of the Missouri Botanical Garden **86**, 472-522 (1999).

21   Manchester, S. R. & Shuang-Xing, G. *Palaeocarpinus* (extinct Betulaceae) from

northwestern China: new evidence for Paleocene floristic continuity between Asia, North

America, and Europe. *Int. J. Plant Sci.* **157**, 240-246 (1996).

22   Marincovich, L., Brouwers, E. M., Hopkins, D. M. & McKenna, M. C. in *The Arctic*

Ocean Region   (ed A. Grantz, Johnson, L., and Sweeney, J.) 403–426 (Geological

Society of America, 1990).

23   Jiang, D., Klaus, S., Zhang, Y.-P., Hillis, D. M. & Li, J.-T. Asymmetric biotic

interchange across the Bering land bridge between Eurasia and North America. *National*

Science Review **6**, 739-745 (2019).

24   Gladenkov, A. Y. Onset of connections between the Pacific and Arctic Oceans through

the Bering Strait in the Neogene. *Stratigraphy and Geological Correlation* **12**, 175-187

(2004).

25   Rybczynski, N. *et al.* Mid-Pliocene warm-period deposits in the High Arctic yield insight

into camel evolution. *Nature Communications* **4**, 1-9 (2013).

26   Hutchison, J. H. & Harington, C. A peculiar new fossil shrew (Lipotyphla, Soricidae)

793   from the High Arctic of Canada. *Canadian Journal of Earth Sciences* **39**, 439-443

794   (2002).

795   27   Wang, X., Rybczynski, N., Harington, C. R., White, S. C. & Tedford, R. H. J. S. r. A

796   basal ursine bear (*Protarctos abstrusus*) from the Pliocene High Arctic reveals Eurasian

797   affinities and a diet rich in fermentable sugars. **7**, 17722 (2017).

798   28   Hulbert, J., Richard C & Harington, C. R. An early Pliocene hipparionine horse from the

799   Canadian Arctic. *Palaeontology* **42**, 1017-1025 (1999).

800   29   Kender, S. *et al.* Closure of the Bering Strait caused mid-Pleistocene transition cooling.

801   *Nature Communications* **9**, 5386 (2018).

802   30   Knudson, K. P. & Ravelo, A. C. North Pacific Intermediate Water circulation enhanced

803   by the closure of the Bering Strait. *Paleoceanography* **30**, 1287-1304 (2015).

804   31   Koblmüller, S. *et al.* Whole mitochondrial genomes illuminate ancient intercontinental

805   dispersals of grey wolves (*Canis lupus*). *J. Biogeogr.* **43**, 1728-1738 (2016).

806   32   Salis, A. T. *et al.* Lions and brown bears colonized North America in multiple

807   synchronous waves of dispersal across the Bering Land Bridge. *Mol. Ecol.* **31**, 6407-6421

808   (2022).

809   33   Running Horse Collin, Y. *et al.* Sustainability insights from Late Pleistocene climate

810   change and horse migration patterns. *Science* **388**, 748-755,

811   doi:doi:10.1126/science.adr2355 (2025).

812   34   Brikiatis, L. The De Geer, Thulean and Beringia routes: key concepts for understanding

813   early Cenozoic biogeography. *J. Biogeogr.* **41**, 1036-1054 (2014).

814   35   McKenna, M. C. Holarctic landmass rearrangement, cosmic events, and Cenozoic

815   terrestrial organisms. *Annals of the Missouri Botanical Garden* **70**, 459-489 (1983).

816   36   McKenna, M. C. in *Structure and Development of the Greenland-Scotland Ridge: New*

817   *Methods and Concepts* Vol. 8 (ed M.H.P. Bott, Saxov, S., Talwani, M., Thiede, J.) 351-

818   399 (Springer, 1983).

819   37   Tiffney, B. H. The Eocene North Atlantic land bridge: its importance in Tertiary and

820   modern phytogeography of the Northern Hemisphere. *Journal of the Arnold Arboretum*

821   **66**, 243-273 (1985).

822   38   Szalay, F. S. & McKenna, M. C. Beginning of the age of mammals in Asia: the late

823   Paleocene Gashato fauna, Mongolia. Bulletin of the AMNH; v. 144, article 4. *Bulletin of*

824   *the American Museum of Natural History* **144**, 269-318 (1971).

825   39   Tucholke, B. E. & McCoy, F. W. in *The Western North Atlantic Region*   (eds Peter R.

826   Vogt & Brian E. Tucholke) 589-602 (Geological Society of America, 1986).

827   40   Blythe, A. E. & Kleinspehn, K. L. Tectonically versus climatically driven Cenozoic

828   exhumation of the Eurasian plate margin, Svalbard: Fission track analyses. *Tectonics* **17**,

829   621-639 (1998).

830   41   Dawson, M. R., West, R. M., Ramaekers, P. & Hutchison, J. H. New evidence on the

831   palaeobiology of the Eureka Sound Formation, Arctic Canada. *Arctic* **28**, 110-116 (1975).

832   42   Müller, R. D. & Spielhagen, R. F. Evolution of the Central Tertiary Basin of Spitsbergen:

833   towards a synthesis of sediment and plate tectonic history. *Palaeogeogr.,*

834   *Palaeoclimatol., Palaeoecol.* **80**, 153-172 (1990).

43   Lee, C., Lehnert, O. & Nowlan, G. Sedimentology, stratigraphy, and clast biostratigraphy

of Cretaceous and Tertiary strata, northeastern Ellesmere Island, Nunavut. *Geological*

Survey of Canada Bulletin **592**, 115-167 (2008).

44   McKenna, M. C. Fossil mammals and early Eocene North Atlantic land continuity.

Annals of the Missouri Botanical Garden **62**, 335-353 (1975).

45   Strauch, F. Die Thule-Landbrücke als Wanderweg und Faunenscheide zwischen Atlantik

und Skandik im Tertiär. *Geologische Rundschau* **60**, 381-417 (1970).

46   Axelrod, D. I. Biogeography of oaks in the Arcto-Tertiary province. *Annals of the*

Missouri Botanical Garden **70**, 629-657 (1983).

47   Saunders, A. D., Fitton, J.G., Kerr, A.C., Norry, M.J. & Kent, R.W. in *Large igneous*

provinces: continental, oceanic, and planetary flood volcanism (ed M.F. Coffin J.J.

Mahoney) 45-93 (American Geophysical Union, 1997).

48   Hohbein, M. W., Sexton, P. F. & Cartwright, J. A. Onset of North Atlantic Deep Water

production coincident with inception of the Cenozoic global cooling trend. *Geology* **40**,

255-258 (2012).

49   Smallwood, J. R. & Gill, C. E. The rise and fall of the Faroe–Shetland Basin: evidence

from seismic mapping of the Balder Formation. *Journal of the Geological Society* **159**,

627-630 (2002).

50   Shaw Champion, M., White, N., Jones, S. & Lovell, J. Quantifying transient mantle

convective uplift: An example from the Faroe□Shetland basin. *Tectonics* **27** (2008).

51   Hartley, R. A., Roberts, G. G., White, N. & Richardson, C. Transient convective uplift of

an ancient buried landscape. *Nature Geoscience* **4**, 562-565 (2011).

52   Beard, K. C. & Dawson, M. R. Intercontinental dispersal of Holarctic land mammals near

the Paleocene/Eocene boundary; paleogeographic, paleoclimatic and biostratigraphic

implications. *Bull. Soc. Geol. Fr.* **170**, 697-706 (1999).

53   Gingerich, P. D. in *Causes and consequences of globally warm climates in the Early*

Paleogene   (eds SL Wing, PD Gingerich, B Schmitz, & E Thomas)   463-478 (Geological

Society of America, 2003).

54   Bowen, G. J., Koch, P. L., Meng, J., Ye, J. & Ting, S. Age and correlation of fossiliferous

late Paleocene–early Eocene strata of the Erlian Basin, Inner Mongolia, China. *American*

Museum Novitates **2005**, 1-26 (2005).

55   Smith, T., Rose, K. D. & Gingerich, P. D. Rapid Asia–Europe–North America

geographic dispersal of earliest Eocene primate Teilhardina during the Paleocene–Eocene

thermal maximum. *Proceedings of the National Academy of Sciences* **103**, 11223-11227

(2006).

56   Savage, D. E. The Sparnacian-Wasatchian mammalian fauna, Early Eocene, of Europe

and North America. *Abh. Hess. Landesamt. Bodenf* **60**, 154-158 (1971).

57   Dawson, M. R. Early Eocene rodents (Mammalia) from the Eureka Sound Group of

Ellesmere Island, Canada. *Canadian Journal of Earth Sciences* **38**, 1107-1116,

doi:10.1139/cjes-38-7-1107 (2001).

58   Dawson, M. R., McKenna, M. C., Beard, K. C. & Hutchison, J. H. An early Eocene

plagiomenid mammal from Ellesmere and Axel Heiberg Islands, Arctic Canada. *Kaupia*

3, 179-192 (1993).

59   Robertson, P. & Mason, G. D. Shatter cones from Haughton Dome, Devon Island,

Canada. *Nature* **255**, 393-394 (1975).

60   Jessberger, E. K. Ar-40-Ar-39 dating of the Haughton impact structure. *Meteoritics* **23**,

233-234 (1988).

61   Omar, G. *et al.* Fission-track dating of Haughton astrobleme and included biota, Devon

Island, Canada. *Science* **237**, 1603-1605 (1987).

62   Sherlock, S. C. *et al.* Re□evaluating the age of the Haughton impact event. *Meteoritics &*

Planetary Science **40**, 1777-1787 (2005).

63   Young, K. E. *et al.* Impact thermochronology and the age of Haughton impact structure,

Canada. *Geophys. Res. Lett.* **40**, 3836-3840 (2013).

64   Erickson, T. M. *et al.* Resolving the age of the Haughton impact structure using coupled

40Ar/39Ar and U-Pb geochronology. *Geochim. Cosmochim. Acta* **304**, 68-82 (2021).

65   Hickey, L. J., Johnson, K. R. & Dawson, M. R. The stratigraphy, sedimentology, and

fossils of the Haughton Formation: a post□impact crater□fill, Devon Island, NWT,

Canada. *Meteoritics* **23**, 221-231 (1988).

66   Dawson, M. R. *Oreolagus* and other Lagomorpha (Mammalia) from the Miocene of

Colorado, Wyoming, and Oregon. *University Colorado Studies Earth Sciences* **1**, 1-36

(1965).

67   Martin, J. E. A new and unusual shrew (Soricidae) from the Miocene of Colorado and

South Dakota. *J. Paleontol.* **3**, 636-641 (1978).

68   Rybczynski, N., Dawson, M. R. & Tedford, R. H. J. N. A semi-aquatic Arctic

mammalian carnivore from the Miocene epoch and origin of Pinnipedia. **458**, 1021

(2009).

69 Rabi, M., Bastl, K., Botfalvai, G., Evanics, Z. & Peigné, S. A new carnivoran fauna from

the late Oligocene of Hungary. *Palaeobiodiversity and Palaeoenvironments* **98**, 509-521

(2018).

70 Whitlock, C. & Dawson, M. R. Pollen and Vertebrates of the Early Neogene Haughton

Formation, Devon Island, Arctic Canada. *Arctic* **43**, 324-330 (1990).

71 Frisch, T. & Thorsteinsson, R. Haughton astrobleme: A mid-Cenozoic impact crater

Devon Island, Canadian arctic archipelago. *Arctic*, 108-124 (1978).

72 Dinerstein, E. *The return of the unicorns: the natural history and conservation of the*

greater one-horned rhinoceros. (Columbia University Press, 2003).

73 Havmøller, R. G. *et al.* Will current conservation responses save the critically endangered

Sumatran rhinoceros Dicerorhinus sumatrensis? *Oryx* **50**, 355-359 (2016).

74 Khan, M. K. B. M. & Strien, N. J. v. *Asian rhinos: status survey and conservation action*

plan. Vol. 32 (IUCN, 1997).

75 Rookmaaker, K. & Antoine, P.-O. New maps representing the historical and recent

distribution of the African species of rhinoceros: *Diceros bicornis*, *Ceratotherium simum*

and *Ceratotherium cottoni*. *Pachyderm* **52**, 91-96 (2012).

76 Antoine, P.-O., Becker, D., Pandolfi, L. & Geraads, D. in *Rhinos of the World: Ecology,*

Conservation and Management 31-48 (Springer, 2025).

77 Stadler, T. Sampling-through-time in birth–death trees. *J. Theor. Biol.* **267**, 396-404

(2010).

78 Wright, A. M., Bapst, D. W., Barido-Sottani, J. & Warnock, R. C. Integrating fossil

observations into phylogenetics using the fossilized birth–death model. *Annual Review of*

Ecology, Evolution, and Systematics **53**, 251-273 (2022).

79   Höhna, S., Landis, M. J. & Heath, T. A. Phylogenetic inference using RevBayes. *Current*

protocols in bioinformatics **57**, 6.16. 11-16.16. 34 (2017).

80   Lu, X.-K., Deng, T. & Pandolfi, L. Reconstructing the phylogeny of the hornless

rhinoceros Aceratheriinae. *Frontiers in Ecology and Evolution* **11**, 1005126 (2023).

81   Goloboff, P. A. & Morales, M. E. TNT version 1.6, with a graphical interface for MacOS

and Linux, including new routines in parallel. *Cladistics* **39**, 144-153 (2023).

82   Bapst, D. W. paleotree: an R package for paleontological and phylogenetic analyses of

evolution. *Methods in Ecology and Evolution* **3**, 803-807 (2012).

83   Matzke, N. J. Probabilistic historical biogeography: new models for founder-event

speciation, imperfect detection, and fossils allow improved accuracy and model-testing.

Frontiers in Biogeography **5**, 242-248 (2013).

84   Matzke, N. J. Model Selection in Historical Biogeography Reveals that Founder-event

Speciation is a Crucial Process in Island Clades. *Syst. Biol.* **63**, 951-970 (2014).

85   Stochastic mapping under biogeographical models (2016).

86   Ree, R. H. & Smith, S. A. Maximum Likelihood Inference of Geographic Range

Evolution by Dispersal, Local Extinction, and Cladogenesis. *Syst. Biol.* **57**, 4-14,

doi:10.1080/10635150701883881 (2008).

87   Owen, R. *On the archetype and homologies of the vertebrate skeleton.* (author, 1848).

88   Owen, R. Odontography. London. *UK: Hippolyte Bailliere* (1845).

89   Gray, J. E. On the natural arrangement of vertebrate mammals. *London medical*

repository **15**, 296-310 (1821).

90   Abel, O. *Kritische Untersuchungen über die paläogenen Rhinocerotiden Europas*.

(Hölder, 1910).

91   Uhlig, U. Die Rhinocerotoidea (Mammalia) aus der unteroligozänen Spaltenfüllung

Möhren 13 bei Treuchtlingen in Bayern. Abhandlungen der Bayerischen Akademie der

Wissenschafen, Mathematisch-Naturwissenschaftliche Klasse, Neue Folge. *Munich* **170**,

1254 (1999).

92   Böhme, M. *et al.* Na Duong (northern Vietnam)-an exceptional window into Eocene

ecosystems from Southeast Asia. *Zitteliana. Reihe A, Mitteilungen der Bayerischen*

Staatssammlung für Paläontologie und Geologie **53**, 121-167 (2013).

93   Becker, D., Antoine, P.-O. & Maridet, O. A new genus of Rhinocerotidae (Mammalia,

Perissodactyla) from the Oligocene of Europe. *Journal of Systematic Palaeontology* **11**,

947-972 (2013).

94   Tissier, J., Antoine, P.-O. & Becker, D. New material of *Epiaceratherium* and a new

species of *Mesaceratherium* clear up the phylogeny of early Rhinocerotidae

(Perissodactyla). *Royal Society Open Science* **7**, 200633 (2020).

95   Antoine, P.-O. *et al.* A revision of *Aceratherium blanfordi* Lydekker, 1884 (Mammalia:

Rhinocerotidae) from the Early Miocene of Pakistan: postcranials as a key. *Zool. J. Linn.*

Soc. **160**, 139-194 (2010).

96   Antoine, P.-O. *et al.* A new rhinoceros clade from the Pleistocene of Asia sheds light on

mammal dispersals to the Philippines. *Zool. J. Linn. Soc.* **194**, 416-430 (2022).

97   Liu, S. *et al.* Ancient and modern genomes unravel the evolutionary history of the

rhinoceros family. *Cell* **184**, 4874-4885. e4816 (2021).

98   Paterson, R. S. *et al.* Phylogenetically informative proteins from an Early Miocene

rhinocerotid. *Nature* **643**, 719-724, doi:10.1038/s41586-025-09231-4 (2025).

99   Antoine, P.-O., Duranthon, F. & Welcomme, J.-L. *Alicornops* (Mammalia,

970   Rhinocerotidae) dans le Miocène supérieur des collines Bugti (Balouchistan, Pakistan):

971   implications phylogénétiques. *Geodiversitas* **25**, 575-603 (2003).

972   100   Eberle, J. *et al.* The First Tertiary fossils of mammals, turtles, and fish from Canada's

973   Yukon. *American Museum Novitates* **2019**, 1-28 (2019).

974   101   Prothero, D. R. *The evolution of North American rhinoceroses*. (Cambridge University

975   Press, 2005).

976   102   Licht, A. *et al.* Balkanatolia: The insular mammalian biogeographic province that partly

977   paved the way to the Grande Coupure. *Earth-Sci. Rev.* **226**, 103929 (2022).

978   103   Pandolfi, L., Pierre-Olivier, A., Bukhsianidze, M., Lordkipanidze, D. & Rook, L.

979   Northern Eurasian rhinocerotines (Mammalia, Perissodactyla) by the Pliocene–

980   Pleistocene transition: phylogeny and historical biogeography. *Journal of Systematic*

981   *Palaeontology* **19**, 1031-1057 (2021).

982   104   Ward, C. T., Crowley, B. E. & Secord, R. Enamel carbon, oxygen, and strontium isotopes

983   reveal limited mobility in an extinct rhinoceros at Ashfall Fossil Beds, Nebraska, USA.

984   *Scientific Reports* **15**, 11651 (2025).

985   105   Sun, D., Deng, T. & Wang, S. The first record of the genus *Prosantorhinus*

986   (Perissodactyla: Rhinocerotidae) of East Asia. *Zool. J. Linn. Soc.* **202**, zlad183 (2024).

987   106   Dawson, M. R. Bering down : Miocene dispersals of land mammals between North

988   America and Europe. *The Miocene Land Mammals of Europe* (1999).

989   107   Grímsson, F. & Denk, T. Fagus from the Miocene of Iceland: systematics and

990   biogeographical considerations. *Rev. Palaeobot. Palynol.* **134**, 27-54 (2005).

108 Stepien, C. A. & Haponski, A. E. in *Biology and Culture of Percid Fishes: Principles*

and Practices (ed Konrad Dabrowski Patrick Kestemont, Robert C. Summerfelt) 3-60

(Springer, 2015).

109 Haponski, A. E. & Stepien, C. A. Phylogenetic and biogeographical relationships of the

Sander pikeperches (Percidae: Perciformes): patterns across North America and Eurasia.

Biol. J. Linn. Soc. **110**, 156-179 (2013).

110 Stepien, C., Behrmann-Godel, J, Bernatchez, L. in *Biology of perch* (ed Moyer G

Couture P) Ch. 2, 7–46 (CRC Press, 2015).

111 Straume, E. O., Gaina, C., Medvedev, S. & Nisancioglu, K. H. Global Cenozoic

paleobathymetry with a focus on the Northern Hemisphere oceanic gateways. *Gondwana*

Research **86**, 126-143 (2020).

112 Smedsrud, L. H. *et al.* The role of the Barents Sea in the Arctic climate system. *Rev.*

Geophys. **51**, 415-449 (2013).

113 Lasabuda, A. P. *et al.* Paleobathymetric reconstructions of the SW Barents Seaway and

their implications for Atlantic–Arctic ocean circulation. *Communications Earth &*

Environment **4**, 231 (2023).

114 Jakobsson, M. *et al.* The early Miocene onset of a ventilated circulation regime in the

Arctic Ocean. *Nature* **447**, 986-990 (2007).

115 Jokat, W., Lehmann, P., Damaske, D. & Nelson, J. B. Magnetic signature of North-East

Greenland, the Morris Jesup Rise, the Yermak Plateau, the central Fram Strait:

constraints for the rift/drift history between Greenland and Svalbard since the Eocene.

Tectonophysics **691**, 98-109 (2016).

116   Ehlers, B.-M. & Jokat, W. Paleo-bathymetry of the northern North Atlantic and

consequences for the opening of the Fram Strait. *Marine Geophysical Research* **34**, 25-43

(2013).

117   Engen, Ø., Faleide, J. I. & Dyreng, T. K. Opening of the Fram Strait gateway: A review

of plate tectonic constraints. *Tectonophysics* **450**, 51-69 (2008).

118   Kristoffersen, Y. in *Geological History of the Polar Oceans: Arctic Versus Antarctic* 63–

76 (Springer, 1990).

119   Knies, J. & Gaina, C. Middle Miocene ice sheet expansion in the Arctic: Views from the

Barents Sea. *Geochemistry, Geophysics, Geosystems* **9** (2008).

120   Myhre, A., Thiede, J., Firth, J., Ahagon, N., Black, K., Bloemendal, J., Brass, G.,

Bristow, J., Chow, N., Cremer,M. Site 909. *Proceedings of the Ocean Drilling Program.*

Initial Reports. Ocean Drilling Program. **151**, 159–220 (1995).

121   Beard, K. C. The oldest North American primate and mammalian biogeography during

the Paleocene–Eocene Thermal Maximum. *Proceedings of the National Academy of*

Sciences **105**, 3815-3818, doi:doi:10.1073/pnas.0710180105 (2008).

122   Denk, T., Grímsson, F. & Zetter, R. Episodic migration of oaks to Iceland: Evidence for a

North Atlantic "land bridge" in the latest Miocene. *Am. J. Bot.* **97**, 276-287 (2010).

123   Wold, C. Palaeobathymetric reconstruction on a gridded database: The northern North

Atlantic and southern Greenland-Iceland-Norwegian Sea. *Geological Society, London,*

Special Publications **90**, 271-302 (1995).

124   Poore, H., Samworth, R., White, N., Jones, S. & McCave, I. Neogene overflow of

northern component water at the Greenland☐Scotland Ridge. *Geochemistry, Geophysics,*

Geosystems **7** (2006).

125   Ramsay, A. T., Smart, C. W. & Zachos, J. C. A model of early to middle Miocene deep

ocean circulation for the Atlantic and Indian Oceans. *Geological Society, London, Special*

Publications **131**, 55-70 (1998).

126   Denk, T. *et al.* in *Late Cainozoic floras of Iceland: 15 million years of vegetation and*

climate history in the northern North Atlantic Topics in Geobiology (ed Friðgeir

Grimsson Thomas Denk , Reinhard Zetter , Leifur A. Símonarson) 647-668 (Springer,

2011).

127   Poore, R. H. *Neogene Epeirogeny and the Iceland Plume* PhD thesis, University of

Cambridge, (2008).

128   Eldholm, O., Myhre, A. M., Thiede, J. in *Cenozoic plants and climates of the Arctic* Vol.

127 *NATO ASI Series* (ed H. C. Fisher M. C. Boulter) 35-55 (Springer,

Berlin/Heidelberg, 1994).

129   Thiede, J., Eldholm, O. in *Structure and development of the Greenland-Scotland Ridge:*

New methods and concepts   (ed S. Saxow M. H. P. Bott, M. Talwani, J.Thiede) 445-456

(Plenum, 1983).

130   Trettin, H. in *Geology of the Innuitian Orogen and Arctic Platform of Canada and*

Greenland Vol. 3 *Geology of Canada Series* (ed HP Trettin) 59-66 (Geological Survey

of Canada, 1991).

131   Funck, T., Jackson, H. R., Louden, K. E. & Klingelhöfer, F. Seismic study of the

transform□rifted margin in Davis Strait between Baffin Island (Canada) and Greenland:

What happens when a plume meets a transform. *Journal of Geophysical Research: Solid*

Earth **112** (2007).

Oakey, G. N. & Chalmers, J. A. A new model for the Paleogene motion of Greenland relative to North America: Plate reconstructions of the Davis Strait and Nares Strait regions between Canada and Greenland. *Journal of Geophysical Research: Solid Earth* **117** (2012).

van Strien, N. J. *Dicerorhinus sumatrensis* (Fischer) the Sumatran or two-horned Asiatic Rhinoceros. A study of literature. *Mededelingen Landbouwhogeschool (Netherlands)* **74**, 1-82 (1974).

Miller, F. L., Barry, S. J. & Calvert, W. A. Sea-ice crossings by caribou in the south-central Canadian Arctic Archipelago and their ecological importance. *Rangifer*, 77-88 (2005).

Cuyler, C. *et al.* Muskox status, recent variation, and uncertain future. *Ambio* **49**, 805-819 (2020).

Peeters, B. *et al.* Sea ice loss increases genetic isolation in a high Arctic ungulate metapopulation. *Global Change Biol.* **26**, 2028-2041 (2020).

Zachos, J., Pagani, M., Sloan, L., Thomas, E. & Billups, K. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* **292**, 686-693 (2001).

Zachos, J. C., Dickens, G. R. & Zeebe, R. E. An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* **451**, 279-283 (2008).

Stein, R. The late Mesozoic☐Cenozoic Arctic Ocean climate and sea ice history: A challenge for past and future scientific ocean drilling. *Paleoceanography and Paleoclimatology* **34**, 1851-1894 (2019).

Polyak, L. *et al.* History of sea ice in the Arctic. *Quaternary Science Reviews* **29**, 1757–1778 (2010).

1081   141   Brice, K. L., Arthur, M. A. & Marincovich Jr, L. Late Paleocene Arctic Ocean

1082   shallow☐marine temperatures from mollusc stable isotopes. *Paleoceanography* **11**, 241-

1083   249 (1996).

1084   142   O'Brien, C. L. *et al.* The enigma of Oligocene climate and global surface temperature

1085   evolution. *Proceedings of the National Academy of Sciences* **117**, 25302-25309 (2020).

1086   143   Tripati, A. K. *et al.* Evidence for glaciation in the Northern Hemisphere back to 44 Ma

1087   from ice-rafted debris in the Greenland Sea. *Earth and Planetary Science Letters* **265**,

1088   112-122 (2008).

1089   144   Thiede, J. *et al.* Late Cenozoic history of the Polar North Atlantic: results from ocean

1090   drilling. *Quaternary Science Reviews* **17**, 185-208 (1998).

1091   145   Gruetzner, J., Matthiessen, J., Geissler, W. H., Gebhardt, A. C. & Schreck, M. A revised

1092   core-seismic integration in the Molloy Basin (ODP Site 909): Implications for the history

1093   of ice rafting and ocean circulation in the Atlantic-Arctic gateway. *Global Planet.*

1094   *Change* **215**, 103876, doi:https://doi.org/10.1016/j.gloplacha.2022.103876 (2022).

1095   146   Backman, J. *et al.* Age model and core☐seismic integration for the Cenozoic Arctic

1096   Coring Expedition sediments from the Lomonosov Ridge. *Paleoceanography* **23** (2008).

1097   147   Moran, K. *et al.* The Cenozoic palaeoenvironment of the Arctic Ocean. *Nature* **441**, 601-

1098   605 (2006).

1099   148   Darby, D. A. Ephemeral formation of perennial sea ice in the Arctic Ocean during the

1100   middle Eocene. *Nature Geoscience* **7**, 210-213 (2014).

1101   149   St. John, K. Cenozoic ice☐rafting history of the central Arctic Ocean: Terrigenous sands

1102   on the Lomonosov Ridge. *Paleoceanography* **23** (2008).

1103   150   Stickley, C. E. *et al.* Evidence for middle Eocene Arctic sea ice from diatoms and ice-

1104   rafted debris. *Nature* **460**, 376-379 (2009).

1105   151   Westerhold, T. *et al.* An astronomically dated record of Earth's climate and its

1106   predictability over the last 66 million years. *Science* **369**, 1383-1387 (2020).

1107   152   Eberle, J. J. & Greenwood, D. R. Life at the top of the greenhouse Eocene world—a

1108   review of the Eocene flora and vertebrate fauna from Canada's High Arctic. *Bulletin* **124**,

1109   3-23 (2012).

1110   153   Pound, M. J., Haywood, A. M., Salzmann, U. & Riding, J. B. Global vegetation

1111   dynamics and latitudinal temperature gradients during the Mid to Late Miocene (15.97–

1112   5.33 Ma). *Earth-Sci. Rev.* **112**, 1-22 (2012).

1113   154   Klein, D. R. Comparative ecological and behavioral adaptations of Ovibos moschatus and

1114   Rangifer tarandus. *Rangifer* **12**, 47-55 (1992).

1115   155   Hansen, B. & Aanes, R. Kelp and seaweed feeding by High-Arctic wild reindeer under

1116   extreme winter conditions. *Polar Res.* **31**, 17258 (2012).

1117   156   Bryant, J. P. & Kuropat, P. J. Selection of winter forage by subarctic browsing

1118   vertebrates: the role of plant chemistry. *Annu. Rev. Ecol. Syst.* **11**, 261-285 (1980).

1119   157   Hassan, M. Y. & Haller, J. *Tertiary faunas from Kap Brewster, East Greenland: de*

1120   *Danske ekspeditioner til Østgrønland 1947-52*. (Reitzel, 1953).

1121   158   Wappler, T. *et al.* Before the 'Big Chill': A preliminary overview of arthropods from the

1122   middle Miocene of Iceland (Insecta, Crustacea). *Palaeogeogr., Palaeoclimatol.,*

1123   *Palaeoecol.* **401**, 1-12 (2014).

1124   159   Grímsson, F. & Símonarson, L. A. Upper Tertiary non-marine environments and climatic

1125   changes in Iceland. *Jökull* **58**, 303-314 (2008).

1126   160   Símonarson, L. A. Fyrstu landspendýraleifarnar úr íslenskum tertíerlögum.

1127   *Natturufraedingurinn* **59**, 189 (1990).

1128   161   Wang, X., Flynn, L. J. & Fortelius, M. *Fossil mammals of Asia: Neogene biostratigraphy*

1129   *and chronology*. (Columbia University Press, 2013).

1130   162   Behrensmeyer, A. K., Western, D. & Boaz, D. E. D. New perspectives in vertebrate

1131   paleoecology from a recent bone assemblage. *Paleobiology* **5**, 12-21,

1132   doi:doi:10.1017/S0094837300006254 (1979).

1133   163   Behrensmeyer, A. K., Kidwell, S. M. & Gastaldo, R. A. Taphonomy and paleobiology.

1134   *Paleobiology* **26**, 103-147, doi:10.1666/0094-8373(2000)26[103:tap]2.0.co;2 (2000).

1135   164   Antoine, P.-O., Alférez, F. & Iñigo, C. A new elasmotheriine (Mammalia,

1136   Rhinocerotidae) from the Early Miocene of Spain. *Comptes Rendus Palevol* **1**, 19-26

1137   (2002).

1138   165   Short, R. A., Wallace, S. C. & Emmert, L. G. A new species of Teleoceras (Mammalia,

1139   Rhinocerotidae) from the late Hemphillian of Tennessee. *Bull. Fla. Mus. Nat. Hist.* **56**,

1140   183-260 (2019).

1141   166   Höhna, S. *et al.* RevBayes: Bayesian phylogenetic inference using graphical models and

1142   an interactive model-specification language. *Syst. Biol.* **65**, 726-736 (2016).

1143   167   Höhna, S. *et al.* Probabilistic graphical model representation in phylogenetics. *Syst. Biol.*

1144   **63**, 753-771 (2014).

1145   168   Barido-Sottani, J. *et al. Estimating a time-calibrated phylogeny of fossil and extant taxa*

1146   *using RevBayes*. .5.2:1--5.2:23 (No commercial publisher, 2020).

169   Heath, T. A., Huelsenbeck, J. P. & Stadler, T. The fossilized birth–death process for

coherent calibration of divergence-time estimates. *Proceedings of the National Academy*

of Sciences **111**, E2957-E2966 (2014).

170   Stadler, T., Gavryushkina, A., Warnock, R. C., Drummond, A. J. & Heath, T. A. The

fossilized birth-death model for the analysis of stratigraphic range data under different

speciation modes. *J. Theor. Biol.* **447**, 41-55 (2018).

171   Drummond, A. J., Ho, S. Y. W., Phillips, M. J. & Rambaut, A. Relaxed phylogenetics

and dating with confidence. *PLoS Biol.* **4**, e88 (2006).

172   Antoine, P.-O. Phylogénie et évolution des Elasmotheriina. *Mémoires du Muséum*

national d'Histoire naturelle **188**, 5-350 (2002).

173   Tobe, S. S., Kitchener, A. C. & Linacre, A. M. Reconstructing mammalian phylogenies:

a detailed comparison of the cytochrome b and cytochrome oxidase subunit I

mitochondrial genes. *PloS one* **5**, e14156 (2010).

174   Bremer, K. r. Branch support and tree stability. *Cladistics* **10**, 295-304 (1994).

175   Cohen, K. M., Finney, S.C., Gibbard, P.L. and Fan, J.-X. The ICS International

Chronostratigraphic Chart. *Episodes* **36**, 199-204 (2013).

176   Bapst, D. W. A stochastic rate-calibrated method for time-scaling phylogenies of fossil

taxa. *Methods in Ecology and Evolution* **4**, 724-733 (2013).

177   Geraads, D., Cerdeño, E., García Fernandez, D., Pandolfi, L., Billia, E., Athanassiou, A.,

Albayrak, E., Codrea, V., Obada, T., Deng, T., Tong, H., Lu, X., Pícha, S., Marciszak,

A., Jovanovic, G., Becker, D., Zervanova, J., Chaïd Saoudi, Y., Bacon, A.-M., Sévêque,

N., Patnaik, R., Brezina, J., Spassov, N., and Uzunidis, A.   (2021).

178 Ree, R. H., Moore, B. R., Webb, C. O. & Donoghue, M. J. A likelihood framework for

inferring the evolution of geographic range on phylogenetic trees. *Evolution* **59**, 2299-

2311, doi:10.1111/j.0014-3820.2005.tb00940.x (2005).

179 Ronquist, F. Dispersal-vicariance analysis: a new approach to the quantification of

historical biogeography. *Syst. Biol.* **46**, 195-203 (1997).

180 Landis, M. J., Matzke, N. J., Moore, B. R. & Huelsenbeck, J. P. Bayesian analysis of

biogeography when the number of areas is large. *Syst. Biol.* **62**, 789-804 (2013).

181 Dupin, J. *et al.* Bayesian estimation of the global biogeographical history of the

Solanaceae. *J. Biogeogr.* **44**, 887-899 (2017).

1178

1179

Acknowledgements

The Canadian Museum of Nature (CMN) resides on the traditional, unceded territory of the Anishinābe Algonquin people who have stewarded the land for thousands of years. We acknowledge that the CMN's scientific research occurs across Canada—from coast to coast to coast—on the territories of the Métis and First Nations people and in Inuit Nunangat. This work was supported by a Natural Sciences and Engineering Research Council of Canada Grant to D.F. (NSERC RGPIN-2018-05305) and The W. Garfield Weston Foundation to N.R. The field work would not have been possible without the support from the Polar Continental Shelf Program, the Nunavut Planning Commission, and the Nunavut Impact Review Board. This research was also supported by palaeontology permits from the Government of Nunavut, Department of Culture, Language, Elders and Youth, and with the permission of the Qikiqtani Inuit Association, especially Grise Fiord. Guidance on Inuktitut species name was provided by Jarloo Kiguktak. The work benefitted from discussions and fieldwork collaboration of J. Gosse. Field research was supported by the CMN and the Carnegie Museum of Natural History and a National Geographic grant to M.R.D. Collections support was provided by K. Shepherd, M. Currie, S. Rufolo, and S. Swan. 3D models were created and edited by A. Tirabasso and A. McDonald. We thank K. Johnson for helpful discussion. We also thank Jérémy Tissier for productive comments that greatly improved the systematics components of manuscript.

Author Contributions

D.F. described the specimen, took measurements and photos, performed the phylogenetic analyses, performed the biogeographic analyses, and wrote the manuscript. N.R. was field leader in 2007, 2008, 2009, and 2010, described the specimen, took measurements and photos, and wrote the manuscript. M.G. was part of the field team in 2008, 2009, and 2010, described the

specimen, took measurements and photos, and wrote the manuscript. M.R.D was field leader in

1986 when the majority of the rhinocerotid specimen was collected, described the specimen,

took measurements, and participated in 2007 and 2008 field work.

Competing Interests

The authors declare no competing interests.

TABLES

TABLE 1. Model fit statistics for the various biogeographic models.

NALB closure	Dispersal model	LnL	numparams	d	e	j	AICc	ΔAICc	AICc_wt
Unstratified	**BAYAREALIKE+J**	**-104.0**	**3**	**0.003**	**1.00E-07**	**0.088**	**214.4**	**0.0**	**1.0000**
	DEC+J	-111.7	3	0.006	1.00E-12	0.094	229.8	15.4	0.0005
	DIVALIKE+J	-111.7	3	0.007	1.00E-12	0.076	229.9	15.5	0.0004
	DIVALIKE	-125.0	2	0.015	1.00E-12	-	254.3	39.9	2.20E-09
	DEC	-132.6	2	0.014	0.008	-	269.4	55.0	1.10E-12
	BAYAREALIKE	-147.4	2	0.013	0.070	-	299.1	84.7	4.10E-19
21 Ma	**BAYAREALIKE+J**	**-109.9**	**3**	**0.004**	**0.002**	**0.092**	**226.3**	**0.0**	**0.9900**
	DIVALIKE+J	-115.4	3	0.009	1.00E-12	0.064	237.2	10.9	0.0042
	DEC+J	-115.5	3	0.007	1.00E-12	0.091	237.5	11.2	0.0036
	DIVALIKE	-125.0	2	0.016	1.00E-12	-	254.2	27.9	8.90E-07
	DEC	-132.2	2	0.014	0.006	-	268.7	42.4	6.30E-10
	BAYAREALIKE	-147.9	2	0.014	0.073	-	300.0	73.7	9.70E-17
35 Ma	**BAYAREALIKE+J**	**-115.7**	**3**	**0.005**	**0.005**	**0.100**	**237.8**	**0.0**	**0.7900**
	DIVALIKE+J	-117.2	3	0.010	1.00E-12	0.065	240.8	3.0	0.1800
	DEC+J	-118.8	3	0.008	0.002	0.092	244.2	6.4	0.0330
	DIVALIKE	-125.8	2	0.017	1.00E-12	-	255.9	18.1	9.60E-05
	DEC	-132.7	2	0.015	0.005	-	269.6	31.8	1.00E-07
	BAYAREALIKE	-149.0	2	0.014	0.075	-	302.2	64.4	8.40E-15

1210

1211   **FIGURES**

1212

1213   **FIGURE 1. 3D scan of the skull elements of *Epiaceratherium itjilik* sp. nov. (CMNFV**

1214   **59632) in right lateral view.**

1215

1216

1217

1218

1219

1220

1221

FIGURE 2. Maxillary elements of *Epiaceratherium itjilik* **sp. nov. (CMNFV 59632).**

Shown in labial (top), occlusal (middle), and lingual (bottom) views is the right maxilla (**a**) with

P2-M3 and left maxilla (**b**) with P2-M3.

1222

1223

1224

1225

1226

1227 **FIGURE 3. Mandibular elements of *Epiaceratherium itjilik* sp. nov. (CMNFV 59632).**

1228 Shown in labial (top), occlusal (middle), and lingual (bottom) views, is the right mandible (**a**)

1229 with p1-4, m1, and partial m2-3 as well as the right and left tusks and left mandible (**b**) and

1230 dentition with p3-4, and m2.

FIGURE 4. **Dispersals between North America and Europe occurred as late as the**

Miocene. Ancestral character estimation of biogeographic regions for rhinocerotids based on a

NALB opening date of 21 Ma (dotted line).

1236 **FIGURE 5. Dispersals between North America and Europe occurred with relatively high**

1237 **frequency.** Mean counts of dispersal events under an unstratified model (**A**), a model limiting

1238 NALB dispersal to before 21 Ma (**B**), and a model limiting NALB dispersal to before 35 Ma (**C**).

1239 Analysis performed on the Fossilized Birth-Death tree. Error bars represent standard deviations

1240 of counts. Black stars highlight dispersal between North America and Europe..

1241