At the onset of active whisking, the input layer of barrel cortex exhibits a 24 h window of increased excitability that depends on prior experience.

Author names and affiliations:

Molly C. Shallow[1*†], Lucy Tian[1*‡], Hudson Lin[1], Katheryn B Lefton[1,2], Siyu Chen[3], Joseph D. Dougherty[3], Joe P. Culver[4], Mary E. Lambo[1], Keith B. Hengen[1]

* These authors contributed equally to this work.

[1] Department of Biology, Washington University in Saint Louis

[2] Department of Neuroscience, Washington University in Saint Louis

[3] Department of Genetics, Washington University in Saint Louis

[4] Department of Radiology, Washington University in Saint Louis

[†] Current address: University of Oregon

[‡] Current address: University of Washington

Abstract

The development of motor control over sensory organs is a critical milestone in sensory processing, enabling active exploration and shaping of the sensory environment. However, whether the onset of sensory organ motor control directly influences the development of corresponding sensory cortices remains unknown. Here, we exploit the late onset of whisking behavior in mice to address this question in the somatosensory system. Using ex vivo electrophysiology, we discovered a transient increase in the intrinsic excitability of excitatory neurons in layer IV of the barrel cortex, which processes whisker input, precisely coinciding with the onset of active whisking at postnatal day 14 (P14). This increase in neuronal gain was specific to layer IV, independent of changes in synaptic strength, and required prior sensory experience. Strikingly, the effect was not observed in layer II/III of the barrel cortex or in the visual cortex upon eye opening, suggesting a unique interaction between the development of active sensing and the thalamocortical input layer in the somatosensory system. Predictive modeling indicated that changes in active membrane conductances alone could reliably distinguish P14 neurons in control but not whisker-deprived hemispheres. Our findings demonstrate an experience-dependent, lamina-specific refinement of neuronal excitability tightly linked to the emergence of active whisking. This transient increase in the gain of the thalamic input layer coincides with a critical period for synaptic plasticity in downstream layers, suggesting a role in facilitating cortical maturation and sensory processing. Together, our results provide evidence for a direct interaction between the development of motor control and sensory cortex, offering new insights into the experience-dependent development and refinement of sensory systems. These findings have broad implications for understanding the interplay between motor and sensory development, and how the mechanisms of perception cooperate with behavior.

Introduction

Cortical microcircuits give rise to conscious perception [1, 2, 3], adaptable interaction with stimuli [4, 5, 6, 7], and rapidly adjust to the environment into which an individual is born [8, 9, 10, 11, 12]. As a result, understanding the combination of internal processes and external influences that contribute to the development of cortical networks is a key question with broad consequences, both neurobiological [13] and societal [14, 15] in form. However, it is challenging to identify and control the features of the external world that drive specific cortical circuits. As a result, it can be difficult to pinpoint the mechanisms by which cortical development interacts with the environment. With this in mind, the mouse whisker-barrel system is an ideal model system for studying the postnatal development of cortex, specifically the interaction of experience with internal programs. The whisker-barrel system is characterized by the emergence of topographically organized maps whose architecture is notably accessible for two reasons; *first*, under normal conditions, the stereotyped rows and columns of whisker pads on the face are distinctly visible in the cellular architecture in layer IV (i.e., "barrels" in S1 [16]), and *second*, it is easy to manipulate highly specific inputs by simply removing the desired subset of whiskers— each of which map onto a single barrel [17]. Because of this accessibility, the whisker-barrel system has been utilized to uncover broad principles of development, plasticity, sensory coding, and cortical microcircuitry [18, 19, 20, 21, 22, 23].

While the structural development of barrels in the first days of life has been extensively studied [24, 25], the cortical impact of another powerful feature of the whisker-barrel system is largely unknown: the sudden onset of active whisking at postnatal day 14 (P14) [26]. The active manipulation of sensory organs to explore the world (such as sniffing with the nose), is a prominent feature of most sensory systems [27, 28, 29, 30]. This is true of tactile information; from the first moments of life, cutaneous somatosensation is driven by limb movement and locomotion [31]. In contrast to the rest of the body, however, an animal's whiskers are not actively controlled during the first two weeks of life [32, 33, 34]. While the barrels are formed and innervated prior to and shortly after birth [35, 36, 32, 18, 20], passive input through the whiskers in the first ∼two weeks establishes connectivity from the barrels (layer IV) to layers II/III [37]. A few days before the onset of active whisking, there is a brief period of "symmetric whisking", signaling an impending change from passive to active sensing [32]. Then, analogous to the binary shift that comprises eye opening, at P14 mice commence control of individual whiskers for exploration of objects and the environment. Perhaps unsurprisingly, the sudden emergence of complex whisking behavior coincides with significant Hebbian modification in layers II/III in and after the third week of life [33, 38, 39].

Despite phenomenological parallels with eye opening, the onset of active whisking differs in at least one crucial aspect: by P14, the critical period for barrel formation and plasticity is largely understood to be closed [24, 35, 40]. In contrast, the critical period for ocular dominance plasticity begins ∼10 d after eye opening [41]. This delay between the maturation of sensory transduction and the behavioral control of whiskers provides a unique opportunity to isolate the role of motor control in sensory system development. In other words, the barrel cortex on P14 is distinctive— it presents a structurally mature circuit exposed to a developmentally-timed onset of organized input [16, 42]. The default possibility is that, at P14, active whisking simply introduces complex stimuli into established circuitry. In this case, motor control of sensory organs indirectly refines sensory cortex by determining stimuli— largely reflected by plasticity in layers II/III [33, 38, 39]. Alternatively, sensory cortices could cooperatively interact with key events in motor development. In this case, the onset of sensory organ movement should accompany a parallel change in sensory cortices that is positioned to facilitate information processing. Whether the barrel cortex actively cooperates with the onset of whisking behavior and consequent complex input is unknown [43].

Based on prior work, it is reasonable to suggest that a plastic change in layer IV barrels is unlikely on P14. Barrel structure and synaptic connectivity are established prior to P14 [40], possibly leaving little room for significant synaptic modification of layer IV at this age— the relative immutability of sensory cortices after their respective critical periods is a general phenomenon [44]. While plasticity isn't limited to the synapse—intrinsic plasticity is regularly observed in many systems —non-synaptic changes almost always accompany synaptic plasticity [45, 46, 47, 48, 49]. In this context, we sought to answer a simple question: is there evidence of a plastic interaction between layer IV barrel cortex and the onset of whisker motor control at P14?

To address this, we used ex vivo slice recordings in layers II/III and IV of barrel cortex as well as in layer IV in primary visual cortex. We measured intrinsic excitability and synaptic strength throughout this key developmental time period to assess the possibility of plastic change in the barrels at the onset of active whisking. We identify a robust, non-synaptic form of experience-dependent plasticity that occurs at the onset of active whisking. This transient event is specific to excitatory neurons in layer IV barrel cortex.

Results

To explore the possibility that neurons in the barrel fields of layer IV S1 ($S1_{BF}$) show evidence of a plastic change at the onset of active whisking (P14), we took acute ex vivo brain slices (350 μm) containing layer IV barrels from mice (N = 82 animals; Fig. 1A,B). We conducted whole-cell patch-clamp recordings from excitatory spiny stellate neurons, identified by morphological characteristics and confirmed by post hoc imaging ([50, 51], Fig. 1B). To capture possible changes at P14, slices were made intermittently between P11 and P19 (Fig. 1C). We assessed intrinsic excitability via standard F/I curves (firing rate per current step; [49]), and we estimated the distribution of post-synaptic strengths by measuring miniature excitatory post-synaptic potentials (mEPSCs; [52]). In addition to stellate neurons in the barrels, we recorded excitatory pyramidal neurons in layers II/III of S1 (dorsal to the barrels; [53]), as well as two populations of NKX2.1-positive inhibitory interneurons—somatostatin (SST) and parvalbumin (PV) subtypes [54, 55].

Excitability of barrel excitatory neurons is increased for ∼24 h on P14

A fundamental property determining whether and how a neuron converts inputs into spikes is the intrinsic excitability of its membrane. A defining feature of cortical physiology in early postnatal life is a progressive reduction in neuronal intrinsic excitability. This typically unfolds over weeks, and is characterized by increasing thresholds for action potential generation and decreasing resting membrane potentials. As cortical neurons grow and develop more complex dendritic trees and form more synaptic connections, their input resistance typically decreases, which reduces their overall excitability (Fig. S1)[56, 57, 58]. Membrane excitability is assayed by injecting an increasing series of positive current steps and counting the number of action potentials generated by each step. The slope of the resultant curve—the F/I curve— thus describes the intrinsic excitability of a neuron.

Given the progressive decrease in excitability across development, the sudden onset of active whisking at P14 might reasonably expected to drive no meaningful change in intrinsic excitability beyond that which is already unfolding. Alternatively, increased sensory behavior might result in a compensatory decrease in intrinsic excitability, as cortical sensory neurons homeostatically regulate their excitability to counterbalance significant changes in input [45]. To evaluate this, we patched spiny stellate neurons in visually identified barrels at P11, 12, 14, 16, 18, and 19. We delivered ten 500 ms currents between 0 and 180 pA in steps of 20 pA (Fig. 1D). For quantification, we fit a linear model to each neuron's F/I curve between 20 and 100 pA (all cells were active and not saturated in that range) and extracted the model slope as an estimate of intrinsic excitability. There was a significant main effect of age ($p = 1.26e - 05$, mixed-effects analysis of variance:

slope $\sim$ age + animal_id, where animal ID is included as a fixed effect). Posthoc comparisons of P14 to all other ages yielded p values between 0.0000064 and 0.013 (Fig. 1E). Examination of passive properties—tau, membrane resistance, capacitance, and resting membrane potential—revealed either no change or a progressive decline with age, as expected. There were no significant alterations in any passive properties on P14 (Fig. S1A-D). These results suggest that the intrinsic excitability of excitatory neurons in the input layer of $S1_{BF}$ support an unexpected, transient increase in intrinsic excitability coincident with the onset of active whisking.

Intrinsic excitability plasticity is often tightly linked with congruent changes in synaptic strength [45, 46, 47, 48, 49]. Thus, we reasoned that elevated excitability on P14 might indicate a generalized increase in synaptic strength—as would be expected of a modifiable circuit subjected to increased input. Alternatively, increased input might be expected to drive a compensatory decrease in synaptic strength [52, 59, 60, 61]. To test these possibilities, we evaluated mEPSC amplitudes in $S1_{BF}$ excitatory neurons between P11 and P18— note that mEPSC amplitude is reflective of both homeostatic and Hebbian plasticity [52, 49, 11, 62]. Surprisingly, we found no evidence of any change in synaptic strength on P14 (Fig. 1F). Across the range of ages in question, mEPSC amplitude trended towards a decrease over time ($p = 0.08$; mixed-effects analysis of variance: mini amplitude $\sim$ age + animal_id), which is consistent with postnatal cortical maturation [63]. However, in contrast to intrinsic excitability, there was no evidence of a change in mEPSC amplitude specific to P14 (P12 vs P14 $p = 0.8915$; P16 vs P14 $p = 0.8515$). mEPSC frequency, often taken as an indicator of presynaptic properties [64], also showed no change over time ($p = 0.20$) and no specific change on P14 (P12 vs P14 $p = 0.9934$; P16 vs P14 $p = 0.8216$).

In addition to mEPSC amplitude, glutamatergic plasticity can alter subunit composition [65]. To evaluate this possibility, we examined mEPSC the waveform, as the specifics of channel composition determine waveform shape [66, 67]. While AMPA receptor subunit composition is expected to change across postnatal development [68], and thus influence mEPSC waveform shape, we sought to test whether the kinetics of glutamate-mediated synaptic transmission change abruptly at the onset of active whisking. We trained a classifier to predict age based on mEPSC waveform—should there be distinct, consistent differences in any aspect of the mEPSC waveform on P14, a classifier would learn this in a data-driven manner. We computed the mean mEPSC waveform for each neuron recorded from P11 to P18 and extracted the first four principal components, which encompassed the majority of the variance in the data (82.5%), allowing us to describe waveform characteristics without substantial information loss (Fig. S1E,F). We then trained a class-balanced logistic regression model to predict the age group based on the principal component scores (Stratified K-Fold cross-validation with 5 splits and 200 repeats). Between P11 and P18, the overall balanced accuracy was 48% (chance = 20%), an effect driven by classifier performance on P11 (62.7%) and P16 (77.5%) (Fig. S1G), consistent with a gradual, time-dependent shift in GluA subunit composition across development. However, on P14, performance was below chance (18.8%), suggesting that there is no discernible shift in the dynamics of glutamatergic synaptic transmission driven by or in anticipation of onset of active whisking. Examination of PC1 loadings indicates that the differentiability of ages (i.e., P11 and P18) was largely driven by changes in the falling phase of the mEPSC (Fig. S1H).

Taken together, these data suggest that, at the onset of active whisking, there is a plastic event in layer IV barrel cortical excitatory neurons. Excitatory neurons in the input layer of barrel cortex exhibit a transient ($\sim$24 h) increase in intrinsic excitability, despite the fact that synaptic strength and kinetics are unaffected. This raises the possibility of a brief, developmentally-timed increase in the gain of input from VPM thalamus to cortex that cooperates with the initiation of a related motor program. However, it is important to assess the cell-type specificity of this phenomenon, as

an equivalent change in inhibitory neurons in layer IV S1$_{BF}$ would mitigate increased excitatory gain. In other words, a generalized, network-wide increase in excitability could be self-stabilizing.

Barrel inhibitory neurons are unaffected by the onset of active whisking

To investigate the possibility that increased intrinsic excitability is a general feature of barrel cortex at P14, we sought to record F/I curves from parvalbumin-positive inhibitory interneurons (PV cells), as they represent the most common inhibitory subclass in the cortex [59, 69, 70, 71, 72]. However, the parvalbumin protein begins expression at approximately P14 [73], making it challenging to identify PV interneurons in animals around the onset of active whisking. To circumvent this, and label a subset of inhibitory interneurons in the cortex, we performed a genetic cross between two transgenic mouse lines. The first line, C57BL/6J-Tg(Nkx2-1-cre)2Sand/J [74], expresses Cre recombinase under the control of the Nkx2-1 promoter, which targets forebrain GABAergic interneurons. The second line, TRAP (EGFP/RPL10A [75]), expresses EGFP in the presence of Cre recombinase. This combination results in EGFP expression in Nkx2.1 positive cells. Crucially, Nkx2.1 is a transcription factor that participates in the differentiation of PV and somatostatin (SST) interneurons [54, 55] (Fig. 2A). Within the Nkx2.1-positive population, PV and SST interneurons can be differentiated by assessing the presence/absence of SST immunoreactivity (Fig. 2B) and by electrophysiological properties (Fig. 2C) [76].

We recorded F/I curves in Nkx2.1-positive inhibitory interneurons on P12, P14, and P16 in a total of 15 mice. In comparison to excitatory neurons, Nkx2.1-positive neurons had higher spike thresholds and continued to increase their activity at current injections above 200 pA without depolarization blockade. Thus, inhibitory F/I curves were assessed from 0 and 500 pA in steps of 20 pA (Fig. 2D inset). We divided putative PV and putative SST subpopulations based on accommodation (a progressive reduction in firing during a sustained stimulus) and firing rate per current injection. In stark contrast to excitatory neurons, putative PV interneurons showed no sign of increased excitability on P14 (2D), and there were no signs of specific changes in passive properties on P14 (Fig. S2). These data suggest that the plasticity observed in excitatory neurons is not only distinct in its singular regulation of membrane excitability, but is also cell-type specific.

A subset of electrophysiological features contribute to intrinsic plasticity at P14

Broadly, mechanisms of intrinsic excitability can be considered in three categories. First, passive conductances comprise ion channels that are constantly open (at rest) and allow ions to flow based on their electrochemical gradients. These channels maintain the resting membrane potential and contribute to the overall resistance and capacitance of the membrane, affecting how the cell responds to incoming stimuli without directly participating in the generation of action potentials. Second, active conductances comprise ion channels that open and close in response to changes in membrane potential. These conductances actively regulate the flow of ions across the membrane. They include voltage-gated channels like Na$^+$, K$^+$, and Ca^{2+} channels, which generate dynamic responses to depolarization. Third, spiking properties define the output signals of neurons and include action potential threshold, the frequency and pattern of firing, and the adaptability of response to sustained inputs.

To parse candidate mechanisms underlying the increase in intrinsic excitability in barrel excitatory neurons on P14, we extracted measures reflecting passive conductances (resting membrane potential, input resistance, membrane time constant—tau, and capacitance) (Fig. 3A-E), active conductances (dynamic input resistance and time to first spike) (Fig. 3F,G), and spiking features (after hyperpolarization, spike frequency adaptation, spike threshold, and instantaneous frequency)(Fig. 3H-J). We employed a predictive modeling approach to ask which, if any, of these three feature sets was sufficient to accurately classify the age of withheld neurons, with particular

attention to performance on P14. In simple terms, a computational model attempted to learn to predict an animal's age based only on a small set of electrophysiological features from a neuron. Trained on active conductances, logistic regression models accurately classified withheld P14 neurons at $76.86 \pm 1.02\%$ accuracy (chance = 16.7%) (Fig. 3K). Trained on passive properties, models accurately identified P14 neurons at $35.1.\pm 1.16\%$, and trained on spiking features, models performed at $40.9.\pm 1.2\%$ on P14 (Fig. 3L,M). The highest performance across all three models was P14 active conductances. Passive properties were most effective at predicting P11 (52%), while spiking features' effectiveness peaked on P16 at 56%. Undoubtedly, ion channel expression changes across development and as a feature of experience— assuming a stereotyped developmental timeline, it is unsurprising that a classifier can learn signatures to discriminate cells collected only 24 h apart. However, it is noteworthy that the only set of features that meaningfully differentiates neurons on P14 is the active conductances, suggesting that the underlying channels interact with the onset of whisker motor control.

Region- and layer- specificity of intrinsic excitability plasticity

Our findings thus far could be explained by two competing hypotheses. 1) The first posits that the increased excitability observed on P14 is an indirect response to enhanced sensory input. In other words, increased input through the whiskers drives feed-forward, Hebbian changes throughout the barrel cortex, and increased excitability is a component of this response [46, 47]. If this were the case, we would expect to see significant changes in synaptic strength throughout the $S1_{BF}$ microcircuit. However, such changes are notably absent in the synapses of layer IV (Fig. 2). 2) Alternatively, the increased excitability on P14 could reflect a coordinated developmental program that proactively shapes the functional properties of the barrel cortex input layer to coincide with the onset of active sensory input. Such an adaptation might serve to optimize information processing as the sensory environment becomes more complex. To distinguish between these possibilities, we addressed two key questions. First, do similar changes in excitability occur in layer II/III of the barrel cortex? and, second, does the sudden introduction of complex stimuli in another sensory modality, such as vision, elicit comparable plasticity? If either of these scenarios were confirmed, it would support the hypothesis that the observed changes are an indirect response to enhanced sensory input.

We recorded F/I curves in excitatory pyramidal neurons in S1 layer II/III (immediately dorsal to the layer IV barrels) on P12, P14, and P16 (N = 11 mice). There was a significant main effect of age ($p = 0.026$; Fig. 4A), however this was explained by a decrease in excitability from P12 to P14 ($p = 0.026$) and P12 to P16 ($p = 0.128$). These data suggest that the effects observed in layer IV do not generalize to the rest of the barrel cortical circuit.

The visual system provides a similar opportunity to study the sudden introduction of complex stimuli into a cortical sensory network, as animals transition from a state of closed eyes to open in a relatively tight temporal window between P12 and P14 [77]. To ask if the intrinsic excitability plasticity observed in layer IV $S1_{BF}$ excitatory neurons is a general phenomenon of a such a binary shift, we recorded F/I curves in layer IV excitatory pyramidal neurons in binocular primary visual cortex ($V1_b$) at P12, 13, 14, 16, 19, and 20 (N = 19 mice). In this context, there was a significant main effect of age ($p = 0.029$), although none of the post hoc pairwise comparisons reached significance. P14 vs P20 was the closest contrast to significance ($p = 0.075$); the mean F/I slope on P14 was lower than on P20.

Because eye opening can occur over a range of days within the window of our recordings, we aligned $V1_b$ neurons based on each animal's eye-opening status to understand if this, not age, could recapitulate the results from $S1_{BF}$. F/I curve slopes did not differ significantly as a function of the

status of eye-opening ($p = 0.173$; Fig. 4C).

Taken together, these results demonstrate a) increased excitability on P14 is not a general feature of cortical excitatory neurons, b) the sudden onset of organized input into $S1_{BF}$ does not cause plasticity of intrinsic excitability in layers II/III, and c) the sudden onset of organized visual information does not drive intrinsic excitability plasticity in excitatory layer IV neurons in ($V1_b$).

Intrinsic plasticity on P14 is experience dependent

Critical periods for postnatal cortical plasticity are profoundly influenced by sensory experiences [78]. To investigate whether passive whisker input prior to P14 is required for the observed intrinsic plasticity in layer IV of the $S1_{BF}$, we conducted unilateral whisker plucking at P10, 12, 14, 16, and 18 across a cohort of 32 mice. Subsequent F/I curves were recorded on P11, P12, P14, P16, P18, and P19, as depicted in (Fig. 5A). Importantly, whisker plucking preserves the follicles and underlying whisker musculature, maintaining the integrity of the sensory apparatus while isolating the effects of tactile input, and the contralateral whiskers are unperturbed [17].

In the context of sensory deprivation, layer IV $S1_{BF}$ excitatory F/I curves revealed a significant main effect of age ($p = 3.86e - 07$). This comprised a lessening of intrinsic excitability across development, exemplified by the contrast of P19 (the least excitable) to P11 (the most excitable; $p = 0.0000001$; Fig. 5B). Comparing intact to deprived, there was a near-significant main effect of sensory condition (0.083), and a significant interaction of sensory condition and age ($p = 0.006$). Given the focused hypothesis on P14, a two-sample t-test was employed to compare firing rate at 100 pA between intact and deprived sensory conditions at this age— whisker plucking significantly reduced the excitability of layer IV barrel excitatory neurons on P14 ($p = 0.040$; Fig. 5C). In the full analysis (ANOVA), the only intact vs deprived comparison that was near significant was on P11 ($p = 0.069$). This is remarkably different than the increased excitability observed on P14 with intact whiskers (Fig. 1E).

The effect on deprived P11 in consistent with a rapid homeostatic response to offset the impact of deprivation [52, 49, 59]. Remarkably, despite the increase in excitability, we observed no concomitant change in mEPSC amplitude on P11 after 24 h of deprivation (p = 0.9976; Fig. S3). Taken together, these data suggest three things; first, that the intrinsic plasticity on P14 requires prior experience, second, that whisker deprivation drives an acute homeostatic increase in excitability, and third, there is a notable absence of synaptic homeostasis in post-critical-period layer IV $S1_{BF}$.

We next evaluated the impact of whisker deprivation in layer II/III $S1_{BF}$ excitatory neurons on P12, P14, and P16 in N = 12 mice. Neurons in the deprived hemisphere exhibited a significant main effect of age ($p = 0.00124$) in which intrinsic excitability declined over time, as observed under intact conditions— there was no evidence of an aberration at P14. In layer II/III, the main effect of sensory condition was also near near-significant ($p = 0.085$), and consistent with a homeostatic increase in excitability following whisker plucking. In contrast to layer IV, there was no interaction of age and sensory condition ($p = 0.545$), suggesting that the effect of whisker plucking was consistent from P12-16 in layers II/III. Across layers II/III and IV, the only condition in which intact excitability was greater than deprived excitability was on P14 in layer IV (Fig. 5C,D).

These data raise the question of how experience modifies neuronal excitability. One possibility is that the mechanisms underlying the layer IV plasticity on P14— active conductances (Fig. 3K-M)—are suppressed in the absence of passive input beginning at P10. In this case, we would expect that a classifier trained on active conductances would fail to meaningfully identify P14. Alternatively, it is possible that whisker plucking mutes the P14 effect by reducing other features, such as passive properties. In this case, P14 should still be readily identifiable in terms of active

conductances, but a corresponding, counteractive change in another feature set should also emerge. To disambiguate these, we extracted measurements of passive, active, and spiking features from F/I recordings in deprived hemispheres between P11 and P19. Trained on active conductances, age was predicted with a balanced accuracy of $34.84 \pm 0.35\%$, and P14 was classified with $30.4 \pm 0.948\%$ accuracy (Fig. 5E). At P11, which was remarkable for its apparent homeostatic increase following 24 h of deprivation, active-conductances provided 64% classification accuracy. Passive and spiking features supported balanced accuracies of $34.31 \pm 00.34\%$ and $33.67 \pm 0.37\%$, respectively, although in each case performance on P14 was below chance ($6.83 \pm 0.58\%$ and $8.35 \pm 0.64\%$, respectively; Fig. 5F).

Taken together, these data suggest that plasticity of active conductances on P14 requires experience, and although driven by an opposite sensory manipulation, the homeostatic response to deprivation acts through a similar set of underlying conductances.

Discussion

At P14, mice acutely transition from passive to active whisker use for exploration of objects and the environment [32, 79]. As a result, study of the barrel cortex at this key behavioral shift stands to provide insight into the interplay of primary sensory cortical circuits and the development of the motor programs. The motor control of sensory organs is a fundamental component of sensation that impacts on the structure of information arriving in the cortex. As a result, the sudden onset of sensory organ motor control might be hypothesized to cooperate with plastic events in corresponding sensory cortices. Here, we recorded synaptic and intrinsic properties of neurons in the barrel cortical microcircuit throughout a period of postnatal development encompassing the onset of active whisking. We describe a non-synaptic form of plasticity in excitatory neurons of the input layer (IV) to S1$_{BF}$, whose locus is consistent with rapid regulation of active conductances. Summarily, this population of neurons, whose inputs include excitatory afferents from VPM and POm thalamus [72, 80, 81], significantly increases their input/output ratio for approximately 24 h on P14.

While the critical period for layer IV barrel structure is closed by P14 [82, 83, 18, 24], the critical period for Hebbian modification of intracortical processing in layers II/III opens after the barrels are formed and persists through the onset of active whisking [22, 84, 33] (Fig. 1B). Our finding that layer IV synaptic properties are unchanged at P14 (Fig. 1F) is consistent with layer IV barrels having already exited their critical period. In other words, at P14, the anatomical structure and synaptic architecture of the barrels are established, and active whisking is poised to drive refinement of intracortical connectivity in layers II/III—a minimum of two synapses away from the thalamus [80]. This sets up an interesting problem for developing sensory circuits. External input is necessary to drive experience-dependent circuit refinement, but cortical gain control and inhibition by ongoing activity constrains the impact of whisker-driven input in upper layers [85, 86, 87, 88]. Speculatively, the elevated excitability of excitatory (but not inhibitory) neurons in layer IV on P14 is well positioned to briefly increase the gain of feedforward input to layers II/III, precisely when motor-controlled whisker input is required to drive Hebbian change.

A plastic change in intrinsic properties without a corresponding shift in synaptic strength is uncommon. Intrinsic plasticity is a key feature of both homeostatic and Hebbian plasticity [52, 45, 49, 46, 89]. However, intrinsic plasticity typically co-occurs with a variety of synaptic changes including synaptic scaling, LTP, and LTD. Put simply, when intrinsic plasticity is observed, synaptic modifications also occur as part of a broader adaptive response to environmental or experimental conditions. Against this backdrop, our results suggest that these two loci of plastic change—synaptic and intrinsic—are dissociable and capable of independent regulation. We offer two plausible scenarios

that could give rise to our data. First, at P14, the lack of synaptic change implies active constraint or compensation. Alternatively, the threshold for activating intrinsic plasticity might simply be lowered at P14, while developmental programs and changing external input are insufficient to drive synaptic change. Our data provide possible clues to resolve this.

Prior work demonstrates that whisker plucking/trimming drives homeostatic synaptic scaling in layers II/III [60], layer IV [90], and layer V [91] of barrel cortex. However, these observations involve animals older than P28, past the closure of all critical periods in barrel cortex. In our data, whisker plucking failed to elicit a homeostatic response in synaptic strength, but prominently increased intrinsic excitability in the first 24 h, consistent with a compensatory response. Changes in intrinsic excitability are a component of the classical homeostatic response to perturbation, which normally includes synaptic compensation [52, 49]. Taken with the exclusively intrinsic plasticity at P14 and prior observations of synaptic homeostasis after P28, these data suggest that synaptic modifiability is temporarily suppressed in layer IV $S1_{BF}$ excitatory neurons around the time period wherein animals begin active whisking.

A comparison of the increased excitability at P14 (under intact conditions) and the compensatory increase in excitability at P11 (following 24 h of deprivation) is worthwhile. Each observation lacks measurable synaptic change. Further, electrophysiologically, it appears that active conductances are the primary mechanism by which excitability is modified in each event. However, the stimulus-response logic is inverted between the two. In the case of whisker plucking, diminished input drives a compensatory increase in excitability— this reflects the principles of maintaining a set-point in the face of perturbation. In the case of the onset of active whisking, however, increased excitability is poised to amplify input in the context of increased complexity and frequency. This suggests that, in the second week of life, intrinsic properties are actively regulated by the barrel cortical circuit for both homeostatic (negative feed back) and non-homeostatic ends.

A central question that we are unable to resolve in our data is how the intrinsic plasticity on P14 is initiated. One possibility is anticipatory. In this case, increased excitability at P14 is the result of a developmental program whose timing requires prior input. This explanation is parallel to the timing of key plastic events in the visual system by prior experience [78]. Alternatively, the intrinsic plasticity on P14 is responsive. In this case, a highly plastic set of conductances react to active whisking-driven input via positive feedback, resulting in increased excitability. Future work could endeavor to disambiguate these explanations by plucking whiskers at P13, potentially allowing for the assembly and normal timing of an anticipatory program. Alternatively, experimental imposition of complex whisker stimulation prior to P14 would be expected to engage a reactive mechanism.

Two further limitations should be taken into consideration. First, in addition to thalamic afferents, excitatory neurons in layer IV $S1_{BF}$ receive substantial input from other layer IV excitatory neurons [92]. As a result, the stimulus-related impact of increased intrinsic excitability in this population cannot be ascertained in ex vivo slice recordings. While our data rule out parallel changes in inhibitory neurons, the computational impact of increased excitability in the spiny stellate (excitatory) population is likely not linear as it impacts both feedforward and local recurrent connections. And second, without preventing the intrinsic plasticity on P14, it is only possible to speculate on possible developmental roles. While disrupting specific developmental events can be challenging, (see, for example, visual critical periods), modeling and careful perturbations can offer empirical insight. From this perspective, it will be interesting to ask whether P14 plasticity is evident in transgenic models of neurodevelopmental disorders.

We sought to test the possibility of a interaction between the development of sensory organ motor control and corresponding primary sensory cortices. Our results suggest an unusual decoupling

of synaptic and intrinsic plasticity that results in increased gain in layer IV excitatory neurons. This effect is transient, specific to cell type, layer, and region, and requires prior experience. Together, our data indicate that motor control development may influence sensory cortical development via plastic regulation of gain control.

Acknowledgments

This work was supported by NIH BRAIN Initiative 1R01NS118442-01 (KBH), NIH/NINDS R00NS089800 (KBH), R01NS099429 (JPC), and R01MH124808 (JDD).

We thank Drs. Arianna Maffei and Steve Mennerick for their insights and conversations about this research.

Author Contributions

MCS, KL, and KBH developed the conceptualization and methodology of the experiments. MCS, LT, HL, SC, and MEL collected the data. LT, MCS, and KBH analyzed the data. KBH, JDD, and JPC provided resources. LT, MCS, and KBH produced the figures and wrote the paper.

Methods

Animals

All surgical techniques and experimental procedures were conducted in accordance with protocols approved by the Washington University in St. Louis Institutional Animal Care and Use Committee, following the National Institutes of Health guidelines for the care and use of research animals. Mouse lines used in this study are: C57BL/6, C57BL/6J-Tg(Nkx2-1-cre)2Sand/J [74], and TRAP (EGFP/RPL10A) [75].

Whisker Deprivation

Postnatal day 10 (P10) mice were anesthetized under transient isoflurane and whiskers on one side of the face were gently removed using forceps similar to previously described [18]. Whisker plucking is a robust manipulation that removes the entire whisker but does not cause damage to the follicles. Removal of whisker growth was repeated every other day until the day of recording.

Slice preparation

Mice aged postnatal day 11 (P11) to P20 of either sex were anesthetized with isoflurane and decapitated. The brain was rapidly removed and submerged in ice cold, oxygenated artificial CSF (aCSF). aCSF was continuously oxygenated and contained the following (in mM): 126 NaCl, 3 KCl, 2 MgSO4, 1 NaPO4, 25 NaHCO3, 2 CaCl2 and 14 dextrose, oxygenated with 5% CO_2/ 95% O_2. For barrel cortex recordings, coronal brain slices (325 μm) containing S1 were taken in the "across-row" plane, oriented at a 50°angle toward coronal from the midsagittal plane. With this technique, each slice containing the posteromedial barrel subfield (PMBSF) includes one barrel column from each of the five whisker barrel rows (A through E), allowing for each column to be identified in slices and used for targeting the electrophysiological recordings [93, 94].

For visual cortex recordings, coronal brain slices (325 μm) containing V1 were taken as previously described [49]. Slices were incubated at 37 °C for 30 min and then kept at room temperature and oxygenated with 95/5 until recording (0.5-7hrs).

Electrophysiology

Recordings were performed at 33 °C, with constant oxygenation (95/5 O_2/CO_2) and circulation of aCSF. Borosilicate glass recording pipettes with tip resistances of 3-6 MΩ were pulled using a Sutter P-2000 micropipette puller. Standard internal solution contained (mM): 20 KCl, 100 K-gluconate, 10 HEPES, 4 Mg-ATP, 0.3 Na-GTP, 10 phosphocreatine, and 0.4% biocytin. Barrel cortex recordings were performed within slice regions where PMBSF structure could be clearly visualized under a 10x air objective. Neurons were visualized with a 40x immersion objective using infrared differential contrast imaging (IR-DIC). Excitatory cells in layers IV and II/III were identified based on morphology (stellate in layer IV, pyramidal in layer II/III) [26, 95]. PV+ and SST+ interneurons were identified using EGFP expression driven by the NXK2.1-cre allele [96, 97]. V1 was identified using the shape and morphology of ventricles and the white matter, and recordings were obtained from layer IV pyramidal neurons based on morphology [95]. F/I curves were recorded in standard aCSF containing the channel blockers APV (50 μM), DNQX (20 μM) and picrotoxin (20 μM) in current-clamp mode. A small DC bias current was injected to maintain Vm at -70 mV between depolarizations. For excitatory F/I curves, current injections of increasing magnitude were injected from 0-180 pA with steps of 20 pA. For inhibitory F/I curves, a firing threshold was first determined using brief current injection. Current injections were then injected in 20 pA steps from threshold to 600 pA. For AMPA miniature EPSC (mEPSC) recordings, neurons were voltage clamped at -70 mV in standard aCSF containing the channel blockers APV (50 μm), picrotoxin (20 μM) and TTX (0.2 μM). Neurons were not included in analyses if the resting membrane potential was more positive than -55 mV, input resistance was <80 MΩ, series resistance was >20 MΩ, or if any of these parameters changed >20% during the recording. Morphology, as well as location in the layer was confirmed by post hoc reconstruction of biocytin fills. The first 100 individual mEPSCs were isolated from recordings using the Event Search function of pClamp. All data were further analyzed using custom Python scripts.

Histology

Recorded excitatory and inhibitory neurons were filled with biocytin (0.4% in internal solution) and slices were fixed overnight in 4% paraformaldehyde PBS. Slices were then washed in PBS, incubated in 5% NGS, 1% Triton X-100 block solution for 2 h, and incubated overnight at 4°C with AlexaFluor-568 streptavidin (1:2000, Invitrogen) and anti-somatostatin antibody (1:500, Abcam) in block solution. On the second day, slices were washed in PBS and incubated with Goat Anti-Rabbit AlexaFluor-405 (1:500, Abcam) in block solution for 1 h, 0.1% TO-PRO-3 in PBS solution for 30 minutes, then mounted on glass slides with Fluoromount and coverslipped. Slices were imaged with a fluorescent microscope (Thunder, Leica).

Logistic Regression Classification

To predict an animal's age based on electrophysiological features and assess the impact of sensory deprivation, we performed logistic regression analysis using Python. The analysis was conducted separately for three feature sets: active conductances (dynamic input resistance and time to first spike), passive properties (membrane resistance, membrane potential, capacitance, and tau), and spiking features (afterhyperpolarization, spike frequency adaptation, and instantaneous frequency).

Data Preprocessing: The dataset was first grouped by age and balanced by randomly sampling an equal number of cells (n) from each age group to ensure class balance. The balanced dataset was then split into training (80%) and testing (20%) sets using stratified sampling based on age. The features were standardized using the StandardScaler from the sklearn library to ensure all features had zero mean and unit variance.

Model Training and Evaluation: A multinomial logistic regression model (LogisticRegression from the sklearn library) was trained on the standardized training set. The model's performance was evaluated on the held-out test set using 1) F1 score (macro average): The harmonic mean of precision and recall, calculated for each class and then averaged, providing an overall measure of the model's accuracy, and 2) Balanced accuracy: The average of the proportion of correct predictions within each class, taking into account class imbalance.

The logistic regression analysis was repeated for 500 iterations (reps) to ensure robustness of the results. In each iteration, the dataset was resampled, and the model was retrained and evaluated. The mean and standard error of the metrics across iterations were reported.

Comparison of Sensory Conditions: The logistic regression analysis was performed separately for intact and deprived sensory conditions. The accuracy in predicting age 14 (the onset of active whisking) was compared between the two conditions to assess the impact of sensory deprivation on the model's performance. The mean accuracy and standard error for age 14 were calculated across iterations.

Statistics

Statistical analyses were performed using R and packages tidyverse (version 2.0.0) and stats (version 4.2.3).

Data Preparation: Variables 'animal id', 'age', and 'cell id' were converted to factors. Data were filtered to include a range of current injections in which cells were active and not saturated (e.g., between 20 and 100 pA for excitatory layer IV $S1_{BF}$).

Slope Calculation: In the context of F/I curves, for each neuron, the slope of the frequency-current relationship was calculated using linear regression (lm function from the stats package) with 'frequency' as the dependent variable and 'current' as the independent variable.

Analysis of Variance (ANOVA): To assess the effect of age on the frequency-current slope, a mixed-effects ANOVA was performed using the aov function from the stats package. The model included 'age' and 'animal id' as fixed factors, with 'slope' as the dependent variable (or a neuronal property, such as capacitance where appropriate). The ANOVA model was specified as follows:

$slope\ factor(age) + factor(animalid)$

Post-hoc Tests: If the ANOVA indicated a significant main effect of, e.g., age, post-hoc tests were performed using Tukey's Honest Significant Difference (HSD) test.

Figures

Figure 1: At the onset of active whisking, spiny stellate neurons in layer 4 barrel cortex exhibit a transient increase in excitability. A) Experimental approach. Ex vivo slices were taken from the barrel field of primary somatosensory cortex (SS_{BF}) from postnatal day 11 (P11) through P19. B) Layer IV barrels were visualized under differential interference contrast imaging (DIC) at 10x and individual excitatory cells were chosen under 40x. After recording, cell morphology and location were visualized using streptavidin stains and confocal microscopy. C) Timeline of experiment. Outline of relevant critical periods for structural and synaptic plasticity as well as the timeline of recordings and the onset of active whisking. D) Location of intracellular patch recordings in SS_{BF}. E) F/I curves reveal a one day, transient, increase in the intrinsic excitability of Layer IV excitatory cells on P14 that coincides with the onset of active whisking, right. Example traces from each age, left. F) mEPSC recorded from Layer IV excitatory cells do not show an accompanying synaptic strength change, right. Example mEPSC traces from each age, left. Printed p values are for the main effect of age. For F/I curves N=26 mice, N=108 cells. For mEPSCs N=22 mice, N=94 cells. * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001

Figure 2: Increased excitability at the onset of active whisking is specific to excitatory neurons.
A) Experimental approach. To fluorescently label a subset of isocortical inhibitory interneurons, Nkx2.1-Cre mice were crossed with TRAP mice, which express EGFP-tagged ribosomal protein L10a under a Cre-dependent promoter. (Top) In this context, Nkx2.1 positive neurons in Layer IV barrels were recorded on postnatal day 12 (p12) through p16. (Bottom) Subsequent staining for somatostatin (SST) protein allowed for the separation of three cell types in our dataset: excitatory, parvalbumin (NKX positive, SST negative), and SST (NKX and SST positive). B. Example triple label image demonstrating the separation of the three putative cell types. Streptavidin is loaded in the recording electrode and thus labels cells that were patched (intensity reflects the duration of patch). Nkx2.1 labels PV and SST interneurons in the isocortex. Somatostatin label is used to define SST interneurons. Merge demonstrates the molecular logic of putative cell type identification. C. PV and SST subpopulations of Nkx2.1 positive neurons are differentiable based on electrophysiological properties, accommodation and the peak instantaneous firing rate. (Top) Kernel density estimate and scatter of Nkx2.1 positive neurons' accommodation and max firing, colored by PV/SST classification. (Bottom) Example traces from an excitatory neuron (E) and putative PV interneuron in response to a 160 pA current step. D. F/I curve of putative PV interneurons in Layer IV barrel cortex as a function of age reveals no increase in intrinsic excitability on p14. For F/I curves, N=15 mice, N=40 cells.

Figure 3: Plasticity in active conductances underlies increased excitability on P14. A) Illustration of recording from excitatory neurons in Layer 4 of the barrel fields of primary somatosensory cortex (SS_{BF}). Legend indicates recorded ages and applies to the B-J. B-E) Passive properties of SS_{BF} L4 excitatory neurons as a function of age: B) resting membrane potential, C) input resistance (measured with a small hyperpolarizing current step), D) membrane time constant (tau), and E) membrane capacitance. F) Dynamic input resistance as a function of age. Note that this is a measure of the change in membrane resistance in response to a small positive current, in this case a 40 pA pulse. The voltage deflection is divided by the current (40 pA)— higher values suggest a less leaky membrane. G) Time to first spike as a function of age. H) Afterhyperpolarization (AHP) amplitude as a function of age. I) Spike threshold as a function of age. J) Spike frequency adaptation as a function of age— this refers to a decrease in firing rate over the course of a square wave. K-M) Logistic regression was used to predict animal age as a function of three features sets measured in each neuron: active conductances (dynamic input resistance and time to first spike), passive properties (membrane resistance, membrane potential, capacitance, tau), and spiking features (AHP, spike frequency adaptation, and instantaneous frequency). K) Confusion matrix— correctly classified neurons fall on the diagonal— for active conductances. Note that chance is 16.67% and that P14 is robustly identifiable. L) Same as K but for passive properties. M) Same as K but for spiking features.

Figure 4: Increased excitability at P14 is not a general property of sensory cortices A) Inset: Illustration of recording from excitatory neurons in Layer 2/3 of primary somatosensory cortex. Excitatory F/I curves for ages P12, P14, and P16. At P14, L2/3 neurons exhibit reduced excitability compared to P12. B) Excitatory F/I curve of excitatory L4 neurons in the binocular portion of primary visual cortex ($V1_b$) for ages P12, P13, P14, P16, P19, and P20. At P14, neuronal excitability in $V1_b$ is unremarkable. C) Mouse eye opening occurs between P10 and P14. F/I curve showing the data presented in B as a function of whether an animal's eyes were yet opened.For Layer 2/3 F/I curves, N=11 mice, N=47 cells. For V1 F/I curves, N=19 mice, N=168 cells.

Figure 5: Increase in intrinsic excitability is experience dependent and is blocked by early whisker deprivation. A) Timeline of whisker deprivation. Mice were unilaterally whisker plucked starting at 10 days postnatal (P10) and plucking continued every other day until the day of recording. Ex vivo slices were taken from the barrel field of primary somatosensory cortex (SS_{BF}) from postnatal day 11 (P11) through P19 B) F/I curves from Layer IV excitatory cells in the deprived hemisphere do not show the transient increase in intrinsic excitability indicating that this change is experience dependent, insets show the location of intracellular recordings, left, and current injection traces, right. C) Comparing F/I curves for the intact vs deprived hemisphere at each age shows a one day increase in intrinsic excitability in the deprived hemisphere on P11, top left. These comparisons also show that intrinsic excitability in the deprived hemisphere is lower than in the intact hemisphere, further suggesting this change is experience dependent. * indicates p ¡ 0.05; two-sample t-test of firing frequency at 100 pA current injection. D) F/I curves from Layer II/III excitatory cells in the deprived hemisphere do not show an increase in excitability on P14. Intrinsic excitability in deprived hemisphere Layer II/III cells is slightly higher than in the control hemisphere, and overall excitability decreases with age. E) Confusion matrix showing the ability of a logistic regression classifier to predict an animal's age based on active conductances in the deprived hemisphere (active conductances comprise dynamic input resistance and time to first spike; correct classifications fall along the diagonal). Note that chance is 16.67% and that P14 is minimally identifiable. F) Classifier balanced accuracy as a function of feature set (gray: active conductances, teal: passive properties, green: spiking features) and condition (intact: solid, deprived: open). Chance is indicated by the dashed red line. For Layer 4 F/I curves, N=32 mice, N=142 cells. For Layer 2/3 F/I curves, N=12 mice, N=40 cells.

Figure S1: Passive membrane properties and mEPSC waveform analysis of layer IV excitatory neurons in the barrel cortex. (A-D) Passive membrane properties of layer IV excitatory neurons as a function of age: (A) membrane time constant (tau), (B) input resistance, (C) membrane capacitance, and (D) resting membrane potential. (E) Mean mEPSC waveforms from layer IV excitatory neurons across development (P11-P18). (F) Kernel density estimate plot showing the the first two principal components of PCA run on the mean waveforms shown in E. Color indicates age as in E. (G) Confusion matrix for a logistic regression model trained to predict age based on the first three principal components of mEPSC waveforms. (H) Loadings for the first principal component (PC1), illustrating the temporal features of the mEPSC waveform that contribute most to the variance across development.

Figure S2: Passive membrane properties of putative parvalbumin (PV) and somatostatin (SST) interneurons in layer IV of the barrel cortex. Passive membrane properties of putative PV and SST interneurons as a function of age (P12, P14, and P16): (Left) membrane time constant (tau), (Center) input resistance, and (Right) membrane capacitance.

Figure S3: Whisker deprivation does not induce homeostatic synaptic scaling in layer IV excitatory neurons at P11. Comparison of mEPSC amplitudes between intact and deprived hemispheres in layer IV excitatory neurons at P11, 24 h after the onset of unilateral whisker deprivation. No significant difference was observed between the two conditions (p = 0.9976), indicating an absence of homeostatic synaptic scaling at this age and time point.

References

[1]   A. Berti et al. "Shared Cortical Anatomy for Motor Awareness and Motor Control." In: *Science* 309.5733 (July 2005), pp. 488–491. ISSN: 1095-9203. DOI: 10.1126/science.1110625. URL: http://dx.doi.org/10.1126/science.1110625.

[2]   Charles D. Gilbert and Wu Li. "Top-down influences on visual processing." In: *Nature Reviews Neuroscience* 14.5 (Apr. 2013), pp. 350–363. ISSN: 1471-0048. DOI: 10.1038/nrn3476. URL: http://dx.doi.org/10.1038/nrn3476.

[3]   Stanislas Dehaene and Jean-Pierre Changeux. "Experimental and Theoretical Approaches to Conscious Processing." In: *Neuron* 70.2 (Apr. 2011), pp. 200–227. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2011.03.018. URL: http://dx.doi.org/10.1016/j.neuron.2011.03.018.

[4]   Michael N. Shadlen and William T. Newsome. "Neural Basis of a Perceptual Decision in the Parietal Cortex (Area LIP) of the Rhesus Monkey." In: *Journal of Neurophysiology* 86.4 (Oct. 2001), pp. 1916–1936. ISSN: 1522-1598. DOI: 10.1152/jn.2001.86.4.1916. URL: http://dx.doi.org/10.1152/jn.2001.86.4.1916.

[5]   Timothy J. Buschman et al. "Synchronous Oscillatory Neural Ensembles for Rules in the Prefrontal Cortex." In: *Neuron* 76.4 (Nov. 2012), pp. 838–846. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2012.09.029. URL: http://dx.doi.org/10.1016/j.neuron.2012.09.029.

[6]   Zhengyu Ma et al. "Cortical Circuit Dynamics Are Homeostatically Tuned to Criticality In Vivo." In: *Neuron* 104.4 (Nov. 2019). PMCID: PMC6934140, 655–664.e4. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2019.08.031. URL: http://dx.doi.org/10.1016/j.neuron.2019.08.031.

[7]   Yifan Xu et al. "Sleep restores an optimal computational regime in cortical networks." In: *Nature Neuroscience* (Jan. 2024). ISSN: 1546-1726. DOI: 10.1038/s41593-023-01536-9. URL: http://dx.doi.org/10.1038/s41593-023-01536-9.

[8]   Collin Blakemore and Grahame F. Cooper. "Development of the Brain depends on the Visual Environment." In: *Nature* 228.5270 (Oct. 1970), pp. 477–478. ISSN: 1476-4687. DOI: 10.1038/228477a0. URL: http://dx.doi.org/10.1038/228477a0.

[9]   Takao K. Hensch and Michael P. Stryker. "Columnar Architecture Sculpted by GABA Circuits in Developing Cat Visual Cortex." In: *Science* 303.5664 (Mar. 2004), pp. 1678–1681. ISSN: 1095-9203. DOI: 10.1126/science.1091031. URL: http://dx.doi.org/10.1126/science.1091031.

[10]   Michela Fagiolini and Takao K. Hensch. "Inhibitory threshold for critical-period activation in primary visual cortex." In: *Nature* 404.6774 (Mar. 2000), pp. 183–186. ISSN: 1476-4687. DOI: 10.1038/35004582. URL: http://dx.doi.org/10.1038/35004582.

[11]   Keith B. Hengen et al. "Firing Rate Homeostasis in Visual Cortex of Freely Behaving Rodents." In: *Neuron* 80.2 (Oct. 2013), pp. 335–342. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2013.08.038. URL: http://dx.doi.org/10.1016/j.neuron.2013.08.038.

[12]   Keith B. Hengen et al. "Neuronal Firing Rate Homeostasis Is Inhibited by Sleep and Promoted by Wake." In: *Cell* 165.1 (Mar. 2016), pp. 180–191. ISSN: 0092-8674. DOI: 10.1016/j.cell.2016.01.046. URL: http://dx.doi.org/10.1016/j.cell.2016.01.046.

[13]   Alessandro Sale, Nicoletta Berardi, and Lamberto Maffei. "Environment and Brain Plasticity: Towards an Endogenous Pharmacotherapy." In: *Physiological Reviews* 94.1 (Jan. 2014), pp. 189–234. ISSN: 1522-1210. DOI: 10.1152/physrev.00036.2012. URL: http://dx.doi.org/10.1152/physrev.00036.2012.

[14]   Martha J. Farah. "The Neuroscience of Socioeconomic Status: Correlates, Causes, and Consequences." In: *Neuron* 96.1 (Sept. 2017), pp. 56–71. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2017.08.034. URL: http://dx.doi.org/10.1016/j.neuron.2017.08.034.

[15]   Joan Luby et al. "The Effects of Poverty on Childhood Brain Development: The Mediating Effect of Caregiving and Stressful Life Events." In: *JAMA Pediatrics* 167.12 (Dec. 2013), p. 1135. ISSN: 2168-6203. DOI: 10.1001/jamapediatrics.2013.3139. URL: http://dx.doi.org/10.1001/jamapediatrics.2013.3139.

[16]   Thomas A. Woolsey and Hendrik Van der Loos. "The structural organization of layer IV in the somatosensory region (S I) of mouse cerebral cortex." In: *Brain Research* 17.2 (Jan. 1970), pp. 205–242. ISSN: 0006-8993. DOI: 10.1016/0006-8993(70)90079-x. URL: http://dx.doi.org/10.1016/0006-8993(70)90079-x.

[17]   Thomas A. Woolsey and Janice R. Wann. "Areal changes in mouse cortical barrels following vibrissal damage at different postnatal ages." In: *Journal of Comparative Neurology* 170.1 (Nov. 1976), pp. 53–66. ISSN: 1096-9861. DOI: 10.1002/cne.901700105. URL: http://dx.doi.org/10.1002/cne.901700105.

[18]   K Fox. "A critical period for experience-dependent synaptic plasticity in rat barrel cortex." In: *The Journal of Neuroscience* 12.5 (May 1992), pp. 1826–1838. ISSN: 1529-2401. DOI: 10.1523/jneurosci.12-05-01826.1992. URL: http://dx.doi.org/10.1523/JNEUROSCI.12-05-01826.1992.

[19]   M E Diamond, M Armstrong-James, and F F Ebner. "Experience-dependent plasticity in adult rat barrel cortex." In: *Proceedings of the National Academy of Sciences* 90.5 (Mar. 1993), pp. 2082–2086. ISSN: 1091-6490. DOI: 10.1073/pnas.90.5.2082. URL: http://dx.doi.org/10.1073/pnas.90.5.2082.

[20]   K Fox. "The cortical component of experience-dependent synaptic plasticity in the rat barrel cortex." In: *The Journal of Neuroscience* 14.12 (Dec. 1994), pp. 7665–7679. ISSN: 1529-2401. DOI: 10.1523/jneurosci.14-12-07665.1994. URL: http://dx.doi.org/10.1523/JNEUROSCI.14-12-07665.1994.

[21]   K Fox et al. "Glutamate receptor blockade at cortical synapses disrupts development of thalamocortical and columnar organization in somatosensory cortex." In: *Proceedings of the National Academy of Sciences* 93.11 (May 1996), pp. 5584–5589. ISSN: 1091-6490. DOI: 10.1073/pnas.93.11.5584. URL: http://dx.doi.org/10.1073/pnas.93.11.5584.

[22]   Daniel E. Feldman and Michael Brecht. "Map Plasticity in Somatosensory Cortex." In: *Science* 310.5749 (Nov. 2005), pp. 810–815. ISSN: 1095-9203. DOI: 10.1126/science.1115807. URL: http://dx.doi.org/10.1126/science.1115807.

[23]   J. Lubke. "Morphometric Analysis of the Columnar Innervation Domain of Neurons Connecting Layer 4 and Layer 2/3 of Juvenile Rat Barrel Cortex." In: *Cerebral Cortex* 13.10 (Oct. 2003), pp. 1051–1063. ISSN: 1460-2199. DOI: 10.1093/cercor/13.10.1051. URL: http://dx.doi.org/10.1093/cercor/13.10.1051.

[24] Li-Jen Lee et al. "Neonatal whisker trimming causes long-lasting changes in structure and function of the somatosensory system." In: *Experimental Neurology* 219.2 (Oct. 2009), pp. 524–532. ISSN: 0014-4886. DOI: 10.1016/j.expneurol.2009.07.012. URL: http://dx.doi.org/10.1016/j.expneurol.2009.07.012.

[25] Frank Lambert Rice et al. "AComparative analysis of the development of the primary somatosensory cortex: Interspecies similarities during barrel and laminar development." In: *Journal of Comparative Neurology* 236.4 (June 1985), pp. 477–495. ISSN: 1096-9861. DOI: 10.1002/cne.902360405. URL: http://dx.doi.org/10.1002/cne.902360405.

[26] Miguel Maravall, Edward A. Stern, and Karel Svoboda. "Development of Intrinsic Properties and Excitability of Layer 2/3 Pyramidal Neurons During a Critical Period for Sensory Maps in Rat Barrel Cortex." In: *Journal of Neurophysiology* 92.1 (July 2004), pp. 144–156. ISSN: 1522-1598. DOI: 10.1152/jn.00598.2003. URL: http://dx.doi.org/10.1152/jn.00598.2003.

[27] David Souto and Dirk Kerzel. "Visual selective attention and the control of tracking eye movements: a critical review." In: *Journal of Neurophysiology* 125.5 (May 2021), pp. 1552–1576. ISSN: 1522-1598. DOI: 10.1152/jn.00145.2019. URL: http://dx.doi.org/10.1152/jn.00145.2019.

[28] Roland S. Johansson and J. Randall Flanagan. "Coding and use of tactile signals from the fingertips in object manipulation tasks." In: *Nature Reviews Neuroscience* 10.5 (Apr. 2009), pp. 345–359. ISSN: 1471-0048. DOI: 10.1038/nrn2621. URL: http://dx.doi.org/10.1038/nrn2621.

[29] Joel Mainland and Noam Sobel. "The Sniff Is Part of the Olfactory Percept." In: *Chemical Senses* 31.2 (Dec. 2005), pp. 181–196. ISSN: 0379-864X. DOI: 10.1093/chemse/bjj012. URL: http://dx.doi.org/10.1093/chemse/bjj012.

[30] Jeffrey D. Moore et al. "Hierarchy of orofacial rhythms revealed through whisking and breathing." In: *Nature* 497.7448 (Apr. 2013), pp. 205–210. ISSN: 1476-4687. DOI: 10.1038/nature12076. URL: http://dx.doi.org/10.1038/nature12076.

[31] M. J. Z. Hartmann. "Active touch, exploratory movements, and sensory prediction." In: *Integrative and Comparative Biology* 49.6 (Nov. 2009), pp. 681–690. ISSN: 1557-7023. DOI: 10.1093/icb/icp107. URL: http://dx.doi.org/10.1093/icb/icp107.

[32] Hiroyuki Arakawa and Reha S. Erzurumlu. "Role of whiskers in sensorimotor development of C57BL/6 mice." In: *Behavioural Brain Research* 287 (July 2015), pp. 146–155. ISSN: 0166-4328. DOI: 10.1016/j.bbr.2015.03.040. URL: http://dx.doi.org/10.1016/j.bbr.2015.03.040.

[33] Reha S. Erzurumlu and Patricia Gaspar. "Development and critical period plasticity of the barrel cortex." In: *European Journal of Neuroscience* 35.10 (May 2012), pp. 1540–1553. ISSN: 1460-9568. DOI: 10.1111/j.1460-9568.2012.08075.x. URL: http://dx.doi.org/10.1111/j.1460-9568.2012.08075.x.

[34] Margo Landers and H. Philip Zeigler. "Development of rodent whisking: Trigeminal input and central pattern generation." In: *Somatosensory & Motor Research* 23.1–2 (Jan. 2006), pp. 1–10. ISSN: 1369-1651. DOI: 10.1080/08990220600700768. URL: http://dx.doi.org/10.1080/08990220600700768.

[35] Jenq-Wei Yang et al. "Thalamic Network Oscillations Synchronize Ontogenetic Columns in the Newborn Rat Barrel Cortex." In: *Cerebral Cortex* 23.6 (May 2012), pp. 1299–1316. ISSN: 1047-3211. DOI: 10.1093/cercor/bhs103. URL: http://dx.doi.org/10.1093/cercor/bhs103.

[36] S. An et al. "Long-Term Potentiation in the Neonatal Rat Barrel Cortex In Vivo." In: *Journal of Neuroscience* 32.28 (July 2012), pp. 9511–9516. ISSN: 1529-2401. DOI: 10.1523/jneurosci.1212-12.2012. URL: http://dx.doi.org/10.1523/JNEUROSCI.1212-12.2012.

[37] Kevin J. Bender, Juliana Rangel, and Daniel E. Feldman. "Development of Columnar Topography in the Excitatory Layer 4 to Layer 2/3 Projection in Rat Barrel Cortex." In: *The Journal of Neuroscience* 23.25 (Sept. 2003), pp. 8759–8770. ISSN: 1529-2401. DOI: 10.1523/jneurosci.23-25-08759.2003. URL: http://dx.doi.org/10.1523/JNEUROSCI.23-25-08759.2003.

[38] Chiaki Itami and Fumitaka Kimura. "Developmental Switch in Spike Timing-Dependent Plasticity at Layers 4–2/3 in the Rodent Barrel Cortex." In: *The Journal of Neuroscience* 32.43 (Oct. 2012), pp. 15000–15011. ISSN: 1529-2401. DOI: 10.1523/jneurosci.2506-12.2012. URL: http://dx.doi.org/10.1523/JNEUROSCI.2506-12.2012.

[39] Daniel E Feldman. "Timing-Based LTP and LTD at Vertical Inputs to Layer II/III Pyramidal Cells in Rat Barrel Cortex." In: *Neuron* 27.1 (July 2000), pp. 45–56. ISSN: 0896-6273. DOI: 10.1016/s0896-6273(00)00008-8. URL: http://dx.doi.org/10.1016/s0896-6273(00)00008-8.

[40] Jenq-Wei Yang et al. "Development of the whisker-to-barrel cortex system." In: *Current Opinion in Neurobiology* 53 (Dec. 2018), pp. 29–34. ISSN: 0959-4388. DOI: 10.1016/j.conb.2018.04.023. URL: http://dx.doi.org/10.1016/j.conb.2018.04.023.

[41] Yoshiaki Tagawa et al. "Multiple periods of functional ocular dominance plasticity in mouse visual cortex." In: *Nature Neuroscience* 8.3 (Feb. 2005), pp. 380–388. ISSN: 1546-1726. DOI: 10.1038/nn1410. URL: http://dx.doi.org/10.1038/nn1410.

[42] Reha S Erzurumlu and Peter C Kind. "Neural activity: sculptor of 'barrels' in the neocortex." In: *Trends in Neurosciences* 24.10 (Oct. 2001), pp. 589–595. ISSN: 0166-2236. DOI: 10.1016/s0166-2236(00)01958-5. URL: http://dx.doi.org/10.1016/S0166-2236(00)01958-5.

[43] Ben Mitchinson et al. "Active vibrissal sensing in rodents and marsupials." In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 366.1581 (Nov. 2011), pp. 3037–3048. ISSN: 1471-2970. DOI: 10.1098/rstb.2011.0156. URL: http://dx.doi.org/10.1098/rstb.2011.0156.

[44] D. H. Hubel and T. N. Wiesel. "The period of susceptibility to the physiological effects of unilateral eye closure in kittens." In: *The Journal of Physiology* 206.2 (Feb. 1970), pp. 419–436. ISSN: 1469-7793. DOI: 10.1113/jphysiol.1970.sp009022. URL: http://dx.doi.org/10.1113/jphysiol.1970.sp009022.

[45] Niraj S. Desai, Lana C. Rutherford, and Gina G. Turrigiano. "Plasticity in the intrinsic excitability of cortical pyramidal neurons." In: *Nature Neuroscience* 2.6 (June 1999), pp. 515–520. ISSN: 1546-1726. DOI: 10.1038/9165. URL: http://dx.doi.org/10.1038/9165.

[46] Wei Zhang and David J. Linden. "The other side of the engram: experience-driven changes in neuronal intrinsic excitability." In: *Nature Reviews Neuroscience* 4.11 (Nov. 2003), pp. 885–900. ISSN: 1471-0048. DOI: 10.1038/nrn1248. URL: http://dx.doi.org/10.1038/nrn1248.

[47] Gaël Daoudal and Dominique Debanne. "Long-Term Plasticity of Intrinsic Excitability: Learning Rules and Mechanisms." In: *Learning & Memory* 10.6 (Nov. 2003), pp. 456–465. ISSN: 1549-5485. DOI: 10.1101/lm.64103. URL: http://dx.doi.org/10.1101/lm.64103.

[48] Yuan Fan et al. "Activity-dependent decrease of excitability in rat hippocampal neurons through increases in Ih." In: *Nature Neuroscience* 8.11 (Oct. 2005), pp. 1542–1551. ISSN: 1546-1726. DOI: 10.1038/nn1568. URL: http://dx.doi.org/10.1038/nn1568.

[49] Mary E. Lambo and Gina G. Turrigiano. "Synaptic and Intrinsic Homeostatic Mechanisms Cooperate to Increase L2/3 Pyramidal Neuron Excitability during a Late Phase of Critical Period Plasticity." In: *The Journal of Neuroscience* 33.20 (May 2013), pp. 8810–8819. ISSN: 1529-2401. DOI: 10.1523/jneurosci.4502-12.2013. URL: http://dx.doi.org/10.1523/JNEUROSCI.4502-12.2013.

[50] Joachim Lübke et al. "Columnar Organization of Dendrites and Axons of Single and Synaptically Coupled Excitatory Spiny Neurons in Layer 4 of the Rat Barrel Cortex." In: *The Journal of Neuroscience* 20.14 (July 2000), pp. 5300–5311. ISSN: 1529-2401. DOI: 10.1523/jneurosci.20-14-05300.2000. URL: http://dx.doi.org/10.1523/JNEUROSCI.20-14-05300.2000.

[51] Guanxiao Qi and Dirk Feldmeyer. "Cell-Type Specific Neuromodulation of Excitatory and Inhibitory Neurons via Muscarinic Acetylcholine Receptors in Layer 4 of Rat Barrel Cortex." In: *Frontiers in Neural Circuits* 16 (Feb. 2022). ISSN: 1662-5110. DOI: 10.3389/fncir.2022.843025. URL: http://dx.doi.org/10.3389/fncir.2022.843025.

[52] Gina G. Turrigiano et al. "Activity-dependent scaling of quantal amplitude in neocortical neurons." In: *Nature* 391.6670 (Feb. 1998), pp. 892–896. ISSN: 1476-4687. DOI: 10.1038/36103. URL: http://dx.doi.org/10.1038/36103.

[53] Aslı Ayaz et al. "Layer-specific integration of locomotion and sensory information in mouse barrel cortex." In: *Nature Communications* 10.1 (June 2019). ISSN: 2041-1723. DOI: 10.1038/s41467-019-10564-8. URL: http://dx.doi.org/10.1038/s41467-019-10564-8.

[54] Qing Xu et al. "Origins of Cortical Interneuron Subtypes." In: *The Journal of Neuroscience* 24.11 (Mar. 2004), pp. 2612–2622. ISSN: 1529-2401. DOI: 10.1523/jneurosci.5667-03.2004. URL: http://dx.doi.org/10.1523/JNEUROSCI.5667-03.2004.

[55] Simon J.B. Butt et al. "The Requirement of Nkx2-1 in the Temporal Specification of Cortical Interneuron Subtypes." In: *Neuron* 59.5 (Sept. 2008), pp. 722–732. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2008.07.031. URL: http://dx.doi.org/10.1016/j.neuron.2008.07.031.

[56] D A McCormick and D A Prince. "Post-natal development of electrophysiological properties of rat cerebral cortical pyramidal neurones." In: *The Journal of Physiology* 393.1 (Dec. 1987), pp. 743–762. ISSN: 1469-7793. DOI: 10.1113/jphysiol.1987.sp016851. URL: http://dx.doi.org/10.1113/jphysiol.1987.sp016851.

[57] Anne-Marie M. Oswald and Alex D. Reyes. "Maturation of Intrinsic and Synaptic Properties of Layer 2/3 Pyramidal Neurons in Mouse Auditory Cortex." In: *Journal of Neurophysiology* 99.6 (June 2008), pp. 2998–3008. ISSN: 1522-1598. DOI: 10.1152/jn.01160.2007. URL: http://dx.doi.org/10.1152/jn.01160.2007.

[58] S. J. Etherington and S. R. Williams. "Postnatal Development of Intrinsic and Synaptic Properties Transforms Signaling in the Layer 5 Excitatory Neural Network of the Visual Cortex." In: *Journal of Neuroscience* 31.26 (June 2011), pp. 9526–9537. ISSN: 1529-2401. DOI: 10.1523/jneurosci.0458-11.2011. URL: http://dx.doi.org/10.1523/JNEUROSCI.0458-11.2011.

[59] Melanie A. Gainey and Daniel E. Feldman. "Multiple shared mechanisms for homeostatic plasticity in rodent somatosensory and visual cortex." In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 372.1715 (Mar. 2017), p. 20160157. ISSN: 1471-2970. DOI: 10.1098/rstb.2016.0157. URL: http://dx.doi.org/10.1098/rstb.2016.0157.

[60] Stanislaw Glazewski, Stuart Greenhill, and Kevin Fox. "Time-course and mechanisms of homeostatic plasticity in layers 2/3 and 5 of the barrel cortex." In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 372.1715 (Mar. 2017), p. 20160150. ISSN: 1471-2970. DOI: 10.1098/rstb.2016.0150. URL: http://dx.doi.org/10.1098/rstb.2016.0150.

[61] Alejandro Torrado Pacheco et al. "Sleep Promotes Downward Firing Rate Homeostasis." In: *Neuron* 109.3 (Feb. 2021), 530–544.e6. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2020.11.001. URL: http://dx.doi.org/10.1016/j.neuron.2020.11.001.

[62] Melanie A. Gainey et al. "Synaptic Scaling Requires the GluR2 Subunit of the AMPA Receptor." In: *The Journal of Neuroscience* 29.20 (May 2009), pp. 6479–6489. ISSN: 1529-2401. DOI: 10.1523/jneurosci.3753-08.2009. URL: http://dx.doi.org/10.1523/JNEUROSCI.3753-08.2009.

[63] Niraj S. Desai et al. "Critical periods for experience-dependent synaptic scaling in visual cortex." In: *Nature Neuroscience* 5.8 (June 2002), pp. 783–789. ISSN: 1546-1726. DOI: 10.1038/nn878. URL: http://dx.doi.org/10.1038/nn878.

[64] Yildirim Sara et al. "An Isolated Pool of Vesicles Recycles at Rest and Drives Spontaneous Neurotransmission." In: *Neuron* 45.4 (Feb. 2005), pp. 563–573. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2004.12.056. URL: http://dx.doi.org/10.1016/j.neuron.2004.12.056.

[65] Jeremy M. Henley and Kevin A. Wilkinson. "Synaptic AMPA receptor composition in development, plasticity and disease." In: *Nature Reviews Neuroscience* 17.6 (Apr. 2016), pp. 337–350. ISSN: 1471-0048. DOI: 10.1038/nrn.2016.37. URL: http://dx.doi.org/10.1038/nrn.2016.37.

[66] Siqiong June Liu and R. Suzanne Zukin. "Ca2+-permeable AMPA receptors in synaptic plasticity and neuronal death." In: *Trends in Neurosciences* 30.3 (Mar. 2007), pp. 126–134. ISSN: 0166-2236. DOI: 10.1016/j.tins.2007.01.006. URL: http://dx.doi.org/10.1016/j.tins.2007.01.006.

[67] Laurence Cathala et al. "Changes in synaptic structure underlie the developmental speeding of AMPA receptor–mediated EPSCs." In: *Nature Neuroscience* 8.10 (Sept. 2005), pp. 1310–1318. ISSN: 1546-1726. DOI: 10.1038/nn1534. URL: http://dx.doi.org/10.1038/nn1534.

[68] Christian Lohmann and Helmut W. Kessels. "The developmental stages of synaptic plasticity." In: *The Journal of Physiology* 592.1 (Nov. 2013), pp. 13–31. ISSN: 1469-7793. DOI: 10.1113/jphysiol.2012.235119. URL: http://dx.doi.org/10.1113/jphysiol.2012.235119.

[69] Melanie A. Gainey, Joseph W. Aman, and Daniel E. Feldman. "Rapid Disinhibition by Adjustment of PV Intrinsic Excitability during Whisker Map Plasticity in Mouse S1." In: *The Journal of Neuroscience* 38.20 (Apr. 2018), pp. 4749–4761. ISSN: 1529-2401. DOI: 10.1523/jneurosci.3628-17.2018. URL: http://dx.doi.org/10.1523/JNEUROSCI.3628-17.2018.

[70] Bernardo Rudy et al. "Three groups of interneurons account for nearly 100% of neocortical GABAergic neurons." In: *Developmental Neurobiology* 71.1 (Dec. 2010), pp. 45–61. ISSN: 1932-846X. DOI: 10.1002/dneu.20853. URL: http://dx.doi.org/10.1002/dneu.20853.

[71] Yazan N. Billeh et al. "Systematic Integration of Structural and Functional Data into Multi-scale Models of Mouse Primary Visual Cortex." In: *Neuron* 106.3 (May 2020), 388–403.e18. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2020.01.040. URL: http://dx.doi.org/10.1016/j.neuron.2020.01.040.

[72] B Semihcan Sermet et al. "Pathway-, layer- and cell-type-specific thalamic input to mouse barrel cortex." In: *eLife* 8 (Dec. 2019). ISSN: 2050-084X. DOI: 10.7554/elife.52665. URL: http://dx.doi.org/10.7554/eLife.52665.

[73] JoséAntonio del Rio et al. "The development of parvalbumin-immunoreactivity in the neocortex of the mouse." In: *Developmental Brain Research* 81.2 (Sept. 1994), pp. 247–259. ISSN: 0165-3806. DOI: 10.1016/0165-3806(94)90311-5. URL: http://dx.doi.org/10.1016/0165-3806(94)90311-5.

[74] Qing Xu, Melissa Tam, and Stewart A. Anderson. "Fate mapping Nkx2.1-lineage cells in the mouse telencephalon." In: *Journal of Comparative Neurology* 506.1 (Nov. 2007), pp. 16–29. ISSN: 1096-9861. DOI: 10.1002/cne.21529. URL: http://dx.doi.org/10.1002/cne.21529.

[75] Rebecca Ouwenga et al. "Transcriptomic Analysis of Ribosome-Bound mRNA in Cortical Neurites In Vivo." In: *The Journal of Neuroscience* 37.36 (Aug. 2017), pp. 8688–8705. ISSN: 1529-2401. DOI: 10.1523/jneurosci.3044-16.2017. URL: http://dx.doi.org/10.1523/JNEUROSCI.3044-16.2017.

[76] Nathan W. Gouwens et al. "Classification of electrophysiological and morphological neuron types in the mouse visual cortex." In: *Nature Neuroscience* 22.7 (June 2019), pp. 1182–1195. ISSN: 1546-1726. DOI: 10.1038/s41593-019-0417-0. URL: http://dx.doi.org/10.1038/s41593-019-0417-0.

[77] Jennifer L. Hoy and Cristopher M. Niell. "Layer-Specific Refinement of Visual Cortex Function after Eye Opening in the Awake Mouse." In: *The Journal of Neuroscience* 35.8 (Feb. 2015), pp. 3370–3383. ISSN: 1529-2401. DOI: 10.1523/jneurosci.3174-14.2015. URL: http://dx.doi.org/10.1523/JNEUROSCI.3174-14.2015.

[78] George D. Mower. "The effect of dark rearing on the time course of the critical period in cat visual cortex." In: *Developmental Brain Research* 58.2 (Feb. 1991), pp. 151–158. ISSN: 0165-3806. DOI: 10.1016/0165-3806(91)90001-y. URL: http://dx.doi.org/10.1016/0165-3806(91)90001-y.

[79] W.I. Welker. "Analysis of Sniffing of the Albino Rat 1)." In: *Behaviour* 22.3–4 (1964), pp. 223–244. ISSN: 1568-539X. DOI: 10.1163/156853964x00030. URL: http://dx.doi.org/10.1163/156853964X00030.

[80] Ingrid Bureau, Francisca von Saint Paul, and Karel Svoboda. "Interdigitated Paralemniscal and Lemniscal Pathways in the Mouse Barrel Cortex." In: *PLoS Biology* 4.12 (Nov. 2006). Ed. by Ford Ebner, e382. ISSN: 1545-7885. DOI: 10.1371/journal.pbio.0040382. URL: http://dx.doi.org/10.1371/journal.pbio.0040382.

[81] Hanno S Meyer et al. "Number and laminar distribution of neurons in a thalamocortical projection column of rat vibrissal cortex." en. In: *Cereb. Cortex* 20.10 (Oct. 2010), pp. 2277–2286.

[82] Herbert P. Killackey and Gary R. Belford. "The formation of afferent patterns in the somatosensory cortex of the neonatal rat." In: *Journal of Comparative Neurology* 183.2 (Jan. 1979), pp. 285–303. ISSN: 1096-9861. DOI: 10.1002/cne.901830206. URL: http://dx.doi.org/10.1002/cne.901830206.

[83] Daniel J. Simons and Peter W. Land. "Early experience of tactile stimulation influences organization of somatic sensory cortex." In: *Nature* 326.6114 (Apr. 1987), pp. 694–697. ISSN: 1476-4687. DOI: 10.1038/326694a0. URL: http://dx.doi.org/10.1038/326694a0.

[84] Kevin Fox and Rachel O.L. Wong. "A Comparison of Experience-Dependent Plasticity in the Visual and Somatosensory Systems." In: *Neuron* 48.3 (Nov. 2005), pp. 465–477. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2005.10.013. URL: http://dx.doi.org/10.1016/j.neuron.2005.10.013.

[85] Adam M. Packer and Rafael Yuste. "Dense, Unspecific Connectivity of Neocortical Parvalbumin-Positive Interneurons: A Canonical Microcircuit for Inhibition?" In: *The Journal of Neuroscience* 31.37 (Sept. 2011), pp. 13260–13271. ISSN: 1529-2401. DOI: 10.1523/jneurosci.3131-11.2011. URL: http://dx.doi.org/10.1523/JNEUROSCI.3131-11.2011.

[86] Shawn R. Olsen et al. "Gain control by layer six in cortical circuits of vision." In: *Nature* 483.7387 (Feb. 2012), pp. 47–52. ISSN: 1476-4687. DOI: 10.1038/nature10835. URL: http://dx.doi.org/10.1038/nature10835.

[87] Peter Y. Borden et al. "Thalamic bursting and the role of timing and synchrony in thalamocortical signaling in the awake mouse." In: *Neuron* 110.17 (Sept. 2022), 2836–2853.e8. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2022.06.008. URL: http://dx.doi.org/10.1016/j.neuron.2022.06.008.

[88] Carl C. H. Petersen et al. "Interaction of sensory responses with spontaneous depolarization in layer 2/3 barrel cortex." In: *Proceedings of the National Academy of Sciences* 100.23 (Oct. 2003), pp. 13638–13643. ISSN: 1091-6490. DOI: 10.1073/pnas.2235811100. URL: http://dx.doi.org/10.1073/pnas.2235811100.

[89] Dominique Debanne, Yanis Inglebert, and Michaël Russier. "Plasticity of intrinsic neuronal excitability." In: *Current Opinion in Neurobiology* 54 (Feb. 2019), pp. 73–82. ISSN: 0959-4388. DOI: 10.1016/j.conb.2018.09.001. URL: http://dx.doi.org/10.1016/j.conb.2018.09.001.

[90] Qian-Quan Sun. "Experience-Dependent Intrinsic Plasticity in Interneurons of Barrel Cortex Layer IV." In: *Journal of Neurophysiology* 102.5 (Nov. 2009), pp. 2955–2973. ISSN: 1522-1598. DOI: 10.1152/jn.00562.2009. URL: http://dx.doi.org/10.1152/jn.00562.2009.

[91] Stuart David Greenhill, Adam Ranson, and Kevin Fox. "Hebbian and Homeostatic Plasticity Mechanisms in Regular Spiking and Intrinsic Bursting Cells of Cortical Layer 5." In: *Neuron* 88.3 (Nov. 2015), pp. 539–552. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2015.09.025. URL: http://dx.doi.org/10.1016/j.neuron.2015.09.025.

[92] Sandrine Lefort et al. "The Excitatory Neuronal Network of the C2 Barrel Column in Mouse Primary Somatosensory Cortex." In: *Neuron* 61.2 (Jan. 2009), pp. 301–316. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2008.12.020. URL: http://dx.doi.org/10.1016/j.neuron.2008.12.020.

[93] Peter W. Land and Daniel J. Simons. "Cytochrome oxidase staining in the rat smI barrel cortex." In: *Journal of Comparative Neurology* 238.2 (Aug. 1985), pp. 225–235. ISSN: 1096-9861. DOI: 10.1002/cne.902380209. URL: http://dx.doi.org/10.1002/cne.902380209.

[94] Gerald T. Finnerty, Langdon S. E. Roberts, and Barry W. Connors. "Sensory experience modifies the short-term dynamics of neocortical synapses." In: *Nature* 400.6742 (July 1999), pp. 367–371. ISSN: 1476-4687. DOI: 10.1038/22553. URL: http://dx.doi.org/10.1038/22553.

[95] Federico Scala et al. "Layer 4 of mouse neocortex differs in cell types and circuit organization between sensory areas." In: *Nature Communications* 10.1 (Sept. 2019). ISSN: 2041-1723. DOI: 10.1038/s41467-019-12058-z. URL: http://dx.doi.org/10.1038/s41467-019-12058-z.

[96] Shilpi Minocha et al. "Nkx2.1 regulates the generation of telencephalic astrocytes during embryonic development." In: *Scientific Reports* 7.1 (Mar. 2017). ISSN: 2045-2322. DOI: 10.1038/srep43093. URL: http://dx.doi.org/10.1038/srep43093.

[97] Laura A.B. Elias, Gregory B. Potter, and Arnold R. Kriegstein. "A Time and a Place for Nkx2-1 in Interneuron Specification and Migration." In: *Neuron* 59.5 (Sept. 2008), pp. 679–682. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2008.08.017. URL: http://dx.doi.org/10.1016/j.neuron.2008.08.017.