The GnRH pulse generator activity in mouse models of polycystic ovary syndrome

Ziyue Zhou[1], Su Young Han[1], Maria Pardo-Navarro[1], Ellen G. Wall[1], Reena Desai[2], Szilvia Vas[1], David J. Handelsman[2], Allan E. Herbison[1]

[1]Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge CB2 3EG, United Kingdom, [2]ANZAC Research Institute, University of Sydney, Sydney, Australia.

Correspondence should be addressed to Allan E. Herbison (email: aeh36@cam.ac.uk)

Disclosures: The authors have nothing to disclose and no competing interests.

Abstract

One in five women worldwide suffer from polycystic ovary syndrome (PCOS) that, alongside subfertility and hyperandrogenism, typically presents with increased luteinizing hormone (LH) pulsatility. As such, it is suspected that the arcuate kisspeptin (ARN^{KISS}) neurons that represent the GnRH pulse generator are dysfunctional in PCOS. We used here *in vivo* GCaMP fiber photometry and other approaches to examine the behavior of the GnRH pulse generator in two mouse models of PCOS. We began with the peripubertal androgen (PPA) mouse model of PCOS but found that it had a reduction in the frequency of ARN^{KISS} neuron synchronization events (SEs) that drive LH pulses. Examining the prenatal androgen (PNA) model of PCOS, we observed highly variable patterns of pulse generator activity with no significant differences detected in ARN^{KISS} neuron SEs, pulsatile LH secretion, or serum testosterone, estradiol, and progesterone concentrations. However, an unsupervised machine learning approach identified that the ARN^{KISS} neurons of acyclic PNA mice continued to exhibit cyclical patterns of activity similar to that of normal mice. The frequency of ARN^{KISS} neuron SEs was significantly increased in algorithm-identified "diestrous stage" PNA mice compared to controls. In addition, ARN^{KISS} neurons exhibited reduced feedback suppression to progesterone in PNA mice and their pituitary gonadotrophs were also less sensitive to GnRH. These observations demonstrate the importance of understanding GnRH pulse generator activity in mouse models of PCOS. The unexpected functional disassociation of cyclical GnRH pulse generator activity in the acyclic PNA mouse indicates

the presence of a complex phenotype with deficits at multiple levels of the hypothalamo-
pituitary-gonadal axis.

Introduction

Polycystic ovary syndrome (PCOS) is one of the most common causes of female infertility
with variable reproductive and metabolic phenotypes (Bozdag, Mumusoglu et al. 2016,
Lizneva, Suturina et al. 2016). Many women with PCOS have an elevated body mass index
but up to 30% of patients do not, resulting in PCOS being further classified into obese and
lean subtypes (Nestler and Jakubowicz 1997, Kumari, Haq et al. 2005, Toosy, Sodi et al.
2018). Although androgen excess, oligo-menorrhea/anovulation, and the existence of
polycystic ovaries are the key diagnostic indicators for PCOS, a common secondary feature
is increased luteinizing hormone (LH) pulsatility (Waldstreicher, Santoro et al. 1988, The
Rotterdam ESHRE/ASRM-sponsored PCOS consensus workshop group 2004). Basal and
pulsatile LH secretion can be up to threefold higher in both lean and obese PCOS women
and is considered to be a significant contributor to their subfertility (Morales, Laughlin et al.
1996, Dam, Roelfsema et al. 2002, Phylactou, Clarke et al. 2021). A presumed increase in
the frequency of pulsatile gonadotropin-releasing hormone (GnRH) secretion in PCOS
women results in an increased LH/follicle-stimulating hormone (FSH) ratio (Wildt, Häusler et
al. 1981, Dalkin, Haisenleder et al. 1989). This is thought to drive excess ovarian androgen
production that, in turn, impairs estrogen and progesterone feedback regulation of GnRH
secretion (Pastor, Griffin-Korf et al. 1998, Chhabra, McCartney et al. 2005, Moore, Prescott
et al. 2013, Moore, Prescott et al. 2015).

It has become clear that the arcuate nucleus kisspeptin (ARN^{KISS}) neurons represent the
mammalian GnRH pulse generator (Clarkson, Han et al. 2017, Herbison 2018, Goodman,
Herbison et al. 2022). These neurons exhibit brief episodes of synchronized activity that
release kisspeptin on the distal projections of GnRH neurons to drive pulsatile GnRH
secretion into the portal vasculature (Herbison 2021, Liu, Yeo et al. 2021). Recent studies
indicate that the synchronized activity amongst ARN^{KISS} neurons required for pulsatile
hormone secretion is generated by local glutamatergic transmission that is subsequently
modulated by neurokinin B and dynorphin (Qiu, Nestor et al. 2016, Han, Morris et al. 2023,
Morris and Herbison 2023). While the ARN^{KISS} neurons are the primary site of estrogen
negative feedback suppressing pulsatile GnRH secretion (McQuillan, Clarkson et al. 2022),
the mechanism of progesterone feedback remains unclear but could also involve the
ARN^{KISS} neurons (Goodman, Holaskova et al. 2011). As such, it is thought that dysfunction
within the ARN^{KISS} neuron pulse generator may be responsible for the elevated LH pulse

frequency observed in PCOS (Walters, Gilchrist et al. 2018, Coutinho and Kauffman 2019,

Rodriguez Paris and Bertoldo 2019, Ruddenklau and Campbell 2019).

We have developed GCaMP fiber photometry methodologies that allow the activity of the

ARN^{KISS} neuron pulse generator to be monitored at high temporal resolution in freely

behaving mice (Han, Kane et al. 2019, McQuillan, Han et al. 2019, Vas, Wall et al. 2024).

Using this and other approaches, we report here a detailed investigation into the pulse

generator activity in two widely used mouse models of PCOS including the peripubertal

androgen (PPA) model of "obese PCOS", and prenatal androgen (PNA) model of "lean

PCOS" (Stener-Victorin, Padmanabhan et al. 2020).

Results

Peripubertal androgen (PPA) mice

1. Peripubertal dihydrotestosterone (DHT)-treated animals exhibit significantly

increased body weight with acyclic estrous cycles

Female 129S6Sv/Ev C57BL/6 *Kiss1$^{Cre/+}$*,GCaMP6s mice exposed to DHT from 3 weeks of

age (PPA: n=4) started to show a significant increase in weight 4 weeks after capsule

implantation compared to controls with empty capsules (control: n=5) (Sidak's multiple

comparisons test; Fig.1-Suppl. Fig.1A). The PPA animals were completely acyclic with

vaginal cytology only ever showing the diestrous stage (Fig.1-Suppl. Fig.1B-D).

2. Peripubertal exposure to DHT results in a slowed GnRH pulse generator

Abrupt and large increases in arcuate nucleus kisspeptin (ARN^{KISS}) neurons activity, termed

synchronization events (SEs), are observed with GCaMP fiber photometry in mice (Clarkson,

Han et al. 2017, Han, Kane et al. 2019, McQuillan, Han et al. 2019). Both adult PPA and

control mice recorded in diestrus for 24 hours exhibited ARN^{KISS} neuron SEs (Fig.1A,B). The

frequency of SEs in PPA mice (n=4) was approximately half that of controls (n=5) with SE

frequency/hour showing an inhibitory trend (P = 0.0714, Mann-Whitney U test; Fig.1C) and

SE interval significantly increased (P = 0.0159, Mann-Whitney U test; Fig.1D). The SE

interval frequency distribution pattern revealed a shift to the right for PPA mice with a tight

clustering within the 70-90-minute bin centers (Fig.1E). To compare the dynamics of SEs

from control and PPA animals, continuous (10 Hz) photometry recordings were performed in

diestrus to obtain high temporal resolution profiles (Fig.1F). These profiles were the same between control (n=5) and PPA (n=4) animals with SE duration, and both upswing and downswing durations calculated from the full-width half maximum (FWHM) points being not significantly different (Fig.1G-H).

Prenatal androgen (PNA) mice

1. The PNA mouse model exhibits highly disordered estrous cyclicity

Female mice originating from dams treated with DHT on gestational days 16, 17, and 18 were found to have a significantly increased anogenital distance (n=10; P < 0.001, Mann-Whitney U test) compared with controls (n=9; Fig.2A). Body weight was not different between PNA and control animals at 10 weeks of age (n=12 per group; P = 0.6602, Mann-Whitney U test; Fig.2B). The PNA mice exhibited very disordered estrous cycles remaining mostly in persistent diestrus (71 ± 3%) with occasional excursions to metestrus. The proestrus stage was never encountered and estrus was observed rarely (Fig.2C-E).

2. Circulating gonadal steroid hormone levels are not different between control and PNA animals.

As androgen excess is one of the hallmarks of PCOS, we initially measured serum testosterone levels in control and PNA animals using a commercial testosterone enzyme-linked immunosorbent assay (ELISA) (Risal, Manti et al. 2021). This did not detect any difference in serum testosterone levels between control (n=13) and PNA (n=20) animals in diestrus (P = 0.8203, Mann-Whitney U test; Fig.2F). Given concerns over the sensitivity and specificity of the steroid hormone ELISAs (Handelsman, Jimenez et al. 2015), we progressed to using ultrasensitive liquid chromatography-mass spectrometry (LC-MS) (Handelsman, Gibson et al. 2020) to measure the circulating levels of testosterone, androstenedione, estradiol, and progesterone in diestrous PNA and control mice (n=6 per group). No significant differences (Mann-Whitney U tests) were detected for androstenedione, estradiol, or progesterone while all samples were below the limit of detection for testosterone (< 0.01 ng/mL) (Fig.2G-I).

3. Kisspeptin, neurokinin B, and dynorphin immunoreactivity are relatively unchanged within the ARN of PNA mice

Previous studies have found variable effects of PNA treatment on kisspeptin, neurokinin B
(NKB), and dynorphin protein and mRNA expression within the ARN (Barnes, Rosenfield et
al. , Yan, Yuan et al. 2014, Osuka, Iwase et al. 2016, Gibson, Jaime et al. 2021, McCarthy,
Dischino et al. 2021, Moore, Lohr et al. 2021). To re-examine this issue, we assessed levels
of immunofluorescence for all three peptides in the ARN of control (n=4-8) and PNA (n=4-6)
mice (Fig.2-Suppl. Fig.1A-C). This revealed a significant but modest elevation in kisspeptin
immunofluorescence within the rostral ARN (P = 0.0187, Sidak's multiple comparisons test)
in PNA mice but not elsewhere in the ARN (Fig.2-Suppl. Fig.1A). No significant changes
were detected for NKB or dynorphin immunoreactivity in any region of the ARN (Fig.2-Suppl.
Fig.1B,C).

4. PNA animals exhibit highly variable patterns of ARN^{KISS} neuron SEs

Both control and PNA mice exhibited robust SEs across 24-hour recording periods (Fig.3A-
D). Control mice recorded in diestrus had approximately 0.8 SEs per hour (Fig.3E) with an
SE interval of 74.75 ± 5.71 min (Fig.3F) and a normal-like distribution of SE intervals
(Fig.3G). Unexpectedly, diestrous PNA mice exhibited highly variable patterns of ARN^{KISS}
neuron activity that could appear similar to diestrous controls (Fig.3C) or very different
(Fig.3D) with episodes of multi-peak SEs (mpSEs). However, overall SE frequency and
intervals were not different between PNA (n=6) and control (n=7) animals (Fig.3E,F; Mann-
Whitney U tests). The frequency distribution of inter-SE intervals in PNA mice exhibited a
peak interval cluster around the 10-minute bin center (39%) with a gradual tail off in SE
intervals (Fig.3G). Continuous (10 Hz) photometry recordings in diestrus did not reveal any
difference in the profiles of single SEs between controls (n=7) and PNA (n=6) animals
(Fig.3H-J).

5. PNA mice have elevated mean LH secretion but normal LH pulse frequency

We have previously found a near perfect relationship between ARN^{KISS} neuron SEs and
pulsatile LH secretion in male and female mice (Han, Kane et al. 2019, McQuillan, Han et al.
2019). To assess this relationship in PNA mice, ARN^{KISS} neuron recordings were
commenced, and tail-tip blood samples taken at 12-minute intervals until an SE was
detected after which the blood sampling rate was switched to 6-minute intervals. We
observed a perfect correlation between SEs and pulsatile LH secretions in both control and
PNA mice: every SE was followed by an LH pulse, and no LH pulses were detected without
a preceding SE (Fig.4A,C). Temporal analysis of SEs and LH release indicated that the SE
peak precedes each LH peak by 7.7 ± 0.6 minutes and 6.6 ± 0.4 minutes in control (n=7

from 4 mice) and PNA (n=7 from 5 mice) animals (P = 0.190, Mann-Whitney U test),
respectively (Fig.4B,D).

The absence of significant differences in ARN^{KISS} neuron SE frequency in PNA mice and
their tight correlation with pulsatile LH secretion indicated that pulsatile LH secretion may not
be different between PNA and control mice. Prior studies have found conflicting data with
one finding no difference in LH pulse frequency (McCarthy, Dischino et al. 2021) and
another reporting increased LH pulsatility (Moore, Prescott et al. 2015). To re-address this
we undertook 4-hour tail-tip blood sampling at 10-minute intervals to assess pulsatile LH
secretion in wild-type C57BL6/J diestrous-stage PNA (n=8) and control (n=7) mice. Mean LH
levels were found to be significantly elevated in PNA mice (P = 0.0205, Mann-Whitney U test;
Fig.4E-G) but the variable LH pulse frequency and amplitude were unchanged compared to
controls (Fig.4H,I).

To evaluate whether the response of pituitary gonadotrophs to GnRH was altered, PNA and
control mice were given GnRH (200 ng/kg, i.p.) at the end of the 4-hour tail-tip bleeding, and
two further tail-tip blood samples collected for LH measurement at 10-minute intervals. Both
control (n=7) and PNA (n=11) mice showed elevated LH levels in response to exogenous
GnRH (Fig.4J). However, PNA mice showed a small but significantly lower level of LH 20
minutes after GnRH injection (P=0.0152, Sidak's multiple comparisons test). The total
GnRH-evoked increment in LH release was significantly lower in PNA mice compared to
controls (P = 0.0028, Mann-Whitney U test, Fig.4K).

6. The pulse generator exhibits cyclical activity in PNA mice

The patterns of ARN^{KISS} neuron SEs recorded from acyclic "diestrous" PNA mice were highly
variable and could exhibit typical diestrous-like activity but also characteristics such as
mpSEs (Fig.3D) that are only usually observed during estrus and metestrus (Vas, Wall et al.
2024). This suggested that, despite the acyclical state of the PNA mice, the pulse generator
may still be undergoing cycles of activity. To assess this, we conducted long-duration
photometry recordings in both PNA and control mice for four consecutive days starting on
proestrus for controls and any day for the acyclic PNA mice. We have recently found that k-
means clustering, an unsupervised machine learning-based approach, provides a very
sensitive characterization of ARN^{KISS} neuron activity across the estrous cycle (Vas, Wall et al.
2024). This involves analysis of SEs based on five parameters: the frequency of regular
single-peak SEs; the standard deviation of inter-SE intervals for regular SEs; and the
number, duration, and profiles of mpSEs (Fig.5A).

All control animals showed typical cycling patterns of ARN^{KISS} neuron activity over the four-
day recording period (n=5, two representative examples shown in Fig.5C). The most
pronounced features of this cyclicity are the suppression of all SE activity during late
proestrus to early estrus (represented by Cluster_0 activity) with a gradual return in SEs
during the late estrus (Cluster_1). This is often followed by the appearance of mpSEs
(Cluster_3/4) during the estrus to metestrus transition. The activity patterns then become the
typical diestrous single SEs with variable frequencies (Cluster_1 & _2) (Fig.5C). Surprisingly,
we observed that the ARN^{KISS} neurons in PNA mice also exhibited variable but shifting SE
profiles over the four-day recordings despite vaginal smears remaining in metestrus or
diestrus throughout (n=4, two examples shown in Fig.5D). These PNA mice exhibited a
cyclical pattern of activity similar to that of control mice with a period of slow Cluster_0
activity directly followed by an mpSEs period (Cluster_3/4) and then regular diestrous activity
(Cluster_1 & _2) (Fig.5D).

Overall, quantification of the time spent in each cluster pattern over the four days was
remarkably similar between control (n=5) and PNA (n=4) mice (Fig.5B). The proportion of
Cluster_0 (PNA 21%; control 21%) and Cluster_1 (PNA 44%; control 42%) activity was
almost identical with the major trend being a nonsignificant increased time ($P > 0.05$, Sidak's
multiple comparisons test) spent in mpSE Cluster_3/4 activity (PNA 20%; control 12%) at the
expense of the fast regular SE Cluster_2 pattern (PNA 15%; control 25%) (Fig.5B).

7. PNA mice have increased diestrous-like GnRH pulse generator frequency

Given that our earlier recordings from PNA mice (Fig.3C,D) were unavoidably taken from
random times of the pulse generator cycle, we used the unsupervised machine learning
algorithm above to identify "diestrous-like" (Cluster_1&_2) episodes of ARN^{KISS} neuron SE
activity in PNA mice and compared this with diestrous controls. In this case, the frequency of
SEs was 48% greater in PNA mice (n=6) compared to control animals (n=7; $P = 0.0029$,
Mann-Whitney U test; Fig.6C) as also reflected in a significantly lower inter-SE interval ($P =$
0.0047, Mann-Whitney U test; Fig.6D). The frequency distribution of inter-SE intervals in
PNA mice was slightly shifted to the left compared to control mice so that majority of PNA
SE intervals were in 10-70 minute bin compared to the 30-90 minute bin centers in control
mice (Fig.6E).

8. PNA mice have impaired progesterone negative feedback

Progesterone negative feedback is thought to be abnormal in women with PCOS (Pastor,
Griffin-Korf et al. 1998, Chhabra, McCartney et al. 2005) and PNA mice (Moore, Prescott et
al. 2015). We assessed the response of ARN^{KISS} neuron activity to vehicle or progesterone
(4 mg/kg, i.p.) in PNA and control mice with diestrous activity patterns. As expected
(McQuillan, Han et al. 2019), we found a marked suppression of ARN^{KISS} neuron SEs that
lasted for up to 6 hours following progesterone in control mice (n=3, $P < 0.001$, Sidak's
multiple comparisons test, Fig.7A,B,E). The response in PNA mice was much less consistent,
and overall, no significant differences in SE activity detected at any time point (n=3,
Fig.7C,D,F).

9. No differences exist in ARN^{KISS} neuron activity in ovariectomized (OVX) PNA and
control females

Gonadal steroids play a key role in modulating the activity of the ARN^{KISS} neurons
(McQuillan, Han et al. 2019, McQuillan, Clarkson et al. 2022). To assess whether any
differences exist in ARN^{KISS} neuron activity in PNA mice in the absence of gonadal steroids,
we undertook 24-hour fiber photometry recordings in mice that had been ovariectomized for
at least 3 weeks. Removal of gonadal steroids results in increased SE frequency and the
emergence of clusters of tightly coupled individual synchronizations as previously described
(McQuillan, Clarkson et al. 2022) (Fig.8A,B). We observed no significant differences (Mann-
Whitney U tests) between SE frequency (Fig. 8C) or intervals (Fig. 8D,E) in PNA and control
animals (n=4 per group). Similarly, high-resolution recordings at 10 Hz did not reveal any
differences in SE profiles between OVX control and OVX PNA mice (Fig.8F,H).

Discussion

It is important to establish animal models that faithfully recapitulate the symptoms and
features of women with PCOS to be able to make progress in treating the disorder (Stener-
Victorin, Padmanabhan et al. 2020). Increased LH pulse frequency is observed in ~75% of
women with PCOS and is considered to be a significant contributor to their sub-fertility
(Taylor, McCourt et al. 1997). We provide here a detailed assessment of the GnRH pulse
generator in two commonly used mouse models of PCOS. The obese PPA mouse model
was found to have a pulse generator that actually operates at a slower frequency than
normal. In contrast, the PNA mouse model has a pulse generator that is operating faster
than normal but, remarkably, maintains cyclical-like patterns of activity despite being in
reproductively acyclic mice.

Pulse generator activity in PPA mice

The chronic administration of DHT to female mice from postnatal day 21 generates an
androgen receptor-dependent mouse model of PCOS with infertility and metabolic
disturbances (Caldwell, Middleton et al. 2014, Caldwell, Edwards et al. 2017, Stener-Victorin,
Padmanabhan et al. 2020, Kerbus, Decourt et al. 2024). In agreement with those prior
studies, we found that PPA mice start to exhibit increased body weight around four weeks
after DHT exposure and are acyclic. Fiber photometry revealed that the frequency of
ARN[KISS] neuron SEs was reduced by ~50% in PPA mice compared to controls. This clearly
indicates that the PPA mouse does not provide an appropriate PCOS model of elevated
pulse generator activity. Given this, we did not proceed to examine pulsatile LH secretion in
PPA mice but note that a recent study has reported pulsatile LH secretion to be unchanged
in this animal model (Coyle, Prescott et al. 2022). It is very likely that chronic excess
androgen activation in this model provides additional negative feedback suppression of
ARN[KISS] neurons. The treatment of adult OVX female mice with the same DHT regimen is
reported to decrease LH pulse frequency and amplitude (Esparza, Terasaka et al. 2020).

Variable reproductive parameters of the PNA mouse model

The PNA mouse model has been studied by many different research groups with some
features being identified in a highly consistent manner while others are not (Stener-Victorin,
Padmanabhan et al. 2020). Common characteristics, also present in the current study, are
those of increased anogenital distance, normal body weight, and highly disrupted estrous
cycles (Sullivan and Moenter 2004, Silva, Prescott et al. 2018, Stener-Victorin,
Padmanabhan et al. 2020, Gibson, Jaime et al. 2021, McCarthy, Dischino et al. 2021). In
contrast, reported differences in circulating gonadal steroid concentrations in PNA mice have
been inconsistent among studies. Using ELISAs, some studies have demonstrated PNA
mice to have elevated testosterone levels (Sullivan and Moenter 2004, Moore, Prescott et al.
2015, Silva, Prescott et al. 2018) while others (Roland and Moenter 2011), including
ourselves now, find no differences. Concerns over the specificity and selectivity of ELISAs
for measuring gonadal steroids in mice (Handelsman and Wartofsky 2013, Auchus 2014,
Wierman, Auchus et al. 2014, Handelsman, Jimenez et al. 2015) led us to re-assess
gonadal steroid hormone levels using gold-standard ultrasensitive LC-MS (Handelsman,
Gibson et al. 2020). We previously observed that testosterone levels were usually below the
limit of detection in C57/BL6 female mice (Wall, Desai et al. 2023) but considered that any
elevated concentrations in PNA mice might become detectable. Unfortunately, this was not

the case, and it remains unclear if testosterone levels are elevated in PNA mice. We were,

however, able to detect estradiol and progesterone concentrations and find that neither is

different between control and PNA mice. A prior ELISA assessment also reported that

estradiol and progesterone levels were unchanged in PNA animals (Moore, Prescott et al.

2015).

Another area of considerable inconsistency among studies involves the effects of PNA

treatment on both mRNA and protein expression of kisspeptin, neurokinin B, and dynorphin

in the ARN (Yan, Yuan et al. 2014, Osuka, Iwase et al. 2016, Gibson, Jaime et al. 2021,

McCarthy, Dischino et al. 2021, Moore, Lohr et al. 2021). We re-investigated this question by

quantifying the immunoreactive levels of these three proteins and observed an increase in

only kisspeptin expression and only in rostral ARN of PNA animals. While there are

important roles for these neuropeptides in modulating ARN[KISS] neuron synchronizations

(Han, Morris et al. 2023, Morris and Herbison 2023) and their subsequent activation of

GnRH secretion (Liu, Yeo et al. 2021), these generally inconsistent findings indicate that any

PNA-induced changes in the expression of the neuropeptides are subtle. Indeed, the

principal determinant of kisspeptin, neurokinin B, and dynorphin expression in ARN[KISS]

neurons is estradiol (Navarro, Gottsch et al. 2009) and we demonstrate here that circulating

estradiol levels are not altered in PNA mice.

Pulse generator activity and pulsatile LH in the PNA mouse model

One of the key features of the PNA mouse model is thought to be that of increased pulsatile

LH secretion (Stener-Victorin, Padmanabhan et al. 2020). However, this has only been

reported in a single study with PNA mice exhibiting an approximately 40% increase in LH

pulse frequency (Moore, Prescott et al. 2015). A recent paper undertaking the same tail-tip

pulse bleeding approach found that LH pulse frequency was not significantly altered in PNA

mice (McCarthy, Dischino et al. 2021). Our present observations may explain these

discrepancies. As the pulse generator continues to cycle in the acyclic PNA mouse, it is

essentially impossible to define the "pulse generator stage" of a PNA mouse at the time of

bleeding and consequently wide variations in LH pulse frequency would be expected. We

find here that LH pulse frequency recorded from such random "pulse generator" cycling wild-

type PNA mice is not significantly different to that of controls. Nevertheless, with the benefit

of being able to measure ARN[KISS] neuron activity directly, it has been possible to show that

"diestrous-stage" PNA mice do have an increased rate of pulse generator frequency

compared with controls. As such, the PNA mouse model does to some degree replicate the

increased pulse frequency seen in many women with PCOS.

The most striking feature of pulse generator activity in PNA mice is its ability to retain cyclical
patterns of activity within a reproductively acyclic mouse. The pulse generator of PNA mice
exhibits clear transitions from periods of "estrous-like" Cluster_0 activity through to mpSE-
dominated Cluster_3/4 phase and onto regular "diestrous-like" Cluster_1&_2 patterns. The
percentage of time PNA mice spend in each pattern of activity was not found to be
significantly different to controls. We also note that the dynamics of individual ARN[KISS]
neuron SEs remain unchanged under all conditions in PNA mice, suggesting that the
fundamental synchronization mechanism and numbers of contributing neurons is not altered
(Han, Morris et al. 2023). This is consistent with the observation that PNA treatment does
not alter ARN[KISS] neuron firing activity in adult acute brain slices (Gibson, Jaime et al. 2021).
Together, these observations indicate that the acyclic reproductive phenotype of PNA mice
does not result from an absence of cyclical activity in the pulse generator. Equally, a
significant impact of PNA treatment on the activity of the GnRH neuron can be excluded
(Jaime and Moenter 2022) and, at least in mice, pulse generation occurs autonomously at
the GnRH neuron distal dendron with no role for GnRH neuron cell bodies (Wang, Guo et al.
2020, Herbison 2021). The degree to which cyclical GnRH pulse generator activity in PNA
mice reflects women with PCOS is unknown. However, a very wide range of LH pulse
frequency is found in anovulatory women with PCOS (Taylor, McCourt et al. 1997),
suggesting the intriguing possibility that the GnRH pulse generator activity may also continue
to cycle in women with PCOS.

Gonadal steroid feedback in PNA mice

Pulse generator activity was found to be identical in ovariectomized PNA and control mice.
This indicates that, despite androgen receptors being expressed by ARN[KISS] neurons in the
perinatal period (Watanabe, Fisher et al. 2023), excess DHT exposure at this time does not
engender any fundamental differences in their synchronization behavior in the absence of
gonads as adults. This suggests that the subtle increase in pulse generator frequency
detected here in "diestrous-like" PNA mice arises from the effects of circulating gonadal
steroids. As gonadal steroid levels are normal in PNA mice, it is likely that perinatal DHT
exposure programs long-term changes in gonadal steroid receptor function in adult PNA
mice. While direct estrogen feedback is critical in maintaining the suppressed activity of the
pulse generator throughout the estrous cycle (McQuillan, Clarkson et al. 2022), ARN[KISS]
neuron ESR1 expression was not found to be altered in PNA mice (Moore, Lohr et al. 2021).

Reduced progesterone negative feedback is widely considered to be a major factor
underlying enhanced LH pulsatility in women with PCOS (Pastor, Griffin-Korf et al. 1998,
Chhabra, McCartney et al. 2005) and the ability of progesterone to suppress LH secretion is
reduced in OVX PNA mice (Moore, Prescott et al. 2015). Whereas progesterone exerts a
robust and prolonged suppression of ARN^{KISS} neuron SEs in normal (McQuillan, Han et al.
2019) and control mice, we find this to be much less effective in PNA females. Precisely how
perinatal DHT treatment desensitizes ARN^{KISS} neurons to progesterone is unknown and
could occur directly or indirectly (Moore, Lohr et al. 2021). Reported effects of PNA on
progesterone receptors in the ARN are inconsistent with studies reporting either reduced or
unchanged expression (Moore, Prescott et al. 2015, Gibson, Jaime et al. 2021, Moore, Lohr
et al. 2021), although one recent study found a 30% reduction in progesterone receptor
mRNA expression by ARN^{KISS} neuron in PNA mice (Moore, Lohr et al. 2021).
417
Although we find that the PNA mouse has reduced progesterone feedback at the level of the
pulse generator, as is suspected for women with PCOS, it is unclear how this might pattern
ARN^{KISS} neuron activity. Control and PNA mice exhibit similar proportions of time with halted
or slow pulse generator activity (Cluster_0). This pattern of activity is thought to occur in
response to increased progesterone level following the LH surge (McQuillan, Han et al. 2019,
Vas, Wall et al. 2024). Thus, it is somewhat perplexing to find clear Cluster_0 activity in PNA
mice that have reduced sensitivity to progesterone negative feedback. This might suggest
that in other mechanisms can become responsible for luteal phase slowing of pulse
generator activity in PNA mice. Equally, is unclear how reduced progesterone feedback
could elevate the "diestrous-like" pattern of ARN^{KISS} neuron activity, although roles for
progesterone outside the luteal phase have been suggested (Leipheimer, Bona-Gallo et al.
1984).
430
Previous studies have demonstrated that the estrogen-activated LH surge mechanism is
normal in PNA mice (Moore, Prescott et al. 2013) and, despite proestrous smears never
being observed in PNA mice, copra lutea can be found suggesting that LH surges and
ovulation can occur (Moore, Prescott et al. 2013, Lei, Ding et al. 2017, McCarthy, Dischino et
al. 2021). This raises the intriguing possibility that the GnRH surge generator may also be
operational in PNA mice but masked by abnormal downstream signalling. In that regard, we
note that pituitary gonadotroph sensitivity to GnRH is blunted in PNA mice as previously
reported (Silva, Desroziers et al. 2019). Given that pulse generator activity is the same in
ovariectomized PNA and control mice, reduced gonadotroph sensitivity would explain the
reduced rise in LH following ovariectomy in PNA mice observed in other studies (Moore,
Prescott et al. 2013, Moore, Prescott et al. 2015). Nevertheless, we note that the PNA

mouse clearly does not model women with PCOS that show a much-elevated LH response
to GnRH (Barnes, Rosenfield et al. 1989, Batrinos 1993, Morales, Laughlin et al. 1996, Patel,
Coffler et al. 2003)
445
In summary, we report here that the GnRH pulse generator operates at a lower frequency in
PPA mice, but at a higher frequency when "diestrous-like" pulse generator activity was
analyzed in PNA mice. From the perspective of understanding enhanced pulsatile LH
secretion in women with PCOS, the PNA model is clearly the most appropriate model.
However, we note that many features of PNA mice such as their pulse generator free-
running behavior in the ovariectomized state and gonadal steroid hormone concentrations
are normal. Remarkably, we find evidence that the pulse generator continues to exhibit
estrous cycle-like patterns of activity within an acyclic PNA mouse model. Thus, the
anovulatory phenotype of PNA mice arises in a complex manner from deficits likely at many
levels of the hypothalamo-pituitary-ovarian axis. The degree to which the PNA mouse model
reflects women with PCOS remains an important, albeit very difficult, question to address.
457
458
459 **Methods**
460
461 ***Mice***
462
Female 129S6Sv/Ev C57BL/6 *Kiss1*[Cre/+]mice (Yeo, Kyle et al. 2016) were crossed with the
Ai162 (TIT2L-GC6s-ICL-tTA2)-D Cre-dependent GCaMP6s males (JAX stock #031562)
(Daigle, Madisen et al. 2018) to generate *Kiss1*-Cre/+,*Ai126D* mice as previously
characterized (Han, Morris et al. 2023). Mice were group-housed in conventional cages with
environmental enrichment under conditions of controlled temperature ($22 \pm 2\ ^0C$) and lighting
(12-hour light/12-hour dark cycle; lights on at 05:30 or 07:00) with *ad libitum* access to food
(RM3 (E), Special Diets Services, UK) and water. All animal experimental protocols were
approved by the University of Cambridge, UK (P174441DE). Where required, mice were
bilaterally ovariectomized (OVX) under isoflurane anesthesia at least 3 weeks prior to
experimentation.
473
474 ***Peripubertal androgen (PPA) model***
475
Peripubertal female mice were exposed to DHT as previously described (van Houten,
Kramer et al. 2012, Caldwell, Edwards et al. 2017). Briefly, peripubertal female mice (21-23
days old) were implanted subcutaneously (s.c.) with either an empty (control) or

dihydrotestosterone (DHT)-filled (PPA) 1-cm silastic implants (id, 1.47 mm; od, 1.95 mm,
Dow Corning Corp, catalog no. 508–006). These silastic implants containing ~10 mg DHT
provide steady-state DHT release for at least 6 months (Singh, O'Neill et al. 1995). The body
weight of these animals was monitored weekly until optic fiber implantation at the 10 weeks
of age.

Prenatal androgen (PNA) model

Adult females were paired with males and checked for copulatory plugs, indicating day 1 of
gestation. Pregnant dams were given s.c. injections of sesame oil vehicle (100 µL) alone or
containing 250 µg of DHT on gestational days 16, 17, and 18 (Sullivan and Moenter 2004).
Female offspring from androgen-treated (PNA) and vehicle-treated (control) dams were used
for experiments.

Estrous cycle determination

The estrous cycle stage was determined by the relative ratio of leukocytes, cornified, and
nucleated cells observed in the vaginal epithelial cell smear samples on the morning of
collection (Ajayi and Akhigbe 2020). The smears were collected using 5 µL of sterile
phosphate-buffered saline (PBS) which was transferred to a glass slide to air dry before
staining with filtered Giemsa (1:1 in MilliQ water) and examined under a light microscope.

Testosterone ELISA measurements and analysis

Blood samples for testosterone measurement were collected in the afternoon of diestrus
through a tail-tip blood sample prior to ovariectomy or from the inferior vena cava before
perfusion. The samples were then spun at room temperature at 3,000 rpm for 10 minutes.
Serum was collected and kept at –20°C until the day of hormone measurement. Serum
testosterone levels were measured using the testosterone rat/mouse ELISA kit according to
the manufacturer's instructions (Demeditec Diagnostics, GmbH, DEV9911, Kiel, Bermany).
The assay sensitivity for mouse testosterone ELISA was 0.066 ng/mL, and intra-assay
coefficient of variation was 6.5%.

Liquid Chromatography–Mass Spectrometry (LC-MS)

Terminal blood samples were collected from the inferior vena cava of anaesthetized control
and PNA animals at 10:00 on diestrus exactly as described previously (Wall, Desai et al.

2023). Plasma levels of estradiol, progesterone, testosterone, and androstenedione were measured using liquid chromatography-mass spectrometry (LC-MS) (Handelsman, Gibson et al. 2020).

Quantification of kisspeptin/neurokinin B/dynorphin expression in ARN

Adult diestrous PNA and control female mice were given a lethal dose of pentobarbital (3 mg/100µL, intraperitoneal, i.p.) and perfused transcardially with 4% paraformaldehyde (PFA). Brains were post-fixed in 4% PFA for 1 hour at 4°C and transferred to 30% sucrose at 4°C until sunk. Forty-µm-thick sections were collected with microtomes and processed for kisspeptin and neurokinin B (NKB) immunofluorescence and dynorphin (DYN) Nickel-3,3′-Diaminobenzidine (DAB) staining. For kisspeptin staining, brain sections were incubated with rabbit anti-kisspeptin (1:5,000, AC566; a gift from Dr. Alain Caraty) followed by goat anti-rabbit Alexa Fluor 568 (1:500, Molecular Probes). For NKB staining, brain sections were incubated with guinea pig anti-NKB (1:5,000, IS-3/61, a gift from Dr. Philippe Ciofi) followed by biotinylated goat anti-guinea pig IgG (1:1,000, Vector Laboratories) and DyLight Streptavidin 488 (1:500, BioLegend). Images were taken using a Leica SP8 Laser Scanning Confocal Microscope (Leica Microsystems) at the Cambridge Advanced Imaging Center and analyzed using ImageJ. Brain sections for DYN Nickel DAB labeling were incubated with rabbit anti-DYN B (1:12,000, a gift from Dr. Philippe Ciofi) (Griffond, Deray et al. 1993) followed by biotinylated goat anti-rabbit IgG (1:200, Vector Laboratories). Brain sections were further incubated in avidin-biotin peroxidase solution before reacting with Nickel DAB developing solution. Images were taken using an Olympus BX43 upright microscope with DP74 digital camera (Olympus Life Science) and analyzed using Image J.

Stereotaxic surgery and GCaMP fiber photometry

Adult female PNA, PPA, and their respective control *Kiss1*-Cre/+,*Ai126D* mice (10-14 weeks old) were anesthetized with 2% isoflurane, given meloxicam (5 mg/kg, s.c.), buprenorphine (0.05 mg/kg, s.c.), and dexamethasone (10 mg/kg, s.c.) and placed in a stereotaxic apparatus. A small hole was drilled in the skull and a unilateral indwelling optical fiber (400 µm diameter; 0.48 NA, Doric LenSEs, Quebec, Canada) was implanted directly above the mid-caudal ARN (2.0 mm posterior to bregma, 3.0 mm lateral to the superior sagittal sinus, 5.93 mm deep). Following optic fiber implantation, mice were monitored through recovery for 5 days and received post-operative analgesics (meloxicam, 5 mg/kg; orally) for up to two days. Then mice were handled daily for at least 3 weeks and habituated to the fiber

photometry recording setup before fluorescence signals were recorded from freely behaving
mice in their home cages.

Fiber photometry was undertaken as detailed previously (Clarkson, Han et al. 2017).
Photometry systems were built using Doric components (Doric Lenses, QC, Canada) and
National Instrument data acquisition board (TX, USA) based on a previous design (Lerner,
Shilyansky et al. 2015). Blue (465-490 nm) and violet (405 nm) LED lights were sinusoidally
modulated at frequencies of 531 and 211 Hz respectively. Both lights were focused onto a
single fiber optic connected to the mouse. The light intensity at the tip of the fiber was 30-80
micro watts. Emitted fluorescence signal from the brain was collected via the same fiber,
passed through a 500-550 nm emission filter, before focused onto a fluorescence detector
(Doric, QC, Canada). The emissions were collected at 10 Hz and the two GCaMP6
emissions were recovered by demodulating the 465-490 nm signals (calcium-dependent)
and 405 nm (calcium independent) signals.

Fluorescence signals were sampled using either a scheduled mode (5/15 seconds on/off) or
a continuous mode of light emission with custom software (Tussock Innovation). Twenty-
four-hour recordings were made from PNA, PPA, and their respective control female mice in
the diestrous stage starting 4 hours after lights-on. Four-day recordings were started on
proestrus for control mice and on any diestrous or metestrous stage for PNA mice.
Continuous data acquisition mode was used to provide high temporal resolution
comparisons of SEs. Signals from each SE were normalized to its peak (intact: -120 to +120
seconds; OVX: -120 to +200 seconds from the peak time). The full widths at half maximum
(FHWM) of SE were determined at half-maximal amplitude of the normalized SEs. Averaged
SE parameters for each animal were calculated before computing the means for each
treatment group.

Analysis was performed in MATLAB with the subtraction of 405 signal from 465-490 signal to
extract the calcium-dependent fluorescence signal. An exponential fit algorithm was used to
correct for baseline shift. The signal was calculated in dF/F (%) with the equation dF/F =
($F_{fluorescence}$-$F_{baseline}$)/ $F_{baseline}$) x 100. The Findpeaks algorithm was used to detect SEs with
peaks in dF/F greater than 50% of the maximum signal in each recording. Inter-SE intervals
represent the time between peak values of each SEs. Multi-peak SE in PNA animals and
cluster SE in OVX were analyzed as one SE and SE intervals were calculated from the first
peak.

Serial blood sampling, GnRH injection, and LH measurement

Mice were handled daily for at least 3 weeks to habituate for tail-tip bleeding. To examine the

relationship between ARN^{KISS} neuron SEs and pulsatile LH secretion, freely behaving mice

were attached to the fiber photometry system, and 4 µL blood samples were taken every 6

or 12 min from the tail-tip over a period of 2 to 4 hours during recording. For standard serial

blood sampling experiments, LH secretion was measured by obtaining 4 µL blood samples

every 10 minutes for 4 hours (between 12:00 and 17:00). At the end of the bleeding period,

200 ng/kg body weight of GnRH (Bachem, catalog No. 4033013, Switzerland) was injected

i.p., and two more blood samples were collected at 10-minute intervals. Whole blood

samples were diluted 1:15 in 1x PBS with 0.05% Tween 20 (PBS-T) for LH ELISA (Steyn,

Wan et al. 2013), an assay sensitivity of 0.04 ng/mL, as well as intra-assay and interassay

coefficients of variation of 9.3% and 10.5%, respectively. Pulses were defined and analyzed

by PULSAR with the settings optimized for intact female mice: G1 = 3.5, G2 = 2.6, G3 = 1.9,

G4 = 1.5, and G5 = 1.2 (Porteous, Haden et al. 2021).

K-means clustering

Data analysis and unsupervised k-means clustering analysis were performed without vaginal

cytology information as previously described (Vas, Wall et al. 2024). Synchronization events

with multiple peaks occurring within 160s were defined as mpSEs. In brief, 4-day recordings

were segmented into 24-hour periods before applying a customized MATLAB code to extract

features of the population activity of the ARN^{KISS} neurons including the (i) number of SEs

within a given time period; (ii) prevalence of different type of SEs (single- and mpSEs); (iii)

inter-SE intervals, variation in the inter-SE intervals (reflecting regularity of activity patterns);

and (iv) shape of the SEs (height, variation in height, half-duration). These 24-hour periods

were further fragmented using a 7-hour window size with 1-hour sliding steps. The first and

the last 3 hours of each 24-hour recording were extrapolated from the assignment of the 4th

and 21st hours, respectively. The following parameters were used to describe the activity

pattern of ARN^{KISS} neurons: (i) number of SEs, where mpSEs were counted as a single SE;

(ii) standard deviation (SD) of inter-SE intervals; (iii) number of mpSEs; (iv) average number

of peaks within an mpSEs; and (v) average duration of mpSEs. All these parameter values

were then linearly rescaled to values between 0 and 1. To perform k-means clustering on the

data set, we used RapidMiner Studio's (https://rapidminer.com, educational license) built-in

module (k = 5, number of runs with random initiations = 100, measure types =

"NumericalMeasures", numerical measure = "EuclideanDistance", max optimization steps =

100). Cluster centroid data and cluster labels (to each 7-hour time window) were exported

from RapidMiner Studio and were visualized in R (using "ggplot2", "readr" and "fmsb" libraries).

Progesterone injection

Progesterone stock solution (800 µg/mL) was prepared by dissolving 8 mg of progesterone (Sigma, Catalog No. P0130) in 1 mL ethanol followed by 1:10 dilution in sesame oil. Diestrous female mice were recorded for 3 hours before an i.p. injection of progesterone (4 mg/kg body weight) or vehicle (100 µL, sesame oil). A previous study has shown that i.p. injection of progesterone at 8 mg/kg body weight to C57BL/6 mice results in peak circulating progesterone concentrations of 110 ng/mL before dropping to 34 ng/mL 1 hour later (Wong, Ray et al. 2012). Circulating progesterone levels at proestrus in C57BL/6 mice are 22 ± 4 ng/mL (Wall, Desai et al. 2023), so we employed half of this dose to test the sensitivity of progesterone negative feedback to ARN^{KISS} neurons.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 10 software. Depending on different experimental designs, and the nature of data distribution, data were analyzed by Mann-Whitney U tests, Kruskal-Wallis test followed by Dunn's multiple comparison tests, two-way ANOVA followed by Sidak's multiple comparisons test, or two-way repeated-measures ANOVA followed by Sidak's multiple comparisons test. The threshold level for statistical significance was set at $P < 0.05$ with data presented as mean ± SEM with each dot representing one animal.

Acknowledgement

This research was supported by the Wellcome Trust (212242/Z/18/Z) to AEH and Medical Research Council (MR N013433-1) and Harding Distinguished Postgraduate Scholars Programme Leverage Scheme to ZZ.

Figures legends

*Figure 1 – figure supplement 1. **Peripubertal androgen (PPA) treatment causes an increase in body weight and a loss of estrous cyclicity.***

(**A**) Weekly body weight of animals (mean ± SEM) with blank (control: n=5) or dihydrotestosterone (PPA: n=4) capsules. Asterisks above each weekly data point denote significance from Sidak's multiple comparisons test. Asterisks between lines denote the mean effect of the capsule over 7 weeks. Two-way repeated-measure ANOVA with Sidak's multiple comparisons test. (**B**) The proportion of time spent at each stage of the estrous cycle over 21 days (n=4-5 per group). Two-way ANOVA followed by Sidak's multiple comparisons tests. (**C-D**) Representative graphs of estrous cycle pattern over 21 days in (**C**) control and (**D**) PPA females. P, proestrus; E, estrus; M, metestrus; D, diestrus. Data show mean ± SEM. Each circle is an individual animal. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure 1. **Slower pulse generator activity in PPA animals.**
Representative 24-hour photometry recordings showing synchronization events (SEs) observed in diestrous (**A**) control and (**B**) PPA females with the light-off period (19:00-07:00) represented by the shaded area and expanded views of the traces (13:00-17:00) from (A-B). Triangles point to identified SEs. (**C**) SE frequency and (**D**) SE intervals in control (n=5) and PPA (n=4) mice. Mann-Whitney U tests. (**E**) Frequency histograms showing relative SE frequencies occurring in 20-minute bins, calculated separately for controls (white, n=114, 5 mice) and PPA (orange, n=55, 4 mice). X-axis represents the bin centers. (**F**) Continuous recordings at 10 Hz sampling rate showing normalized profile of SE overlaid from control (black, 7 SEs from 5 animals) and PPA (orange, 5 SEs from 4 animals). 'a' and 'b' give values of full width at half maximum (FWHM) upswing and downswing respectively. (**G**) FWHM length of control (n=5) and PPA (n=4) animals. Mann-Whitney U test. (**H**) Durations of upswing and downswing for control (n=5) and PPA (n=4) animals, respectively. Kruskal-Wallis test followed by Dunn's multiple comparisons test. Data show mean ± SEM. Each circle is an individual animal. * $P < 0.05$

Figure 2. **Prenatal androgen exposure leads to increased anogenital distance and disrupted estrous cyclicity.**
(**A**) Anogenital distance (control: n=9; PNA: n=10) and (**B**) Body weight of control and PNA animals (n=12 per group). Mann-Whitney U tests. (**C**) Proportion of time spent at each estrous cycle stage over 21 days (n=7 per group). Two-way ANOVA followed by Sidak's post-hoc tests. (**D-E**) Representative graphs of estrous cycle pattern in (**D**) control and (**E**) PNA females. P, proestrus; E, estrus; M, metestrus; D, diestrus. (**F**) Serum testosterone level of control (n=13) and PNA (n=20) animals in diestrus measured by enzyme-linked immunosorbent assay (ELISA). The limit of detection for testosterone ELISA is 0.066 ng/mL. (**G-I**) Plasma levels of (**G**) androstenedione, (**H**) estradiol, and (**I**) progesterone in control and PNA animals measured using liquid chromatography-mass spectrometry (LC-MS). Blood

samples were collected at 10:00 diestrus (n=6 per group). Dotted lines represent the limit of
detection for each hormone. Androstenedione: 0.05 ng/mL; estradiol: 0.50 pg/mL;
progesterone: 0.05 ng/mL. All samples were below limit of detection for testosterone LC-MS
measurement (< 0.01 ng/mL). Mann-Whitney U tests. Data show mean ± SEM. Each circle
is an individual animal. ** $P < 0.01$, *** $P < 0.001$.

*Figure 2 – figure supplement 1. **Slightly increased kisspeptin immunofluorescence in***
the rostral arcuate nucleus of PNA mice.
Representative arcuate nucleus images of (**A**) Kisspeptin (KISS), (**B**) Neurokinin B (NKB)
immunofluorescent staining, and (**C**) Dynorphin (DYN) Nickel-3,3′-Diaminobenzidine (DAB)
staining from control and PNA animals, along with the quantified immunoreactive level in
rostral, middle, and caudal arcuate nucleus (rARN, mARN, cARN) in control (n=4-8) and
PNA (n=4-6) animals. Scale bars represents 100 μm. Two-way ANOVA followed by Sidak's
multiple comparisons test. Data show mean ± SEM. Each circle is an individual animal.
Asterisk represents significant multiple comparisons results. * $P < 0.05$

*Figure 3. **PNA animals exhibit highly variable patterns of ARN^KISS neuron SEs***
Representative 24-hour photometry recordings showing SEs observed in (**A,B**) two control
and (**C,D**) two PNA females with the light-off period (17:30-05:30 or 19:00-07:00)
represented by the shaded area. Recordings were performed when vaginal cytology
indicated diestrus. (**E**) SE frequency and (**F**) SE intervals in control (n=7) and PNA (n=6)
mice. Mann-Whitney U tests. (**G**) Frequency histograms showing relative SE frequencies
occurring in 20-minute bins, calculated separately for controls (black, n=123, 7 mice) and
PNA (red, n=98, 6 mice). X-axis represents the bin centers. Multi-peak SE (mpSE) with
peaks occurring within 160 seconds was considered as one SE with interval calculated from
the first peak. (**H**) Continuous (10 Hz) recording showing normalized mean profile of SE
overlaid from control (black, 22 SEs from 7 animals) and PNA (red, 20 SEs from 6 animals).
(**I**) FWHM length (upswing + downswing) in control (n=7) and PNA (n=6) animals. Mann-
Whitney U test. (**J**) Durations of FWHM upswing and downswing for control (n=7) and PNA
(n=6) animals, respectively. Kruskal-Wallis test followed by Dunn's multiple comparisons test.
Data show mean ± SEM. Each circle is an individual animal.

*Figure 4. **Increased total LH concentration but normal LH pulse frequency in PNA***
animals.
Representative examples from (**A**) control and (**C**) PNA mice showing the relationship of
SEs (black) with LH secretions (red). Triangles and red asterisks indicate identified SEs and
LH pulses, respectively. Normalized increase in LH plotted against the SEs in (**B**) control

and (**D**) PNA animals. The amplitudes of SEs and serum LH levels were normalized to their

peaks, with time 0 being the peak of LH in control (n=7 from 4 mice) and PNA (n=7 from 5

mice) animals. Representative examples of LH pulse profiles in (**E**) control and (**F**) PNA

animals. Red asterisks indicate identified LH pulses. (**G**) Total LH concentration, (**H**) LH

pulse frequency per hour, and (**I**) LH pulse amplitude in diestrous control (n=7, with 13 LH

pulses) and PNA (n=8, with 21 LH pulses) mice during 4-hour sampling at 10-minute

intervals. Mann-Whitney U tests. (**J**) Change in LH levels following an intraperitoneal

injection of GnRH (200 ng/kg, i.p.) in diestrous control (n=7) and PNA (n=11) mice. An

asterisk above the 20 minutes data point denote significance between LH levels in control

and PNA from two-way repeated-measure ANOVA followed by Sidak's post-hoc tests. (**K**)

GnRH-evoked changes in LH levels in control (n=7) and PNA (n=11) mice were calculated

by the difference in LH levels before (-10 minutes and 0 minutes) and after (10 minutes and

20 minutes) GnRH injections. Mann-Whitney U test. Data show mean ± SEM. Each circle is

an individual animal. * $P < 0.05$, ** $P < 0.01$

*Figure 5. **PNA animals exhibit cycling pulse generator activity.***

(**A**). Cluster centroid values for normalized parameters used in k-means clustering with a

schematic plot of corresponding photometry recordings patterns. The parameters used for

cluster assignments are labelled in Cluster_0 with the following axes: (top) standard

deviation (SD) of SE intervals, (top right) number of multi-peak SEs (mpSEs), (bottom right)

number of single SEs, (bottom left) number of peaks in mpSEs, (top left) duration of mpSEs.

All axes have a minimum value of 0 and maximum value of 1. (**B**) Bar graph indicating

proportion of the cluster-assignments for 4-day recordings in control (n=5) and PNA (n=4)

animals. Data show mean ± SEM. (**C-D**) Representative examples of four-day consecutive

recording of ARN^{KISS} activity and corresponding hourly k-means cluster assignments in (**C**)

two control animals starting in proestrus on day 1 and transiting to diestrus on day 4

according to vaginal smears; (**D**) two PNA animals remaining in either metestrus or diestrus

for four days. Light-off periods (19:00-07:00) are represented by the shaded area in the

photometry recording. Color fields represent the assigned clusters for the center (4th hour) of

the 7-hour moving time windows. Estrous stages of the mice determined by vaginal lavage

at the beginning and the end of each 24-hour recording were labelled above the

corresponding k-means cluster assignments: proestrus to estrus (PE), estrus to metestrus

(EM), metestrus to diestrus (MD), stuck in metestrus (MM), or stuck in diestrus (DD). K-

means clustering was performed without vaginal cytology information.

*Figure 6. **Faster ARN^{KISS} neuron synchronization events in "diestrous-like" PNA***

animals.

Representative 24-hour photometry recordings showing ARN^{KISS} neuron SEs observed in (**A**)
diestrous control and (**B**) algorithm-identified diestrous PNA females. The light-off period
(17:30-05:30) is represented by the shaded area. (**C**) SE frequency and (**D**) SE intervals in
control (n=7) and PNA (n=6) mice. Mann-Whitney U tests. (**E**) Histogram showing
percentage of SE interval frequencies occurring in 20-minute bins, calculated separately for
controls (black, n=119, 7 mice) and PNA (red, n=151, 6 mice) animals. X-axis represents the
bin centers.
779
*Figure 7. **Defective progesterone negative feedback in PNA animals**.*
(**A-D**) Examples of 15-hour photometry recordings of the pulse generator activity in (**A-B**)
control and (**C-D**) PNA animals receiving (**A**,**C**) vehicle (Veh) or (**B**,**D**) progesterone (P4, 4
mg/kg) i.p. injections. Dashed line represents when i.p. injection took place. Normalized
number of SEs in (**E**) control (n=3) and (**F**) PNA (n=3) animals before and after vehicle
(black) and P4 (red) injections. Data were analyzed in 3-hour bins and normalized to the
number of SEs before injections (-3-0 h) for each mouse. The asterisks above the data point
denote significant effects at indicated time periods compared to the pre-treatment period (-3-
0 h) by Sidak's multiple comparisons test. Data represent mean ± SEM. Two-way repeated-
measure ANOVA with Sidak's multiple comparisons tests. *** $P < 0.001$.
790
*Figure 8. **Similar ARN^{KISS} activity and SE profiles in control and PNA animals following***
ovariectomy.
Representative 24-hour photometry recordings showing ARN^{KISS} neuron SEs observed in (**A**)
control and (**B**) PNA females 3 weeks after ovariectomy (OVX) with the light-off period
(19:00-07:00) represented by the shaded area and expanded views of the traces (13:00-
15:00) from (**A-B**). (**C**) SE frequency and (**D**) SE intervals in control and PNA mice (n=4),
Mann-Whitney U tests. (**E**) Histogram showing relative SE frequencies occurring in 5-minute
bins, calculated separately for controls (n=212, 4 mice) and PNA (n=233, 4 mice) animals.
X-axis represents the bin centers. Each SE cluster was analyzed as a single SE with the
interval calculated from the first peak. (**F**) Continuous recordings at 10 Hz sampling rate
showing normalized profile of SEs following OVX overlaid from control (black, 14 SEs from 4
animals) and PNA (red, 10 SEs from 4 animals). (**G**) FWHM length of OVX control and PNA
animals (n=4). Mann-Whitney U test. (**H**) Durations of upswing and downswing for control
and PNA animals (n=4), respectively. $P > 0.05$, Kruskal-Wallis test followed by Dunn's
multiple comparisons test. Data show mean ± SEM. Each circle is an individual animal.
806

References

Ajayi, A. F. and R. E. Akhigbe (2020). "Staging of the estrous cycle and induction of estrus in experimental rodents: an update." Fertil Res Pract **6**: 5.

Auchus, R. J. (2014). "Steroid assays and endocrinology: best practices for basic scientists." Endocrinology **155**(6): 2049-2051.

Barnes, R. B., R. L. Rosenfield, S. Burstein and D. A. Ehrmann (1989). "Pituitary-ovarian responses to nafarelin testing in the polycystic ovary syndrome." N Engl J Med **320**(9): 559-565.

Barnes, R. B., R. L. Rosenfield, S. Burstein and D. A. Ehrmann (1989). "Pituitary-Ovarian Responses to Nafarelin Testing in the Polycystic Ovary Syndrome." New England Journal of Medicine **320**(9): 559-565.

Batrinos, M. L. (1993). "Diagnostic Dilemmas in Polycystic Ovarian Syndrome." Annals of the New York Academy of Sciences **687**(1): 230-234.

Bozdag, G., S. Mumusoglu, D. Zengin, E. Karabulut and B. O. Yildiz (2016). "The prevalence and phenotypic features of polycystic ovary syndrome: a systematic review and meta-analysis." Human Reproduction **31**(12): 2841-2855.

Caldwell, A. S., L. J. Middleton, M. Jimenez, R. Desai, A. C. McMahon, C. M. Allan, D. J. Handelsman and K. A. Walters (2014). "Characterization of reproductive, metabolic, and endocrine features of polycystic ovary syndrome in female hyperandrogenic mouse models." Endocrinology **155**(8): 3146-3159.

Caldwell, A. S. L., M. C. Edwards, R. Desai, M. Jimenez, R. B. Gilchrist, D. J. Handelsman and K. A. Walters (2017). "Neuroendocrine androgen action is a key extraovarian mediator in the development of polycystic ovary syndrome." Proc Natl Acad Sci U S A **114**(16): E3334-e3343.

Chhabra, S., C. R. McCartney, R. Y. Yoo, C. A. Eagleson, R. J. Chang and J. C. Marshall (2005). "Progesterone Inhibition of the Hypothalamic Gonadotropin-Releasing Hormone Pulse Generator: Evidence for Varied Effects in Hyperandrogenemic Adolescent Girls." The Journal of Clinical Endocrinology & Metabolism **90**(5): 2810-2815.

Clarkson, J., S. Y. Han, R. Piet, T. McLennan, G. M. Kane, J. Ng, R. W. Porteous, J. S. Kim, W. H. Colledge, K. J. Iremonger and A. E. Herbison (2017). "Definition of the hypothalamic GnRH pulse generator in mice." Proc Natl Acad Sci U S A **114**(47): E10216-e10223.

Coutinho, E. A. and A. S. Kauffman (2019). "The Role of the Brain in the Pathogenesis and Physiology of Polycystic Ovary Syndrome (PCOS)." Med Sci (Basel) **7**(8).

Coyle, C. S., M. Prescott, D. J. Handelsman, K. A. Walters and R. E. Campbell (2022). "Chronic androgen excess in female mice does not impact luteinizing hormone pulse frequency or putative GABAergic inputs to GnRH neurons." Journal of Neuroendocrinology **34**(4): e13110.

Daigle, T. L., L. Madisen, T. A. Hage, M. T. Valley, U. Knoblich, R. S. Larsen, M. M. Takeno, L. Huang, H. Gu, R. Larsen, M. Mills, A. Bosma-Moody, L. A. Siverts, M. Walker, L. T. Graybuck, Z. Yao, O. Fong, T. N. Nguyen, E. Garren, G. H. Lenz, M. Chavarha, J. Pendergraft, J. Harrington, K. E. Hirokawa, J. A. Harris, P. R. Nicovich, M. J. McGraw, D. R. Ollerenshaw, K. A. Smith, C. A. Baker, J. T. Ting, S. M. Sunkin, J. Lecoq, M. Z. Lin, E. S. Boyden, G. J. Murphy, N. M. da Costa, J. Waters, L. Li, B. Tasic and H. Zeng (2018). "A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-Type Targeting and Functionality." Cell **174**(2): 465-480.e422.

Dalkin, A. C., D. J. Haisenleder, G. A. Ortolano, T. R. Ellis and J. C. Marshall (1989). "The frequency of gonadotropin-releasing-hormone stimulation differentially regulates gonadotropin subunit messenger ribonucleic acid expression." Endocrinology **125**(2): 917-924.

Dam, E. W. C. M. V., F. Roelfsema, J. D. Veldhuis, F. M. Helmerhorst, M. Frölich, A. E. Meinders, H. M. J. Krans and H. Pijl (2002). "Increase in daily LH secretion in response to short-term calorie restriction in obese women with PCOS." American Journal of Physiology-Endocrinology and Metabolism **282**(4): E865-E872.

Esparza, L. A., T. Terasaka, M. A. Lawson and A. S. Kauffman (2020). "Androgen Suppresses In Vivo and In Vitro LH Pulse Secretion and Neural Kiss1 and Tac2 Gene Expression in Female Mice." Endocrinology **161**(12).

Gibson, A. G., J. Jaime, L. L. Burger and S. M. Moenter (2021). "Prenatal Androgen Treatment Does Not Alter the Firing Activity of Hypothalamic Arcuate Kisspeptin Neurons in Female Mice." eNeuro **8**(5).

Goodman, R. L., A. E. Herbison, M. N. Lehman and V. M. Navarro (2022). "Neuroendocrine control of gonadotropin-releasing hormone: Pulsatile and surge modes of secretion." Journal of Neuroendocrinology **34**(5): e13094.

Goodman, R. L., I. Holaskova, C. C. Nestor, J. M. Connors, H. J. Billings, M. Valent, M. N. Lehman and S. M. Hileman (2011). "Evidence that the Arcuate Nucleus Is an Important Site of Progesterone Negative Feedback in the Ewe." Endocrinology **152**(9): 3451-3460.

Griffond, B., A. Deray, D. Fellmann, P. Ciofi, D. Croix and C. Bugnon (1993). "Colocalization of prolactin- and dynorphin-like substances in a neuronal population of the rat lateral hypothalamus." Neurosci Lett **156**(1-2): 91-95.

Han, S. Y., G. Kane, I. Cheong and A. E. Herbison (2019). "Characterization of GnRH Pulse Generator Activity in Male Mice Using GCaMP Fiber Photometry." Endocrinology **160**(3): 557-567.

Han, S. Y., P. G. Morris, J.-C. Kim, S. Guru, M. Pardo-Navarro, S.-H. Yeo, H. J. McQuillan and A. E. Herbison (2023). "Mechanism of kisspeptin neuron synchronization for pulsatile hormone secretion in male mice." Cell Reports **42**(1): 111914.

Handelsman, D. J., E. Gibson, S. Davis, B. Golebiowski, K. A. Walters and R. Desai (2020). "Ultrasensitive Serum Estradiol Measurement by Liquid Chromatography-Mass Spectrometry in Postmenopausal Women and Mice." J Endocr Soc **4**(9): bvaa086.

Handelsman, D. J., M. Jimenez, G. K. S. Singh, J. Spaliviero, R. Desai and K. A. Walters (2015). "Measurement of Testosterone by Immunoassays and Mass Spectrometry in Mouse Serum, Testicular, and Ovarian Extracts." Endocrinology **156**(1): 400-405.

Handelsman, D. J. and L. Wartofsky (2013). "Requirement for mass spectrometry sex steroid assays in the Journal of Clinical Endocrinology and Metabolism." J Clin Endocrinol Metab **98**(10): 3971-3973.

Herbison, A. E. (2018). "The Gonadotropin-Releasing Hormone Pulse Generator." Endocrinology **159**(11): 3723-3736.

Herbison, A. E. (2021). "The dendron and episodic neuropeptide release." Journal of Neuroendocrinology **33**(11): e13024.

Jaime, J. and S. M. Moenter (2022). "GnRH Neuron Excitability and Action Potential Properties Change with Development But Are Not Affected by Prenatal Androgen Exposure." eNeuro **9**(6).

Kerbus, R. I., C. Decourt, M. A. Inglis, R. E. Campbell and G. M. Anderson (2024). "Androgen receptor actions on AgRP neurons are not a major cause of reproductive and metabolic impairments in peripubertally androgenized mice." Journal of Neuroendocrinology **n/a**(n/a): e13370.

Kumari, A. S., A. Haq, R. Jayasundaram, L. O. Abdel-Wareth, S. A. al Haija and M. Alvares
(2005). "Metformin monotherapy in lean women with polycystic ovary syndrome."
Reproductive BioMedicine Online **10**(1): 100-104.

Lei, L., L. Ding, J. Su, M. Liu, Q. Shi, J. Zhou, H. Sun and G. Yan (2017). "Attenuated expression
of MTR in both prenatally androgenized mice and women with the hyperandrogenic
phenotype of PCOS." PLoS One **12**(12): e0187427.

Leipheimer, R. E., A. Bona-Gallo and R. V. Gallo (1984). "The influence of progesterone and
estradiol on the acute changes in pulsatile luteinizing hormone release induced by
ovariectomy on diestrus day 1 in the rat." Endocrinology **114**(5): 1605-1612.

Lerner, T. N., C. Shilyansky, T. J. Davidson, K. E. Evans, K. T. Beier, K. A. Zalocusky, A. K. Crow,
R. C. Malenka, L. Luo, R. Tomer and K. Deisseroth (2015). "Intact-Brain Analyses Reveal
Distinct Information Carried by SNc Dopamine Subcircuits." Cell **162**(3): 635-647.

Liu, X., S.-H. Yeo, H. J. McQuillan, M. K. Herde, S. Hessler, I. Cheong, R. Porteous and A. E.
Herbison (2021). "Highly redundant neuropeptide volume co-transmission underlying
episodic activation of the GnRH neuron dendron." eLife **10**: e62455.

Lizneva, D., L. Suturina, W. Walker, S. Brakta, L. Gavrilova-Jordan and R. Azziz (2016).
"Criteria, prevalence, and phenotypes of polycystic ovary syndrome." Fertility and Sterility
106(1): 6-15.

McCarthy, E. A., D. Dischino, C. Maguire, S. Leon, R. Talbi, E. Cheung, C. D. Schteingart, P. J.
M. Rivière, S. D. Reed, R. A. Steiner and V. M. Navarro (2021). "Inhibiting Kiss1 Neurons With
Kappa Opioid Receptor Agonists to Treat Polycystic Ovary Syndrome and Vasomotor
Symptoms." The Journal of Clinical Endocrinology & Metabolism **107**(1): e328-e347.

McQuillan, H. J., J. Clarkson, A. Kauff, S. Y. Han, S. H. Yip, I. Cheong, R. Porteous, A. K.
Heather and A. E. Herbison (2022). "Definition of the estrogen negative feedback pathway
controlling the GnRH pulse generator in female mice." Nature Communications **13**(1): 7433.

McQuillan, H. J., J. Clarkson, A. Kauff, S. Y. Han, S. H. Yip, I. Cheong, R. Porteous, A. K.
Heather and A. E. Herbison (2022). "Definition of the estrogen negative feedback pathway
controlling the GnRH pulse generator in female mice." Nat Commun **13**(1): 7433.

McQuillan, H. J., S. Y. Han, I. Cheong and A. E. Herbison (2019). "GnRH Pulse Generator
Activity Across the Estrous Cycle of Female Mice." Endocrinology **160**(6): 1480-1491.

Moore, A. M., D. B. Lohr, L. M. Coolen and M. N. Lehman (2021). "Prenatal Androgen
Exposure Alters KNDy Neurons and Their Afferent Network in a Model of Polycystic Ovarian
Syndrome." Endocrinology **162**(11).

Moore, A. M., M. Prescott and R. E. Campbell (2013). "Estradiol Negative and Positive
Feedback in a Prenatal Androgen-Induced Mouse Model of Polycystic Ovarian Syndrome."
Endocrinology **154**(2): 796-806.

Moore, A. M., M. Prescott, C. J. Marshall, S. H. Yip and R. E. Campbell (2015). "Enhancement
of a robust arcuate GABAergic input to gonadotropin-releasing hormone neurons in a model
of polycystic ovarian syndrome." Proc Natl Acad Sci U S A **112**(2): 596-601.

Morales, A., G. Laughlin, T. Bützow, H. Maheshwari, G. Baumann and S. Yen (1996). "Insulin,
somatotropic, and luteinizing hormone axes in lean and obese women with polycystic ovary
syndrome: common and distinct features." The Journal of Clinical Endocrinology &
Metabolism **81**(8): 2854-2864.

Morales, A. J., G. A. Laughlin, T. Bützow, H. Maheshwari, G. Baumann and S. S. Yen (1996).
"Insulin, somatotropic, and luteinizing hormone axes in lean and obese women with
polycystic ovary syndrome: common and distinct features." The Journal of Clinical
Endocrinology & Metabolism **81**(8): 2854-2864.

Morris, P. G. and A. E. Herbison (2023). "Mechanism of Arcuate Kisspeptin Neuron
Synchronization in Acute Brain Slices From Female Mice." Endocrinology 164(12).

Navarro, V. M., M. L. Gottsch, C. Chavkin, H. Okamura, D. K. Clifton and R. A. Steiner (2009).
"Regulation of gonadotropin-releasing hormone secretion by
kisspeptin/dynorphin/neurokinin B neurons in the arcuate nucleus of the mouse." J
Neurosci 29(38): 11859-11866.

Nestler, J. E. and D. J. Jakubowicz (1997). "Lean women with polycystic ovary syndrome
respond to insulin reduction with decreases in ovarian P450c17 alpha activity and serum
androgens." J Clin Endocrinol Metab 82(12): 4075-4079.

Osuka, S., A. Iwase, T. Nakahara, M. Kondo, A. Saito, Bayasula, T. Nakamura, S. Takikawa, M.
Goto, T. Kotani and F. Kikkawa (2016). "Kisspeptin in the Hypothalamus of 2 Rat Models of
Polycystic Ovary Syndrome." Endocrinology 158(2): 367-377.

Pastor, C. L., M. L. Griffin-Korf, J. A. Aloi, W. S. Evans and J. C. Marshall (1998). "Polycystic
Ovary Syndrome: Evidence for Reduced Sensitivity of the Gonadotropin-Releasing Hormone
Pulse Generator to Inhibition by Estradiol and Progesterone1." The Journal of Clinical
Endocrinology & Metabolism 83(2): 582-590.

Patel, K., M. S. Coffler, M. H. Dahan, R. Y. Yoo, M. A. Lawson, P. J. Malcom and R. J. Chang
(2003). "Increased luteinizing hormone secretion in women with polycystic ovary syndrome
is unaltered by prolonged insulin infusion." J Clin Endocrinol Metab 88(11): 5456-5461.

Phylactou, M., S. A. Clarke, B. Patel, C. Baggaley, C. N. Jayasena, T. W. Kelsey, A. N.
Comninos, W. S. Dhillo and A. Abbara (2021). "Clinical and biochemical discriminants
between functional hypothalamic amenorrhoea (FHA) and polycystic ovary syndrome
(PCOS)." Clinical Endocrinology 95(2): 239-252.

Porteous, R., P. Haden, E. C. R. Hackwell, A. Singline, M. K. Herde, R. Desai, D. J. Handelsman,
D. R. Grattan and A. E. Herbison (2021). "Reformulation of PULSAR for Analysis of Pulsatile
LH Secretion and a Revised Model of Estrogen-Negative Feedback in Mice." Endocrinology
162(11).

Qiu, J., C. C. Nestor, C. Zhang, S. L. Padilla, R. D. Palmiter, M. J. Kelly and O. K. Rønnekleiv
(2016). "High-frequency stimulation-induced peptide release synchronizes arcuate
kisspeptin neurons and excites GnRH neurons." eLife 5: e16246.

Risal, S., M. Manti, H. Lu, R. Fornes, H. Larsson, A. Benrick, Q. Deng, C. E. Cesta, M. A.
Rosenqvist and E. Stener-Victorin (2021). "Prenatal androgen exposure causes a sexually
dimorphic transgenerational increase in offspring susceptibility to anxiety disorders."
Translational Psychiatry 11(1): 45.

Rodriguez Paris, V. and M. J. Bertoldo (2019). "The Mechanism of Androgen Actions in PCOS
Etiology." Medical Sciences 7(9): 89.

Roland, A. V. and S. M. Moenter (2011). "Prenatal androgenization of female mice programs
an increase in firing activity of gonadotropin-releasing hormone (GnRH) neurons that is
reversed by metformin treatment in adulthood." Endocrinology 152(2): 618-628.

Ruddenklau, A. and R. E. Campbell (2019). "Neuroendocrine Impairments of Polycystic Ovary
Syndrome." Endocrinology 160(10): 2230-2242.

Silva, M. S., M. Prescott and R. E. Campbell (2018). "Ontogeny and reversal of brain circuit
abnormalities in a preclinical model of PCOS." JCI Insight 3(7).

Silva, M. S. B., E. Desroziers, S. Hessler, M. Prescott, C. Coyle, A. E. Herbison and R. E.
Campbell (2019). "Activation of arcuate nucleus GABA neurons promotes luteinizing
hormone secretion and reproductive dysfunction: Implications for polycystic ovary
syndrome." EBioMedicine 44: 582-596.

Singh, J., C. O'Neill and D. J. Handelsman (1995). "Induction of spermatogenesis by androgens in gonadotropin-deficient (hpg) mice." Endocrinology **136**(12): 5311-5321.

Stener-Victorin, E., V. Padmanabhan, K. A. Walters, R. E. Campbell, A. Benrick, P. Giacobini, D. A. Dumesic and D. H. Abbott (2020). "Animal Models to Understand the Etiology and Pathophysiology of Polycystic Ovary Syndrome." Endocr Rev **41**(4).

Steyn, F. J., Y. Wan, J. Clarkson, J. D. Veldhuis, A. E. Herbison and C. Chen (2013). "Development of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult male mice." Endocrinology **154**(12): 4939-4945.

Sullivan, S. D. and S. M. Moenter (2004). "Prenatal androgens alter GABAergic drive to gonadotropin-releasing hormone neurons: implications for a common fertility disorder." Proc Natl Acad Sci U S A **101**(18): 7129-7134.

Taylor, A. E., B. McCourt, K. A. Martin, E. J. Anderson, J. M. Adams, D. Schoenfeld and J. E. Hall (1997). "Determinants of Abnormal Gonadotropin Secretion in Clinically Defined Women with Polycystic Ovary Syndrome." The Journal of Clinical Endocrinology & Metabolism **82**(7): 2248-2256.

The Rotterdam ESHRE/ASRM-sponsored PCOS consensus workshop group (2004). "Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS)." Human Reproduction **19**(1): 41-47.

Toosy, S., R. Sodi and J. M. Pappachan (2018). "Lean polycystic ovary syndrome (PCOS): an evidence-based practical approach." J Diabetes Metab Disord **17**(2): 277-285.

van Houten, E. L. A. F., P. Kramer, A. McLuskey, B. Karels, A. P. N. Themmen and J. A. Visser (2012). "Reproductive and Metabolic Phenotype of a Mouse Model of PCOS." Endocrinology **153**(6): 2861-2869.

Vas, S., E. Wall, Z. Zhou, L. Kalmar, S. Y. Han and A. E. Herbison (2024). "Long-term recordings of arcuate nucleus kisspeptin neurons across the mouse estrous cycle." Endocrinology.

Waldstreicher, J., N. F. Santoro, J. E. Hall, M. Filicori and W. F. Crowley, Jr. (1988). "Hyperfunction of the hypothalamic-pituitary axis in women with polycystic ovarian disease: indirect evidence for partial gonadotroph desensitization." J Clin Endocrinol Metab **66**(1): 165-172.

Wall, E. G., R. Desai, Z. Khant Aung, S. H. Yeo, D. R. Grattan, D. J. Handelsman and A. E. Herbison (2023). "Unexpected Plasma Gonadal Steroid and Prolactin Levels Across the Mouse Estrous Cycle." Endocrinology **164**(6).

Walters, K. A., R. B. Gilchrist, W. L. Ledger, H. J. Teede, D. J. Handelsman and R. E. Campbell (2018). "New Perspectives on the Pathogenesis of PCOS: Neuroendocrine Origins." Trends in Endocrinology & Metabolism **29**(12): 841-852.

Wang, L., W. Guo, X. Shen, S. Yeo, H. Long, Z. Wang, Q. Lyu, A. E. Herbison and Y. Kuang (2020). "Different dendritic domains of the GnRH neuron underlie the pulse and surge modes of GnRH secretion in female mice." eLife **9**: e53945.

Watanabe, Y., L. Fisher, R. E. Campbell and C. L. Jasoni (2023). "Defining potential targets of prenatal androgen excess: Expression analysis of androgen receptor on hypothalamic neurons in the fetal female mouse brain." Journal of Neuroendocrinology **35**(6): e13302.

Wierman, M. E., R. J. Auchus, D. J. Haisenleder, J. E. Hall, D. Handelsman, S. Hankinson, W. Rosner, R. J. Singh, P. M. Sluss and F. Z. Stanczyk (2014). "Editorial: The new instructions to authors for the reporting of steroid hormone measurements." J Clin Endocrinol Metab **99**(12): 4375.

Wildt, L., A. Häusler, G. Marshall, J. S. Hutchison, T. M. Plant, P. E. Belchetz and E. Knobil
(1981). "Frequency and amplitude of gonadotropin-releasing hormone stimulation and
gonadotropin secretion in the rhesus monkey." Endocrinology **109**(2): 376-385.
Wong, R., D. Ray and D. A. Kendall (2012). "Progesterone pharmacokinetics in the mouse:
implications for potential stroke therapy." Journal of Pharmacy and Pharmacology **64**(11):
1614-1620.
Yan, X., C. Yuan, N. Zhao, Y. Cui and J. Liu (2014). "Prenatal androgen excess enhances
stimulation of the GNRH pulse in pubertal female rats." Journal of Endocrinology **222**(1): 73-
85.
Yeo, S. H., V. Kyle, P. G. Morris, S. Jackman, L. C. Sinnett-Smith, M. Schacker, C. Chen and W.
H. Colledge (2016). "Visualisation of Kiss1 Neurone Distribution Using a Kiss1-CRE Transgenic
Mouse." J Neuroendocrinol **28**(11).
1051