The genetic architecture of floral trait divergence between hummingbird- and self-pollinated monkeyflower (*Mimulus*) species

Hongfei Chen[1,5], Colette S. Berg[2], Matthew Samuli[2], V. Alex Sotola[3,4], Andrea L. Sweigart[3], Yao-Wu Yuan[1*], Lila Fishman[2*]

[1] Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT 06269 USA

[2] Division of Biological Sciences, University of Montana, Missoula MT 59812 USA

[3] Department of Genetics, University of Georgia, Athens, GA 30602, USA.

[4]Current address: Biology Department, State University of New York at Oneonta, Oneonta, NY 13820 USA

[5] Current address: Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06520 USA

*For correspondence

SUMMARY

(1) Pollination syndromes are a key component of flowering plant diversification, prompting questions

about the architecture of single traits and genetic coordination among traits. Here, we investigate the

genetics of extreme floral divergence between naturally hybridizing monkeyflowers *Mimulus parishii*

(self-pollinated) and *M. cardinalis* (hummingbird-pollinated).

(2) We mapped quantitative trait loci (QTLs) for 18 pigment, pollinator reward/handling, and dimensional

traits in parallel sets of F_2 hybrids plus recombinant inbred lines and generated nearly isogenic lines

(NILs) for two dimensional traits, pistil length and corolla size.

(3) Our multi-population approach revealed a highly polygenic basis (n = 190 QTLs total) for pollination

syndrome divergence, capturing minor QTLs even for pigment traits with leading major loci. There was

significant QTL overlap within pigment and dimensional categories. Nectar volume QTLs clustered with

those for floral dimensions, suggesting a partially shared module. The NILs refined two pistil length

QTLs, only one of which has tightly correlated effects on other dimensional traits.

(4) An overall polygenic architecture of floral divergence is partially coordinated by genetic modules

formed by linkage (pigments) and likely pleiotropy (dimensions plus nectar). This work illuminates

pollinator syndrome evolution in a model radiation and generates a robust framework for molecular and

ecological genomics.

KEYWORDS – genetic architecture, QTL mapping, modularity, pollination syndrome, polygenic trait,

pigment., floral evolution

Introduction

Across flowering plants, distantly related taxa often show similarities in a suite of floral phenotypes that can be recognized as pollination syndromes (Fenster *et al.*, 2004; Dellinger, 2020), while switches between pollination syndromes are common even among closely related species. For example, the evolution from bee- to hummingbird-pollination, which is characterized by red color, copious nectar, and stigma and anthers exerted beyond a large bill-accommodating corolla, has happened more than 10 times independently in *Penstemon* alone (Wilson *et al.*, 2007; Wessinger & Hileman, 2016). Similarly, autogamous self-pollination, which is associated with inconspicuous coloration, reduced nectar rewards, and reduced anther-stigma separation (Sicard & Lenhard, 2011), has evolved countless times within animal-pollinated lineages (Stebbins, 1970; Barrett, 2002; Goodwillie *et al.*, 2005). Both convergence and divergence in pollination syndromes requires the correlated evolution of multiple traits to maintain floral phenotypic integration and reproductive fitness throughout the entire evolutionary path. Three non-exclusive genetic mechanisms may contribute to such coordinated evolution of pollination syndromes and other complex multi-trait strategies. At one extreme, natural selection on floral traits may be strong enough to restrict successful plants to a few discrete adaptive peaks even in the face of gene flow (Bleiweiss, 2001), building stereotypical multi-trait pollination syndromes from variation at multiple unlinked loci (Wessinger *et al.*, 2023). At the other extreme, pleiotropy among floral traits (Troth *et al.*, 2018) may enforce coordinated evolution of trait modules during pollination syndrome divergence (Smith, 2016; Wessinger & Hileman, 2016). Finally, genome architectures that suppress recombination in heterozygotes can package genes for functionally distinct traits into adaptive supergenes (Lowry & Willis, 2010; Hermann *et al.*, 2013; Edwards *et al.*, 2021; Liang *et al.*, 2023). Distinguishing among these explanations reveals very different barriers to traversing the phenotypic landscape as flowers evolve coordinately from one multi-phenotypic optimum to another.

Over the past three decades, quantitative trait locus (QTL) mapping has revealed the genetic architecture of pollination syndrome divergence between numerous closely-related pairs of plant species, including three sections of *Mimulus* monkeyflowers (Bradshaw *et al.*, 1998; Fishman *et al.*, 2002, 2013, 2015; Stankowski *et al.*, 2023), *Petunia* (Stuurman *et al.*, 2004), *Ipomoea* (Rifkin *et al.*, 2021; Liao *et al.*, 2021) and many others. Across angiosperm diversity from monocots to diverse eudicots, these genome-wide approaches reveal two broad patterns. First, divergence in pollinator-attraction and reward traits such as flower color, scent, or nectar volume is often controlled by few loci, each of

moderate to large effect, whereas divergence in dimensional traits (corolla or reproductive organ size

and shape) often involves more loci, each of small effect. This pattern may in part reflect focus on

attraction/reward traits in studies of transitions among distinct animal pollination syndromes vs. a primary

focus on floral dimensions in shifts to self-pollination. However, it may also reflect consistent differences

in the underlying genetic variation and patterns of selection for different categories of trait. Second, co-

localization of QTLs for at least some floral traits suggests floral integration/modularity through

pleiotropy and/or the adaptive evolution of supergene architecture to facilitate trait packaging in the

face of gene flow (Yeaman & Whitlock, 2011). Floral integration/modularity has been considered a

plausible mechanism that facilitates rapid evolution of pollination syndromes (Diggle, 2014; Wessinger *et*

al., 2014; Smith, 2016; Dellinger *et al.*, 2019; Kostyun *et al.*, 2019). However, few studies have clearly

identified intra-floral evolutionary modules, and the pattern of integration and modularity across sets of

floral traits remains an open question. Understanding both the build-up (integration) and the breakdown

(modularity) of trait correlations to generate complex floral strategies requires a locus- and gene-scale

understanding of the genomic bases of pollination syndromes.

Although QTL mapping has robustly advanced understanding of the genetic architecture of divergence

in multi-factorial trait syndromes, connecting their effects to the underlying genes remains a challenge.

First, especially when traits are polygenic and the underlying loci have small effects, scans of single

experimental hybrid mapping populations capture the overall architecture of genetic variation, but not

individual loci and their effects. Thus, replication of mapping experiments tests QTL robustness to

environmental variation and increases confidence in shared QTLs. Second, even major QTLs often

contain 10s to 100s of genes, confounding pleiotropy and linkage as causes of genetic correlation and

QTL coincidence. Similarly, fine-scale identification of causal variants benefits from isolation from

segregating background effects. Construction of near-isogenic lines (NILs) allows both detection and

isolation, as in the identification of major flower color and scent loci in *Petunia* (e.g. (Klahre *et al.*, 2011;

Berardi *et al.*, 2021) and *Mimulus* (Bradshaw & Schemske, 2003; Yuan *et al.*, 2013b; Byers *et al.*, 2014;

Yuan *et al.*, 2016; Liang *et al.*, 2023). However, because NIL construction generally involves strong

selection for resemblance to the introgressing parent for a focal trait combined with opposing selection

on other traits, it may not capture the evolutionary contributions of more complex genetic architectures

to trait (co-)variation. Thus, a combination of genome-wide QTL characterization plus targeted NIL

construction is a powerful approach to understand both the genome-wide architecture and gene-scale

causes of divergence in complex pollination syndromes.

Here, we employ an integrated approach to characterize the genetic architecture of pollination

syndrome divergence and the detailed genetics of key traits between closely related monkeyflowers

Mimulus cardinalis and *M. parishii* (Phyrmaceae, section *Erythranthe*). These taxa, which are

hummingbird-pollinated and self-pollinated, respectively (Fig. 1), are the most florally divergent

members of a recent adaptive radiation (Nelson *et al.*, 2021b). Nevertheless, they hybridize in areas of

range overlap, producing genomic signatures of recent introgression (Nelson *et al.*, 2021b). Along with

bee-pollinated *M. lewisii*, which likely resembles the common ancestor of all three taxa, these closely

related species are a model system for understanding the genetics of floral divergence and speciation

(Yuan, 2019). Work in this system has provided key insights into the evolution and molecular biology of

divergent pollination syndromes and their roles in local adaptation and speciation (Hiesey *et al.*, 1971;

Bradshaw *et al.*, 1995; Schemske & Bradshaw, 1999; Ramsey *et al.*, 2003; Fishman *et al.*, 2013; Yuan *et*

al., 2013b, 2014; Stathos & Fishman, 2014; Fishman *et al.*, 2015; Yuan *et al.*, 2016; Peng *et al.*, 2017;

Nelson *et al.*, 2021b,a; Liang *et al.*, 2023). New resources, including chromosome-scale reference

genomes (www.Mimubase.org), dense linkage maps (Sotola *et al.*, 2023), and stable transformation

protocols (Yuan, 2019), now enable genome-wide mapping, gene-scale dissection, and molecular

characterization of the loci underlying *Erythranthe* floral diversity. Notably, because *Mimulus parishii* x *M.*

cardinalis hybrids segregate more freely than other crosses in the group (Fishman *et al.*, 2013, 2015;

Sotola *et al.*, 2023), they were key to the recent genetic and molecular dissection of a novel speciation

supergene (Liang *et al.*, 2023). With the current study, we take a major step toward a similarly detailed

understanding of the full suite of floral traits contributing to pollination syndrome divergence.

To robustly characterize the genetic architecture of extreme floral divergence, we examine patterns of

floral trait (co)inheritance and map QTLs in one extensively phenotyped focal F_2 population, then map

QTLs for a subset of traits in independent F_2 and RIL growouts to capture additional minor QTLs in

distinct environmental and genetic backgrounds. We assess patterns of QTL size and coincidence within

and across trait categories and propose three hypotheses about the genetic architecture: (i):

Divergence of floral traits produced by relatively simple biochemical pathways such as flower color are

controlled by few loci with moderate to large effects, whereas dimensional traits involve more loci of

small effects; (ii) minor QTLs are more subject to stochasticity and differences in environment and

genetic background (for RILs) among our mapping populations; and (iii) traits within a category (e.g.,

pigment or dimensions) are controlled by integrated sets of overlapping QTLs (modules) while overlap

between categories is relatively low. Finally, we characterize independent nearly isogenic lines (NILs) for

two highly polygenic dimensional traits, flower size and pistil length; these NILs provide proof of concept

(and sound some useful cautionary notes) on the dissection of floral dimensional QTLs. Our multi-trait

and multi-generation approach provides a broad and deep characterization of the genetics of floral

divergence and opens paths toward understanding both its molecular bases and effects on patterns of

mating and introgression in wild populations.

Fig. 1 Floral phenotypes. (a) The parental line *M. cardinalis* CE10, *M. parishii* PAR, and their F1 hybrid. (b) Representative F2 progeny. (c) The main floral traits measured in this study, using *M. parishii* as examples. CTL: corolla tube length; KTL: calyx tube length; KTW: calyx tube width; STL: stamen length; PIL: pistil length; CTW: corolla tube width; VPL: ventral petal length; VPW: ventral petal width; CLL: corolla limb length; CLW: corolla limb width. Scale bars, 3 mm.

Materials and Methods

Study system, mapping populations, and phenotyping

Hummingbird-pollinated *M. cardinalis* (Phrymaceae) is a perennial herb of low-elevation seeps and riverbanks from northern Baja California to southern Oregon (Angert & Schemske, 2005; Angert, 2009). It has red flowers with long tubular corolla, copious nectar, and exserted stigma and anthers (Fig. 1a). *M. parishii* is an annual self-pollinating herb generally found along ephemeral streams in southern California. *M. parishii* has small pale pink flowers, little stigma–anther separation, and no nectar (Fig. 1a, Table 1). All hybrids in this study were generated from two highly inbred parental lines: Sierran CE10 for *M. cardinalis* (Yuan *et al.*, 2013b) and PAR for *M. parishii* (Fishman *et al.*, 2015; Nelson *et al.*, 2021b; Liang *et al.*, 2023). F_1 hybrids were generated with PAR as the seed parent and selfed to generate F_2 seeds, while recombinant inbred lines (RILs) were generated by single-seed-descent from F_2 individuals through 3-6 generations of self-fertilization (Sotola *et al.*, 2023). F_2 hybrids (Fig. 1b) were grown in two separate greenhouse common gardens at the University of Connecticut (UC_F_2) and the University of Montana (UM_F_2), and the RILs were grown at the University of Georgia (UGA), as detailed in Supplementary Methods S1.

In the UC_F_2 growout, we measured three pigment, two pollinator reward/handling and nine dimensional floral traits, plus flowering time, on F_2s (n = 253) plus parental lines and F_1 hybrids (n = 8 each) (Fig. 1). We scanned the ventral petal of each flower to quantify petal lobe anthocyanin (PLA) and carotenoid (PLC) pigment intensity. The proportion of red (R), green (G), and blue (B) pixels in a square area of the same size of the adaxial surface of the petal were estimated from scanned images using Image J (http://rsbweb.nih.gov/ij/). The relative petal lobe anthocyanin concentration was estimated using the equation "[(R + B)/2] – G", a simple and effective approach previously used for genetic mapping of anthocyanin content variation (Yuan *et al.*, 2013b) and independently verified in other plant species (Valle *et al.*, 2018). Similarly, the relative carotenoid concentration was estimated by the equation "[(R + G)/2] – B", which also proved effective as our QTL mapping successfully located the previously characterized carotenoid locus *YELLOW UPPER* (*YUP*) (see Results). Nectar guide anthocyanin (NGA) and nectar guide trichome length (NGT) were visually scored in F_1 and F_2 hybrids on semi-quantitative scales defined by the parental extremes (CARD = 9 and 7, respectively, PAR = 1 for both). Nectar volume (NEV) was measured for two flowers per individual on their first day of opening, using a pipette accurate to 1.5µL. To reduce environmental effects, nectar was measured at 4:00 PM-7:00 PM after watering at

12:00-1:00PM each day. Floral dimensions (Fig. 1c) were measured on one of the second pair of open

flowers using a digital caliper, and stigma-anther distance calculated as pistil length – stamen length. F_2

trait distributions were tested for normality with a Shapiro–Wilk test implemented in R v. 3.6.0. Because

some traits were non-normally distributed, we calculated pairwise Spearman's correlation coefficients (r)

for phenotypic correlations using the *psych::corr.test* function in R v. 3.6.0; we calculated broad-sense

heritability for each trait and genotypic correlations following (Fishman *et al.*, 2002).

Overlapping subsets of key floral traits were measured using parallel methods in the UM_F_2 (n = 278)

and UGA_RIL growouts (n = 145) (Supplementary Methods S2). Traits with a shared abbreviation

represent the same floral dimension, except for pistil length (PIL) and stigma-anther separation (SAS),

which included the stigma lobes in the UC_F_2 and UGA_RILs (Fig. 1c) but not the UM_F_2s. In the UM_F_2

growout, we characterized an additional pollinator handling trait associated with pollination syndrome

divergence in *Mimulus*, touch-sensitive stigma closure (Friedman *et al.*, 2017). The bilobed stigmas of *M.*

cardinalis close rapidly (<5s; like a tiny venus flytrap) when touched, while *M. parishii* stigmas are

insensitive and/or non-closing (Fishman *et al.*, 2024). Prior to the other floral measurements, a single

tester touched each stigma head-on with a pencil eraser to mimic pollinator contact and scored stigma

closure speed on a 4-point scale (0 = no closure = PAR-like, 3 = fast closure = CE10-like, 1 and 2 =

slower and faster intermediates, respectively).

Genetic context and QTL mapping

We previously constructed a joint linkage map of the two F_2 populations and a separate map of the RILs

using windowed genotypes from ddRAD sequences aligned to the CE10 *M. cardinalis* reference genome

(Sotola *et al.*, 2023). The dense F_2 and RIL linkage maps are generally highly collinear with each other

and the physical genome assemblies (www.Mimubase.org). However, an *M. cardinalis*-specific reciprocal

translocation involving portions of Chromosomes 6 and 7 (Fishman *et al.*, 2013; Stathos & Fishman,

2014) causes inter-chromosome linkage (i.e. they form a single linkage group in F_2s: LG6&7), excess

heterozygosity, and underdominant hybrid sterility. In addition, a gametophytic Dobzhansky-Muller

incompatibility involving Chr4 (~7-8 Mb) and Chr8 (~12-40 Mb) eliminates three genotypic classes in F_2

and later hybrids (Sotola *et al.*, 2023).

We conducted QTL mapping in QTL Cartographer (Wang *et al.*, 2005) in parallel on the three

populations using composite interval mapping (model 6, with forward-backward regression to choose 10

cofactors, window size 10 cM). LOD significance thresholds for QTL detection for each trait were set with

1000 permutations. Due to substantial retained heterozygosity in the RILs (Sotola *et al.*, 2023), we used

F_2 rather than RIL population settings to allow full estimation of QTL effects using all individuals. To

evaluate QTL coincidence across populations and traits, as well as define physical bounds, we defined a

1.5 LOD-drop confidence interval (CI) around each peak. For the two F_2 populations, which share a

linkage map, overlap was directly determined. For comparing F_2s and RILs, we translated QTL peaks and

intervals to the physical positions of boundary markers. We assigned QTL numbers within traits across

populations based on CI overlap (Table S1). We tested whether the mean effect size (r^2) of all 144 unique

(at level of trait) QTLs differed among the trait categories using ANOVA in JMP 18. For the nine traits

measured in all three populations (94 named QTLs), we similarly tested whether QTLs detected in one (n

= 55), two (n = 32) or three (n = 7) populations were, on average, of different magnitude. Using the

physical positions of all QTLs in Table S1, we calculated the degree of QTL overlap between each pair

using the Jaccard index, following (Liao *et al.*, 2021) (Supplementary Methods S3), then calculated the

mean (and standard error) of QTL overlap within and between trait categories. We tested whether there

was greater overlap within than between trait categories using 1000 permutations in which traits were

randomly assigned to categories. Because some non-dimensional traits may plausibly share a partial

genetic basis with flower size, we also specifically assessed the overlap of nectar volume and flowering

time QTLs with those in the three multi-trait categories.

Construction and characterization of nearly isogenic lines (NILs)

NILs were constructed via phenotypic selection prior to QTL mapping, and thus provide an independent

approach to dissecting the genetics of dimensional traits. To construct pistil length (PIL) NILs in the *M.*

parishii genetic background, we chose an F_2 individual with overall similarity to *M. parishii* in both floral

and vegetative traits, but with conspicuously longer pistil, for serial backcrossing (with NIL as pollen

donor) to *M. parishii.* From each backcross growout of ~95 plants, we selected one individual that

closely resembled *M. parishii* but with longer pistil for the next round. Bulked segregant analysis in a

BC_2S_1 population (two rounds of backcrosses followed by one round of selfing) and subsequent

genotyping in the same population using markers within the identified fragments revealed two

chromosomal regions (Chr 4: 0-4.1 Mb; Chr 6: 42-52 Mb) introgressed from *M. cardinalis* that co-

segregate with pistil length. Further genotyping of a BC_3S_1 population narrowed the chromosome 6

locus to a genomic interval at 42.85 Mb-51.76 Mb (Supplementary Methods S4). Selfing a BC_3S_1

individual heterozygous for both fragments generated nine genotypes across the two loci, which also

allowed us to decompose the BC$_3$ NIL into two NILs. A similar crossing approach was used to generate a

corolla limb length (CLL) NIL representing flower size (Supplementary Methods S4).

Results

Floral trait divergence, distributions, and correlations in F$_2$ hybrids

In the focal UC_F$_2$ grow-out, the CE10 *M. cardinalis* and PAR *M. parishii* parental lines were highly

differentiated for all traits, with F$_1$ and F$_2$ means always intermediate (Table 1). Floral dimensions were

normally distributed in F$_2$s, but petal carotenoid values (PLC) were bimodally distributed while petal lobe

anthocyanins (PLA) were skewed toward CE10 and nectar volume (NEV) toward PAR (Fig. S1).

Quantitative traits other than NEV, which had negative H^2 estimates due to its extreme skew, had high

broad-sense heritability (H^2 > 0.49). The UM_F$_2$ and RILs had similar distributions for each shared trait,

but mean nectar volume was much higher in the RILs (Fig. S1). All floral dimensions other than stigma-

anther separation (SAS) were positively correlated both phenotypically (r$_P$) and genetically (r$_G$) (Fig. S2).

The key mating system trait of stigma-anther separation was most highly correlated with pistil length (r$_G$

= 0.57), less with the other length metrics (r$_G$ = 0.25 - 0.31), and uncorrelated with width metrics. Floral

dimensional traits were only moderately correlated with flowering time (FLT) but flower length traits (KTL,

CTL, STL, PIL) were highly correlated with nectar volume (r$_P$= 0.58-0.70, r$_G$ not calculable for NEV due to

negative H^2) and petal lobe carotenoids were strongly correlated with pistil length and stamen length

(both r$_G$ > 0.6).

Genetic architecture - QTL mapping of individual traits in multiple mapping populations

We identified 190 floral QTLs, which define 144 QTL locations if collapsed (within traits) across the three

mapping populations (Fig. 2, Table S1).

Pigment traits – As expected from previous work, petal lobe carotenoids (PLC) were primarily affected by

a fully shared *M. cardinalis*-recessive major QTL on LG4 (coincident with *YUP*; PLC4.1 in Table S1). We

also detected two smaller carotenoid loci in the F$_2$s, and two more in the RILs. For petal lobe

anthocyanins (PLA), four loci were detected: PLA4.1 was detected in all three growouts and coincident

with the *YUP-SOLAR-PELAN* supergene, PLA4.2 and PLA3 were found in both F$_2$s but not the RILs, and

PLA6&7 was found in both RILs and UC_F$_2$. Nectar guide anthocyanins (NGA) were under the control of

two major loci, NGA3 (r^2 = 0.21) and NGA4 (r^2 = 0.31), and two additional small QTLs.

Table 1. Floral trait variation (means +/- SE for *M. cardinalis* (CE10), *M. parishii* (PAR), and their F_1 and F_2 hybrids in UC_F_2 growout. Broad-sense heritability (H^2) was calculated following Fishman *et al.* (2002). Traits marked ** were also measured in both UM_F_2 and RIL mapping populations, while nectar volume (NEV; *) was also characterized in RILs.

Trait	*M. cardinalis* CE10 (n=8)	*M. parishii* PAR (n=8)	F_1 hybrid (n=8)	F_2 hybrids (n=253)	H^2
Pigment					
Petal lobe carotenoid (PLC)**	99.38 ± 0.44	2.29 ± 0.50	13.89 ± 0.48	48.01 ± 2.12	1.00
Petal lobe anthocyanin (PLA)**	74.39 ± 1.11	21.46 ± 1.93	64.15 ± 1.02	52.38 ± 1.23	0.96
Nectar guide anthocyanin (NGA)	9	1	5	5.20 ± 0.12	NA
Pollinator reward/handling					
Nectar volume (NEV)*	45.95 ± 3.83	0 ± 0	3.28 ± 0.17	5.30± 0.29	-0.35
Nectar guide trichome (NGT)	7	1	4	4.40 ± 0.10	NA
Dimension					
Corolla limb length (CLL)	36.45 ± 0.69	8.18 ± 0.09	19.97 ± 0.49	20.21 ± 0.26	0.89
Corolla limb width (CLW) **	20.23 ± 1.62	8.94 ± 0.09	18.61 ± 0.43	18.30 ± 0.22	0.49
Ventral petal width (VPW)	12.46 ± 0.39	3.28 ± 0.07	7.27 ± 0.12	7.52 ± 0.09	0.82
Ventral petal length (VPL)	10.43 ± 0.21	2.77 ± 0.06	5.51 ± 0.10	6.11 ± 0.07	0.91
Corolla tube width (CTW)**	8.29 ± 0.27	4.43 ± 0.27	7.69 ± 0.15	7.26 ± 0.08	0.84
Calyx tube length (KTL)	22.30 ± 0.17	9.82 ± 0.28	16.17 ± 0.29	16.90 ± 0.14	0.90
Calyx tube width (KTW)	7.68 ± 0.13	2.98 ± 0.08	5.06 ± 0.08	5.34 ± 0.06	0.91
Corolla tube length (CTL)**	32.85 ± 0.35	11.74 ± 0.31	21.68 ± 0.23	22.46 ± 0.20	0.93
Pistil length (PIL)**	45.88 ± 0.43	11.76 ± 0.21	25.89 ± 0.29	27.20 ± 0.28	0.96
Stamen length (STL)**	42.93 ± 0.28	13.31 ± 0.47	24.91 ± 0.27	25.30 ± 0.24	0.94
Stigma-anther separation (SAS)**	2.95 ± 0.28	-1.55 ± 0.34	0.99 ±0.16	1.90 ± 0.09	0.71
Flowering time (FLT)**	79.38 ± 0.84	53.5 ± 1.07	55.13 ± 0.30	66.68 ± 0.51	0.94

Reward and handling traits – We detected four nectar volume QTLs in the UC_F_2 and two completely

non-overlapping ones in the RILs. RIL QTLs NEV5 and NEV8 had absolutely ~4x larger effects than the

largest F_2 one (NEV6&7; r^2 = 0.20), which explained only ~1/7 of the parental difference in NEV. The four

largest NEV QTLs (RIL pair, plus NEV2 and NEV6&7) were each coincident with dense clusters of floral

dimension QTLs (see below). Nectar guide trichomes (NGT) and stigma closure speed (SCS) QTLs, which

were scored on semi-quantitative scales, had relatively low explanatory power in the segregating F_2

populations (all r^2: 0.04-0.11). However, QTLs for these traits explained from 20% (each of the two *M.*

parishii-recessive stigma closure QTLs: SCS5 and SCS6&7) to 40% of the parental difference (additive

NGT8). Thus, they provide key targets for understanding the genetic underpinnings of these important

but understudied components of floral syndrome evolution.

Fig. 2. Quantitative trait loci (QTLs) for floral traits associated with pollination syndrome divergence between selfer *Mimulus parishii* and hummingbird pollinated *M. cardinalis*, as detected in three mapping populations. Bars show QTL 1.5 LOD drop confidence intervals and arrows indicate QTL peak position (up = QTL effect is in direction expected from parental divergence, down = opposite). The UC_F_2 QTL bars (all traits other than stigma closure) are unbordered, UM_F_2 QTL bars are bordered in gray, and RIL QTL bars are bordered in black. Relative QTL magnitude is indicated by the color-intensity of the QTL bar. The x-axis is position in centiMorgans (cM) on each of the seven F_2 linkage groups; these correspond to eight chromosomes due to a reciprocal translocation between Chr 6 and Chr 7 in *M. cardinalis* vs. *M. parishii* that generates inter-chromosomal linkage (Fishman *et al.*, 2013; Stathos & Fishman, 2014; Sotola *et al.*, 2023)

Dimensional traits and flowering time – In the UC_F_2, floral size was polygenic (71 dimensional QTLs).

QTL sizes were correspondingly small, with the leading QTL for each trait explaining from 11% (CLL,

CTL, KTW) to ~20% (VPL, VPW) of the F_2 variance. All primary size QTLs in this F_2 population moved trait

values in the direction expected from the parental difference. Consistent with the transgressive

segregation of stigma-anther separation in the UC_F_2s (Fig. S1), 2 of the 6 QTLs for this composite trait

had opposite effects from expectation (Table S1). For the six shared dimensional traits, we mapped 43,

39, and 29 QTLs in the UC_F_2, UM_F_2 and RILs, respectively, and ~1/3 (35/111) were shared across two

or more populations. For flowering time, FLT4.1 (all three mapping populations) and FLT8.1 (UC_F_2 and

RILs) were shared, but the other 7 QTLs were each found in only a single population. Flowering time

QTLs are not particularly small in absolute terms (all $2a > 6$ days), so this variation may reflect true

genotype x environment interactions for phenology.

Patterns across trait categories – genetic architecture, repeatability, modularity, and directionality

Overall, QTLs for pigment traits were nearly twice as large as dimensional QTLs (0.135 vs.0.075, p =

0.003), while flowering time and handling/reward QTLs were intermediate (Fig. 3a). For shared traits,

pigment QTLs and larger ones were significantly more likely to be detected in all three mapping

populations (both P < 0.005). However, QTLs detected in one or two populations were equally small

(0.074 vs. 0.077, P = 0.94 by Tukey's HSD). This pattern of moderate repeatability suggests that each

mapping population stochastically detected only a subset of minor loci from the larger (shared) pool of

variants influencing each polygenic trait.

Fig. 3. Summary of QTL effects by trait category. (a) Effect size (r^2) for QTLs in pigment, pollinator reward/handling, and dimensional categories, plus flowering time. (b) QTL overlap (Jaccard index ± 1SE) within and between the three multi-trait categories, plus overlap of each with nectar volume and flowering time.

Overlap of QTLs within both pigment (Jaccard index = 0.25) and dimension (0.30) categories were

significantly greater than null expectation (p = 0.036 and 0.0001, respectively), suggesting that each

forms a distinct intra-floral evolutionary module (Fig. 3b). Much lower overlap (0.04) within the pollinator

reward/ handling set is not surprising, given its grab-bag of traits. However, nectar volume QTLs strongly

overlapped with those for floral dimensions (0.29), suggesting a joint evolutionary module with flower

size, while overlap of flowering time and dimension QTLs was intermediate (Fig. 3b). Overall, only 7%

(10/144) unique QTLs had additive effects opposite to those expected from the parental difference,

suggesting consistent divergent natural selection (Orr, 1998). Notable exceptions were flowering time

and the composite floral trait of stigma-anther separation, with >1/4 and 1/3 (respectively) of their QTLs

opposite to expectation.

Dissection of floral dimension QTLs with NILs

(a)

C; C   C; P   C; H

H; C   H; P   H; H

P; C   P; H   P; P

(b) 0.0 Mb

65.5 Mb

M. parishii   *PIL 4.1* NIL   *PIL 6&7.2* NIL   *M. cardinalis*

Fig. 4 Pistil NILs. (a) Phenotypes of 9 genotypic combinations of the *PIL4.1* and *PIL6&7.2* loci. Genotypic designations: C, homozygous for *M. cardinalis*; H, heterozygous; P, homozygous for *M. parishii*. Under each image, the *PIL4.1* genotype is followed by the *PIL6&7.2* genotype. The colored dashed lines indicate the positions of the stigmas and long anthers, respectively. Scale bar, 3 mm. (b) Graphical genotypes of the *PIL4.1* NIL and the *PIL6&7.2* NIL. The colored bars represent the 8 chromosomes.

The long-pistil (PIL) and flower size (CLL) NILs isolated *M. cardinalis* alleles in the *M. parishii* background using repeated rounds of backcrossing and selfing with phenotypic selection (Supplemental Methods S3). The long-pistil NIL contained two unlinked regions introgressed from *M. cardinalis*, corresponding to QTLs *PIL4.1* (0-4.1 Mb on Chr4) and *PIL6&7.2* (42.85 Mb-51.76 Mb on Chr6) (Fig. 4). In a BC$_3$S$_1$ population segregating for all genotypic combinations at these two loci, both QTLs exhibited primarily additive effects on pistil length, consistent with QTL effects. The two NIL-isolated QTLs both had absolute effects smaller than in the F$_2$ population but maintained their relative sizes: *PIL4.1* (2a = 5.68mm in F2s, 3mm in NIL) vs. *PIL6&7.2* (2a = 4.24mm in F2s, 1mm in NIL). There was no evidence of an epistatic interaction between

these two loci, as the double NIL was 4mm larger than *M. parishii* (Table S2). The CE10-homozygous

PIL4.1 NIL also increased stamen and corolla tube lengths, but not our measure of overall flower size

(CLL), relative to *M. parishii* (Table 2). In contrast, the *PIL6&7.2* NIL had little effect on stamen or corolla

tube length (Table S2), suggesting that it contains a gene that specifically regulates pistil length.

After selfing and phenotypic selection (Supplemental Methods S3), the final BC$_3$S$_1$ NIL for corolla limb

length (CLL) captured a large region including *CLL6&7.2* (Fig. 5). This NIL was heterozygous across much

of Chr6 (43.54Mb-60.3Mb) and Chr7 (0-29.34Mb), but *M. cardinalis* homozygous near the LG6&7

translocation breakpoint (60.32Mb-61.12Mb on Chr6; 29.34Mb-29.82Mb on Chr7). Unlike the more

tightly localized *PIL6&7.2* introgression (which it includes in heterozygous state), the *CLL6&7.2* NIL has

increased pistil, stamen, and corolla tube length, as well as greater CLL, relative to *M. parishii* (Fig. 5a,

Table S3). Because the *CLL6&7.2* NIL includes a recombination-suppressed translocation breakpoint

associated with underdominant pollen sterility (Sotola *et al.*, 2023), its multiple phenotypic effects may

reflect *PIL6&7.2*, additional linked genes, and pleiotropic effects of sterility per se (Fishman et al. 2015).

(a)

M. parishii   *CLL6&7.2* NIL

(b) 0.0 Mb

65.5 Mb

CLL6&7.2 NIL

Fig. 5 Flower size NIL. (a) Flowers of *M. parishii* and the *CLL6&7.2* NIL. Scale bar, 4mm. (b) Graphical genotype of the *CLL6&7.2* NIL.

Discussion

We used QTL mapping in three hybrid growouts, as well as NIL construction, to investigate the genetic

architecture of pollination syndrome divergence between hummingbird-pollinated *Mimulus cardinalis*

and self-pollinated *M. parishii*. Despite some post-mating barriers (Sotola *et al.* 2023), these taxa are

good models for understanding the early stages of speciation: hybrids between *M. parishii* and *M.*

cardinalis are florally diverse (Fig. 1), more fit than their respective hybrids with bee-pollinated *M. lewisii*

(Fishman *et al.*, 2013, 2015; Stathos & Fishman, 2014), and subject to ongoing introgression in areas of

range overlap (Nelson *et al.*, 2021b). Along with directly illuminating the quantitative genetic basis of

floral evolution, this work provides a foundation for understanding the molecular genetic basis,

evolutionary history, and speciation consequences of complex pollination syndromes.

Overall, we identified a notably complex genetic basis for divergence between multi-trait pollination

syndromes. Our findings largely confirm the initial hypotheses of a highly polygenic and thus less

repeatably-mapped genetic basis for dimensional traits relative to pigment traits. Further, elevated

genetic integration (shared QTL positions) within both pigment and dimensional trait categories

(compared to low QTL overlap across categories) suggests trait modularity, as predicted. However, we

also mapped new minor QTLs even for pigment traits influenced by known major supergenes (Liang *et*

al., 2023), and characterized a complex (and background/environment-dependent) genetic basis for the

key reward trait of nectar volume. This genetic architecture contrasts with both the highly oligogenic

and tightly integrated (including across pigment, reward, and dimension traits) genetic architecture of

divergence between *M. cardinalis* and bee-pollinated *M. lewisii* (Bradshaw *et al.*, 1995, 1998; Fishman *et*

al., 2013) and highly polygenic dimensional divergence between *M. parishii* and *M. lewisii* (Fishman *et*

al., 2015)or selfer *M. nasutus* and bee-pollinated *M. guttatus* (Fishman *et al.*, 2002). Along with

chromosome-scale genomes and new functional genetic tools (Yuan, 2019), this complexity and diversity

of genetic architectures reinforces the value of the *M. cardinalis* species complex for understanding the

ecological, molecular, and evolutionary mechanisms of floral syndrome divergence.

A long walk to dramatic floral divergence – genetic basis of individual traits

Although they show evidence of hybridization and ongoing introgression in areas of current range

overlap (Nelson *et al.*, 2021b), hummingbird pollinated *M. cardinalis* and selfer *M. parishii* likely both

evolved from a bee-pollinated *M. lewisii*-like ancestor, which florally resembles their F_1 hybrid (Fig. 1).

Thus, it is not surprising that the genetic architecture of their floral divergence is a composite of patterns

and loci seen in hybrids of each crossed to *M. lewisii*. Furthermore, the overall patterns of QTL size and

directionality for different trait categories largely parallel predictions based on previous empirical work,

as well as the underlying molecular and developmental pathways, where known. Specifically, we

expected major leading QTLs for pigment traits and the reward trait of nectar volume. These key

pollination syndrome traits may offer a limited set of mutational targets and pathways to adaptive

evolution for both biochemical and evolutionary reasons (Bleiweiss, 2001; Wessinger & Rausher, 2012)

and color divergence often maps to major on-off switches (Wessinger & Hileman, 2020). In contrast,

dimensional traits may present near-infinite molecular targets for minor-effect mutations and exhibit high

levels of intra-specific standing variation readily available for rapid polygenic adaptation under novel

directional selection (Roels & Kelly, 2011; Troth *et al.*, 2018). However, as summarized below, we find

minor QTLs for all categories of traits in this wide cross. Although strong bias of QTL effects suggests

that directional natural selection has driven trait divergence, this abundance of genetic contributors to

variation in hybrids may reflect drift and relaxed selection along the lineage leading to selfer *M. parishii*

compared to the more constrained shift between discrete bee- and hummingbird- attraction peaks

(Bleiweiss 2001).

Consistent with expectation and previous genetics in this system, carotenoid (PLC) and anthocyanin

(PLA, NGA) pigment traits each had a leading major QTL ($r^2 > 0.19$) in the core UC_F_2 population. Co-

localized *PLC4.1* and *PLA4.1* correspond to the small RNA locus *YELLOW UPPER* (*YUP*) and the *R2R3-*

MYB gene *PETAL LOBE ANTHOCYANIN* (*PELAN*), respectively (Liang *et al.*, 2023), which form a

pigment supergene novel to *Mimulus* section *Erythranthe* (Liang *et al.*, 2023) along with another

anthocyanin-regulating *MYB, SISTER OF LIGHT AREAS (SOLAR)* (Liang *et al.*, 2022, 2023). It is unknown

whether *PELAN* or *SOLAR* (or another linked gene) underlies the coincident nectar guide anthocyanin

QTL (*NGA4.1*); nonetheless, QTL co-localization here reflects remarkably tight linkage of carotenoid and

anthocyanin pigment variants affecting distinct pathways rather than pleiotropic effects of a single gene.

Beyond the *YUP- SOLAR- PELAN* supergene, however, our multiple maps revealed an unexpectedly

complex and novel genetic basis for pigment traits. The four additional QTLs for each pigment trait had

widely variable effects (Table S1), and only partially overlapped with *Mimulus* pigment loci identified with

alternative approaches. For example, NGA3/PLA3 contains *RED TONGUE* (*RTO*), an R3-MYB gene

previously shown to repress anthocyanin biosynthesis in both petal lobes and nectar guides of *M. lewisii*

flowers (Ding *et al.*, 2020). However, *ROSE INTENSITY (ROI)* (Yuan *et al.*, 2013b), which controls the

reduced anthocyanin of pale pink Sierran *M. lewisii* relative to *M. cardinalis*, was not coincident with any

QTLs in this study, though PLA6&7, a rare wrong-way QTL at which *M. parishii* alleles confer darker

anthocyanin pigmentation, contains several *R3-MYBs* related to *ROI* and *RTO*. Thus, there are clearly a

diversity of mutational paths to complex floral pigment patterns in *Mimulus* flowers, paralleling the layers

of complexity of similar pollination syndrome shifts in *Petunia* (Berardi *et al.*, 2021). Overall, our pigment

QTL data provide an unusually nuanced picture of the divergence of floral attraction traits, a roadmap

for molecular characterization of the underlying genes, and the opportunity to study their effects on

pollination ecology in natural and artificial hybrids.

The pollinator reward and handling traits (nectar volume, stigma closure speed, nectar guide trichomes)

are each essentially lost in selfer *M. parishii* but may have followed distinct evolutionary paths to that

endpoint. High nectar volume (NEV) maintains high hummingbird visitation rates to *M. cardinalis*

(Schemske & Bradshaw, 1999), while touch-sensitive stigma closure (SCS) enhances pollen export in

outcrossing monkeyflowers (Fetscher, 2001) and repeatedly degenerates in selfers (Friedman *et al.*,

2017; Fishman *et al.*, 2024). In contrast, nectar guide trichomes (NGT) may be under relaxed selection in

both *M. cardinalis* and *M. parishii* relative to bee-pollinated *M. lewisii* (Chen & Yuan, 2024) and the

common ancestor of these taxa. Like pigment traits, both nectar volume and nectar guide trichomes are

under the control of major leading QTLs in *M. lewisii* x *M. cardinalis* and *M. parishii* x *M. lewisii* hybrids

respectively, while loss of stigma closure appears moderately polygenic in yellow monkeyflowers

(Fishman et al. 2024). Thus, QTLs for these traits, although not as extensively studied as floral pigments

or dimensions, provide key comparative insight into the genetic architecture and (eventually) molecular

basis of pollinator syndrome divergence.

Both nectar guide trichomes (NGT; UC_F_2 only) and stigma closure (SCS; UM_F_2 only) exhibit genetic

novelty and complexity relative to parallel studies in *Mimulus*. Intriguingly, none of the four moderate

NGT QTLs identified here includes the MYB transcription factor GUIDELESS (1.13 Mb on CE10 Chr6

(Chen & Yuan, 2024), previously inferred to be the major locus causal of NGT loss in *M. parishii* relative

to bee-pollinated *M. lewisii* (Chen & Yuan, 2024) and also implicated as key determinant of nectar guide

formation (both pigment and cell shape) in *M. lewisii* via mutagenesis (Yuan *et al.*, 2013a). This suggests

both a much more complex genetic basis for nectar guide divergence between these two non-bee

species and possibly epistatic interactions masking nonsynonymous mutations inferred as causal of NGT

loss in *M. parishii* (Chen & Yuan, 2024). Similarly, the two QTLs of moderate effect (SCS5 and SCS6&7)

for stigma closure (which together explain only 1/3 of total F_2 variance, suggesting many additional

minor loci) do not map to chromosomal regions syntenic with the five QTLs that fully explain similar shift

between selfer/noncloser *Mimulus nasutus* and fast-closer *M. guttatus* (Fishman *et al.*, 2024). However,

SCS6&7 contains a Mechanosensitive Channel of Small Conductance-like 10 (MSL10) gene homologous

to a highly stigma-expressed candidate mechanosensor identified in the *M. guttatus* complex (Fishman

et al., 2024). Furthermore, as in the *M. guttatus* complex, stigma closure QTLs appear independent of

the other floral trait reductions associated with the evolution of selfing (Fig. 2, Table S1), suggesting

abundant genomic targets for independent losses of this plant movement trait in selfers. Although

pollinator-handling traits have not been as extensively studied as pigments, dimensions, or rewards,

both nectar guide cell shape (e.g. (Glover *et al.*, 1998) and stigma movement (Newcombe, 1922;

Fishman *et al.*, 2024) vary widely across Lamiales (>25,000 species). By revealing a diversity of underlying

438 genetic mechanisms, even just within monkeyflowers, this work is a key step toward understanding the

439 integration (and dis-integration in selfers) of pollination syndromes in diverse taxa with tubular, bilaterally

440 symmetric flowers.

441 Nectar volume, with six small-to-moderate sized QTLs in the UC_F_2s and RILs together, has an even

442 more complex architecture, including polygenicity, epistasis, and gene x environment interactions. This

443 sharply contrasts with the simple genetic architecture invoked for this trait in *M. lewisii* x *M. cardinalis*

444 hybrids, which identified two leading QTLs each >30% of the F_2 variance (Bradshaw *et al.*, 1998). Our

445 largest F_2 QTLs, NEV2 and NEV6&7, were much smaller (r^2 = 0.12- 0.17) and all four summed to only

446 ~35% of the parental difference. Moreover, a model including all 2-way QTL interactions found

447 significant ($P < 0.005$) interactions of NEV6&7 with NEV4 and NEV2; along with the strong skew of NEV

448 phenotypes toward low (*M. parishii*-like) values (Fig. S1), this suggests that epistatic interactions may

449 contribute to parental divergence. The genetics of nectar volume was also context-dependent; the two

450 RIL QTLs were both much larger (10.9-14.3 µLs each) than any found in UC_F_2s and in distinct locations.

451 Differences between greenhouse conditions, as well as postzygotic barriers that further skew allele

452 frequencies in RIL populations (Sotola *et al.*, 2023), may contribute to the lack of repeatability. However,

453 as discussed further below, nectar volume may be a particularly complex composite trait whose

454 divergence encompasses both highly polygenic dimensional traits and simpler biochemical switches.

456 **Causes and consequences of floral integration within and between trait categories**

457 The comparative study of pollination syndromes suggests that integrated evolution of the many floral

458 traits associated with a given pollination syndrome likely involves coordinated change via a smaller

459 number of genetically-correlated floral modules (Smith, 2016). However, although floral modules have

460 been identified from patterns of genetic correlation and QTL overlap in hybrids in several systems, there

461 is no consensus yet about the prevalence of floral modules within and across trait categories or in

462 different evolutionary contexts (e.g. bee-to-hummingbird vs. outcrosser-selfer or adaptation-with-gene

463 flow vs. allopatric divergence). For example, nectar traits and floral dimensions each form tightly intra-

464 correlated but distinct modules in a transition from outcrossing to selfing in *Ipomaea* (Liao *et al.*,

465 2021)while bee-to-hummingbird transitions run the gamut from highly integrated across all traits in

466 *Mimulus* (Bradshaw *et al.*, 1995, 1998; Fishman *et al.*, 2013) to largely un-coordinated except by the

action of natural selection in the face of gene flow (Wessinger *et al.*, 2014, 2023; Kostyun *et al.*, 2019).

Here, we identify significantly intra-correlated but distinct modules for floral color and dimensions, as

well as coordination between the latter module and the reward trait of nectar volume (Fig. 3b). While

supporting a role for trait integration and modularity for pollination syndrome evolution, our findings

also underline key challenges in applying this conceptual framework at the QTL level.

Elevated QTL overlap within the natural category of pigment traits might suggest pleiotropy as the

source of genetic correlation in hybrids as well as trait integration throughout divergent evolution.

However, because the carotenoid and anthocyanin/flavanol pigment pathways are biochemically distinct

(Grotewold, 2006) and a key multi-pigment supergene has been molecularly dissected in our system, we

know that tight integration of pigments traits has causes beyond pleiotropy. In particular, the *YUP-*

SOLAR-PELAN supergene on Chr 4 strongly influences all three pigment traits due to tight linkage of

adjacent genes (Liang *et al.*, 2023). Linkage may also underlie QTL coincidence for the two anthocyanin

traits (PLA and NGA) more broadly. MYB transcription factors, including *PELAN* and *SOLAR*, often occur

in tandem clusters within plant genomes, providing fertile ground for multiple independent (i.e. linked

but potentially non-pleiotropic) mutations affecting anthocyanin production in different tissues. Indeed,

such genomic flexibility is key to the proposed importance of both transcription factors (Romani &

Moreno, 2021) and gene duplicates (Ohno, 1999) as key loci for evolutionary innovation. Thus, although

both pigment and dimensional modules identified in hybrids may reflect pleiotropy, tight linkage is a

particularly plausible alternative source for the former. However, their maintenance as modules

contributing to pollinator syndrome shifts potentially implicates natural selection acting not only on the

individual traits or genes but on trait coordination in the face of gene flow.

Although the tremendous diversity of floral morphologies implies freedom to evolve along many paths,

developmental coordination among floral whorls (e.g. petals and stamens) is expected from their serial

homology. Indeed, strong genetic correlations (Fig. S2) and elevated QTL overlap among dimensional

traits (Figs. 2 & 3) are consistent with a general "flower size" developmental module (Krizek & Anderson,

2013), even if not always due to pleiotropy. Further, coordination of floral dimensions by many multi-

trait QTLs is consistent both with parallel interspecific transitions (Fishman *et al.*, 2002; Goodwillie *et al.*,

2006; Kostyun *et al.*, 2019; Liao *et al.*, 2021) and with a highly pleiotropic and polygenic basis to

standing variation for corolla size traits within *Mimulus* populations (Troth *et al.*, 2018). However, the key

evolutionary steps in pollination syndrome shifts may often require breaking rather than following the

genetic correlations among dimensional traits within populations, and thus may involve rare or novel

uncoordinated variants. In particular, the exertion of sexual parts beyond the corolla in hummingbird

pollination and loss of stigma-anther distance in autogamous selfers entail changes in the *relative*

lengths of different floral whorls, while hummingbird pollination in tubular flowers also involves shifts in

corolla length vs. width (Fig. 1). Thus, although many shared loci may influence overall flower size

differences during pollination syndrome divergence, key evolutionary shifts in the relative size and

position of floral organs must be achieved via specific loci with isolated effects on single traits or via

disproportionate shifts in size at many loci.

Thus, despite significant modularity for floral dimensions, the subset of size QTLs with disproportionate

effects on different whorls (i.e. less integrated) may be particularly important for divergence in pollination

syndromes. For example, the *M. parishii* corolla tube and pistil are the same length, with stamens that

are slightly exerted past both (Table 1, Fig. 1), whereas the *M. cardinalis* style extends 13mm past the

corolla tube and 3mm past the stamens. This dramatic exertion of the style (a key feature or

hummingbird pollination in tubular flowers) implies the action of several PIL-only loci or many multi-trait

size loci with slightly greater effects on pistil length (PIL) than stamen (STL) or corolla tube length (CTL).

Stigma-anther separation, the key trait for self-pollination, requires similar disproportionality. SAS QTLs

exhibit all possible combinations, including joint PIL/STL length QTLs without effects on SAS, regions

affecting all three traits, and QTLs that only affect stigma-anther separation (Fig. 2, Table S1). Further,

genetic dissection of individual pistil length NILs revealed that one multi-trait QTL (*PIL6&7.2)* could be

isolated as a PIL-only factor, while the other (*PIL4.1*) retained parallel effects on multiple length traits.

The former is a promising target for dissecting the genetics of mating system evolution *per se*, while the

latter is a candidate for overall flower size evolution. More generally, our finding of significant but not

particularly high integration of dimensional traits (e.g., QTL overlap indices < ½ those of a similar study

in *Ipomoea* outcrosser-selfer hybrids (Liao *et al.*, 2021), suggests that genetic coordination of floral

dimensions may not be a strong constraint when selection acts on shape as well as size.

In addition to the two modules matching pre-assigned trait categories, our results reveal integration

between categories at both the level of the individual QTL (flowering time) and genome-wide (nectar

volume). Genome-wide QTL and genetic correlations between flowering time and floral dimensions were

not particularly elevated (Fig. 3), but we identified one intriguing genome region that may reflect speed-

size co-ordination. FLT4.1, at which *M. parishii* alleles confer much earlier flowering ($2a$ = 7.8 to 18.6

days), shares a 0.5-1Mb region on Chr4 with a partially overlapping set of 13 floral-size QTLs across all

populations. This co-incidence could be due to linkage among mulitple independent genes, but may

reflect a flowering time locus that pleiotropically mediates tradeoffs between speed and size (i.e., fast

flowering = small flowers), as in *M. guttatus* (Troth *et al.*, 2018). Further genetic dissection of this region

promises a clean test of those alternatives. Nectar volume was even more integrated with dimension

traits (Fig. 3), with particularly high genetic correlations with flower length traits (Fig. S2), QTL overlap as

high as within the dimension module (Fig. 3b) and both NEV QTLs in F_2s coincident with multi-trait size

QTLs (Fig. 2). This result intriguingly contrasts with a recent study of floral integration in *Ipomoea* (Liao *et*

al., 2021), where nectar traits formed an evolutionary module only weakly correlated with flower size and

few QTL positions were shared. While this difference may in part reflect our measurement of only a

single nectar trait, allowing no tests for an even more coordinated nectar-only module, nectar-size

integration may reflect the specifics of floral development in tubular flowers. Some components of

nectar volume variation (e.g., post-development sugar or water provisioning) may be independent of

flower size genes, whereas others (e.g., nectary size) may be directly downstream of developmental

shifts causing reduced corollas in *M. parishii*. Further work with additional nectar traits measured in

multiple environmental conditions and genetic backgrounds (given non-overlapping RIL and F2 QTLs)

will be necessary to tease apart the mechanisms underlying this apparent integration.

Dissecting the genes underlying polygenic flower size variation in hybrids

To understand the molecular mechanisms of floral evolution and trace their history across species

divergence, we must get our hands on the causal variants. NIL generation with phenotypic selection is a

common tool for fine-mapping of focal mutants in *Arabidopsis*, and has been used in *Mimulus* to dissect

major loci controlling pigment divergence (Yuan *et al.*, 2013b, 2016; Liang *et al.*, 2022, 2023). Here, pistil

length and corolla limb length NILs confirm and refine key dimensional QTLs (see above) and provide a

major step towards understanding the molecular basis of poorly understood polygenic traits. This is

particularly important for the key mating system trait of pistil length; mutation and hormonal

manipulation can alter pistil length dramatically and independently of other floral dimensions (Ding *et*

al., 2021), but natural species differences appear highly polygenic and potentially pleiotropic (Tables S1-

S2). Although both pistil length NIL intervals still contain many genes, high recombination on

chromosome ends in these hybrids (Liang *et al.* 2022; 2023; Sotola *et al.* 2023) makes their further

dissection and identification of the causal gene(s) feasible. In contrast, the flower size NIL CLL6&7 (which

overlaps with PIL6&7.2) corresponds to a reciprocal translocation (Fishman *et al.*, 2013; Stathos &

Fishman, 2014; Sotola *et al.*, 2023) resistant to further genetic dissection; complementary approaches,

such as analyses of gene expression networks conducted on segregating NILs with distinct phenotypes

(Langfelder & Horvath, 2008), will be necessary to narrow down functional candidates within this region.

Across all traits, additional targeted fine-mapping of even minor QTLs, along with functional approaches,

promises a detailed understanding of the many contributors to floral trait divergence in monkeyflowers.

Conclusions

Overall, our complementary mapping approaches reveal a polygenic genetic architecture even for

pollinator attraction (pigment) and reward traits with major leading QTLs, as well as shared QTL hotspots

causing strong genetic correlations within pigment and dimensional categories. In addition to enriching

understanding of the genetic architecture and modularity of components of pollination syndromes, this

work creates a strong foundation for further molecular genetic characterization of floral traits and

investigations of the evolutionary genomics of species barriers in this classic model system.

Acknowledgements

We are grateful to T. Nelson and K. Baesen for assistance with the genotyping and phenotyping at the

University of Montana (UM) and to logistical support from the UM ECOR Plant Growth Facility and

Genomics Core. We thank C. Liu, M. Opel, and M. Moriarty for plant care in the UConn Botanical

Conservatory. We are thankful to S. McCann and P. Beardsley for help in generating the RILs. This work

was supported by NSF grants DEB-0846089, DEB-1457763 and OIA-1736249 (to LF), IOS-1827645 (to

ALS and YWY), DEB-1350935 (to ALS), and IOS-2319721 (to YWY).

Competing interests

The authors declare no competing interests.

Author contributions

HC – data collection, analysis, and visualization, drafting of manuscript; CSB – data collection, analysis,

and visualization; MS – data collection; VAS – data collection, analysis, and visualization, drafting of

initial manuscript; ALS – conceptualization, data collection and interpretation, editing of manuscript;

YWY – conceptualization, data collection and interpretation, drafting and editing of manuscript; LF –

conceptualization, data collection, analysis, visualization and interpretation, drafting and editing of

manuscript. All authors read and are accountable for the submitted manuscript.

Data availability

The raw sequence data (PRJNA1003462, PRJNA948041) and individual genotypes

(doi:10.5061/dryad.v6wwpzh1m) used for generating linkage maps are publicly available. The phenotype

matrices used for quantitative genetics and QTL mapping are archived at Dryad

https://datadryad.org/stash/share/lDhHN-mqGkS5zdMy9QIcCWuhDNzqm1npxaDtyjAhO6k and will be

fully released upon publication. The raw sequencing data used to identify the potential DNA fragments

responsible for the long pistils of the PIL NIL and the large flowers of the CLL NIL have been deposited

to NCBI under the accession numbers PRJNA1116746 and PRJNA1117077, respectively, and will be

released upon publication.

References

Angert AL. **2009**. The niche, limits to species' distributions, and spatiotemporal variation in demography across the elevation ranges of two monkeyflowers. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 19693–19698.

Angert AL, Schemske DW. **2005**. The evolution of species' distributions: reciprocal transplants across the elevation ranges of Mimulus cardinalis and M. lewisii. *Evolution* **59**: 1671–1684.

Barrett SCH. **2002**. The evolution of plant sexual diversity. *Nature Reviews Genetics* **3**: 274–284.

Berardi AE, Esfeld K, Jäggi L, Mandel T, Cannarozzi GM, Kuhlemeier C. **2021**. Complex evolution of novel red floral color in Petunia. *The Plant Cell* **33**: 2273–2295.

Bleiweiss R. **2001**. Mimicry on the QT(L): genetics of speciation in Mimulus. *Evolution* **55**: 1706–1709.

Bradshaw HD, Otto KG, Frewen BE, McKay JK, Schemske DW. **1998**. Quantitative Trait Loci Affecting Differences in Floral Morphology Between Two Species of Monkeyflower (Mimulus). *Genetics* **149**: 367–382.

Bradshaw HD, Schemske DW. **2003**. Allele substitution at a flower colour locus produces a pollinator shift in monkeyflowers. *Nature* **426**: 176–178.

Bradshaw HD, Wilbert SM, Otto KG, Schemske DW. **1995**. Genetic mapping of floral traits associated with reproductive isolation in monkeyflowers (Mimulus). *Nature* **376**: 762–765.

Byers KJRP, Vela JP, Peng F, Riffell JA, Bradshaw HD. **2014**. Floral volatile alleles can contribute to pollinator-mediated reproductive isolation in monkeyflowers (Mimulus). *The Plant Journal* **80**: 1031–1042.

Chen H, Yuan Y-W. **2024**. Genetic basis of nectar guide trichome variation between bumblebee- and self-pollinated monkeyflowers (Mimulus): role of the MIXTA-like gene GUIDELESS. *BMC Plant Biology* **24**: 62.

Dellinger AS. **2020**. Pollination syndromes in the 21st century: where do we stand and where may we go? *New Phytologist* **228**: 1193–1213.

Ding B, Li J, Gurung V, Lin Q, Sun X, Yuan Y. **2021**. The leaf polarity factors SGS3 and YABBYs regulate style elongation through auxin signaling in Mimulus lewisii. *New Phytologist* **232**: 2191–2206.

Ding B, Patterson EL, Holalu SV, Li J, Johnson GA, Stanley LE, Greenlee AB, Peng F, Bradshaw HD, Blinov ML, *et al.* 2020. Two MYB Proteins in a Self-Organizing Activator-Inhibitor System Produce Spotted Pigmentation Patterns. *Current Biology* **30**: 802–814.

Edwards MB, Choi GPT, Derieg NJ, Min Y, Diana AC, Hodges SA, Mahadevan L, Kramer EM, Ballerini ES. **2021**. Genetic architecture of floral traits in bee- and hummingbird-pollinated sister species of Aquilegia (columbine). *Evolution* **75**: 2197–2216.

Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. **2004**. Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution, and Systematics* **35**: 375–403.

Fetscher AE. **2001**. Resolution of male-female conflict in an hermaphroditic flower. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **268**: 525–529.

Fishman L, Beardsley P, Stathos A, Williams CF, Hill JP. **2015**. The genetic architecture of traits associated with
the evolution of self-pollination in Mimulus. *New Phytologist* **205**: 907–917.

Fishman L, Kelly AJ, Willis JH. **2002**. Minor quantitative trait loci underlie floral traits associated with mating
system divergence in Mimulus. *Evolution* **56**: 2138–2155.

Fishman L, McIntosh M, Nelson TC, Baesen K, Finseth FR, Stark-dykema E. **2024**. Undoing the 'nasty:
dissecting touch-sensitive stigma movement (thigmonasty) and its loss in self-pollinating monkeyflowers. *Biorxiv*.

Fishman L, Stathos A, Beardsley P, Williams CF, Hill JP. **2013**. Chromosomal rearrangements and the genetics
of reproductive barriers in Mimulus (monkeyflowers). *Evolution* **67**: 2547–2560.

Friedman J, Hart KS, Bakker MC den. **2017**. Losing one's touch: Evolution of the touch-sensitive stigma in the
Mimulus guttatus species complex. *American Journal of Botany* **104**: 335–341.

Glover BJ, Perez-Rodriguez M, Martin C. **1998**. Development of several epidermal cell types can be specified by
the same MYB-related plant transcription factor. *Development* **125**: 3497–3508.

Goodwillie C, Kalisz S, Eckert CG. **2005**. The evolutionary enigma of mixed mating systems in plants:
occurrence, theoretical explanations, and empirical evidence. *Annual Review of Ecology, Evolution, and Systematics*
36: 47–79.

Goodwillie C, Ritland C, Ritland K. **2006**. The genetic basis of floral traits associated with mating system
evolution in Leptosiphon (Polemoniaceae): an analysis of quantitative trait loci. *Evolution* **60**: 491–504.

Grotewold E. **2006**. The genetics and biochemistry of floral pigments. *Annual Review of Plant Biology* **57**: 761–
780.

Hermann K, Klahre U, Moser M, Sheehan H, Mandel T, Kuhlemeier C. **2013**. Tight genetic linkage of
prezygotic barrier loci creates a multifunctional speciation island in Petunia. *Current Biology* **23**: 873–877.

Hiesey W, Nobs M, Bjorkman O. **1971**. *Experimental studies on the nature of species: 5. Biosystematics, genetics*
and physiological ecology of the Erythranthe section of Mimulus. Carnegie Institution.

Klahre U, Gurba A, Hermann K, Saxenhofer M, Bossolini E, Guerin PM, Kuhlemeier C. **2011**. Pollinator
choice in Petunia depends on two major genetic Loci for floral scent production. *Current Biology* **21**: 730–739.

Kostyun JL, Gibson MJS, King CM, Moyle LC. **2019**. A simple genetic architecture and low constraint allow
rapid floral evolution in a diverse and recently radiating plant genus. *New Phytologist* **223**: 1009–1022.

Krizek BA, Anderson JT. **2013**. Control of flower size. *Journal of Experimental Botany* **64**: 1427–1437.

Langfelder P, Horvath S. **2008**. WGCNA: an R package for weighted correlation network analysis. *BMC*
Bioinformatics **9**: 559.

Liang M, Chen W, LaFountain AM, Liu Y, Peng F, Xia R, Bradshaw HD, Yuan Y-W. **2023**. Taxon-specific,
phased siRNAs underlie a speciation locus in monkeyflowers. *Science* **379**: 576–582.

Liang M, Foster CE, Yuan Y-W. **2022**. Lost in translation: Molecular basis of reduced flower coloration in a self-
pollinated monkeyflower (Mimulus) species. *Science Advances* **8**: eabo1113.

Liao IT, Rifkin JL, Cao G, Rausher MD. **2021**. Modularity and selection of nectar traits in the evolution of the
selfing syndrome in Ipomoea lacunosa (Convolvulaceae). *New Phytologist* **233**: 1505–1519.

Lowry DB, Willis JH. **2010**. A widespread chromosomal inversion polymorphism contributes to a major life-
history transition, local adaptation, and reproductive isolation. *PLoS Biology* **8**: e1000500.

Nelson TC, Muir CD, Stathos AM, Vanderpool DD, Anderson K, Angert AL, Fishman L. **2021a**. Quantitative
trait locus mapping reveals an independent genetic basis for joint divergence in leaf function, life-history, and floral
traits between scarlet monkeyflower (Mimulus cardinalis) populations. *American Journal of Botany* **108**: 844–856.

Nelson TC, Stathos AM, Vanderpool DD, Finseth FR, Yuan Y-W, Fishman L. **2021b**. Ancient and recent
introgression shape the evolutionary history of pollinator adaptation and speciation in a model monkeyflower
radiation (Mimulus section Erythranthe). *PLoS Genetics* **17**: e1009095.

Newcombe FC. **1922**. Significance of the Behavior of Sensitive Stigmas. *American Journal of Botany* **9**: 99.

Ohno S. **1999**. Gene duplication and the uniqueness of vertebrate genomes circa 1970–1999. *Seminars in Cell &*
Developmental Biology **10**: 517–522.

Orr HA. **1998**. Testing natural selection vs. genetic drift in phenotypic evolution using quantitative trait locus data.
Genetics **149**: 2099–2104.

Peng F, Byers KJRP, Bradshaw HD. **2017**. Less is more: Independent loss-of-function OCIMENE SYNTHASE
alleles parallel pollination syndrome diversification in monkeyflowers (Mimulus). *American Journal of Botany* **104**:
1055–1059.

Ramsey J, Bradshaw HD, Schemske DW. **2003**. Components of reproductive isolation between the
monkeyflowers Mimulus lewisii and M. cardinalis (Phrymaceae). *Evolution* **57**: 1520–1534.

Rifkin JL, Cao G, Rausher MD. **2021**. Genetic architecture of divergence: the selfing syndrome in Ipomoea
lacunosa. *American Journal of Botany* **108**: 2038–2054.

Roels SAB, Kelly JK. **2011**. Rapid evolution caused by pollinator loss in Mimulus guttatus. *Evolution* **65**: 2541–
2552.

Romani F, Moreno JE. **2021**. Molecular mechanisms involved in functional macroevolution of plant transcription
factors. *New Phytologist* **230**: 1345–1353.

Schemske DW, Bradshaw HD. **1999**. Pollinator preference and the evolution of floral traits in monkeyflowers
(Mimulus). *Proceedings of the National Academy of Sciences of the United States of America* **96**: 11910–11915.

Sicard A, Lenhard M. **2011**. The selfing syndrome: a model for studying the genetic and evolutionary basis of
morphological adaptation in plants. *Annals of Botany* **107**: 1433–1443.

Smith SD. **2016**. Pleiotropy and the evolution of floral integration. *New Phytologist* **209**: 80–85.

Sotola VA, Berg CS, Samuli M, Chen H, Mantel SJ, Beardsley PA, Yuan Y-W, Sweigart AL, Fishman L.
2023. Genomic mechanisms and consequences of diverse postzygotic barriers between monkeyflower species.
Genetics **225**: iyad156.

Stankowski S, Chase MA, McIntosh H, Streisfeld MA. **2023**. Integrating top-down and bottom-up approaches to
understand the genetic architecture of speciation across a monkeyflower hybrid zone. *Molecular Ecology* **32**: 2041–
2054.

Stathos A, Fishman L. **2014**. Chromosomal rearrangements directly cause underdominant F1 pollen sterility in
Mimulus lewisii–Mimulus cardinalis hybrids. *Evolution* **68**: 3109–3119.

Stebbins GL. **1970**. Adaptive radiation of reproductive characteristics in angiosperms I: Pollination mechanisms.
Annual Review of Ecology and Systematics **1**: 307–326.

Stuurman J, Hoballah ME, Broger L, Moore J, Basten C, Kuhlemeier C. **2004**. Dissection of Floral Pollination
Syndromes in Petunia. *Genetics* **168**: 1585–1599.

Troth A, Puzey JR, Kim RS, Willis JH, Kelly JK. **2018**. Selective trade-offs maintain alleles underpinning
complex trait variation in plants. *Science* **361**: 475–478.

Valle JC del, Gallardo-López A, Buide ML, Whittall JB, Narbona E. **2018**. Digital photography provides a fast,
reliable, and noninvasive method to estimate anthocyanin pigment concentration in reproductive and vegetative
plant tissues. *Ecology and Evolution* **8**: 3064–3076.

Wang S, Basten CJ, Zeng Z-B. **2005**. *Windows QTL Cartographer 2.5*. Dept. of Statistics, North Carolina State
Univ.

Wessinger CA, Hileman LC. **2016**. Accessibility, constraint, and repetition in adaptive floral evolution.
Developmental Biology **419**: 175–183.

Wessinger CA, Hileman LC. **2020**. Parallelism in Flower Evolution and Development. *Annual Review of Ecology,*
Evolution, and Systematics **51**: 1–22.

Wessinger CA, Hileman LC, Rausher MD. **2014**. Identification of major quantitative trait loci underlying floral
pollination syndrome divergence in Penstemon. *Philosophical Transactions of the Royal Society B: Biological*
Sciences **369**: 20130349.

Wessinger CA, Katzer AM, Hime PM, Rausher MD, Kelly JK, Hileman LC. **2023**. A few essential genetic loci
distinguish Penstemon species with flowers adapted to pollination by bees or hummingbirds. *PLOS Biology* **21**:
e3002294.

Wessinger CA, Rausher MD. **2012**. Lessons from flower colour evolution on targets of selection. *Journal of*
Experimental Botany **63**: 5741–5749.

Wilson P, Wolfe AD, Armbruster WS, Thomson JD. **2007**. Constrained lability in floral evolution: counting
convergent origins of hummingbird pollination in Penstemon and Keckiella. *New Phytologist* **176**: 883–890.

Yeaman S, Whitlock MC. **2011**. The genetic architecture of adaptation under migration-selection balance.
Evolution **65**: 1897–1911.

Yuan Y-W. **2019**. Monkeyflowers (Mimulus): new model for plant developmental genetics and evo-devo. *New*
Phytologist **222**: 694–700.

Yuan Y-W, Rebocho AB, Sagawa JM, Stanley LE, Bradshaw HD. **2016**. Competition between anthocyanin and
flavonol biosynthesis produces spatial pattern variation of floral pigments between Mimulus species. *Proceedings of*
the National Academy of Sciences of the United States of America **113**: 2448–2453.

Yuan Y-W, Sagawa JM, Frost L, Vela JP, Bradshaw HD. **2014**. Transcriptional control of floral anthocyanin
pigmentation in monkeyflowers (Mimulus). *New Phytologist* **204**: 1013–1027.

Yuan Y-W, Sagawa JM, Stilio VSD, Bradshaw HD. **2013a**. Bulk segregant analysis of an induced floral mutant
identifies a MIXTA-like R2R3 MYB controlling nectar guide formation in Mimulus lewisii. *Genetics* **194**: 523–528.

Yuan Y-W, Sagawa JM, Young RC, Christensen BJ, Bradshaw HD. **2013b**. Genetic dissection of a major
anthocyanin QTL contributing to pollinator-mediated reproductive isolation between sister species of Mimulus.
Genetics **194**: 255–263.

747