*TSC2* **loss in neural progenitor cells suppresses**

**translation of ASD/NDD-associated transcripts**

**in an mTORC1- and MNK1/2-reversible fashion**

Pauline Martin,[1, 4] Krzysztof J. Szkop,[2, 4] Francis Robert,[3, 4] Srirupa Bhattacharyya,[1] Roberta

L. Beauchamp,[1] Jacob Brenner,[1] Nicholas E. Redmond[1], Sidong Huang,[3] Serkan Erdin,[1] Ola

Larsson,[2]* Vijaya Ramesh[1]*[§]

[1]Center for Genomic Medicine, Department of Neurology, Massachusetts General Hospital,

Boston, MA 02114, USA; [2]Department of Oncology-Pathology, Science for Life Laboratory,

Karolinska Institute, Stockholm 171 77, Sweden; [3]Department of Biochemistry and

Goodman Cancer Research Institute, McGill University, Montreal, PQ H3G1Y6, Canada;

[4]These authors contributed equally to this work.

* Co-corresponding senior authors: Ola Larsson (ola.larsson@ki.se), Vijaya Ramesh

(ramesh@helix.mgh.harvard.edu)

[§] **Lead contact:** Vijaya Ramesh, 185 Cambridge Street, Boston, MA 02114. Phone: 617-

724-9733; Email: ramesh@helix.mgh.harvard.edu

**SUMMARY**

Tuberous sclerosis complex (TSC), an inherited neurodevelopmental (ND) disorder with

frequent manifestations of epilepsy and autism spectrum disorder (ASD). TSC is caused by

mutations in *TSC1* or *TSC2* tumor suppressor genes, with encoded proteins hamartin/TSC1

and tuberin/TSC2 forming a functional complex inhibiting mechanistic target of rapamycin

complex-1 (mTORC1) signaling, leading to FDA-approved allosteric mTORC1-selective

rapamycin analogs for TSC tumors. Rapalogs are effective for TSC-associated hamartomas,

however, they are not effective for treating ND manifestations. mTORC1 signaling plays an

essential role in protein synthesis through mTORC1-eIF4F and MNK-eIF4E-mediated mRNA

translation. Further, the effects on mRNA translation by specific mTORC1 and MNK inhibitors

such as RMC-6272 and eFT-508 in TSC have never been explored. Here, employing

CRISPR-modified, isogenic TSC2 patient-derived neural progenitor cells (NPCs), we have

examined mRNA translation upon loss of *TSC2*. Our results reveal dysregulated translation in

*TSC2*-Null NPCs, which significantly overlap with the translatome from *TSC1*-Null NPCs,

which we reported recently. Most notably, numerous non-monogenic ASD- NDD- and

epilepsy-associated genes identified in patients harboring putative loss-of-function mutations,

including protein truncating, or damaging missense variants, were translationally suppressed

in *TSC2*-Null NPCs, and their translation were reversed upon RMC-6272 or eFT-508

treatment. Our study here establishes the importance of mTORC1-eIF4F and MNK-eIF4E-

mediated mRNA translation in TSC, ASD and other neurodevelopmental disorders and lay the

groundwork for evaluating drugs in clinical development that target these pathways as a

treatment strategy for TSC as well as ASD/NDD.

**INTRODUCTION**

Tuberous sclerosis complex (TSC [MIM: 191100 and MIM: 613254]) is an autosomal

dominant neurodevelopmental disorder showing clinical manifestations in multiple organ

systems including the central nervous system (CNS).  TSC is caused by mutations in the *TSC1*

[MIM: 605284] or *TSC2* [MIM: 191092] genes, encoding the tumor suppressor proteins

hamartin (TSC1) and tuberin (TSC2), respectively. In addition to slow growing hamartomas in

many organs, neurodevelopmental (ND) defects such as epilepsy, intellectual disability and

autism spectrum disorders (ASD) are commonly observed in these patients. [1-3]. TSC proteins

function as a complex relaying signal from diverse cellular pathways to control mechanistic

target of rapamycin (mTOR, [MIM: 601231]) complex 1 (mTORC1) signaling. Loss of

TSC1/TSC2 proteins results in aberrant activation of mTORC1, which has led to first-

generation mTORC1 inhibitors, rapamycin analogs ("rapalogs"), emerging as a lifelong

therapy for TSC hamartomas [4-7]. Moreover, recent clinical trials reveal that rapalogs reduce

epilepsy in 40% of TSC patients [8]. In contrast, rapalogs are ineffective in treating TSC-

associated neuropsychiatric disorders (TAND) and autism [9; 10].

Among its many activities, mTORC1 signaling plays an essential role in protein synthesis

by regulating the efficiency whereby mRNAs are translated into proteins. Aberrant activation

of mTORC1, as seen with TSC1/TSC2 loss, leads to hyperphosphorylation of key substrates

controlling translation including eukaryotic translation initiation factor (eIF) 4E-binding proteins

(4E-BPs [MIM: 602223 and MIM: 602224]) and ribosomal protein p70S6 kinase (S6K [MIM:

608938]). In the unphosphorylated state, i.e. when mTORC1 activity is low, 4E-BPs bind and

sequester eIF4E [MIM: 133440]. This leads to reduced formation of the heterotrimeric eIF4F-

complex, a key factor for initiation of translation, comprised of eIF4E, eIF4G [MIM: 600495]

and eIF4A [MIM: 602641]. Activation of mTORC1 leads to phosphorylation of 4E-BPs and

thereby release of eIF4E, which promotes eIF4F-complex formation and stimulates 5' cap-

dependent translation (Fig. 1A). Phosphorylation of S6K protein affects protein synthesis and

ribosomal biogenesis via multiple mechanisms, including release of sequestered eIF4A via

phosphorylation/inactivation of programmed cell death 4 (PDCD4 [MIM: 608610]) (Fig. 1A)

reviewed in [11; 12]. Importantly, mTORC1 modulates translation, and thereby molds the

proteome, in a transcript-selective fashion depending on features in the 5' untranslated regions

(UTRs) of target mRNAs. Three main classes of transcripts showing translation paralleling

mTORC1 activity have been described: mRNAs with very short 5'UTRs, highly structured

5'UTRs and those with the 5′ terminal oligopyrimidine (5′ TOP) motif (reviewed in [13]).

Moreover, activation of mTORC1 is associated with suppressed translation of a set of mRNAs

commonly harboring upstream open reading frames in their 5'UTRs [14]. In addition,

phosphorylation of eIF4E at a single conserved serine (S209) by mitogen-activated protein

kinase-interacting kinase MNK1/2 [MIM: 606724 and MIM: 605069] also modulates translation

in a transcript selective fashion (Fig. 1A) [15; 16].

Although commonly used rapalogs efficiently suppress phosphorylation of S6K1 they only

marginally affect 4E-BP1 phosphorylation in many cell types, which is consistent with rapalogs

only partially inhibiting mTORC1 activity [17-19]. This prompted development of bi-steric, third-

generation mTOR inhibitors that link the mTORC1 affinity of rapamycin with a potent second-

generation active-site mTOR kinase inhibitor, resulting in reduced phosphorylation of both

S6K1 and 4E-BP1 [20]. A more recent third-generation mTOR inhibitor, RMC-6272 (a tool

compound for its clinical counterpart RMC-5552), shows dramatically improved mTORC1

selectivity while still efficiently reducing phosphorylation of both S6K1 and 4E-BP1  (Fig. 1A)

{Burnett, 2023 #1356;Lee, 2021 #1418}. Moreover, we demonstrated that RMC-6272 is

superior to rapamycin in durable inhibition the mTORC1-4E-BP1 axis and suppression of

tumor growth in patient-derived *NF2*-deficient models [21]. Furthermore, RMC-6272 has shown

strong anti-tumor activity either alone or when combined with other treatments in several

preclinical cancer models {Burnett, 2023 #1356;Lee, 2021 #1418}, and clinical trials using

RMC-5552 have offered preliminary data of anti-tumor activity at tolerated doses

(NCT04774952 and NCT05557292; [22]. Similarly, recently developed eFT-508 is a highly

potent and selective MNK1/2 inhibitor that effectively inhibits phosphorylation of eIF4E (Fig.

1A) [23] and is in phase 2 clinical trials for several cancer types (NCT02937675, NCT02605083

and NCT03258398).

While numerous global studies of mRNA levels ("transcriptomes") have been carried out

in patient-derived neuronal models of ASD [MIM 209850], TSC and other ND disorders,

studies of how mRNAs are translation is largely lacking. Furthermore, the effects on mRNA

translation from highly specific mTORC1 and MNK inhibitors such as RMC-6272 and eFT-508

in ND disorders have never been explored. Our recent studies employing polysome-profiling

to examine mRNA translation at a transcriptome-wide level in TSC1 patient-derived neural

progenitor cells (NPCs) uncovered *TSC1*-associated transcript-selective alterations in

translation. Importantly, translation of a subset of mRNAs encoding proteins related to synaptic

regulation or ASD were rapamycin-insensitive but showed RMC-6272-sensitive translation.

Interestingly, unlike rapamycin, RMC-6272 also reversed some of the early ND phenotypic

defects in *TSC1*-Null NPCs [24]. These findings are consistent with a key role of the mTORC1-

4E-BP1-eIF4E axis. Here, we determine the role of this axis in regulating mRNA translation

using CRISPR-modified, isogenic TSC2 patient-derived NPCs. Our results reveal a strong

overlap in dysregulated translation observed in *TSC1*-Null and *TSC2*-Null NPCs. Surprisingly,

genes previously associated with ASD and NDD were translationally suppressed in *TSC2*-Null

NPCs, and their translation was reversed upon RMC-6272 or eFT-508 treatment. These

results establish the importance of mTORC1-eIF4F and MNK-eIF4E-mediated mRNA

translation in TSC, ASD and ND disorders and lay the groundwork for evaluating drugs in

clinical development that target these pathways as a treatment strategy for TAND as well as

ASD/NDD.

**MATERIAL AND METHODS**

***Human induced pluripotent stem cell derivation***

We previously reported collection of post-mortem subungal fibroma (SUF) tissue from a

female sporadic TSC2 patient, which was obtained through the pathology department and

approved by the Institutional Review Board at Massachusetts General Hospital. Detailed

genetic analysis of the patient SUF tissue, including exon-by-exon scanning of the entire

coding sequence of *TSC2* isoform 1 (NCBI accession NM_000548), identified a single

heterozygous (*TSC2*-Het) truncating nonsense mutation in *TSC2* exon 34 (originally reported

as coding exon 33; 4207delG, D1400fs>1410X). Further analysis of the SUF tissue for tumor

clonality and loss-of-heterozygosity (LOH) at multiple polymorphic chromosome 16 or *TSC2*

intragenic markers revealed a clonal tumor with no second-hit mutations or LOH [25]. For

culturing of primary *TSC2*-Het fibroblasts, SUF tissue was rinsed twice in Hank's Balanced

Salt Solution (Sigma) followed by mincing in high glucose DMEM (Gibco) containing 15% fetal

bovine serum (FBS; Sigma), 0.5 µg/ml fungizone (Life Technologies), 50 g/ml gentamicin (Life

Technologies), and 1x penicillin-streptomycin/L-glutamine (Cellgro). The tissue was then

digested in the same medium as above (without gentamycin) containing ~5 units collagenase

III (Fisher Scientific), for 30 min at $37^0$C. Digested tissue was then pelleted by centrifugation,

resuspended in high glucose DMEM (containing 15%FBS, 1x penicillin-streptomycin/L-

glutamine) and expanded for subsequent experiments. *TSC2*-Het cells were reprogrammed

using synthetic modified mRNA-based methods to obtain induced pluripotent stem cells

(iPSCs) as described [26; 27]. Briefly, cells were transfected by nucleofection (Amaxa

Nucleofector I) with *in vitro* transcribed mRNAs encoding *Oct4, Sox2, Klf4, c-Myc* and *Lin28*

(Stemgent). After single clone picking, iPSC colonies were cultured in feeder-free conditions

in Essential 8 medium (ThermoFisher) on plates coated with Geltrex (Gibco) using the "thin

gel method" according to the manufacturer's instructions. Colonies were passaged every 4-6

days with daily medium change.

***CRISPR/Cas9-mediated generation of isogenic iPSC lines***

To generate both the corrected-wildtype (WT) and Null isogenic iPSC lines, CRISPR/Cas9

genome editing was performed employing *TSC2*-Het patient-derived iPSCs. To achieve gene

knock-in to correct the germline *TSC2* mutation, we followed the same cloning and transfection

strategy as we previously reported for *TSC1* corrected-WT iPSCs [28]. Briefly, using web-based

CRISPR-related tools developed by GenScript (https://www.genscript.com), we designed a

single guide RNA (sgRNA): 5'-GGACATCCTCGGGGACCCTG-3' targeting *TSC2* exon 34

(coding exon 33) along with an asymmetric donor sequence: 5'-

CCGAGGCCGACCAGGCAGCACTTTCCCCGTCCAGGGTCCCTGACTGTGACCGGGCCT

TAACCTCAGGGCTCAGCCGGCCCACGTCGGCCTTGTCCCCAGGGTCCCCGAGGATGT

CCTGCAGAGTCTGCAG-3' for homology-directed repair. Colonies were manually isolated,

and genomic DNA was extracted from a portion of each colony followed by PCR amplification

using primers TSC2-ex33F: 5'-CTGACAGGGGTTCTCTTTGG-3' and TSC2-ex33R: 5'-

TCCAGGGTCCCTGACTGTGA-3', and Sanger sequencing was performed using a nested

primer TSC2-ex33-seqF: 5'-TCGTCCTCAGTCTCCAGCC-3'. Screening by Sanger

sequencing identified one clone with successful correction back to WT. For generating the

*TSC2*-Null iPSC lines, we used a service provided by an independent company (Alstem Cell

Advancements). Briefly, *TSC2*-Het iPSCs were co-transfected using the Amaxa nucleo-

electroporation system with a pair of sgRNAs targeting *TSC2* exon 2 (coding exon 1) including

sgRNA-1: 5'-TGTTGGGACTGGGAACACCG-3' and sgRNA-2: 5'-

ACTCACTCTCAGTATTTCCG-3'. A total of 50 single clones were screened by PCR

amplification and Sanger sequencing using primers TSC2-F2: 5'-

CCATGGCCAAACCAACAAG-3' and TSC2-R2: 5'-ACCTGGTGCAAGACCAAA-3', with 3

homozygous *TSC2*-Null cell lines identified, all of which harbored the same 59 bp deletion in

*TSC2* exon 2 resulting in a frameshift/premature truncating mutation.

***Differentiation of iPSCs into NPCs***

To generate neural progenitor cells (NPCs), *TSC2*-iPSCs, including the parental *TSC2*-

Het line, the single *TSC2*-WT clone and two of the three *TSC2*-Null clones (clones 5 and 39)

were differentiated using the directed monolayer differentiation protocol as we previously

reported for TSC1-NPCs [28; 29]. Briefly, iPSCs expressing the pluripotency marker TRA-1-60

were sorted and enriched using the MACS MicroBead cell sorting technology (Miltenyi Biotec).

Cells were cultured in neural induction medium (neurobasal medium supplemented with 1x

neural induction supplement (ThermoFisher)) for 7-9 days, followed by a second sorting for

cells expressing polysialylated-neural cell adhesion molecule (PSA-NCAM). The PSC-NCAM-

positive (+) cells then underwent a double-sorting to enrich for NPCs representing CD271(-)

/CD133(+) cells. NPCs were then cultured in neural expansion medium (1:1 neurobasal

medium:advanced DMEM/F12, 1x penicillin/streptomycin and 1x neural induction

supplements (ThermoFisher)). MAC sorting was performed using microbead kits for TRA-1-

60, PSA-NCAM, CD271 and CD133 (Miltenyi Biotec).

***Immunocytochemistry***

To confirm successful derivation of iPSCs and NPCs, immunofluorescent staining was

carried out using antibodies for iPSC markers Oct4 (Santa Cruz Biotechnologies) and Nanog

(Abcam), or NPC markers Nestin (EMD Millipore) and Sox2 (R&D Systems) as previously

described [28; 29]. Briefly, cells were fixed with 4% PFA and then permeabilized and blocked in

a single step using PBS containing 4% normal goat serum (NGS)/0.1% Triton-X-100. Primary

antibodies were diluted into PBS containing 2% NGS/0.1% Triton-X-100. DAPI was used to

stain nuclei (Invitrogen) at 5 $\mu$g/ml. Coverslips were mounted in ProLong Gold antifade

Mountant (Invitrogen) and images were captured using a Nikon Eclipse TE2000-U microscope

and the NIS-Element BR 3.2 imaging software.

***Immunoblot analysis***

Cells were lysed in RIPA lysis buffer (50 mM Tris (pH7.5) and 150 mM NaCl, 0.1% SDS,

0.5% sodium deoxycholate, 1% NP-40, 1x HALT phosphatase inhibitor cocktail (Pierce) and

1x protease inhibitor cocktail (Sigma)) as previously described [30; 31]. Protein lysates were

resolved on 4-12% or 10-20% Novex WedgeWell Tris-glycine gels (Invitrogen), transferred to

0.45 µm pore nitrocellulose (BioRad) and then incubated with primary antibodies to detect

TSC2, pS6K T389, S6K, pS6 S240/244, S6, p4E-BP1 S65, 4E-BP1, peIF4E S209, and eIF4E

(Cell Signaling Technologies), $\beta$-actin (Santa Cruz Biotechnology) or GAPDH (EMD Millipore).

All immunoblotting data shown are representative of at least 3 biological replicates.

***Cell Size and proliferation***

Cell size and proliferation rate were determined for *TSC2*-WT, Het and Null NPCs treated

for 72 h with 50 nM rapamycin, 10 nM RMC-6272 or DMSO as vehicle control. For both cell

size and proliferation, NPCs were analyzed by trypan blue exclusion using the Countess II

automated cell counter (ThermoFisher) as previously described [24; 28]. Briefly, cells were

detached using Accutase enzyme detachment medium (ThermoFisher), and for each sample,

10 μl of suspended cells were combined with 10 μl of trypan blue, and 10 μl of the resulting

mixture was assessed using the Countess II automated cell counter (ThermoFisher),

according to the manufacturer's instructions. Quantification of viable and dead cells was

performed along with size of viable cells. For proliferation, counting of viable cells collected at

days (D) 1, 2, 3 and 4 after seeding was performed for cells treated with 50 nM rapamycin, 10

nM RMC-6272 or DMSO on D1. All data collected using the Countess II cell counter were

analyzed from 3 independent experiments (3 technical replicates each). Proliferation was

further assessed using flow cytometry employing the proliferation marker Ki-67 as previously

described [24]. Briefly, *TSC2*-WT, Het and Null NPCs were seeded onto Geltrex-coated plates

followed by treatment with 50 nM rapamycin, 10 nM RMC-6272, 50 nM eFT-508 or DMSO as

vehicle control for 72 h (two biological replicates). To assess the Ki-67 proliferation status,

NPCs were fixed with 4% PFA, permeabilized with ice cold 100% methanol to a final

concentration of 90% with gentle vortexing and stained using AlexaFluor488-conjugated Ki-

67 monoclonal antibody or AlexFluor488-conjugated IgG as an isotype control (Cell Signaling

Technologies). Data were acquired from ~$1\times10^4$ cells using a BD LSR II Flow Cytometer (BD

Biosciences), and analysis was carried out using FlowJo 10.8.1 (FlowJo LLC). Percentages

of Ki-67 positive cells (acquired through FITC channel) were determined by gating with respect

to Alexa488-IgG control.

***Neurite outgrowth assay***

NPCs were assessed for neurite outgrowth as previously described [28]. Briefly, NPCs were

seeded on coverslips coated with poly-D-lysine (0.1 mg/ml, Sigma) and fibronectin (5 $\mu$g/ml,

Corning) in growth factor-depleted neural expansion medium containing 1:1 neurobasal

medium:advanced DMEM/F12 (ThermoFisher), 1x penicillin/streptomycin and 0.3x neural

induction supplements (ThermoFisher). Cells were treated with 50 nM rapamycin, 10 nM of

RMC-6272 or DMSO vehicle control for 48 h followed by fixation with 4% PFA prior to

immunostaining. Three independent experiments were performed where images were

acquired from 8 non-overlapping fields and analyzed using HCA-Vision software V.2.2.0

(CSIRO, Canberra, Australia) to determine (i) number of cells; (ii) average neurite

outgrowth/cell (total length), (iii) average number of neurites/cell and (iv) average number of

extremities/cell (number of termination points) [32]. Images were acquired using a Nikon Eclipse

TE2000-U microscope and the NIS-Element BR 3.2 imaging software.

***Polysome fractionation and RNA sequencing***

Lysate were prepared for polysome profiling from three biological replicates as previously

described with minor modifications [33]. Briefly, TSC2-NPC lines were seeded at ~50,000

cells/cm$^2$, with a total of ~$2.4 \times 10^7$ cells seeded per replicate for three drug treatment

conditions, including 10 nM RMC-6272, 50 nM EFT-508 or DMSO control. The day after

seeding, cells were treated for 2 h, rinsed with 1x PBS containing 100 µg/ml cycloheximide

(CHX) (Sigma), harvested by scraping on ice in PBS/CHX and pelleted by centrifugation at

300g for 10 min at + 4C. Cell pellets were first resuspended in hypotonic lysis buffer containing

5 mM Tris-HCl (pH 7.5), 2.5 mM MgCl$_2$ and 1.5 mM KCl, followed by addition of DTT (2 mM

final), CHX (100µg/ml final), sodium deoxycholate (0.5% final) and Triton X-100 (0.5% final).

Lysates were then vortexed for 4 sec, cleared by spinning for 2 min at 13,000 rpm and quickly

frozen on dry ice. When ready for processing, lysates were quickly thawed at 37C and loaded

onto a 10–50% sucrose gradient, followed by centrifugation for 2 h and 15 min at 35,000 rpm

in a SW41 rotor using a Sorvall Discovery 90SE centrifuge. The gradients were fractionated

on a Teledyne ISCO Foxy R1 apparatus while monitoring the OD254nm. Fractions

corresponding to mRNA associated with more than two ribosomes were pooled and the RNA

extracted using TRIzol (ThermoFisher) according to the manufacturer's protocol. 10% of the

lysate was removed from the extract prior to loading on the sucrose gradient, extracted using

TRIzol, and denoted as total RNA. RNA sequencing libraries were prepared from the resulting

samples (both polysome-associated and total RNA) using Illumina v2.5 Kits and sequenced

on an Illumina NextSeq 500 at the Canada's Michael Smith Genome Sciences Centre (BC

Cancer Research Institute, Vancouver, Canada).

***RNA-seq data preprocessing and quality control***

Adapters were removed from RNA sequencing reads using bbduk.sh from the BBTools

software suite [https://www.osti.gov/servlets/purl/1241166] (v. 36.59 with settings: k = 13, ktrim

= n, useshortkmers = t, mink = 5, qtrim = t, trimq = 10, minlength = 20). Paired-end reads were

aligned to hg38 using HISAT2 [34] (ver 2.1.0 with settings: --no-mixed, --no-discordant), and

uniquely mapped reads were quantified using the featureCounts function from the RSubreads

package (v.2.6.4) [35] with RefSeq gene definitions and default settings. Polysome-profiling data

from our previous study of TSC1-dependent translation[24] was quantified using the same gene

definitions to allow a direct comparison to the herein generated dataset. Dataset reproducibility

was assessed using Principal Component Analysis using genes within highest quartile of

standard deviations quantified across all samples as input.

***Analysis of gene expression alterations using anota2seq***

Gene expression quantified from polysome-associated and total mRNA were analyzed using

the anota2seq algorithm (v. 1.20.0) [36]. Batch effects were accounted for in the models by

including sample replicates using the "batchVec" argument in anota2seq. Default threshold

settings were used [settings minSlopeTranslation = - 1, maxSlopeTranslation = 2,

minSlopeBuffering = -2, maxSlopeBuffering = 1, deltaPT = deltaP = deltaTP = deltaT =

log2(1.2), minEff=log2(1.5)] with a significance threshold maxRvmPAdj = 0.15 (corresponding

to an FDR < 0.15). Transcripts were classified into three modes of regulation assessed by

anota2seq (changes in mRNA abundance, changes in translation, and translational offsetting)

using the "anota2seqRegModes' function.

***Analysis of gene signatures using empirical cumulative distribution functions***

Empirical cumulative distribution functions (ECDFs) of log2 fold changes for polysome-

associated and total mRNA were plotted for gene signatures (ASD and NDD genes [37], eIF4E

overexpression [38], mTOR sensitive genes [39] obtained from previous publications or to cross-

compare datasets that were found to be translationally regulated in either dataset. The

Wilcoxon rank-sum test was used to determine whether there was a significant shift between

each signature and the background (i.e. genes not within the signature).

***Analysis of enrichment of gene sets***

Gene Ontology (GO) enrichment analysis was performed using genes classified as

regulated via translation by anota2seq (differentially translated genes "DTGs") and annotation

from GO biological process (GO:BP) terms retrieved from MSigDB database (v. 2023.2.Hs) [40]

using one-tailed Fisher's exact test. In this analysis, only GO:BP terms with at least 10 genes

within in the background set (all detected genes analyzed by anoat2seq) of a particular

comparison were considered.  DTG datasets for GO analysis included *TSC2*-Null vs WT

NPCs, *TSC2*-Null NPCs treated with RMC-6272 vs DMSO control, *TSC2*-Null NPCs treated

with eFT508 vs DMSO control, pairwise comparison between *TSC2*-Null NPCs treated with

RMC-6272 or eFT508 as well as pairwise comparison between *TSC2*-Null vs WT NPCs and

our previously published dataset for *TSC1*-Null vs WT NPCs [24].

Enrichment analysis  for selected DTGs identified by anota2seq and target gene sets,

including ASD-, NDD-, EPI-associated genes [37; 40; 41] and eIF4E-sensitive genes [38], was

performed using one-tailed Fisher's exact test. Statistical significance of overlaps between the

two DTG datasets was also assessed using one-tailed Fisher's exact test. Here, all anota2seq

analyzed genes in a single comparison, or intersection of anoat2seq analyzed genes in paired

comparisons, constituted the background sets. These enrichment analyses were performed

in the R statistical software environment (version 3.4.3) (https://www.R-project.org/) using

custom scripts.

***Validation of differential translation using RT-qPCR***

For validation, polysome fractionations were collected from *TSC2*-Null or *TSC2*-WT NPCs

(three biological replicates), and RNA was isolated as described for NPCs above. cDNA was

prepared using M-MuLV Reverse Transcriptase (New England Biolabs, Ipswich, MA) and

oligo(dT)20 primers using the manufacturer's recommendations. RT-qPCRs were performed

with SsoFast EvaGreen Supermix (Bio-Rad, Hercules, CA) using the CFX96 PCR system (Bio-

Rad Hercules, CA). Primers for RT-qPCR are detailed in Supplemental Table S1. The level of

each mRNA was normalized to β-actin (*ACTB*) and Phosphoglycerate Kinase 1 (*PGK1*) using

the comparative CT method and compared across conditions as indicated in the figure legend.

***Statistical analyses***

Methods for statistics are described above, or in corresponding figure legends.

**RESULTS**

***Generation and characterization of an isogenic set of TSC2 patient-derived neural***

***progenitor* cells**

We generated an isogenic set of TSC2 patient-derived NPCs using a similar approach as

we described previously for TSC1 patient-derived NPCs [24; 28]. Primary fibroblasts cultured from

post-mortem SUF tissue were found to harbor a germline heterozygous 1 bp deletion in *TSC2*

exon 34 (originally reported as coding exon 33) with no loss of heterozygosity [25]. Induced

pluripotent stem cells (iPSCs) were generated from the fibroblasts, followed by CRISPR-

based genome editing to (i) correct the heterozygous mutation back to wildtype, or (ii) knock

out the second allele by introducing a 59 bp deletion in *TSC2* exon 2 (coding exon 1), both of

which were confirmed by Sanger sequencing (data not shown). This resulted in an isogenic

set of iPSCs representing the parental *TSC2*-Het line, a single clone for *TSC2*-corrected

(*TSC2*-WT) and three clones for *TSC2*-Null iPSCs. We then selected *TSC2*-WT, Het and two

iPSC clones (clones 5 and 39) for microbead sorting of neural cell surface protein markers to

obtain NPCs as previously described [28]. Immunostaining for neural markers Nestin and Sox2

confirmed successful NPC derivation (Supplemental Fig. S1A). Loss of TSC2 protein

expression was confirmed in *TSC2*-Null clones when compared with the TSC2-WT. We

observed increased phosphorylation of mTORC1 targets p70S6K (pS6K T389), pS6 S240/244

and p4E-BP1 S65 in both *TSC2*-Null NPC clones compared with *TSC2*-WT NPCs, consistent

with activation of mTORC1 signaling. *TSC2*-Het NPCs revealed only minimal activation of

mTORC1, which is in agreement with many previous reports that employed human iPSC

models of TSC [42-45] (Supplemental Fig. S1B). *TSC2*-Null NPCs demonstrated an enlarged cell

phenotype compared with *TSC2*-WT cells (Fig. 1B, D), which is a hallmark of mTORC1

activation. *TSC2*-Het NPCS exhibited an intermediate cell size phenotype (Fig. 1B, D).

Treatment with rapamycin or RMC-6272 attenuated phosphorylation of S6K at T389, along

with its downstream substrate ribosomal protein S6 (pS6 S240/244) and rescued the enlarged

cell size (Fig. 1C-D). As expected, rapamycin showed only minimal effects on phosphorylation

of 4E-BP1 while RMC-6272 demonstrated complete inhibition (Fig. 1C). These results are

consistent with our observations in TSC1-NPCs [24] and  with previous studies showing that the

mTORC1-S6K signaling axis controls cell size [46].

***Rescue of early neurodevelopmental phenotypes by RMC-6272, but not rapamycin***

We previously demonstrated that defects in early ND phenotypes in *TSC1*-Null NPCs,

including increases in proliferation rate and neurite outgrowth, could be rescued by RMC-6272

treatment but not rapamycin. This suggests dependence, at least in part, on the mTORC1-4E-

BP1 signaling axis [24; 28]. Here, we assessed proliferation by viable cell count in TSC2-NPCs

over the course of four days (D1-4) after seeding, along with 72 h treatment with 50 nM

rapamycin, 10 nM RMC-6272 or DMSO control initiated at D1. We observed genotype-

dependent cell numbers whereby *TSC2*-Null NPCs showed increased proliferation relative to

*TSC2*-Het and *TSC2*-WT NPCs while *TSC2*-Het had augmented proliferation as compared to

*TSC2*-WT NPCs (Fig. 2A, left graph region). When compared with DMSO treatment,

rapamycin showed no significant effect in TSC2-NPCs (Fig. 2A, center); whereas RMC-6272

demonstrated significant reduction by D3 in all three NPC genotypes, with the maximal

decrease observed in *TSC2*-Null NPCs (Fig. 2A, compare left and right graph regions).

Consistently, flow cytometry-based analysis showed a reduction in the percentage of S-

positive cells in RMC-6272 vs DMSO treated *TSC2*-WT (50.9% vs 62.7%), *TSC2*-Het (56.2%

vs 64.9%) and *TSC2*-Null (53.7% vs 68.9%) NPCs. In contrast, similar percentages of Ki-67

positivity were found for rapamycin or DMSO treatment (Supplemental Fig. S2A). Moreover,

treatment with eFT508/tomivosertib, a MNK1/2 inhibitor, which attenuates phosphorylation of

eIF4E at S209 (peIF4E S209) but not 4E-BP1 (p4E-BP1 S65) did not alter Ki-67 positivity in

*TSC2*-Null NPCs (Supplemental Fig. S2A and B).

We next studied neurites in our isogenic set of TSC2-NPCs using MAP2 immunostaining

and HCA-Vision image analysis software to, per cell, quantitate three parameters: (i) length of

neurite outgrowth, (ii) number of neurite roots and (iii) number of extremities (terminal endpoint

of primary and branched outgrowths). In TSC2-NPCs treated for 48 h with DMSO, a significant

increase in neurite length of ~3.2-fold in *TSC2*-Null and an intermediate increase of ~1.4-fold

in *TSC2*-Het was observed relative to *TSC2*-WT NPCs (Fig. 2B-C). *TSC2*-Null NPCs also

revealed a significant increase in neurite number (~1.5-fold) and extremities (~1.7-fold)

relative to *TSC2*-WT NPCs (Fig. 2B and C). Upon RMC-6272 treatment, we observed

decreases in neurite length, number and extremities of 0.73, 0.90 and 0.83, respectively, for

*TSC2*-Het NPCs, and 0.46, 0.68 and 0.64, respectively, for *TSC2*-Null NPCs (Fig. 2D). Similar

to the proliferation assays, rapamycin failed to rescue neurite-related phenotypes, and in fact

showed an opposite trend in several parameters (Fig. 2D). Taken together, these data support

that the third-generation mTORC1 inhibitor RMC-6272 is superior to rapamycin in reversing

the early ND phenotypes proliferation and neurite phenotypes in NPCs and that this depends

on the more potent inhibition of mTORC1 downstream targets including 4E-BP1.

***Abundant mRNA translation alteration in TSC2-Null NPCs largely overlap with those***

***observed in TSC1-Null NPCs***

Our findings above demonstrate activation of mTORC1 signaling and increases in

proliferation and neurite outgrowth in *TSC2*-Null NPCs, compared with WT cells. As mRNA

translation is a major downstream target of mTORC1, it is necessary to study how both mRNA

levels and translation are affected by loss of *TSC2* to disentangle gene expression-dependent

effects on cellular phenotypes. We therefore applied polysome-profiling in *TSC2*-Null and

*TSC2*-WT NPCs. During polysome-profiling, cytoplasmic mRNA is fractionated based on

ribosome-association and the resulting polysome-associated (mRNA associated with >2

ribosomes) and total cytoplasmic mRNA pools are then quantified using RNA sequencing

followed by anota2seq analysis [36; 47]. This allows identification of *TSC2*-associated alterations

in (i) "translation", defined as polysome-associated mRNA alterations not paralleled by

corresponding changes in total mRNA levels; (ii) "abundance", representing congruent

modulation of total and polysome-associated mRNA; and (iii) "offsetting", defined as

alterations in total mRNA not paralleled by corresponding changes in polysome-associated

mRNA. This analysis revealed numerous *TSC2*-associated alterations in mRNA translation

visualized using a scatterplot comparing *TSC2*-dependent changes in polysome-associated

and total mRNA (Fig. 3A, Supplemental Table S4) and densities of p-values and FDRs

obtained from anota2seq analysis (Fig. 3B). These alterations in translation occurred

alongside less abundant changes in mRNA levels and essentially in absence of translational

offsetting (Fig. 3A). Although the analysis is expected to be hampered by the large number of

regulated genes, we used gene ontology analysis to identify cellular functions targeted by

these changes in translation. This revealed that translationally suppressed transcripts encode proteins enriched for annotation to cellular functions including protein DNA complex organization, chromatin remodeling, RNA metabolism, and cell cycle while those involved in a variety of metabolic processed were translationally activated (Fig. 3C).

To assess how these *TSC2*-associated alterations in mRNA translation compares to our previously described *TSC1*-associated changes, we looked for differences in the response to loss of *TSC1* vs *TSC2* relative to their respective control conditions (i.e. the interaction effect) using an anota2seq-like approaches we described previously [48]. Strikingly, only one gene showed a statistically significant difference in translation response upon *TSC1* vs *TSC2* loss (at FDR < 0.15), and this analysis therefore argues for largely overlapping effects on mRNA translation. To further assess this, we generated gene signatures of transcripts that were translationally up or downregulated in *TSC1*-Null or *TSC2*-Null NPCs relative to their controls and visualized how these gene sets were regulated across the two models. This revealed that transcripts whose translation was activated or suppressed in *TSC2*-Null NPCs showed similar regulation in *TSC1*-Null NPCs and, vice versa, transcripts whose translation were activated or suppressed in *TSC1*-Null NPCs showed similar regulation in *TSC2*-Null NPCs (Fig. 3D). Yet, there were differences in mRNA level changes for these subsets (compare Fig. 3D left and right plots) that, together with expected threshold-effects, resulted in a smaller set of 949 translationally suppressed and 189 translationally activated transcripts passed threshold applied in anota2seq for both comparisons (Fig. 3E). As expected, these shared genes were enriched in similar gene ontology categories as those identified upon *TSC2* loss (Fig. 3F). In aggregate, these results indicated extensive remodeling of mRNA translation in *TSC2*-Null NPCs largely recapitulating that observed in *TSC1*-Null NPCs.

***Suppressed translation of mRNAs encoding proteins associated with ASD and NDD upon* TSC2 *loss***

To explore the relationship between altered gene expression associated with *TSC2* loss

and patient phenotypes we reasoned that *TSC2* loss may affect translation of mRNAs

encoding proteins implicated in non-syndromic autism spectrum disorder (ASD), general

neurodevelopmental disorders (NDD) and/or epilepsy.  To assess this, we utilized data from

recent studies that identified high confidence *de novo* mutations in large cohorts of individuals

diagnosed with ASD, NDD as well as epilepsy. For ASD/NDD-related genes, we used datasets

reported by Fu and colleagues, including 72 genes associated with ASD (ASD72) and 373

genes associated with NDD (NDD373) identified at a stringent FDR $\leq$ 0.001 threshold, as well

as 185 ASD-associated genes (ASD185) and 664 NDD-associated genes (NDD664) identified

at reduced stringency (FDR $\leq$ 0.05) [37]. For epilepsy-related genes, we used genes identified

in the Epi25 Collaborative, [41] that consists of 83 genes identified at p $\leq$ 0.005 by whole-exome

sequencing (case-control study) in the full epilepsy cohort (EPI). We  next assessed whether

expression of ASD-, NDD- and epilepsy-associated genes are regulated in *TSC2*-Null NPCs

as determined by anota2seq analysis (i.e. from Fig. 3A). This analysis revealed strong

enrichment of ASD72 (~43% of all ASD72 genes; p = $1.72 \times 10^{-7}$) and NDD373 (~37% of all

NDD273 genes; p = $2.59 \times 10^{-19}$) genes among translationally suppressed transcripts as

compared to background (Table 1). This finding was further substantiated by applying two

approaches that do not use thresholds to identify sets of differentially regulated genes: i) gene

set enrichment analysis (GSEA) [40] (Fig. 4A); and ii) the same method as applied for *TSC1* vs

*TSC2* (i.e. Fig. 3D) gene subset-comparisons (Fig. 4B, top). Both these approaches showed

strong enrichment of ASD72 and NDD373 genes among transcripts translationally suppressed

in *TSC2*-Null as compared to isogenic WT NPCs. Similarly, in *TSC2*-Null NPCs, strong

enrichments among translationally suppressed genes were observed for the larger ASD185

and NDD664 gene sets (Supplemental Table S2A, Supplemental Fig. S3A and B) as well as

genes translationally suppressed in both *TSC2*-Null and *TSC1*-Null NPCs relative to their

respective controls (i.e. genes from Fig. 3E; Supplemental Table S2B). A similar analysis for

EPI-associated genes revealed a smaller enrichment (~27% of all EPI-associated genes; p =

5.82x10$^{-3}$) (Table 1, Supplemental Fig. S3C and D) that was not confirmed using methods that

do not use thresholds to identify regulated sets of genes.

To ensure that this unexpected translational suppression of ASD- and NDD-associated

genes in NPCs occurs in the presence of expected effects on mRNA translation from mTORC1

activation, we applied the same approaches described above to sets of transcripts whose

translation was altered following overexpression of eIF4E [38]. Consistent with activation of the

mTORC1-eIF4F translational axis in *TSC2*-Null NPCs, there was a strong overlap between

genes whose translation was activated upon eIF4E-overexpression and *TSC2* loss

(Supplemental Table S3). A similar result was obtained when only considering genes that

were identified as translationally regulated by anota2seq following both *TSC1* and *TSC2* loss

(i.e. from Fig. 3E) and when a method not relying on thresholds was used (Fig. 4B, middle) as

well as for an additional signature of mTORC1-sensitive translation reported previously [14](Fig.

4B, bottom). Therefore, surprisingly, genes associated with ASD and NDD are largely

translationally suppressed in NPCs following inactivation of *TSC2* and these changes in

translation occur in parallel with activation of mTORC1-sensitive translation.

***5'UTRs of ASD and NDD genes resemble those of mRNAs whose translation is***

***suppressed upon activation of mTORC1***

The above results unveiled that translation of mRNAs encoding proteins associated with

ASD and NDD are affected by the activity of mTORC1. As mTORC1-sensitive translation

largely relies on features in the 5' untranslated region (5'UTR), this suggests that the observed

suppressed translation of ASD- and NDD-associated genes may depend on their 5'UTRs. As

discussed above, mRNAs whose translation parallels activity of mTORC1 often have short

and/or highly structured 5'UTRs and lack upstream open reading frames (uORFs). Strikingly,

ASD- and NDD-associated genes showed an opposite profile for 5'UTR length, GC-content

(which associates with secondary structures of the 5'UTR) and uORFs that was similar to

mRNAs whose translation was suppressed upon *TSC2* loss (Fig. 4C-E). Therefore, ASD- and

NDD-associated genes largely show 5'UTRs that are consistent with translational suppression

following increased activation of mTORC1.

***RMC-6272 and eFT-508 reverse translationally suppressed* TSC2-*dependent***

***transcripts and ASD and NDD genes***

The findings above suggest that suppressed translation of ASD- and NDD-associated

genes following loss of *TSC2* may be important for ND manifestation. Furthermore, this raises

the possibility that inhibition of mTORC1 signaling, or potentially other key pathways

modulating translation, may reverse *TSC2*-dependent translation of ASD- and NDD-

associated genes. To assess this, we performed polysome-profiling in *TSC2*-Null NPCs

treated with the bi-valent mTORC1 inhibitor RMC-6272, or the MNK1/2 inhibitor eFT-508. The

resulting dataset was analyzed as described above using anota2seq to determine how these

agents modulate gene expression. Both treatments resulted in dramatic mRNA translation

changes as visualized in scatterplots comparing effects in polysome-associated and total

mRNA levels as well as p-value and FDR densities (Supplemental Fig. S4A-D). Next, we

generated gene signatures of transcripts that were translationally increased or decreased in

*TSC2*-Null NPCs treated with RMC-6272 relative to DMSO control and compared how these

gene sets were regulated in *TSC2*-Null NPCs treated with eFT-508 vs DMSO control. This

revealed that transcripts whose translation was activated or suppressed by RMC-6272

showed similar regulation upon eFT-508 treatment and, vice versa, transcripts whose

translation was activated or suppressed by eFT-508 showed similar regulation upon RMC-

6272 treatment (Fig. 5A and B). To determine whether the agents reversed *TSC2*-sensitive

translation, we used the same strategy as above when comparing *TSC1*- vs *TSC2*-sensitive

translation (i.e. Fig. 3D). This revealed that both RMC-6272 as well as eFT-508 partially

reversed *TSC2*-sensitive translation (Fig. 5C). Gene ontology analysis of genes overlapping

in *TSC2*-Null NPCs treated with RMC-6272 or eFT-508 revealed significant reversal of some

enriched annotations related to cellular functions that were identified as *TSC2*-dependent

(shown in Fig. 3C), including positive regulation of transcription by RNA polymerase II, positive

regulation of RNA metabolic process and protein DNA complex organization along with

transcripts involved in various metabolic processes (Fig. 5D).

By comparing anota2seq outputs for these analyses we observed that, among all

transcripts whose translation was increased in *TSC2*-Null cells (Fig. 3A), 614 or 936 were

suppressed upon RMC-6272 or eFT-508 treatment, respectively, with an overlap of 513

transcripts; and 1454 or 1177 transcripts that were translationally suppressed in *TSC2*-Null

NPCs were translationally activated upon RMC-6272 or eFT-508 treatment, respectively, with

an overlap of 874 transcripts (all overlaps were statistically significant; Supplemental Fig. S4E,

Supplemental Table S4). Gene ontology analysis of *TSC2*-dependent differentially translated

genes (from Fig. 3A) revealed significant reversal of enriched annotations upon treatment with

RMC-6272 or eFT-508 as well as reversal of key cellular functions shown in Fig. 3C

(Supplemental Fig. S4F and G).

Although the inhibitors led to altered translation of similar numbers of transcripts in *TSC2*-

Null NPCs, anota2seq analysis contrasting the two treatments revealed substantial

differences in terms of affected mRNAs (Supplemental Fig. S5A and B). This was further

substantiated by gene ontology analysis, which revealed that while RMC-6272 e.g. more

efficiently suppressed translation of mRNAs encoding proteins involved in translation and

oxidative phosphorylation, mRNAs showing relatively more suppressed translation after

eFT508 treatment were e.g. annotated to neuronal development (Supplemental Fig. S5C).

Next, we assessed whether translation of mRNAs encoding ASD- and NDD-associated

genes were affected by mTORC1 or MNK1/2 activity. This revealed a striking reversal in

translation of both ASD and NDD genes in *TSC2*-Null NPCs treated with RMC-6272 or eFT-

508 (Fig. 6A). To identify affected genes, we compared outputs from the corresponding

anota2seq analyses. Among the 31 ASD72 genes translationally suppressed in *TSC2*-Null

NPCs, 28 (~90%) and 18 (~58%) were translationally activated upon RMC-6272 (RMC-

responsive) and eFT508 treatment (eFT-responsive), respectively (while all eFT-responsive

genes were identified as RMC-responsive, 10 genes were only responsive to RMC-6272;

Table 1, Fig. 6B). Moreover, for translationally suppressed NDD373 genes, 96 (~70%) and 69

(50%) were RMC- and eFT-responsive (Table 1), respectively (60 were both RMC- and eFT-

responsive; similar results were obtained for ASD185 and NDD664 genes; Supplemental

Table S2A). To validate alterations in translation upon *TSC2* loss as well as reversal by the

inhibitors, we selected five ASD/NDD-associated genes that were translationally suppressed

in *TSC2*-Null NPCs: *SETD1B* [MIM: 611055], *DYRK1A* [MIM: 600855], *CHD8* [MIM: 610528],

*NRXN1* [MIM: 600565], and *TCF7L2* [MIM: 62228], and also five eIF4E-sensitive genes that

were translationally activated in *TSC2*-Null NPCs: *AP2S1* [MIM: 602242], *LZTR1* [MIM:

600574], *WDR54*, *ALDOC* [MIM: 103870], and *EIF2B2* [MIM: 606454]. Quantitative RT-PCR

(RT-qPCR) analysis from polysome-associated mRNA fractions (>2 ribosomes) adjusted for

alterations in total mRNA showed a reduction for all five ASD/NDD-related genes and

increases in all five eIF4E-sensitive genes in *TSC2*-Null vs WT NPCs, confirming the above

polysome-profiling results. Moreover, RMC-6272 or eFT508 reversed *TSC2*-dependent

translationally suppressed ASD/NDD and translationally activated eIF4E-sensitive genes (Fig.

6C).

Taken together, *TSC2*-dependent decreased translation of mRNAs associated with ASD

and NDD can be reversed upon RMC-6272 or eFT508 treatment. These results are intriguing

and   potentially   imply   paradoxically   suppressed   translation   of   a   subset   of

neurodevelopmentally important genes that nonetheless depend on mTORC1-eIF4F pathway

activation.

**DISCUSSION**

The central role of mTORC1 signaling in translation initiation and protein synthesis in

a variety of cell types, including stem cells, as well as in human disease is well established [11;

49; 50]. While many transcriptome studies have been reported in *TSC1/2*-deficient neuronal cells,

how *TSC2* loss affects mRNA translation, particularly in patient-derived cells, is unknown. We

recently reported *TSC1*-dependent alterations in mRNA translation employing an isogenic set

of CRISPR-modified, TSC1 patient-derived NPCs. Although polysome-profiling revealed a

partial reversal of *TSC1*-associated gene expression changes following rapamycin treatment,

most genes related to neural activity/synaptic regulation or ASD remained rapamycin-

insensitive [24]. Here, we describe generation and characterization of an independent isogenic

set of CRISPR-modified NPCs from a TSC patient with a loss-of-function germline mutation

in *TSC2*. We find that *TSC2*-Null NPCs mimic *TSC1* loss in activation of mTORC1 signaling

and early ND phenotypic changes [24]. Rapamycin treatment of *TSC2*-Null NPCs fully inhibited

phosphorylation of S6K and reduced enlarged cell size but had minimal effects on 4E-BP1

phosphorylation and early ND phenotypes. In contrast, the third-generation mTORC1 inhibitor

RMC-6272 revealed efficient inhibition of both S6K and 4E-BP1 phosphorylation and rescued

the ND phenotypes as well as cell size. These data suggest that while enlarged cell size

observed upon *TSC2* loss depends on the mTORC1-S6K axis of the pathway, which is

consistent with previous reports [46], the altered ND phenotypes depend on the mTORC1-4E-

BP1 signaling axis.

We observed dramatic changes in mRNA translation associated with *TSC2* loss in NPCs

that largely overlapped with those observed upon *TSC1* loss. Provided that *TSC1* and *TSC2*

loss-assert similar effects on mTORC1 signaling, a substantial overlap is expected. Importantly

as the cells were derived from independent patients, our results support a robust effect on

mRNA translation across patients with hyperactive mTORC1 activity. In this study, in addition

to RMC-6272 that inhibits mTORC1 activity, we also considered alternative approaches to

modulate activity of the eIF4F complex and therefore  included the MNK1/2 inhibitor

eFT508/tomivosertib to understand the convergence  of mTORC1-4E-BP1 and MNK-eIF4E

signaling pathways on *TSC2*-dependent protein synthesis (see Fig.1A). While mTORC1-

sensitive translation has been studied extensively at the transcriptome-wide level [39; 51-53], the

understanding of how the MNK-eIF4E axis modulates mRNA translation at a global scale is

more limited [16]. This study is the first that includes a direct comparison between these

pathways toward mRNA translation and reveals that a large proportion of translational

alterations observed in *TSC2*-Null NPCs can be reversed by RMC-6272 or eFT-508, strongly

suggesting that these changes are indeed mediated by mTORC1-4E-BP1 and MNK-eIF4E

signaling via active eIF4F complex-dependent protein translation. However, the drugs are not

equivalent since RMC-6272 would be expected to affect both 4E-BP1-dependent and

MNK1/2-dependent translation whereas eFT-508 would be restricted to MNK1/2-dependent

translation. We also observe ample differences between the two drugs with respect to

translation, which are in line with previous studies showing differences in affected phenotypes

as the mTORC1-4E-BP1 axis is a key regulator of e.g. cell cycle progression [19; 54] while

MNK1/2-eIF4E controls e.g. cancer-associated phenotypes including cell migration [55; 56].

As epilepsy and ASD are strongly associated with TSC, we compared transcripts that were

differentially translated in *TSC2*-Null vs WT NPCs to recent genetic datasets available from

large cohorts of individuals with ASD, NDD and epilepsy [37; 41]. Interestingly, numerous non-

monogenic ASD-, NDD- and epilepsy-associated genes identified in patients harboring

putative loss-of-function mutations, including protein truncating or damaging missense

variants, were translationally suppressed in *TSC2*-Null NPCs and overlapped with data from

our previously reported *TSC1*-model. Notably, the results for ASD- and NDD-associated

genes were confirmed using multiple methods, which suggests that for these subsets there is

a strong negative regulatory effect on mRNA translation impacting a large proportion of these

mRNAs. This was not the case for epilepsy-associated genes, in that a smaller fraction of

these genes was translationally suppressed upon *TSC2* loss. Importantly, these decreases in

translation were observed side-by-side with the expected increased translation of mRNAs that

were previously reported to be translationally activated upon mTORC1 pathway activation.

This suppressed translation was, for a majority of identified ASD/NDD genes, reversed by

inhibition of mTORC1 using RMC-6272 or MNK1/2 using eFT-508. A recent elegant large-

scale transcriptome study demonstrated that CRISPR perturbation of several ASD genes

results in differential expression of ASD/NDD-related genes that positively correlate with co-

expression of the perturbed gene, thereby establishing a high level of convergence across

ASD risk genes [57]. Our results here showing translational suppression of many ASD/NDD

genes upon loss of *TSC2* is consistent with the positive correlation and co-expression reported

by Liao and colleagues [57]. However, most notably, our results demonstrate that a majority of

translationally suppressed ASD/NDD/epilepsy genes can be reversed by RMC-6272 or eFT-

508, thereby establishing the roles of mTORC1-4E-BP1 and MNK-eIF4E in regulating

translation of these genes.

Many of the translationally suppressed ASD genes are FMRP targets. FMRP was

originally considered to function as a translational repressor that can bind to numerous mRNAs

that are generally larger than average and enriched for genes associated with ID and ASD [58;

59]. However, recent studies of FMRP-deficiency  document a decrease in translation initiation

rates of many FMRP targets including ASD-related genes, which contrasts the notion that

FMRP is solely a translational repressor and supports a model where FMRP can also function

to increase the translation initiation of many targets [60-63]. Furthermore, in *TSC2*-Null NPCs we

also observe that 32 translationally suppressed ASD/NDD-related genes are associated with

chromatin-remodeling [64; 65]. Additional studies are therefore necessary to fully comprehend

how the ASD- and NDD-associated genes are translationally suppressed upon loss of *TSC2*.

In the literature, a main focus has been towards understanding mechanisms underlying

transcript-selective translation paralleling mTORC1 activity [13] while less attention has been

given to transcript that show decreased translation upon mTORC1 activation. Translation of

mRNAs enriched for 5'UTRs containing uORFs are translationally suppressed upon mTORC1

activation [39]. Consistently, we observed an increased frequency of uORFs in both ASD- and

NDD-associated genes; however, it should be noted that uORFs are also common among

background transcripts, and a relatively large proportion of ASD- and NDD-associated genes

do not have uORFs in their 5'UTRs.

Our results raise the critical question as to how ASD/NDD-related genes are translationally

suppressed upon mTORC1 activation in *TSC2*-Null NPCs. In addition to mTORC1, the

integrated stress response (ISR), which selectively modulates translation of transcripts with

5'UTRs harboring uORFs [66], is a key regulator of mRNA translation. In response to cellular

stress, phosphorylation of eukaryotic translation initiation factor 2 alpha (eIF2$\alpha$) is increased

by stress sensing kinases, which leads to a decrease in global protein synthesis paralleled by

increased translation of mRNAs with 5'-uORFs including the transcription factor ATF4

(reviewed in [67]). A recent study that employed multiplexed enhanced protein dynamics mass

spectrometry has shown that activation of the ISR pathway by cellular stressors and inhibition

of mTORC1 by Torin1 share  translatome changes with approximately 20% of the proteins

synthesized at reduced levels, demonstrating the convergence of the two pathways in

regulating translation of specific proteins [68]. In contrast, our results reveal that mTORC1

inhibition in fact reverses the translational downregulation of ASD/NDD genes upon mTORC1

activation in *TSC2*-deficient NPCs raising a question whether ISR signaling has any role in

TSC. The  involvement of the ISR pathway in our TSC models clearly needs further

investigation.

Another potential mechanism of translational regulation is via microRNAs (miRNAs), which

are involved in ND disorders [69]. Among 31 translationally suppressed transcripts in *TSC2*-Null

NPCs that show strong association with ASD (ASD72, FDR<0.001), 28 are predicted to be

targeted by at least one miRNA, using the miTarBase 2017 library (data not shown).

Interestingly, of the individual miRNAs identified by miTarBase analysis, four (hsa-miR-27b-

3p, hsa-let-7g-3p, hsa-miR-29a-3p and hsa-miR-515-5p) were reported as increased upon

*TSC1*- or *TSC2*-shRNA suppression in human neural stem cells generated from skin

fibroblast-derived iPSCs (p <0.05) [70]; and three (hsa-miR-21-5p, hsa-miR-19b-3p and hsa-

miR-144-3p) were shown to be overexpressed in human autism brain samples [71]. Based on

this, it is tempting to speculate that miRNAs may play a role in suppressing translation in

*TSC2*-Null NPCs, and future studies will address this mechanistic possibility.

In summary, we establish that loss of *TSC2* in patient-derived NPCs results in numerous

changes in mRNA translation, which significantly overlaps with loss of *TSC1*. Loss of *TSC2*

associates with suppressed translation of a large proportion of ASD- and NDD-related genes,

which is reversed by inhibition of mTORC1 or MNK1/2, demonstrating the importance of

eIF4F-dependent mRNA translation in ND disorders. Our work also supports the therapeutic

potential for RMC-5552 (the clinical counterpart of RMC-6272) and eFT508/tomivosertib that

are currently under clinical development for cancer, as a strategy for treatment of TAND as

well as ASD and other neurodevelopmental disorders.

**DECLARATION OF INTERESTS**

The authors declare no competing interests.

**ACKNOWLEDGEMENTS**

This paper is dedicated to the memory of the late Dr. Jerry Pelletier, a distinguished James

McGill Professor in the Department of Biochemistry and Oncology from the Goodman Cancer

Institute at McGill University, Montreal, Canada, who unfortunately passed away during the

completion of this study. This work was supported by NIH R01 NS109540 (V.R.), the Swedish

Brain foundation FO2022-0178 (O.L.); the Swedish research council 2020-01665 (O.L.); the

Wallenberg academy fellow program 2013.0181 (O.L.).

**AUTHOR CONTRIBUTIONS**

Conceptualization and study design: V.R. and O.L.; experimental methods and data

acquisition P.M., K.S., F.R., S.B., R.B., N.R., J.B., S.C. and I.N.; data analysis and

interpretation: P.M., K.S., F.R., S.B., R.B. S.C., I.N., S.H., S.E. and O.L.; manuscript

preparation and editing: P.M., K.S., F.R., S.B., R.B., I.N., S.H., S.E., O.L. and V.R.

**WEB RESOURCES**

OMIM, https://www.omim.org/

GenScript, https://www.genscript.com

BBTools software suite, https://www.osti.gov/servlets/purl/1241166

R statistical software environment, https://www.R-project.org/

**DATA AND CODE AVAILABILITY**

The polysome-profiling datasets supporting the conclusions of this article are included within

the article (and its additional file(s)), and the full datasets will be deposited into Gene

Expression Omnibus (GEO).

**REFERENCES**

1. de Vries, P.J., Heunis, T.M., Vanclooster, S., Chambers, N., Bissell, S., Byars, A.W., Flinn,

J., Gipson, T.T., van Eeghen, A.M., Waltereit, R., et al. (2023). International consensus

recommendations for the identification and treatment of tuberous sclerosis complex-

associated neuropsychiatric disorders (TAND). J Neurodev Disord 15, 32.

2. Henske, E.P., Jozwiak, S., Kingswood, J.C., Sampson, J.R., and Thiele, E.A. (2016).

Tuberous sclerosis complex. Nat Rev Dis Primers 2, 16035.

3. Sahin, M., Henske, E.P., Manning, B.D., Ess, K.C., Bissler, J.J., Klann, E., Kwiatkowski,

D.J., Roberds, S.L., Silva, A.J., Hillaire-Clarke, C.S., et al. (2016). Advances and

Future Directions for Tuberous Sclerosis Complex Research: Recommendations From

the 2015 Strategic Planning Conference. Pediatr Neurol 60, 1-12.

4. Bissler, J.J., McCormack, F.X., Young, L.R., Elwing, J.M., Chuck, G., Leonard, J.M.,

Schmithorst, V.J., Laor, T., Brody, A.S., Bean, J., et al. (2008). Sirolimus for

angiomyolipoma in tuberous sclerosis complex or lymphangioleiomyomatosis. N Engl

J Med 358, 140-151.

5. Davies, D.M., de Vries, P.J., Johnson, S.R., McCartney, D.L., Cox, J.A., Serra, A.L.,

Watson, P.C., Howe, C.J., Doyle, T., Pointon, K., et al. (2011). Sirolimus therapy for

angiomyolipoma in tuberous sclerosis and sporadic lymphangioleiomyomatosis: a

754  phase 2 trial. Clinical cancer research : an official journal of the American Association

755   for Cancer Research 17, 4071-4081.

756 6. Julich, K., and Sahin, M. (2014). Mechanism-based treatment in tuberous sclerosis

757   complex. Pediatric neurology 50, 290-296.

758 7. Krueger, D.A., Care, M.M., Holland, K., Agricola, K., Tudor, C., Mangeshkar, P., Wilson,

759   K.A., Byars, A., Sahmoud, T., and Franz, D.N. (2010). Everolimus for subependymal

760   giant-cell astrocytomas in tuberous sclerosis. The New England journal of medicine

761   363, 1801-1811.

762 8. Overwater, I.E., Rietman, A.B., van Eeghen, A.M., and de Wit, M.C.Y. (2019). Everolimus

763   for the treatment of refractory seizures associated with tuberous sclerosis complex

764   (TSC): current perspectives. Ther Clin Risk Manag 15, 951-955.

765 9. Krueger, D.A., Sadhwani, A., Byars, A.W., de Vries, P.J., Franz, D.N., Whittemore, V.H.,

766   Filip-Dhima, R., Murray, D., Kapur, K., and Sahin, M. (2017). Everolimus for treatment

767   of tuberous sclerosis complex-associated neuropsychiatric disorders. Ann Clin Transl

768   Neurol 4, 877-887.

769 10. Overwater, I.E., Rietman, A.B., Mous, S.E., Bindels-de Heus, K., Rizopoulos, D., Ten

770   Hoopen, L.W., van der Vaart, T., Jansen, F.E., Elgersma, Y., Moll, H.A., et al. (2019).

771   A randomized controlled trial with everolimus for IQ and autism in tuberous sclerosis

772   complex. Neurology 93, e200-e209.

773 11. Saba, J.A., Liakath-Ali, K., Green, R., and Watt, F.M. (2021). Translational control of stem

774   cell function. Nat Rev Mol Cell Biol 22, 671-690.

775 12. Thoreen, C.C. (2017). The molecular basis of mTORC1-regulated translation. Biochem

776   Soc Trans 45, 213-221.

777 13. Masvidal, L., Hulea, L., Furic, L., Topisirovic, I., and Larsson, O. (2017). mTOR-sensitive

778   translation: Cleared fog reveals more trees. RNA Biol 14, 1299-1305.

14. Gandin, V., Masvidal, L., Cargnello, M., Gyenis, L., McLaughlan, S., Cai, Y., Tenkerian,

C., Morita, M., Balanathan, P., Jean-Jean, O., et al. (2016). mTORC1 and CK2

coordinate ternary and eIF4F complex assembly. Nat Commun 7, 11127.

15. Bramham, C.R., Jensen, K.B., and Proud, C.G. (2016). Tuning Specific Translation in

Cancer Metastasis and Synaptic Memory: Control at the MNK-eIF4E Axis. Trends

Biochem Sci 41, 847-858.

16. Karampelias, C., Watt, K., Mattsson, C.L., Ruiz, A.F., Rezanejad, H., Mi, J., Liu, X., Chu,

L., Locasale, J.W., Korbutt, G.S., et al. (2022). MNK2 deficiency potentiates beta-cell

regeneration via translational regulation. Nat Chem Biol 18, 942-953.

17. Choo, A.Y., Yoon, S.O., Kim, S.G., Roux, P.P., and Blenis, J. (2008). Rapamycin

differentially inhibits S6Ks and 4E-BP1 to mediate cell-type-specific repression of

mRNA translation. Proceedings of the National Academy of Sciences of the United

States of America 105, 17414-17419.

18. Fan, Q.W., Nicolaides, T.P., and Weiss, W.A. (2018). Inhibiting 4EBP1 in Glioblastoma.

Clin Cancer Res 24, 14-21.

19. Larsson, O., Morita, M., Topisirovic, I., Alain, T., Blouin, M.J., Pollak, M., and Sonenberg,

N. (2012). Distinct perturbation of the translatome by the antidiabetic drug metformin.

Proc Natl Acad Sci U S A 109, 8977-8982.

20. Rodrik-Outmezguine, V.S., Okaniwa, M., Yao, Z., Novotny, C.J., McWhirter, C., Banaji, A.,

Won, H., Wong, W., Berger, M., de Stanchina, E., et al. (2016). Overcoming mTOR

resistance mutations with a new-generation mTOR inhibitor. Nature 534, 272-276.

21. Bhattacharyya, S., Oblinger, J.L., Beauchamp, R.L., Kosa, L., Robert, F., Plotkin, S.R.,

Chang, L.S., and Ramesh, V. (2024). Preclinical evaluation of the third-generation, bi-

steric mechanistic target of rapamycin complex 1-selective inhibitor RMC-6272 in NF2-

deficient models. Neurooncol Adv 6, vdae024.

22. Burnett, G.L., Yang, Y.C., Aggen, J.B., Pitzen, J., Gliedt, M.K., Semko, C.M., Marquez, A.,

Evans, J.W., Wang, G., Won, W.S., et al. (2023). Discovery of RMC-5552, a Selective

Bi-Steric Inhibitor of mTORC1, for the Treatment of mTORC1-Activated Tumors. J Med

Chem 66, 149-169.

23. Reich, S.H., Sprengeler, P.A., Chiang, G.G., Appleman, J.R., Chen, J., Clarine, J., Eam,

B., Ernst, J.T., Han, Q., Goel, V.K., et al. (2018). Structure-based Design of Pyridone-

Aminal eFT508 Targeting Dysregulated Translation by Selective Mitogen-activated

Protein Kinase Interacting Kinases 1 and 2 (MNK1/2) Inhibition. J Med Chem 61, 3516-

3540.

24. Aksoylu, I.S., Martin, P., Robert, F., Szkop, K.J., Redmond, N.E., Bhattacharyya, S.,

Wang, J., Chen, S., Beauchamp, R.L., Nobeli, I., et al. (2023). Translatome analysis

of tuberous sclerosis complex 1 patient-derived neural progenitor cells reveals

rapamycin-dependent and independent alterations. Mol Autism 14, 39.

25. Niida, Y., Stemmer-Rachamimov, A.O., Logrip, M., Tapon, D., Perez, R., Kwiatkowski,

D.J., Sims, K., MacCollin, M., Louis, D.N., and Ramesh, V. (2001). Survey of somatic

mutations in tuberous sclerosis complex (TSC) hamartomas suggests different genetic

mechanisms for pathogenesis of TSC lesions. Am J Hum Genet 69, 493-503.

26. Schlaeger, T.M., Daheron, L., Brickler, T.R., Entwisle, S., Chan, K., Cianci, A., DeVine, A.,

Ettenger, A., Fitzgerald, K., Godfrey, M., et al. (2015). A comparison of non-integrating

reprogramming methods. Nat Biotechnol 33, 58-63.

27. Warren, L., Manos, P.D., Ahfeldt, T., Loh, Y.H., Li, H., Lau, F., Ebina, W., Mandal, P.K.,

Smith, Z.D., Meissner, A., et al. (2010). Highly efficient reprogramming to pluripotency

and directed differentiation of human cells with synthetic modified mRNA. Cell Stem

Cell 7, 618-630.

28. Martin, P., Wagh, V., Reis, S.A., Erdin, S., Beauchamp, R.L., Shaikh, G., Talkowski, M.,

Thiele, E., Sheridan, S.D., Haggarty, S.J., et al. (2020). TSC patient-derived isogenic

neural progenitor cells reveal altered early neurodevelopmental phenotypes and

rapamycin-induced MNK-eIF4E signaling. Mol Autism 11, 2.

29. Sellgren, C.M., Gracias, J., Watmuff, B., Biag, J.D., Thanos, J.M., Whittredge, P.B., Fu,
T., Worringer, K., Brown, H.E., Wang, J., et al. (2019). Increased synapse elimination
by microglia in schizophrenia patient-derived models of synaptic pruning. Nat Neurosci
22, 374-385.

30. Han, S., Kim, S., Bahl, S., Li, L., Burande, C.F., Smith, N., James, M., Beauchamp, R.L.,
Bhide, P., Diantonio, A., et al. (2012). The E3 Ubiquitin Ligase Protein Associated with
Myc (Pam) Regulates Mammalian/Mechanistic Target of Rapamycin Complex 1
(mTORC1) Signaling in Vivo through N- and C-terminal Domains. The Journal of
biological chemistry 287, 30063-30072.

31. James, M.F., Han, S., Polizzano, C., Plotkin, S.R., Manning, B.D., Stemmer-Rachamimov,
A.O., Gusella, J.F., and Ramesh, V. (2009). NF2/merlin is a novel negative regulator
of mTOR complex 1, and activation of mTORC1 is associated with meningioma and
schwannoma growth. Molecular and cellular biology 29, 4250-4261.

32. Wang, D., Lagerstrom, R., Sun, C., Bishof, L., Valotton, P., and Gotte, M. (2010). HCA-
vision: Automated neurite outgrowth analysis. J Biomol Screen 15, 1165-1170.

33. Gandin, V., Sikstrom, K., Alain, T., Morita, M., McLaughlan, S., Larsson, O., and
Topisirovic, I. (2014). Polysome fractionation and analysis of mammalian translatomes
on a genome-wide scale. J Vis Exp.

34. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol
37, 907-915.

35. Liao, Y., Smyth, G.K., and Shi, W. (2019). The R package Rsubread is easier, faster,
cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic
Acids Res 47, e47.

36. Oertlin, C., Lorent, J., Murie, C., Furic, L., Topisirovic, I., and Larsson, O. (2019). Generally
applicable transcriptome-wide analysis of translation using anota2seq. Nucleic Acids
Res 47, e70.

37. Fu, J.M., Satterstrom, F.K., Peng, M., Brand, H., Collins, R.L., Dong, S., Wamsley, B., Klei,
L., Wang, L., Hao, S.P., et al. (2022). Rare coding variation provides insight into the
genetic architecture and phenotypic context of autism. Nat Genet 54, 1320-1331.

38. Larsson, O., Li, S., Issaenko, O.A., Avdulov, S., Peterson, M., Smith, K., Bitterman, P.B.,
and Polunovsky, V.A. (2007). Eukaryotic translation initiation factor 4E induced
progression of primary human mammary epithelial cells along the cancer pathway is
associated with targeted translational deregulation of oncogenic drivers and inhibitors.
Cancer Res 67, 6814-6824.

39. Gandin, V., Masvidal, L., Hulea, L., Gravel, S.P., Cargnello, M., McLaughlan, S., Cai, Y.,
Balanathan, P., Morita, M., Rajakumar, A., et al. (2016). nanoCAGE reveals 5' UTR
features that define specific modes of translation of functionally related MTOR-
sensitive mRNAs. Genome Res 26, 636-648.

40. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. Proc Natl Acad Sci U S A 102, 15545-15550.

41. Epi, C., Chen, S., Neale, B.M., and Berkovic, S.F. (2023). Shared and distinct ultra-rare
genetic risk for diverse epilepsies: A whole-exome sequencing study of 54,423
individuals across multiple genetic ancestries. medRxiv.

42. Blair, J.D., Hockemeyer, D., and Bateup, H.S. (2018). Genetically engineered human
cortical spheroid models of tuberous sclerosis. Nat Med 24, 1568-1578.

43. Costa, V., Aigner, S., Vukcevic, M., Sauter, E., Behr, K., Ebeling, M., Dunkley, T., Friedlein,
A., Zoffmann, S., Meyer, C.A., et al. (2016). mTORC1 Inhibition Corrects
Neurodevelopmental and Synaptic Alterations in a Human Stem Cell Model of
Tuberous Sclerosis. Cell Rep 15, 86-95.

44. Sundberg, M., Tochitsky, I., Buchholz, D.E., Winden, K., Kujala, V., Kapur, K., Cataltepe,
D., Turner, D., Han, M.J., Woolf, C.J., et al. (2018). Purkinje cells derived from TSC

patients display hypoexcitability and synaptic deficits associated with reduced FMRP

levels and reversed by rapamycin. Mol Psychiatry 23, 2167-2183.

45. Zucco, A.J., Pozzo, V.D., Afinogenova, A., Hart, R.P., Devinsky, O., and D'Arcangelo, G.

(2018). Neural progenitors derived from Tuberous Sclerosis Complex patients exhibit

attenuated PI3K/AKT signaling and delayed neuronal differentiation. Mol Cell Neurosci

92, 149-163.

46. Dowling, R.J., Topisirovic, I., Alain, T., Bidinosti, M., Fonseca, B.D., Petroulakis, E., Wang,

X., Larsson, O., Selvaraj, A., Liu, Y., et al. (2010). mTORC1-mediated cell proliferation,

but not cell growth, controlled by the 4E-BPs. Science 328, 1172-1176.

47. Oertlin, C., Watt, K., Ristau, J., and Larsson, O. (2022). Anota2seq Analysis for

Transcriptome-Wide Studies of mRNA Translation. Methods Mol Biol 2418, 243-268.

48. Sandri, B.J., Masvidal, L., Murie, C., Bartish, M., Avdulov, S., Higgins, L., Markowski, T.,

Peterson, M., Bergh, J., Yang, P., et al. (2019). Distinct Cancer-Promoting Stromal

Gene Expression Depending on Lung Function. Am J Respir Crit Care Med 200, 348-

358.

49. Bartish, M., Abraham, M.J., Goncalves, C., Larsson, O., Rolny, C., and Del Rincon, S.V.

(2023). The role of eIF4F-driven mRNA translation in regulating the tumour

microenvironment. Nat Rev Cancer 23, 408-425.

50. Tahmasebi, S., Khoutorsky, A., Mathews, M.B., and Sonenberg, N. (2018). Translation

deregulation in human disease. Nat Rev Mol Cell Biol 19, 791-807.

51. Lee, B.J., Boyer, J.A., Burnett, G.L., Thottumkara, A.P., Tibrewal, N., Wilson, S.L., Hsieh,

T., Marquez, A., Lorenzana, E.G., Evans, J.W., et al. (2021). Selective inhibitors of

mTORC1 activate 4EBP1 and suppress tumor growth. Nat Chem Biol 17, 1065-1074.

52. Silvera, D., Ernlund, A., Arju, R., Connolly, E., Volta, V., Wang, J., and Schneider, R.J.

(2017). mTORC1 and -2 Coordinate Transcriptional and Translational Reprogramming

in Resistance to DNA Damage and Replicative Stress in Breast Cancer Cells. Mol Cell

Biol 37.

53. Volta, V., Perez-Baos, S., de la Parra, C., Katsara, O., Ernlund, A., Dornbaum, S., and
Schneider, R.J. (2021). A DAP5/eIF3d alternate mRNA translation mechanism
promotes differentiation and immune suppression by human regulatory T cells. Nat
Commun 12, 6979.

54. Mamane, Y., Petroulakis, E., Martineau, Y., Sato, T.A., Larsson, O., Rajasekhar, V.K., and
Sonenberg, N. (2007). Epigenetic activation of a subset of mRNAs by eIF4E explains
its effects on cell proliferation. PLoS One 2, e242.

55. Beggs, J.E., Tian, S., Jones, G.G., Xie, J., Iadevaia, V., Jenei, V., Thomas, G., and Proud,
C.G. (2015). The MAP kinase-interacting kinases regulate cell migration, vimentin
expression and eIF4E/CYFIP1 binding. Biochem J 467, 63-76.

56. Robichaud, N., del Rincon, S.V., Huor, B., Alain, T., Petruccelli, L.A., Hearnden, J.,
Goncalves, C., Grotegut, S., Spruck, C.H., Furic, L., et al. (2015). Phosphorylation of
eIF4E promotes EMT and metastasis via translational control of SNAIL and MMP-3.
Oncogene 34, 2032-2042.

57. Liao, C., Moyses-Oliveira, M., De Esch, C.E.F., Bhavsar, R., Nuttle, X., Li, A., Yu, A., Burt,
N.D., Erdin, S., Fu, J.M., et al. (2023). Convergent coexpression of autism-associated
genes suggests some novel risk genes may not be detectable in large-scale genetic
studies. Cell Genom 3, 100277.

58. Ascano, M., Jr., Mukherjee, N., Bandaru, P., Miller, J.B., Nusbaum, J.D., Corcoran, D.L.,
Langlois, C., Munschauer, M., Dewell, S., Hafner, M., et al. (2012). FMRP targets
distinct mRNA sequence elements to regulate protein expression. Nature 492, 382-
386.

59. Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y., Mele, A., Fraser, C.E., Stone,
E.F., Chen, C., Fak, J.J., Chi, S.W., et al. (2011). FMRP stalls ribosomal translocation
on mRNAs linked to synaptic function and autism. Cell 146, 247-261.

60. Das Sharma, S., Metz, J.B., Li, H., Hobson, B.D., Hornstein, N., Sulzer, D., Tang, G., and

Sims, P.A. (2019). Widespread Alterations in Translation Elongation in the Brain of

Juvenile Fmr1 Knockout Mice. Cell Rep 26, 3313-3322 e3315.

61. Flanagan, K., Baradaran-Heravi, A., Yin, Q., Dao Duc, K., Spradling, A.C., and Greenblatt,

E.J. (2022). FMRP-dependent production of large dosage-sensitive proteins is highly

conserved. Genetics 221.

62. Richter, J.D., and Zhao, X. (2021). The molecular biology of FMRP: new insights into

fragile X syndrome. Nat Rev Neurosci 22, 209-222.

63. Sawicka, K., Hale, C.R., Park, C.Y., Fak, J.J., Gresack, J.E., Van Driesche, S.J., Kang,

J.J., Darnell, J.C., and Darnell, R.B. (2019). FMRP has a cell-type-specific role in CA1

pyramidal neurons to regulate autism-related transcripts and circadian memory. Elife

8.

64. Gabriele, M., Lopez Tobon, A., D'Agostino, G., and Testa, G. (2018). The chromatin basis

of neurodevelopmental disorders: Rethinking dysfunction along the molecular and

temporal axes. Prog Neuropsychopharmacol Biol Psychiatry 84, 306-327.

65. Mossink, B., Negwer, M., Schubert, D., and Nadif Kasri, N. (2021). The emerging role of

chromatin remodelers in neurodevelopmental disorders: a developmental perspective.

Cell Mol Life Sci 78, 2517-2563.

66. Guan, B.J., van Hoef, V., Jobava, R., Elroy-Stein, O., Valasek, L.S., Cargnello, M., Gao,

X.H., Krokowski, D., Merrick, W.C., Kimball, S.R., et al. (2017). A Unique ISR Program

Determines Cellular Responses to Chronic Stress. Mol Cell 68, 885-900 e886.

67. Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljujic, M., Samali, A., and Gorman, A.M. (2016).

The integrated stress response. EMBO Rep 17, 1374-1395.

68. Klann, K., Tascher, G., and Munch, C. (2020). Functional Translatome Proteomics Reveal

Converging and Dose-Dependent Regulation by mTORC1 and eIF2alpha. Mol Cell 77,

913-925 e914.

964 69. Sun, E., and Shi, Y. (2015). MicroRNAs: Small molecules with big roles in

965   neurodevelopment and diseases. Exp Neurol 268, 46-53.

966 70. Pawlik, B., Grabia, S., Smyczynska, U., Fendler, W., Drozdz, I., Liszewska, E., Jaworski,

967   J., Kotulska, K., Jozwiak, S., Mlynarski, W., et al. (2022). MicroRNA Expression Profile

968   in TSC Cell Lines and the Impact of mTOR Inhibitor. Int J Mol Sci 23.

969 71. Mor, M., Nardone, S., Sams, D.S., and Elliott, E. (2015). Hypomethylation of miR-142

970   promoter and upregulation of microRNAs that target the oxytocin receptor gene in the

971   autism prefrontal cortex. Mol Autism 6, 46.

973 **FIGURE TITLES AND LEGENDS**

975 **Figure 1. Treatment with RMC-6272 or rapamycin rescues increased cell size in *TSC2*-**

976 **mutant NPCs.**

977 (A) Schematic representation of how mTORC1 signaling modulates mRNA translation.

978 Inhibitors used to target the pathway axes are indicated in red.

979 (B) Representative bright field images, including enlarged inset images (black boxed regions),

980 are shown for *TSC2*-Het and -Null NPCs compared with *TSC2*-WT cells. Scale bar=100µm.

981 (C) Immunoblotting of an isogenic set of *TSC2*-WT, -Het and -Null (cl. 39) NPCs, following 24

982 h treatment with DMSO (control), rapamycin (50nM) or RMC-6272 (10nM)  are shown for

983 TSC2/tuberin as well as mTORC1 pathway readouts pS6K T389, pS6 S240/244 and p4E-BP1

984 S65 along with respective total proteins.

985 (D) Average cell size for *TSC2*-Het and -Null NPCs relative to *TSC2*-WT cells following 48 h

986 with DMSO (control),  rapamycin or RMC-6272 (same concentrations as in C) was quantitated,

987 and data represents the mean (+SEM) from 3 independent experiments, each with triplicate

988 technical replicates. P-values of comparisons (*$p<0.05$, **$p<0.01$, ***$p<0.001$) from Students

989 t-test are indicated.

**Figure 2. Altered neurodevelopmental phenotypes in *TSC2* mutant NPCs are rescued**

**by RMC-6272, but not rapamycin.**

(A) Proliferation by live cell counting (trypan blue exclusion) was carried out over days D1-4

in *TSC2*-WT, -Het and -Null NPCs under control treatment (left graph region), or treatment

initiated at D1 with rapamycin (50 nM, middle graph region) or RMC-6272 (10 nM, right graph

region). Quantitation for each timepoint is normalized to cell seeding number (D0=1). Data

represents the mean ($\pm$SEM) from 3 independent experiments (3 replicates/cell

line/experiment). Student's t-test p-values (*$p<0.05$, **$p<0.01$, ***$p<0.001$) are indicated in

DMSO-treated NPCs (left graph region) for increasing cell numbers in *TSC2*-Het and Null

NPCs at each timepoint (D1-4) relative to *TSC2*-WT NPCs at the respective timepoint. For

drug-treated NPCs (middle and right graph regions), within each genotype, p-values are

indicated for changes in cell numbers relative to DMSO controls at the corresponding

timepoint.

(B) Representative images of immunofluorescence staining in NPCs using neuronal marker

MAP2 (gray), performed 48 h after initiating nerite outgrowth, are shown along with traced

images generated employing HCA-Vision image quantitation software (green), which traces

the cell body along branching layers of the neurite arborization. Scale bar = 100 µm.

(C and D) Bar plots indicate average neurite length (left panel), average neurite number

(center) and extremities (right) per cell for *TSC2*-Het and -Null NPCs relative to *TSC2*-WT

NPCs under control conditions (C) or for rapamycin (50 nM) or RMC-6272 (10 nM) treatment

in *TSC2*-Het and -Null cells relative to DMSO control (D) . Data represents mean ($\pm$SEM) from

3 independent experiments (8 non-overlapping image fields/cell line/experiment). P-values

from Student's t-test (*$p<0.05$, **$p<0.01$, ***$p<0.001$) are shown.

1014

1015 **Figure 3. Extensive remodeling of mRNA translation in *TSC2*-Null NPCs aligns with**

1016 **those observed following loss of *TSC1*.**

(A) Scatterplot of polysome-associated mRNA versus total mRNA log2 fold changes between *TSC2*-Null (-/-) and WT NPCs. Genes are colored according to their mode of regulation as identified by the anota2seq analysis (FDR < 0.15). The number of mRNAs regulated via translation, mRNA abundance or offsetting is indicated.

(B) Kernel densities of p-values and FDRs (from anota2seq analysis) for the comparison of *TSC2*-Null versus WT NPCs. Densities are shown for analysis of polysome-associated RNA (translated mRNA), total mRNA, translation and offsetting. A shift of the density towards low p-values/FDRs indicates a higher frequency of changes.

(C) Enrichment of top ten gene ontology annotation for biological processes among genes that undergo significant changes in translation (light red – activated translation, dark red – suppressed translation) in *TSC2*-Null versus WT NPCs.

(D) Scatterplots from anota2seq analysis of *TSC2*-Null versus WT NPCs (top left and bottom right panels) and *TSC1*-Null versus WT NPCs (top right and bottom left panels). Transcripts identified as regulated by translation in anota2seq analysis in the *TSC2*-Null versus WT NPC (top left and right panels) or *TSC1*-Null versus WT NPC (bottom left and right panels) comparisons are indicated. Also shown are empirical cumulative distribution functions of the log2 fold changes in translation for the corresponding gene sets. Grey dots and lines correspond to the background (i.e. not in gene sets). Significant shifts of gene sets, relative to control, were determined using Wilcoxon rank-sum test and the resulting p-values are shown.

(E) Venn diagrams comparing translationally regulated genes identified in anota2seq analysis following loss of *TSC1* or *TSC2*.

(F) Enrichments of gene ontology annotation for biological processes among genes identified as showing consistent regulation in both *TSC1*-Null and *TSC2*-Null NPCs relative to their controls (same color codes as in panel C).

**Figure 4. Suppressed translation of transcripts encoding ASD- and NDD-associated proteins in *TSC2*-Null NPCs associates with distinct 5'UTR features.**

(A) GSEA analysis of ASD- and NDD-associated genes and anota2seq analysis of translation

(fold-changes) of the comparison of *TSC2*-Null (-/-) versus WT NPCs. Rankings for ASD- and

NDD-associated genes (ASD72 and NDD373, respectively)  were shifted towards the right

denoting suppressed translation. Normalized enrichment scores (NES) and adjusted p-values

are shown.

(B) Scatterplots from anota2seq analysis for *TSC2*-Null versus WT NPCs. High confidence

ASD- and NDD-associated genes (ASD72 and NDD373, respectively) (top left panel) as well

as eIF4E- (middle left panel) or mTOR-sensitive (bottom left panel) transcripts are indicated.

Shown to the right of each panel are corresponding empirical distribution functions together

with associated statistics (as in Fig. 3D).

(C-D) Boxplots of 5'UTR lengths (C) and GC-contents (D) (according to the RefSeq database,

version 109.20201120) for ASD- and NDD-associated genes as well as mRNAs translationally

regulated in *TSC2*-Null versus WT NPCs (after excluding ASD- and NDD-associated genes;

background corresponds to mRNAs not in any of the subsets). The mean length or GC-

contents for each set are indicated together with p-values from Wilcoxon-tests. (E) The

proportion of mRNAs harboring upstream open reading frames (uORFs) in their 5'UTRs

among subsets of transcripts from C-D. The lower part of the bar graphs corresponds to

uORFs with a strong Kozak context. Differences in proportions of uORFs were assessed using

Fisher's exact tests for the indicated comparisons.

1063

**Figure 5. mTORC1 or MNK1/2 inhibition leads to changes in mRNA translation and**

**reverses translation of suppressed transcripts in *TSC2*-Null NPCs.**

(A and B) Scatterplots from anota2seq analysis of *TSC2*-Null NPCs treated with RMC-6272

versus DMSO (A, top left panel) or eFT-508 versus DMSO (B, top left panel). Transcripts

identified as regulated by translation in anota2seq analysis in the RMC-6272 versus DMSO

treated NPCs (A, left panels) or eFT-508 versus DMSO treated NPCs (B, left panels)

comparisons are indicated. Also shown are empirical cumulative distribution functions (A and

B, right panels) of the log2 fold changes in translation for the corresponding gene sets. Grey

dots and lines correspond to the background (i.e. not in gene sets). Significant shifts of gene

sets, relative to control, were determined using Wilcoxon rank-sum test and the resulting p-

values are shown.

(C) Assessing how transcripts whose translation was altered in *TSC2*-Null vs WT NPCs (top

panel, from Fig. 3A) are modulated upon treatment with RMC-6272 (bottom left panel) or eFT-

508 (bottom right panel).

(D) Enrichments of gene ontology annotation for biological processes among genes identified

as showing reversal in translation overlapping between RMC-6272 and eFT-508 treatment

(same color codes as in Figure 3C).

**Figure 6. mTORC1 or MNK1/2 inhibition reverses translation of suppressed transcripts**

**encoding ASD- and NDD-associated proteins in *TSC2*-Null NPCs.**

(A) Scatterplots for high confidence ASD- and NDD-associated genes (ASD72 and NDD373)

from anota2seq analysis of *TSC2*-Null versus WT NPCs (top panel) as well as RMC-6272

(bottom left) or eFT508 (bottom right) treatment of *TSC2*-Null NPCs are shown. The shape

and position of ellipses correspond to the mean and standard deviation for all genes within

each gene set, respectively.

(B) A Venn diagram comparing ASD associated genes (ASD72) whose translation was

suppressed in *TSC2*-Null NPC relative to *TSC2*-WT NPCs and reversed by treatment with

RMC-6272 or eFT-508.

(C) Plots denote RT-qPCR analysis from the polysome-associated RNA pool (3 biological

replicates) for 5 ASD- or NDD-associated genes (left) as well as 5 eIF4E-sensitive genes

(right) from the indicated cells and treatments. Data were normalized to total mRNA obtained

in parallel as well as beta-actin mRNA expression. P-values from student's t-test test (*p<0.05,

**p<0.01, ***p<0.001) for the indicated comparisons are shown.

| Drug Tx* | Direction of regulation | Geneset name | # Genes | pV | Gene ID |
|---|---|---|---|---|---|
| DMSO | Down (baseline) | ASD72 | 31 | 1.72E-07 | *MED13, NIPBL, ASH1L, KMT2A, CHD2, EHMT1, KMT5B, CHD8, WAC, RFX3, TLK2, SMARCC2, POGZ, KMT2C, ADNP, KCNB1, MEIS2, ASXL3, DYRK1A, SPAST, SETD5, VEZF1, AUTS2, NRXN1, TCF7L2, TAOK1, NF1, ZBTB20, MED23, KDM5B, DNMT3A* |
| RMC | Up | ASD72 | 28 | 1.78E-11 | *MED13, NIPBL, ASH1L, KMT2A, CHD2, EHMT1, KMT5B, CHD8, WAC, RFX3, TLK2, POGZ, KMT2C, ADNP, MEIS2, ASXL3, DYRK1A, SPAST, SETD5, VEZF1, AUTS2, NRXN1, TCF7L2, TAOK1, NF1, MED23, KDM5B, DNMT3A* |
| eFT508 | Up | ASD72 | 18 | 8.63E-06 | *MED13, NIPBL, ASH1L, KMT2A, CHD2, KMT5B, WAC, RFX3, KMT2C, MEIS2, ASXL3, SPAST, VEZF1, AUTS2, NRXN1, TCF7L2, TAOK1, NF1* |
| n/a | Down (baseline) | NDD373 | 138 | 2.59E-19 | *MED13, NIPBL, SMC3, CNOT1, KMT2E, UBTF, KDM3B, TRRAP, USP7, ASH1L, KMT2A, CREBBP, KAT6A, SIN3A, CHD2, ARID1A, SPEN, EHMT1, TLE3, UBR5, CHD7, KMT5B, HIVEP2, CHD8, WAC, ZNF462, NSD1, ARID2, GATAD2B, FBXO11, TAB2, HDAC2, KMT2D, ZNF148, RFX3, TBL1XR1, ZEB2, SETBP1, PHIP, TLK2, RAI1, KIF5C, SMARCC2, SON, STAG1, POGZ, ZMYND11, KIF11, KMT2C, ADNP, KCNB1, COL4A1, KAT6B, SCN8A, ZMYND8, RFX7, PHF12, ZFHX3, SHOC2, TRIO, SOS1, ZNF292, ZFHX4, GNB1, SETD1B, TANC2, KIF2A, ARIH1, SOX9, ACVR2A, TNRC6B, SET, MEIS2, CTNND1, SMARCE1, GLI3, YY1, ASXL3, PDE4D, SETD2, SYNCRIP, BRAF, MAP3K7, DYRK1A, WDR48, TCF4, SRP54, SMAD4, SPAST, SETD5, HNRNPD, PIK3R1, SNAP25, FBXW7, VEZF1, SRSF1, KLF7, TOP2B, ZMIZ1, AUTS2, NRXN1, TRIM8, PPP2CA, CHAMP1, TCF7L2, TAOK1, ATF2, ZBTB10, NF1, SATB1, CSNK2A1, BICD2, CAPRIN1, ZBTB20, CDK13, ZMYM2, GABRB3, CBX5, TCF12, MAP2K1, MYCN, CDK8, SOX11, SYT1, TRPM3, RAC1, MSL2, ABCC9, MED23, DNM1L, KLHL20, TNPO3, KDM5B, KBTBD7, PPM1D, KRAS, DNMT3A, MIB1* |
| RMC | Up | NDD373 | 96 | 4.37E-18 | *MED13, NIPBL, SMC3, CNOT1, KMT2E, UBTF, TRRAP, USP7, ASH1L, KMT2A, CREBBP, KAT6A, SIN3A, CHD2, EHMT1, TLE3, UBR5, CHD7, KMT5B, HIVEP2, CHD8, WAC, ZNF462, NSD1, ARID2, FBXO11, TAB2, KMT2D, ZNF148, RFX3, ZEB2, SETBP1, TLK2, RAI1, KIF5C, SON, STAG1, POGZ, ZMYND11, KIF11, KMT2C, ADNP, COL4A1, KAT6B, SCN8A, ZMYND8, PHF12, ZFHX3, SHOC2, TRIO, SOS1, ZNF292, SETD1B, TANC2, ARIH1, SOX9, ACVR2A, TNRC6B, MEIS2, CTNND1, YY1, ASXL3, PDE4D, SETD2, MAP3K7, DYRK1A, WDR48, SRP54, SPAST, SETD5, PIK3R1, SNAP25, VEZF1, KLF7, ZMIZ1, AUTS2, NRXN1, PPP2CA, CHAMP1, TCF7L2, TAOK1, ZBTB10, NF1, SATB1, CDK13, MAP2K1, MYCN, SYT1, TRPM3, MSL2, ABCC9, MED23, KLHL20, KDM5B, PPM1D, DNMT3A* |
| eFT508 | Up | NDD373 | 69 | 1.48E-10 | *MED13, NIPBL, SMC3, KMT2E, ASH1L, KMT2A, KAT6A, SIN3A, CHD2, SPEN, UBR5, KMT5B, HIVEP2, WAC, ZNF462, ARID2, FBXO11, TAB2, KMT2D, ZNF148, RFX3, ZEB2, KIF5C, STAG1, ZMYND11, KIF11, KMT2C, KAT6B, SCN8A, RFX7, ZFHX3, SHOC2, TRIO, SOS1, ZNF292, TANC2, KIF2A, ARIH1, ACVR2A, MEIS2, ASXL3, PDE4D, SETD2, BRAF, MAP3K7, WDR48, SRP54, SPAST, PIK3R1, SNAP25, VEZF1, ZMIZ1, AUTS2, NRXN1, CHAMP1, TCF7L2, TAOK1, ATF2, ZBTB10, NF1, SATB1, CDK13, GABRB3, SYT1, ABCC9, DNM1L, KLHL20, TNPO3, MIB1* |
| n/a | Down (baseline) | EPI | 22 | 5.82E-03 | *KMT2E, DHX57, TTF1, KDM5B, TTI1, PCDH19, NCOA1, CDKAL1, KANSL1L, DYRK1A, CHD9, ORC2, FAM184A, TUT7, BIRC6, ALKBH8, SETD1B, ANK1, SLC6A16, BPTF, ZSCAN26, PMFBP1* |
| RMC | Up | EPI | 19 | 6.97E-05 | *KMT2E, DHX57, KDM5B, TTI1, PCDH19, NCOA1, KANSL1L, DYRK1A, CHD9, ORC2, FAM184A, TUT7, BIRC6, ALKBH8, SETD1B, SLC6A16, BPTF, ZSCAN26, PMFBP1* |
| eFT508 | Up | EPI | 11 | 3.32E-02 | *KMT2E, PCDH19, NCOA1, CHD9, FAM184A, TUT7, BIRC6, ALKBH8, SLC6A16, BPTF, PMFBP1* |

**Table 1. Gene set enrichment analysis of translationally downregulated genes in *TSC2*-Null NPCs related to ASD, NDD and epilepsy.** *Drug Tx, treatment: DMSO, 10 nM RMC-6272 or 50 nM eFT508; pV, p-value; ASD72, 72 gene variants (false discovery rate (FDR)<0.001) identified in patients with autism spectrum disorder; NDD373, 373 gene variants (FDR<0.001) identified in patients with neurodevelopmental disorders [37]; EPI83, 83 gene variants identified in full epilepsy cohort [41].

# Figure 1

# Figure 2

# Figure 3

# Figure 4

# Figure 5

# Figure 6

**A**

ASD72  NDD373

*TSC2-/-* vs WT

*TSC2-/-* + RMC-6272 vs DMSO

*TSC2-/-* + eFT-508 vs DMSO

**B**

28 RMC responsive
ASD72 genes

18 eFT responsive
ASD72 genes

*ADNP, CHD8,
DNMT3A,
DYRK1A,
EHMT1, KDM5B,
MED23, POGZ,
SETD5, TLK2*

*ASH1L, ASXL3,
AUTS2, CHD2,
KMT2A, KMT2C,
SUV420H1, MED13,
MEIS2, NF1, NIPBL,
NRXN1, RFX3,
SPAST, TAOK1,
TCF7L2, VEZF1,
WAC*

**C**

ASD/NDD-associated genes

eIF4E-sensitive genes

☐ *TSC2*-WT, Vehicle    ▨ *TSC2*-Null, RMC-6272
■ *TSC2*-Null, Vehicle    ▦ *TSC2*-Null, eFT-508