Individual differences in uncertainty evaluation explain opposing exploratory behaviors in anxiety and apathy

Xinyuan Yan [1], R. Becket Ebitz [3], Nicola Grissom [2], David P. Darrow [4†], Alexander B. Herman [1†*]

[1]Department of Psychiatry and Behavioral Sciences, University of Minnesota, Minneapolis, MN 55455, USA

[2]Department of Psychology, University of Minnesota, 75 E River Rd, Minneapolis, MN 55455, USA

[3]Department of Neuroscience, Universite de Montreal, 2900 Edouard Montpetit Blvd, Montreal, Quebec H3T 1J4, Canada

[4]Department of Neurosurgery, University of Minnesota, Minneapolis, MN 55455, USA

* Corresponding author: Alexander B. Herman

† Denotes equal contribution

Email: herma686@umn.edu

Author Contributions: Conceptualization, X.Y, A.B.H, R.B.E, N.G, D.P.D; Methodology, X.Y, R.B.E, A.B.H; Investigation, D.P.D, A.B.H; Formal Analysis, X.Y; Writing – Original Draft, X.Y; Writing - Review & Editing, X.Y, D.P.D, A.B.H; Funding Acquisition, A.B.H and N.G; Supervision, A.B.H, D.P.D.

Competing Interest Statement: The authors declare no competing interests.

This PDF file includes:

Main Text
Figures 1 to 5
Tables 0

Abstract

Navigating uncertain environments is a fundamental challenge for adaptive behavior, and affective states such as anxiety and apathy can profoundly influence an individual's response to uncertainty. Uncertainty encompasses both volatility and stochasticity, where volatility refers to how rapidly the environment changes and stochasticity describes outcomes resulting from random chance. This study investigates how anxiety and apathy modulate perceptions of environmental volatility and stochasticity and how these perceptions impact exploratory behavior. In a large online sample (N = 1001), participants completed a restless three-armed bandit task, and their choices were analyzed using latent state models to quantify the computational processes. We found that anxious individuals attributed uncertainty more to environmental volatility than stochasticity, leading to *increased* exploration, particularly after reward omission. Conversely, apathetic individuals perceived uncertainty as more stochastic than volatile, resulting in *decreased* exploration. The ratio of perceived volatility to stochasticity mediated the relationship between anxiety and exploratory behavior following adverse outcomes. These findings reveal distinct computational mechanisms underlying anxiety and apathy in uncertain environments. Our results provide a novel framework for understanding the cognitive and affective processes driving adaptive and potentially maladaptive behaviors under uncertainty, with implications for the characterization and treatment of neuropsychiatric disorders.

Keywords: apathy, anxiety, uncertainty, volatility, stochasticity, exploration, multi-armed bandit task

Introduction

Life is filled with unexpected challenges. Adapting to an ever-changing environment requires a balance between exploring the unknown and exploiting the known. How individuals interpret the cause of undesirable outcomes (e.g., failure in investment, career plateau, bad weather) in an uncertain environment shapes their subsequent actions [1]. When changes in outcomes are attributed to how quickly the environment is changing (the volatility), individuals may be motivated to explore more to acquire additional information and adapt their behavior. In contrast, attributing adverse outcomes to mere chance or 'bad luck' (i.e., stochasticity) may reduce the motivation to explore, leading some individuals to stick with existing strategies [2].

How individuals perceive and respond to environmental uncertainty can vary significantly, and these differences may be influenced by underlying affective states [3]. Apathy, characterized by a lack of motivation and goal-directed behavior [4,5], is an affective state associated with imprecise beliefs about action outcomes [6] and a tendency to persist with previous choices rather than explore [7]. This suggests that apathetic individuals may view outcomes as primarily stochastic, attributing events more to chance than controllable variables. This bias could discourage exploration and potentially reinforce a cycle of failure and helplessness [8].

In contrast, anxiety, marked by excessive worry and a heightened perception of potential threats [9,10] and uncertainty [11], may lead individuals to overestimate environmental volatility. Consequently, anxious individuals could be driven to seek new information to update their beliefs and reduce uncertainty [12]. However, research on the link between anxiety and exploration has yielded mixed findings,

with some studies showing increased exploration to mitigate uncertainty [13,14] and

others showing reduced exploration to avoid unpredictable feedback under high

anxiety [15,16]. Notably, apathy and anxiety often coexist in clinical populations, such

as Alzheimer's [17], Parkinson's disease [18], and depression [19,20], despite having

distinct neural representations [21,22].

Building on these findings, we propose three fundamental questions to further

elucidate the relationship between affective states and decision-making under

uncertainty. First, we aim to investigate whether apathy and anxiety exhibit distinct

behavioral patterns when individuals are faced with uncertain situations. Second,

we seek to examine how individual differences in levels of apathy and anxiety are

associated with perceptions of different types of uncertainty, specifically volatility

and stochasticity. Finally, we intend to explore how perceived volatility influences

exploratory behavior during decision-making processes.

We posit two competing hypotheses:

(1) Both apathetic and anxious individuals engage in less exploratory behavior but

through different computational mechanisms. Apathetic individuals weigh

stochasticity over volatility and explore less, while anxious individuals

overestimate volatility, leading to a sense that their actions cannot track or learn

from the environment, ultimately leading to exploitation. This may provide a

computational account for learned helplessness [23] and the co-occurrence of apathy

and anxiety in various clinical populations, such as Parkinson's and Alzheimer's

diseases.

(2) Apathetic individuals manifest less exploration, while anxious individuals

engage in more exploration. Apathetic individuals weigh stochasticity over

volatility and explore less, while anxious individuals overestimate volatility but

explore more to reduce their uncertainty. This result would be consistent with

previous findings suggesting that the two affective states have distinct neural

substrates [21,24].

To address these questions, we employed a restless three-armed bandit task (Figure

1A), a well-established paradigm for capturing adaptive learning in volatile

environments [25,26]. We adopted the Hidden Markov Model (HMM) to obtain the

likelihood of individuals switching between exploitation and exploration states

[27,28]. To further investigate how volatility and stochasticity modulate exploration,

we utilized a Kalman filter model, which can dissociate two distinct sources of

noise, volatility (process noise variance) and stochasticity (observation noise

variance), during inference [29]. Together, these methods offer a comprehensive view

of the cognitive mechanisms underlying exploratory behavior and the

manifestation of anxiety and apathy.

Results

We recruited a large gender-balanced online sample consisting of 1,001 adults. The participants, ranging in age from 18 to 54 years (mean $\pm$ SD = 28.446 $\pm$ 10.354 years; gender: 493 female), performed a restless three-armed bandit task, as depicted in Figure 1A. During this task, participants selected among three playing card images, with each card representing a different option. They made their selections by moving their cursor over their chosen card. The probability of receiving a reward from each card deck varied randomly over time. After each choice, feedback was displayed on the screen indicating whether a reward was received. Participants also completed symptom surveys assessing levels of anxiety and apathy (details in Methods and SI Section 1, Table S1). We defined the trial as a *switch* trial if the chosen option was different from the last trial, and a *stay* trial if the choice was the same as the last trial.

Figure 1. Three-armed restless bandit task and distinct behavioral patterns associated with apathy and anxiety.

(A) Three-armed restless bandit task. Participants chose one option from among the three targets to receive reward or non-reward feedback. Each target was associated with a hidden reward probability that randomly and independently changed throughout the task. The lower panel indicates the example choice and reward sequence and the definition of stay and switch. Specifically, stay was defined as choosing the same target as in the previous trial, while switch was defined as choosing a different target. "+1" denotes reward feedback, and "+0" denotes reward omission.

(B) Most participants earned more rewards than expected by chance

(C) Apathy and anxiety correlated positively.

(D) Apathy correlated negatively with switch behaviors, while anxiety correlated

positively with switch behaviors. Anxious individuals were more sensitive to

undesired feedback (no reward) and exhibited more switch behaviors compared to

reward feedback.

(Panels in Figure 1C and 1D utilize binned correlation plots [25 quantile bins based

on the x-axis], with lines representing the standard error (S.E.). N.B. that these may

be smaller than the symbol. Statistical analyses were performed on raw data.)

Apathy and anxiety predicted distinct exploratory behaviors

We first evaluated the performance by comparing the total number of rewarded

trials each participant experienced against the number expected by chance. Out of

the 1001 participants, 985 accrued more rewarded trials than would be statistically

expected by chance, suggesting significant effectiveness in their decision-making

strategies (Figure 1B). As expected, anxiety and apathy showed a significant

positive correlation (r = 0.35, $p<10^{-29}$, Figure 1C), which is consistent with

previous findings on their co-occurrence [17].

To investigate the relationship between apathy and the percentage of switch

behaviors (P(switch)), as well as anxiety and P(switch), we conducted partial

correlations between apathy and exploration while controlling for anxiety, and

between anxiety and exploration while controlling for apathy. We found that

apathy negatively predicted P(switch) (r = -0.16, $p<0.001$) regardless of feedback

type (reward or no-reward), while anxiety positively correlated with P(switch) (r =

0.13, $p<0.001$). Intriguingly, the relationship between anxiety and switch behaviors

was greater after non-reward feedback (r = 0.16, $p<0.001$) compared to reward

feedback (r = 0.07, $p=0.024$) (their difference, z-score = 2.40, $p=0.01$). Though co-

existing in this population, these two affective states predicted distinct switch

behaviors under uncertainty (Figure 1D). The stronger relationship between

anxiety and P(switch) after undesirable feedback indicates that highly anxious

individuals are more sensitive to negative feedback, which may lead them to

disengage.

Next, we fitted the behavior with a Hidden Markov Model (HMM) to decode the

hidden states, "*explore*", and "*exploit*" (Figure 2A) [26,27,30,31]. Each arm is associated

with a hidden reward probability that randomly and independently changes

throughout the task (Figure 2A). In our study, exploration and exploitation states

are considered hidden states underlying the observed choices, such as switching

between decks or repeatedly choosing from the same deck. We calculated the

percentage of explore states, i.e., P(explore). Consistently, apathy correlated

negatively with P(explore) (r = -0.17, $p<0.001$), while anxiety positively correlated

with (P(explore)) (r = 0.11, $p=0.003$) as well as the percentage of exploration after

reward omission(P(explore|0) (r = 0.13, $p<0.001$) (Figure 2C).

The hidden states associated with each observed behavior is important but not

sufficient for fully understanding the dynamic decision-making process. It's also

necessary to examine how these hidden states transition between each other.

Therefore, we investigated how apathy and anxiety manifest in the transition

probability (Figure 2B) between explore and exploit. As predicted, apathy had a

positive correlation with the transition probability from *explore* to *exploit* (r = 0.13,

$p<0.001$) but a negative correlation with the transition probability from *exploit* to

explore (r = -0.08, $p=0.011$). In contrast, anxiety had a negative correlation with

the transition probability from *explore* to *exploit* (r = -0.07, $p=0.035$) but a positive

correlation with the transition probability from *exploit* to *explore* (r = 0.07,

$p<0.022$) (Figure 2D). All significant results reported in the study survived False

Discovery Rate (FDR, $p<0.05$) correction.

Figure 2. Apathy and anxiety have opposing relationships with exploration and explore and exploit state dynamics.

(A) Unrolled structure of the hidden Markov model (HMM) used to infer the explore and exploit states' underlying behavior.

(B) The transition probabilities within and between states in the HMM.

(C) The probability of exploration, plotted as a function of apathy (top left) and anxiety (top right). The probability of exploration following a reward omission

is plotted as a function of apathy (bottom left) and anxiety (bottom right).

(D) The transition probability from explore to exploit, plotted as a function of

apathy (top left) and anxiety (top right); the transition probability from exploit

to explore plotted as a function of apathy (bottom left) and anxiety (bottom

right).

(Panels in Figure 2C and 2D utilize binned correlation plots [25 quantile bins

based on the x-axis], with lines representing the standard error (S.E.). N.B. that

these may be smaller than the symbol. Statistical analyses were performed on

raw data). $* p < 0.05$, $** p < 0.01$, $*** p < 0.001$. All p-values corrected for

multiple comparisons.

Apathy and anxiety are associated with distinct computational processes

underlying exploration.

We then asked whether differing perceptions of the environment might explain the

distinct patterns of exploration predicted by apathy and anxiety we observed.

To address this question, we utilized a Kalman filter model (Figure 3A), which can

dissociate sources of uncertainty into perceived volatility and stochasticity [29].

Kalman filter (KF) models have been widely applied in psychology and

neuroscience to study various aspects of learning and decision-making [32,33] (for

more detailed information about the model, please refer to the Method section).

We also fitted the behavioral data to alternative models including volatile Kalman

filter (VKF) [29], Rescorla-Wagner models single (RW1) [34] and dual learning rates

(RW2) to weigh positive and negative learning rates [35]. We employed Hierarchical

Bayesian inference (HBI) to fit models to choice data. Further, we used Bayesian

model selection (BMS) and protected exceedance probability (PXP) to select the

winning model (Figure 3B). The Kalman filter served as the best model for our

population, and we examined the resulting distribution of volatility and

stochasticity (Figure 3B).

We first conducted correlation analyses using all data points (Figure 3C).

Specifically, we found that apathy was positively correlated with stochasticity ($r =$

0.08, p=0.013) but negatively correlated with volatility ($r = $ -0.08, p=0.008).

Conversely, anxiety showed a negative correlation with stochasticity ($r = $ -0.12,

p=0.001) and a positive correlation with volatility ($r = $ 0.12, p=0.002). These

correlations highlight the distinct cognitive biases associated with apathy and

anxiety in processing environmental uncertainties.

To clearly illustrate and confirm the findings, we categorized participants into

distinct groups based on their apathy and anxiety levels. For apathy, we identified

the high apathy group ($N = 223$) as those scoring in the top 25% on the Apathy

Motivation Index (AMI), which assesses apathy in behavioral and social domains

[36]. Conversely, the low apathy group (N = 251) comprised individuals scoring in

the bottom 25% of apathy scores. Similarly, for anxiety, the high anxiety group (N

= 228) included participants within the top 25% of scores on the GAD-7 scale [37],

while the low anxiety group (N = 250) consisted of those in the bottom 25%. These

classifications allowed for a direct comparison of behaviors and traits between

individuals with varying degrees of apathy and anxiety.

We conducted linear regression analyses using volatility and stochasticity as the

dependent variables with the high versus low anxiety and apathy groups as

predictors. The Methods section provides details of the regression model

specifications.

As hypothesized, apathetic individuals overestimated stochasticity ($t(471) = 3.06$,

$p=0.002$) and underestimated the volatility compared to those with low apathy

($t(471) = -3.24$, $p=0.001$)(Figure 3D). Consequently, apathetic individuals

exhibited a lower learning rate than their low apathy counterparts ($t(471) = -3.11$,

$p=0.002$).

In contrast, individuals with high anxiety levels tended to overestimate volatility

($t(475) = 2.84$, $p=0.004$) and underestimate stochasticity compared to those with

low anxiety ($t(475) = -3.04$, $p=0.002$), resulting in a higher learning rate among the

high anxiety group ($t(475) = 3.21$, $p=0.001$) (Figure 3E). Furthermore,

comparisons showed that anxious individuals had higher volatility estimates than

those with high apathy ($t(449) = 2.75$, $p=0.006$), whereas apathetic individuals had

higher stochasticity estimates than their anxiety counterparts ($t(449) = -3.01$,

$p=0.002$) (SI Section 2, Figure S1).

Figure 3. Apathy and anxiety have opposing relationships with volatility and stochasticity.

(A) The schematic of the Kalman filter model used in our analysis. The diagram illustrates how this model can differentiate between volatility (process noise variance) and stochasticity (observation noise variance), providing insights into the underlying decision-making processes.

(B) Bayesian model comparison and the distribution of volatility, stochasticity

(C) Correlation plots that visually represent the relationships between apathy, anxiety, and the distinct computational processes of volatility and stochasticity.

(D) Highly apathetic individuals overestimated stochasticity but underestimated the volatility, resulting in a lower learning rate.

(E) In contrast, highly anxious individuals overestimated volatility but underestimated stochasticity, resulting in a higher learning rate.

(Panels in Figure 3C utilize binned correlation plots [25 quantile bins based on the x-axis], with lines representing the standard error (S.E.). N.B. that these may be smaller than the symbol. Statistical analyses were performed on raw data). $*p < 0.05$, $** p < 0.01$, $*** p < 0.001$. All p-values corrected for multiple comparisons).

The ratio of volatility to stochasticity distinguished apathy and anxiety

To clarify the differential impacts of apathy and anxiety on decision-making under uncertainty, we computed the ratio of volatility to stochasticity, $/s$, to represent the balance between these two types of uncertainties. A higher v/s indicates a perception of greater volatility relative to stochasticity, while a lower ratio suggests a perception of more stochasticity relative to volatility. We applied a logarithmic transformation to the ratio to manage extreme values (e.g. cases where individuals might perceive very high volatility but very low stochasticity).

Consistently, our findings reveal a clear distinction: v/s correlated negatively with

apathy (r = -0.08, p=0.010) but positively with anxiety (r = 0.13, p<0.001) (Figure

4A).

The ratio of volatility and stochasticity mediated the relationship between anxiety

and the exploration after negative feedback

To determine whether individual differences in the perception of uncertainty

explain the relationship between exploratory behavior and affect, we conducted a

mediation analysis with anxiety, switching after reward omission (P(switch | 0)),

and v/s.The results demonstrate that the relationship between anxiety and the

tendency to switch after receiving no reward is significantly mediated by the v/s

(Figure 4B). This mediation was also significant for the analogous HMM model-

based measures (see SI Section 3, Figure S2). No significant mediation effect was

found for apathy, however, reinforcing the unique pathways through which anxiety

influences exploratory behavior. These results explain why individuals with higher

anxiety might explore more after negative feedback, driven by an overweighting of

perceived volatility relative to stochasticity as a strategy to reduce uncertainty and

manage risks.

Figure 4. Distinctions in apathy and anxiety on the ratio of volatility to stochasticity and its mediation effect.

(A) The ratio of volatility to stochasticity, plotted as a function of apathy (left) and anxiety positively (right).

(B) Mediation analysis, showing the mediating effect of the ratio of volatility to stochasticity on the relationship between anxiety and switch behavior after reward omission.

(Panels in Figure 4A utilize binned correlation plots [25 quantile bins based on the x-axis], with lines representing the standard error (S.E.). N.B. that these may be smaller than the symbol. Statistical analyses were performed on raw data). $* p <$ 0.05, $** p <$0.01, $*** p < 0.001$. All p-values corrected for multiple comparisons).

A low dimensional manifold unifies exploration, perceptions of uncertainty and affective state.

The HMM state-model of exploration-exploitation and the Kalman filter process model of uncertainty estimation represent complementary ways of understanding adaptive behavior that our mediation results suggest are intrinsically related. We hypothesized that a latent structure underlying adaptive behavior on this task might unify these descriptions of behavior. We utilized advanced dimensionality reduction methods to uncover such a latent structure in the raw task behavior. First, we formatted each participant's trial-by-trial task data into sequences of choices to stay (repeat the choice on the last trial) or switch (choose a different option) and reward outcome for two consecutive trials ({choice$_{t-1}$, outcome$_{t-1}$, choice$_t$}, Figure 5A and 5B). The behavioral data for each participant was then transformed into counts for each of these eight unique sequences. Then we applied Uniform Manifold Approximation and Projection (UMAP) [38], a computationally efficient algorithm that can preserve both the local and global distances between

data points in high-dimensional space, to learn the two-dimensional manifold

underlying the eight-dimensional behavioral data (Figure 5C, see Methods for

more algorithm details). Including additional reward history and applying other

dimensionality reduction methods like principle component analysis (PCA), and t-

distributed Stochastic Neighbor Embedding (t-SNE) did not change the results (SI

Section 4, Figure S3, Table S2).

Our analysis using UMAP revealed distinct correlations within the derived

dimensions. Specifically, the dimension 1 score (the horizontal axis) exhibited a

very strong significant negative correlation with exploratory behavior (P(explore)

($r = -0.90$, $p < 10^{-200}$), but it showed no significant relationship with the ratio of

volatility to stochasticity (v/s)($r = 0.01$, $p = 0.74$) (Figure 5D, 5E). In contrast, the

dimension 2 score (the vertical axis) demonstrated a strong negative correlation

with v/s ($r = -0.76$, $p < 10^{-185}$), which was significantly more pronounced than its

correlation with P(explore) ($r = -0.19$, $p < 10^{-10}$) (Figure 5F, 5G). This suggests that

dimension 1 primarily represents exploratory behavior, while dimension 2

primarily reflects the computational factors: volatility and stochasticity. (i.e.,

volatility and stochasticity).

Further, both dimensions also showed correlations with affective states: the

dimension 1 score was positively correlated with apathy ($r = 0.14$, $p < 0.001$), and

negatively correlated with anxiety ($r = -0.11$, $p < 0.001$). Similarly, the dimension 2

score had a positive correlation with apathy ($r = 0.097$, $p = 0.002$) and a negative

correlation with anxiety ($r = -0.088$, $p = 0.004$).

It is worth noting that we only found linear relationship between apathy, anxiety,

and exploration, as well as between these affective states and the ratio of volatility

to stochasticity (our analysis using higher order effects among these variables did

not yield significant results, more details can be found in SI Section 5, Table S3).

To delve deeper into how these factors interact in the low-dimensional space

defined by UMAP, we divided the data manifold into two groups based on the

dimension1 score: a monotonically decreasing group (left part, dimension 1 score

< -0.671, N=390) and a monotonically increasing group (right part, dimension 1

score> -0.671, N=611). The methodology used to identify the turning point

(dimension 1 score = -0.671) that differentiates the monotonically decreasing

group from the monotonically increasing group is detailed in the Methods section.

The analysis revealed that the monotonically decreasing group had relatively

higher levels of anxiety compared to the monotonically increasing group

(t(999)=2.08, p =0.037), while their apathy levels were significantly lower (t(999)=

-3.56, p=0.0003). This segmentation allows us to further explore and understand

the complex interplay between affective states, computational parameters, and

exploratory behaviors within a structured, low-dimensional framework.

Notably, individuals in the left monotonically decreasing group, characterized by

high anxiety and low apathy, generally perceive higher volatility relative to

stochasticity and demonstrate greater exploratory behavior compared to those in

the right monotonically increasing group. Interestingly, within the decreasing

group, higher perceived volatility correlates with reduced exploration. Conversely,

in the increasing group, an increased perception of volatility tends to stimulate

more exploratory actions. These results suggest that while severe anxiety might

suppress exploration due to overwhelming uncertainty, moderate anxiety in the

general population can promote exploration as a coping mechanism to gather

information and reduce anxiety symptoms.

We now address our final research question; what is the relationship between

volatility and exploratory behavior? Considering the parabolic relationship

between the manifold dimension reflecting exploration and the dimension

representing v/s, we predicted that the relationship between exploration and v/s

might be quadratic.

To test this hypothesis, we constructed a regression model as follows:

P(explore) ~ $v/s + v/s^2$ + anxiety + apathy

The results revealed that both the linear and quadratic terms are significant (linear

term, coefficient = 0.02, SE = 0.003, t(996)=5.83, $p<10^{-9}$; quadratic term,

coefficient = 0.005, SE = 7.86×10^{-4}, t(996)=6.59, $p<10^{-9}$), indicating a complex,

non-linear relationship between the ratio of volatility to stochasticity and

exploration (see SI Section 6, Figure S4), which was consistent with the manifold

representation.

Figure 5. Visualizing the complex relationships in decision-making through low-dimensional space.

(A) All possible sequences of choices and rewards that participants could make during the experiment

(B) The frequency distribution of individual decision-making patterns. The black line in the box plot represents each pattern's mean value, highlighting participants' typical behaviors.

(C) Schematic high-dimensional space of participants' decision-making pattern

(D) The two-dimensional space representation of exploration by using the Uniform Manifold Approximation and Projection (UMAP) (Different dimensionality reduction methods such as principal component analysis (PCA), and t-distributed Stochastic Neighbor Embedding (t-SNE) lead to a similar space)

(E) Dimension 1 exclusively represents P(explore) but does not represent the ratio of volatility to stochasticity

(F) The two-dimensional space representation of the ratio of volatility to stochasticity behavior by using UMAP (Different dimensionality reduction methods such as principal component analysis (PCA), and t-distributed Stochastic Neighbor Embedding (t-SNE) lead to a similar space).

(G) Dimension 2 mainly represents the ratio of volatility to stochasticity but not P(explore).

(H) The manifold has been separately dissociated into the monotonically decreasing group (the most left panel) and monotonically increasing group (the most right panel). The monotonically decreasing group was associated with a relatively higher anxiety level than the monotonically increasing group, while the apathy level was significantly lower than the monotonically increasing group. Within the monotonically decreasing group (left part), a higher volatility to stochasticity ratio leads to decreased exploration. In contrast, within the

446 monotonically increasing group (right part), a higher volatility to stochasticity ratio

447 encouraged higher exploration. This exploration serves as a coping strategy to

448 relieve anxious feelings in the environment.

Discussion

We found that apathy and anxiety predicted opposing patterns of exploratory

behavior, which were explained partly by differing perceptions of uncertainty.

Anxiety was associated with increased exploration after reward omission and

greater volatility estimation: the attribution of uncertainty to a rapidly changing

(but still learnable) environment. Apathy, in contrast, predicted decreased

exploration and higher stochasticity estimation: the perception of uncontrollable

randomness. Following a dimensionality reduction of the raw behavioral data,

exploration and perceptions of uncertainty emerged as the dimensions of an

underlying latent structure that unified the different model approaches and the

affective states. These findings elucidate the complex interplay between cognitive

assessments of uncertainty, affective states, and decision-making processes,

offering several key insights into adaptive and maladaptive behaviors under

uncertainty.

The distinct patterns of exploratory behavior observed in anxious and apathetic

individuals highlight the role of affective states in shaping responses to

uncertainty. Anxious individuals, who generally display a heightened sensitivity to

potential threats and environmental changes, exhibited a bias toward perceiving

greater volatility and exploring more after negative outcomes. Our mediation

analysis revealed that the perception of volatility relative to stochasticity partially

mediates the relationship between anxiety and exploratory behavior after reward

omission. This finding is consistent with previous results [39] and offers a

mechanistic explanation for why anxious individuals in a healthy population might

choose to explore more after receiving negative feedback. The perceived

overweighting of volatility relative to stochasticity may drive these individuals to

seek more information, potentially as a strategy to reduce uncertainty and manage

perceived risks more effectively [40]. Although such a strategy may be beneficial for

adaptation in genuinely volatile environments, it may also contribute to excessive

worry and stress, especially if the perceived level of volatility exceeds actual

environmental volatility [9,11]. Consequently, anxious individuals may find

themselves in a prolonged state of heightened arousal and uncertainty, leading to

suboptimal decision-making and diminished well-being.

On the other hand, apathetic individuals, who generally exhibit diminished

motivation and responsiveness [41], tended to attribute outcomes more to

stochasticity in our study. This perception might underlie their reduced exploratory

behavior, reflecting a disengagement from active learning and adaptation. If

outcomes seem random and beyond our control, expending energy to explore may

seem futile, and focusing on what we know seems rational. While this approach

may conserve energy, this inflexibility can perpetuate a cycle of disengagement

and maintain apathetic symptoms [42,43]. Apathetic individuals may fail to recognize

the potential benefits of exploration and remain stuck in suboptimal decision-

making patterns, further reinforcing their disengagement from the environment [4].

The dimensionality reduction of the behavioral sequence data using UMAP

allowed us to examine the relationship between exploration and the estimation of

volatility and stochasticity. Despite the intuitive connection between these two

behavior models, their relationship has not been directly examined. Our results

showed that exploration and uncertainty estimation related closely to the two axes

of a parabolic latent structure of adaptive behavior. As a result, both model-based

metrics were necessary to characterize the spectrum of individual differences fully.

Segmenting the data on the manifold further illuminated the fine-grained interplay

between affective states and exploratory behavior. Individuals with relatively

higher anxiety and lower apathy (the monotonically decreasing group) generally

weighted volatility more and demonstrated greater exploratory behavior compared

to those with lower anxiety and higher apathy (the monotonically increasing

group). However, *within* these groups formed on the manifold, individuals

exhibited opposing relationships between uncertainty and exploration. In the

higher anxiety group, perceived volatility correlates inversely with exploration.

However, in the lower anxiety group, increased volatility perception predicts

greater exploration.

These results reconcile previously inconsistent findings regarding exploratory

behavior in individuals with anxiety, with some studies showing more exploitative

behavior [15,16], and others finding that higher anxiety predicts more exploratory

behaviors [13,14]. The relationship between perceived volatility and exploration is

modulated by the degree of anxiety, with more severe anxiety potentially

suppressing exploration as a form of avoidance. In contrast, milder anxiety may

promote exploration as a coping mechanism to gather more information reduce

uncertainty, and alleviate discomfort. This dual response to perceived volatility

underscores the complex interplay between anxiety levels, environmental

perceptions, and behavioral strategies in managing emotional responses.

Our results have implications for behavioral approaches aimed at improving

adaptive functioning and mental health. Cognitive-behavioral therapies that focus

on challenging and restructuring interpretations of beliefs about uncertainty and

controllability could be personalized by symptom profile [44,45]. For anxious

individuals, interventions that help recalibrate perceptions of volatility and equip

them with coping strategies for managing uncertainty may reduce maladaptive

worry and enhance decision-making efficacy [46,47]. Furthermore, encouraging

anxious individuals to integrate information over longer horizons when undesirable

things happen, rather than attempting to change strategies to these events rapidly,

may prove a beneficial approach to managing anxiety [39]. For apathetic individuals,

strategies aimed at strengthening the locus of control and highlighting the efficacy

of their actions may counteract tendencies towards stochasticity attribution.

Training individuals to view the world as more controllable may be a practical

addition to existing psychotherapeutic approaches, including Behavioral Activation

and Motivational Interviewing [48]. Challenging beliefs about uncontrollability may

promote a more balanced perception of the environment and encourage

exploration. Previous studies have demonstrated that individuals who believed they

received treatment or cognitive training that enhanced prosocial behaviors and

cognitive abilities actually exhibited more prosocial behaviors and improved

performance in standard IQ tests [49,50]. This suggests that manipulating participants'

beliefs about their environmental control through placebo interventions could

enhance their resilience in uncertain situations.

Another important clinical implication involves using an individual's position on

the behavioral manifold (Figure 5) to predict how their behavior might change in

response to treatment based on their symptoms. For example, a patient positioned

in the upper left quadrant before treatment may exhibit higher anxiety, lower

apathy, and increased exploratory behavior. During and after treatment, monitoring

these behavioral shifts may allow us to infer changes in their affective states or

symptoms based on their new manifold position. To develop such a tool, several

questions remain: Do changes within an individual follow a predictable trajectory

on this manifold? Do clinical populations conform to the same manifold, or do

they deviate, projecting into the larger, unoccupied areas of the manifold? The

answers to these questions could enhance the implementation of dimensional

approaches for individualized neuropsychiatric care [51].

Our results must be interpreted in light of notable limitations. First, the study

primarily utilized an online sample, which may not accurately represent the

demographic and clinical characteristics of populations with specific mental health

diagnoses. The potential differences in internet access, motivation, and the self-

report nature of online studies can introduce biases that may differ from clinical

settings. Consequently, the generalizability of our findings to clinical populations

remains to be determined. Second, our results, while robust statistically, are

inherently correlational, limiting our ability to infer causal relationships between

the affective states of apathy and anxiety and their impacts on decision-making

processes. The observed associations provide a strong foundation for

hypothesizing causal mechanisms but do not confirm them. Future studies may

examine clinical samples of conditions known to affect adaptive decision-making

under uncertainty, such as depression, anxiety disorders, and Parkinson's disease,

as well as interventions targeting the physiology of adaptive behavior.

Method

Ethics approval

The experimental procedures of all experiments were in line with the standards set

by the Declaration of Helsinki and were approved by the local Research Ethics

Committee of the University of Minnesota, Twin Cities. Participants provided

written informed consent after the experimental procedure had been fully

explained and were reminded of their right to withdraw at any time during the

study.

Participants

We recruited a sample of 1512 participants via Amazon Mechanical Turk (MTurk)

and Prolific (Prolific. co); exclusion criteria included current or history of

neurological and psychiatric disorders. 1001 participants completed all

questionnaires and the bandit task (age range 18-54, mean $\pm$ SD = 28.446 $\pm$ 10.354

years; gender, 493 female). All participants were compensated for their time in

accordance with minimum wage.

Questionnaire measurement

Participants' anxiety and apathy states were measured by the General Anxiety

Disorder Screener (GAD-7) [37], and the Apathy-Motivation Index [36], respectively.

More specifically, GAD-7 contains 7 items for assessing anxiety severity in the last

two weeks. All items were rated on a 4-point scale, with higher scores indicating

greater anxiety. Participants' apathy level was measured using the 18-item Apathy-

Motivation Index (AMI), which was designed to identify and measure general

apathy, as well as subtypes of apathy in behavioral, social, and emotional domains.

Higher scores on AMI represent greater apathy. For all questionnaire scores, see

SI, Section 1 & Table S1.

Three-armed restless bandit task

We assessed exploration-exploitation behavioral dynamics using a 300-trial three-

armed restless bandit task [27]. Participants were free to choose between three targets

for the potential to earn a reward of 1 point. Each target is associated with a hidden

reward probability that randomly and independently changes throughout the task.

We seeded each participant's reward probability walks randomly to prevent biases

due to particular kinds of environments. We assessed performance by comparing

the total number of rewarded trials to that expected by chance. Out of the 1001

participants, 985 accrued more rewarded trials than would be expected by chance.

Dimensionality reduction method

Popular and valid dimensionality reduction techniques to reveal manifolds include

t-distributed stochastic neighborhood embedding (t-SNE) [52], uniform manifold

approximation and projection (UMAP) [38], and Principal component analysis (PCA)

[53]. However, t-SNE suffers from limitations, including slow computation time and

loss of global data structure, and it is not a deterministic algorithm [54]. The main

drawback of PCA is that it is highly affected by outliers in the dataset [53]. In

contrast, UMAP is a deterministic and efficient algorithm, it also preserves both

local and global structure of original high-dimensional data. Uniform Manifold

Approximation and Projection (UMAP)

UMAP was implemented in the R language. The eight-dimensional datasets from

all participants were passed into the R package *umap*, version 0.2.8.0, available at

https://cran.r-project.org/web/packages/umap/) with default parameter setting as

n_component = 2, n_neighbors = 15, min_dist = 0.3, metric = 'Euclidean'. For

reproducibility reasons, we fixed the random_state in this algorithm. The

hyperparameter *n_neighbors* decide the radius of the search region. Larger values

will include more neighbors, thus forcing this algorithm to consider more global

structure of original n-dimension data. Another important hyper-parameter,

min_dist determines the allowed minimum distance apart for cases in lower-

dimensional space. *metric* defines the way that UMAP is used to measure distances

along the manifold.

Model-free analyses

We adopted some widely used model-free measures, including win-stay and lose-

shift [35,55] as the direct measurement for this learning task.

Win-stay. Win-stay is defined as the percentage of times that the choice in trial *t*-1

was repeated on trial *t* following a reward.

Lose-shift. In contrast, lose-shift equals the percentage of trials that the choice was

shifted or changed when the outcome of trial t-1 was non-reward.

Model free results can be found at SI Section 7, Table S4.

Mediation analyses

Mediation analysis is a statistical method used to examine the underlying

mechanisms by which an independent variable influences a dependent variable

through one or more mediator variables [56]. In our study, we employed the

bootstrapping method to estimate the mediation effect of volatility and

stochasticity on the relationship between affective states (apathy and anxiety) and

exploration. Bootstrapping is a nonparametric approach to effect-size estimation

and hypothesis testing that is increasingly recommended for many types of

analyses, including mediation [57]. This method involves repeatedly resampling from

the available data to generate an empirical approximation of the sampling

distribution of the indirect effect (i.e., the effect of the independent variable on the

dependent variable through the mediator). We used this distribution to calculate p-

values and construct confidence intervals based on 5,000 resamples. Bootstrapping

is preferred over other methods, such as the Sobel test because it does not assume

the normality of the sampling distribution and provides more accurate confidence

intervals that are bias-corrected and accelerated [56,57]. This approach offers a robust

and powerful way to test mediation hypotheses, particularly in cases where the

sample size is relatively small or the data violate assumptions of normality [58].

Hidden Markov Model

We fit a Hidden Markov Model (HMM) to the behavior, to decode the hidden state

of each trial for each participant. Fundamentally, the HMM has two layers, the

hidden layer (i.e., state) and the observable layer. The hidden dimension should

satisfy the Markov property. That is, the current hidden state only depends on the

previous state but not any past model history. The observable dimension entirely

depends on the current hidden states and is independent of other observations.

Parameters of the hidden Markov model can be represented as $\Omega \rightarrow (T, O, c)$.

Specifically, T is the transition probabilities matrix, O is the observation

probabilities matrix, or emissions matrix, and c refers to a vector with initial

probabilities for each hidden state. Here, we have two hidden states, an

"exploration" state and an "exploitation" state.

The transition probability between exploration and exploitation states can be

represented by:

681
$$T = \begin{pmatrix} t_{11} & \cdots & t_{1m} \\ \vdots & \ddots & \vdots \\ t_{m1} & \cdots & t_{mm} \end{pmatrix}$$

683
$$t_{explore,exploit} = P(q_k = s_{exploit}|q_{k-1} = s_{explore})$$

685 Where $t_{explore,exploit}$ refers to the transition probability from hidden state $s_{exploit}$

686 to another hidden state $s_{explore}$

687 k = time instant, m = state sequence length

689 Then matrix O represents the transition probabilities between hidden and

690 observable states.

692
$$O = \begin{pmatrix} o_{11} & \cdots & o_{1n} \\ \vdots & \ddots & \vdots \\ o_{m1} & \cdots & o_{mn} \end{pmatrix}$$

694
$$o_{explore,exploit} = P(\lambda_k = r_{exploit}|q_{k-1} = s_{explore})$$

k = time instant, n = observation sequence length

Both matrix O and T satisfy the principle that the sum along the rows must be

equal to 1.

c is an m-dimensional row vector that refers to the initial probability distribution.

In our current study, the initial probability was fixed and equal to the available

choices.

We fit HMM via expectation-maximization using the Baum-Welch algorithm and

decode hidden states from observed choice sequences by the Viterbi algorithm [59].

Model results of HMM can be found at SI Section 7, Table S5.

Kalman filter

The Kalman filter (KF) model has been widely applied in psychology and

neuroscience to study various aspects of learning and decision-making [32,33].

In the Kalman filter model for a multi-armed bandit task, *process noise* and

observation noise refer to two distinct sources of uncertainty that affect the

learning and decision-making process.

Process noise represents the uncertainty in the evolution of the hidden state

(reward mean) over time. It accounts for how the true state evolves from one point

in time to the next. In mathematical terms, process noise is part of the state

transition equation in the Kalman Filter:

717
$$x_t = x_{t-1} + e_t$$

x_t is the state at time t

e_t is the process noise t, which is assumed to be drawn from a normal distribution

with zeros mean and **process noise variance** v. Where the $e_t \sim N(0, v)$.

The process noise captures the idea that the reward-means for each arm can change

from one trial to the next, even in the absence of any observations. A higher

process noise variance v indicates a more volatile environment, where the reward

means are expected to change more rapidly.

In contrast, **observation noise** represents the uncertainty in the observed rewards,

given the current hidden state (reward mean). Which is assumed to be Gaussian

with zero mean and a fixed variance σ^2.

The observation noise captures the idea that the observed rewards are noisy and

can deviate from the true reward mean due to random fluctuations or measurement

errors.

A higher measurement noise variance indicates a more stochastic environment,

where the observed rewards are less reliable and informative about the underlying

reward means.

The Kalman Filter operates optimally when the statistical properties of the process

noise and the measurement noise are accurately known.

When observation noise variance (σ^2) is high relative to the process noise variance

(v), the Kalman gain will be small, and the model will rely more on its prior beliefs

and less on noisy observations. Conversely, when the observation noise variance

(v), is high relative to the process noise variance (σ^2), the Kalman gain will be

large, and the model will update its beliefs more strongly based on the observed

rewards.

Extended Kalman filter for three-armed bandit task

The Kalman filter model can be extended to capture the effects of both volatility

and stochasticity in a multi-armed bandit task [29,60].

In the current study, process noise variance (v) and observation noise variance (σ^2)

represent volatility and stochasticity, respectively.

A traditional assumption of the Kalman filter is that the process noise variance, v,

as well as the observation noise variance, σ^2 are constant.

Reward means update:

$$m_t = m_{t-1} + k_t(O_t - m_{t-1})$$

Where m_t is the estimated mean or value of the chosen arm at time t

and O_t is the observed reward at time t.

The mean update is driven by the prediction error, which is the difference between

the observed reward and the previous estimate.

Kalman gain is defined as:

$$k_t = (w_{t-1} + v)/(w_{t-1} + v + \sigma^2)$$

Here, k_t represents the Kalman gain or learning rate, which adjusts the weight given to new information based on the relative uncertainty of the prior estimate (w_{t-1}) and the total noise $(v + \sigma^2)$. When the stochasticity (σ^2) is high relative to the volatility (v), the Kalman gain (learning rate) will be small, and the model will rely more on its prior beliefs and less on the observations. Conversely, when the volatility (v), is high relative to the stochasticity (σ^2), the Kalman gain (learning rate) will be large, and the model will update its beliefs more strongly based on the observed rewards.

Variance update equation:

$$w_t = (1 - k_t)(w_{t-1} + v)$$

This equation updates the posterior variance (w_t), which represents the estimate's uncertainty after observing O_t.

Volatile Kalman filter for three-armed bandit task

The key difference between a standard Kalman filter and a volatile Kalman filter (VKF) is the variance of the process noise, a stochastic variable that changes with time. In other words, the VKF introduces parameters to handle the volatility in the process noise. Specifically, it allows the process noise variance v to vary with the observed prediction errors, reflecting changes in environmental volatility. Our approach here is essentially the same as that taken by Piray and Daw [29]. Here, we briefly described the model details as follows.

788 Kalman gain:

$$k_t = (w_t + v_{t-1})/(w_{t-1} + v_{t-1} + \sigma^2)$$

790 Update for the reward means:

$$m_t = m_{t-1} + k_t(O_t - m_{t-1})$$

794 Update for posterior variance w_t:

$$w_t = (1 - k_t)(w_{t-1} + v_{t-1})w_{t-1,t} = (1 - k_t)w_{t-1}$$

798 Update for volatility:

$$v_t = v_{t-1} + \lambda((m_t - m_{t-1})^2 + w_{t-1} + w_t - 2w_{t-1,t} - v_{t-1})$$

Rescorla-Wagner models

We also fitted the data to the classical Rescorla-Wagner model. Successful

adaptation in a dynamic situation requires the appropriate feedback-based learning

process where individuals integrate the feedback (reward or non-reward) into the

stimulus-outcome association [61]. The basic reinforcement learning model, the

Rescorla-Wagner model [34] can address this process well. So the first model (RW1)

was defined as:

$$v_t = v_{t-1} + a \times (R_{t-1} - v_{t-1})$$

where v_t is the value of the option on trial t.

a represents the general learning rate from feedback.

To verify whether participants employed distinct or shared computational

responses to positive and negative feedback, we built another model with two

learning rates, one for positive feedback and the other for negative feedback [35].

This model (RW2) can be defined as:

820
821
$$v_t = v_{t-1} + \alpha^{pos} \times (R_{t-1} - v_{t-1}), positive\ feedback$$
$$v_t = v_{t-1} + \alpha^{neg} \times (R_{t-1} - v_{t-1}), negative\ feedback$$

Where v_t is the value of the option on trial t. α^{pos} and α^{neg} represent the learning

rates from positive and negative feedback, respectively.

For these two models, $R_{t-1} \in \{0,1\}$ represents the feedback received in response

to participants' choice on trial t-1. And $R_{t-1} - v_{t-1}$ represents prediction error in

trial t-1.

We used a softmax choice function to map the value into choice. The softmax

function for these four models can be defined as:

833
$$P^t = \frac{exp^{(\beta V_{,t})}}{exp^{(\beta V_{t,1} + \beta V_{t,2} + \beta V_{t,3})}}$$

Where the β represents the inverse temperature with choice value.

Model fitting and comparison

Hierarchical Bayesian inference (HBI) is a powerful method for model fitting and

comparison in group studies [62]. Unlike traditional approaches such as maximum

likelihood estimation (MLE) or maximum a posteriori (MAP) estimation, which fit

models to each subject independently, HBI simultaneously fits models to all

subjects while constraining individual fits based on group-level statistics (i.e.,

empirical priors). This approach yields more robust and reliable parameter

estimates, particularly when individual subject data is noisy or limited.

In our study, we employed HBI to fit models to choice data. The method quantifies

group-level mean parameters and their corresponding hierarchical errors. To

ensure that parameter estimates remain within appropriate bounds during the fitting

process, we used the sigmoid function to transform parameters bounded in the unit

range or with an upper bound and the exponential function to transform parameters

bounded to positive values. The initial parameters of all models were obtained

using a MAP procedure, with the initial prior mean and variance for all parameters

set to 0 and 6.25, respectively, based on previous research [29]. This initial variance

allows parameters to vary widely without substantial influence from the prior.

For model comparison, we used Bayesian model selection, specifically employing

the protected exceedance probability (PXP) to select the winning model. The PXP

quantifies the probability that a given model is more frequent in the population

than all other models under consideration while accounting for the possibility that

the observed differences in model evidence may be due to chance [63]. The model

with the highest PXP is selected as the winning model. This approach inherently

penalizes model complexity, favoring models that balance goodness-of-fit and

parsimony. Model performance (log-likelihood) can be found in SI Section 8,

Table 6.

Turning point to divide the manifold into monotonically increasing and

decreasing group

To divide the manifold into monotonically increasing and decreasing groups, we

sorted the scores for dimension 1 in ascending order. Initially, we fitted a linear

model using the first three data points located on the upper left of the manifold. We

then expanded this model by sequentially including one additional data point from

dimension 1, continuing this process until we incorporated the last score (i.e., the

maximum dimension 1 score, situated on the upper right of the manifold).

Throughout this procedure, we monitored the t-statistic of the dimension1

coefficient to assess the statistical significance of dimension1 as a predictor.

Notably, a dimension 1 score of -0.671 marked the most significant negative

coefficient, after which the relationship between dimension 1 and dimension2

gradually shifted to become positive (see SI Section 9, Figure S5)

False discovery rate correction

We adopted FDR (False Discovery Rate) correction, which was introduced by

Benjamini and Hochberg (1995) [64] to control the expected proportion of false

positives (Type I errors).

Acknowledgments

This work was supported by NIMH under award #R21MH127607, NIDA under award #K23DA050909, and the University of Minnesota's MnDRIVE (Minnesota's Discovery, Research and Innovation Economy) initiative. Natural Sciences and Engineering Council of Canada (Discovery Grant RGPIN-2020-05577), the Research Corporation for Science Advancement & Frederick Gardner Cottrell Foundation (Project #29087), and a Research Fellowship from the Jacobs Foundation to R.B.E.

We thank A. David Redish, Cathy Chen, Seth D Koenig, and Brian Sweiss for helpful comments on the manuscript.

Code and data availability.

The data, and source code used to do data analysis, as well as generate main and supplementary Figures, are published at

Reference

1. Soltani, A. & Izquierdo, A. Adaptive learning under expected and unexpected uncertainty. *Nat. Rev. Neurosci.* **20**, 635–644 (2019).

2. Piray, P. & Daw, N. D. A model for learning based on the joint estimation of stochasticity and volatility. *Nat. Commun.* **12**, 6587 (2021).

3. Pulcu, E. & Browning, M. The Misestimation of Uncertainty in Affective Disorders. *Trends Cogn. Sci.* **23**, 865–875 (2019).

4. Husain, M. & Roiser, J. P. Neuroscience of apathy and anhedonia: a transdiagnostic approach. *Nat. Rev. Neurosci.* **19**, 470–484 (2018).

5. Marin, R. S. Apathy: Concept, Syndrome, Neural Mechanisms, and Treatment. *Semin. Clin. Neuropsychiatry* **1**, 304–314 (1996).

6. Hezemans, F. H., Wolpe, N. & Rowe, J. B. Apathy is associated with reduced precision of prior beliefs about action outcomes. *J. Exp. Psychol. Gen.* **149**, 1767–1777 (2020).

7. Scholl, J., Trier, H. A., Rushworth, M. F. S. & Kolling, N. The effect of apathy and compulsivity on planning and stopping in sequential decision-making. *PLoS Biol.* **20**, e3001566 (2022).

8. Huys, Q. J. M. & Dayan, P. A Bayesian formulation of behavioral control. *Cognition* **113**, 314–328 (2009).

9. Browning, M., Behrens, T. E., Jocham, G., O'Reilly, J. X. & Bishop, S. J.

Anxious individuals have difficulty learning the causal statistics of aversive

environments. *Nat. Neurosci.* **18**, 590–596 (2015).

10. Uhr, K. & Dugas, M. J. The role of fear of anxiety and intolerance of

uncertainty in worry: an experimental manipulation. *Behav Res Ther* **47**, 215–

223 (2009).

11. Gagne, C., Zika, O., Dayan, P. & Bishop, S. J. Impaired adaptation of learning

to contingency volatility in internalizing psychopathology. *Elife* **9**, e61387

(2020).

12. Hirsh, J. B., Mar, R. A. & Peterson, J. B. Psychological entropy: a framework

for understanding uncertainty-related anxiety. *Psychol. Rev.* **119**, 304–320

(2012).

13. Aberg, K. C., Toren, I. & Paz, R. A neural and behavioral trade-off between

value and uncertainty underlies exploratory decisions in normative anxiety.

Mol. Psychiatry **27**, 1573–1587 (03/2022).

14. Witte, K., Wise, T., Huys, Q. J. & Schulz, E. Exploring the Unexplored: Worry

as a Catalyst for Exploratory Behavior in Anxiety and Depression. (2024)

doi:10.31234/osf.io/td8xh.

15. Fan, H., Gershman, S. J. & Phelps, E. A. Trait somatic anxiety is associated

with reduced directed exploration and underestimation of uncertainty. *Nat Hum*

Behav 1–12 (2022).

16. Smith, R. *et al.* Lower Levels of Directed Exploration and Reflective Thinking

Are Associated With Greater Anxiety and Depression. *Front. Psychiatry* **12**,

(2022).

17. Holthoff, V. A. *et al.* Regional cerebral metabolism in early Alzheimer's

disease with clinically significant apathy or depression. *Biol. Psychiatry* **57**,

412–421 (2005).

18. Wen, M.-C., Chan, L. L., Tan, L. C. S. & Tan, E. K. Depression, anxiety, and

apathy in Parkinson's disease: insights from neuroimaging studies. *Eur. J.*

Neurol. **23**, 1001–1019 (2016).

19. Steffens, D. C., Fahed, M., Manning, K. J. & Wang, L. The neurobiology of

apathy in depression and neurocognitive impairment in older adults: a review

of epidemiological, clinical, neuropsychological and biological research.

Transl. Psychiatry **12**, 1–16 (2022).

20. Kalin, N. H. The Critical Relationship Between Anxiety and Depression. *Am.*

J. Psychiatry **177**, 365–367 (2020).

21. Dan, R. *et al.* Separate neural representations of depression, anxiety and apathy

in Parkinson's disease. *Sci. Rep.* **7**, 12164 (2017).

22. Tinaz, S. *et al.* Distinct neural circuits are associated with subclinical

neuropsychiatric symptoms in Parkinson's disease. *J. Neurol. Sci.* **423**, 117365

(2021).

23. Seligman, M. E. P. Helplessness: On depression, development, and death. *A

series of books in psychology.* **250**, (1975).

24. Oosterwijk, C. S., Vriend, C., Berendse, H. W., van der Werf, Y. D. & van den

Heuvel, O. A. Anxiety in Parkinson's disease is associated with reduced

structural covariance of the striatum. *J. Affect. Disord.* **240**, 113–120 (2018).

25. Bonanno, G. A., Chen, S., Bagrodia, R. & Galatzer-Levy, I. R. Resilience and

Disaster: Flexible Adaptation in the Face of Uncertain Threat. *Annu. Rev.

Psychol.* **75**, 573–599 (2024).

26. Chen, C. S., Knep, E., Han, A., Ebitz, R. B. & Grissom, N. M. Sex differences

in learning from exploration. *Elife* **10**, e69748 (2021).

27. Ebitz, R. B., Albarran, E. & Moore, T. Exploration Disrupts Choice-Predictive

Signals and Alters Dynamics in Prefrontal Cortex. *Neuron* **97**, 450-461.e9

(01/2018).

28. Kaske, E. A. *et al.* Prolonged Physiological Stress Is Associated With a Lower

Rate of Exploratory Learning That Is Compounded by Depression. *Biol

Psychiatry Cogn Neurosci Neuroimaging* (2022)

doi:10.1016/j.bpsc.2022.12.004.

29. Piray, P. & Daw, N. D. A simple model for learning in volatile environments.

PLoS Comput. Biol. **16**, e1007963 (2020).

30. Hampton, A. N., Bossaerts, P. & O'Doherty, J. P. The role of the ventromedial

prefrontal cortex in abstract state-based inference during decision making in

humans. *J. Neurosci.* **26**, 8360–8367 (2006).

31. Schlagenhauf, F. *et al.* Striatal dysfunction during reversal learning in

unmedicated schizophrenia patients. *Neuroimage* **89**, 171–180 (2014).

32. Cheng, S. *et al.* Uncertainty-aware and multigranularity consistent constrained

model for semi-supervised hashing. *IEEE Trans. Circuits Syst. Video Technol.*

32, 6914–6926 (2022).

33. Dayan, P., Kakade, S. & Montague, P. R. Learning and selective attention. *Nat.*

Neurosci. **3 Suppl**, 1218–1223 (2000).

34. Saffron, D. *The Rescorla - Wagner Interpretation of Blocking and*

Overshadowing in Pavlovian Conditioning. (University of Sydney, 1979).

35. den Ouden, H. E. M. *et al.* Dissociable effects of dopamine and serotonin on

reversal learning. *Neuron* **80**, 1572 (2013).

36. Ang, Y.-S., Lockwood, P., Apps, M. A. J., Muhammed, K. & Husain, M.

Distinct Subtypes of Apathy Revealed by the Apathy Motivation Index. *PLoS*

One **12**, e0169938 (2017).

37. Löwe, B. *et al.* Validation and standardization of the Generalized Anxiety

Disorder Screener (GAD-7) in the general population. *Med. Care* **46**, 266–274

(2008).

38. McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold

Approximation and Projection. *Journal of Open Source Software* **3**, 861

(2018).

39. Aylward, J. *et al.* Altered learning under uncertainty in unmedicated mood and

anxiety disorders. *Nat Hum Behav* **3**, 1116–1123 (2019).

40. Grupe, D. W. & Nitschke, J. B. Uncertainty and anticipation in anxiety: an

integrated neurobiological and psychological perspective. *Nat. Rev. Neurosci.*

14, 488–501 (2013).

41. Fahed, M. & Steffens, D. C. Apathy: Neurobiology, assessment and treatment.

Clin. Psychopharmacol. Neurosci. **19**, 181–189 (2021).

42. Pagonabarraga, J., Kulisevsky, J., Strafella, A. P. & Krack, P. Apathy in

Parkinson's disease: clinical features, neural substrates, diagnosis, and

treatment. *Lancet Neurol.* **14**, 518–531 (2015).

43. Pessiglione, M., Vinckier, F., Bouret, S., Daunizeau, J. & Le Bouc, R. Why not

try harder? Computational approach to motivation deficits in neuro-psychiatric

diseases. *Brain* **141**, 629–650 (2018).

44. Heimberg, R. G., Brozovich, F. A. & Rapee, R. M. *A Cognitive-Behavioral*

Model of Social Anxiety Disorder: Update and Extension. (2010).

45. Hofmann, S. G., Asnaani, A., Vonk, I. J. J., Sawyer, A. T. & Fang, A. The

efficacy of cognitive behavioral therapy: A review of meta-analyses. *Cognit.*

Ther. Res. **36**, 427–440 (2012).

46. Beck, J. S. *Cognitive Behavior Therapy: Basics and Beyond*. (Guilford Press,

2011).

47. Craske, M. G. & Mystkowski, J. L. Exposure therapy and extinction: Clinical

studies. in *Fear and learning: From basic processes to clinical implications*

217–233 (American Psychological Association, Washington, 2007).

48. Alexopoulos, G. S. & Arean, P. A model for streamlining psychotherapy in the

RDoC era: the example of "Engage." *Mol. Psychiatry* **19**, 14–19 (2014).

49. Yan, X., Yong, X., Huang, W. & Ma, Y. Placebo treatment facilitates social

trust and approach behavior. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 5732–5737

(2018).

50. Foroughi, C. K., Monfort, S. S., Paczynski, M., McKnight, P. E. &

Greenwood, P. M. Placebo effects in cognitive training. *Proc. Natl. Acad. Sci.*

U. S. A. **113**, 7470–7474 (2016).

51. Cuthbert, B. N. & Insel, T. R. Toward the future of psychiatric diagnosis: the

seven pillars of RDoC. *BMC Med.* **11**, 126 (2013).

52. Hinton, G. E. & Roweis, S. Stochastic neighbor embedding. *Adv. Neural Inf.*

Process. Syst. **15**, (2002).

53. Jolliffe, I. T. & Cadima, J. Principal component analysis: a review and recent

developments. *Philos. Trans. A Math. Phys. Eng. Sci.* **374**, 20150202 (2016).

54. Rhys, H. I. *Machine Learning with R, the Tidyverse, and Mlr*. (Simon and

Schuster, New York, NY, 2020).

55. Bari, A. *et al.* Serotonin modulates sensitivity to reward and negative feedback

in a probabilistic reversal learning task in rats. *Neuropsychopharmacology* **35**,

1290–1301 (2010).

56. Hayes, A. F. *Introduction to Mediation, Moderation, and Conditional Process*

Analysis: A Regression-Based Approach. Xvii. vol. 507 (Guilford Press, New

York, NY, 2017).

57. Mackinnon, D. P., Lockwood, C. M. & Williams, J. Confidence Limits for the

Indirect Effect: Distribution of the Product and Resampling Methods.

Multivariate Behav. Res. **39**, 99 (2004).

58. Preacher, K. J. & Hayes, A. F. Asymptotic and resampling strategies for

assessing and comparing indirect effects in multiple mediator models. *Behav.*

Res. Methods **40**, 879–891 (2008).

59. Coelho, J. P., Pinho, T. M. & Boaventura-Cunha, J. *Hidden Markov Models:*

Theory and Implementation Using Matlab ®. (CRC Press, Boca Raton, FL :

CRC Press, 2019. | "A science publishers book.," 2019).

60. Chakroun, K., Mathar, D., Wiehler, A., Ganzer, F. & Peters, J. Dopaminergic

modulation of the exploration/exploitation trade-off in human decision-

making. *Elife* **9**, (2020).

61. Forstmann, B. U., Ratcliff, R. & Wagenmakers, E.-J. Sequential sampling

models in cognitive neuroscience: Advantages, applications, and extensions.

Annu. Rev. Psychol. **67**, 641–666 (2016).

62. Piray, P., Dezfouli, A., Heskes, T., Frank, M. J. & Daw, N. D. Hierarchical

Bayesian inference for concurrent model fitting and comparison for group

studies. *PLoS Comput. Biol.* **15**, e1007043 (2019).

63. Rigoux, L., Stephan, K. E., Friston, K. J. & Daunizeau, J. Bayesian model

selection for group studies — Revisited. *Neuroimage* **84**, 971–985 (2014).

64. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical

and powerful approach to multiple testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).