**Individual differences in uncertainty evaluation explain opposing exploratory**

**behaviors in anxiety and apathy**

Xinyuan Yan [1], R. Becket Ebitz [3], Nicola Grissom [2], David P. Darrow [4†], Alexander

B. Herman [1†*]

[1]Department of Psychiatry and Behavioral Sciences, University of Minnesota,

Minneapolis, MN 55455, USA

[2]Department of Psychology, University of Minnesota, 75 E River Rd, Minneapolis,

MN 55455, USA

[3]Department of Neuroscience, Universite de Montreal, 2900 Edouard Montpetit

Blvd, Montreal, Quebec H3T 1J4, Canada

[4]Department of Neurosurgery, University of Minnesota, Minneapolis, MN 55455,

USA

* Corresponding author: Alexander B. Herman

† Denotes equal contribution

**Email:** herma686@umn.edu

**Author Contributions:** Conceptualization, X.Y, A.B.H, R.B.E, N.G, D.P.D;

Methodology, X.Y, R.B.E, A.B.H; Investigation, D.P.D, A.B.H; Formal Analysis,

X.Y; Writing – Original Draft, X.Y; Writing - Review & Editing, X.Y, D.P.D,

A.B.H; Funding Acquisition, A.B.H and N.G; Supervision, A.B.H, D.P.D.

**Competing Interest Statement:** The authors declare no competing interests.

**This PDF file includes:**

Main Text
Figures 1 to 5
Tables 0

**Abstract**

Navigating uncertain environments is a fundamental challenge for adaptive behavior, and

affective states such as anxiety and apathy can profoundly influence an individual's response to

uncertainty. Uncertainty encompasses both volatility and stochasticity, where volatility refers to

how rapidly the environment changes and stochasticity describes outcomes resulting from

random chance. This study investigates how anxiety and apathy modulate perceptions of

environmental volatility and stochasticity and how these perceptions impact exploratory

behavior. In a large online sample (N = 1001), participants completed a restless three-armed

bandit task, and their choices were analyzed using latent state models to quantify the

computational processes. We found that anxious individuals attributed uncertainty more to

environmental volatility than stochasticity, leading to *increased* exploration, particularly after

reward omission. Conversely, apathetic individuals perceived uncertainty as more stochastic than

volatile, resulting in *decreased* exploration. The ratio of perceived volatility to stochasticity

mediated the relationship between anxiety and exploratory behavior following adverse outcomes.

These findings reveal distinct computational mechanisms underlying anxiety and apathy in

uncertain environments. Our results provide a novel framework for understanding the cognitive

and affective processes driving adaptive and potentially maladaptive behaviors under

uncertainty, with implications for the characterization and treatment of neuropsychiatric

disorders.

**Keywords:** apathy, anxiety, uncertainty, volatility, stochasticity, exploration, multi-armed bandit

task

**Introduction**

Life is filled with unexpected challenges. Adapting to an ever-changing environment requires a balance between exploring the unknown and exploiting the known. How individuals interpret the cause of undesirable outcomes (e.g., failure in investment, career plateau, bad weather) in an uncertain environment shapes their subsequent actions [1]. When changes in outcomes are attributed to how quickly the environment is changing (the volatility), individuals may be motivated to explore more to acquire additional information and adapt their behavior. In contrast, attributing adverse outcomes to mere chance or 'bad luck' (i.e., stochasticity) may reduce the motivation to explore, leading some individuals to stick with existing strategies [2].

How individuals perceive and respond to environmental uncertainty can vary significantly, and these differences may be influenced by underlying affective states [3]. Apathy, characterized by a lack of motivation and goal-directed behavior [4,5], is an affective state associated with imprecise beliefs about action outcomes [6] and a tendency to persist with previous choices rather than explore [7]. This suggests that apathetic individuals may view outcomes as primarily stochastic, attributing events more to chance than controllable variables. This bias could discourage exploration and potentially reinforce a cycle of failure and helplessness [8].

In contrast, anxiety, marked by excessive worry and a heightened perception of potential threats [9,10] and uncertainty [11], may lead individuals to overestimate environmental volatility. Consequently, anxious individuals could be driven to seek new information to update their beliefs and reduce uncertainty [12]. However, research on the link between anxiety and exploration has yielded mixed findings, with some studies showing increased exploration to mitigate uncertainty [13,14] and others showing reduced exploration to avoid unpredictable feedback under high anxiety [15,16]. Notably, apathy and anxiety often coexist in clinical populations, such as Alzheimer's [17], Parkinson's disease [18], and depression [19,20], despite having distinct neural representations [21,22].

Building on these findings, we propose three fundamental questions to further elucidate the

relationship between affective states and decision-making under uncertainty. First, we aim to

investigate whether apathy and anxiety exhibit distinct behavioral patterns when individuals are

faced with uncertain situations. Second, we seek to examine how individual differences in levels

of apathy and anxiety are associated with perceptions of different types of uncertainty,

specifically volatility and stochasticity. Finally, we intend to explore how perceived volatility

influences exploratory behavior during decision-making processes.

We posit two competing hypotheses:

(1) Both apathetic and anxious individuals engage in less exploratory behavior but through

different computational mechanisms. Apathetic individuals weigh stochasticity over volatility

and explore less, while anxious individuals overestimate volatility, leading to a sense that their

actions cannot track or learn from the environment, ultimately leading to exploitation. This may

provide a computational account for learned helplessness [23] and the co-occurrence of apathy and

anxiety in various clinical populations, such as Parkinson's and Alzheimer's diseases.

(2) Apathetic individuals manifest less exploration, while anxious individuals engage in more

exploration. Apathetic individuals weigh stochasticity over volatility and explore less, while

anxious individuals overestimate volatility but explore more to reduce their uncertainty. This

result would be consistent with previous findings suggesting that the two affective states have

distinct neural substrates [21,24].

To address these questions, we employed a restless three-armed bandit task (Figure 1A), a well-

established paradigm for capturing adaptive learning in volatile environments [25,26]. We adopted

the Hidden Markov Model (HMM) to obtain the likelihood of individuals switching between

exploitation and exploration states [27,28]. To further investigate how volatility and stochasticity

modulate exploration, we utilized a Kalman filter model, which can dissociate two distinct

sources of noise, volatility (process noise variance) and stochasticity (observation noise

variance), during inference [29]. Together, these methods offer a comprehensive view of the

cognitive mechanisms underlying exploratory behavior and the manifestation of anxiety and

apathy.

## Results

We recruited a large gender-balanced online sample consisting of 1,001 adults. The participants, ranging in age from 18 to 54 years (mean ± SD = 28.446 ± 10.354 years; gender: 493 female), performed a restless three-armed bandit task, as depicted in Figure 1A. During this task, participants selected among three playing card images, with each card representing a different option. They made their selections by moving their cursor over their chosen card. The probability of receiving a reward from each card deck varied randomly over time. After each choice, feedback was displayed on the screen indicating whether a reward was received. Participants also completed symptom surveys assessing levels of anxiety and apathy (details in Methods and SI Section 1, Table S1). We defined the trial as a *switch* trial if the chosen option was different from the last trial, and a *stay* trial if the choice was the same as the last trial.

**Figure 1. Three-armed restless bandit task and distinct behavioral patterns associated with apathy and anxiety.**

(A) Three-armed restless bandit task. Participants chose one option from among the three targets

to receive reward or non-reward feedback. Each target was associated with a hidden reward

probability that randomly and independently changed throughout the task. The lower panel

indicates the example choice and reward sequence and the definition of stay and switch.

Specifically, stay was defined as choosing the same target as in the previous trial, while switch

was defined as choosing a different target. "+1" denotes reward feedback, and "+0" denotes

reward omission.

(B) Most participants earned more rewards than expected by chance

(C) Apathy and anxiety correlated positively.

(D) Apathy correlated negatively with switch behaviors, while anxiety correlated positively with

switch behaviors. Anxious individuals were more sensitive to undesired feedback (no reward)

and exhibited more switch behaviors compared to reward feedback.

(Panels in Figure 1C and 1D utilize binned correlation plots [25 quantile bins based on the x-

axis], with lines representing the standard error (S.E.). N.B. that these may be smaller than the

symbol. Statistical analyses were performed on raw data.)

***Apathy and anxiety predicted distinct exploratory behaviors***

We first evaluated the performance by comparing the total number of rewarded trials each

participant experienced against the number expected by chance. Out of the 1001 participants,

985 accrued more rewarded trials than would be statistically expected by chance, suggesting

significant effectiveness in their decision-making strategies (Figure 1B). As expected, anxiety

and apathy showed a significant positive correlation (r = 0.35, $p<10^{-29}$, Figure 1C), which is

consistent with previous findings on their co-occurrence [17].

To investigate the relationship between apathy and the percentage of switch behaviors

(P(switch)), as well as anxiety and P(switch), we conducted partial correlations between apathy

and exploration while controlling for anxiety, and between anxiety and exploration while

controlling for apathy. We found that apathy negatively predicted P(switch) (r = -0.16, $p<0.001$)

regardless of feedback type (reward or no-reward), while anxiety positively correlated with

P(switch) (r = 0.13, $p<0.001$). Intriguingly, the relationship between anxiety and switch

behaviors was greater after non-reward feedback (r = 0.16, $p<0.001$) compared to reward

feedback (r = 0.07, $p=0.024$) (their difference, z-score = 2.40, $p=0.01$). Though co-existing in

this population, these two affective states predicted distinct switch behaviors under uncertainty

(Figure 1D). The stronger relationship between anxiety and P(switch) after undesirable feedback

indicates that highly anxious individuals are more sensitive to negative feedback, which may

lead them to disengage.

Next, we fitted the behavior with a Hidden Markov Model (HMM) to decode the hidden states,

"*explore*", and "*exploit*" (Figure 2A) [26,27,30,31]. Each arm is associated with a hidden reward

probability that randomly and independently changes throughout the task (Figure 2A). In our

study, exploration and exploitation states are considered hidden states underlying the observed

choices, such as switching between decks or repeatedly choosing from the same deck. We

calculated the percentage of explore states, i.e., P(explore). Consistently, apathy correlated

negatively with P(explore) (r = -0.17, *p*<0.001), while anxiety positively correlated with

(P(explore)) (r = 0.11, *p*=0.003) as well as the percentage of  exploration after reward

omission(P(explore|0) (r = 0.13, *p*<0.001) (Figure 2C).

The hidden states associated with each observed behavior is important but not sufficient for fully

understanding the dynamic decision-making process. It's also necessary to examine how these

hidden states transition between each other. Therefore, we investigated how apathy and anxiety

manifest in the transition probability (Figure 2B) between explore and exploit. As predicted,

apathy had a positive correlation with the transition probability from *explore* to *exploit* (r = 0.13,

*p*<0.001) but a negative correlation with the transition probability from *exploit* to *explore* (r = -

0.08, *p*=0.011). In contrast, anxiety had a negative correlation with the transition probability

from *explore* to *exploit* (r = -0.07, *p*=0.035) but a positive correlation with the transition

probability from *exploit* to *explore* (r = 0.07, *p*<0.022) (Figure 2D). All significant results

reported in the study survived False Discovery Rate (FDR, *p*<0.05) correction.

**Figure 2. Apathy and anxiety have opposing relationships with exploration and explore and exploit state dynamics.**

(A) Unrolled structure of the hidden Markov model (HMM) used to infer the explore and exploit states' underlying behavior.

(B) The transition probabilities within and between states in the HMM.

(C) The probability of exploration, plotted as a function of apathy (top left) and anxiety (top right). The probability of exploration following a reward omission is plotted as a function of apathy (bottom left) and anxiety (bottom right).

(D) The transition probability from explore to exploit, plotted as a function of apathy (top

left) and anxiety (top right); the transition probability from exploit to explore plotted as a

function of apathy (bottom left) and anxiety (bottom right).

(Panels in Figure 2C and 2D utilize binned correlation plots [25 quantile bins based on the x-

axis], with lines representing the standard error (S.E.). N.B. that these may be smaller than

the symbol. Statistical analyses were performed on raw data).  * $p < 0.05$, ** $p < 0.01$, *** $p$

$< 0.001$. All p-values corrected for multiple comparisons.

*Apathy and anxiety are associated with distinct computational processes underlying*
*exploration.*

We then asked whether differing perceptions of the environment might explain the distinct patterns of exploration predicted by apathy and anxiety we observed.

To address this question, we utilized a Kalman filter model (Figure 3A), which can dissociate sources of uncertainty into perceived volatility and stochasticity [29]. Kalman filter (KF) models have been widely applied in psychology and neuroscience to study various aspects of learning and decision-making [32,33] (for more detailed information about the model, please refer to the Method section).

We also fitted the behavioral data to alternative models including volatile Kalman filter (VKF) [29], Rescorla-Wagner models single (RW1) [34] and dual learning rates (RW2) to weigh positive and negative learning rates [35]. We employed Hierarchical Bayesian inference (HBI) to fit models to choice data. Further, we used Bayesian model selection (BMS) and protected exceedance probability (PXP) to select the winning model (Figure 3B). The Kalman filter served as the best model for our population, and we examined the resulting distribution of volatility and stochasticity (Figure 3B).

We first conducted correlation analyses using all data points (Figure 3C). Specifically, we found that apathy was positively correlated with stochasticity (r = 0.08, $p$=0.013) but negatively correlated with volatility (r = -0.08, $p$=0.008). Conversely, anxiety showed a negative correlation with stochasticity (r = -0.12, $p$=0.001) and a positive correlation with volatility (r = 0.12, $p$=0.002). These correlations highlight the distinct cognitive biases associated with apathy and anxiety in processing environmental uncertainties.

To clearly illustrate and confirm the findings, we categorized participants into distinct groups based on their apathy and anxiety levels. For apathy, we identified the high apathy group (N = 223) as those scoring in the top 25% on the Apathy Motivation Index (AMI), which assesses apathy in behavioral and social domains [36]. Conversely, the low apathy group (N = 251) comprised individuals scoring in the bottom 25% of apathy scores. Similarly, for anxiety, the high anxiety group (N = 228) included participants within the top 25% of scores on the GAD-7 scale [37], while the low anxiety group (N = 250) consisted of those in the bottom 25%. These classifications allowed for a direct comparison of behaviors and traits between individuals with varying degrees of apathy and anxiety.

We conducted linear regression analyses using volatility and stochasticity as the dependent variables with the high versus low anxiety and apathy groups as predictors. The Methods section provides details of the regression model specifications.

As hypothesized, apathetic individuals overestimated stochasticity (t(471) = 3.06, $p$=0.002) and underestimated the volatility compared to those with low apathy (t(471) = -3.24, $p$=0.001)(Figure 3D). Consequently, apathetic individuals exhibited a lower learning rate than their low apathy counterparts (t(471) = -3.11, $p$=0.002).

In contrast, individuals with high anxiety levels tended to overestimate volatility (t(475) = 2.84, $p$=0.004) and underestimate stochasticity compared to those with low anxiety (t(475) = -3.04, $p$=0.002), resulting in a higher learning rate among the high anxiety group (t(475) = 3.21, $p$=0.001) (Figure 3E). Furthermore, comparisons showed that anxious individuals had higher volatility estimates than those with high apathy (t(449) = 2.75, $p$=0.006), whereas apathetic individuals had higher stochasticity estimates than their anxiety counterparts (t(449) = -3.01, $p$=0.002) (SI Section 2, Figure S1).

**Figure 3. Apathy and anxiety have opposing relationships with volatility and stochasticity.**

(A) The schematic of the Kalman filter model used in our analysis. The diagram illustrates how this model can differentiate between volatility (process noise variance) and stochasticity (observation noise variance), providing insights into the underlying decision-making processes.

(B) Bayesian model comparison and the distribution of volatility, stochasticity

(C) Correlation plots that visually represent the relationships between apathy, anxiety, and the distinct computational processes of volatility and stochasticity.

(D) Highly apathetic individuals overestimated stochasticity but underestimated the volatility, resulting in a lower learning rate.

(E) In contrast, highly anxious individuals overestimated volatility but underestimated stochasticity, resulting in a higher learning rate.

(Panels in Figure 3C utilize binned correlation plots [25 quantile bins based on the x-axis], with lines representing the standard error (S.E.). N.B. that these may be smaller than the symbol. Statistical analyses were performed on raw data). $*p < 0.05$, $** p < 0.01$, $*** p < 0.001$. All p-values corrected for multiple comparisons).

*The ratio of volatility to stochasticity distinguished apathy and anxiety*

To clarify the differential impacts of apathy and anxiety on decision-making under uncertainty, we computed the ratio of volatility to stochasticity, $/s$ , to represent the balance between these two types of uncertainties. A higher $v/s$ indicates a perception of greater volatility relative to stochasticity, while a lower ratio suggests a perception of more stochasticity relative to volatility. We applied a logarithmic transformation to the ratio to manage extreme values (e.g. cases where individuals might perceive very high volatility but very low stochasticity).

Consistently, our findings reveal a clear distinction: $v/s$ correlated negatively with apathy (r = -0.08, $p$=0.010) but positively with anxiety (r = 0.13, $p$<0.001) (Figure 4A).

*The ratio of volatility and stochasticity mediated the relationship between anxiety and the exploration after negative feedback*

To determine whether individual differences in the perception of uncertainty explain the relationship between exploratory behavior and affect, we conducted a mediation analysis with anxiety, switching after reward omission (P(switch | 0)), and $v/s$.The results demonstrate that

the relationship between anxiety and the tendency to switch after receiving no reward is

significantly mediated by the $v/s$ (Figure 4B). This mediation was also significant for the

analogous HMM model-based measures (see SI Section 3, Figure S2). No significant mediation

effect was found for apathy, however, reinforcing the unique pathways through which anxiety

influences exploratory behavior. These results explain why individuals with higher anxiety might

explore more after negative feedback, driven by an overweighting of perceived volatility relative

to stochasticity as a strategy to reduce uncertainty and manage risks.

**Figure 4. Distinctions in apathy and anxiety on the ratio of volatility to stochasticity and its**

**mediation effect.**

(A) The ratio of volatility to stochasticity, plotted as a function of apathy (left) and anxiety

positively (right).

(B) Mediation analysis, showing the mediating effect of the ratio of volatility to stochasticity on

the relationship between anxiety and switch behavior after reward omission.

(Panels in Figure 4A utilize binned correlation plots [25 quantile bins based on the x-axis], with

lines representing the standard error (S.E.). N.B. that these may be smaller than the symbol.

Statistical analyses were performed on raw data). \* $p < 0.05$, \*\* $p < 0.01$, \*\*\* $p < 0.001$. All p-

values corrected for multiple comparisons).

**A low dimensional manifold unifies exploration, perceptions of uncertainty and affective**

**state.**

The HMM state-model of exploration-exploitation and the Kalman filter process model of

uncertainty estimation represent complementary ways of understanding adaptive behavior that

our mediation results suggest are intrinsically related. We hypothesized that a latent structure

underlying adaptive behavior on this task might unify these descriptions of behavior. We utilized

advanced dimensionality reduction methods to uncover such a latent structure in the raw task

behavior.

First, we formatted each participant's trial-by-trial task data into sequences of choices to stay

(repeat the choice on the last trial) or switch (choose a different option) and reward outcome for

two consecutive trials ({$choice_{t-1}$, $outcome_{t-1}$, $choice_t$}, Figure 5A and 5B). The behavioral data

for each participant was then transformed into counts for each of these eight unique sequences.

Then we applied Uniform Manifold Approximation and Projection (UMAP) [38], a

computationally efficient algorithm that can preserve both the local and global distances between

data points in high-dimensional space, to learn the two-dimensional manifold underlying the

eight-dimensional behavioral data (Figure 5C, see Methods for more algorithm details).

Including additional reward history and applying other dimensionality reduction methods like

principle component analysis (PCA), and t-distributed Stochastic Neighbor Embedding (t-SNE)

did not change the results (SI Section 4, Figure S3, Table S2).

Our analysis using UMAP revealed distinct correlations within the derived dimensions.

Specifically, the dimension 1 score (the horizontal axis) exhibited a very strong significant

negative correlation with exploratory behavior (P(explore)) ($r = -0.90$, $p<10^{-200}$), but it showed no

significant relationship with the ratio of volatility to stochasticity ($v/s$)($r = 0.01$, $p=0.74$) (Figure

5D, 5E). In contrast, the dimension 2 score (the vertical axis) demonstrated a strong negative

correlation with $v/s$ ($r = -0.76$, $p<10^{-185}$), which was significantly more pronounced than its

correlation with P(explore) ($r = -0.19$, $p<10^{-10}$) (Figure 5F, 5G). This suggests that dimension 1

primarily represents exploratory behavior, while dimension 2 primarily reflects the

computational factors: volatility and stochasticity. (i.e., volatility and stochasticity).

Further, both dimensions also showed correlations with affective states: the dimension 1 score

was positively correlated with apathy ($r= 0.14$, $p<0.001$), and negatively correlated with anxiety

(r = - 0.11, $p<0.001$). Similarly, the dimension 2 score had a positive correlation with apathy (r =

0.097, $p=0.002$) and a negative correlation with anxiety (r = -0.088, $p=0.004$).

It is worth noting that we only found linear relationship between apathy, anxiety, and

exploration, as well as between these affective states and the ratio of volatility to stochasticity

(our analysis using higher order effects among these variables did not yield significant results,

more details can be found in SI Section 5, Table S3).

To delve deeper into how these factors interact in the low-dimensional space defined by UMAP,

we divided the data manifold into two groups based on the dimension1 score: a monotonically

decreasing group (left part, dimension 1 score < -0.671, $N=390$) and a monotonically increasing

group (right part, dimension 1 score> -0.671, $N=611$). The methodology used to identify the

turning point (dimension 1 score = -0.671) that differentiates the monotonically decreasing group

from the monotonically increasing group is detailed in the Methods section. The analysis

revealed that the monotonically decreasing group had relatively higher levels of anxiety

compared to the monotonically increasing group (t(999)=2.08, $p$ =0.037), while their apathy

levels were significantly lower (t(999)= -3.56, $p=0.0003$). This segmentation allows us to further

explore and understand the complex interplay between affective states, computational

parameters, and exploratory behaviors within a structured, low-dimensional framework.

Notably, individuals in the left monotonically decreasing group, characterized by high anxiety

and low apathy, generally perceive higher volatility relative to stochasticity and demonstrate

greater exploratory behavior compared to those in the right monotonically increasing group.

Interestingly, within the decreasing group, higher perceived volatility correlates with reduced

exploration. Conversely, in the increasing group, an increased perception of volatility tends to

stimulate more exploratory actions. These results suggest that while severe anxiety might

suppress exploration due to overwhelming uncertainty, moderate anxiety in the general

population can promote exploration as a coping mechanism to gather information and reduce

anxiety symptoms.

We now address our final research question; what is the relationship between volatility and

exploratory behavior? Considering the parabolic relationship between the manifold dimension

reflecting exploration and the dimension representing $v/s$, we predicted that the relationship

between exploration and $v/s$ might be quadratic.

To test this hypothesis, we constructed a regression model as follows:

*P(explore) ~ $v/s + v/s^2$ + anxiety + apathy*

The results revealed that both the linear and quadratic terms are significant (linear term,

coefficient = 0.02, SE = 0.003, t(996)=5.83, $p<10^{-9}$; quadratic term, coefficient = 0.005, SE =

$7.86\times10^{-4}$, t(996)=6.59, $p<10^{-9}$), indicating a complex, non-linear relationship between the ratio

of volatility to stochasticity and exploration (see SI Section 6, Figure S4), which was consistent

with the manifold representation.

**Figure 5. Visualizing the complex relationships in decision-making through low-**

**dimensional space.**

(A) All possible sequences of choices and rewards that participants could make during the

experiment

(B) The frequency distribution of individual decision-making patterns. The black line in the box

plot represents each pattern's mean value, highlighting participants' typical behaviors.

(C) Schematic high-dimensional space of participants' decision-making pattern

(D) The two-dimensional space representation of exploration by using the Uniform Manifold

Approximation and Projection (UMAP) (Different dimensionality reduction methods such as

principal component analysis (PCA), and t-distributed Stochastic Neighbor Embedding (t-SNE)

lead to a similar space)

(E) Dimension 1 exclusively represents P(explore) but does not represent the ratio of volatility to

stochasticity

(F) The two-dimensional space representation of the ratio of volatility to stochasticity behavior

by using UMAP (Different dimensionality reduction methods such as principal component

analysis (PCA), and t-distributed Stochastic Neighbor Embedding (t-SNE) lead to a similar

space).

(G) Dimension 2 mainly represents the ratio of volatility to stochasticity

but not P(explore).

(H) The manifold has been separately dissociated into the monotonically decreasing group (the

most left panel) and monotonically increasing group (the most right panel). The monotonically

decreasing group was associated with a relatively higher anxiety level than the monotonically

increasing group, while the apathy level was significantly lower than the monotonically

increasing group. Within the monotonically decreasing group (left part), a higher volatility to

stochasticity ratio leads to decreased exploration. In contrast, within the monotonically

increasing group (right part), a higher volatility to stochasticity ratio encouraged higher

exploration. This exploration serves as a coping strategy to relieve anxious feelings in the

environment.

**Discussion**

We found that apathy and anxiety predicted opposing patterns of exploratory behavior, which

were explained partly by differing perceptions of uncertainty. Anxiety was associated with

increased exploration after reward omission and greater volatility estimation: the attribution of

uncertainty to a rapidly changing (but still learnable) environment. Apathy, in contrast, predicted

decreased exploration and higher stochasticity estimation: the perception of uncontrollable

randomness. Following a dimensionality reduction of the raw behavioral data, exploration and

perceptions of uncertainty emerged as the dimensions of an underlying latent structure that

unified the different model approaches and the affective states. These findings elucidate the

complex interplay between cognitive assessments of uncertainty, affective states, and decision-

making processes, offering several key insights into adaptive and maladaptive behaviors under

uncertainty.

The distinct patterns of exploratory behavior observed in anxious and apathetic individuals

highlight the role of affective states in shaping responses to uncertainty. Anxious individuals,

who generally display a heightened sensitivity to potential threats and environmental changes,

exhibited a bias toward perceiving greater volatility and exploring more after negative outcomes.

Our mediation analysis revealed that the perception of volatility relative to stochasticity partially

mediates the relationship between anxiety and exploratory behavior after reward omission. This

finding is consistent with previous results [39] and offers a mechanistic explanation for why

anxious individuals in a healthy population might choose to explore more after receiving

negative feedback. The perceived overweighting of volatility relative to stochasticity may drive

these individuals to seek more information, potentially as a strategy to reduce uncertainty and

manage perceived risks more effectively [40]. Although such a strategy may be beneficial for

adaptation in genuinely volatile environments, it may also contribute to excessive worry and

stress, especially if the perceived level of volatility exceeds actual environmental volatility [9,11].

Consequently, anxious individuals may find themselves in a prolonged state of heightened

arousal and uncertainty, leading to suboptimal decision-making and diminished well-being.

On the other hand, apathetic individuals, who generally exhibit diminished motivation and

responsiveness [41], tended to attribute outcomes more to stochasticity in our study. This

perception might underlie their reduced exploratory behavior, reflecting a disengagement from

active learning and adaptation. If outcomes seem random and beyond our control, expending

energy to explore may seem futile, and focusing on what we know seems rational. While this

approach may conserve energy, this inflexibility can perpetuate a cycle of disengagement and

maintain apathetic symptoms [42,43]. Apathetic individuals may fail to recognize the potential

benefits of exploration and remain stuck in suboptimal decision-making patterns, further

reinforcing their disengagement from the environment [4].

The dimensionality reduction of the behavioral sequence data using UMAP allowed us to

examine the relationship between exploration and the estimation of volatility and stochasticity.

Despite the intuitive connection between these two behavior models, their relationship has not

been directly examined. Our results showed that exploration and uncertainty estimation related

closely to the two axes of a parabolic latent structure of adaptive behavior. As a result, both

model-based metrics were necessary to characterize the spectrum of individual differences fully.

Segmenting the data on the manifold further illuminated the fine-grained interplay between

affective states and exploratory behavior. Individuals with relatively higher anxiety and lower

apathy (the monotonically decreasing group) generally weighted volatility more and

demonstrated greater exploratory behavior compared to those with lower anxiety and higher

apathy (the monotonically increasing group). However, *within* these groups formed on the

manifold, individuals exhibited opposing relationships between uncertainty and exploration. In

the higher anxiety group, perceived volatility correlates inversely with exploration. However, in

the lower anxiety group, increased volatility perception predicts greater exploration.

These results reconcile previously inconsistent findings regarding exploratory behavior in

individuals with anxiety, with some studies showing more exploitative behavior [15,16], and others

finding that higher anxiety predicts more exploratory behaviors [13,14]. The relationship between

perceived volatility and exploration is modulated by the degree of anxiety, with more severe

anxiety potentially suppressing exploration as a form of avoidance. In contrast, milder anxiety

may promote exploration as a coping mechanism to gather more information reduce uncertainty,

and alleviate discomfort. This dual response to perceived volatility underscores the complex

interplay between anxiety levels, environmental perceptions, and behavioral strategies in

managing emotional responses.

Our results have implications for behavioral approaches aimed at improving adaptive functioning

and mental health. Cognitive-behavioral therapies that focus on challenging and restructuring

interpretations of beliefs about uncertainty and controllability could be personalized by symptom

profile [44,45]. For anxious individuals, interventions that help recalibrate perceptions of volatility

and equip them with coping strategies for managing uncertainty may reduce maladaptive worry

and enhance decision-making efficacy [46,47]. Furthermore, encouraging anxious individuals to

integrate information over longer horizons when undesirable things happen, rather than

attempting to change strategies to these events rapidly, may prove a beneficial approach to

managing anxiety [39]. For apathetic individuals, strategies aimed at strengthening the locus of

control and highlighting the efficacy of their actions may counteract tendencies towards

stochasticity attribution. Training individuals to view the world as more controllable may be a

practical addition to existing psychotherapeutic approaches, including Behavioral Activation and

Motivational Interviewing [48]. Challenging beliefs about uncontrollability may promote a more

balanced perception of the environment and encourage exploration. Previous studies have

demonstrated that individuals who believed they received treatment or cognitive training that

enhanced prosocial behaviors and cognitive abilities actually exhibited more prosocial behaviors

and improved performance in standard IQ tests [49,50]. This suggests that manipulating

participants' beliefs about their environmental control through placebo interventions could

enhance their resilience in uncertain situations.

Another important clinical implication involves using an individual's position on the behavioral

manifold (Figure 5) to predict how their behavior might change in response to treatment based

on their symptoms. For example, a patient positioned in the upper left quadrant before treatment

may exhibit higher anxiety, lower apathy, and increased exploratory behavior. During and after

treatment, monitoring these behavioral shifts may allow us to infer changes in their affective

states or symptoms based on their new manifold position. To develop such a tool, several

questions remain: Do changes within an individual follow a predictable trajectory on this

manifold? Do clinical populations conform to the same manifold, or do they deviate, projecting

into the larger, unoccupied areas of the manifold? The answers to these questions could enhance

the implementation of dimensional approaches for individualized neuropsychiatric care [51].

Our results must be interpreted in light of notable limitations.  First, the study primarily utilized

an online sample, which may not accurately represent the demographic and clinical

characteristics of populations with specific mental health diagnoses. The potential differences in

internet access, motivation, and the self-report nature of online studies can introduce biases that

may differ from clinical settings. Consequently, the generalizability of our findings to clinical

populations remains to be determined. Second, our results, while robust statistically, are

inherently correlational, limiting our ability to infer causal relationships between the affective

states of apathy and anxiety and their impacts on decision-making processes. The observed

associations provide a strong foundation for hypothesizing causal mechanisms but do not

confirm them. Future studies may examine clinical samples of conditions known to affect

adaptive decision-making under uncertainty, such as depression, anxiety disorders, and

Parkinson's disease, as well as interventions targeting the physiology of adaptive behavior.

**Method**

*Ethics approval*

The experimental procedures of all experiments were in line with the standards set by the

Declaration of Helsinki and were approved by the local Research Ethics Committee of the

University of Minnesota, Twin Cities. Participants provided written informed consent after the

experimental procedure had been fully explained and were reminded of their right to withdraw at

any time during the study.

*Participants*

We recruited a sample of 1512 participants via Amazon Mechanical Turk (MTurk) and Prolific

(Prolific. co); exclusion criteria included current or history of neurological and psychiatric

disorders. 1001 participants completed all questionnaires and the bandit task (age range 18-54,

mean $\pm$ SD = 28.446 $\pm$ 10.354 years; gender, 493 female). All participants were compensated for

their time in accordance with minimum wage.

*Questionnaire measurement*

Participants' anxiety and apathy states were measured by the General Anxiety Disorder Screener

(GAD-7) [37], and the Apathy-Motivation Index [36], respectively. More specifically, GAD-7

contains 7 items for assessing anxiety severity in the last two weeks. All items were rated on a 4-

point scale, with higher scores indicating greater anxiety. Participants' apathy level was

measured using the 18-item Apathy-Motivation Index (AMI), which was designed to identify

and measure general apathy, as well as subtypes of apathy in behavioral, social, and emotional

domains. Higher scores on AMI represent greater apathy. For all questionnaire scores, see SI,

Section 1 & Table S1.

*Three-armed restless bandit task*

We assessed exploration-exploitation behavioral dynamics using a 300-trial three-armed restless

bandit task [27]. Participants were free to choose between three targets for the potential to earn a

reward of 1 point. Each target is associated with a hidden reward probability that randomly and

independently changes throughout the task. We seeded each participant's reward probability

walks randomly to prevent biases due to particular kinds of environments. We assessed

performance by comparing the total number of rewarded trials to that expected by chance. Out of

the 1001 participants, 985 accrued more rewarded trials than would be expected by chance.

*Dimensionality reduction method*

Popular and valid dimensionality reduction techniques to reveal manifolds include t-distributed

stochastic neighborhood embedding (t-SNE) [52], uniform manifold approximation and projection

(UMAP) [38], and Principal component analysis (PCA) [53]. However, t-SNE suffers from

limitations, including slow computation time and loss of global data structure, and it is not a

deterministic algorithm [54]. The main drawback of PCA is that it is highly affected by outliers in

the dataset [53]. In contrast, UMAP is a deterministic and efficient algorithm, it also preserves both

local and global structure of original high-dimensional data. Uniform Manifold Approximation

and Projection (UMAP)

UMAP was implemented in the R language. The eight-dimensional datasets from all participants

were passed into the R package *umap*, version 0.2.8.0, available at https://cran.r-

project.org/web/packages/umap/) with default parameter setting as n_component = 2,

n_neighbors = 15, min_dist = 0.3, metric = 'Euclidean'. For reproducibility reasons, we fixed the

random_state in this algorithm. The hyperparameter *n_neighbors* decide the radius of the search

region. Larger values will include more neighbors, thus forcing this algorithm to consider more

global structure of original n-dimension data. Another important hyper-parameter, *min_dist*

determines the allowed minimum distance apart for cases in lower-dimensional space. *metric*

defines the way that UMAP is used to measure distances along the manifold.

*Model-free analyses*

We adopted some widely used model-free measures, including win-stay and lose-shift [35,55] as the

direct measurement for this learning task.

*Win-stay*. Win-stay is defined as the percentage of times that the choice in trial *t*-1 was repeated

on trial *t* following a reward.

*Lose-shift*. In contrast, lose-shift equals the percentage of trials that the choice was shifted or

changed when the outcome of trial t-1 was non-reward.

Model free results can be found at SI Section 7, Table S4.

*Mediation analyses*

Mediation analysis is a statistical method used to examine the underlying mechanisms by which

an independent variable influences a dependent variable through one or more mediator variables

[56]. In our study, we employed the bootstrapping method to estimate the mediation effect of

volatility and stochasticity on the relationship between affective states (apathy and anxiety) and

exploration. Bootstrapping is a nonparametric approach to effect-size estimation and hypothesis

testing that is increasingly recommended for many types of analyses, including mediation [57].

This method involves repeatedly resampling from the available data to generate an empirical

approximation of the sampling distribution of the indirect effect (i.e., the effect of the

independent variable on the dependent variable through the mediator). We used this distribution

to calculate p-values and construct confidence intervals based on 5,000 resamples. Bootstrapping

is preferred over other methods, such as the Sobel test because it does not assume the normality

of the sampling distribution and provides more accurate confidence intervals that are bias-

corrected and accelerated [56,57]. This approach offers a robust and powerful way to test mediation

hypotheses, particularly in cases where the sample size is relatively small or the data violate

assumptions of normality [58].

*Hidden Markov Model*

We fit a Hidden Markov Model (HMM) to the behavior, to decode the hidden state of each trial

for each participant. Fundamentally, the HMM has two layers, the hidden layer (i.e., state) and

the observable layer. The hidden dimension should satisfy the Markov property. That is, the

current hidden state only depends on the previous state but not any past model history. The

observable dimension entirely depends on the current hidden states and is independent of other

observations. Parameters of the hidden Markov model can be represented as $\Omega \rightarrow (T, O, c)$.

Specifically, $T$ is the transition probabilities matrix, $O$ is the observation probabilities matrix, or

emissions matrix, and $c$ refers to a vector with initial probabilities for each hidden state. Here,

we have two hidden states, an "exploration" state and an "exploitation" state.

The transition probability between exploration and exploitation states can be represented by:

$$T = \begin{pmatrix} t_{11} & \cdots & t_{1m} \\ \vdots & \ddots & \vdots \\ t_{m1} & \cdots & t_{mm} \end{pmatrix}$$

$$t_{explore,exploit} = P(q_k = s_{exploit}|q_{k-1} = s_{explore})$$

609      Where $t_{explore,exploit}$ refers to the transition probability from hidden state $s_{exploit}$ to another

610      hidden state $s_{explore}$

611      $k$ = time instant, $m$ = state sequence length

613      Then matrix $O$ represents the transition probabilities between hidden and observable states.

$$O = \begin{pmatrix} o_{11} & \cdots & o_{1n} \\ \vdots & \ddots & \vdots \\ o_{m1} & \cdots & o_{mn} \end{pmatrix}$$

$$o_{explore,exploit} = P(\lambda_k = r_{exploit}|q_{k-1} = s_{explore})$$

$k$ = time instant, $n$ = observation sequence length

Both matrix $O$ and $T$ satisfy the principle that the sum along the rows must be equal to 1.

$c$ is an m-dimensional row vector that refers to the initial probability distribution. In our current

study, the initial probability was fixed and equal to the available choices.

We fit HMM via expectation-maximization using the Baum-Welch algorithm and decode hidden

states from observed choice sequences by the Viterbi algorithm [59]. Model results of HMM can be

found at SI Section 7, Table S5.

*Kalman filter*

The Kalman filter (KF) model has been widely applied in psychology and neuroscience to study

various aspects of learning and decision-making [32,33].

In the Kalman filter model for a multi-armed bandit task, *process noise* and *observation noise*

refer to two distinct sources of uncertainty that affect the learning and decision-making process.

Process noise represents the uncertainty in the evolution of the hidden state (reward mean) over

time. It accounts for how the true state evolves from one point in time to the next. In

mathematical terms, process noise is part of the state transition equation in the Kalman Filter:

$$x_t = x_{t-1} + e_t$$

$x_t$ is the state at time $t$

$e_t$ is the process noise $t$, which is assumed to be drawn from a normal distribution with zeros

mean and **process noise variance** $v$. Where the $e_t \sim N(0, v)$.

The process noise captures the idea that the reward-means for each arm can change from one

trial to the next, even in the absence of any observations. A higher process noise variance $v$

indicates a more volatile environment, where the reward means are expected to change more

rapidly.

In contrast, **observation noise** represents the uncertainty in the observed rewards, given the

current hidden state (reward mean). Which is assumed to be Gaussian with zero mean and a

fixed variance $\sigma^2$.

The observation noise captures the idea that the observed rewards are noisy and can deviate from

the true reward mean due to random fluctuations or measurement errors.

A higher measurement noise variance indicates a more stochastic environment, where the

observed rewards are less reliable and informative about the underlying reward means.

The Kalman Filter operates optimally when the statistical properties of the process noise and the

measurement noise are accurately known.

When observation noise variance ($\sigma^2$) is high relative to the process noise variance ($v$), the

Kalman gain will be small, and the model will rely more on its prior beliefs and less on noisy

observations. Conversely, when the observation noise variance ($v$), is high relative to the process

noise variance ($\sigma^2$), the Kalman gain will be large, and the model will update its beliefs more

strongly based on the observed rewards.

*Extended Kalman filter for three-armed bandit task*

The Kalman filter model can be extended to capture the effects of both volatility and

stochasticity in a multi-armed bandit task [29,60].

In the current study, process noise variance ($v$) and observation noise variance ($\sigma^2$) represent

volatility and stochasticity, respectively.

A traditional assumption of the Kalman filter is that the process noise variance, $v$, as well as the

observation noise variance, $\sigma^2$ are constant.

Reward means update:

$$m_t = m_{t-1} + k_t(O_t - m_{t-1})$$

670 Where $m_t$ is the estimated mean or value of the chosen arm at time $t$

671 and $O_t$ is the observed reward at time $t$.

672 The mean update is driven by the prediction error, which is the difference between the observed

673 reward and the previous estimate.

675 Kalman gain is defined as:

677
$$k_t = (w_{t-1} + v)/(w_{t-1} + v + \sigma^2)$$

Here, $k_t$ represents the Kalman gain or learning rate, which adjusts the weight given to new

information based on the relative uncertainty of the prior estimate ($w_{t-1}$) and the total noise ($v +$

$\sigma^2$). When the stochasticity ($\sigma^2$) is high relative to the volatility ($v$), the Kalman gain (learning

rate) will be small, and the model will rely more on its prior beliefs and less on the observations.

Conversely, when the volatility ($v$), is high relative to the stochasticity ($\sigma^2$), the Kalman gain

(learning rate) will be large, and the model will update its beliefs more strongly based on the

observed rewards.

Variance update equation:

688
689
$$w_t = (1 - k_t)(w_{t-1} + v)$$

This equation updates the posterior variance ($w_t$), which represents the estimate's uncertainty

after observing $O_t$.

***Volatile Kalman filter for three-armed bandit task***

The key difference between a standard Kalman filter and a volatile Kalman filter (VKF) is the

variance of the process noise, a stochastic variable that changes with time. In other words, the

VKF introduces parameters to handle the volatility in the process noise. Specifically, it allows

the process noise variance $v$ to vary with the observed prediction errors, reflecting changes in

environmental volatility.

Our approach here is essentially the same as that taken by Piray and Daw [29]. Here, we briefly

described the model details as follows.

Kalman gain:

$$k_t = (w_t + v_{t-1})/(w_{t-1} + v_{t-1} + \sigma^2)$$

Update for the reward means:

$$m_t = m_{t-1} + k_t(O_t - m_{t-1})$$

Update for posterior variance $w_t$:

$$w_t = (1 - k_t)(w_{t-1} + v_{t-1})w_{t-1,t} = (1 - k_t)w_{t-1}$$

Update for volatility:

$$v_t = v_{t-1} + \lambda((m_t - m_{t-1})^2 + w_{t-1} + w_t - 2w_{t-1,t} - v_{t-1})$$

***Rescorla-Wagner models***

We also fitted the data to the classical Rescorla-Wagner model. Successful adaptation in a dynamic

situation requires the appropriate feedback-based learning process where individuals integrate the

feedback (reward or non-reward) into the stimulus-outcome association [61]. The basic

reinforcement learning model, the Rescorla-Wagner model [34] can address this process well. So the

first model (RW1) was defined as:

$$v_t = v_{t-1} + a \times (R_{t-1} - v_{t-1})$$

where $v_t$ is the value of the option on trial $t$.

$a$ represents the general learning rate from feedback.

To verify whether participants employed distinct or shared computational responses to positive

and negative feedback, we built another model with two learning rates, one for positive feedback

and the other for negative feedback [35]. This model (RW2) can be defined as:

$$v_t = v_{t-1} + \alpha^{pos} \times (R_{t-1} - v_{t-1}), positive\ feedback$$
$$v_t = v_{t-1} + \alpha^{neg} \times (R_{t-1} - v_{t-1}), negative\ feedback$$
735

Where $v_t$ is the value of the option on trial $t$. $\alpha^{pos}$ and $\alpha^{neg}$ represent the learning rates from

positive and negative feedback, respectively.

For these two models, $R_{t-1} \in \{0,1\}$ represents the feedback received in response to participants'

choice on trial $t$-1. And $R_{t-1} - v_{t-1}$ represents prediction error in trial $t$-1.

We used a softmax choice function to map the value into choice. The softmax function for these

four models can be defined as:

$$P^t = \frac{exp^{(\beta V, t)}}{exp^{(\beta V_{t,1} + \beta V_{t,2} + \beta V_{t,3})}}$$

Where the $\beta$ represents the inverse temperature with choice value.

***Model fitting and comparison***

Hierarchical Bayesian inference (HBI) is a powerful method for model fitting and comparison in

group studies [62]. Unlike traditional approaches such as maximum likelihood estimation (MLE) or

maximum a posteriori (MAP) estimation, which fit models to each subject independently, HBI

simultaneously fits models to all subjects while constraining individual fits based on group-level

statistics (i.e., empirical priors). This approach yields more robust and reliable parameter

estimates, particularly when individual subject data is noisy or limited.

In our study, we employed HBI to fit models to choice data. The method quantifies group-level

mean parameters and their corresponding hierarchical errors. To ensure that parameter estimates

remain within appropriate bounds during the fitting process, we used the sigmoid function to

transform parameters bounded in the unit range or with an upper bound and the exponential
function to transform parameters bounded to positive values. The initial parameters of all models
were obtained using a MAP procedure, with the initial prior mean and variance for all parameters
set to 0 and 6.25, respectively, based on previous research [29]. This initial variance allows
parameters to vary widely without substantial influence from the prior.
For model comparison, we used Bayesian model selection, specifically employing the protected
exceedance probability (PXP) to select the winning model. The PXP quantifies the probability
that a given model is more frequent in the population than all other models under consideration
while accounting for the possibility that the observed differences in model evidence may be due
to chance [63]. The model with the highest PXP is selected as the winning model. This approach
inherently penalizes model complexity, favoring models that balance goodness-of-fit and
parsimony. Model performance (log-likelihood) can be found in SI Section 8, Table 6.
771
***Turning point to divide the manifold into monotonically increasing and decreasing group***
To divide the manifold into monotonically increasing and decreasing groups, we sorted the
scores for dimension 1 in ascending order. Initially, we fitted a linear model using the first three
data points located on the upper left of the manifold. We then expanded this model by
sequentially including one additional data point from dimension 1, continuing this process until
we incorporated the last score (i.e., the maximum dimension 1 score, situated on the upper right
of the manifold). Throughout this procedure, we monitored the t-statistic of the dimension1
coefficient to assess the statistical significance of dimension1 as a predictor. Notably, a
dimension 1 score of -0.671 marked the most significant negative coefficient, after which the
relationship between dimension 1 and dimension2 gradually shifted to become positive (see SI
Section 9, Figure S5)
783
784 ***False discovery rate correction***
785 We adopted FDR (False Discovery Rate) correction, which was introduced by Benjamini and
786 Hochberg [64] to control the expected proportion of false positives (Type I errors).
787
788

**Acknowledgments**

This work was supported by NIMH under award #R21MH127607, NIDA under award

#K23DA050909, and the University of Minnesota's MnDRIVE (Minnesota's Discovery,

Research and Innovation Economy) initiative. Natural Sciences and Engineering Council of

Canada (Discovery Grant RGPIN-2020-05577), the Research Corporation for Science

Advancement & Frederick Gardner Cottrell Foundation (Project #29087), and a Research

Fellowship from the Jacobs Foundation to R.B.E.

We thank A. David Redish, Cathy Chen, Seth D Koenig, and Brian Sweiss for helpful comments

on the manuscript.

**Code and data availability.**

The source scripts used to do data analysis are published at

https://github.com/psywalkeryanxy/apathy_anxiety_uncertainty

Data is available upon request.

**Reference**

1.  Soltani, A. & Izquierdo, A. Adaptive learning under expected and unexpected uncertainty. *Nat. Rev. Neurosci.* **20**, 635–644 (2019).

2.  Piray, P. & Daw, N. D. A model for learning based on the joint estimation of stochasticity and volatility. *Nat. Commun.* **12**, 6587 (2021).

3.  Pulcu, E. & Browning, M. The Misestimation of Uncertainty in Affective Disorders. *Trends Cogn. Sci.* **23**, 865–875 (2019).

4.  Husain, M. & Roiser, J. P. Neuroscience of apathy and anhedonia: a transdiagnostic approach. *Nat. Rev. Neurosci.* **19**, 470–484 (2018).

5.  Marin, R. S. Apathy: Concept, Syndrome, Neural Mechanisms, and Treatment. *Semin. Clin. Neuropsychiatry* **1**, 304–314 (1996).

6.  Hezemans, F. H., Wolpe, N. & Rowe, J. B. Apathy is associated with reduced precision of prior beliefs about action outcomes. *J. Exp. Psychol. Gen.* **149**, 1767–1777 (2020).

7.  Scholl, J., Trier, H. A., Rushworth, M. F. S. & Kolling, N. The effect of apathy and compulsivity on planning and stopping in sequential decision-making. *PLoS Biol.* **20**, e3001566 (2022).

8.  Huys, Q. J. M. & Dayan, P. A Bayesian formulation of behavioral control. *Cognition* **113**, 314–328 (2009).

9.  Browning, M., Behrens, T. E., Jocham, G., O'Reilly, J. X. & Bishop, S. J. Anxious individuals have difficulty learning the causal statistics of aversive environments. *Nat. Neurosci.* **18**, 590–596 (2015).

10. Uhr, K. & Dugas, M. J. The role of fear of anxiety and intolerance of uncertainty in worry: an experimental manipulation. *Behav Res Ther* **47**, 215–223 (2009).

11. Gagne, C., Zika, O., Dayan, P. & Bishop, S. J. Impaired adaptation of learning to

contingency volatility in internalizing psychopathology. *Elife* **9**, e61387 (2020).

12. Hirsh, J. B., Mar, R. A. & Peterson, J. B. Psychological entropy: a framework for

understanding uncertainty-related anxiety. *Psychol. Rev.* **119**, 304–320 (2012).

13. Aberg, K. C., Toren, I. & Paz, R. A neural and behavioral trade-off between value and

uncertainty underlies exploratory decisions in normative anxiety. *Mol. Psychiatry* **27**, 1573–

1587 (03/2022).

14. Witte, K., Wise, T., Huys, Q. J. & Schulz, E. Exploring the Unexplored: Worry as a Catalyst

for Exploratory Behavior in Anxiety and Depression. (2024) doi:10.31234/osf.io/td8xh.

15. Fan, H., Gershman, S. J. & Phelps, E. A. Trait somatic anxiety is associated with reduced

directed exploration and underestimation of uncertainty. *Nat Hum Behav* 1–12 (2022).

16. Smith, R. *et al.* Lower Levels of Directed Exploration and Reflective Thinking Are

Associated With Greater Anxiety and Depression. *Front. Psychiatry* **12**, (2022).

17. Holthoff, V. A. *et al.* Regional cerebral metabolism in early Alzheimer's disease with

clinically significant apathy or depression. *Biol. Psychiatry* **57**, 412–421 (2005).

18. Wen, M.-C., Chan, L. L., Tan, L. C. S. & Tan, E. K. Depression, anxiety, and apathy in

Parkinson's disease: insights from neuroimaging studies. *Eur. J. Neurol.* **23**, 1001–1019

(2016).

19. Steffens, D. C., Fahed, M., Manning, K. J. & Wang, L. The neurobiology of apathy in

depression and neurocognitive impairment in older adults: a review of epidemiological,

clinical, neuropsychological and biological research. *Transl. Psychiatry* **12**, 1–16 (2022).

20. Kalin, N. H. The Critical Relationship Between Anxiety and Depression. *Am. J. Psychiatry*

**177**, 365–367 (2020).

21. Dan, R. *et al.* Separate neural representations of depression, anxiety and apathy in

Parkinson's disease. *Sci. Rep.* **7**, 12164 (2017).

22. Tinaz, S. *et al.* Distinct neural circuits are associated with subclinical neuropsychiatric

symptoms in Parkinson's disease. *J. Neurol. Sci.* **423**, 117365 (2021).

23. Seligman, M. E. P. Helplessness: On depression, development, and death. *A series of books*

*in psychology.* **250**, (1975).

24. Oosterwijk, C. S., Vriend, C., Berendse, H. W., van der Werf, Y. D. & van den Heuvel, O.

A. Anxiety in Parkinson's disease is associated with reduced structural covariance of the

striatum. *J. Affect. Disord.* **240**, 113–120 (2018).

25. Bonanno, G. A., Chen, S., Bagrodia, R. & Galatzer-Levy, I. R. Resilience and Disaster:

Flexible Adaptation in the Face of Uncertain Threat. *Annu. Rev. Psychol.* **75**, 573–599

(2024).

26. Chen, C. S., Knep, E., Han, A., Ebitz, R. B. & Grissom, N. M. Sex differences in learning

from exploration. *Elife* **10**, e69748 (2021).

27. Ebitz, R. B., Albarran, E. & Moore, T. Exploration Disrupts Choice-Predictive Signals and

Alters Dynamics in Prefrontal Cortex. *Neuron* **97**, 450-461.e9 (01/2018).

28. Kaske, E. A. *et al.* Prolonged Physiological Stress Is Associated With a Lower Rate of

Exploratory Learning That Is Compounded by Depression. *Biol Psychiatry Cogn Neurosci*

*Neuroimaging* (2022) doi:10.1016/j.bpsc.2022.12.004.

29. Piray, P. & Daw, N. D. A simple model for learning in volatile environments. *PLoS Comput.*

*Biol.* **16**, e1007963 (2020).

30. Hampton, A. N., Bossaerts, P. & O'Doherty, J. P. The role of the ventromedial prefrontal

cortex in abstract state-based inference during decision making in humans. *J. Neurosci.* **26**,

8360–8367 (2006).

31. Schlagenhauf, F. *et al.* Striatal dysfunction during reversal learning in unmedicated

schizophrenia patients. *Neuroimage* **89**, 171–180 (2014).

32. Cheng, S. *et al.* Uncertainty-aware and multigranularity consistent constrained model for

semi-supervised hashing. *IEEE Trans. Circuits Syst. Video Technol.* **32**, 6914–6926 (2022).

33. Dayan, P., Kakade, S. & Montague, P. R. Learning and selective attention. *Nat. Neurosci.* **3**

**Suppl**, 1218–1223 (2000).

34. Saffron, D. *The Rescorla - Wagner Interpretation of Blocking and Overshadowing in*

*Pavlovian Conditioning*. (University of Sydney, 1979).

35. den Ouden, H. E. M. *et al.* Dissociable effects of dopamine and serotonin on reversal

learning. *Neuron* **80**, 1572 (2013).

36. Ang, Y.-S., Lockwood, P., Apps, M. A. J., Muhammed, K. & Husain, M. Distinct Subtypes

of Apathy Revealed by the Apathy Motivation Index. *PLoS One* **12**, e0169938 (2017).

37. Löwe, B. *et al.* Validation and standardization of the Generalized Anxiety Disorder Screener

(GAD-7) in the general population. *Med. Care* **46**, 266–274 (2008).

38. McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold

Approximation and Projection. *Journal of Open Source Software* **3**, 861 (2018).

39. Aylward, J. *et al.* Altered learning under uncertainty in unmedicated mood and anxiety

disorders. *Nat Hum Behav* **3**, 1116–1123 (2019).

40. Grupe, D. W. & Nitschke, J. B. Uncertainty and anticipation in anxiety: an integrated

neurobiological and psychological perspective. *Nat. Rev. Neurosci.* **14**, 488–501 (2013).

41. Fahed, M. & Steffens, D. C. Apathy: Neurobiology, assessment and treatment. *Clin.*

*Psychopharmacol. Neurosci.* **19**, 181–189 (2021).

42. Pagonabarraga, J., Kulisevsky, J., Strafella, A. P. & Krack, P. Apathy in Parkinson's

disease: clinical features, neural substrates, diagnosis, and treatment. *Lancet Neurol.* **14**,

518–531 (2015).

43. Pessiglione, M., Vinckier, F., Bouret, S., Daunizeau, J. & Le Bouc, R. Why not try harder?

Computational approach to motivation deficits in neuro-psychiatric diseases. *Brain* **141**,

629–650 (2018).

44. Heimberg, R. G., Brozovich, F. A. & Rapee, R. M. *A Cognitive-Behavioral Model of Social*

*Anxiety Disorder: Update and Extension*. (2010).

45. Hofmann, S. G., Asnaani, A., Vonk, I. J. J., Sawyer, A. T. & Fang, A. The efficacy of

cognitive behavioral therapy: A review of meta-analyses. *Cognit. Ther. Res.* **36**, 427–440

(2012).

46. Beck, J. S. *Cognitive Behavior Therapy: Basics and Beyond*. (Guilford Press, 2011).

47. Craske, M. G. & Mystkowski, J. L. Exposure therapy and extinction: Clinical studies. in

*Fear and learning: From basic processes to clinical implications* 217–233 (American

Psychological Association, Washington, 2007).

48. Alexopoulos, G. S. & Arean, P. A model for streamlining psychotherapy in the RDoC era:

the example of "Engage." *Mol. Psychiatry* **19**, 14–19 (2014).

49. Yan, X., Yong, X., Huang, W. & Ma, Y. Placebo treatment facilitates social trust and

approach behavior. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 5732–5737 (2018).

50. Foroughi, C. K., Monfort, S. S., Paczynski, M., McKnight, P. E. & Greenwood, P. M.

Placebo effects in cognitive training. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 7470–7474 (2016).

51. Cuthbert, B. N. & Insel, T. R. Toward the future of psychiatric diagnosis: the seven pillars

of RDoC. *BMC Med.* **11**, 126 (2013).

52. Hinton, G. E. & Roweis, S. Stochastic neighbor embedding. *Adv. Neural Inf. Process. Syst.*

**15**, (2002).

53. Jolliffe, I. T. & Cadima, J. Principal component analysis: a review and recent developments.

*Philos. Trans. A Math. Phys. Eng. Sci.* **374**, 20150202 (2016).

54. Rhys, H. I. *Machine Learning with R, the Tidyverse, and Mlr*. (Simon and Schuster, New

York, NY, 2020).

55. Bari, A. *et al.* Serotonin modulates sensitivity to reward and negative feedback in a

probabilistic reversal learning task in rats. *Neuropsychopharmacology* **35**, 1290–1301

(2010).

56. Hayes, A. F. *Introduction to Mediation, Moderation, and Conditional Process Analysis: A*

*Regression-Based Approach. Xvii*. vol. 507 (Guilford Press, New York, NY, 2017).

57. Mackinnon, D. P., Lockwood, C. M. & Williams, J. Confidence Limits for the Indirect

Effect: Distribution of the Product and Resampling Methods. *Multivariate Behav. Res.* **39**,

99 (2004).

58. Preacher, K. J. & Hayes, A. F. Asymptotic and resampling strategies for assessing and

comparing indirect effects in multiple mediator models. *Behav. Res. Methods* **40**, 879–891

(2008).

59. Coelho, J. P., Pinho, T. M. & Boaventura-Cunha, J. *Hidden Markov Models: Theory and*

*Implementation Using Matlab* ®. (CRC Press, Boca Raton, FL : CRC Press, 2019. | "A

science publishers book.," 2019).

60. Chakroun, K., Mathar, D., Wiehler, A., Ganzer, F. & Peters, J. Dopaminergic modulation of
the exploration/exploitation trade-off in human decision-making. *Elife* **9**, (2020).

61. Forstmann, B. U., Ratcliff, R. & Wagenmakers, E.-J. Sequential sampling models in
cognitive neuroscience: Advantages, applications, and extensions. *Annu. Rev. Psychol.* **67**,
641–666 (2016).

62. Piray, P., Dezfouli, A., Heskes, T., Frank, M. J. & Daw, N. D. Hierarchical Bayesian
inference for concurrent model fitting and comparison for group studies. *PLoS Comput.*
*Biol.* **15**, e1007043 (2019).

63. Rigoux, L., Stephan, K. E., Friston, K. J. & Daunizeau, J. Bayesian model selection for
group studies — Revisited. *Neuroimage* **84**, 971–985 (2014).

64. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and
powerful approach to multiple testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).