Title

Spatial localization of hippocampal replay requires dopamine signaling

Authors

Matthew R. Kleinman[1] and David J. Foster[1,2]

Affiliations

1 Helen Wills Neuroscience Institute and Department of Psychology, University of California, Berkeley, CA 94720, USA

2 Lead contact

Correspondence

davidfoster@berkeley.edu, mattrkleinman@berkeley.edu

Abstract

Sequenced reactivations of hippocampal neurons called replays, concomitant with sharp-wave ripples in the local field potential, are critical for the consolidation of episodic memory, but whether replays depend on the brain's reward or novelty signals is unknown. Here we combined chemogenetic silencing of dopamine neurons in ventral tegmental area (VTA) and simultaneous electrophysiological recordings in dorsal hippocampal CA1, in freely behaving male rats experiencing changes to reward magnitude and environmental novelty. Surprisingly, VTA silencing did not prevent ripple increases where reward was increased, but caused dramatic, aberrant ripple increases where reward was unchanged. These increases were associated with increased reverse-ordered replays. On familiar tracks this effect disappeared, and ripples tracked reward prediction error, indicating that non-VTA reward signals were sufficient to direct replay. Our results reveal a novel dependence of hippocampal replay on dopamine, and a role for a VTA-independent reward prediction error signal that is reliable only in familiar environments.

Introduction

Spatial information is encoded in the firing of hippocampal place cells, which are thought to provide a cognitive map to support memory and navigation (O'Keefe and Dostrovsky, 1971; O'Keefe and Nadel, 1978). During pauses in locomotion, place cells participate in structured population bursts of activity, representing temporally-compressed trajectories through experienced locations, a phenomenon termed replay (Diba and Buzsáki, 2007; Foster and Wilson, 2006). Replay sequences can occur in the same order as experience, called forward replay, or in the reverse order of experience, called reverse replay (Diba and Buzsáki, 2007; Foster and Wilson, 2006). Replay appears after just one experience in a novel environment (Berners-Lee et al., 2022), and is preferentially generated towards goals in goal-directed tasks (Pfeiffer and Foster, 2013; Widloski and Foster, 2022). Experimentally interrupting or lengthening replay-associated sharp wave ripples (SWR) in the local field potential (LFP) disrupts and enhances learning of a spatial memory task, respectively (Fernández-Ruiz et al., 2019; Jadhav et al., 2012). Replay is thus thought to support memory consolidation and online planning (Buzsáki, 2015; Foster, 2017).

Intriguingly, reward drives increased rates of SWR (Singer and Frank, 2009), and
only reverse replay, not forward, is increased at highly rewarding locations (Ambrose et
al., 2016). Theoretical work suggests replay functions to update spatial representations
of value to influence behavior and optimize reward receipt (Mattar and Daw, 2018).
These findings hint that replay may be strongly modulated by reward-processing areas
in the brain, such as the midbrain dopamine system (Fields et al., 2007). Dopamine
neuron activity in the ventral tegmental area (VTA) is consistent with coding of reward
prediction error (RPE), with increased activity at unexpected rewards and decreased
activity with omission of expected rewards (Schultz et al., 1997). Subsequent work
investigated dopamine release in spatial tasks, finding it ramps towards large rewards
(Guru et al., 2020; Howe et al., 2013) in a manner consistent with encoding of RPE for a
value function over space (Kim et al., 2020).
Besides the well-established role of midbrain dopamine neurons in reward
processing, dopamine release in hippocampus has been implicated in stabilizing place
fields (Kentros et al., 2004), gating the increase in plasticity in dorsal CA1 synapses by
novel experiences (Li et al., 2003), and improving memory retention via increasing
replay (McNamara et al., 2014). Furthermore, VTA activity is increased in novel
environments (Guru et al., 2020; McNamara et al., 2014; Takeuchi et al., 2016),
suggesting the hippocampus and VTA may coordinate to signal spatial novelty and
induce learning in new environments (Lisman and Grace, 2005). However, recent work
implicates locus coeruleus (LC) as the dominant source of dopaminergic input to dorsal
CA1 and show its necessity and sufficiency for novelty-mediated episodic memory
consolidation (Guru et al., 2020; McNamara et al., 2014; Takeuchi et al., 2016), leaving
the role of VTA unclear.
We therefore tested whether VTA dopamine neurons are required for reward-
related modulation of SWR and replay. We expressed an inhibitory DREADD
(Armbruster et al., 2007) in VTA dopamine neurons and implanted a tetrode microdrive
above hippocampus. We could then inhibit VTA dopamine signaling and simultaneously
record neural activity in the dorsal CA1 region of hippocampus while rats collected
rewards in familiar and novel environments. If VTA dopamine signaling is required for
coordinating replay to valuable locations, we expected to see deficits in the capacity for
reward to recruit SWR and replay. Additionally, if VTA dopamine is critical for inducing
plasticity in CA1 in novel environments, novelty may significantly increase the effect of
VTA inactivation on hippocampal replay.
81
Results
VTA inactivation in a simple spatial task with reward changes
We combined tetrode recordings in dorsal CA1 (dCA1) and chemogenetic
silencing of VTA dopamine neurons to determine whether reward-related changes in
hippocampal ripples and replay required VTA dopamine signaling. Transgenic rats
expressing cre-recombinase under the tyrosine hydroxylase (TH) promoter were
stereotactically injected with cre-dependent virus containing the inhibitory DREADD
hM4Di (Experimental, n=4) or mCherry-only control (Control, n=3) into bilateral VTA
(Figure 1A). We observed widespread expression across the extent of VTA and co-
localization with TH (Figure 1B, Figure 1 – supplement 1), enabling specific and
reversible inactivation of VTA dopamine signaling. Post-experiment histology confirmed

overlapping virus expression and TH-positive neurons in putative VTA near the injection
site (-5.6 mm AP from bregma), as well as approximately 0.5 mm anterior and posterior
(-5 to -6 mm AP). Recording microdrives containing 6-32 independently adjustable
96

97
Figure 1. Experimental design and linear track behavior.
(A) TH-cre rats underwent stereotactic surgery to inject virus bilaterally into VTA and implant a tetrode
microdrive above dorsal CA1. **(B)** Co-expression of mCherry (red) and TH (green) in VTA from three
example animals. Left panel, mCherry-only virus, scale bar 600 µm; middle panel, hM4Di-mCherry, scale
bar 150 µm; right panel, hM4Di-mCherry, scale bar 75 µm. **(C)** Intraperitoneal injection of saline or CNO
(1-4 mg/kg) preceded recording sessions by at least 10 minutes. Rats were placed at one end of a linear
track and collected liquid chocolate reward from wells at each end. Each epoch lasted 10-20 laps and
reward changes were unsignaled to the animal. For each session, the Incr. end was defined as the
reward end with 4X reward in Epoch 2, and the Unch. end was defined as the reward end with 1X reward
in Epoch 2. **(D)** During stopping periods at reward ends, LFP was bandpass filtered in the ripple band
(150-250 hz) and SWR events were detected. **(E)** Three example ripple-filtered LFP traces from one lap
(two stopping periods) are shown. **(F)** Cumulative distribution of reward end stopping periods at the Unch.
reward end in Epoch 1 and 2 for experimental rats (left panel) and control rats (right panel). See also
Figure 1 – Supplement 2. **(G)** The duration of Unch. reward end stopping periods decreased from Epoch
1 to Epoch 2. The mean stopping durations in each condition were calculated per session, then analyses
performed across sessions. Mean ± standard error, Exp Saline, Epoch 1: 6.67±0.61, Epoch 2: 5.13±0.53;
Exp CNO, Epoch 1: 7.766±0.61, Epoch 2: 6.04±0.42. Con Saline, Epoch 1: 6.96±0.4, Epoch 2:
4.77±0.27; Con CNO, Epoch 1: 6.59±0.29, Epoch 2: 4.43±0.15. Mixed-effects GLM with epoch, drug,
animal group, and all interactions, with individual animal as a random effect: epoch, z=-3.37, p<0.001; all

other terms, n.s. **(H)** Cumulative distribution of reward end stopping periods at the Incr. reward end in
Epoch 1 and 2 for experimental rats (left panel) and control rats (right panel). **(I)** The duration of Incr.
reward end stopping periods increased from Epoch 1 to Epoch 2. The mean stopping durations in each
condition were calculated per session, then analyses performed across sessions. Mean ± standard error,
Exp Saline, Epoch 1: 6.64±0.54, Epoch 2: 10.87±0.8; Exp CNO, Epoch 1: 7.28±0.66, Epoch 2:
12.31±0.82. Con Saline, Epoch 1: 7.44±0.78, Epoch 2: 11.29±0.5; Con CNO, Epoch 1: 6.47±0.35, Epoch
2: 10.34±0.27. Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual
animal as a random effect: epoch, $z=4.62$, $p<10^{-5}$; all other terms, n.s. *, $p<0.05$; **, $p<0.01$.

tetrodes (bilateral 32 tetrode, n=4; unilateral 20 tetrode, n=2; unilateral 6 tetrode, n=1)
were implanted above dCA1 (Fig. 1A). Tetrodes were lowered to the pyramidal cell
layer of dCA1, using the presence of sharp-wave ripples with upward deflections in the
LFP, recording depth characteristic of dCA1, and spatially-restricted firing of place cells
to confirm the recording location.
Before each experimental session, rats were given intraperitoneal injection of
CNO, to activate hM4Di receptors and suppress VTA dopamine neuron activity, or
saline, then performed a simple task on linear tracks (1.5 to 2.5 m in length), collecting
liquid chocolate rewards from each end (Fig. 1C). Each session began with equal 0.1 ml
reward volume at each end (Epoch 1) for 10-20 laps (1 lap was comprised of reward
collection at both ends; mean, 16 laps). This was followed by unsignaled quadrupling of
reward at one end to 0.4 ml (Incr. end), while reward at the other end remained
unchanged (Unch. end), for 10-20 laps (Epoch2; mean, 16.7 laps). Finally, reward was
equalized again to 0.1 ml at both ends (Epoch 3) for up to 20 laps (mean, 11.6 laps).
Each animal performed this task on familiar linear tracks (>2 sessions on track;
total session count for each condition: Experimental rats: 36 saline, 34 CNO; Control
rats: 23 saline, 25 CNO) and novel linear tracks (1st or 2nd session on track;
Experimental rats: 9 saline, 10 CNO; Control rats: 12 saline, 12 CNO). During stopping
periods at either end of the track (velocity ≤ 8 cm/s, position ≤ 10 cm from end), SWR
were identified as peaks in the ripple band (150-250 hz) in LFP (Fig. 1D-E; see
Methods).
Gross behavior was largely unaffected by VTA suppression (e.g., all reward
consumed on each lap), but CNO in experimental animals systematically affected
stopping period duration. Visits to the Unch. end in Epoch 2 were significantly shorter
than in Epoch 1, despite unchanged reward volume there, and this reduction was
present in both saline and CNO sessions (Fig. 1F-G). CNO did not affect control
animals (Fig. 1F-G). Visits to the Incr. end were significantly longer in Epoch 2 than in
Epoch 1 in all conditions, owing to the increased reward consumption time (Fig. 1H-I).
Changes in stopping period duration in Epoch 3 were similar across all conditions:
Unch. visit duration increased from Epoch 2 to Epoch 3 (Fig. 1 – Supplement 2E), while
Incr. visit duration decreased (Fig. 1 – Supplement 2F). Separate analysis of novel and
familiar sessions revealed the pattern of shorter duration Unch. visits in Epoch 2
compared to Epoch 1 did not depend on novelty (Fig. 1 – Supplement 2A-D). However,
the main effect of CNO in experimental rats of prolonging stopping periods occurred in
novel sessions (Fig. 1 – Supplement 2A-B), not in familiar sessions (Fig. 1 –
Supplement 2C-D). Rats consistently ran slightly faster towards the Incr. end than the
Unch. end in Epoch 2, across all conditions (Fig. 1 – Supplement 3).

We interpret the reduction in visit duration as a behavioral signature of the Unch.
end becoming relatively less valuable during Epoch 2, when the reward volume is larger
at the Incr. end. Though visit durations were slightly longer in CNO sessions in
experimental rats, particularly in novel track sessions, this behavioral effect of a relative
value decrease remained, indicating VTA inactivation did not prevent rats from
recognizing a devalued location.
170
Reward-related modulation of SWR rate is mediated by novelty and VTA
We analyzed the rate of SWR occurrence (number of SWR / total duration rat
was stationary) during the first 10 s of each stopping period, when rats were consuming
reward. In individual sessions, SWR rate increased robustly in all conditions shortly after
stopping at the reward wells and beginning reward consumption (Fig. 2A). Relative to
Epoch 1, SWR rate during Epoch 2 tended to increase dramatically at Incr. end visits
and decrease at Unch. end visits. During Epoch 3, SWR rate at the Incr. end dropped
precipitously relative to Epoch 2, while often increasing at the Unch. end (Fig. S4A).
Surprisingly, during novel sessions, VTA inactivation often led to increased SWR
rate at both reward ends (Fig. 2A, right). SWR rate still increased in Epoch 2 at the Incr.
end even without normal VTA signaling, indicating reward sensitivity per se was not
abolished, but suggesting the localization of this increased SWR rate to where reward
increased was disrupted.
Pooling across sessions revealed this dramatic increase in SWR rate at the
Unch. end was typical with CNO in novel sessions, and further suggested there was a
reduction in the difference in SWR rate between the Incr. and Unch. ends in Epoch 2 in
both familiar and novel experiences (Fig. 2B). We therefore used a Poisson generalized
linear model (GLM) to quantify the changes in SWR rate across reward end, epoch,
drug condition, and novelty (see Methods). In experimental rats, CNO and reward both
influenced SWR rate, with significant effects for the CNO main effect ($z=3.19$, $p<0.01$),
the interaction between Incr. end and Epoch 2 ($z=9.02$, $p<10^{-10}$) and the three-way
interaction between Incr. end, Epoch 2, and CNO ($z=-2.06$, $p<0.05$). Control animals
showed no apparent effect of CNO (Fig. 2C). The same Poisson GLM fit to control rat
data confirmed this, with significant coefficients only for Incr. end ($z=-2.42$, $p<0.05$) and
the interaction between Incr. end and Epoch 2 ($z=7.64$, $p<10^{-10}$). SWR duration was
reduced in novel sessions, consistent with replays becoming longer with increased
familiarity (Berners-Lee et al., 2021), as well as in Epoch 2, but was otherwise
unaffected by reward or drug (Fig. 2 – Supplement 4).
To assess the interaction between VTA inactivation and novelty, we fit the
Poisson GLM separately to novel and familiar sessions, then used bootstrapping to
generate distributions of SWR rates for each reward end and condition under the null
hypothesis that CNO had no effect (see Methods). We found the actual difference in
SWR rates between saline and CNO sessions in experimental animals during Epoch 2
was significantly greater than chance (Fig. 2 – Supplement 1A) in novel sessions for
both the Unch. end (one-tailed test, CNO>saline, $p<0.001$) and the Incr. end (one-tailed
test, saline>CNO, $p<0.01$), as well as at the Unch. end in familiar sessions (one-tailed
test, CNO>saline, $p<0.01$). There was no significant difference between saline and CNO
in control animals at either reward end in either familiar or novel sessions (Fig. 2 –
Supplement 1B; one-tailed tests, all n.s.).

A potential functional role for the reward-related changes in SWR rate is to
strengthen downstream representations of particularly rewarding locations at the
expense of less rewarding locations. We tested whether VTA suppression blunted the
SWR rate difference between reward ends in Epoch 2. Across both familiar and novel
214

Figure 2. Modulation of SWR rate by reward, novelty, and VTA inactivation.
(A) SWR rate as a function of time in stopping period in Epoch 1 and 2 for four example sessions in
experimental rats; from left to right, saline on familiar track, saline on novel track, CNO on familiar track,
and CNO on novel track. In each panel, visits to the Incr. end are on the left and visits to the Unch. end
are on the right. Relative to Epoch 1 (black lines), in Epoch 2 (red lines) SWR rate increased at Incr. end
and decreased at Unch. end in all conditions except for CNO on a novel track (far right), where SWR rate
increased at both ends in Epoch 2. SWR rate was binned in 0.25 s windows and smoothed with a two-bin
Gaussian. Line, mean; shading, standard error. **(B)** SWR rate in experimental rats as a function of epoch,
drug (saline in solid lines, CNO in dashed lines), reward end (Unch. in black, Incr. in green), and novelty
(familiar in left panel, novel in right panel). A mixed-effects Poisson generalized linear model (GLM) was
fit to predict changes in SWR rate across reward end, epoch, drug condition, and novelty, with animal
identity as a random effect. Significant coefficients: CNO ($z=3.19$, $p<0.01$), the two-way interaction Incr.
end × Epoch 2 ($z=9.02$, $p<10^{-10}$), and the three-way interaction between Incr. end × Epoch 2 × CNO ($z=$-
2.06, $p<0.05$). *, $p<0.05$; **, $p<0.01$. **(C)** SWR rate in control rats as a function of epoch, drug (saline in
solid lines, CNO in dashed lines), reward end (Unch. in black, Incr. in green), and novelty (familiar in left
panel, novel in right panel). The same Poisson GLM fit to control rat data had significant coefficients: Incr.
end ($z=-2.42$, $p<0.05$) and the two-way interaction Incr. end × Epoch 2 ($z=7.64$, $p<10^{-10}$). **(D)** Difference
between SWR rate at Incr. and Unch. ends in Epoch 2 in Experimental rats. Full stopping period, left
panel. Trimmed stopping period, with first 1 s and last 1 s of visit excluded to eliminate all slow
approaching/leaving movement, right panel. Saline, gray bars; CNO, white bars. Mean and standard
error. A mixed-effects model with drug, novelty, and their interaction, and animal-specific intercepts ("full
model") was compared to reduced model lacking the drug terms ("reduced model"). Full stopping periods:
full model, intercept, $z=6.05$, $p<10^{-5}$; other terms, n.s. $AIC_{reduced}-AIC_{full}=5.22$. Trimmed stopping periods:
full model, intercept, $z=3.5$, $p<0.001$; other terms, n.s. $AIC_{reduced}-AIC_{full}=4.55$. **(E)** Difference between
SWR rate at Incr. and Unch. ends in Epoch 2 in Control rats, as in (D). Full stopping periods: full model,
intercept, $z=3.95$, $p<10^{-5}$; other terms, n.s. $AIC_{reduced}-AIC_{full}=-3.54$. Trimmed stopping periods: full model,
novelty, $z=2.31$, $p<0.05$; other terms, n.s. $AIC_{reduced}-AIC_{full}=-3.1$. **(F)** In experimental rats, the difference in
SWR rates at each reward end (Incr. – Unch.) in Epoch 2, after subtracting the mean rates in Epoch 1,
averaged over a 5-lap sliding window within Epoch 2. Blue lines, novel sessions. Gray lines, familiar
sessions. Blue and gray asterisks denote the centers of sliding windows in which the difference in SWR
rate was significantly greater than 0 in novel and familiar sessions, respectively (one-sample t-test,
$p<0.05$, uncorrected for multiple comparisons). Shading denotes 95% confidence interval. See also
Figure S4. **(G)** As in (F), but for control animals.

environments, CNO reduced the difference in SWR rate between the Incr. end and
Unch. end in experimental rats but not in control rats (Fig. 2D-E, left panels;
experimental group, full mixed-effects model with drug terms strongly outperformed
reduced model lacking drug terms, $AIC_{reduced}-AIC_{full}=5.22$; control group, reduced model
lacking drug terms strongly outperformed the full model, $AIC_{reduced}-AIC_{full}=-3.54$). We
saw little difference in this measure when comparing sessions that occurred first in each
day to those that occurred later (Fig. 2 – Supplement 3), and little behavioral evidence
from running velocity or number of laps completed that motivation systematically varied
across session number. To control for the possibility that VTA inactivation caused
changes in locomotor or other non-consummatory behavior at reward wells that might
affect SWR emission, we omitted the first and last 1 s of each stopping period to isolate
the reward consumption period. The effect of CNO remained in experimental rats,
reducing SWR rate discrimination between Incr. and Unch. ends (Fig. 2D-E, right
panels; experimental group, full mixed-effects model with drug terms strongly
outperformed reduced model lacking drug terms, $AIC_{reduced}-AIC_{full}=4.55$; control group,
reduced model lacking drug terms strongly outperformed the full model, $AIC_{reduced}-$
$AIC_{full}=-3.1$).

We next looked for within-epoch changes in SWR rate to determine whether VTA
inactivation altered the dynamics of the response to reward changes. We calculated the
difference in SWR rate at the Incr. and Unch. reward ends (each with its Epoch 1 mean
subtracted) in a 5-lap sliding window. In all time windows and conditions except novel
CNO sessions in experimental rats, the SWR rate at the Incr. end was significantly
greater than at the Unch. end (Fig. 2F-G; Incr. – Unch. significantly greater than 0, one-
sample t-test, $p<0.05$, uncorrected for multiple comparisons).
In novel sessions, VTA inactivation did not prevent an initially larger increase in
SWR rate at the Incr. end than the Unch. end, but caused that difference to diminish
over laps (Fig. 2F). By the middle of the epoch, there was no statistically-significant
difference in reward modulation between the reward ends (one-sample t-test, $p>0.05$,
for 5-lap windows centered on laps 8-13 and 15), consistent with an initial appropriately-
localized reaction to reward change that eventually spread across the track. We found a
similar deficit in Epoch 3, with SWR rate decreasing significantly more compared to
Epoch 2 at the Incr. end than the Unch. end (Incr. – Unch. significantly below 0, one-
sample t-test, $p<0.05$, uncorrected for multiple comparisons) for almost every task
condition and timepoint except in novel CNO sessions in experimental rats (Fig. 2 –
Supplement 2B-C). This suggests VTA inactivation may disrupt the normal magnitude
or time course of the SWR response to negative value changes as well.
Overall, VTA inactivation spared the capacity for increased reward to modulate
SWR rate but led to decreased differentiation of low and high value locations,
particularly in novel environments where SWR rate increased spatially-indiscriminately.
290
SWR rate is correlated with RPE even with VTA inactivation
Taken together, the above results demonstrate VTA inactivation caused changes
in the normal dynamics of the response of SWR rate to positive and negative changes
in reward value (Fig. 2). However, because each session had only two timepoints when
reward value changed by fixed amounts, our experiment was not optimized to probe the
precise relationship between SWR rate and reward changes. Additionally, the effect of
VTA inactivation was particularly prominent with novelty, when both SWR rate and its
modulation by reward changes were greater, raising the possibility that large SWR rates
and fluctuations, rather than novelty per se, depend on VTA dopamine signaling.
To address these questions, we designed a volatile reward schedule (Experiment
2) with frequent, large reward changes at one end of the linear track, and tested
whether VTA inactivation impacted the capacity for SWR rate to track value (Fig. 3A,
top). The "stable end" delivered 0.2 ml every lap, while the "volatile end" reward volume
was drawn pseudorandomly from 0, 0.1, 0.2, 0.4, and 0.8 ml (mean 0.37 ml; blocks of
20 laps were comprised of 3 laps x 0 ml, 4 laps x 0.1 ml, 3 laps x 0.2 ml, 4 laps x 0.4 ml,
and 6 laps x 0.8 ml). The position of the stable and volatile ends randomly varied across
sessions.
This reward schedule also allowed us to test whether SWR rate was correlated
with value, RPE, or neither. We expected SWR rate at the volatile end would be
predominantly determined by the current reward volume there, but potentially also
modulated by previous reward volumes (Fig. 3A, bottom). If SWR rate is correlated with
value, then for a given current volume, larger reward volumes at the last visit will lead to
higher SWR rates compared to when the last visit was a smaller reward volume.

Conversely, if SWR rate is correlated with RPE, the opposite modulation by last reward
volume will be observed: the larger the previous reward volume, the lower the current
SWR rate.
A subset of rats performed sessions of the modified reward schedule (mean 53.4
laps per session; Total sessions per condition: Experimental rats 2 and 4: 6 saline, 7
CNO; Control rats 1-3: 16 saline, 18 CNO). Each rat completed 1-2 saline sessions
before any CNO sessions and all sessions were on the same linear track, meaning
almost all CNO sessions took place on a familiar track. As expected, SWR rate at the
volatile end was predominantly determined by the current volume (Fig. 3B-C). There
was little obvious difference between saline and CNO in either experimental (Fig. 3B) or
control rats (Fig. 3C). SWR rate at the stable end was largely stable across laps,
although there was a trend towards higher SWR rate if the most recent volatile end visit
was lower volume, consistent with lap-by-lap changes in the relative value of the stable
end (Fig. 3 – Supplement 1).
We next investigated how SWR rate at the volatile end varied as a function of
both current and immediately previous volatile end volume in experimental rats (Fig. 3D-
E, top panels). For each current volume, we subtracted the mean SWR rate across all
previous volumes, and examined how previous volume affected the mean-subtracted
SWR rates across all current volumes (Fig. 3D-E, middle panels).
The mean-subtracted SWR rate was modestly negatively correlated with the
previous volume (Pearson correlation: saline, r=-0.076, p=0.177; CNO, r=-0.109,
p<0.05). A GLM found the mean-subtracted SWR rate was significantly affected by
previous volume (z=-2.3, p<0.05), but not drug (z=-0.24, p=0.81) or current volume
(z=0.196, p=0.844). There was no similar behavioral effect: for each current volume, we
subtracted the mean reward end visit duration, and found no correlation between the
previous reward volume and mean-subtracted visit duration (Pearson correlation; saline,
r=0.011, p=0.844; CNO, r=-0.075, p=0.175).
We next separated visits to the volatile end using a median split based on the
recent volumes (mean of previous 3 visits). SWR rates were higher for a given current
reward volume when the recent reward history was low (Fig. 3D-E, bottom panels). A
Poisson GLM predicting SWR rate as a function of current volume, drug condition, and
whether reward history was low or high (relative to the median for the session) revealed
significant effects of current volume (z=13.86, $p<10^{-10}$), as expected, and reward history
split (z=-2.23, p<0.05), but not drug condition (z=-1.05, p=0.29).
Finally, we separated combinations of current and previous volume into those
with negative RPE (current < previous) and positive RPE (current > previous) and found
mean-subtracted SWR rate was significantly affected by RPE sign (Fig. 3F; two-way
ANOVA with drug and RPE sign, RPE sign: F[1,518]=6.42, p<0.05), but not drug
(F[1,518]=0.3, p=0.582; drug X RPE sign, F[1,518]=0.07, p=0.785).
Given the lack of an effect of drug in experimental rats, we pooled all sessions
(both animal groups, both drug conditions) in Experiment 2 to maximize experimental
power and found similar results as in just experimental rats (Fig. 3 – Supplement 2). On
top of a large increase in SWR rate with current volume at the volatile end (Fig. 3 –
Supplement 2A-B), SWR rate was also significantly negatively correlated with the
previous volatile volume, both at the volatile end (Fig. 3 – Supplement 2C) and at the
stable end (Fig. 3 – Supplement 2E). Accordingly, SWR rates were significantly lower

Figure 3. Frequent reward changes modulated SWR rate.

(A) Recording sessions in the volatile reward task were preceded by intraperitoneal injection of saline or CNO by at least 10 minutes. Rats were placed on the stable end to begin each session, which delivered 0.2 ml reward at each visit, while the volatile end delivered 0, 0.1, 0.2, 0.4, or 0.8 ml, pseudorandomly chosen on each lap. Bottom panel, schematic of how value and RPE would modulate SWR. Given a particular current volume, value coding predicts a positive correlation between SWR rate and previous volume, while RPE coding predicts a negative correlation. **(B)** SWR rate as a function of reward volume and time in end visit in example rat, experimental rat 4. Left panel, saline. Right panel, CNO. In stable panel, traces are colored based on previous volatile end visit volume. In volatile panel, traces are colored based on current volatile volume. See also Figure 3 – Supplement 1 and 2. **(C)** SWR rate as a function of reward volume and time in end visit in example control rat 3, as in (B). **(D)** Top panel, SWR rate at volatile end as a function of current and previous volatile volume, for saline sessions in experimental rats. Middle panel, SWR rate for each non-zero volatile volume plotted as a function of previous volume, with the mean SWR rate for that current volume subtracted. Unfilled symbols, mean of previous volume across all current volumes. Thick dashed line, linear fit to mean values. Pearson correlation between (ripple rate – mean) and previous volume, r=-0.076, p=0.177. Error bars, standard error. Bottom panel, SWR rate as a function of reward volume, separated by recent reward history (median split on average of last 3 visits). Black, recent history below median; red, recent history above median. **(E)** Same as (D), for CNO sessions in experimental rats. Middle panel, Pearson correlation between (ripple rate – mean) and previous volume, r=-0.109, p<0.05. GLM fitting SWR rate as a function of drug, current volume, and previous volume: previous volume, z=-2.31, p<0.05; drug and current volume, both p>0.8. Bottom panel, Poisson GLM fitting ripple rate as a function of volume, drug condition, and reward history (above/below median), with animal-specific intercept as random effect: volume, z=13.86, p<10^{-10}; history, z=-2.23, p<0.05; drug, z=-1.05, p=0.29. **(F)** The RPE of volatile end visits were calculated by subtracting the previous volatile volume from the current volume. Two-way ANOVA with drug and RPE sign (+/-): drug (F[1,518]=0.3, p=0.582), RPE sign (F[1,518]=6.42, p<0.05), drug X RPE sign (F[1,518]=0.07, p=0.785).

for negative RPE than for positive/non-negative RPE (Fig. 3 – Supplement 2D-F). Finally, recent reward history at the volatile end significantly affected SWR rate (Fig. 3 – Supplement 2G), with higher SWR rate when recent rewards were lower.

Taken together, SWR rate was modulated by reward volume changes consistent with RPE-like coding. This modulation did not require normal VTA dopamine signaling, at least in familiar environments. The lack of effect of VTA inactivation, even with frequent, large swings in value and SWR rate, corroborates the results from Experiment 1 that novel experiences are particularly susceptible to disruption, indicating VTA dopamine release is critical when learning new reward locations.

Rate of reverse replay is increased with reward in novel environments only with intact VTA signaling

Previous work discovered the incidence rate of reverse replay, but not forward replay, was increased at locations with increased reward (Ambrose et al., 2016). We therefore analyzed single unit data collected in Experiment 1 (excluding experimental rat 2, who had a 6-tetrode recording drive) to determine whether this modulation of replay required VTA dopamine signaling. As previously observed (e.g., Ambrose et al., 2016), place cells in dCA1 had directional fields, such that the location a neuron was active while the rat moved in one direction on the track (e.g., "upward") was often distinct from its activity when the rat moved in the other direction (e.g., "downward"). This directionality was apparent in both familiar and novel sessions, including in experimental rats with either saline or CNO (Fig. 4A). We found no effect of CNO on within-session field reliability, but significantly less reliability in novel compared to familiar sessions (Fig. 4 – Supplement 1A). Field similarity across running directions

was slightly but significantly increased by both CNO and novelty (Fig. 4 – Supplement
1B).
Two place fields were defined for each neuron, one for each running direction,
permitting Bayesian decoding methods to estimate both position and direction from
neural activity (Fig. 4B-C). Sessions with accurate position and direction decoding
during run, primarily due to sufficiently high cell yield (cell count, mean±sem; included
sessions: 26.1±1.1, excluded sessions: 9.5±1.6; two-sample t-test, t(133)=5.3, p<10^{-5})
and also marginally smaller average place field size (mean place field size, defined as
number of spatial bins with >1 hz firing rate, mean±sem: included sessions: 47.7±1.3,
excluded sessions: 57.7±5.8; two-sample t-test, t(133)=-2.33, p<0.05), were included for
replay analysis (Fig. 4 – Supplement 2; total sessions included: Experimental rats: novel
saline, n=8; novel CNO, n=8; familiar saline, n=18; familiar CNO, n=23; Control rats:
novel saline, n=12; novel CNO, n=11; familiar saline, n=16; familiar CNO, n=17).
Candidate replay events were periods of high population spiking activity while the
rat was not running (z-scored spike count>3, minimum duration of 50 ms, rat velocity≤8
cm/s). We used a memory-less Bayesian decoder, with 40 ms decoding windows
advancing by 5 ms steps, to estimate position and direction from neural activity.
Replays were defined as candidate events with spatial trajectories meeting a threshold
for motion and minimum total posterior in one running direction map (Ambrose et al.,
2016) (see Methods), with the running direction with greater posterior probability used to
classify replay directionality, described below.
Forward replays were spatial trajectories moving across the track in the same direction
as the rat when fields were calculated, e.g., moving "downward" with posterior
probability in the "downward" place field map (Fig. 4B, left panel). Reverse replays were
trajectories that moved in the opposite direction as the rat, e.g., moving "upward" with
posterior probability in the "downward" place field map or vice versa (Fig. 4B, middle
and right panels). We found forward and reverse replays occurred in all conditions,
including in novel sessions with saline (Fig. 4B) or CNO (Fig. 4C). We therefore asked
how novelty and drug condition influenced the effect of reward change on rates of
reverse and forward replay.
Consistent with previous work (Ambrose et al., 2016), the rate of reverse replay
was strongly modulated by the reward volumes on the track. Excluding novel CNO
sessions for the moment, in all other conditions in experimental rats, when reward was
larger at the Incr. end than the Unch. end (unequal reward), reverse replay was
significantly increased at the Incr. end relative to when rewards were equal (novel
saline: equal reward, 0.0019±0.0009 replay/s; unequal reward, 0.0121±0.004 replay/s;
two-sample t-test, t(420)=3.235, p=0.0013; familiar saline: equal reward, 0.0033±0.0012
replay/s; unequal reward, 0.0128±0.0025 replay/s; two-sample t-test, t(907)=3.822,
p=0.0001; familiar CNO: equal reward, 0.0033±0.001 replay/s; unequal reward,
0.0139±0.0022 replay/s; two-sample t-test, t(1153)=5.234, p<10^{-6}). The rate of reverse
replay was not significantly increased at the Unch. end with unequal rewards in any of
these conditions (all p>0.05). This led to a bias for reverse replay to preferentially occur
at the Incr. end when rewards were unequal (Fig. 4D-E). In control rats, reward changes
caused similar changes to the balance of reverse replay (Fig. 4H-I), with a significantly
larger swing in reverse replay bias in novel sessions (mixed-effects model with drug,
novelty, and replay directionality: novelty X directionality, z=2.18, p<0.05; all other

terms, n.s.). The lack of effect of reward in familiar sessions in control rats (Figure 4I)
was likely due to a dearth of replay in these sessions, with mean reverse replay rates at

Figure 4. Replay recruitment by reward change in novel sessions requires VTA signaling.

(A) Place cells exhibit directional place fields on the linear track. Fields calculated from movement in a particular direction ("right" fields and "left" fields), ordered based on field center location in either running direction ("right" order and "left" order). Color denotes z-scored firing rate, from high (yellow) to low (blue).

Example saline session and CNO session from experimental rat 3. See also Figure 4 – Supplement 1 and
2. **(B)** Three example replays from Epoch 2 of a novel saline session from experimental rat 3. Red,
posterior in upwards map; blue, posterior in downwards map. Title indicates reward end (Incr., Unch.) and
replay direction (Reverse, Forward). The horizontal black line indicates rat position. **(C)** Three example
replays from Epoch 2 of a novel CNO session from experimental rat 3, as in (B). **(D)** The difference in rate
of reverse replay at each end (Incr. – Unch.) in novel sessions in experimental rats. Error bars, standard
error of the mean. Reward condition is indicated by color (equal reward, epoch 1 and 3, gray; unequal
reward, epoch 2, orange), and drug condition is indicated on the x-axis. The difference in replay rate
between equal and unequal reward conditions was assessed with a mixed-effects linear model with drug,
novelty, and replay directionality, and animal-specific intercepts as random effect: drug X novelty X
directionality, z=-2.27, p<0.05; novelty, z=-2.01, p<0.05; novelty X directionality, z=2.15, p<0.05; all other
terms, n.s. **(E)** Same as (D), but for familiar sessions. **(F)** Same as (D), but for forward replay. **(G)** Same
as (F), but for familiar sessions. **(H)** Same as (D), but for control rats. The difference in replay rate
between equal and unequal reward conditions was assessed with a mixed-effects linear model with drug,
novelty, and replay directionality, and animal-specific intercepts as random effect: novelty X directionality,
z=2.18, p<0.05; all other terms, n.s. **(I)** Same as (H), but for familiar sessions. **(J)** Same as (H), but for
forward replay. **(K)** Same as (J), but for familiar sessions.
484
485
both reward ends under 0.008 replays/s during Epoch 2, compared to greater than
0.013 replays/s at the Incr. end in Epoch 2 in novel sessions. Reward changes caused
no consistent effects in the rates of forward replay (Fig. 4F-G, Fig. 4J-K), nor in the
rates of replay that began far from the animal (non-local replay, Fig. 4 – Supplement 3).
Conversely, in novel CNO sessions in experimental rats, reverse replay rate
failed to be biased towards the larger reward location (Fig. 4D). With unequal reward,
the rate of reverse replay did increase at the Incr. end (equal reward, 0.0053±0.0016
replay/s; unequal reward, 0.0116±0.0029 replay/s; two-sample t-test, t(332)=2.043,
p<0.05), but also increased somewhat at the Unch. end (equal reward, 0.0173±0.004
replay/s; unequal reward, 0.0249±0.0068; two-sample t-test, t(319)=1.019, n.s.), leading
to no consistent change in the difference at the two reward ends. This effect of VTA
inactivation on the bias of replay between the reward ends when reward contingencies
changed was specific to novel sessions and reverse replay (mixed-effects model with
drug, novelty, and replay directionality: drug X novelty X directionality, z=-2.27, p<0.05;
novelty, z=-2.01, p<0.05; novelty X directionality, z=2.15, p<0.05; all other terms, n.s).
VTA dopamine signaling was therefore required to direct reward-related changes in
reverse replay, specifically in novel environments.
503
Discussion
Here we demonstrated a critical role for VTA dopamine signaling in driving
hippocampal SWR and reverse replay selectively to locations with increased reward.
Surprisingly, we found this was true only in novel environments, with only modest
effects of VTA inactivation on SWR rates and no discernible effect on replay rates in
environments that had been explored several times before. We additionally recorded
activity in a modified task that allowed us to differentiate SWR rate modulation by value
and RPE. While SWR rate was modulated by RPE, VTA inactivation had little effect on
this RPE-like modulation, suggesting that at least in familiar environments normal VTA
dopamine signaling is dispensable for this reward-related hippocampal activity.
Our work has several limitations. First, we did not directly measure the
suppression of VTA neurons after CNO injection. Previous work in other brain areas

found hM4Di activation suppressed firing rates to around 60% of baseline (Mahler et al.,
2014; Chang et al., 2015), in addition to diminished synaptic transmission even when
spikes occurred (Stachniak et al., 2014). Combined with the incomplete expression of
hM4Di in TH-positive neurons in our animals, we expect VTA activity was significantly
but not completely suppressed. Because our results depend only on any degree of
blunting differences in dopamine release at different reward locations, rather than the
total absence of dopamine signaling, measuring the magnitude of suppression was not
essential for our conclusions. Second, we chose simple behavioral tasks specifically to
isolate the effect of reward changes per se on hippocampal replay, distinct from learning
or performance effects. In more complex tasks, replay overrepresents high value
locations (Pfeiffer and Foster, 2013; Widloski and Foster, 2022), which may aid in
consolidating memory of and planning routes to those locations. In our hands, VTA
inactivation reduced the SWR rate difference between high and low value locations in
novel environments by about 2/3 (Figure 2D), which would greatly reduce any effect of
reward-biased replay in learning more complex environments or tasks. Finally, we found
no effect of VTA inactivation on SWR rates in familiar environments even with large and
frequent reward changes (Figure 3), but replicating this result with more animals and
probing the interaction between volatile reward changes and environmental novelty are
important future directions.
Why is VTA inactivation particularly disruptive during novel experiences?
Dopamine neuron firing rates are elevated in novel environments (McNamara et al.,
2014; Takeuchi et al., 2016). More specifically, early in experience dopamine neuron
activity ramps while mice run towards both larger and smaller rewards and this ramping
activity declines over experience until modest ramps persist only towards the larger
reward (Guru et al., 2020). Activation of VTA projections to dorsal CA1 improves
retention of spatial learning of a novel maze configuration, while also promoting replay-
related reactivation (McNamara et al., 2014), while inactivation of VTA causes an
increase in spatial working memory-related errors in novel, but not familiar,
environments (Martig et al., 2009). The results presented here support the hypothesis
that VTA is critically involved in learning in new environments, as its inactivation
prevents the selective recruitment of replay-associated planning or memory
consolidation mechanisms to high value locations.
VTA is not the sole source of dopamine release in hippocampus, with recent
work demonstrating locus coeruleus (LC) axons likely provide the bulk of dopamine to
dorsal CA1 and can be necessary for novelty-related spatial learning (Kempadoo et al.,
2016; Takeuchi et al., 2016). LC axons in CA1 are active in locations immediately
preceding a new reward location in a familiar environment but not in a novel one,
despite similar behavior in both cases indicating mice had learned the reward locations
(Kaufman et al., 2020). This result, coupled with findings that LC neurons are modulated
by reward-predicting stimuli similarly to substantia nigra dopamine neurons (Bouret et
al., 2012; Bouret and Richmond, 2015), suggests LC activity can convey reward-related
information and thereby compensate for VTA inactivation, but only in familiar
environments where it is not signaling more general novelty. Altogether, our work adds
to the body of evidence that VTA can directly or indirectly mediate hippocampal
plasticity and spatial learning and memory (Gasbarri et al., 1996; McNamara et al.,
2014; Rosen et al., 2015; Rossato et al., 2009), and suggests an intriguing distinction

between the function of VTA and LC dopamine release in hippocampus (Duszkiewicz et
al., 2019).
These results also support the hypothesis that reverse replay is intimately
involved in reward learning (Ambrose et al., 2016; Foster and Wilson, 2006; Mattar and
Daw, 2018). By activating representations associated with the location of the current
reward and then progressing sequentially to earlier positions that preceded reward,
reverse replay may provide a neural eligibility trace by which spatial positions can be
associated with their proximity to reward (Foster and Wilson, 2006; Sutton and Barto,
1998). Dopamine release at reward detection and consumption would then couple a
temporal gradient of dopamine concentration with the temporally-extended, reverse
sequential activation of states that led to that reward. Indeed, VTA dopamine neurons
are activated when SWR and replay occur during a spatial working memory task, but
not during subsequent sleep (Gomperts et al., 2015), indicating close coordination
specifically during reward learning. CA1, VTA, and medial prefrontal cortex neurons are
jointly coupled via oscillatory mechanisms during spatial working memory (Fujisawa and
Buzsáki, 2011), suggesting downstream targets of both replay (Berners-Lee et al., 2021)
and VTA dopamine neurons (Lammel et al., 2008) may receive temporally-precise
conjunctive input from them. We expect future work aimed at untangling under what
conditions VTA and replay influence each other and coordinate to provide downstream
areas with sequential activity in the presence of dopamine to be particularly fruitful in
understanding how reward drives spatial learning.
583
584 **Author Contributions**
585 M.R.K. and D.J.F. conceived of and designed the study. M.R.K. acquired the data and
586 performed the analyses. M.R.K. and D.J.F. wrote the manuscript.
587

Acknowledgements
We thank Stanford Gene Vector and Virus Core and Karl Deisseroth for viral constructs
and the Biological Imaging Facility at University of California, Berkeley for assistance
with tissue imaging. This work was supported by NIH grant NS113557. Animal use
conformed to NIH guidelines and was approved by the UC Berkeley Animal Care and
Use Committee.
594

595 **Declaration of Interests**
596 The authors declare no competing interests.
597
598
599
600
601 **Methods**
602
603 **Lead contact**
604   Further information and requests for resources and reagents should be directed
605 to and will be fulfilled by the Lead Contact, David Foster (davidfoster@berkeley.edu).
606
607 **Materials availability**

608   This study did not generate any unique reagents.
609
610 **Data and code availability**
611   All data and code are available from the Lead Contact upon request. Custom
612 analysis code is publicly available at the DOI listed in the key resource table. Any
613 additional information required to reanalyze the data reported in this work is available
614 from the Lead Contact upon request.
615
Experimental model and subject details
All experimental procedures were performed in accordance with the University of
California Berkeley Animal Care and Use Committee and US National Institutes of
Health guidelines. A total of twelve adult male Sprague Dawley TH-cre knock-in rats
(inotiv, HsdSage:SD-TH$^{em1(IRES-Cre)Sage}$, age 3-10 months, 300-750 g) were used in this
experiment, of which seven contributed data to the present report. Two were excluded
from further analysis due to lack of virus expression evident in post-mortem
immunohistochemistry, two were excluded due to faulty recording hardware, and one
was excluded due to non-performance of behavioral tasks. Animals were housed on a
standard, non-inverted 12-h light cycle. Rats were pair-housed with littermates prior to
the start of experiments, after which they were single-housed.
627
Behavioral pre-training
Adult male Sprague Dawley TH-cre knock-in rats were fed ad lib and handled
daily prior to experimental training. They were then food restricted to 85-90% of
baseline weight and trained to collect liquid chocolate reward (0.1 ml) from each end of
a single linear track (200 cm length) for at least 15 sessions. 3-6 other linear tracks
were present in the room during this pre-training, with positions constant for the duration
of experiments with each animal.
635
Surgical procedures
Each rat underwent virus injection and drive implantation in one surgery (control
rats 1 and 2) or two surgeries spaced 4-20 days apart (experimental rats 1-4, control rat
3).
Virus injection. All virus was obtained from the Stanford Gene Vector and Virus
Core under material transfer agreement with the laboratory of Karl Deisseroth.
Experimental rats were injected with AAV-DJ-EF1a-DIO-hM4D(Gi)-mCherry (GVVC-
AAV-129) and control rats were injected with AAV-DJ-EF1a-DIO-mCherry (GVVC-AAV-
14), with 1 µl of virus delivered stereotactically to VTA in each hemisphere (-5.6 mm
posterior, ± 0.7 mm lateral, and -8 mm ventral, all from bregma and skull surface). Data
collection began four weeks after virus injection to allow for expression.
Recording drive implantation. Each rat was implanted with a recording
microdrive, targeting bilateral (32 tetrodes, ~40 g, n = 4 rats) or unilateral (20 tetrodes,
~35 g, n = 2 rats; 6 tetrodes, ~20 g, n = 1 rat) dorsal CA1. Each tetrode bundle of four
platinum iridium wires (Neuralynx) was independently adjustable and electroplated with
gold to an impedance of 150-300 kΩ. Tetrodes were advanced over the course of 1-3
weeks to the pyramidal cell layer. Rats were reintroduced to the pre-training linear track
after several days of post-surgical recovery.

654
Tissue processing and immunohistochemistry
Eight weeks after virus injection, rats were deeply anesthetized with isoflurane
and transcardially perfused with phosphate-buffered saline (PBS) and then 4%
paraformaldehyde (PFA) in PBS. Brains were stored in 4% PFA for >24 hours, then
30% sucrose dissolved in PBS for >7 days for cryoprotection. 20-40 µm sections were
made in a cryostat and mounted on slides. For tyrosine hydroxylase (TH) staining, all
steps were performed at room temperature in a dark container on a slow orbital shaker.
Sections were rinsed three times for 10 minutes each in PBS, then incubated for 2
hours in blocker (3% normal donkey serum and 0.3% Triton-X in PBS). Sections were
then kept for 16-20 hours in blocking buffer with primary antibody (1:200, rabbit α-TH,
EMD Millipore 657012, or sheep α-TH, Abcam ab113). After three 10-minute washes in
PBS, sections were incubated with secondary antibody in blocking buffer for 2 hours
(1:200, Alexa Fluor 488-conjugated α-rabbit, Invitrogen ThermoFisher R37118, or Alexa
Fluor 488-conjugated α-sheep, Abcam ab150177). Imaging was performed at the
Biological Imaging Facility at the University of California Berkeley using a Zeiss
AxioImager M2.
671
Drug delivery
At least 10 minutes prior to beginning a recording session (individual animal
means of 12 to 17.4 minutes, except for experimental rat 1, with an average of 4
minutes before recording session), rats were injected intraperitoneally (IP) with saline or
1-4 mg/kg clozapine N-oxide (CNO) solution (2-4 mg/ml in diH$_2$O with 50-100 µl
dimethyl sulfoxide). In Experiment 1 (described below), 2 mg/kg CNO was delivered in
all control animal CNO sessions and in most CNO sessions (22/30 sessions), including
all novel track CNO sessions, in 3 out of 4 experimental animals. Experimental rat 3
received 1 mg/kg CNO in all sessions, motivated by our desire to have enough low
dosage data in one animal in case 2 mg/kg CNO had an effect in control animals
(Gomez et al., 2017). However, this was not the case, and we saw no difference
between CNO doses, consistent with some previous reports in the literature (Mahler et
al., 2014; Vazey and Aston-Jones, 2014), and therefore combined all data. In
Experiment 2, animals received 1mg/kg (n=6 sessions), 2 mg/kg (n=14 sessions), or 4
mg/kg (n=5 sessions), with the higher dose included to probe whether it could reveal a
role for VTA in the absence of novelty. As in Experiment 1, we saw little difference
between doses and combined all data. 1-4 sessions (mean±s.d., 1.5±0.65) were
completed each day, with at least 1.5 hours between injections. Altogether, we were
guided in the injection timing and drug doses by previous work in the field (reviewed in
Smith et al., 2016), with drug delivery 0-30 minutes before behavioral testing (Kane et
al., 2017), significant neuronal suppression occurring within 5-10 minutes and returning
to baseline 70 minutes post-injection (Chang et al., 2015; all recording sessions were
completed by 60 minutes post injection), and CNO doses between 1 and 10 mg/kg
(Smith et al., 2016).
To prevent the possibility of carry over effects of CNO, saline sessions never
followed CNO sessions in the same day (except for 3 recording days in experimental rat
3, when CNO preceded saline sessions by > 4 hours). Besides this restriction, any other
ordering of saline and CNO sessions was permitted. In days with two sessions, for

example, saline -> saline, saline -> CNO, and CNO -> CNO all occurred. Nevertheless,
we attempted to keep the drug X session number of the day close to balanced. In
particular, in Experiment 1 novel sessions (described below), each animal had equal
numbers of saline and CNO 1st and 2nd sessions of the day.
704
Task design
In Experiment 1, animals progressed through three epochs. In Epoch 1, animals
collected 0.1 ml rewards from each end for 10-20 laps. Then, unsignaled to the rat, the
session entered Epoch 2, where reward at one end (Incr. end) was increased to 0.4 ml
while the other (Unch. end) remained at 0.1 ml. The assignment of track ends to be Incr.
and Unch. randomly varied session to session. After 10-20 laps in Epoch 2, the reward
changed again unsignaled to the rat, with both reward ends again delivering 0.1 ml in
Epoch 3. Rats completed up to 20 laps in Epoch 3, before being removed and placed
back into a rest box. This same task was repeated on distinct linear tracks that varied
based on position in the room, material of construction, color, length, orientation, and
reward well size and position. Sessions were classified as either "novel" (the 1st or 2nd
experience on a particular linear track) or "familiar" (3rd or later experience on a specific
track). The track used for pre-training was used first for both saline and CNO sessions.
Then, each novel track was used for 2-6 sessions, with all sessions consisting of only
saline or CNO (excluding one track each in experimental rats 2 and 3 that had both
saline and CNO sessions). The assignment of saline or CNO to each novel track was
varied across rats.
In Experiment 2, reward at the stable end (0.2 ml) remained fixed throughout the
session, while at the volatile end it varied pseudorandomly every lap between 0 and 0.8
ml (mean 0.37 ml; blocks of 20 laps were comprised of 3 laps x 0 ml, 4 laps x 0.1 ml, 3
laps x 0.2 ml, 4 laps x 0.4 ml, and 6 laps x 0.8 ml). Rats were allowed to continue
running until sated. Which track end was assigned to be stable and volatile varied
randomly session by session. Each rat performed saline and CNO sessions of this task
on the same linear track. The linear track was initially novel (except for in experimental
rat 3). However, 1-2 saline sessions preceded the first CNO session, rendering it
familiar for almost all CNO sessions and most saline sessions. In each rat, all
Experiment 2 sessions were completed after all Experiment 1 sessions.
732
Data acquisition
Rat position was monitored at 30 frames/s using overhead camera and LEDs on
the recording drive, then tracked using automated software (Spike Gadgets). Two-
dimensional position and velocity were smoothed using a 7-bin median average,
followed by a 5 bin Gaussian filter. Linearized position was then used for further
analysis. Neural data was collected using a 128-channel wireless HH128 headstage
(Spike Gadgets). LFP was sampled at 30 kHz and spikes extracted based on threshold
crossing of 40-60 µv (Trodes software, Spike Gadgets). Individual units were
differentiated based on manual clustering of spike waveform peak amplitudes using
custom software (xclust2, M. A. Wilson, MIT).
743
744 **Behavioral analysis**

745   Reward end visits were defined as periods when the rat was within 10 cm of the
746   end of the track (approximately ~3-5 cm from the reward well, depending on the track).
747   When analyzing visit durations, we excluded a small number of outliers (< 15 total
748   across all sessions) that were longer than 60 s.
749
LFP analysis
For each sesson, 2-5 tetrodes with visible sharp wave ripples (SWR) were
selected for SWR analysis. LFP from one channel from each tetrode was band-pass
filtered between 150 and 250 Hz, then the smoothed (Gaussian kernel, 12 ms s.d.),
absolute value of the Hilbert transform was averaged across tetrodes. For detecting
SWR, we examined periods when the rat position was within 10 cm of the reward wells
with velocity ≤ 8 cm/s. SWR were classified as local peaks when the average ripple
power exceeded 4 s.d. above the mean, with start and end points defined as the time
ripple power reached the mean before and after the peak, with a minimum start to end
duration of 150 ms and maximum of 1 s. SWR rate for each reward end visit was then
calculated as the number of SWR detected divided by the total duration of rat velocity ≤
8 cm/s during that end visit. During Experiment 1, all analyses considered SWR rate
only during the first 10 s of each end visit to isolate the reward consumption-related
period and exclude occasional longer task-disengaged resting periods. Results shown
in Figure 2 were similar if we extended this to examine up to the first 20 s of each
reward visit. In Experiment 2, we included the first 20 s of each end visit to allow for the
longer consumption time required for 0.8 ml.
In Experiment 2, we defined mean-subtracted SWR rate:
768   Mean-subtracted rate (x,y) = rate (x,y) – rate (x)
769   Where x and y are current and previous volatile volumes, respectively.
770
Single unit analysis
Place fields were calculated for each neuron based on spiking activity while the
rat velocity exceeded 8 cm/s. Position was binned into 2 cm bins and directional place
fields were calculated as the histogram of spike counts in each position bin (smoothed
with Gaussian kernel, 4 cm s.d.) normalized by the animal's occupancy in each bin,
separately for periods when the rat was moving in each direction on the linear track
(e.g., left and right). We calculated several properties of place fields to determine
whether novelty or VTA inactivation affected them. The map direction correlation was
defined as the Pearson correlation between the place field calculated in each running
direction, such that a value of 1 indicates perfectly reliable firing dependent only on
position, not running direction. The lap to session correlation was defined for each
neuron only for the running direction with higher max firing rate. For that running
direction, a place field was calculated independently for each lap, smoothed (Gaussian
kernel, 4 cm s.d.), correlated to the directional field calculated from the entire session,
and the average correlation coefficient taken across all laps.
786
Replay analysis
Because decoding quality relies on sufficient spiking and SWR and population
spike density events do not always co-occur, our replay analysis used population
spiking to determine candidate events. Candidate replay events were determined based

on population activity while the rat was not running (velocity ≤ 8 cm/s) and near the
reward wells (10 cm away at most). Population spike density was binned into 1 ms bins
and smoothed (Gaussian kernel, 20 ms s.d.) and candidate events defined as local
peaks when the population rate exceeded the mean by 3 s.d. and that lasted at least 50
ms, with start and end defined as the nearest times the rate crossed the mean before
and after the peak. A memory-less Bayesian decoding algorithm was used to classify
both position and running direction during candidate events, as in previous work 13.
Replay position in each running direction was estimated in time windows of 40 ms,
beginning at the start of the event, and advancing in 5 ms steps. The start/end of
putative trajectories within a candidate event were determined by removing bins at
either end of the event that contained zero spikes or had a position difference > 50% of
the track length from the next/previous window. Candidate events with a remaining
length of at least 5 time bins, an absolute weighted correlation (Wu and Foster, 2014)
exceeding 0.5, and at least 55% of the posterior probability in one of the running
directions (Ambrose et al., 2016) were classified as replay. Replays were classified as
forward or reverse by comparing the direction of replay movement across the track with
the running direction map containing the majority of the posterior probability: if they
matched (e.g., the replay moved upward and used upward fields), the replay was
classified as forward, and otherwise it was classified as reverse.
Only sessions with sufficiently accurate behavioral position decoding accuracy
during run were included. Bayesian decoding using the directional place fields was
applied to 250 ms non-overlapping windows covering the entirety of each session. For
all time bins with mean animal velocity >20 cm/s, position >20 cm from reward wells,
and >5 total spikes from any neurons, actual and decoded position and running
direction were compared, yielding a position decoding error (distance in cm) and
direction decoding match (same or different). Sessions with mean decoding error >35
cm or direction match <60% were excluded, which totaled 4/64 sessions in experimental
rats and 12/72 sessions in control rats.
819
820 **Statistics**
821   A mixed effects Poisson generalized linear model was used to test which
822 experimental variables affected SWR rate using the Matlab *fitglme* function
823 (Mathworks), similarly to previous work 13. Animal ID was modeled as a random effect,
824 allowing baseline SWR rate to vary across rats. The full model was as follows:
825
826 SWR rate = exp[b0 + b1 × (Incr. reward end) + b2 × (Epoch 2) + b3 × (CNO) + b4 ×
827 (Incr. reward end) × (Epoch 2) + b5 ×(Incr. reward end) × (CNO) + b6 × (Epoch 2) ×
828 (CNO) + b7 × (Incr. reward end) × (Epoch 2) × (CNO)]
829
Where "Incr. reward end" is a dummy variable indicating the rat is at the Incr. reward
end, "Epoch 2" is a dummy variable indicating the visit is occurring in Epoch 2, and
"CNO" is a dummy variable indicating it is a session with CNO injected. The coefficient
for each term corresponds to the log multiplicative change in SWR rate from the
reference condition (animal-specific rate at Unch. end, not in Epoch 2, of a saline
session). The offset term log(duration) for each stopping period accounted for the total
time the rat was stationary at each reward well visit, so the model fit SWR rate, rather

than SWR count. Experimental and control rats were fit separately with this model, as
were familiar and novel sessions.
Bootstrapping was used to assess the effect of drug on SWR rate in each
experimental condition. For each combination of novelty, reward end, and epoch, drug
identity was shuffled 5,000 times, generating a distribution of the chance difference
between the SWR rate in saline vs. CNO sessions. P-values were determined using
one-tailed tests, under the hypothesis that CNO would cause lower SWR rates at the
Incr. end and higher SWR rates at the Unch. end when compared to saline.
In analyses used to assess the effect of various experimental variables on
behavioral and neural measurements, the following variables were consistently defined:
"animal group" was a dummy variable indicating experimental rats, "drug" was a dummy
variable indicating CNO, "novelty" was a dummy variable indicating novel session,
"epoch" was a categorical variable indicating epoch number, "reward end" was a
dummy variable indicating Incr. reward end, "RPE sign" was a dummy variable
indicating a positive RPE, and "previous volume" was a categorical variable indicating
the volatile volume at the previous visit.

Supplemental Figure Legends

Figure 1 – Supplement 1. Additional histology examples.
 (A) Additional example section from experimental rat with evident virus expression. Scale bar 300 μm. Dashed line marks approximate location of midline. mCherry puncta (top panel) and TH-positive dopaminergic neurons (bottom panel) indicated virus expression in the VTA. **(B)** Additional example section from control rat with evident virus expression. Scale bar 150 μm. **(C)** Example sections from 1st rat excluded due to lack of virus expression. Scale bar 600 μm. Complete lack of mCherry puncta (top panel) indicated failure of virus expression. **(D)** Example sections from 2nd rat excluded due to lack of virus expression. Scale bar 300 μm.

Figure 1 – Supplement 2. Behavioral effects of novelty and VTA inactivation.
(A) In novel sessions, Unch. visit duration decreased from Epoch 1 to Epoch 2, while CNO additionally led to longer visit duration in experimental rats. Mean ± standard error, Exp Saline, Epoch 1: 7.551±1.11, Epoch 2: 4.98±0.44; Exp CNO, Epoch 1: 12.14±1.7, Epoch 2: 6.75±0.64. Con Saline, Epoch 1: 7.76±0.65, Epoch 2: 4.86±0.35; Con CNO, Epoch 1: 7.46±0.59, Epoch 2: 4.96±0.25. Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual animal as a random effect: epoch, $z=-2.99$, $p<0.01$; group X drug, $z=3.29$, $p<0.01$; all other terms, n.s. **(B)** In novel sessions, Incr. visit duration increased from Epoch 1 to Epoch 2, while CNO additionally led to longer visit duration in experimental rats. Mean ± standard error, Exp Saline, Epoch 1: 7.52±1.32, Epoch 2: 10.05±0.88; Exp CNO, Epoch 1: 11.55±2, Epoch 2: 13.39±1.18. Con Saline, Epoch 1: 7.77±0.58, Epoch 2: 10.97±0.55; Con CNO, Epoch 1: 6.77±0.38, Epoch 2: 10.46±0.22. Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual animal as a random effect: epoch, $z=2.82$, $p<0.01$; group X drug, $z=2.91$, $p<0.01$; all other terms, n.s. **(C)** In familiar sessions, Unch. visit duration decreased from Epoch 1 to Epoch 2, with only a modest effect of CNO compared to novel sessions. Mean ± standard error, Exp Saline, Epoch 1: 6.45±0.72, Epoch 2: 5.17±0.66; Exp CNO, Epoch 1: 6.47±0.42, Epoch 2: 5.841±0.51. Con Saline, Epoch 1: 6.54±0.5, Epoch 2: 4.72±0.37; Con CNO, Epoch 1: 6.18±0. 3, Epoch 2: 4.17±0.17. Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual animal as a random effect: epoch, $z=-2.37$, $p<0.05$; all other terms, n.s. **(D)** In familiar sessions, Incr. visit duration increased from Epoch 1 to Epoch 2. Mean ± standard error, Exp Saline, Epoch 1: 6.41±0.59, Epoch 2: 11.1±0.97; Exp CNO, Epoch 1: 6.03±0.18, Epoch 2: 11.99±1. Con Saline, Epoch 1: 7.27±0.66, Epoch 2: 11.46±0.71; Con CNO, Epoch 1: 6.33±0.48, Epoch 2: 10.28±0.4. Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual animal as a random effect: epoch, $z=3.97$, $p<0.001$; all other terms, n.s. **(E)** Unch. visit duration increased from Epoch 2 to Epoch 3. Mean ± standard error, Exp Saline, Epoch 2: 5.13±0.53, Epoch 3: 7.95±0.6; Exp CNO, Epoch 2: 6.04±0.42, Epoch 3: 7.89±0.68. Con Saline, Epoch 2: 4.77±0.27, Epoch 3: 6.23±0.37; Con CNO, Epoch 2: 4.43±0.15, Epoch 3: 5.96±0.19. Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual animal as a random effect: epoch, $z=2.32$, $p<0.05$; all other terms, n.s. **(F)** Incr. visit duration decreased from Epoch 2 to Epoch 3. Mean ± standard error, Exp Saline, Epoch 2: 10.87±0.8, Epoch 3: 9.08±1.01; Exp CNO, Epoch 2: 12.31±0.82, Epoch 3: 9.92±0.95. Con Saline, Epoch 2:

11.29±0.5, Epoch 3: 6.48±0.28; Con CNO, Epoch 2: 10.34±0.27, Epoch 3: 6.42±0.29.
Mixed-effects GLM with epoch, drug, animal group, and all interactions, with individual
animal as a random effect: epoch, z=-4.77, $p<10^{-5}$, group X epoch, z=2.31, p<0.05; all
other terms, n.s.

Figure 1 – Supplement 3. Effect of reward change on running velocity.
Running speed towards the Incr. end in Epoch 2 was consistently significantly faster
than towards the Unch. end, across all conditions. The median running speed in all non-
zero velocity timepoints while the animal was located outside of the reward end zones in
each epoch and running direction was calculated for each session. Mean and standard
error across sessions are shown here. Mixed-effects model predicting the velocity
difference in Epoch 2, Incr. – Unch., as a function of drug, novelty, and their interaction,
with animal-specific intercepts. Experimental group : intercept, z=2.99, p<0.01; drug,
z=3.18 , p<0.01; all other terms, n.s. Control group: intercept, z=7.85, $p<10^{-10}$; all other
terms, n.s. Filled symbol, saline; unfilled symbol, CNO.

Figure 2 – Supplement 1. Modulation of SWR rate by reward increase.
(A) In experimental rats, a mixed effects Poisson GLM was fit to the data and 5,000
drug identity shuffles. The difference between model-predicted SWR rate in saline and
CNO sessions at each reward end (Unch. top row, Incr. bottom row) and novelty
condition (familiar left column, novel right column), in data (red lines) and in bootstrap
shuffles (histogram). Significance values reflect one-tailed hypothesis test, with
hypotheses that Unch. saline < Unch. CNO and Incr. saline > Incr. CNO. **(B)** A mixed
effects GLM with bootstrap, as in (A), but for control animals.

Figure 2 – Supplement 2. SWR rate in Epoch 3.
(A) SWR rate as a function of time in stopping period in Epoch 2 and 3 for four example
sessions in experimental rats, as in Figure 2a. Epoch 2 (red lines), Epoch 3 (dashed
gray lines). SWR rate was binned in 0.25 s windows and smoothed with a 2 bin
Gaussian. Line, mean; shading, standard error. **(B)** Same as Figure 2F, but for Epoch 3.
(C) Same as Figure 2G, but for Epoch 3.

Figure 2 – Supplement 3. Similar results independent of session number of the
day. The dataset for Experiment 1 was split into two, with one part including only
sessions that occurred 1st in any given day ("1st of day") and the other including all
sessions that were not the 1st in any given day ("2nd+ of day"). Although the low
resultant session counts in each group precluded statistical analysis, the main effect of
CNO in reducing SWR rate difference between the reward ends was very similar.

Figure 2 – Supplement 4. Ripple duration is increased with familiarity. The duration
of SWR was examined similarly to the analysis on SWR rate. A mixed-effects Poisson
generalized linear model (GLM) was fit to predict changes in SWR duration across
reward end, epoch, drug condition, and novelty, with animal identity as a random effect.
Significant coefficients: CNO (z=-5.86, $p<10^{-5}$) and Epoch 2 (z=-2.1, p<0.05).

Figure 3 – Supplement 1. SWR rate at stable end in experimental rats.

(A) At stable end visits in saline sessions, SWR rate was not significantly modulated by
the previous volatile end visit reward volume. Pearson correlation between SWR rate
and previous volatile volume, r=-0.0643, p=0.21. Two sample t-test between volatile
volume ≤ 2 and volatile volume > 2, t(380)=1.465, p=0.144. **(B)** At stable end visits in
CNO sessions, SWR rate was not significantly modulated by the previous volatile end
visit reward volume. Pearson correlation between SWR rate and previous volatile
volume, r=-0.0645, p=0.205. Two sample t-test between volatile volume ≤ 2 and volatile
volume > 2, t(386)=1.137, p = 0.256. Two-way ANOVA with drug and previous volatile
volume ≤ 2: drug (F[1,766]=6.43, p<0.05), volume ≤ 2 (F[1,766]=3.36, p=0.067), drug X
volume (F[1,766]=0.03, p=0.853). Error bars, standard error.
985
Figure 3 – Supplement 2. SWR rate in all sessions in volatile reward task.
(A) SWR rate as a function of reward volume and time in end visit, as in Figure 3B, for
all sessions combined (including saline and CNO sessions in experimental and control
rats). Left panel, stable reward end. Right panel, volatile reward end. In stable panel,
traces are colored based on previous volatile end visit volume. In volatile panel, traces
are colored based on current volatile volume. **(B)** SWR rate at volatile end as a function
of current and previous volatile volume, as in Figure 3D, for all volatile reward task
sessions. **(C)** SWR rate for each non-zero volatile volume plotted as a function of
previous volume, with the mean SWR rate for that current volume subtracted. Unfilled
symbols, mean of previous volume across all current volumes. Thick dashed line, linear
fit to mean values. Pearson correlation between (ripple rate – mean) and previous
volume, r=-0.07, p<0.01, consistent with RPE coding. Error bars, standard error. **(D)**
Positive RPE caused significantly greater ripple rate than negative RPE (two-sample t-
test, t[1661]=2.741, p<0.01). **(E)** SWR rate at the stable end was significantly negatively
correlated with the most recent volatile volume (r=-0.06, p<0.01). **(F)** SWR rate at the
stable end was significantly greater when the most recent volatile end volume was less
than or equal in volume (≤ 2) than when it was greater (two-sample t-test,
t[2485]=2.582, p<0.01). **(G)** SWR rate at the volatile end was significantly higher if
recent reward history was lower than the average. Reward volume at the 3 previous
visits was averaged, then split above and below the median. Poisson GLM with two
terms, current volume and reward history (above/below median): current volume,
z=22.21, p<10^{-10}; history, z=-2.03, p<0.05).
1008
Figure 4 – Supplement 1. Effect of novelty and VTA inactivation on place cell
properties.
(A) Correlation between single lap place fields and session averaged field. Three-way
ANOVA with drug, novelty, and animal group: novelty (F[1,3249]=6.75, p<0.01), novelty
X group (F[1,3249]=15.76, p<0.01), all others, p>0.2. **(B)** Correlation between
unidirectional fields calculated separately in each running direction. Three-way ANOVA
with drug, novelty, and animal group: drug (F[1,2816]=5.76, p<0.05), novelty
(F[1,2816]=28.21, p<10^{-10}), drug X novelty (F[1,2816]=5.52, p<0.05), novelty X group
(F[1,2816]=6.56, p<0.05), all others, p>0.17.
1018
1019 **Figure 4 – Supplement 2. Run decoding accuracy in replay analysis sessions.**

(A) Mean decoding error during run. Position and running direction were decoded during periods of strong locomotion (animal velocity >20 cm/s and position >20 cm from the reward wells) in 250 ms bins. Sessions with >35 cm mean decoding error were excluded from analysis. A mixed-effects model predicting position decoding error as a function of drug, novelty, recorded neuron count, and mean place field size, found more recorded cells and smaller mean field size in both groups led to smaller decoding errors. Experimental rats: drug, z=-2.48, p<0.05; cell count, z=-3.48, p<0.01; mean field size, z=5.36, p<10^{-5}; other terms, n.s. Control rats: cell count, z=-4.07, p<0.01; mean field size, z=4.04, p<0.01; other terms, n.s. Filled and unfilled symbols are saline and CNO sessions, respectively. Error bars, standard error. **(B)** Mean fraction of bins where actual and decoded running direction were the same. Sessions with <60% match were excluded from analysis. Symbols as in (A). A mixed-effects model predicting probability of matching real and decoded run direction as a function of drug, novelty, recorded neuron count, and mean place field size, found more recorded cells in both groups led to higher match probability. Experimental rats: drug, z=2.33, p<0.05; novelty, z=-2.96, p<0.01; cell count, z=6.81, p<10^{-5}; other terms, n.s. Control rats: cell count, z=6.23, p<10^{-5}; other terms, n.s.

Figure 4 – Supplement 3. Non-local replay was unaffected by experimental manipulations.
(A) The difference in rate of reverse replay at each end (Incr. – Unch.) in novel sessions in experimental rats. Error bars, standard error of the mean. Reward condition is indicated by color (equal reward, epoch 1 and 3, gray; unequal reward, epoch 2, orange), and drug condition is indicated on the x-axis. The difference in replay rate between equal and unequal reward conditions was assessed with a mixed-effects linear model with drug, novelty, and replay directionality, and animal-specific intercepts as random effect: all terms, n.s. **(B)** Same as (A), but for familiar sessions. **(C)** Same as (A), but for forward replay. Unequal reward, epoch 2, purple. **(D)** Same as (C), but for familiar sessions.

1066 **Supplemental Figures**
1067

1068
1069 **Figure 1 – Supplement 1**
1070

1071
1072　**Figure 1 – Supplement 2**

1073
1074 **Figure 1 – Supplement 3**

Figure 2 – Supplement 1

Figure 2 – Supplement 2

1161
1162 **Figure 2 – Supplement 3**

1195
1196   **Figure 2 – Supplement 4**

1211
1212 **Figure 3 – Supplement 1**

1239
1240 **Figure 3 – Supplement 2**

Figure 4 – Supplement 1

1287
1288 **Figure 4 – Supplement 2**

1321
1322 **Figure 4 – Supplement 3**

References

Ambrose RE, Pfeiffer BE, Foster DJ. 2016. Reverse Replay of Hippocampal Place Cells Is Uniquely Modulated by Changing Reward. *Neuron* **91**:1124–1136. doi:10.1016/j.neuron.2016.07.047

Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL. 2007. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proceedings of the National Academy of Sciences* **104**:5163–5168. doi:10.1073/pnas.0700293104

Berners-Lee A, Feng T, Silva D, Wu X, Ambrose ER, Pfeiffer BE, Foster DJ. 2022. Hippocampal replays appear after a single experience and incorporate greater detail with more experience. *Neuron* **110**:1829-1842.e5. doi:10.1016/j.neuron.2022.03.010

Berners-Lee A, Wu X, Foster DJ. 2021. Prefrontal Cortical Neurons Are Selective for Non-Local Hippocampal Representations during Replay and Behavior. *The Journal of Neuroscience* **41**:5894–5908. doi:10.1523/JNEUROSCI.1158-20.2021

Bouret S, Ravel S, Richmond BJ. 2012. Complementary neural correlates of motivation in dopaminergic and noradrenergic neurons of monkeys. *Front Behav Neurosci* **6**:40. doi:10.3389/fnbeh.2012.00040

Bouret S, Richmond BJ. 2015. Sensitivity of Locus Ceruleus Neurons to Reward Value for Goal-Directed Actions. *The Journal of Neuroscience* **35**:4005–4014. doi:10.1523/JNEUROSCI.4553-14.2015

Buzsáki G. 2015. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus* **25**:1073–1188. doi:10.1002/hipo.22488

Chang, C. Y., Esber, G. R., Marrero-Garcia, Y., Yau, H.-J., Bonci, A., & Schoenbaum, G. (2015). Brief optogenetic inhibition of dopamine neurons mimics endogenous negative reward prediction errors. Nature Neuroscience, 19(1), 1–8. https://doi.org/10.1038/nn.4191

Diba K, Buzsáki G. 2007. Forward and reverse hippocampal place-cell sequences during ripples. *Nat Neurosci* **10**:1241–2. doi:10.1038/nn1961

Duszkiewicz AJ, McNamara CG, Takeuchi T, Genzel L. 2019. Novelty and Dopaminergic Modulation of Memory Persistence: A Tale of Two Systems. *Trends Neurosci*. doi:10.1016/j.tins.2018.10.002

Fernández-Ruiz A, Oliva A, Fermino de Oliveira E, Rocha-Almeida F, Tingley D, Buzsáki G. 2019. Long-duration hippocampal sharp wave ripples improve memory. *Science (1979)* **364**:1082–1086. doi:10.1126/science.aax0758

Fields HL, Hjelmstad GO, Margolis EB, Nicola SM. 2007. Ventral tegmental area neurons in learned appetitive behavior and positive reinforcement. *Annu Rev Neurosci* **30**:289–316. doi:10.1146/annurev.neuro.30.051606.094341

Foster DJ. 2017. Replay Comes of Age. *Annu Rev Neurosci* **40**:581–602. doi:10.1146/annurev-neuro-072116-031538

Foster DJ, Wilson M a. 2006. Reverse replay of behavioural sequences in hippocampal place cells during the awake state. *Nature* **440**:680–3. doi:10.1038/nature04587

Fujisawa S, Buzsáki G. 2011. A 4 Hz oscillation adaptively synchronizes prefrontal, VTA, and hippocampal activities. *Neuron* **72**:153–65. doi:10.1016/j.neuron.2011.08.018

Gasbarri A, Sulli A, Innocenzi R, Pacitti C, Brioni JD. 1996. Spatial memory impairment
induced by lesion of the mesohippocampal dopaminergic system in the rat.
Neuroscience **74**:1037–1044. doi:10.1016/S0306-4522(96)00202-3

Gomez, J. L., Bonaventura, J., Lesniak, W., Mathews, W. B., Sysa-Shah, P., Rodriguez,
L. A., Ellis, R. J., Richie, C. T., Harvey, B. K., Dannals, R. F., Pomper, M. G., Bonci,
A., & Michaelides, M. (2017). Chemogenetics revealed: DREADD occupancy and
activation via converted clozapine. Science, 357(6350), 503–507.
https://doi.org/10.1126/science.aan2475

Gomperts SN, Kloosterman F, Wilson M a. 2015. VTA neurons coordinate with the
hippocampal reactivation of spatial experience. *Elife* **4**:1–22.
doi:10.7554/eLife.05360

Guru A, Seo C, Post RJ, Kullakanda DS, Schaffer JA, Warden MR. 2020. Ramping
activity in midbrain dopamine neurons signifies the use of a cognitive map. *BioRxiv*.
doi:doi.org/10.1101/2020.05.21.108886

Howe MW, Tierney PL, Sandberg SG, Phillips PEM, Graybiel AM. 2013. Prolonged
dopamine signalling in striatum signals proximity and value of distant rewards.
Nature **500**:575–579. doi:10.1038/nature12475

Jadhav SP, Kemere C, German PW, Frank LM. 2012. Awake hippocampal sharp-wave
ripples support spatial memory. *Science (1979)* **336**:1454–8.
doi:10.1126/science.1217230

Kane, G. A., Vazey, E. M., Wilson, R. C., Shenhav, A., Daw, N. D., Aston-Jones, G., &
Cohen, J. D. (2017). Increased locus coeruleus tonic activity causes
disengagement from a patch-foraging task. Cognitive, Affective and Behavioral
Neuroscience, 17(6), 1073–1083. https://doi.org/10.3758/s13415-017-0531-y

Kaufman AM, Geiller T, Losonczy A. 2020. A Role for the Locus Coeruleus in
Hippocampal CA1 Place Cell Reorganization during Spatial Reward Learning.
Neuron **105**:1018-1026.e4. doi:10.1016/j.neuron.2019.12.029

Kempadoo KA, Mosharov E V, Choi SJ, Sulzer D, Kandel ER. 2016. Dopamine release
from the locus coeruleus to the dorsal hippocampus promotes spatial learning and
memory. *Proc Natl Acad Sci U S A* **113**:201616515. doi:10.1073/pnas.1616515114

Kentros CG, Agnihotri NT, Streater S, Hawkins RD, Kandel ER. 2004. Increased
attention to spatial context increases both place field stability and spatial memory.
Neuron **42**:283–295. doi:10.1016/S0896-6273(04)00192-8

Kim HGR, Malik AN, Mikhael JG, Bech P, Tsutsui-Kimura I, Sun F, Zhang Y, Li Y,
Watabe-Uchida M, Gershman SJ, Uchida N. 2020. A Unified Framework for
Dopamine Signals across Timescales. *Cell* **183**:1600-1616.e25.
doi:10.1016/j.cell.2020.11.013

Lammel S, Hetzel A, Häckel O, Jones I, Liss B, Roeper J. 2008. Unique Properties of
Mesoprefrontal Neurons within a Dual Mesocorticolimbic Dopamine System.
Neuron **57**:760–773. doi:10.1016/j.neuron.2008.01.022

Li S, Cullen WK, Anwyl R, Rowan MJ. 2003. Dopamine-dependent facilitation of LTP
induction in hippocampal CA1 by exposure to spatial novelty. *Nat Neurosci* **6**:526–
531. doi:10.1038/nn1049

Lisman JE, Grace A a. 2005. The hippocampal-VTA loop: controlling the entry of
information into long-term memory. *Neuron* **46**:703–13.
doi:10.1016/j.neuron.2005.05.002

Mahler, S. v., Vazey, E. M., Beckley, J. T., Keistler, C. R., Mcglinchey, E. M., Kaufling, J., Wilson, S. P., Deisseroth, K., Woodward, J. J., & Aston-Jones, G. (2014). Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine seeking. Nature Neuroscience, 17(4), 577–585. https://doi.org/10.1038/nn.3664

Martig AK, Jones GL, Smith KE, Mizumori SJY. 2009. Context dependent effects of ventral tegmental area inactivation on spatial working memory. *Behavioural Brain Research* **203**:316–320. doi:10.1016/j.bbr.2009.05.008

Mattar MG, Daw ND. 2018. Prioritized memory access explains planning and hippocampal replay. *Nat Neurosci* **21**:1609–1617. doi:10.1038/s41593-018-0232-z

McNamara CG, Tejero-Cantero Á, Trouche S, Campo-Urriza N, Dupret D. 2014. Dopaminergic neurons promote hippocampal reactivation and spatial memory persistence. *Nat Neurosci* **17**:1658–1660. doi:10.1038/nn.3843

O'Keefe J, Dostrovsky J. 1971. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res* **34**:171–175. doi:10.1016/0006-8993(71)90358-1

O'Keefe J, Nadel L. 1978. The hippocampus as a cognitive map. Clarendon Press.

Pfeiffer BE, Foster DJ. 2013. Hippocampal place-cell sequences depict future paths to remembered goals. *Nature* **497**:1–8. doi:10.1038/nature12112

Rosen ZB, Cheung S, Siegelbaum SA. 2015. Midbrain dopamine neurons bidirectionally regulate CA3-CA1 synaptic drive. *Nat Neurosci* **18**:1763–1771. doi:10.1038/nn.4152

Rossato JI, Bevilaqua LRM, Izquierdo I, Medina JH, Cammarota M. 2009. Dopamine Controls Persistence of Long-Term Memory Storage. *Science (1979)* **325**:1017–1020. doi:10.1126/science.1172545

Schultz W, Dayan P, Montague PR. 1997. A neural substrate of prediction and reward. *Science (1979)* **275**:1593–1599. doi:10.1126/science.275.5306.1593

Singer AC, Frank LM. 2009. Rewarded Outcomes Enhance Reactivation of Experience in the Hippocampus. *Neuron* **64**:910–921. doi:10.1016/j.neuron.2009.11.016

Smith, K. S., Bucci, D. J., Luikart, B. W., & Mahler, S. v. (2016). Behavioral Neuroscience DREADDs: Use and Application in Behavioral Neuroscience DREADDs: Use and Application in Behavioral Neuroscience. Behavioral Neuroscience, 130(2), 137–155. https://doi.org/10.1037/bne0000135

Stachniak, T. J., Ghosh, A., & Sternson, S. M. (2014). Chemogenetic Synaptic Silencing of Neural Circuits Localizes a Hypothalamus→Midbrain Pathway for Feeding Behavior. Neuron, 82(4), 797–808. https://doi.org/10.1016/j.neuron.2014.04.008

Sutton RS, Barto AG. 1998. Reinforcement learning: an introduction. MIT Press.

Takeuchi T, Duszkiewicz AJ, Sonneborn A, Spooner P a., Yamasaki M, Watanabe M, Smith CC, Fernández G, Deisseroth K, Greene RW, Morris RGM. 2016. Locus coeruleus and dopaminergic consolidation of everyday memory. *Nature* 1–18. doi:10.1038/nature19325

Vazey, E. M., & Aston-Jones, G. (2014). Designer receptor manipulations reveal a role of the locus coeruleus noradrenergic system in isoflurane general anesthesia. Proceedings of the National Academy of Sciences, 111(10), 3859–3864. https://doi.org/10.1073/pnas.1310025111

Widloski J, Foster DJ. 2022. Flexible rerouting of hippocampal replay sequences around
changing barriers in the absence of global place field remapping. *Neuron* **110**:1547-
1558.e8. doi:10.1016/j.neuron.2022.02.002
Wu X, Foster DJ. 2014. Hippocampal replay captures the unique topological structure of
a novel environment. *Journal of Neuroscience* **34**:6459–69.
doi:10.1523/JNEUROSCI.3414-13.2014
1501