Development and Optimization of Genetic Manipulation Systems in Group I *Clostridium botulinum*

Sho Amatsu[1,2#], Kazuki Saito[1], Shigeru Miyata[3], Masahiko Zuka[2], Hirofumi Nariya[4#], and Yukako Fujinaga[1#]

[1]Department of Bacteriology, Graduate School of Medical Sciences, Kanazawa University, Ishikawa, Japan

[2]Department of Forensic Medicine and Pathology, Graduate School of Medical Sciences, Kanazawa University, Ishikawa, Japan

[3]Department of Food and Nutritional Sciences, College of Bioscience and Biotechnology, Chubu University, Aichi, Japan

[4]Laboratory of Food Microbiology, Graduate School of Human Life Sciences Food and Nutritional Sciences, Jumonji University, Saitama, Japan

For correspondence: Sho Amatsu, amatsu@med.kanazawa-u.ac.jp; Hirofumi Nariya, nariya@jumonji-u.ac.jp; Yukako Fujinaga, yukafuji@med.kanazawa-u.ac.jp

Keywords: *Clostridium botulinum*, gene knockout, allele-coupled exchange, xylose-inducible mazF, heat shock, conjugal transformation

Abstract

Clostridium botulinum causes the disease botulism, and nontoxigenic *Clostridium sporogenes* is closely related to group I *C. botulinum*. Despite its pathogenicity, *C. botulinum* remains poorly characterized. Genetic manipulation is critical for understanding bacterial physiology and disease. We compared the conjugal transformation efficiencies of seven strains, including group I *C. botulinum* (strains 62A, 7I03-H, Okra, Osaka05, 111) and *C. sporogenes* (strains JCM 1416^T, ATCC 15579), and our results showed that few or no transformants were obtained in certain strains. In the present study, we demonstrate that our optimized protocol increases the efficiency of DNA transfer from *E. coli* donor cells to recipient strains. In addition, we developed a novel conjugal suicide vector pXMTL that contains xylose-inducible *mazF* as a counter-selection marker, and can be transferred into *Clostridium* spp. by conjugation. The allele-coupled exchange (ACE) system using pXMTL provides a rapid method for precise, markerless and scarless genome editing in group I *C. botulinum* and *C. sporogenes*.

Importance

Group I *C. botulinum* and *C. sporogenes* exhibit low transformation efficiencies, and few or no transformants are yielded by some strains. In this study, we optimized the conjugation protocol to improve transformation efficiency. In addition, we developed a novel suicide vector pXMTL harboring a xylose-inducible *mazF* marker, and can be transferred into *Clostridium* spp. by conjugation. The combination of pXMTL and the optimized conjugation protocol provides a powerful tool for genetic manipulation of group I *C. botulinum* and *C. sporogens*.

Introduction

Clostridium botulinum and related species cause a severe paralytic illness, botulism. *C.*

botulinum is a gram-positive, spore-forming, obligate anaerobe, and is divided into four

genetically and physiologically distinct groups (I to IV). The organism produces

botulinum neurotoxin (BoNT) that has been traditionally classified into seven serotypes

(A to G) (1). Serotypes A, B, E and F toxins produced by group I and II *C. botulinum* are

mainly associated with human botulism, and those of serotypes C and D produced by

group III *C. botulinum* are mainly associated with animal botulism. No botulism cases

related to the serotype G toxin produced by group IV *C. botulinum* (also known as

Clostridium argentinense) have been reported (1). *Clostridium sporogenes* is closely

related to group I *C. botulinum* genetically and physiologically, whereas most strains of

them do not produce BoNT. The organism is the only gut symbiont known to produce

indolepropionic acid (IPA) (2), and obtains its energy by reductive Stickland metabolism

in the gut (3). Development and expansion of genetic manipulation in these organisms

would provide windows into mechanisms of intoxication/infection and gut microbiota-

dependent modulation of energy metabolism. However, Group I *C. botulinum* and *C.*

sporogenes exhibit low transformation efficiency (Fig. 1) and inefficient homologous

recombination (4). A lack of suitable genome editing systems limits analysis using these

strains.

Three approaches to genome editing in *C. botulinum* and related species have been

developed to overcome such limitations (5). First, the ClosTron system disrupts the target

genes of *C. botulinum* and *C. sporogenes* by group II intron insertion into the genomic

target (4). Mutants can be selected using a retrotransposition-activated marker (RAM)

integrated into the genome. Second, in a CRISPR-Cas9-mediated system, target genes

bearing a protospacer adjacent motif (PAM) are cleaved. To achieve precise gene editing,

double-strand DNA breaks (DSBs) are repaired by homology-directed recombination

(HDR) using donor templates. These markerless and scarless gene editing systems were

developed in *C. sporogenes* (6–8), group I *C. botulinum* (8) and group II *C. botulinum*

(9). Last, the two-step allele-coupled exchange (ACE) system also enables markerless

and scarless gene editing at unlimited target sites by double crossover recombination

without DSBs of the bacterial genome. In the ACE systems of *Clostridium* species (spp.),

targeted mutants have been successfully obtained using counterselection markers: *e.g.*, an

orotate phosphoribosyltransferase gene (*pyrE*) from *C. sporogenes* for *C. difficile* and

Clostridium acetobutylicum (5, 10), a cytosine deaminase gene (*codA*) from *Escherichia*

coli for *C. difficile* (11) and a *mazF* gene (mRNA interferase) from *E. coli* for *C. difficile*

(12). However, the *pyrE* and *codA* homolog genes are carried by *C. botulinum*, *C.*

sporogenes and *Clostridium perfringens* (10, 11), indicating that these genes are not

applicable for these organisms. The *mazF* gene is one of the suitable counterselection

markers, whereas a lactose-inducible promoter P_{bgaL} is slightly 'leaky' in *C.*

acetobutylicum (12). Recently, we developed a xylose-inducible *mazF*-based suicide

vector pXM (13). This plasmid can be introduced into *C. perfringens* by electroporation

transformation, in which the xylose-inducible promoter tightly regulates MazF

expression. However, DNA transfer efficiencies by electroporation or conjugation

methods vary between species and strains. Conjugative transfer of clostridial shuttle

vectors from *E. coli* CA434 donor cells to *Clostridium* spp. is highly efficient for *C.*

difficile and other clostridial recipients (5, 14). In the present study, we developed a novel

conjugal *mazF*-based suicide vector pXMTL that contains a xylose-inducible promoter.

In addition, we optimized the conjugal transformation protocol for group I *C. botulinum*

and *C. sporogenes*, resulting in higher transformation efficiencies than the original

protocol. The ACE system using pXMTL provides a rapid method for precise, markerless

and scarless genome editing in group I *C. botulinum* and *C. sporogenes*.

Materials and Methods

Bacterial strains

C. botulinum strain 62A (group I, serotype A1-HA), 7I03-H (group I, serotype A2-

OrfX), Okra (group I, serotype B1-HA), Osaka05 (group I, serotype B6-HA), 111 (group

I, serotype B2-HA/X-OrfX) and *C. sporogenes* strain JCM 1416^{T} (group I; NT, no toxin;

RIKEN BRC) and ATCC 15579 (group I, NT, American Type Culture Collection) were

cultured in TPGY broth (5% Difco^{TM} Tryptone Peptone (Thermo Fisher Scientific), 0.5%

Bacto^{TM} Peptone (Thermo Fisher Scientific), 0.5% Yeast Extract (Merck), 0.1% Glucose

(FUJIFILM Wako Chemicals), 0.1% Sodium Thioglycolate (FUJIFILM Wako

Chemicals) or on TPGY agar plates (1.5% agar) at 37°C in an anaerobic chamber

(HIRASAWA) or anaeropack system (Sugiyama-gen), unless otherwise described. All

bacterial experiments using *C. botulinum* were conducted in the Fujinaga Lab, approved

by the biosafety committee of Kanazawa University and the Minister of Health, Labour

and Welfare of Japan.

Conjugal transformation

Conjugal transformation was performed according to a standard conjugation protocol

(original protocol) (7, 14). Chemically competent cells (15) of *E. coli* CA434 (14) were

transformed with pMTL83153 (16) harboring a codon-optimized *mCherry* (GeneScript)

gene (pMTL83153-mCherry-opt) using the heat shock method at 42°C for 1 min. The

cells were cultured in Luria-Bertani (LB) broth with 30 μg/mL kanamycin and 30 μg/mL

chloramphenicol (Kan/Cm) at 37°C under aerobic conditions. 1 mL of overnight cultures

of CA434 with pMTL83153-mCherry-opt was pelleted by centrifugation and washed

with 1 mL of Nutrient Broth (0.5% Bacto™ Peptone, 0.3% Yeast Extract, 0.5% NaCl).

The pellets were resuspended in 200 μL of an overnight culture of *C. botulinum* or *C.*

sporogenes and 50 μL (5 × 10 μL) of the suspensions were spotted onto TPGY agar plates.

After anaerobic culture at 37°C for 12-24 hr, the cells were harvested in 1 mL of TPGY

broth. 100 μL/plate of the conjugated cell suspensions were inoculated onto TPGY agar

plates with 15 μg/mL thiamphenicol and 250 μg/mL D-cycloserine (Tm/Cs). The plates

were incubated at 37°C for 1-3 days.

For some experiments, this original protocol was adopted as follows. (I) The pre-

culture of CA434 cells harboring pMTL83153-mCherry-opt was harvested at the early-

log phase (OD600 ~ 0.1-0.2), mid-log phase (OD600 ~ 0.5-1.0) or stationary phase

(overnight culture; 12-16 hr). (II) CA434 cells were cultured in LB, Terrific Broth (TB),

or SOC media. (III) CA434 cell pellets were suspended in group I cell cultures and spotted

onto TPGY, BHIS (1 × Pearlcore Brain Heart Infusion agar (Eiken Chemical), 0.5%

Yeast Extract (Merck), 0.2% Fructose (FUJIFILM Wako Chemicals), 0.1% L-Cysteine

(Peptide Institute)) or LB agar plates.

Heat-shock treatment

500 μL of the group I cell cultures were transferred to 1.5 mL O-ring Seal Screw Cap

Tube (WATSON). The tightly capped tubes were incubated at 45°C for 15 min outside

an anaerobic chamber.

pXMTL plasmid construction

The DNA sequence of plasmids pXM (13) and pMTL83151 (16) was amplified by

PCR   using   primers   (pXM: GTTTTAAcacagaaaggatgattgttatg/

GTTGGGTCGtgcaggggggcccgatcg,   and   pMTL: CCCTGCAcgacccaacactctcctac/

TTCTGTGttaaaacaaggatttttccttg). The sequence containing *mazF*, *xylR* and multiple

cloning site (MCS) of pXM and the sequence containing the *traJ*, replication origin of *E.*

coli and *catP* of pMTL83151 were ligated using Gibson Assembly kit (New England

Biolabs) (Fig. 4A). The ligated plasmid (termed pXMTL) was transformed into

chemically competent *E. coli* DH5α cells (TaKaRa Bio). Cm-resistance and xylose-

inducible *mazF* expression of the plasmid were confirmed by culturing the transformants

on LB agar plates with Cm (LB-Cm) and 1% xylose (LB-Cm/Xyl), respectively.

For gene knockout of *bontA*, each 1,500 bp fragment upstream from the second codon

of *bontA* (termed LHA: left-hand arm) and downstream from the second to last codon of

that (termed RHA: right-hand arm) were amplified from genomic DNA extracted from *C.*

botulinum   7I03-H   or   62A   with   primers   (ΔbontA_LHA-BamHI_for:

ataGGATCCgaaattctaatatatcatctgataac,   ΔbontA_LHA-SOE_rev:

AATTACAGtggcatatttaacacctc,   ΔbontA/7I03_RHA-SOE_for:

ATATGCCActgtaattaatctcaaactatatgaac,   ΔbontA/7I03_RHA-PstI_rev:

cttCTGCAGaagcataccacaatgcatatc,   ΔbontA/62A_RHA-SOE_for:

ATATGCCActgtaattaatctcaaactacatg,   ΔbontA/62A_RHA-PstI_rev:

cttCTGCAGtcaataggtttggacgttctatag). The LHA-RHA overlapping sequences were

amplified using splicing by overlap extension PCR (SOE-PCR) and cloned into the

*Bam*HI/*Pst*I site of pXMTL (pXMTL-bontA).

Gene knockout

For the first recombination, conjugal transformation was performed using the

optimized protocol. Briefly, CA434 cells harboring pXMTL-bontA were grown to mid-

log or stationary phases in LB-Kan/Cm medium. The CA434 cells and an overnight

culture of *C. botulinum* were mated on BHIS agar plates. Conjugants were selected on

TPGY agar plates with 7.5 μg/mL thiamphenicol and 250 μg/mL D-cycloserine

(Tm7.5/Cs) for strain 7I03-H or Tm/Cs for strain 62A. Resultant colonies of

transformants were suspended in 100 μL of TE (10 mM Tris pH 8.0, 1 mM EDTA) and

microwaved at 700 W for 2 min. To check the first single crossover recombination, PCR

was performed with 1 μL of the cell suspensions and specific primers (pXMTL-IN_for:

ACAGCCATAACCCTTTAAATTATACATTC, ΔbontA/7I03-check_rev:

TTGCCATATACCACACTTTC, ΔbontA/62A-check_rev:

TATTGTTCTTATCTGTACGAAAC). Clones replicated on TPGY-Tm7.5/Cs or TPGY-

Tm/Cs agar plates were cultured in TPGY broth without antibiotics, and the cultures were

inoculated onto TPGY agar plates with 1% xylose (TPGY-Xyl). The second double

crossover recombination was checked by colony PCR with specific primers

(ΔbontA/7I03-check_for: AAAATCAATATTAGCTCAAGAAAC, ΔbontA/7I03-

check_rev, ΔbontA/62A-check_for: ATCCAAATATATCTATGTGTATCTC,

ΔbontA/62A-check_rev) and by direct Sanger DNA sequencing. Clones replicated on

TPGY-Xyl agar plates were cultured in TPGY broth, and stored in 18% (w/v) glycerol at

-80°C.

Western blotting

C. botulinum 7I03-H cells were cultured in 1 mL of TPGY broth, pelleted by

centrifugation, and lysed in 100 μL of B-PER™ Complete Bacterial Protein Extraction

Reagent (Thermo Fisher Scientific). After heating at 95°C for 5 min, the proteins were

separated by SDS-PAGE and transferred to PVDF membranes (Bio-Rad). The

membranes were blocked with 5% skim milk/PBS-T, and incubated with a rabbit anti-

A16S (62A) polyclonal antibody, followed by an HRP-conjugated anti-rabbit IgG

antibody. Chemiluminescence of the membranes was detected using ECL-select (Cytiva)

and LAS4000 mini (Cytiva).

Results and Discussion

Transformation efficiency of group I *C. botulinum* and *C. sporogenes*

We first assessed the conjugal transformation efficiency of group I *C. botulinum* strains

(62A, 7I03-H, Okra, Osaka05, 111) and *C. sporogenes* strains (JCM 1416^T, ATCC

15579) using pMTL83153-mCherry-opt. These cells were conjugated with *E. coli* CA434

cells harboring a plasmid by the original conjugation protocol (7, 14). Strains 62A, 7I03-

H, Osaka05, 111, JCM 1416^T and ATCC 15579 exhibited transformation efficiencies of

approximately 4.5×10^3, 23, 9.0×10^3, 2.5×10^4, 6.0×10^2 and 8.0×10^2 transformants

per plate, respectively (Fig. 1). No transformants were obtained in strain Okra (Fig. 1).

The *C. botulinum* strains produce BoNTs (62A, serotype A1 (17); 7I03-H, serotype A2

(18); Okra, serotype B1 (19); Osaka05, serotype B6 (20); 111, serotype B2 and X) (21,

22), but the *C. sporogenes* strains do not. The genes encoding BoNT are located on

chromosomes or plasmids (23): the chromosome in strains 62A and 7I03-H, the plasmid

in strains Okra and Osaka05, both in strain 111. There are two types of *bont* gene clusters

(23): the *ha*-type gene cluster in strains 62A, Okra and Osaka05; the *orfX*-type gene

cluster in strain 7I03-H; both in strain 111. Strain 62A is a long-standing laboratory strain.

Strain Okra was isolated from foodborne botulism cases, and strains 7I03-H (24),

Osaka05 (18) and 111 (25) were related to infant botulism cases. According to core gene

phylogenetic analysis, strains 62A, 7I03-H (genetically similar to strain Kyoto-F) and

Okra belong to the group I *C. botulinum* clade. In contrast, strains Osaka05 and ATCC

15579 belong to the *C. sporogenes* clade (26). These data suggest that transformation

efficiencies are not associated with these strains' phylogenetic or pathogenic

characteristics. It has been reported that CLSPO_c06750-knockout cells of *C. sporogenes*

NCIMB 10969 (JCM 1416^T), which is a mutant deficient in the Type IV restriction

system, has a 10-fold higher transformation efficiency than wild-type (WT) (7); *i.e.,*

overcoming the restriction barrier increases transformation efficiency. According to the

REBASE database (27), these strains used in this study employ diverse restriction-

modification (RM) systems (Fig. S1). These defense systems against foreign DNA might

affect the transformation efficiencies of these strains.

Optimizing the transformation protocol

To expand the range of strains capable of being transformed, we optimized the conjugal

transformation method (Fig. 2A). First, we assessed the effect of the culture media for the

donor CA434 cells on the efficiencies. Donor cells cultured in LB medium tended to yield

higher efficiency of DNA transfer to *C. botulinum* 62A compared to TB medium,

although the difference was not statistically significant (Fig. 2B). Second, we assessed

the effect of growth phase of the CA434 cells. Donor cells in the mid-log and stationary

(overnight culture) phases yielded higher efficiency than those in the early-log phase (Fig.

2C). Lastly, the mating was conducted on TPGY, BHIS or LB agar plates. It is known

that culture media during mating influence the transformation efficiency of *C. difficile*

(28). Consistent with this, mating conducted on BHIS agar plates exhibited a 10-fold

higher transformation efficiency of *C. botulinum* 62A than on TPGY agar plates (Fig.

2D). Collectively, the transformation protocol is optimized as below: CA434 cells are

pre-cultured in (I) LB medium, and harvested at (II) the stationary phase (overnight

culture). The cells were mated on (III) <u>BHIS agar plates</u>. This optimized protocol confers

higher transformation efficiencies for strains 62A, 7I03-H, Osaka05, 111 and JCM 1416^T

than the original protocol: 6.3-fold (62A), 5.6-fold (Osaka05), 6.5-fold (111) and 4.2-fold

(JCM 1416^T) (Fig. 3).

It is known that exposing recipient cells to heat shock at 52°C for 5 min before mixing

with donor cells significantly increases the transformation efficiency of *C. difficile* (28)

and other bacteria (29–31). Consistent with this, pre-treatment by heat shock at 45°C for

15 min tended to increase the transformation efficiency of certain strains of group I *C.*

botulinum, albeit not significantly (Fig. 3).

Gene knockout in *C. botulinum* 7I03-H

We established a markerless mutant generation system by ACE using pXM in *C.*

perfringens (13). pXM is a *mazF*-based suicide vector that is counter-selectable for xylose

sensitivity and is introduced into *C. perfringens* strains by electroporation (13). Two

methods are routinely pursued to enable DNA transfer to clostridial recipients:

electroporation and conjugation. Plasmids are capable of being introduced into *C.*

botulinum Hall (group I, A1-HA) by electroporation at ~10^3 transformants/μg of DNA

(32). Consistent with this study, we obtained a similar efficiency with strain 62A by

electroporation, whereas no transformants were obtained with strain 7I03-H (data not

shown). pMTL80000 vectors are transferred at high efficiency by conjugation from *E.*

coli CA434 (HB101 carrying R702) donor strain to *Clostridium* spp. (14, 16). We

developed a novel conjugal suicide vector (termed pXMTL) derived from pXM (13) and

pMTL83151 (16) (Fig. 4A). pXMTL contains a MCS, *catP*, replication origin of *E. coli*,

traJ, *mazF* and *xylR*. The plasmid is a xylose-inducible *mazF*-based suicide vector (Fig.

4A) and can be transferred to *Clostridium* spp. by conjugation. To delete the *bontA* gene

in *C. botulinum* 7I03-H, LHA and RHA were designed and cloned into the MCS of

pXMTL (Fig. 4). LHA contains 1,500 bp of sequence upstream from the second codon

of *bontA*. RHA contains the same length (1,500 bp) of sequence downstream from the

second to last codon (the codon before the stop codon). The first single crossover event

integrates the suicide plasmid into the bacterial chromosome, resulting in

chloramphenicol resistance (Fig. 4B). Due to lack of a replication origin of *Clostridium*

spp. (Fig. 4A), single crossover events will occur in Cm-resistant isolates. It is known

that the length of the homology arms is critical for the frequency of homologous

recombination in *C. difficile* (33). Our preliminary experiments obtained few or no first

recombinants using pXMTL-bontA with shorter homology arms (500 or 1,000 bp, data

not shown). pXMTL with 1,500 bp arms resulted in N- and C-crossover events in strain

7I03-H (Fig. 5A). The first recombinants were induced by xylose to undergo second

crossover events, yielding knockout (KO) and revertant (RT) cells (Fig. 4B, 5). As a result,

8 of 15 clones (53.3%) were *bontA*-KO cells (Fig. 5A). The precise genome editing and

the loss of BoNT/A expression were confirmed by Sanger DNA sequencing (Fig. 5B) and

western blotting (Fig. 5C), respectively. The ACE system also deleted the target genes

using pXMTL in *C. botulinum* 62A and *C. sporogenes* JCM 1416^{T} (data not shown). This

system provides a rapid and markerless gene mutagenesis method for group I *C.*

botulinum and *C. sporogenes*.

Conclusions

C. botulinum causes severe poisoning by producing BoNT (1), and *C. sporogenes*, a

bacterium closely related to group I *C. botulinum*, is a key contributor to the production

of bioactive metabolites in the human gut (2, 3). The ecologies of these organisms,

including sporulation, germination, outgrowth and infection, remain elusive and functions

of their genes are largely unknown. One reason might be because these bacteria exhibit

low transformation efficiencies (Fig. 1). In the present study, we optimized the conjugal

transformation protocol and yielded higher efficiencies than the original protocol. In

addition, we developed a novel conjugal *mazF*-based suicide vector pXMTL harboring a

xylose-inducible promoter, and provided a rapid method for precise, markerless and

scarless genome editing by ACE. This system will facilitate the analysis of gene function

in group I *C. botulinum* and *C. sporogenes.*

Acknowledgments

We thank Shunji Kosaki, Tomoko Kohda, Kaoru Umeda, Shin-ichi Nakamura for

generous gifts of *C. botulinum* strains. We thank Editage (www.editage.com) for English

language editing. We also thank Sachiyo Akagi, Kazumi Kuraoka, Yuki Konoshita for

technical assistance and all members of the Fujinaga Lab. for valuable discussions.

Conflicts of interest

The authors declare no conflicts of interest.

Source of funding

This work was supported by JSPS KAKENHI (18H02654 to YF and 20J23183 to KS).

Contribution of authors

SA and YF conceived and designed the experiments. SA performed the experiments and

data analysis. KS and HN assisted in plasmid construction. SA and YF wrote the original

draft of the manuscript. All authors have discussed the results and approved the

manuscript.

Abbreviations

ACE, allele-coupled exchange; CRISPR, clustered regularly interspaced short

palindromic repeats; Cas9, CRISPR-associated protein 9; BoNT, botulinum neurotoxin;

HA, hemagglutinin; NT, no toxin; LB, Luria-Bertani; TB, Terrific broth; SOC, super

optimal broth with catabolite repression; TPGY, tryptone-peptone-glucose-yeast extract;

BHIS, brain heart infusion-supplemented; MCS, multiple cloning site; Kan, kanamycin;

Cm, chloramphenicol; Tm, thiamphenicol; Cs, D-cycloserine; Xyl, xylose; LHA, left-

hand arm; RHA, right-hand arm; SOE-PCR, splicing by overlap extension PCR; WT,

wild-type; RT, revertant; KO, knockout.

References

1. Collins MD, East AK. 1998. Phylogeny and taxonomy of the food-borne pathogen
Clostridium botulinum and its neurotoxins. J Appl Microbiol 84:5–17.

2. Dodd D, Spitzer MH, Treuren WV, Merrill BD, Hryckowian AJ, Higginbottom
SK, Le A, Cowan TM, Nolan GP, Fischbach MA, Sonnenburg JL. 2017. A gut
bacterial pathway metabolizes aromatic amino acids into nine circulating
metabolites. 7682. Nature 551:648–652.

3. Liu Y, Chen H, Van Treuren W, Hou B-H, Higginbottom SK, Dodd D. 2022.
Clostridium sporogenes uses reductive Stickland metabolism in the gut to generate
ATP and produce circulating metabolites. 5. Nat Microbiol 7:695–706.

4. Heap JT, Pennington OJ, Cartman ST, Carter GP, Minton NP. 2007. The
ClosTron: A universal gene knock-out system for the genus Clostridium. J
Microbiol Methods 70:452–464.

5. Minton NP, Ehsaan M, Humphreys CM, Little GT, Baker J, Henstra AM, Liew F,
Kelly ML, Sheng L, Schwarz K, Zhang Y. 2016. A roadmap for gene system
development in Clostridium. Anaerobe 41:104–112.

6. Guo C-J, Allen BM, Hiam KJ, Dodd D, Treuren WV, Higginbottom S, Nagashima
K, Fischer CR, Sonnenburg JL, Spitzer MH, Fischbach MA. 2019. Depletion of
microbiome-derived molecules in the host using Clostridium genetics. Science
366.

7. Woods C, Humphreys CM, Rodrigues RM, Ingle P, Rowe P, Henstra AM, Köpke
M, Simpson SD, Winzer K, Minton NP. 2019. A novel conjugal donor strain for
improved DNA transfer into Clostridium spp. Anaerobe 59:184–191.

8. Cañadas IC, Groothuis D, Zygouropoulou M, Rodrigues R, Minton NP. 2019.
RiboCas: A Universal CRISPR-Based Editing Tool for Clostridium. ACS Synth
Biol 8:1379–1390.

9. Mertaoja A, Nowakowska MB, Mascher G, Heljanko V, Groothuis D, Minton NP,
Lindström M. 2021. CRISPR-Cas9-Based Toolkit for Clostridium botulinum
Group II Spore and Sporulation Research. Front Microbiol 12:617269.

10. Ng YK, Ehsaan M, Philip S, Collery MM, Janoir C, Collignon A, Cartman ST,
Minton NP. 2013. Expanding the Repertoire of Gene Tools for Precise
Manipulation of the Clostridium difficile Genome: Allelic Exchange Using pyrE
Alleles. PLOS ONE 8:e56051.

11. Cartman ST, Kelly ML, Heeg D, Heap JT, Minton NP. 2012. Precise Manipulation
of the Clostridium difficile Chromosome Reveals a Lack of Association between
the tcdC Genotype and Toxin Production. Appl Environ Microbiol 78:4683–4690.

12. Al-Hinai MA, Fast AG, Papoutsakis ET. 2012. Novel System for Efficient

Isolation of Clostridium Double-Crossover Allelic Exchange Mutants Enabling
Markerless Chromosomal Gene Deletions and DNA Integration. Appl Environ
Microbiol 78:8112–8121.

13. Nariya H, Sawairi S, Tamai E, Yano C T, Shimamoto T, Miyata S. Development
of a novel method for efficient and precise genome editing with mazF in
Clostridium perfringens. in preparation.

14. Purdy D, O'Keeffe TAT, Elmore M, Herbert M, McLeod A, Bokori-Brown M,
Ostrowski A, Minton NP. 2002. Conjugative transfer of clostridial shuttle vectors
from Escherichia coli to Clostridium difficile through circumvention of the
restriction barrier. Mol Microbiol 46:439–452.

15. Inoue H, Nojima H, Okayama H. 1990. High efficiency transformation of
Escherichia coli with plasmids. Gene 96:23–28.

16. Heap JT, Pennington OJ, Cartman ST, Minton NP. 2009. A modular system for
Clostridium shuttle plasmids. J Microbiol Methods 78:79–85.

17. Wentz TG, Yao K, Schill KM, Reddy NR, Skinner GE, Morrissey TR, Wang Y,
Muruvanda T, Manickam G, Pillai CA, Thirunavukkarasu N, Hoffmann M,
Hammack TS, Brown EW, Allard MW, Sharma SK. 2018. Closed Genome
Sequence of Clostridium botulinum Strain CFSAN064329 (62A). Genome
Announc 6:e00528-18.

18. Umeda K, Seto Y, Kohda T, Mukamoto M, Kozaki S. 2009. Genetic
Characterization of Clostridium botulinum Associated with Type B Infant
Botulism in Japan. J Clin Microbiol 47:2720–2728.

19. Smith TJ, Hill KK, Foley BT, Detter JC, Munk AC, Bruce DC, Doggett NA,
Smith LA, Marks JD, Xie G, Brettin TS. 2007. Analysis of the Neurotoxin
Complex Genes in Clostridium botulinum A1-A4 and B1 Strains: BoNT/A3, /Ba4
and /B1 Clusters Are Located within Plasmids. PLOS ONE 2:e1271.

20. Sakaguchi Y, Hosomi K, Uchiyama J, Ogura Y, Umeda K, Sakaguchi M, Kohda
T, Mukamoto M, Misawa N, Matsuzaki S, Hayashi T, Kozaki S. 2014. Draft
Genome Sequence of Clostridium botulinum Type B Strain Osaka05, Isolated
from an Infant Patient with Botulism in Japan. Genome Announc 2:e01010-13.

21. Hosomi K, Sakaguchi Y, Kohda T, Gotoh K, Motooka D, Nakamura S, Umeda K,
Iida T, Kozaki S, Mukamoto M. 2014. Complete nucleotide sequence of a plasmid
containing the botulinum neurotoxin gene in Clostridium botulinum type B strain
111 isolated from an infant patient in Japan. Mol Genet Genomics 289:1267–1274.

22. Zhang S, Masuyer G, Zhang J, Shen Y, Lundin D, Henriksson L, Miyashita S-I,
Martínez-Carranza M, Dong M, Stenmark P. 2017. Identification and
characterization of a novel botulinum neurotoxin. Nat Commun 8:1–10.

23. Hill KK, Smith TJ. 2013. Genetic Diversity Within Clostridium botulinum
Serotypes, Botulinum Neurotoxin Gene Clusters and Toxin Subtypes. Curr Top
Microbiol Immunol 364:1–20.

24. Tabita K, Sakaguchi S, Kozaki S, Sakaguchii G. 1990. Comparative studies on
Clostridium botulinum type A strains associated with infant botulism in Japan and
in California, USA. Jpn J Med Sci Biol 43:219–231.

25. Kakinuma H, Maruyama H, Takahashi H, Yamakawa K, Nakamura S. 1996. The
first case of type B infant botulism in Japan. Acta Paediatr Jpn 38:541–543.

26. Weigand MR, Pena-Gonzalez A, Shirey TB, Broeker RG, Ishaq MK,
Konstantinidis KT, Raphael BH. 2015. Implications of Genome-Based
Discrimination between Clostridium botulinum Group I and Clostridium
sporogenes Strains for Bacterial Taxonomy. Appl Environ Microbiol 81:5420–
5429.

27. Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for DNA
restriction and modification: enzymes, genes and genomes. Nucleic Acids Res
43:D298–D299.

28. Kirk JA, Fagan RP. 2016. Heat shock increases conjugation efficiency in
Clostridium difficile. Anaerobe 42:1–5.

29. Sun F-H, Luo D, Shu D, Zhong J, Tan H. 2014. Development of an Intergeneric
Conjugal Transfer System for Xinaomycins-Producing Streptomyces noursei
Xinao-4. 7. Int J Mol Sci 15:12217–12230.

30. Zeng X, Ardeshna D, Lin J. 2015. Heat Shock-Enhanced Conjugation Efficiency
in Standard Campylobacter jejuni Strains. Appl Environ Microbiol 81:4546–4552.

31. Zhang S, Chen T, Jia J, Guo L, Zhang H, Li C, Qiao R. 2019. Establishment of a
highly efficient conjugation protocol for Streptomyces kanamyceticus
ATCC12853. MicrobiologyOpen 8:e00747.

32. Zhou Y, Johnson EA. 1993. Genetic transformation of Clostridium botulinum hall
a by electroporation. Biotechnol Lett 15:121–126.

33. Faulds-Pain A, Wren BW. 2013. Improved Bacterial Mutagenesis by High-
Frequency Allele Exchange, Demonstrated in Clostridium difficile and
Streptococcus suis. Appl Environ Microbiol 79:4768–4771.

419
420

FIG 1 Transformation efficiency of group I *C. botulinum* and *C. sporogenes* by the

original conjugation protocol.

According to the original conjugation protocol (7, 14), CA434 cells harboring

pMTL83153-mCherry-opt cultured overnight in LB medium were mated with *C.*

botulinum strains (62A, Okra, 7I03-H, 111 and Osaka05) and *C. sporogenes* strains (JCM

1416^T and ATCC 15579) on TPGY agar plates. The transformants were selected on

TPGY agar plates with thiamphenicol and D-cycloserine (Tm/Cs). Values are means ±

SD of triplicates. ND, not detected.

FIG 2 Optimization of conjugation protocol for group I *C. botulinum*.

(**A**) Schematic model of the standard (original) conjugation protocol.

(**B-D**) Transformation efficiency of *C. botulinum* 62A with pMTL83153-mCherry-opt

containing a chloramphenicol (Cm) resistance gene. The conjugation protocol (**A**) was

performed with three modifications: culture media of donor cells (**B**) and growth

phases of donor cells (**C**) and culture media during mating (**D**). Bold font indicates the

original protocol. Values are means ± SD of triplicates. ND, not detected. *$P < 0.05$,

**$P < 0.01$, ***$P < 0.001$; by ANOVA with Tukey's post-hoc HSD test.

FIG 3 Transformation efficiency of group I *C. botulinum* and *C. sporogenes* by the

optimized conjugation protocol.

Plasmid pMTL83153-mCherry-opt was transferred from CA434 cells to *C. botulinum*

strains (62A, Okra, 7I03-H, 111 and Osaka05) and *C. sporogenes* strains (JCM 1416[T] and

ATCC 15579). Transformants were selected on TPGY-Tm/Cs agar plates. Gray, white,

and hatched bars represent transformation efficiencies by the original protocol, the

optimized protocol (BHIS), and the optimized protocol with heat shock, respectively.

Values are means ± SD of triplicates. ND, not detected. *$P < 0.05$, **$P < 0.01$, ***$P <$

0.001; by unpaired, two-tailed Student's *t*-test.

FIG 4 Markerless mutagenesis by allele-coupled exchange (ACE) using a conjugal suicide vector pXMTL.

(**A**) Map of the engineered plasmid pXMTL with homology arms. pXMTL harbors *mazF*, *xylR* and MCS of pXM and *catP*, the replication origin of *E. coli* and *traJ* of pMTL83151.

(**B**) Schematic protocol of ACE using pXMTL, which deletes the *bontA* gene on the chromosome of *C. botulinum* 7I03-H. (1) Left homology arm (LHA) and right homology arm (RHA) targeting *bontA* were cloned into MCS of pXMTL (pXMTL-bontA). (2) At the first recombination step, the recombinant in which the pXMTL-bontA plasmid is crossed is selected on TPGY-Tm7.5/Cs agar plates. (3) At the second recombination step, the plasmid is cured by exposing 1% xylose, resulting in *bontA*-KO mutants (KO) or wild-type revertants (RT).

FIG 5 *bontA* gene knockout in *C. botulinum* 7I03-H.

(**A**) PCR analysis of the first (upper panel) and second (lower panel) recombinants of *C. botulinum* 7I03-H.

(**B**) DNA sequence of interest of *bontA*-KO (#1-4, #2-2, #3-3) cells was confirmed by Sanger DNA sequencing. Sequence chromatograms were visualized using SnapGene Viewer 6.0.2.

(**C**) Protein expression of BoNT/A in WT, *bontA*-KO (#1-4, #2-2, #3-3) and RT (#3-4) cells was confirmed by western blotting.

Fig. S1 Restriction and modification systems in group I *C. botulinum* and *C. sporogenes*. Restriction modification (RM) systems of strain 62A, Kyoto-F (genetically similar to 7I03-H), Okra, Osaka05, 111 and DSM 795[T] (JCM 1416[T]) listed in the REBASE database (27). Brown, orange and cyan colors indicate type I, II and IV RM systems, respectively.