Multi-omic phenotyping of iPSC-derived neurons harboring the *MAPT* V337M mutation reveals

tau hypophosphorylation and perturbed axon morphology pathways

Gregory A. Mohl*[1,2], Gary Dixon*[1], Emily Marzette[1], Justin McKetney[3,4,5], Avi J. Samelson[1],

Carlota Pereda Serras[1,6], Julianne Jin[1], Nabeela Ariqat[1], Andrew Li[1], Steven C. Boggess[1],

Danielle L. Swaney[3,4,5], Martin Kampmann[1,7]

Affiliations

1. Institute for Neurodegenerative Diseases, University of California San Francisco, San

Francisco, CA, USA

2. Department of Neurology, University of California San Francisco, San Francisco, CA, USA

3. Gladstone Data Science and Biotechnology Institute, The J. David Gladstone Institutes, San

Francisco, CA, USA

4. Quantitative Bioscience Institute, University of California San Francisco, San Francisco, CA,

USA

5. Department of Cellular and Molecular Pharmacology, University of California San Francisco,

San Francisco, CA, USA

6. Bakar Computational Health Sciences Institute, University of California San Francisco, San

Francisco, CA USA

7. Department of Biochemistry and Biophysics, University of California San Francisco, San

Francisco, CA, USA

* Contributed equally.

23
24 Correspondence: martin.kampmann@ucsf.edu

Abstract

Tau aggregation is a hallmark of several neurodegenerative diseases, including Alzheimer's disease and frontotemporal dementia. There are disease-causing variants of the tau-encoding gene, *MAPT*, and the presence of tau aggregates is highly correlated with disease progression. However, the molecular mechanisms linking pathological tau to neuronal dysfunction are not well understood. This is in part due to an incomplete understanding of the normal functions of tau in development and aging, and how the associated molecular and cellular processes change in the context of causal disease variants of tau. To address these questions in an unbiased manner, we conducted multi-omic characterization of iPSC-derived neurons harboring the *MAPT* V337M mutation or *MAPT* knockdown. RNA-seq and phosphoproteomics revealed that both V337M mutation and tau knockdown perturbed levels of transcripts and phosphorylation of proteins related to axonogenesis or axon morphology. Surprisingly, we found that neurons with V337M tau had much lower tau phosphorylation than neurons with WT tau. Functional genomics screens uncovered regulators of tau phosphorylation in neurons and found that factors involved in axonogenesis modified tau phosphorylation in both *MAPT* WT and *MAPT* V337M neurons. Intriguingly, the p38 MAPK pathway specifically modified tau phosphorylation in *MAPT* V337M neurons. We propose that V337M tau perturbs tau phosphorylation and axon morphology pathways that are relevant to the normal function of tau, which could contribute to previously reported cognitive changes in preclinical *MAPT* variant carriers.

Introduction

Neurodegenerative diseases are a growing public health burden and remain very challenging to treat because we lack a complete understanding of the underlying disease mechanisms. A common theme in many neurodegenerative diseases is the aggregation of pathological proteins [1]. Tau aggregation is a hallmark of neurodegenerative diseases collectively called tauopathies, including Alzheimer's disease and frontotemporal dementia. In Alzheimer's disease, tau aggregation and phosphorylation changes correlate better with disease progression than amyloid beta pathology [2] despite clear genetic evidence linking amyloid beta to the disease [3]. In frontotemporal dementia, rare causal variants of tau that are fully penetrant for the disease prove a direct role for tau in disease pathogenesis [4].

Tremendous progress has been made in revealing the diverse molecular and cellular mechanisms that are disrupted by pathogenic tau. Recent work in human induced pluripotent stem cell (iPSC)-derived neurons has shown that pathogenic variants of tau sensitize neurons to different types of cellular stress and that this effect can be rescued by lowering tau levels via autophagy [5]. Other groups have shown that tau interferes with RNA splicing and stress granules homeostasis [6-9], disrupts the nuclear envelope [10-12], perturbs axonal trafficking [13, 14] or disrupts mitochondrial dynamics [15]. Acetylated tau has also been shown to disrupt chaperone mediated autophagy, rerouting tau and other clients to be degraded by other mechanisms [16]. Pathogenic tau has also been shown to perturb plasticity of the axon initial segment and cause changes to neuronal excitability [17] and has been implicated in driving excitotoxicity [9, 18-20]. Many of these data support a tau toxic gain-of-function model, and tau lowering has been successfully shown to be beneficial in cultured neurons and animal models [5, 21]. In fact, tau lowering is currently being tested in the clinic by antibodies and ASOs [22].

These focused studies have linked tau to diverse cellular processes that go awry in

neurodegeneration. However, there are few unbiased and comprehensive studies that examine

phenotypes on multiple intracellular levels or with respect to normal tau function, leaving many

open questions about the direct effects of pathogenic tau and how diverse cellular phenotypes

interact.

To characterize the earliest changes that pathogenic tau causes in human neurons and to

understand mechanistically how pathogenic tau causes human disease, we used a multi-omic

approach to unbiasedly determine the cellular phenotypes linked to pathogenic tau. We modeled

pathogenic tau by using human iPSC-derived neurons with the *MAPT* V337M mutation, a known

cause of frontotemporal dementia. We used two sets of iPSCs, one from a healthy donor

(WTC11) and one from a patient with the *MAPT* V337M mutation (GIH6C1) [17, 23].

Our RNA-seq, ATAC-seq, proteomics and phosphoproteomics results all point to

changes in axonogenesis due to the *MAPT* V337M mutation. Recently published mouse

phosphoproteomics datasets in tau knockout mice and P301S mice strongly support the link

between tau and axonogenesis factors and intriguingly suggest that these effects are due to tau

loss of function [24, 25]. We have found that tau knockdown and *MAPT* V337M mutation have

overlapping effects on the levels and phosphorylation of proteins relevant to axonogenesis and

axon morphology, suggesting that the mutation perturbs a normal function of tau. *MAPT* V337M

neurons have hypophosphorylated tau, which is recapitulated by artificially overexpressing

V337M tau but not WT or R406W tau in neurons with endogenous tau knockdown. Unbiased

CRISPR screens for regulators of tau phosphorylation uncovered axonogenesis-related regulators

of tau phosphorylation and show that the p38 MAPK pathway may play a role in modifying tau

phosphorylation specifically in V337M neurons. We propose that V337M tau perturbs tau

phosphorylation and axon morphology pathways that are relevant to the normal function of tau,

which could contribute to previously reported cognitive changes in preclinical *MAPT* variant

carriers.

Results

MAPT V337M and MAPT knockdown perturb the transcription of axonogenesis-related genes

iPSCs generated from a healthy individual (WTC11, referred to as *MAPT* WT [26]) or an

FTD patient with the *MAPT* V337M mutation (GIH6C1, referred to as **MAPT* Het [23]) were

edited in previous work [17, 23] with Cas9 to generate isogenic pairs either introducing or

correcting the *MAPT* V337M mutation (Fig. 1A). The *MAPT* WT iPSCs were edited with Cas9

to generate a heterozygous *MAPT* V337M/WT clone (*MAPT* Het) and homozygous *MAPT*

V337M/V337M clone (*MAPT* Hom). The **MAPT* Het iPSCs were corrected with Cas9 to

generate a *MAPT* WT/WT clone (**MAPT* WT). We engineered the GIH6C1 lines to introduce a

doxycycline-inducible Ngn2 for neuronal differentiation and CRISPRi machinery. We

transduced the iPSCs with lentiviral sgRNAs targeting *MAPT* to knockdown tau or non-targeting

control (NTC) sgRNAs for further mechanistic characterization (Figure S1A-C).

RNA-seq of neurons harvested at 2 and 4 weeks of differentiation revealed overlap

between effects in *MAPT* Het neurons and *MAPT* WT tau knockdown neurons (Figure 1B).

Genes that were differentially expressed in *MAPT* Het neurons and *MAPT* WT tau knockdown

neurons were significantly enriched for regulators of axonogenesis (Figure 1C). Knocking down

tau in *MAPT* Het neurons resulted in only five differentially expressed genes (Figure S1D).

Differentially expressed genes in *MAPT* Hom and **MAPT* Het neurons compared to isogenic

controls were also significantly enriched for regulators of axonogenesis, even at one week of

differentiation (Figure S1E-G). While many of the same transcripts relevant to axonogenesis are

perturbed in *MAPT* Het and *MAPT* Het, we did not see a high level of concordance in direction

of change (Figure S1E). On the other hand, the changes in *MAPT* Het and *MAPT* Hom are

extremely similar, suggesting high concordance between distinct clones in the same genetic

background.

ATAC-seq at 2 and 4 weeks of differentiation showed similar patterns as the RNA-seq

(Figure 1D, Figure S2A), and genes with differentially accessible peaks proximal to their

transcription start site (TSS) were enriched for axon-related genes (Figure S2B). Transcription

factor motif analysis showed that motifs for the AP-1 Transcription factor network, which

includes the cJun family of transcription factors, were consistently more accessible in *MAPT* Het

and *MAPT* WT tau knockdown neurons compared to controls (Figure 1E, Figure S2C).

Supporting the validity of the ATAC-seq results, we found that both p-cJun and cJun are

increased in *MAPT* Het, *MAPT* Hom and *MAPT* Het neurons vs. isogenic controls (Figure 1F-

H). *MAPT* V337M and tau knockdown induce overlapping changes in chromatin accessibility

and transcription of axonogenesis-related genes, suggesting that some phenotypes in *MAPT*

V337M neurons are relevant to normal tau function.

MAPT V337M and tau knockdown perturb phosphorylation of axonogenesis-related proteins

We hypothesized that changes in cJun and p-Jun may reflect broad changes in

intracellular signaling caused by V337M tau. To identify shifts in signaling occurring at

relatively early stages of axonogenesis, we determined the total proteome and phosphoproteome

of 1 week neurons with *MAPT* V337M and/or tau knockdown by mass spectrometry.

Phosphoproteomic analysis of *MAPT* V337M neurons confirmed elevated p-cJun levels while

also uncovering differential phosphorylation of proteins regulating neuron projection

development and splicing (Figure 2A,C). There was significant overlap in the proteins with

differential phosphorylation between *MAPT* Hom, *MAPT* Het and *MAPT* Het neurons vs.

isogenic controls (Figure 2B), though the identities of the differential phosphosites varied

between conditions (Figure S3A). Gene set enrichment analysis for the 56 conserved proteins

with changes in phosphorylation in *MAPT* V337M neurons showed that the top enriched terms

were related to neuron projection development (Figure 2C). The total protein levels for many of

these factors were not significantly changed, suggesting that these changes are due to specific

signaling events altering phosphorylation patterns, rather than just changes in protein levels

(Figure S3B-D).

We next compared our phosphoproteomic datasets to recently published mouse

phosphoproteomic datasets using tau knockout mice [24] or P301S tau mice [25]. We found

significant overlap for proteins with differential phosphorylation in our data and the tau knockout

mice but not with the P301S mice (Figure 2D). However, we also noted that there was significant

overlap between the tau knockout mice and the P301S mice. Gene set enrichment analysis

identified substantial enrichment of axonogenesis-related protein phosphorylation changes in the

45 conserved proteins with differential phosphorylation in *MAPT* V337M neurons, tau knockout

mice, and P301S tau mice (Figure 2E). When we determined an even more focused set of

proteins that also have differential phosphorylation in *MAPT* V337M homozygous and *MAPT*

V337M heterozygous neurons from patient iPSCs, we found a core network of proteins in highly

related pathways regulating neuron morphogenesis and polarity (Figure 2F), including *ANK3* and

MAPRE3. *ANK3* and *MAPRE3* were recently identified to be important for V337M tau-induced

defects in axon initial segment plasticity [17].

We observed two patterns of protein phosphorylation changes due to tau knockdown

(Figure 2G). Many phosphorylation changes were specific to either tau knockdown in the *MAPT*

WT neurons or the *MAPT* V337M neurons. When we performed gene set enrichment analysis on

proteins with differential phosphorylation in *MAPT* WT tau knockdown neurons, the only

significantly enriched term was "Regulation of microtubule-based process," with many of these

proteins being involved in axonogenesis (Figure S3E). Gene set enrichment analysis of proteins

with differential phosphorylation in *MAPT* V337M tau knockdown compared to *MAPT* V337M

showed that splicing factors were predominantly affected, whereas cytoskeletal and

axonogenesis proteins were not perturbed (Figure S3F).

V337M tau is hypophosphorylated in neurons

We observed that *MAPT* V337M neurons had lower tau phosphorylation compared to

WT across all domains of the protein at many sites (Figure 3A and 3B) and validated these

changes by western blot in all sets of neurons (Figure 3C-E). Many of the differential

phosphorylation sites are known to be hyperphosphorylated in Alzheimer's disease and other

tauopathies [27-29] (Figure S4A).

To further explore how V337M tau may have decreased phosphorylation in neurons, we

overexpressed WT tau, V337M tau or R406W tau in *MAPT* WT neurons with endogenous tau

knocked down. Consistent with our phosphoproteomics results, V337M tau had decreased

phosphorylation at numerous sites despite having similar tau levels to WT tau and R406W tau

(Figure 3F,G). Intriguingly, R406W tau only had decreased phosphorylation at some of these

sites. These data suggest tau variants affect tau phosphorylation in neurons via distinct

mechanisms.

Extensive work has been done to characterize tau phosphorylation sites and map them to

their kinases [30-35]. Proline-directed phosphorylation sites were decreased in *MAPT* V337M

neurons, many of which serve as priming sites for additional sites of decreased tau

phosphorylation (Figure S4B). Leveraging our global view of phosphorylation changes in *MAPT*

V337M neurons, we predicted which kinases may have changes in activity based on known

kinase-substrate relationships (Figure 3H). Kinases in the p38 MAPK pathway such as *MAP2K3*

and *MAP2K6* were predicted to have increased activity in *MAPT* V337M neurons (Figure 3I).

MAPK11 and *MAPK14* targets had increased phosphorylation specifically in *MAPT* V337M

neurons with tau knockdown, whereas *MAPK12* substrates had decreased phosphorylation

specifically in *MAPT* WT neurons with tau knockdown. Known tau kinases with well-

documented roles in tauopathy were also predicted to have differential activity, including

GSK3B, *CDK5,* and *CDK5R1*. CDK5 and p38 MAPKs are both proline-directed kinases that are

known to phosphorylate tau at several sites that had decreased phosphorylation in *MAPT* V337M

neurons.

CRISPR screens uncover regulators of tau phosphorylation in neurons

To directly test which kinases perturb tau phosphorylation in *MAPT* WT and *MAPT*

V337M neurons, we employed CRISPRi and CRISPRa screens to test the effects of gene

knockdown or overexpression on tau phosphorylation using the AT8 antibody, which detects the

tau pS202/pT205 phosphoepitope (Figure 4A, Figure S5A-D). We transduced iPSCs with a

lentiviral sgRNA library targeting 2,325 genes encoding kinases, phosphatases and other proteins

in the "druggable genome"[36]. Two weeks after differentiation, neurons were fixed and stained

with AT8 and sorted based on AT8 signal. Next generation sequencing identified genes that

causally regulate AT8 levels. We filtered out hits for enrichment analysis that also modified T22

levels in previously published work (Figure S5E) [37]. Cytoskeleton genes and genes involved in

neuron projection development modified tau phosphorylation in both *MAPT* WT and *MAPT*

V337M neurons (Figure 4A,B) without altering T22 levels (Figure S5E). Intriguingly, several

kinases in the p38 MAPK pathway altered tau phosphorylation specifically in *MAPT* V337M

neurons. Other kinases predicted to have differential activity that may have regulated tau

phosphorylation in *MAPT* V337M neurons did not affect tau pS202/pT205 levels, including

CDK5, *CDK5R1*, and *GSK3B* (Figure S5F). We mapped the detected tau phosphorylation sites in

our neurons to their known kinases based on the literature, overlaying phosphorylation sites that

were differential in *MAPT* V337M (blue) with kinases whose knockdown or overexpression

modified tau phosphorylation at S202/T205 (red) (Figure 4D). The overlap between differential

tau phosphorylation and kinases that regulate pS202/pT205 in neurons (purple) narrows the list

down to a few candidate kinases. Overexpression of *MARK1*, a kinase that phosphorylates tau in

the microtubule binding domain and regulates tau's interaction with microtubules, caused

increased tau phosphorylation in *MAPT* WT neurons (Figure 4C). This is consistent with

previous work showing that phosphorylation at S262, S324 and S356 affects phosphorylation

sites distal from the microtubule binding domain, such as S202/T205 [38].

Discussion

We have discovered that an FTD-causing variant of tau leads to tau hypophosphorylation

and perturbs axonogenesis pathways in differentiating neurons, overlapping at least in part with

effects seen in tau knockdown. These findings are surprising because disease-associated tau is

typically associated with increased tau phosphorylation and would not be expected to have

shared phenotypic overlap with tau loss. Other groups have shown in mice or in primary neurons

that reducing tau can have varying effects on axonogenesis. Acute tau ablation in mouse neurons

in vitro prevents axonogenesis by inhibiting polarization [39, 40] and tau knockout in primary

hippocampal neurons and human iPSC-derived neurons reduces neurite outgrowth and branching

[41, 42].

The question remains whether a loss of tau function (potentially caused by a tau

mutation) would have adverse effects to a disease variant carrier throughout life and with respect

to disease. The function of normal healthy tau is unclear and has been debated for many years.

This is in large part due to the many conflicting studies, both in physiological and pathogenic

contexts. Given the earlier results in showing the importance of tau for axonogenesis, it was

expected that knocking out tau in mice would be lethal and that tau would be essential for

neurodevelopment. Early mouse studies showed that tau knockout was surprisingly well

tolerated [43]. There were no obvious defects in polarization or gross morphology, but

microtubules in small caliber axons were destabilized. *Map1a* was upregulated in tau knockout

mice, suggesting that the mice were compensating for tau loss. This could explain the difference

in phenotypes as compared to the acute depletion of tau with ASOs. Knocking out tau and

Map1b, another microtubule-associated protein, leads to much more severe phenotypes than

either knockout individually [44]. Dawson et al. disputed the findings of Harada et al. due to

poor WT data [45]. In their work, they found that indeed tau knockout did cause a delay in

neurite outgrowth and axonogenesis.

Biswas and Kalil showed that tau knockout neurons had altered microtubule dynamics in

growth cones, resulting in a change in overall growth cone morphology [46]. Microtubules were

less bundled, and microtubule polymerization directionality as measured by EB3 was more

dispersed in tau knockout neurons. There also was a reduction in tyrosinated tubulin projecting

into the filopodia of the peripheral domain. Another paper showed that tau knockout increased

Fyn mobility in dendrites and lowered Fyn localization in dendrites and spines [47]. Intriguingly,

expressing P301L tau had the opposite effect and anchored or trapped Fyn in dendritic spines.

Many motor and behavioral phenotypes have been observed in tau knockout mice. Tau

knockout mice or mice with acute tau reduction with antisense oligonucleotides have

consistently shown resistance to seizures [21, 48-51]. Another consistent theme is that there are

often behavioral and learning changes in tau knockout mice, including hyperactivity, fear

conditioning, and memory [52-57]. There is more controversy over the effect of tau knockout on

motor function. Some groups report motor deficits in tau knockout mice [52, 58, 59], while

others claim there are no significant changes in tau knockout mice to motor function [48, 49, 57].

One group showed that tau is essential for long term depression in the hippocampus [60], while

another showed that tau knockout only perturbs long term potentiation [57]. Tau phosphorylation

has also been shown to be required for long term depression [61].

Considering our data in the context of these other findings, we expect that loss-of-

function phenotypes would coincide with the onset of tau expression and axon extension. Tau

loss of function may precede human disease onset by many decades, occurring during

development and continuing through adulthood via perturbed synaptic plasticity. A study showed

that mice with the *MAPT* P301L mutation show early cognitive changes before tau pathology is

detectable [62]. A Parkinson's disease GWAS study found that *MAPT* was a significant risk

locus for Parkinson's disease that is uncoupled from the age of onset [63]. Ye and colleagues

proposed that tau may drive changes during development or early in life that then increase risk

for Parkinson's disease decades later [64]. Two studies have also identified cognitive differences

between *MAPT* carriers and non-carrier siblings decades before expected disease onset [65, 66].

Our work also emphasized the importance of having iPSCs from multiple individuals and

multiple clones paired with appropriate controls, such as tau knockdown and knockout.

Furthermore, comparisons to other published data sets revealed previously underappreciated

relationships, including overlapping molecular phenotypes between *MAPT* knockout and *MAPT*

P301S mice. It will be fascinating to uncover the mechanisms of these shared signaling pathway

changes and to determine if they are due a shared stress response, or if downstream phenotypes

converge despite unique upstream perturbations. Previous work using different differentiation

protocols and much longer time scales showed that FTD-causing tau variants cause tau

hyperphosphorylation in more mature neurons, suggesting that there is a complex, time-

dependent effect of *MAPT* mutations on developmental tau phosphorylation patterns. There was

substantial overlap between our RNA-seq findings and a recent paper using *MAPT* V337M

neurons in an organoid model, which is interesting because of the observed tau

hyperphosphorylation at the later timepoints in their model [9]. Intriguingly, one other group

reported decreased tau phosphorylation in organoid-derived iPSC neurons with *MAPT* R406W.

[67] Earlier work showed that fetal tau was highly phosphorylated during development, but the

precise mechanisms and functions of this process are still unknown.[32, 68-70] Our findings

suggest that at earlier timepoints different tau mutations may behave in unexpected ways and

have complex effects on cellular pathways. [5, 9, 71]

We acknowledge that there are limitations to our study. Our neurons under the conditions

we used only express a single isoform of tau, the fetal isoform 0N3R. Understanding how

different tau isoforms are regulated and how they contribute tau function in health and disease is

an open question. Additionally, it will be intriguing to understand how different disease variants

of tau perturb neurons. Our data showing phosphorylation differences between WT tau, V337M

tau and R406W tau joins a growing body of literature showing that different mutations have

different effects on tau properties, including microtubule binding, microtubule polymerization,

and fibril formation [72-78]. Our data suggests a potential tau loss of function caused by the

V337M mutation at an early timepoint, but we have not functionally validated if the mutation

causes a change in tau function in neurons.

Beyond the findings presented here, we expect that the data sets we have generated will

continue to be useful to the field as we resolve the plethora of molecular and cellular phenotypes

driven by pathogenic tau in a variety of contexts. Similarly, although much is still to be learned

about the consequences of dysregulated tau phosphorylation (both loss and gain), our functional

genomic screens could inform the design of tau phosphorylation modulators, perhaps even for

therapy.

Conclusions

Our study aims to characterize WT, V337M tau and tau knockdown neurons to

understand how tau loss or mutation perturbs neuron biology. We show that V337M tau and tau

knockdown have conserved effects in RNA-seq, ATAC-seq and phosphoproteomics.

Surprisingly, we found that V337M tau causes tau hypophosphorylation. We performed

functional genomics screens to uncover the regulators of tau phosphorylation in WT and V337M

tau neurons. V337M tau perturbs axon morphology pathways similarly to tau loss and causes tau

hypophosphorylation, which could contribute to the previously reported cognitive changes in

preclinical *MAPT* variant carriers.

Materials and Methods

Human iPSC culture and neuronal differentiation (Adapted from Tian et al. 2021)

Human iPSCs from the WTC11 background were cultured in StemFlex Medium

(GIBCO/Thermo Fisher Scientific; Cat. No. A3349401). Human iPSCs from the GIH6C1

background were cultured in mTeSR Plus medium (StemCell Technologies; Cat. No. 100-0276).

iPSCs were grown in plates or dishes coated with Growth Factor Reduced, Phenol Red-Free,

LDEV-Free Matrigel Basement Membrane Matrix (Corning; Cat. No. 356231) diluted 1:100 in

Knockout DMEM (GIBCO/Thermo Fisher Scientific; Cat. No. 10829-018). StemFlex Medium

was replaced daily. When cells reached 80-90% confluency, cells were dissociated with StemPro

Accutase Cell Dissociation Reagent (GIBCO/Thermo Fisher Scientific; Cat. No. A11105-01) at

37ºC for 5 min, centrifuged at 200 g for 5 min, resuspended in StemFlex Medium supplemented

with 10 nM Y-27632 dihydrochloride ROCK inhibitor (Tocris; Cat. No. 125410) and placed

onto Matrigel-coated plates or dishes. Studies at UCSF with human iPSCs were approved by the

Human Gamete, Embryo, and Stem Cell Research (GESCR) Committee.

For individual gene knockdown in CRISPRi iPSCs, sgRNAs were introduced into iPSCs via

lentiviral delivery. Cells were selected by 1 µg/ml puromycin for 2-4 days and recovered for 2-4

days. Phenotypes were evaluated 5-7 days after infection.

The WTC11 CRISPRi iPSC lines were previously engineered to express mNGN2 under a

doxycycline-inducible system in the AAVS1 safe harbor locus. The GIH6C1 iPSC lines were

engineered in this work to express Ngn2 under a doxycycline-inducible promoter in the AAVS1

safe harbor locus. For their neuronal differentiation, we followed our previously described

protocol [79]. Briefly, iPSCs were pre-differentiated in matrigel-coated plates or dishes in N2

Pre-Differentiation Medium containing the following: Knockout DMEM/F12 (GIBCO/Thermo

Fisher Scientific; Cat. No. 12660-012) as the base, 1X MEM Non-Essential Amino Avids

(GIBCO/Thermo Fisher Scientific; Cat. No. 17502-048), 10 ng/mL NT-3 (PeproTech; Cat. No.

450-03), 10 ng/mL BDNF (PeproTech; Cat. No. 450-02), 1µg/mL Mouse Laminin (Thermo

Fisher Scientific; Cat. No. 23017-015), 10 nM ROCK inhibitor and 2µg/mL doxycycline to

induce the expression of Ngn2. After three days, or Day 0, pre-differentiated cells were

dissociated with accutase and plated into BioCoat Poly-D-Lysine-coated plates or dishes

(Corning; assorted Cat. No.) in Classic N2 neuronal medium or BrainPhys Neuronal Medium.

Classic N2 neuronal medium contained the following: half DMEM/F12 (GIBCO/Thermo Fisher

Scientific; Cat. No. 11320-033) and half Neurobasal-A (GIBCO/Thermo Fisher Scientific; Cat.

No. 10888-022) as the base, 1X MEM Non-Essential Amino Acids, 0.5X GlutaMAX

Supplement (GIBCO/Thermo Fisher Scientific; Cat. No. 35050-061), 0.5X N2 Supplement, 0.5X

B27 Supplement (GIBCO/Thermo Fisher Scientific; Cat. No. 17504-044), 10 ng/mL NT-3, 10

ng/mL BDNF and 1 µg/mL Mouse Laminin. BrainPhys Neuronal Medium was comprised of the

following: BrainPhys Neuronal Medium (StemCell Technologies; Cat. No. 05791) as the base,

0.5x N2 Supplement, 0.5X B27 Supplement, 10 ng/mL NT-3, 10ng/mL BDNF, and 1 µg/mL

Mouse Laminin. Neuronal medium was fully changed on day 3 post differentiation and then

half-replaced on day 7 and weekly thereafter.

GIHC1 iPSC cell line generation

GIH6C1 and GIH6C1Δ1E11 were obtained from NeuraCell [23]. iPSCs were transfected with

pC13N-dCas9-BFP-KRAB and TALENS targeting the human CLYBL intragenic safe harbor

locus (between exons 2 and 3) (pZT-C13-R1 and pZT-C13-L1, Addgene #62196, #62197) [80]

using DNA In-Stem (VitaScientific). At the same time, the iPSCs were also transfected with

pUCM-AAVS1-TO_hNGN2 (Addgene #105840) [81] and TALENS targeting the human

AAVS1 intragenic safe harbor locus (pTALdNC□AAVS1_T1, pTALdNC□AAVS1_T2). [82]

After two weeks, BFP-positive iPSCs (CRISPRi+/mNGN2-), mCherry-positive iPSCs

(CRISPRi-/mNGN2+) and BFP/mCherry-positive iPSCs (CRISPRi+/mNGN2+) were isolated

via FACS sorting. Cells were plated sparsely in a 10 cm dish (5,000-10,000 per dish) and

allowed to grow up until they formed large colonies. Homogenous BFP+/mCherry+ colonies

were picked with a pipette tip and placed into a 24 well plate for expansion and characterization.

Cre mRNA was then transfected into the iPSCs to remove the selection marker and mCherry.

Cells were sorted for mCherry negativity, and then mCherry negative colonies were picked and

genotyped.

Western blots

Neurons were washed 3 times with ice-cold PBS. Ice-cold RIPA with protease and phosphatase

inhibitors was added to cells. Lysates were incubated on ice for 2 minutes and then scraped

down. Lysates were centrifuged at 12500xg for 10 minutes at 4 ºC. The supernatants were

collected, and the concentrations were measured with the BCA assay (Thermo Fisher Scientific;

Cat No. 23225). 10-20 µg protein were loaded onto 4-12% Bis-Tris polyacrylamide gel (Thermo

Fisher Scientific; Cat No. NP0336BOX) Nitrocellulose (BioRad, Cat. No. 1620146) or PVDF

membranes were used to transfer the protein in a BioRad Transblot for 11 minutes at 25 V, 25 A.

Membranes were then blocked for 1 hour with Licor Intercept blocking buffer (Licor, Cat. No.

927-60001) at room temperature. Primary antibody was added in Licor Intercept block overnight

at 4 ºC. Blots were washed 3 times for 5 minutes with TBST at room temperature. Secondary

antibodies were added in Licor Intercept block for 1 hour at room temperature. Blots were

washed 3 times for 5 minutes with TBST at room temperature and imaged on a Licor Odyssey.

Immunoblots were quantified by intensity using ImageStudio (Licor).

Bulk RNA sequencing sample preparation

RNA was harvested from day 7, day 14 and day 28 post differentiation neurons using a Zymo

microprep kit (Zymo Research, Cat No. R2062). The library was prepared by first depleting

ribosomal RNA (New England BioLabs, Cat No. E7405L). cDNA synthesis was then performed

on all remaining RNAs (New England BioLabs, Cat. No. E7765S). Paired-end (PE65)

sequencing was performed at the Chan Zuckerberg Biohub and the UCSF Center for Advanced

Technology.

ATAC-seq sample preparation

Omni-ATAC-seq was performed as previously described.[83] In short, nuclei from 50,000

neurons were resuspended with Tn5 transposase (to tag and cleave open chromatin with PCR

adapters) and incubated at 37 C for 30 minutes on a thermomixer at 1,000 rpm. DNA was then

extracted using the Qiagen MinElute Reaction Cleanup Kit (Cat#28204). Tagged sequences were

amplified using Illumina/Nextera i5 common adapter and i7 index adapters. DNA libraries were

purified using AMPure XP beads (A63880), and paired-end (PE65) sequencing was performed at

the Chan Zuckerberg Biohub and the UCSF Center for Advanced Technology.

Proteomics sample preparation

Briefly, neurons were scraped off 15 cm dishes at day 7 of differentiation and flash frozen in

liquid nitrogen. Cell pellet was lysed by adding 1 ml of 6 M GnHCl, 100mM Tris pH 8 and

boiling at 95 C for 5 minutes two times with 5 min rest in between. DNA was sheared three

times via probe sonication at 20% amplitude for 10 s., followed by 10 s of rest. Following

sonication, samples were allowed to solubilize on ice for 20 mins before clearing cell debris by

centrifugation at 16,000 x g for 10 mins and determining protein concentration was using Protein

Thermo Scientific 660 assay. Enough lysate for 1 mg of protein was aliquoted and Tris 2-

carboxyethyl phosphine (TCEP) and chloroacetamide (CAA) were added to each sample to a

final concentration of 40 mM and 10 mM respecitively, before incubating for 10 min at 45 C

with shaking. Guanidine was then diluted at least 1:5 with 100 mM Tris pH 8. Trypsin and LysC

(Promega) were added at a 1:100 (enzyme:protein w:w) ratio (total protease:protein ratio of

1:50) and digested overnight at 37°C with shaking. Following digestion, 10% trifluoroacetic acid

(TFA) was added to each sample to a final pH ~2. Samples were desalted under vacuum using

Sep Pak tC18 cartridges (Waters). Each cartridge was activated with 1 mL 80% acetonitrile

(ACN)/0.1% TFA, then equilibrated with 3×1 mL of 0.1% TFA. Cartridges were then washed

with 4×1 mL of 0.1% TFA, and samples were eluted with 0.8 mL 50% ACN/0.25% formic acid

(FA). 20 μg of each sample was kept for protein abundance measurements, and the remainder

was used for phosphopeptide enrichment. Samples were dried by vacuum centrifugation.

 Phosphopeptide enrichment

For phosphopeptide enrichment of samples for phosphoproteomics, IMAC beads (Fe-IMAC

from Cube Biotech) were prepared by washing 3x with washing buffer (0.1% TFA, 80%

ACN). Dry, digested peptide samples were resuspended in washing buffer and incubated for 15

mins at 37 C with shaking. Peptides were enriched for phosphorylated peptides using a King

Fisher Flex (KFF). A more detailed KFF protocol can be provided. Briefly, after resuspension

peptides were mixed with beads and bound peptides were washed three times with wash buffer

before being eluted from beads using 50% ACN, 2.5 % NH4OH solution. Enriched

phosphorylated peptide samples were acidified using 75% ACN, 10% FA (at a ratio of 5:3

elution buffer:acid buffer), and filtered by centrifugation through NEST tips.

Mass spectrometry data acquisition

Digested samples were analyzed on an Orbitrap Exploris 480 mass spectrometry system (Thermo

Fisher Scientific) equipped with either an Easy nLC 1200 or Neo Vanquish ultra-high pressure

liquid chromatography system (Thermo Fisher Scientific) interfaced via a Nanospray Flex

source. Separation was performed using a 15 cm long PepSep column with a 150 um inner

diameter packed with 1.5um Reprosil C18 particles. Mobile phase A consisted of 0.1% FA, and

mobile phase B consisted of 0.1% FA/80% ACN. Abundance samples were separated by an

organic gradient from 4% to 30% mobile phase B over 62 minutes followed by an increase to

45% B over 10 minutes, then held at 90% B for 8 minutes at a flow rate of 600 nL/minute.

Phosphoproteomics samples were separated by an organic gradient from 2% to 25% mobile

phase B over 62 minutes followed by an increase to 40% B over 10 minutes, then held at 95% B

for 8 minutes at a flow rate of 600 nL/minute. To expand the spectral library, two samples from

each set of replicates was acquired in a data dependent manner. Data dependent analysis (DDA)

was performed by acquiring a full scan over a m/z range of 350-1100 in the Orbitrap at 60,000

resolving power (@200 m/z) with a normalized AGC target of 300%, an RF lens setting of 40%,

and a maximum ion injection time of "Auto". Dynamic exclusion was set to 45 seconds, with a

10 ppm exclusion width setting. Peptides with charge states 2-6 were selected for MS/MS

interrogation using higher energy collisional dissociation (HCD), with 20 MS/MS scans per

cycle. MS/MS scans were analyzed in the Orbitrap using isolation width of 1.6 m/z, normalized

HCD collision energy of 30%, normalized AGC of 200% at a resolving power of 15,000 with a

22 ms maximum ion injection time. Similar settings were used for data dependent analysis of

phosphopeptide-enriched and abundance samples. Data-independent analysis (DIA) was

performed on all samples. An MS scan at 60,000 resolving power over a scan range of 350-1100

m/z, a normalized AGC target of 300%, an RF lens setting of 40%, and the maximum injection

time set to "Auto", followed by DIA scans using 20 m/z isolation windows over 350-1100 m/z

with a 2 m/z overlap at a normalized HCD collision energy of 30%.

Antibodies used in this study

cJun (CST, #9165)

p-cJun (CST, #91952)

Tau13 (Santa Cruz Biotechnology, sc-21796)

AT8 (Invitrogen, MN1020)

AT100 (Invitrogen, MN1060)

AT180 (Invitrogen, MN1040)

Tau pT217 (Invitrogen, 44-744)

Tau pS396 (Invitrogen, 44-752G)

GAPDH (Santa Cruz Biotechnology, sc-47724)

β-Actin (CST, #4967)

Molecular Cloning:

Overexpression constructs were generated using our previously described PSAP expression

vector [84] as a backbone. This vector expressed PSAP fused to a c-terminal mScarlett. We

cloned emGFP-BRD2 into this vector (deleting PSAP-mScarlett) and then used XhoI and AgeI

restriction enzyme sites to clone in 0N3R tau. We then cloned a gene block for ORF-BamHI-

(GS)4-exFlag-T2A-mApple into the vector using the AgeI and EcoRI sites [37]. We then

mutated WT 0N3R tau to V337M and R406W to generate the final overexpression constructs.

CRISPR screening:

45 million iPSCs were infected with lentivirus encoding for the H1 sublibrary (Horlbeck et al

Elife) at an MOI of ~0.3 and selected with 1ug/mL puromycin until 100% BFP positive.

Lentivirus preparation was performed as described

(https://dx.doi.org/10.17504/protocols.io.8dfhs3n, [37], [79]). For CRISPRa screens, TMP was

added at a final concentration of 50uM for all cultures after selection. Cells were then

differentiated and cultured as previously described (dx.doi.org/10.17504/protocols.io.bcrjiv4n,

[79]). Upon differentiation, pre-differentiated cells were plated on three 15cm PDL-coated dishes

at a density of 15 million cells per plate. Neurons were then matured for two weeks. At two

weeks of age, neurons were lifted with papain and zinc fixed as previously described [37]. On

the day of sorting, preparation for FACS was performed as described [37] using the AT8

antibody (Thermo MN1020) at a concentration of 1:200. After sorting, cells were pelleted at

200xg for 20 minutes, the supernatant was removed and the pellet was frozen at -20. Genomic

DNA was extracted with the NucleoSpin Blood L kit. sgRNA cassettes were amplified, pooled,

and sequenced as described [79]. CRISPR screens were analyzed using MAGeCK-iNC as

previously described [79]. Briefly, raw sequencing reads were cropped and aligned using custom

scripts that are publicly available (https://kampmannlab.ucsf.edu/resources). Raw phenotype

scores and p-values were calculated for target genes and negative control genes using a Mann-

Whitney U-test.

Data Analysis

RNA-seq analysis

Sequencing data was aligned to the human reference genome hg38. Rbowtie2 was used to align

and count the number of transcripts from aligned reads. Differentially expressed genes were

determined using DEseq2.

ATAC-seq analysis

Sequencing data was aligned to the human reference genome hg38 using Rbowtie2. Peak calling

was performed with MACS2. Differential ATACseq was performed using DEseq2, and motif

analysis was performed with the motifDB and motifmatchr packages. Differential motif analysis

was performed with the chromVar package.

Gene set enrichment analysis

Enrichr was used to perform gene set enrichment analysis on RNA-seq, proteomics and

phosphoproteomics datasets [85].

Proteomics and Phosphoproteomics Analysis

Raw files were searched using the directDIA+ feature in Spectronaut, with DDA files provided

as supplementary search files against a full human proteome from Uniprot (reviewed entries

only, isoforms included). Phosphosites were extracted from the PTMsites output table from

Spectronaut, and collapsed using the Tukey's median polish functionality of MSstats in R.

Abbreviations

MAPT: Microtubule Associated Protein Tau

iPSC: induced pluripotent stem cell

MAPK: Mitogen-Activated Protein Kinase

MAP2K: Mitogen-Activated Protein Kinase Kinase

ASOs: Anti-sense oligonucleotides

FTD: Frontotemporal Dementia

WT: wild type

MAPT Het: *MAPT* heterozygous (*MAPT* V337M/WT)

MAPT Hom: *MAPT* homozygous (*MAPT* V337M/V337M)

mNGN2: mouse Neurogenin2

AAVS1: adeno-associated virus integration site 1, safe harbor locus.

CLYBL: Citramalyl-CoA Lyase, here refers to an intergenic safe harbor locus.

sgRNA: single guide RNA

NTC: non-targeting control

ANK3: Ankyrin 3

MAPRE3: Microtubule-Associated Protein RP/EB Family Member 3, or EB3

GSK3B: Glycogen Synthase Kinase 3 Beta

CDK5: Cyclin Dependent Kinase 5

CDK5R1: Cyclin Dependent Kinase 5 Regulatory Subunit 1

MARK1: Microtubule Affinity Regulating Kinase 1

Map1a: Microtubule-Associated Protein 1A

Map1b: Microtubule-Associated Protein 1B

LDEV: Lactate Dehydrogenase Elevating Virus

DMEM: Dulbecco's Modified Eagle Medium

ROCK: Rho Associated Coiled-Coil Containing Protein Kinase

BDNF: Brain-Derived Neurotrophic Factor

KRAB: Krüppel-associated box

TALENS: transcription activator-like effector nucleases

PBS: Phosphate-buffered saline

TBST: Tris-buffered saline + 0.1% Tween20

TCEP: Tris 2-carboxyethyl phosphine

CAA: chloroacetamide

TFA: trifluoroacetic acid

ACN: acetonitrile

FA: formic acid

KFF: King Fisher Flex

DDA: Data dependent analysis

AGC: Automated gain control

DIA: Data-independent analysis

MS: Mass spectrometry

Ppm: parts per million

PTM: Post translational modification

MS/MS: Tandem mass spectrometry

HCD: Higher energy collisional dissociation

GAPDH: glyceraldehyde-3-phosphate dehydrogenase

PSAP: Prosaposin

emGFP: emerald green fluorescent protein

BRD2: Bromodomain Containing 2

ORF: open reading frame

TMP: trimethoprim

PDL: poly-D lysine

FACS: fluorescence-activated cell sorting

MAGeCK-iNC: Model-based Analysis of Genome-wide CRISPR/Cas9 Knockout – including

negative controls

Declarations:

Ethics approval and consent to participate

Studies at UCSF with human iPSCs were approved by the Human Gamete, Embryo, and Stem

Cell Research (GESCR) Committee.

Consent for publication

All authors have approved the contents of this study and provided consent for publication.
595

Competing interests

598 M.K. is a co-scientific founder of Montara Therapeutics and serves on the Scientific Advisory

599 Boards of Engine Biosciences, Casma Therapeutics, Alector, and Montara Therapeutics, and is

600 an advisor to Modulo Bio and Theseus Therapeuties. M.K. is an inventor on US Patent

601 11,254,933 related to CRISPRi and CRISPRa screening, and on a US Patent application on *in*

602 *vivo* screening methods.

Funding

M.K. was supported by the Rainwater Charitable Foundation/Tau consortium, the Chan

Zuckerberg Initiative Ben Barres Early Career Acceleration Award, and NIH grants R01

AG062359, R01 AG082141, U54 NS100717. M.K. and D.L.S were supported by U54

NS123746. G.D. was supported by the A.P. Gianni Foundation. A.J.S was supported by NIH F32

AG063487 and NIH K99 AG080116. S.C.B. was supported by the Alzheimer's Association

Research Fellowship 23AARF-1027616.

Author's contributions

613 G.A.M., G.D. and M.K. conceptualized and led the project and wrote the manuscript with input

614 from all co-authors. G.A.M, G.D., E.M., A.J.S., C.P.S, J.J., N.A., A.L., and S.C.B. contributed to

615 making cell lines, performing experiments, and data analysis. J.M. and D.P.S. performed and

616 analyzed the mass spectrometry.

Acknowledgements

We would like to thank all members of the Kampmann lab for helpful feedback and support. We

would like to thank Dr. Ruilin Tian, Dr. Carlo Condello, Dr. Noam Teyssier, Dr. Parker

Grosjean, Dr. Olivia Teter and Ian Steele for helpful discussions and contributions to preliminary

studies. We would like to thank the Tau Consortium Stem Cell Group for sharing the GIH6C1

and GIH6C1Δ1E11 iPSC cell lines and Dr. Li Gan for sharing the WTC11 V337M iPSC cell

lines.

Authors' Information

Gregory A. Mohl and Gary Dixon have contributed equally to this work.

628
Figure 1: RNA-seq and ATAC-seq in neurons reveal conserved effects of *MAPT* V337M

knockdown on axonogenesis pathways (A) iPSCs from a healthy donor (WTC11, here called

MAPT WT) or a patient with the heterozygous *MAPT* V337M mutation (GIH6C1, here called

*MAPT Het) were edited with Cas9 previously to generate a heterozygous *MAPT* V337M clone

(*MAPT* Het), a homozygous *MAPT* V337M clone (*MAPT* Hom) and a healthy isogenic control

(GIH6C1Δ1E11, here called *MAPT WT). These cells were engineered to express dox-inducible

mNGN2 in AAVS1 and CRISPRi machinery in CLYBL. We transduce the iPSCs with lentivirus

for sgRNA/BFP expression. **(B)** Heatmap comparing changes in gene expression based on RNA-

seq in *MAPT* Het NTC and *MAPT* WT MAPT KD vs. *MAPT* WT NTC at 2 and 4 weeks post

differentiation. Three independent wells of neurons for each genotype/sgRNA combination were

harvested at each timepoint. **(C)** Gene Ontology (GO) term enrichment analysis of the RNA-seq

experiment in (B). Genes that are differentially expressed in both *MAPT* Het and *MAPT* WT

MAPT KD vs. *MAPT* WT NTC were analyzed with Enrichr, and top terms with minimal overlap

were plotted. Pathways related to axonogenesis and neuron morphology are colored magenta.

(D) Heatmap summarizing ATAC-seq differential peaks at 4 weeks of differentiation. Two

independent wells of neurons for each genotype/sgRNA combination were harvested. **(E)**

Heatmap summarizing ATAC-seq transcription factor motif analysis at 4 weeks of

differentiation from the same experiment in (D). Clusters were analyzed for pathway enrichment

using Enrichr, and major pathways are annotated ("AP-1 Transcription Factor Network" and

"Neuron Differentiation"). **(F)** Western blot measuring p-cJun and cJun levels in neurons at 1

week of differentiation. Two independent wells of neurons for each genotype/sgRNA

combination were harvested. **(G-H)** Quantification of cJun (G) and p-cJun (H) from the western

blot in (F). Significance was calculated using one-way ANOVA with Dunnett's multiple

comparison test, and comparisons were restricted within the donor background.

Figure 2: Proteomics uncovers altered phosphorylation of axonogenesis-related proteins in neurons with the *MAPT* V337M mutation. (A) Volcano plot showing changes in protein phosphorylation in *MAPT* Hom vs. *MAPT* WT neurons using mass spectrometry. Four independent 150mm dishes of neurons for each condition were harvested after one week of differentiation. Dots represent individual phosphorylation sites. **(B)** Proteins with differential phosphorylation between *MAPT* Het, *MAPT* Hom neurons, and **MAPT* Het vs. controls. Significance was calculated using multiple t-tests adjusted with Šidák single-step correction. Proteins with differential phosphorylation in all three datasets were filtered to identify 56 conserved proteins. **(C)** GO term enrichment of the 56 proteins from (B). Neuron morphology term bars are magenta, and splicing term bars are green. **(D)** Proteins with differential phosphorylation between *MAPT* Het vs. *MAPT* WT and two published mouse phosphoproteomics datasets, including tau KO mice and P301S mice vs. WT mice. Significance was calculated using multiple t-tests adjusted with Šidák single-step correction. Proteins with differential phosphorylation in all three datasets were filtered to identify 45 conserved proteins. **(E)** GO term enrichment of the 45 proteins from (D). Annotations are consistent with (C). **(F)** String-DB protein-protein interaction network of proteins with differential phosphorylation in five datasets: *MAPT* Het, *MAPT* Hom, **MAPT* Het vs isogenic controls, and tau KO mice and P301S mice vs. controls. The inner circle is colored based on the protein $\log_2$ fold change, and the outer circles are colored based on the $\log_2$ fold change for the indicated phosphorylation site. **(G)** Heatmap of phosphoproteomics data comparing *MAPT* KD vs. isogenic controls. Phosphosites that are decreased in *MAPT* Het vs. *MAPT* WT but that are rescued by tau knockdown in *MAPT* Het neurons are labeled in purple as a tau "gain-of-function" signature.

677   Phosphosites that are changed in the same direction in *MAPT* WT *MAPT* KD and *MAPT* Het vs.

678   WT are labeled in orange as a tau "loss-of-function" signature.

Figure 3: Tau phosphorylation is reduced in neurons with the *MAPT* V337M mutation. (A)
Heatmap of tau phosphorylation from *MAPT* Hom or *MAPT* Het vs. *MAPT* WT. Phosphosites
that were not detected in more than half of the replicates in both samples are marked in grey, and
statistically significant phosphorylation changes are marked with an asterisk. **(B)** Protein domain
map of 0N3R tau with detected phosphorylation sites labeled. Decreased phosphorylations
detected in either *MAPT* Hom or *MAPT* Het neurons are labeled in blue. When
phosphoproteomics could not distinguish between multiple potential phosphosites, all are
included. **(C)** Western blot validating decreased tau phosphorylation in neurons with V337M tau.
Two independent wells of neurons were harvested after one week of differentiation. AT8 was
used to label tau pS202/pT205, and Tau13 was used to label total tau. **(D-E)** Quantification of
total tau levels (D) or pS202/pT205 levels (E) from the western blot in (C). One way ANOVA
with Šidák's correction and comparisons within donor backgrounds was used to test for
significance. **(F)** WT, V337M and R406W tau were overexpressed via lentivirus in *MAPT* WT
MAPT KD iPSCs. Three independent wells of neurons were harvested after one week of
differentiation. pTau and total tau levels were analyzed by Western blot. **(G)** Quantification of
the western blot in (F). Band intensities were normalized to actin and to the WT tau
overexpression line. Significance was calculated using two-way ANOVA with Dunnet's multiple
comparisons test. **(H)** Kinase activity analysis from phosphorylation changes in neurons after
one week of differentiation with the homozygous *MAPT* V337M mutation (*MAPT* Hom) vs.
isogenic controls (*MAPT* WT). The $\log_2$ fold change of phosphopeptide abundance for annotated
kinase substrates is plotted. The range is represented by the thin lines, the box represents the
IQR, and the median is represented by a thick line. **(I)** Heatmap for kinase activity scores from
all five phosphoproteomic datasets vs. isogenic controls.

Figure 4: CRISPR screens elucidate regulators of tau phosphorylation in neurons. (A)

Pooled genetic screening workflow for tau pS202/pT205 levels in neurons. *MAPT* WT CRISPRi

NGN2 iPSCs, *MAPT* Het CRISPRi NGN2 iPSCs or *MAPT* WT CRISPRa NGN2 iPSCs were

transduced by lentivirus with a pooled "druggable genome" sgRNA library targeting 2,318 genes

enriched for kinases and phosphatases. iPSCs were differentiated into neurons. After two weeks,

neurons were fixed, stained with AT8 and sorted for high or low AT8 staining. The high AT8

and low AT8 samples were sequenced to determine which sgRNAs were enriched in either

fraction. **(B)** Scatter plot comparing CRISPRi screens in *MAPT* Het neurons vs. *MAPT* WT

neurons. AT8 non-hits are labeled with grey circles, AT8 hits that also modify tau levels (using

the T22 antibody as a surrogate for total tau levels) are labeled with "+", and AT8-specific hits

are labeled with black circles. Top genotype-specific hits are labeled in black, and key pathways

are labeled in green (cytoskeleton), purple (Neuron projection development) and blue (p38

MAPK pathway). **(C)** Rank plot showing the results of the CRISPRa screen in *MAPT* WT

neurons. AT8 non-hits are labeled in grey and AT8 hits are labeled in black. Hits that are

cytoskeleton-related genes are labeled in green and hits that are related to neuron projection

development are labeled in purple. **(D)** Detected tau phosphosites are mapped to the phenotype

of known tau kinases from the CRISPRi/a screens. Tau phosphosites that were significantly

different in either *MAPT* Hom or *MAPT* Het neurons vs. *MAPT* WT are indicated with blue

text/boxes. Kinases whose knockdown or overexpression perturb tau phosphorylation at

S202/T205 are indicated by red text/boxes. Overlap between significant kinases and differential

phosphorylations are indicated by purple boxes. Grey boxes indicate phosphorylations by known

tau kinases that are not significantly differential or involved in tau pS202/T205. References for

known tau kinase activity are indicated by the kinase name.

References

1.   Alzheimer, A., *Uber eigenartige Erkrankung der Hirnrinde.* All Z Psychiatr, 1907.
64: p. 146–148.
2.   Malpetti, M., R. Joie, and G.D. Rabinovici, *Tau Beats Amyloid in Predicting Brain*
Atrophy in Alzheimer Disease: Implications for Prognosis and Clinical Trials. J
Nucl Med, 2022. **63**(6): p. 830–832.
3.   Wu, L., et al., *Early-onset familial Alzheimer's disease (EOFAD).* Can J Neurol
Sci, 2012. **39**(4): p. 436–45.
4.   Wang, Y. and E. Mandelkow, *Tau in physiology and pathology.* Nat Rev Neurosci,
2016. **17**(1): p. 5–21.
5.   Silva, M.C., et al., *Human iPSC-Derived Neuronal Model of Tau-A152T*
Frontotemporal Dementia Reveals Tau-Mediated Mechanisms of Neuronal
Vulnerability. Stem Cell Reports, 2016. **7**(3): p. 325–340.
6.   Ash, P.E.A., et al., *TIA1 potentiates tau phase separation and promotes*
generation of toxic oligomeric tau. Proc Natl Acad Sci U S A, 2021. **118**(9).
7.   Jiang, L., et al., *Interaction of tau with HNRNPA2B1 and N(6)-methyladenosine*
RNA mediates the progression of tauopathy. Mol Cell, 2021. **81**(20): p. 4209–
4227 e12.
8.   Lester, E., et al., *Tau aggregates are RNA-protein assemblies that mislocalize*
multiple nuclear speckle components. Neuron, 2021. **109**(10): p. 1675–1691 e9.
9.   Bowles, K.R., et al., *ELAVL4, splicing, and glutamatergic dysfunction precede*
neuron loss in MAPT mutation cerebral organoids. Cell, 2021. **184**(17): p. 4547–
4563 e17.
10.   Dickson, J.R., M.P. Frosch, and B.T. Hyman, *Altered localization of nucleoporin*
98 in primary tauopathies. Brain Commun, 2023. **5**(1): p. fcac334.
11.   Frost, B., F.H. Bardai, and M.B. Feany, *Lamin Dysfunction Mediates*
Neurodegeneration in Tauopathies. Curr Biol, 2016. **26**(1): p. 129–36.
12.   Paonessa, F., et al., *Microtubules Deform the Nuclear Membrane and Disrupt*
Nucleocytoplasmic Transport in Tau-Mediated Frontotemporal Dementia. Cell
Rep, 2019. **26**(3): p. 582–593 e5.
13.   Stamer, K., et al., *Tau blocks traffic of organelles, neurofilaments, and APP*
vesicles in neurons and enhances oxidative stress. J Cell Biol, 2002. **156**(6): p.
1051–63.
14.   Vossel, K.A., et al., *Tau reduction prevents Abeta-induced axonal transport*
deficits by blocking activation of GSK3beta. J Cell Biol, 2015. **209**(3): p. 419–33.
15.   DuBoff, B., J. Gotz, and M.B. Feany, *Tau promotes neurodegeneration via DRP1*
mislocalization in vivo. Neuron, 2012. **75**(4): p. 618–32.
16.   Caballero, B., et al., *Acetylated tau inhibits chaperone-mediated autophagy and*
promotes tau pathology propagation in mice. Nat Commun, 2021. **12**(1): p. 2238.
17.   Sohn, P.D., et al., *Pathogenic Tau Impairs Axon Initial Segment Plasticity and*
Excitability Homeostasis. Neuron, 2019. **104**(3): p. 458–470 e5.

18.   Chang, C.W., et al., *Tau reduction affects excitatory and inhibitory neurons*
differently, reduces excitation/inhibition ratios, and counteracts network
hypersynchrony. Cell Rep, 2021. **37**(3): p. 109855.

19.   Ittner, L.M., et al., *Dendritic function of tau mediates amyloid-beta toxicity in*
Alzheimer's disease mouse models. Cell, 2010. **142**(3): p. 387–97.

20.   Miyamoto, T., et al., *Phosphorylation of tau at Y18, but not tau-fyn binding, is*
required for tau to modulate NMDA receptor-dependent excitotoxicity in primary
neuronal culture. Mol Neurodegener, 2017. **12**(1): p. 41.

21.   Roberson, E.D., et al., *Reducing endogenous tau ameliorates amyloid beta-*
induced deficits in an Alzheimer's disease mouse model. Science, 2007.
316(5825): p. 750–4.

22.   Chang, C.W., E. Shao, and L. Mucke, *Tau: Enabler of diverse brain disorders and*
target of rapidly evolving therapeutic strategies. Science, 2021. **371**(6532).

23.   Karch, C.M., et al., *A Comprehensive Resource for Induced Pluripotent Stem*
Cells from Patients with Primary Tauopathies. Stem Cell Reports, 2019. **13**(5): p.
939–955.

24.   Andres-Benito, P., et al., *Deregulated Transcription and Proteostasis in Adult*
mapt Knockout Mouse. Int J Mol Sci, 2023. **24**(7).

25.   Ferrer, I., et al., *Dysregulated Protein Phosphorylation in a Mouse Model of*
FTLD-Tau. J Neuropathol Exp Neurol, 2022. **81**(9): p. 696–706.

26.   Miyaoka, Y., et al., *Isolation of single-base genome-edited human iPS cells*
without antibiotic selection. Nat Methods, 2014. **11**(3): p. 291–3.

27.   Dujardin, S., et al., *Tau molecular diversity contributes to clinical heterogeneity in*
Alzheimer's disease. Nat Med, 2020. **26**(8): p. 1256–1263.

28.   Wesseling, H., et al., *Tau PTM Profiles Identify Patient Heterogeneity and Stages*
of Alzheimer's Disease. Cell, 2020. **183**(6): p. 1699–1713 e13.

29.   Wegmann, S., J. Biernat, and E. Mandelkow, *A current view on Tau protein*
phosphorylation in Alzheimer's disease. Curr Opin Neurobiol, 2021. **69**: p. 131–
138.

30.   Sergeant, N., et al., *Biochemistry of Tau in Alzheimer's disease and related*
neurological disorders. Expert Rev Proteomics, 2008. **5**(2): p. 207–24.

31.   Yoshimura, Y., T. Ichinose, and T. Yamauchi, *Phosphorylation of tau protein to*
sites found in Alzheimer's disease brain is catalyzed by Ca2+/calmodulin-
dependent protein kinase II as demonstrated tandem mass spectrometry.
Neurosci Lett, 2003. **353**(3): p. 185–8.

32.   Derkinderen, P., et al., *Tyrosine 394 is phosphorylated in Alzheimer's paired*
helical filament tau and in fetal tau with c-Abl as the candidate tyrosine kinase. J
Neurosci, 2005. **25**(28): p. 6584–93.

33.   Scales, T.M., et al., *Tyrosine phosphorylation of tau by the SRC family kinases*
lck and fyn. Mol Neurodegener, 2011. **6**: p. 12.

34.   Abreha, M.H., et al., *TBK1 interacts with tau and enhances neurodegeneration in*
tauopathy. J Biol Chem, 2021. **296**: p. 100760.

35.   Roth, A., et al., *Comprehensive Characterization of CK1delta-Mediated Tau*
Phosphorylation in Alzheimer's Disease. Front Mol Biosci, 2022. **9**: p. 872171.

36.   Horlbeck, M.A., et al., *Compact and highly active next-generation libraries for*
CRISPR-mediated gene repression and activation. Elife, 2016. **5**.

37.   Samelson, A.J., et al., *CRISPR screens in iPSC-derived neurons reveal*
principles of tau proteostasis. bioRxiv, 2023.

38.   Nishimura, I., Y. Yang, and B. Lu, *PAR-1 kinase plays an initiator role in a*
temporally ordered phosphorylation process that confers tau toxicity in
Drosophila. Cell, 2004. **116**(5): p. 671–82.

39.   Caceres, A., S. Potrebic, and K.S. Kosik, *The effect of tau antisense*
oligonucleotides on neurite formation of cultured cerebellar macroneurons. J
Neurosci, 1991. **11**(6): p. 1515–23.

40.   Caceres, A. and K.S. Kosik, *Inhibition of neurite polarity by tau antisense*
oligonucleotides in primary cerebellar neurons. Nature, 1990. **343**(6257): p. 461–
3.

41.   Morris, S.L. and S.T. Brady, *Tau phosphorylation and PAD exposure in regulation*
of axonal growth. Front Cell Dev Biol, 2022. **10**: p. 1023418.

42.   Buchholz, S., et al., *The tau isoform 1N4R confers vulnerability of MAPT*
knockout human iPSC-derived neurons to amyloid beta and phosphorylated tau-
induced neuronal dysfunction. Alzheimers Dement, 2025: p. e14403.

43.   Harada, A., et al., *Altered microtubule organization in small-calibre axons of mice*
lacking tau protein. Nature, 1994. **369**(6480): p. 488–91.

44.   Takei, Y., et al., *Defects in axonal elongation and neuronal migration in mice with*
disrupted tau and map1b genes. J Cell Biol, 2000. **150**(5): p. 989–1000.

45.   Dawson, H.N., et al., *Inhibition of neuronal maturation in primary hippocampal*
neurons from tau deficient mice. J Cell Sci, 2001. **114**(Pt 6): p. 1179–87.

46.   Biswas, S. and K. Kalil, *The Microtubule-Associated Protein Tau Mediates the*
Organization of Microtubules and Their Dynamic Exploration of Actin-Rich
Lamellipodia and Filopodia of Cortical Growth Cones. J Neurosci, 2018. **38**(2): p.
291–307.

47.   Padmanabhan, P., et al., *Frontotemporal dementia mutant Tau promotes*
aberrant Fyn nanoclustering in hippocampal dendritic spines. Elife, 2019. **8**.

48.   Tan, D.C.S., et al., *Generation of a New Tau Knockout (tauDeltaex1) Line Using*
CRISPR/Cas9 Genome Editing in Mice. J Alzheimers Dis, 2018. **62**(2): p. 571–
578.

49.   Li, Z., et al., *Seizure resistance without parkinsonism in aged mice after tau*
reduction. Neurobiol Aging, 2014. **35**(11): p. 2617–2624.

50.   Gheyara, A.L., et al., *Tau reduction prevents disease in a mouse model of Dravet*
syndrome. Ann Neurol, 2014. **76**(3): p. 443–56.

51.   DeVos, S.L., et al., *Antisense reduction of tau in adult mice protects against*
seizures. J Neurosci, 2013. **33**(31): p. 12887–97.

52.   Velazquez, R., et al., *Acute tau knockdown in the hippocampus of adult mice*
causes learning and memory deficits. Aging Cell, 2018. **17**(4): p. e12775.

53.   Oberrauch, S., et al., *Reward motivation and cognitive flexibility in tau null-*
mutation mice. Neurobiol Aging, 2021. **100**: p. 106–117.

54.   Ikegami, S., A. Harada, and N. Hirokawa, *Muscle weakness, hyperactivity, and*
impairment in fear conditioning in tau-deficient mice. Neurosci Lett, 2000. **279**(3):
p. 129–32.

55.   Goncalves, R.A., et al., *Behavioral Abnormalities in Knockout and Humanized*
Tau Mice. Front Endocrinol (Lausanne), 2020. **11**: p. 124.

56.   Cheng, J.S., et al., *Tau reduction diminishes spatial learning and memory deficits*
after mild repetitive traumatic brain injury in mice. PLoS One, 2014. **9**(12): p.
e115765.

57.   Ahmed, T., et al., *Cognition and hippocampal synaptic plasticity in mice with a*
homozygous tau deletion. Neurobiol Aging, 2014. **35**(11): p. 2474–2478.

58.   Morris, M., et al., *Age-appropriate cognition and subtle dopamine-independent*
motor deficits in aged tau knockout mice. Neurobiol Aging, 2013. **34**(6): p. 1523–
9.

59.   Lei, P., et al., *Motor and cognitive deficits in aged tau knockout mice in two*
background strains. Mol Neurodegener, 2014. **9**: p. 29.

60.   Kimura, T., et al., *Microtubule-associated protein tau is essential for long-term*
depression in the hippocampus. Philos Trans R Soc Lond B Biol Sci, 2014.
369(1633): p. 20130144.

61.   Regan, P., et al., *Tau phosphorylation at serine 396 residue is required for*
hippocampal LTD. J Neurosci, 2015. **35**(12): p. 4804–12.

62.   Boekhoorn, K., et al., *Improved long-term potentiation and memory in young tau-*
P301L transgenic mice before onset of hyperphosphorylation and tauopathy. J
Neurosci, 2006. **26**(13): p. 3514–23.

63.   Blauwendraat, C., et al., *Parkinson's disease age at onset genome-wide*
association study: Defining heritability, genetic loci, and alpha-synuclein
mechanisms. Mov Disord, 2019. **34**(6): p. 866–875.

64.   Ye, H., et al., *Genetics and Pathogenesis of Parkinson's Syndrome.* Annu Rev
Pathol, 2023. **18**: p. 95–121.

65.   Finger, E., et al., *Neurodevelopmental effects of genetic frontotemporal dementia*
in young adult mutation carriers. Brain, 2023. **146**(5): p. 2120–2131.

66.   Geschwind, D.H., et al., *Dementia and neurodevelopmental predisposition:*
cognitive dysfunction in presymptomatic subjects precedes dementia by decades
in frontotemporal dementia. Ann Neurol, 2001. **50**(6): p. 741–6.

67.   Nakamura, M., et al., *Pathological Progression Induced by the Frontotemporal*
Dementia-Associated R406W Tau Mutation in Patient-Derived iPSCs. Stem Cell
Reports, 2019. **13**(4): p. 684–699.

68.   Brion, J.P., et al., *Developmental changes in tau phosphorylation: fetal tau is*
transiently phosphorylated in a manner similar to paired helical filament-tau
characteristic of Alzheimer's disease. J Neurochem, 1993. **61**(6): p. 2071–80.

69.   Goedert, M., et al., *The abnormal phosphorylation of tau protein at Ser-202 in*
Alzheimer disease recapitulates phosphorylation during development. Proc Natl
Acad Sci U S A, 1993. **90**(11): p. 5066–70.

70.   Naseri, N.N., et al., *Specific Combinations of Physiological Tau Phosphorylation*
Regulate Tau-Microtubule Interactions in Developing Neurons. bioRxiv, 2025.

71.   Ehrlich, M., et al., *Distinct Neurodegenerative Changes in an Induced Pluripotent*
Stem Cell Model of Frontotemporal Dementia Linked to Mutant TAU Protein.
Stem Cell Reports, 2015. **5**(1): p. 83–96.

72.   Mutreja, Y., B. Combs, and T.C. Gamblin, *FTDP-17 Mutations Alter the*
Aggregation and Microtubule Stabilization Propensity of Tau in an Isoform-
Specific Fashion. Biochemistry, 2019. **58**(6): p. 742–754.

73.   Hong, M., et al., *Mutation-specific functional impairments in distinct tau isoforms*
of hereditary FTDP-17. Science, 1998. **282**(5395): p. 1914–7.
74.   Hasegawa, M., M.J. Smith, and M. Goedert, *Tau proteins with FTDP-17*
mutations have a reduced ability to promote microtubule assembly. FEBS Lett,
1998. **437**(3): p. 207–10.
75.   DeTure, M., et al., *Missense tau mutations identified in FTDP-17 have a small*
effect on tau-microtubule interactions. Brain Res, 2000. **853**(1): p. 5–14.
76.   Combs, B. and T.C. Gamblin, *FTDP-17 tau mutations induce distinct effects on*
aggregation and microtubule interactions. Biochemistry, 2012. **51**(43): p. 8597–
607.
77.   Combs, B., K. Voss, and T.C. Gamblin, *Pseudohyperphosphorylation has*
differential effects on polymerization and function of tau isoforms. Biochemistry,
2011. **50**(44): p. 9446–56.
78.   Barghorn, S., et al., *Structure, microtubule interactions, and paired helical*
filament aggregation by tau mutants of frontotemporal dementias. Biochemistry,
2000. **39**(38): p. 11714–21.
79.   Tian, R., et al., *CRISPR Interference-Based Platform for Multimodal Genetic*
Screens in Human iPSC-Derived Neurons. Neuron, 2019. **104**(2): p. 239–255
e12.
80.   Cerbini, T., et al., *Transcription activator-like effector nuclease (TALEN)-mediated*
CLYBL targeting enables enhanced transgene expression and one-step
generation of dual reporter human induced pluripotent stem cell (iPSC) and
neural stem cell (NSC) lines. PLoS One, 2015. **10**(1): p. e0116032.
81.   Fernandopulle, M.S., et al., *Transcription Factor-Mediated Differentiation of*
Human iPSCs into Neurons. Curr Protoc Cell Biol, 2018. **79**(1): p. e51.
82.   Oceguera-Yanez, F., et al., *Engineering the AAVS1 locus for consistent and*
scalable transgene expression in human iPSCs and their differentiated
derivatives. Methods, 2016. **101**: p. 43–55.
83.   Corces, M.R., et al., *An improved ATAC-seq protocol reduces background and*
enables interrogation of frozen tissues. Nat Methods, 2017. **14**(10): p. 959–962.
84.   Tian, R., et al., *Genome-wide CRISPRi/a screens in human neurons link*
lysosomal failure to ferroptosis. Nat Neurosci, 2021. **24**(7): p. 1020–1034.
85.   Chen, E.Y., et al., *Enrichr: interactive and collaborative HTML5 gene list*
enrichment analysis tool. BMC Bioinformatics, 2013. **14**: p. 128.

Figure S1: V337M tau and tau knockdown perturb gene expression of axonogenesis-related genes. (A) Normalized RNA counts of *MAPT* from the RNA-seq experiment described in Figure 1B showing tau knockdown in *MAPT* WT and *MAPT* Het neurons. **(B)** Western blot measuring tau knockdown in *MAPT* WT and *MAPT* Het neurons. Two replicates (individual wells) of neurons were harvested after two weeks of differentiation. **(C)** Quantification of the western blot in (B). **(D)** Bar plot showing the number of differentially expressed genes due to *MAPT* KD in either *MAPT* WT or *MAPT* Het neurons. **(E)** Heatmap of RNA-seq from *MAPT* Het, *MAPT* Hom and *MAPT Het neurons vs. isogenic controls at 1 week or 2 weeks of differentiation.

Differentially expressed genes related to axon guidance or axonogenesis are labeled. **(F)** GO

term enrichment analysis of one-week neurons from the RNA-seq experiment in (A). Genes that

are differentially expressed in both *MAPT* Het and *MAPT* Hom vs. *MAPT* WT were analyzed

with Enrichr, and top terms with minimal overlap were plotted. Pathways related to

axonogenesis and neuron morphology are colored magenta. (G) GO term enrichment analysis of

two-week old neurons from the RNA-seq experiment in (A). Genes that are differentially

expressed in both *MAPT* Het, *MAPT* Hom and **MAPT* Het vs. their isogenic *MAPT* WT controls

were analyzed with Enrichr, and top terms with minimal overlap were plotted. Pathways related

to axonogenesis and neuron morphology are colored magenta.

Figure S2: V337M tau and tau knockdown perturb chromatin accessibility of AP-1

transcription factor network motifs. (A) PCA plot of ATAC-seq differential peaks at 2 and 4

weeks of differentiation. Two replicates (individual wells) of neurons were harvested at each

timepoint. **(B)** GO term enrichment analysis using Cellular Component on genes in 2-week

neurons with differential ATAC-seq peaks in both *MAPT* WT *MAPT* KD and *MAPT* Het NTC

vs. *MAPT* WT NTC. Non-significant terms are labeled in grey. **(C)** Heatmap showing the

relative deviation of transcription factor motifs with significantly different accessibility in *MAPT*

979   WT and *MAPT* Het neurons +/- tau knockdown. Two replicates (individual wells) of neurons

980   were harvested at two weeks of differentiation. GO term enrichment analysis was used on

981   clusters of transcription factors to categorize clusters.

Figure S3: V337M tau and tau knockdown cause phosphorylation changes in axonogenesis

and splicing proteins. (A) (*Top*) Overlap between differential phosphosites in neurons derived

from iPSCs edited to introduce the homozygous *MAPT* V337M mutation (*MAPT* Hom)vs.

isogenic controls (*MAPT* WT). Four replicates (independent 150mm dishes) of neurons for each

genotype/sgRNA combination were harvested after one week of differentiation, and the

phosphoproteome was measured using mass spectrometry. Significance was calculated using

Fisher's Exact Test. (*Bottom*) Overlap between proteins with differential phosphorylation in both

datasets. Significance was calculated using Fisher's Exact Test. **(B)** Overlap between proteomic

changes in *MAPT* Hom neurons, neurons derived from iPSCs edited to have the heterozygous

MAPT V337M mutation (*MAPT* Het) and neurons derived from patient iPSCs with the

heterozygous *MAPT* V337M mutation (**MAPT* Het) vs. isogenic controls (*MAPT* WT or **MAPT*

WT). Four replicates (independent 150mm dishes) of neurons for each genotype/sgRNA

combination were harvested after one week of differentiation, and the total proteome was

measured using mass spectrometry. Significance was calculated using multiple t-tests adjusted

with Šidák single-step correction. Significantly differential proteins in all three datasets were

filtered to identify 145 conserved proteins. **(C)** GO term enrichment of the 145 proteins with

differential abundance in *MAPT* Hom, *MAPT* Het and **MAPT* Het neurons compared to isogenic

controls. Top terms with minimal overlap are shown. Term names are colored to match relevant

gene names in the heatmap in (C). **(D)** Heatmap showing the Log_2 fold change of protein

abundance for the 145 proteins with differential abundance in *MAPT* Hom, *MAPT* Het and

**MAPT* Het neurons vs. isogenic *MAPT* WT neurons. Proteins within enriched GO terms are

labeled and colored according to the shared pathways. **(E)** GO term analysis of phospoproteins

with differential phosphorylation in *MAPT* Het NTC and *MAPT* WT *MAPT* KD vs. *MAPT* WT

NTC. Non-significant terms are labeled by grey bars. Regulation of Microtubule-based process is

labeled by a magenta bar due to its overlap with axon-related terms. **(F)** GO term analysis of

phosphoproteins with differential phosphorylation in *MAPT* Het NTC vs. *MAPT* WT NTC and

MAPT Het *MAPT* KD vs. *MAPT* Het NTC. Terms related to RNA processing and splicing are

marked by green bars.

1016

Figure S4: Neurons with V337M tau have decreased tau phosphorylation at disease-

associated phosphorylation sites. (A) Protein domain map of 2N4R tau. Phosphosites detected

in this study are labeled, with disease-associated phosphorylation sites from AD labeled in

orange. Phosphosites not detected in this study are marked with a small black line and are

unlabeled. Domain abbreviations are as follows: N-terminal inserts (N1,N2), proline rich domain

(PRD), microtubule binding repeats (R1, R2, R3, R4). **(B)** Consensus sequences for tau kinases.

Detected tau phosphosites are shown with their sequence context. Phosphorylation sites that are

differential between either *MAPT* V337M heterozygous (*MAPT* Het) or *MAPT* V337M

homozygous (*MAPT* Hom) are labeled blue with an asterisk, and detected phosphosites are

labeled with an open circle. Kinase consensus sequences are annotated with colored boxes, with

priming sites marked with a "P" in an orange circle. V337 is labeled with a bold/underlined red

V. The domains abbreviated as follows: N-terminal projection domain (NTD), proline rich

domain (PRD), Microtubule binding repeats (R1, R3, R4), C-terminal domain (CTD).

Figure S5: Functional genomics uncovers regulators of tau phosphorylation in *MAPT* WT

and *MAPT* V337M neurons. (A-D) Bar plots showing the median intensity of AT8 (A), AT180

(B), AT100 (C) and pS396 (D) in 2-week *MAPT* WT, *MAPT* KD and *MAPT* Het neurons. AT8

was selected for CRISPR screening due to high reproducibility across timepoints and AT8

detection in both *MAPT* WT and *MAPT* Het neurons at 2 weeks of differentiaton. **(E)** Heatmap

of hits from the CRISPRi and CRISPRa AT8 screens and the CRISPRi T22 screen. Many of the

AT8 hits from the three screens do not modify T22 levels and are therefore unlikely to be due to

modifying tau levels [37]. Genes related to cytoskeleton, neuron projection development or the

p38 MAPK pathway are annotated. **(F)** Heatmap of AT8 and T22 screens with the kinases

predicted to have differential activity in Figure 3I. Selected kinases predicted to have differential

activity in *MAPT* V337M neurons with particular disease relevance that did not have a

phenotype in the AT8 screens are highlighted with grey boxes. *MARK1* is annotated with a

yellow box.

1044

cJun - 800 channel

GAPDH - 700 channel

p-cJun - 800 channel

1045
1046
1047   **Source data for Fig. 1F**
1048
1049

tau13

tau pS202/pT205 (AT8)

Actin

1050

1051 **Source data for Fig. 3C**

1052

1053

1054 **Source data for Fig. 3F**