Title

From Soil to Surface: Green Infrastructure Enhances Microbial Communities in the Built Environment

Authors

Malin Mcgonigal[1,2] and Kohei Ito [1*]

Affiliation

1.　BIOTA Inc., Tokyo, 101-0022, Japan

2.　Department of Biosciences, Durham University, DH1 3LE, England

*Corresponding author

e-mail: kohei@biota.ne.jp

Abstract

Background: High microbial diversity offers extensive benefits to both the environment and human health,
contributing to ecosystem stability, nutrient cycling, and pathogen suppression. In built environments, factors
such as building design, human activity, and cleaning protocols influence microbial communities. This study
investigates the impact of landscape design on microbial diversity and function within the "Visionary Lab"
exhibition at the National Museum of Emerging Science and Innovation (Miraikan) in Tokyo, Japan, using 16s
rRNA amplicon sequencing and shallow shotgun sequencing.

Results: The study found that the Visionary Lab samples exhibited higher microbial diversity compared to other
museum areas, as indicated by alpha diversity metrics. Beta diversity analysis revealed distinct microbial
community structures correlated with sampling locations. Despite this, no consistent patterns were observed in
virulence factors or antimicrobial resistance genes across the samples. Metabolic function analysis showed
varied profiles, suggesting diverse ecological interactions influenced by the curated landscape.

Conclusions: The curated landscape design enhanced microbial diversity, highlighting its potential to create
healthier and more sustainable built environments. However, the lack of consistent patterns in virulence factors
and antimicrobial resistance genes underscores the complexity of microbial community dynamics. Future
research should increase sample size, employ advanced sequencing techniques, and explore various built
environments to develop evidence-based guidelines for microbial-friendly and health-promoting indoor spaces.

Keywords

Landscape design, Metagenomics, Microbiome of the built environment, Microbial diversity, Sustainability

Background

The benefits of high microbial diversity are extensive [1, 2]. In the environment, diverse microbial

communities contribute to the breakdown of organic matter and nutrient cycling, thereby maintaining ecosystem

stability and productivity [3]. Soils with greater biodiversity levels are more resistant to environmental

disturbances and more resilient than those with reduced biodiversity, allowing ecosystems to better withstand

and recover from disturbances such as pollution, climate change and habitat destruction [4]. Moreover, a

complex soil microbial community increases competition for nutrients, inhibiting the development or

persistence of pathogens in the soil. Thus, soil microbial diversity can result in direct benefits for human health

by suppressing disease-causing organisms, providing clean air, water, and food, and serving as a source of

producing antibiotics [4]. In human health, it has been reported that 'a healthy microbiota community often

demonstrated high taxonomic diversity, high microbial gene richness and stable core microbiota' [5]. Such

healthy microbiota is associated with improved immune function, reduced inflammation, improved energy and

nutrient extraction, and a lower incidence of allergies and autoimmune diseases [5].

In built environments, such as homes, offices, and public spaces, microbial communities are influenced

by multiple factors, including building design, ventilation systems, occupancy and human activity, types of

construction materials, cleaning protocols, and external environmental conditions [6–8]. These elements interact

to shape the microbial communities found in indoor spaces, affecting everything from air quality to human

health. Understanding how these factors influence microbial diversity is crucial for designing healthier and more

sustainable built environments [6]. The design and integration of landscape elements within built environments

is one factor that can significantly influence microbial communities. Plants, in particular, have been shown to

enhance microbial abundance and diversity within indoor environments [9, 10]. However, the extent to which

plants influence taxonomic diversity, including genes associated with various metabolic functions or

antimicrobial resistance, remains inadequately explored.

From April 2022 until August 2023, the National Museum of Emerging Science and Innovation

('Miraikan') in Tokyo, Japan, held an exhibition titled "Visionary Lab; Microbes Actually Are All Around". The

exhibition featured a full-scale futuristic living space with an adjoining planting area of purposefully curated

floral and shrubbery arrangements. It was designed to encourage visitors to contemplate future lifestyles where

humans and microbes exist plentifully in harmony. Miraikan's planting area integrates two key concepts:

multispecies sustainability and human augmentation of ecosystems. Multispecies sustainability emphasises the interconnectedness of humans, plants, and microbes, advocating for practices that support the health and well-being of all species [11]. Human augmentation of ecosystems involves the deliberate enhancement of natural environments to improve ecosystem services and resilience [12].

The Visionary Lab, with its landscape design, serves as an ideal setting for investigating the effects of these concepts. This study aimed to elucidate the effects of the Visionary Lab's landscape design by comparing microbial samples from within the exhibition space to those from areas of the Miraikan museum, using 16S rRNA as well as shallow shotgun sequencing to supplement our understanding of the different microbiomes. Our objectives were to ascertain the specific effects of controlled landscape design on the microbial ecosystem. It is hypothesised that the specific landscape design of the Visionary Lab, particularly through the strategic integration of plant life and landscape elements, cultivates a unique and rich microbial ecosystem that markedly contrasts with other areas within the museum. This study is both useful and necessary in contributing to informing urban design and public health strategies. The findings from this research may guide the development of built environments that enhance microbial diversity, contributing to improved air quality, reduced pathogen transmission, and overall public health. This approach aligns with the broader goal of sustainable urban design, which seeks to create living spaces that are resilient, ecologically balanced, and conducive to human well-being.

Methods

Study design, sample collection, and DNA extraction

The study was conducted at the National Museum of Emerging Science and Innovation ('Miraikan'), located in 2--3-6 Aomi, Koto-ku, Tokyo 135-0064. Microbial samples were collected via vacuum collection or surface swabbing from different surfaces and locations, as listed in Table 1, and mapped in Figure 1. For vacuum collection, a Makita vacuum cleaner (CL107FDSH) equipped with a DUSTREAM Collector (DU-ST-1) and DUSTREAM Filter (DU-FL-1) from Indoor Biotechnologies was used. For surface swabbing, a sterile cotton-tipped swab (ESwab™; Copan Diagnostics, Brescia, Italy) was used to swab surfaces for 3 minutes, before they were stored in tubes containing Liquid Amies Medium solution. Care was taken to ensure that collection methods did not contaminate or alter the microbial communities present. The samples were promptly sealed, stored at low temperatures, and refrigerated until DNA extraction. Extraction was performed

89 on the 9 samples using the DNeasy® PowerSoil® Pro Kit (QIAGEN, Germany) according to the manufacturer's

90 protocol.

Library preparation and sequencing

For the 16S rRNA amplicon sequencing, sequencing libraries were generated, and indexes were

incorporated to attribute sequences to individual samples. PCR amplification of the targeted V3-V4 regions was

performed using specific primers, CCTAYGGGRBGCASCAG and GGACTACNNGGGTATCTAAT. PCR

products with proper size were selected via 2% agarose gel electrophoresis. An equal amount of PCR products

from each sample was pooled, end-repaired, A-tailed, and then ligated with Illumina adapters. The libraries were

sequenced on a paired-end Illumina platform, producing 250 base pair paired-end raw reads. Library quality was

evaluated and quantified using qPCR. Quantified libraries were then pooled and sequenced on Illumina

platforms according to the effective library concentration and the data amount required. DNA samples, library

preparation, and amplicon sequencing were performed using 250 base paired-end sequencing on the Illumina

NovaSeq 6000 system at Novogene (Beijing, China).

For the shallow shotgun metagenomic sequencing, sequencing libraries were prepared, and quality

control (QC) was performed at each step to ensure data reliability. The DNA samples were randomly

fragmented into shorter sequences, which were end-repaired, A-tailed, and ligated with Illumina adapters. These

adapter-linked fragments were size-selected, PCR-amplified (or left PCR-free if specified), and purified. The

libraries were quantified using Qubit and qPCR, with their size distribution assessed using a fragment analyzer.

Quantified libraries were then pooled and sequenced using the Illumina NovaSeq 6000 platform (Illumina Inc.,

San Diego, CA, USA) according to the effective library concentration and the required data amount, producing a

150-bp paired-end indexed recipe at Novogene (Beijing, China). Low-quality and adapter-containing reads were

filtered out to ensure only high-quality clean reads were used for downstream analysis.

Microbiome analysis of 16S amplicon sequences

Microbiome analysis was performed based on previous studies [13, 14]. To summarise, raw FASTQ

data files were imported into the QIIME2 platform (2022.8) as qza files [15] for quality control and denoising

using DADA2. The sequences were organised into amplicon sequence variants (ASVs), which were then

classified against the SILVA SSU 138 database using the QIIME feature-classifier classification scikit-learn

package.[16, 17]. ASVs classified as mitochondria, chloroplast, or unassigned were excluded from subsequent

analysis. 82,274 reads were subsampled to reduce bias due to differences in read depth between samples.

Shannon diversity and the number of observed features was calculated, as well as the relative species

abundance. The weighted and unweighted UniFrac beta diversity indices were calculated, and the microbial

community structure differences were visualised with principal coordinate analysis plots (PCoA). Data were

visualised using R (version 4.0.4), ggplot2 (version 3.4.3), and ggprism (version 1.0.4).

Microbiome analysis of shallow shotgun metagenomic sequences

Upon completion of sequencing, the Trim Galore pipeline (V0.6.1) [18] was employed for adapter

trimming and quality filtering for the paired-end reads, with a quality threshold of 25 and minimum read length

of 50bp. Metagenomic assembly was carried out using Megahit (v1.2.9) [19]. Taxonomic annotations were

assigned to FASTQ reads using Kraken2 (v.2.1.3) against the standard Kraken2 database, made up of NCBI

taxonomic information, as well as the complete genomes in RefSeq [20]. Sequences classified as Homo sapiens

were removed from the resulting taxonomy table. Metabolic functional trait profiling was performed on the

assembled genomes using the METABOLIC software (v. 4.0) [21], using the METABOLIC-G implementation.

Prokka (v. 1.14.5) [22] was used to annotate the prokaryotic genome, and the outputs were used by

SHORTBRED (v. 0.9.4) [23] in conjunction with pre-computed shortbread markers [24] to determine the

presence and absence of virulence factors and antimicrobial resistance genes across the samples against VFDB

[25] and CARD [26]. Data were visualised using R (version 4.04), ggplot2 (version 3.4.2), and complexheatmap

(version 3.19).

Results

Museum and soil environments characterised by distinct microbial communities

To gain an initial understanding of the taxonomic composition across different environments, we

classified our samples at the genus level (16s rRNA amplicon) and at species level (shallow shotgun). This

revealed that the microbial compositions between the museum environment (MIRAI1-6, MIRAI9) and the soil

environment (MIRAI7-8) were distinctly different, as revealed by both 16s rRNA amplicon sequencing and

shallow shotgun sequencing. The amplicon sequencing results, displayed in Figure 2, indicated that the museum

environment was dominated by genera such as *Acinetobacter* (9.73%), *Enhydrobacter* (11.31%),

Staphylococcus (11.31%), *Bacillus* (4.41%), *Streptomyces* (1.36%), *Corynebacterium* (1.51%)*, Cutibacterium*

(1.48%), and *Brevibacterium* (0.71%), while the soil environment exhibited a different microbial profile with

Pseudonocardia (4.02%), *Streptomyces* (6.67%), *Mycobacterium* (2.81%), *Nocardioides* (1.82%), and

Steroidobacter (0.95%) being the most prevalent. Shallow shotgun sequencing provided a more granular view,

identifying species-level differences within the genera detected by amplicon sequencing (Table 2). In the

museum environment, species such as *Moraxella osloensis* (8.03%), *Cutibacterium acnes* (7.81%),

Staphylococcus aureus (3.68%), *Micrococcus luteus* (3.60%), *Brevibacterium casei* (2.616%), and *Dermacoccus*

nishinomiyaensis (1.98%) were particularly abundant. Conversely, in the soil environment, species like

Pseudonocardia sp. DSM 110487 (2.14%), *Nocardioides sp. NBC_00368* (1.01%), *Streptomyces bathyalis*

(0.796%), *Variovorax paradoxus* (0.70%), *Pseudonocardia sp. HH130630-07* (0.70%), and *Mycolicibacterium*

smegmatis (0.68%) were dominant. The shallow shotgun sequencing approach, with its species-level resolution,

unveiled specific species within the genera, providing a more detailed understanding of the microbial diversity

in both environments.

Planting areas contributed to increasing microbial diversity within the exhibition

In order to generate further insights into the effectiveness of landscape design on microbial diversity,

we analysed the alpha diversity metrics. The alpha diversity indices, as shown in Table 3, reveal notable

differences between the Visionary Lab samples (MIRAI2, MIRAI3, MIRAI4, MIRAI5, MIRAI7, and MIRAI8)

and the non-Visionary-Lab samples (MIRAI1, MIRAI6, and MIRAI9). The Shannon diversity indices for the

Visionary Lab samples tend to be higher, ranging from 9.281 to 10.488, compared to the non-Visionary-Lab

samples, which range from 4.877 to 7.893. This indicates a greater diversity of microbial species in the

Visionary Lab environment. Similarly, the number of observed features is markedly higher in the Visionary Lab

samples, with counts between 1,556 and 2,217, while the non-Visionary-Lab samples show lower counts

ranging from 384 to 1092. The phylogenetic diversity (Faith PD) also appears to reflect this trend, with the

Visionary Lab samples exhibiting values from 85.999 to 150.135, higher than the non-Visionary-Lab samples,

which range from 38.279 to 69.428. Additionally, the Pielou evenness values are generally higher for the

Visionary Lab samples, mostly above 0.87, indicating a more even distribution of species within these microbial

communities, whereas the non-Visionary-Lab samples display lower evenness values ranging from 0.568 to

0.782. Within all samples and across all metrics, MIRAI8 (a soil sample from the visionary lab) shows the

highest overall diversity. These results underscore the impact of the Visionary Lab's curated landscape design on

enhancing microbial diversity, richness, and evenness, as compared to other areas within the museum.

Location-Based Microbial Clustering in Museum Samples

To further understand the differences in microbial community structures, we performed beta diversity

analysis using weighted UniFrac Principal Coordinate Analysis on samples analysed by 16s rRNA amplicon

sequencing. The results, displayed in Figure 3, reveal clear clustering of microbial communities that correspond

to their specific sampling locations within the museum, as mapped in Figure 1.

The PCoA plot can be separated into three main clusters. The first cluster includes samples MIRAI6

and 9, taken from the "Future Backward Thinking" and "Calculator and Nature" exhibits, respectively. These

exhibits are located close to each other, as shown on the map, which corresponds to their similar microbial

community structures. The second cluster consists of samples MIRAI2 and MIRAI1, which again are located

close to one another (whilst MIRAI2 was taken from within the Visionary Lab exhibition, it was towards the

entrance/exit). The third cluster, comprising the remaining samples, were all taken from various locations within

the main area of the visionary lab exhibit, and this proximity is reflected in their similar microbial community

compositions.

No consistent patterns observed in the presence of virulence factors or antimicrobial resistance genes

Having understood the structure of the microbiome of our samples; in order to gain further insights into

the implications, we performed analysis of virulence factors and antimicrobial resistance. Table 4 presents a

comparative analysis of virulence factors and antimicrobial resistance between samples taken from within the

Visionary Lab (MIRAI2-8), where landscape design was implemented, and an external sample (MIRAI1).

Sample MIRAI1, collected outside the premises, exhibited a moderate presence of virulence factors with notable

counts in elements such as fumarate hydratase (VFDBID: VFG038025) and transposase (virulence|961).

However, no antibiotic resistance genes were detected in this sample. Among the Visionary Lab samples,

sample MIRAI3 shows a significant presence of multiple virulence factors with higher counts and hits, such as

biopolymer transporter ExbD (gi 406029600), an uncharacterised protein previously found in *Penicillium*

rubens (gi 15839882) and M28 family peptidase (gi 120401443). Sample M3 also contained antibiotic

resistance genes RbpA and Mycobacterium. Interestingly, sample M7, collected from the soil around the

planting area, shows a presence of virulence factors like phosphoribosyl-AMP cyclohydrolase (VFDB ID:

VFG028663), UV excision repair protein RAD23 homolog B isoform X1 (VFDB ID: VFG009731), and the

same uncharacterised protein identified in MIRAI3 (gi 15839882). It also shared the same antibiotic resistance

gene as MIRAI3- *RbpA* (ADV91011). Sample MIRAI8, also from the soil around the planting area, shows no

detectable virulence factors or antibiotic resistance genes.

Planting areas contribute to microbiomes with a larger array of metabolic functions

Having not been able to elucidate any concrete patterns in virulence or antimicrobial resistance, we

next strived to further understand the functional landscape of the microbiomes. The heatmap in Figure 4

illustrates the presence or absence of genes coding for various metabolic functions across different samples

using a shallow shotgun sequencing approach. Overall, the samples exhibit some common patterns but also

display distinct differences. Categories like "Metal reduction" and "Chlorite reduction" appear consistently

across all samples, indicating these metabolic functions are widely shared. However, other categories, such as

"Complex carbon degradation" and "Nitrogen cycling," show significant variation between samples, suggesting

differing metabolic capabilities. Sample MIRAI3, with a high presence of blue, indicates many metabolic

functions, suggesting it might be the most diverse in terms of metabolic activity. Conversely, samples MIRAI1

and MIRAI2 have the least amount of blue, indicating fewer metabolic functions and suggesting limited

metabolic diversity. Samples MIRAI7 and MIRAI8, collected from the soil around the planting area, show

moderate metabolic diversity, with MIRAI7 exhibiting a broad range of metabolic capabilities, including

complex carbon degradation and nitrogen cycling, pointing to a rich microbial community.

Discussion

Microbial composition and diversity

The distinct microbial profiles observed between the museum and soil samples can be attributed to the

differential ecological niches and associated activities within these settings- they reflect the influence of

different environmental contexts. The museum environment, characterised by genera such as *Acinetobacter*,

Enhydrobacter, *Staphylococcus*, *Bacillus*, *Streptomyces*, *Corynebacterium*, *Cutibacterium*, and *Brevibacterium*,

predominantly harbours human-associated microbes (Figure 2). These taxa are commonly found on human skin,

surfaces frequently touched by humans, and in indoor air, reflecting the high level of human activity and

interaction within the museum environment [27]. For example, *Cutibacterium acnes*, which was notably

abundant in the museum samples, is a well-known skin commensal- the most prevalent bacteria residing in

human, healthy, sebum-rich skin areas such as the face and back [28]. Similarly, *Staphylococcus aureus and*

Staphylococcus hominis, are also human-associated bacteria that thrive in indoor environments, and its presence

in the museum samples align with previous studies reporting that staphylococci, usual colonisers of the human

skin and upper respiratory tract, are commonly spread from humans to built environment surface and air

[29–31]. *Moraxella osloensis* was reported to be the most abundant species across the exhibition samples, and is

gram-negative aerobic bacterium that is considered to be an opportunistic human-associated pathogen,

previously isolated from the upper respiratory tract, blood, genitourethral speciments, and other sites in humans

[32–34]. Again, *Moraxella osloensis* has been found in several locations within the indoor environment where

humans frequent, including sinks and laundry rooms [33, 35]. *Corynebacterium* is another predominant genus in

the skin microbiome, with *Corynebacterium tuberculostearicum* being a ubiquitous skin-colonising bacterium

which has been found to play a role in skin health and disease [36, 37]. These findings are consistent with

previous studies that have shown a high prevalence of human-associated microbes in indoor environments with

significant human traffic ([27, 38, 39].

In contrast, the soil samples from the Visionary Lab's planting area exhibited a different microbial

profile dominated by environment-associated genera such as *Pseudonocardia, Streptomyces, Mycobacterium,*

Nocardioides, and *Steroidobacter* (Figure 2). These taxa are typically associated with soil and plant materials,

indicating the influence of natural environmental conditions in shaping microbial community composition. For

example, members of the genus *Pseudonocardia* are commonly found in soil, plants and the wider environment

[40]. In particular, *Pseudonocardia sp. DSM 110487* and *Pseudonocardia sp. HH 130630-07,* two species

identified by shallow shotgun sequencing to be abundant in the soil samples, are known for their capabilities in

organic matter degradation and their symbiotic relationships with insects, respectively, contributing to the soil's

health and ecological balance [41, 42]. Regarding the genera *Streptomyces,* of which *Streptomyces bathyalis* was

identified as an abundant species within our soil microbiome, studies support our findings, showing that that the

bacteria belonging to this genus are mainly found in soil but are also occasionally isolated from manure and

other environmental sources [43]. The presence of *Mycobacterium* and *Nocardioides* further supports the

environmental origin of our samples, as species from these genera are again frequently isolated from soil and are

involved in nutrient cycling and biodegradation processes [44, 45]. By understanding the distinct microbial

compositions in different contexts, this study underscores the importance of considering both human and

environmental factors in designing and managing indoor spaces to enhance microbial diversity and promote

ecological balance.

The small discrepancies observed between the results obtained from 16s rRNA amplicon sequencing
and shallow shotgun sequencing regarding the most abundant genera and species can likely be attributed to the
limitations and biases inherent in each method. While amplicon sequencing provided a broad overview of the
dominant genera, it was subject to PCR biases that can skew the representation of microbial communities [46].
Additionally, the lack of comprehensive reference databases for soil microbes likely resulted in the
underrepresentation of many species under both sequencing methods [47]. Whilst shallow shotgun sequencing
offered higher resolution at the species level, uncovering specific species within the detected genera, this
method may also have been subjected to potential biases in DNA extraction and library preparation steps. These
findings underscore the importance of using complementary sequencing methods to achieve a comprehensive
characterization of microbial communities. By integrating data from both 16S rRNA amplicon and shallow
shotgun sequencing, we can obtain a more accurate and nuanced picture of the microbial ecosystems present in
different environments [48, 49]. Furthermore, understanding the distinct microbial profiles associated with
human-dominated indoor settings versus natural soil environments can inform strategies for managing microbial
diversity and maintaining healthy indoor microbiomes.

The findings from this study highlight the definitive impact of landscape design on microbial diversity
within the Visionary Lab compared to other areas of the Miraikan museum. The alpha diversity indices,
consistently indicate higher microbial diversity in the Visionary Lab samples compared to the
non-Visionary-Lab samples; underscoring the impact of the curated landscape design on enhancing microbial
diversity (Table 3). Higher Shannon diversity indices in the Visionary Lab compared to the non-Visionary-Lab
suggest a more complex microbial community [50]. This higher diversity is further supported by the increased
number of observed features and higher phylogenetic diversity values, indicating a broader range of microbial
species and greater evolutionary variety in the Visionary Lab, which can enhance ecosystem multifunctionality
[51]. The higher Pielou evenness values in the Visionary Lab samples indicate a more balanced distribution of
species, suggesting that the environment supports a wide range of microbial taxa without dominance by a few
species [52]. Higher evenness within microbial communities is often linked to enhanced ecosystem stability and
productivity [53]. This evenness may be attributed to the diverse plant life and landscape elements designed to
promote a harmonious coexistence of microbes. The notably high diversity observed in sample MIRAI8, a soil
sample from the Visionary Lab, underscores the effectiveness of the landscape design in enhancing microbial
richness and evenness. This finding supports existing literature which consistently suggests that soil, a critical
component of landscape design, plays a crucial role in supporting a diverse microbial ecosystem [54]. Similarly,

a study by van der Heijden et al. (2008) found that plant diversity enhances ecosystem productivity and stability,

partly through its effects on soil microbial communities [55]

The samples outside the Visionary Lab exhibit lower microbial diversity, as evidenced by their alpha

diversity indices (Table 3). Several factors contribute to this. Firstly, the lack of intentional landscape design

outside the exhibition means there are fewer plants and less environmental complexity to support a diverse

microbial community. As discussed, studies have shown that plant diversity and structural complexity provide a

range of niches and resources, which enhance microbial diversity. Without these elements, it is expected that

microbial communities will be less diverse. Secondly, human activity in these areas might introduce

human-associated microbes while reducing environmental microbes - something we can confirm through our

taxonomic assessment (Figure 2). These high-traffic areas can experience frequent cleaning and disinfection,

which can diminish microbial diversity by selectively eliminating certain taxa [56]. Additionally, our taxonomic

assessment revealed that there were few human-associated microbes which dominated in the non-Visionary-Lab

environments, leading to lower overall diversity (reflected in the observed features metric). Thirdly, the

environmental conditions outside the Visionary Lab might not be as conducive to microbial diversity. Factors

such as lower humidity, less organic matter, and fewer plant-derived resources can limit microbial growth and

diversity [57].

The beta diversity analysis reveals significant insights into the spatial organisation and transfer

mechanisms of microbial communities within the museum. The observed clustering of microbial communities,

corresponding to their sampling locations, underscores the influence of spatial proximity on microbial diversity

(Figure 1 and Figure 3). These findings highlight the importance of environmental features and the physical

layout in shaping microbial ecosystems. Mechanisms of microbial transfer play a crucial role in explaining the

formation of these distinct clusters. Human activity is a primary driver of microbial dispersal within built

environments [6]. As visitors move through the museum, they inadvertently carry and deposit microbes through

skin, clothing, and respiratory emissions, contributing to microbial transfer [7] This human-mediated transfer is

particularly evident in areas with high foot traffic, such as the entrances and exits of exhibits. For example, the

clustering of samples MIRAI2 and MIRAI1 can be attributed to the high visitor traffic near the Visionary Lab

entrance, facilitating the homogenization of microbial communities in these regions (Figures 1 & 3). Surface

contact represents another critical mechanism [6]. Surfaces such as exhibit materials, floors, and interactive

displays serve as reservoirs for microbial communities. Visitors touching these surfaces enhance microbial

exchange, particularly in areas with interactive exhibits. Human activity is a significant vector for microbial
transfer, as people can carry microbes on their skin, clothing, and personal items [6]. The interaction between
visitors and the plant-rich environment of the Visionary Lab likely facilitates the transfer and dissemination of
diverse microbial species. Previous studies have shown that environmental conditions, such as moisture levels,
nutrient availability, and vegetation cover, can significantly impact microbial diversity and composition [58, 59].
The clustering of samples within specific exhibits suggests that the unique conditions and interactions within
these microenvironments drive distinct microbial assemblages. Although in this given study we did not collect
data on airflow or ventilation, it is known that airborne dispersal can also contribute to microbial transfer from
different surfaces - for example transfer from soil samples to non-soil samples [60]. Soil and plant surfaces
themselves are reservoirs of microbial life, and their interaction with the surrounding environment can lead to
the spread of microbes, where air currents and ventilation systems lift microbes from soil into the area and
deposit them onto nearby surfaces [61]. This could be the case in the Visionary Lab, where the strategic
placement of environmental features, such as plants, create microhabitats that support diverse microbial
populations, contributing to the unique microbial ecosystems observed within the Visionary Lab samples.

The clustering patterns observed in the beta diversity analysis are consistent with the theory that
proximity to curated environmental features significantly influences microbial composition [62]. This
correlation reinforces the idea that strategic environmental design can be leveraged to shape microbial
ecosystems in desired ways. By understanding the mechanisms of microbial transfer, such as human activity,
airflow, and surface contact, we can optimise the design of built environments to promote beneficial microbial
communities. In conclusion, the distinct community structures revealed by the beta diversity analysis highlight
the importance of spatial organisation and environmental design in shaping microbial ecosystems within the
museum.

Despite the valuable insights gained, it must be acknowledged that due to the small sample size (n=9),
of which only 5 were sequenced by shallow shotgun methods, it may not have been possible to fully capture the
variability and complexity of microbial communities within the Visionary Lab and the Miraikan museum. A
larger sample size would provide a more comprehensive understanding of the microbial diversity in these
environments and allow for further statistical tests to validate the significance of findings.

Virulence factors, antimicrobial resistance, and metabolic function profiles

The comparative analysis of virulence factors and antimicrobial resistance between samples revealed

no clear consistent patterns, highlighting the complexity of microbial community dynamics on human health

(Table 4). The prevalence of virulence factors in sample MIRAI1 suggests a baseline level of microbial

pathogenicity in the external environment, whilst the absence of antibiotic resistance genes in this sample

indicates a potentially lower risk of antibiotic-resistant infections in these settings. The significant presence of

virulence factors in MIRAI3 suggests that microbes in this controlled environment have adapted mechanisms to

enhance their survival and pathogenicity. ExbD, for instance, is involved in nutrient uptake and energy

transduction, which could enhance microbial fitness in nutrient variable conditions. It has been identified before

from *Verrucomicrobiota* bacterium**,** which in turn has been commonly isolated from environmental

environments such as marine and soil locations, as well as the human microbiome such as human faeces [63].

This makes sense in the context of our results, given it was found in sample MIRAI3 which is a sample taken

from a surface- so lots of contact with human environments, but close to the planting area- so transfer from the

soil samples. The detection of antibiotic resistance genes, such as RbpA and Mycobacterium in this sample also

raises concerns about the potential for these environments to harbour and disseminate resistant microbes. RbpA

is known to provide resistance to rifampicin, a critical antibiotic for treating tuberculosis [64]. The presence of

this gene indicates that even in curated environments, there is a risk of developing and spreading antimicrobial

resistance, likely due to selective pressures and microbial interactions. MIRAI 7, collected close to MIRAI 3

from the soil around the planting area, shared several virulence factors and the RbpA resistance gene with

MIRAI3. This suggests that the landscape design, which includes soil and plant materials, creates microhabitats

that support diverse microbial communities with both pathogenic and resistant strains. Phosphoribosyl-AMP

cyclohydrolase, detected in MIRAI7, is crucial for nucleotide biosynthesis and microbial proliferation,

indicating that soil microbes are well-equipped for growth and survival in these enriched environments [65].

The absence of detectable virulence factors and antibiotic resistance genes in MIRAI8 could be due to localised

differences in soil composition or microbial interactions that limit the growth of pathogenic and resistant strains.

A small number of studies [66–69] have concentrated on public settings in built-up environments, but

the majority of these studies have demonstrated that AMR resistance is highly abundant in public settings across

various species [70]. One study revealed that highly maintained environments (subject to intensive cleaning and

disinfection processes) exhibit a higher diversity of antimicrobial resistance genes and reduced microbial

diversity [71]. Based on this we hypothesised that the soil samples taken from the planting area within the

visionary lab would show the least AMR genes and virulence factors (due to its high microbial diversity), and

those with more human contact (samples taken from surfaces) would have more AMR genes and virulence

factors. However, overall, these results suggest that the influence of landscape design on virulence factors and

antimicrobial resistance is complex and does not follow a clear, consistent pattern. This may be due to the

limited sample size and scope of our study not fully capturing the broader patterns and interactions within the

microbial communities. PROKKA and SHORTBRED are tools which depend on the quality and completeness

of the reference databases and the contigs used in the analysis. Incomplete or fragmented contigs may lead to

underestimation or misidentification of genes, affecting the results. The variability observed emphasised the

need for further research to understand the multifaceted interactions between environmental conditions,

microbial communities, and human activity in shaping the presence and distribution of virulence factors and

antimicrobial resistance genes in built environments.

The distinct metabolic profiles highlight how controlled landscape design can shape microbial

communities, fostering unique ecological interactions and functions (Figure 4). Samples which share genes that

code for common functions can be attributed to these functions' broad applications in various settings, both

environmental and human-related. Within the group of functions which exhibit notable variation within the

samples (function present in 30-70% of samples), the most common pattern is for the function to be present in

MIRAI3, MIRAI7 and MIRAI8, and not present within MIRAI1 and MIRAI2. For example, the complex

carbon degradation category follows this pattern. This pattern of distribution of these genes across the samples

can be explained due to the functions within the category being central in an environment with plant material

(e.g. cellulose degrading, hemicellulose-debranching, etc…) [72]. As a result, genes which code for these

functions are more likely to be found within the soil/plant samples (MIRAI7 and MIRAI8). The reason why

these genes are then also found to be present in sample MIRAI3, could be attributed to microbial transfer over

short distances- as MIRAI3 was taken from a location close to the planting area where MIRAI7 and MIRAI8

were taken. This same explanation can be applied to virtually all the remaining functions within this category of

variation within samples. Fermentation is another function that demonstrates significant variation across

samples. Genes associated with fermentation are predominantly found in plant/soil samples (MIRAI7 and

MIRAI8) and are largely absent in samples from human-associated environments (MIRAI1 and MIRAI2).

Fermentation processes are crucial for decomposing plant materials and contributing to soil nutrient cycles [73].

The presence of fermentation genes in these samples underscores the metabolic specialisation of soil microbial

communities in plant-rich environments. In contrast, fermentation-associated genes are less likely to be found in

human samples due to the different metabolic needs and environmental conditions. Human-associated microbial

communities typically rely more on aerobic respiration processes for energy production [74]. The human skin, for example, is an aerobic environment where microbes engage primarily in oxidative phosphorylation rather than fermentation. The absence of plant material and the different nutrient composition on human surfaces make fermentation less relevant, reducing the prevalence of genes associated with this metabolic function in human-associated samples.

This analysis focuses solely on the presence or absence of genes coding for specific functions, which presents certain limitations. For example, in the context of 'nitrogen cycling', the pattern of presence or absence of genes varies across the different 5 functions, leading to the question of why one sample contains one type of gene while lacking genes for closely related functions in the same nitrogen cycle. These ambiguities may arise from biological factors or contamination, complicating interpretation. Thus, more comprehensive analyses considering specific gene abundance are needed to better understand the genetic landscape and its functional implications. Nonetheless, the broad application of certain metabolic functions across various settings underscores their ecological importance, while the variability in others points to localised environmental influences.

Conclusion

This study comprehensively outlines the impact of green infrastructure on microbial diversity, in the context of the Visionary Lab compared to other areas of the Miraikan museum. The curated landscape design, incorporating diverse plant life, supports a more complex and balanced microbial community, contributing to enhanced microbial richness, evenness, and ecosystem multifunctionality. While the Visionary Lab samples exhibited higher microbial diversity, the study did not reveal consistent patterns in virulence factors or antimicrobial resistance genes. This variability underscores the complexity of microbial community dynamics and the influence of localised environmental conditions. Future studies should increase the sample size and variety of sampling locations within and outside the Visionary Lab, as well as employ further advanced sequencing techniques (such as deep metagenomic sequencing) to enhance the robustness and generalizability of results. Additionally, exploring the impact of different environmental variables, such as humidity, temperature, and light exposure, on microbial communities, and conducting longitudinal studies to track microbial changes over time would further elucidate the dynamics of microbial ecosystems. Using this comprehensive approach to build upon the findings from this study would contribute to the development of evidence-based guidelines for

creating microbial-friendly and health-promoting indoor environments, aligning with the broader goal of sustainable urban design.

Declarations

Ethics approval and consent to participate:

Not Applicable.

Consent for publication

Not Applicable

Competing interests

K.I. is a board member at BIOTA Inc., Tokyo, Japan. M.M. is employed by BIOTA Inc. as a research and development intern.

Funding

This work was primarily self-funded, though Novogene provided financial support for 16s rRNA amplicon and shallow shotgun sequencing.

Authors' contributions

K.I. designed the study and analysed the 16s rRNA Amplicon data. M.M. interpreted the 16s rRNA Amplicon data and analysed and interpreted the shallow shotgun data. M.M was the major contributor in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to extend our sincere gratitude to Marin Yamaguchi and Keisuke Otsuka for their diligent work in collecting the samples. Our thanks go to the National Museum of Emerging Science and Innovation (Miraikan) for hosting the "Visionary Lab; Microbes Actually Are All Around" exhibition between April 2022 and August 2023 and for allowing us to collect samples within their facility (https://www.miraikan.jst.go.jp/news/press/202202152332.html). We also thank the Japan Science and Technology Agency (JST) for their collaboration and support in the Visionary Lab exhibition. This study would not have been possible without their collaboration and support.

Figures and Tables

Figure 1: Illustration of the spatial layout of the Visionary Lab and the specific locations where microbial samples were collected. Footprints and coloured paths indicate the movement flow within the exhibit.

Figure 2: Relative abundance of most abundant microbial genera across the samples from A) the museum environment in comparison to those from B) the soil samples (within the Visionary Lab).

Figure 3: Weighted unifrac Principal Coordinate Analysis (PCoA) for samples analysed by 16S rRNA amplicon sequencing. Samples are colour-coded by location and shaped according to surface type.

Figure 4: Heatmap illustrating the presence or absence of genes coding for various metabolic functions across samples sequenced by a shallow shotgun approach. The legend on the right categorises each function by its metabolic category, aiding in the interpretation of metabolic diversity and specialisation among the samples.

Table 1: Summary table regarding samples collected for genomic analysis, and data regarding the number of sequences processed for each sample using the two different sequencing methods. Some samples only have data for 16s rRNA amplicon sequencing as indicated by a dash in the corresponding shallow shotgun column.

Table 2: Table displaying the relative abundance (%) of the 25 most abundant microbial species identified in samples from museum surfaces and soil samples. The 'remainder' row represents the cumulative relative abundance of all other species not listed in the top 25 for each sample.

Table 3: Alpha diversity metrics for all samples. Metrics include Shannon diversity (measure of species richness and evenness), observed features (number of distinct features observed), Faith's Phylogenetic Diversity (measure of biodiversity that incorporates phylogenetic differences between species), and Pielou's Evenness (measure of how evenly the species are distributed).

Table 4: Virulence factors and antibiotic resistance genes identified in samples sequenced by shallow shotgun sequencing. 'Family' represents the database family and gene identifier, 'count' is the total abundance count of each gene, 'hits' is the number of times the gene was detected, and 'marker length' is the length of the gene marker.

References

1. Hu J, Amor DR, Barbier M, Bunin G, Gore J. Emergent phases of ecological diversity and dynamics mapped in microcosms. Science. 2022;378:85–9.

2. Huelsmann M, Ackermann M. Community instability in the microbial world. Science. 2022;378:29–30.

3. Nath A. The Significant Role of Microbial Diversity on Earth's Ecosystems. Applied Microbiology: Open Access. 2023;9:0–1.

4. Bertola M, Ferrarini A, Visioli G. Improvement of Soil Microbial Diversity through Sustainable Agricultural
Practices and Its Evaluation by -Omics Approaches: A Perspective for the Environment, Food Quality and
Human Safety. Microorganisms. 2021;9.

5. Hou K, Wu Z-X, Chen X-Y, Wang J-Q, Zhang D, Xiao C, et al. Microbiota in health and diseases. Signal
Transduct Target Ther. 2022;7:135.

6. Gilbert JA, Stephens B. Microbiology of the built environment. Nat Rev Microbiol. 2018;16:661–70.

7. National Academies of Sciences, Engineering, and Medicine, National Academy of Engineering, Division on
Engineering and Physical Sciences, Health and Medicine Division, Division on Earth and Life Studies, Board
on Infrastructure and the Constructed Environment, et al. The Built Environment and Microbial Communities.
In: Microbiomes of the Built Environment: A Research Agenda for Indoor Microbiology, Human Health, and
Buildings. National Academies Press (US); 2017.

8. Ito K. Understanding the characteristics of the microbiomes of the built environment and their effects on the
human body. Indoor Environment. 2023;26(1):29-42.

9. Bruno A, Fumagalli S, Ghisleni G, Labra M. The Microbiome of the Built Environment: The Nexus for
Urban Regeneration for the Cities of Tomorrow. Microorganisms. 2022;10:2311.

10. Mahnert A, Moissl-Eichinger C, Berg G. Microbiome interplay: plants alter microbial abundance and
diversity within the built environment. Front Microbiol. 2015;6:153475.

11. Rupprecht CDD, Vervoort J, Berthelsen C, Mangnus A, Osborne N, Thompson K, et al. Multispecies
sustainability. Global Sustainability. 2020;3:e34.

12. Funabashi M. Human augmentation of ecosystems: objectives for food production and science by 2045. NPJ
Sci Food. 2018;2:16.

13. Niwa R, Chen D, Seong Y ah, Jo K, Ito K. Direct contact of fermented rice bran beds promotes food-to-hand
transmission of lactic acid bacteria. FEMS Microbiol Lett. 2023;370:fnad120.

14. Ito K, Niwa R, Kobayashi K, Nakagawa T, Hoshino G, Tsuchida Y. A dark matter in sake brewing: Origin of
microbes producing a Kimoto-style fermentation starter. Front Microbiol. 2023;14:1112638.

15. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive,
scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol. 2019;37:852–7.

16. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene
database project: improved data processing and web-based tools. Nucleic Acids Res. 2012;41:D590–6.

17. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing taxonomic
classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome.
2018;6:1–17.

18. FelixKrueger/TrimGalore: v0.6.10 - add default decompression path.
https://doi.org/10.5281/zenodo.7598955.

19. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for large and
complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 2015;31:1674–6.

20. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol. 2019;20:1–13.

21. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, et al. METABOLIC: high-throughput profiling
of microbial genomes for functional traits, metabolism, biogeochemistry, and community-scale functional
networks. Microbiome. 2022;10:1–22.

22. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30:2068–9.

23. Kaminski J, Gibson MK, Franzosa EA, Segata N, Dantas G, Huttenhower C. High-Specificity Targeted
Functional Profiling in Microbial Communities with ShortBRED. PLoS Comput Biol. 2015;11:e1004557.

24. Website. https://huttenhower.sph.harvard.edu/shortbred/.

25. VFDB: Virulence Factors of Bacterial Pathogens. http://www.mgc.ac.cn/VFs/main.htm. Accessed 27 May
2024.

26. McArthur AG, Wright GD. The Comprehensive Antibiotic Resistance Database. https://card.mcmaster.ca/.
Accessed 27 May 2024.

27. Adams RI, Bateman AC, Bik HM, Meadow JF. Microbiota of the indoor environment: a meta-analysis.
Microbiome. 2015;3:1–18.

28. Rozas M, Hart de Ruijter A, Fabrega MJ, Zorgani A, Guell M, Paetzold B, et al. From Dysbiosis to Healthy
Skin: Major Contributions of to Skin Homeostasis. Microorganisms. 2021;9.

29. D'Accolti M, Soffritti I, Bini F, Mazziga E, Mazzacane S, Caselli E. Pathogen Control in the Built
Environment: A Probiotic-Based System as a Remedy for the Spread of Antibiotic Resistance. Microorganisms.
2022;10.

30. Cave R, Misra R, Chen J, Wang S, Mkrtchyan HV. Comparative Genomics Analysis Demonstrated a Link
Between Staphylococci Isolated From Different Sources: A Possible Public Health Risk. Front Microbiol.
2021;12:576696.

31. Madsen AM, Moslehi-Jenabian S, Islam MZ, Frankel M, Spilak M, Frederiksen MW. Concentrations of
Staphylococcus species in indoor air as associated with other bacteria, season, relative humidity, air change rate,
and S. aureus-positive occupants. Environ Res. 2018;160:282–91.

32. Tan L, Grewal PS. Pathogenicity of Moraxella osloensis, a bacterium associated with the nematode
Phasmarhabditis hermaphrodita, to the slug Deroceras reticulatum. Appl Environ Microbiol. 2001;67:5010–6.

33. Kubota H, Mitani A, Niwano Y, Takeuchi K, Tanaka A, Yamaguchi N, et al. Moraxella species are primarily
responsible for generating malodor in laundry. Appl Environ Microbiol. 2012;78:3317–24.

34. Tan L, Grewal PS. Endotoxin activity of Moraxella osloensis against the grey garden slug, Deroceras
reticulatum. Appl Environ Microbiol. 2002;68:3943–7.

35. Zhang T, Liu M, Zhou D, Ma Z, Chen L, Wu D, et al. Environmental factors and particle size shape the
community structure of airborne total and pathogenic bacteria in a university campus. Front Public Health.
2024;12:1371656.

36. Ahmed NM, Joglekar P, Deming C, NISC Comparative Sequencing Program, Lemon KP, Kong HH, et al.
Genomic characterization of the species complex, a ubiquitous member of the human skin microbiome. bioRxiv.
2023. https://doi.org/10.1101/2023.06.16.545375.

37. Altonsy MO, Kurwa HA, Lauzon GJ, Amrein M, Gerber AN, Almishri W, et al. Corynebacterium
tuberculostearicum, a human skin colonizer, induces the canonical nuclear factor-κB inflammatory signaling
pathway in human skin cells. Immun Inflamm Dis. 2020;8:62–79.

38. Gupta M, Lee S, Bisesi M, Lee J. Indoor Microbiome and Antibiotic Resistance on Floor Surfaces: An
Exploratory Study in Three Different Building Types. Int J Environ Res Public Health. 2019;16.

39. National Academies of Sciences, Engineering, and Medicine, National Academy of Engineering, Division
on Engineering and Physical Sciences, Health and Medicine Division, Division on Earth and Life Studies,
Board on Infrastructure and the Constructed Environment, et al. Microorganisms in Built Environments:
Impacts on Human Health. In: Microbiomes of the Built Environment: A Research Agenda for Indoor
Microbiology, Human Health, and Buildings. National Academies Press (US); 2017.

40. Riahi HS, Heidarieh P, Fatahi-Bafghi M. Genus Pseudonocardia: What we know about its biological
properties, abilities and current application in biotechnology. J Appl Microbiol. 2022;132:890–906.

41. Safaei N, Bunk B, Spröer C, Wink J, Mast Y. Complete Genome Sequences of Two Novel Species from the
Family Isolated from the Persian Gulf. Microbiol Resour Announc. 2021;10:e0091821.

42. Bruner-Montero G, Wood M, Horn HA, Gemperline E, Li L, Currie CR. Symbiont-Mediated Protection of

Leaf-Cutter Ants from the Entomopathogenic Fungus Metarhizium anisopliae. MBio. 2021;12:e0188521.

43. STREPTOMYCES. 1999;:2134–8.

44. Ma Y, Wang J, Liu Y, Wang X, Zhang B, Zhang W, et al. : "Specialists" for Hard-to-Degrade Pollutants in
the Environment. Molecules. 2023;28.

45. Walsh CM, Gebert MJ, Delgado-Baquerizo M, Maestre FT, Fierer N. A Global Survey of Mycobacterial
Diversity in Soil. Appl Environ Microbiol. 2019;85.

46. Straub D, Blackwell N, Fuentes AL, Peltzer A, Nahnsen S, Kleindienst S. Interpretations of microbial
community studies are biased by the selected 16S rRNA gene amplicon sequencing pipeline. bioRxiv.
2019;:2019.12.17.880468.

47. Choi J, Yang F, Stepanauskas R, Cardenas E, Garoutte A, Williams R, et al. Strategies to improve reference
databases for soil microbiomes. ISME J. 2017;11:829–34.

48. La Reau AJ, Strom NB, Filvaroff E, Mavrommatis K, Ward TL, Knights D. Shallow shotgun sequencing
reduces technical variation in microbiome analysis. Sci Rep. 2023;13:1–8.

49. Hillmann B, Al-Ghalith GA, Shields-Cutler RR, Zhu Q, Gohl DM, Beckman KB, et al. Evaluating the
Information Content of Shallow Shotgun Metagenomics. mSystems. 2018;3.

50. Kitikidou K, Milios E, Stampoulidis A, Pipinis E, Radoglou K. Using Biodiversity Indices Effectively:
Considerations for Forest Management. Ecologies. 2024;5:42–51.

51. Maherali H, Klironomos JN. Influence of phylogeny on fungal community assembly and ecosystem
functioning. Science. 2007;316:1746–8.

52. Jost L. The Relation between Evenness and Diversity. Diversity . 2010;2:207–32.

53. Wittebolle L, Marzorati M, Clement L, Balloi A, Daffonchio D, Heylen K, et al. Initial community evenness
favours functionality under selective stress. Nature. 2009;458:623–6.

54. Delgado-Baquerizo M, Eldridge DJ, Ochoa V, Gozalo B, Singh BK, Maestre FT. Soil microbial
communities drive the resistance of ecosystem multifunctionality to global change in drylands across the globe.
Ecol Lett. 2017;20:1295–305.

55. van der Heijden MGA, Bardgett RD, van Straalen NM. The unseen majority: soil microbes as drivers of
plant diversity and productivity in terrestrial ecosystems. Ecol Lett. 2008;11:296–310.

56. Chen C-H, Tu C-C, Kuo H-Y, Zeng R-F, Yu C-S, Lu HH-S, et al. Dynamic change of surface microbiota
with different environmental cleaning methods between two wards in a hospital. Appl Microbiol Biotechnol.
2017;101:771–81.

57. Gonzalez JM, Aranda B. Microbial Growth under Limiting Conditions-Future Perspectives.
Microorganisms. 2023;11.

58. Martiny JBH. Dispersal and the Microbiome. Microbe Wash DC. 2015;10:191–6.

59. Fierer N, Jackson RB. The diversity and biogeography of soil bacterial communities. Proc Natl Acad Sci U
S A. 2006;103:626–31.

60. Tong X, Leung MHY, Wilkins D, Lee PKH. City-scale distribution and dispersal routes of mycobiome in
residences. Microbiome. 2017;5:1–13.

61. Gupta A, Gupta R, Singh RL. Microbes and Environment. In: Principles and Applications of Environmental
Biotechnology for a Sustainable Future. Singapore: Springer Singapore; 2017. p. 43–84.

62. Junkins EN, McWhirter JB, McCall L-I, Stevenson BS. Environmental structure impacts microbial
composition and secondary metabolism. ISME Commun. 2022;2:15.

63. Taxonomy browser (Verrucomicrobiota bacterium).

https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=2026799. Accessed 27 May 2024.

64. Hu Y, Morichaud Z, Chen S, Leonetti J-P, Brodolin K. Mycobacterium tuberculosis RbpA protein is a new type of transcriptional activator that stabilizes the σ A-containing RNA polymerase holoenzyme. Nucleic Acids Res. 2012;40:6547–57.

65. Fisher G, Pečaver E, Read BJ, Leese SK, Laing E, Dickson AL, et al. Catalytic Cycle of the Bifunctional Enzyme Phosphoribosyl-ATP Pyrophosphohydrolase/Phosphoribosyl-AMP Cyclohydrolase. ACS Catal. 2023. https://doi.org/10.1021/acscatal.3c01111.

66. Cave R, Misra R, Chen J, Wang S, Mkrtchyan HV. Whole genome sequencing revealed new molecular characteristics in multidrug resistant staphylococci recovered from high frequency touched surfaces in London. Sci Rep. 2019;9:9637.

67. Kang K, Ni Y, Li J, Imamovic L, Sarkar C, Kobler MD, et al. The Environmental Exposures and Inner- and Intercity Traffic Flows of the Metro System May Contribute to the Skin Microbiome and Resistome. Cell Rep. 2018;24:1190–202.e5.

68. Roberts MC, Soge OO, No D. Comparison of Multi-Drug Resistant Environmental Methicillin-Resistant Staphylococcus aureus Isolated from Recreational Beaches and High Touch Surfaces in Built Environments. Front Microbiol. 2013;4:74.

69. Shen C, Feng S, Chen H, Dai M, Paterson DL, Zheng X, et al. Transmission of mcr-1-Producing Multidrug-resistant Enterobacteriaceae in Public Transportation in Guangzhou, China. Clin Infect Dis. 2018;67:S217–24.

70. Surveillance and prevalence of antimicrobial resistant bacteria from public settings within urban built environments: Challenges and opportunities for hygiene and infection control. Environ Int. 2021;157:106836.

71. Mahnert A, Moissl-Eichinger C, Zojer M, Bogumil D, Mizrahi I, Rattei T, et al. Man-made microbial resistances in built environments. Nat Commun. 2019;10:1–12.

72. Yu T, Wu W, Liang W, Wang Y, Hou J, Chen Y, et al. Anaerobic degradation of organic carbon supports uncultured microbial populations in estuarine sediments. Microbiome. 2023;11:81.

73. Vassileva M, Malusà E, Sas-Paszt L, Trzcinski P, Galvez A, Flor-Peregrin E, et al. Fermentation Strategies to Improve Soil Bio-Inoculant Production and Quality. Microorganisms. 2021;9.

74. Fischbach MA, Sonnenburg JL. Eating for two: how metabolism establishes interspecies interactions in the gut. Cell Host Microbe. 2011;10:336–47.

Sample	Collection protocol	Surface type	Place	Reads processed	
				16s rRNA amplicon	Shallow shotgun
MIRAI1	Vacuum	Floor	Near the 'Earth'	114,111	21,916,203
MIRAI2	Vacuum	Floor	Visionary Lab	107,616	30,083,584
MIRAI3	Vacuum	Floor	Visionary Lab near the planting area	109,227	39,829,873
MIRAI4	Swab	Wall	Visionary Lab	121,281	-
MIRAI5	Swab	Desk	Visionary Lab	110,168	-
MIRAI6	Swab	Wall	Calculator and nature	117,977	-
MIRAI7	Swab	Soil	Visionary Lab planting area	104,243	21,768,309
MIRAI8	Swab	Soil	Visionary Lab planting area	82,274	27,126,330
MIRAI9	Swab	Desk	Future backward thinking	115,917	-

Museum microbiome				Soil microbiome		
	Relative abundance				**Relative abundance**	
25 most abundant species	MIRAI1	MIRAI2	MIRAI3	**25 most abundant species**	MIRAI7	MIRAI8
Moraxella osloensis	16.88	7.19	0.02	*Pseudonocardia sp. DSM 110487*	3.94	0.33
Cutibacterium acnes	14.03	9.35	0.04	*Nocardioides sp. NBC_00368*	1.95	0.07
Staphylococcus aureus	0.26	10.79	0.00	*Streptomyces bathyalis*	1.56	0.03
Micrococcus luteus	9.35	1.44	0.02	*Variovorax paradoxus*	0.48	1.02
Nocardioides sp. NBC_00368	0.26	6.47	2.64	*Pseudonocardia sp. HH130630-07*	0.15	1.25
Brevibacterium casei	7.79	0.00	0.06	*Mycolicibacterium smegmatis*	1.30	0.07
Dermacoccus nishinomiyaensis	5.19	0.72	0.02	*Kribbella sp. NBC_00482*	1.19	0.03
Janibacter sp. CX7	0.78	4.32	0.50	*Pseudonocardia sp. AL041005-10*	0.15	1.05
Corynebacterium tuberculostearicum	4.68	0.72	0.00	*Pseudoxanthomonas suwonensis*	0.93	0.26
Janibacter limosus	0.52	4.32	0.32	*Stackebrandtia nassauensis*	1.10	0.03
Staphylococcus hominis	4.16	0.72	0.00	*Mycolicibacterium moriokaense*	1.02	0.07
Agrococcus carbonis	0.26	4.32	0.11	*Streptomyces xanthii*	1.04	0.03
Sphingomonas sp. FARSPH	3.12	1.44	0.06	*Ramlibacter tataouinensis*	0.13	0.89
Janibacter terrae	0.52	3.60	0.26	*Enhydrobacter sp.*	0.26	0.69
Pseudonocardia sp. DSM 110487	0.00	1.44	2.26	*Pseudonocardia sp. HH130629-09*	0.11	0.79
Brachybacterium sp. SGAir0954	0.26	2.16	1.18	*Sorangium cellulosum*	0.37	0.46
Janibacter melonis	0.52	2.16	0.88	*Streptomyces nigra*	0.78	0.03
Nocardioides sp. JQ2195	0.00	2.16	1.38	*Usitatibacter rugosus*	0.09	0.72
Corynebacterium afermentans	3.38	0.00	0.00	*Variovorax sp. HW608*	0.35	0.46
Dermacoccus sp. PAMC28757	2.60	0.72	0.02	*Luteitalea sp. TBR-22*	0.17	0.62
Janibacter sp. A1S7	0.26	2.88	0.07	*Croceibacterium sp. D39*	0.37	0.43
Brachybacterium sp. NBEC-018	0.00	2.16	0.97	*Lysobacter soli*	0.56	0.23
Janibacter indicus	1.30	1.44	0.19	*Luteitalea pratensis*	0.15	0.56
Mycolicibacterium smegmatis	0.00	0.72	1.79	*Kribbella sp. NBC_00709*	0.67	0.03
Nocardioides sp. zg-1228	0.00	1.44	0.95	*Lacunisphaera limnophila*	0.26	0.43
Remainder	23.90	27.34	86.26	Remainder	80.92	89.43

Sample	Alpha diversity metrics			
	Shannon diversity	Observed features	Faith PD	Pielou evenness
MIRAI1	7.893	1,092	69.428	0.782
MIRAI2	9.281	1,597	112.933	0.872
MIRAI3	9.736	1,694	105.363	0.908
MIRAI4	9.955	1,616	88.141	0.934
MIRAI5	9.368	1,556	85.999	0.883
MIRAI6	4.877	384	38.279	0.568
MIRAI7	9.991	1,702	86.597	0.568
MIRAI8	10.488	2,217	150.135	0.944
MIRAI9	5.702	524	44.321	0.631

Sample	Virulence Factor Database				Complex Antibiotic Resistance Database			
	Family	Count	Hits	Marker length	Family	Count	Hits	Marker length
MIRAI1	VFDB|VFG038025(gi_384141332)	350.086	1	42				
	virulence|961|vfid|962|vsiid|961|ssid|transposase	231.307	1	50	-			
MIRAI2	Victors|gi|15926930|ref|NP_374463_1|	468.345	1	27				
	virulence|18940|vfid|42469|vsiid|63306|ssid|hypothetical	20.314	1	168	-			
MIRAI3	VFDB|VFG030641(gi_406029600)	140.943	3	30	gb|AAK44936_1|ARO_3003395|Mycobacterium	368.619	9	102
	VFDB|VFG031723(gi_15839882)	84.071	2	23	gb|ADV91011_1|ARO_3000245|RbpA	199.669	1	48
	VFDB|VFG022109(gi_120401443)	72.607	2	22				
	VFDB|VFG022671(gi_333990322)	44.371	1	28				
MIRAI7	VFDB|VFG028663(gi_433629379)	98.400	1	24	gb|ADV91011_1|ARO_3000245|RbpA	418.201	1	48
	VFDB|VFG009731(gi_118618107)	176.085	1	14				
	VFDB|VFG031723(gi_15839882)	176.085	2	23				
MIRAI8	-				-			

Backward Future Thinking

MIRAI 9

Calculator and Nature

MIRAI 6

Near the Earth

MIRAI 1

Visionary Lab

MIRAI 4

MIRAI 5

MIRAI 7, 8

MIRAI 2

MIRAI 3