Title

From Soil to Surface: Exploring the Impact of Green Infrastructure on Microbial Communities in the Built

Environment

Authors

Malin Mcgonigal[1,2] and Kohei Ito [1*]

Affiliation

1.   BIOTA Inc., Tokyo, 101-0022, Japan

2.   Department of Biosciences, Durham University, Durham, DH1 3LE, England

*Corresponding author

e-mail: kohei@biota.ne.jp

Abstract

High microbial diversity offers extensive benefits to both the environment and human health, contributing to

ecosystem stability, nutrient cycling, and pathogen suppression. In built environments, factors such as building

design, human activity, and cleaning protocols influence microbial communities. This study investigates the

impact of landscape design on microbial diversity and function within the "Visionary Lab" exhibition in Tokyo,

Japan, using 16S rRNA gene amplicon sequencing and shallow shotgun sequencing. Despite the limited sample

size, the study suggests that the Visionary Lab samples may exhibit higher microbial diversity compared to

other museum areas. Potential distinct microbial community structures may be correlated with sampling

locations. However, despite this, no consistent patterns were observed in virulence factors or antimicrobial

resistance genes across the samples. Metabolic function analysis showed varied profiles, suggesting diverse

ecological interactions influenced that may be by the curated landscape. This suggest that the curated landscape

design may have the potential to enhance microbial diversity, highlighting a possible avenue to create healthier

and more sustainable built environments. However, the lack of consistent patterns in virulence factors and

antimicrobial resistance genes underscores the complexity of microbial community dynamics.

Keywords

Landscape design, Metagenomics, Microbiome of the built environment, Microbial diversity, Sustainability

Background

Recent research has increasingly recognized the importance of microbial communities in built environments, particularly their potential impact on human health and well-being. This growing field of study sits at the intersection of microbiology, architecture, and public health, offering new perspectives on how we design and manage our indoor spaces. The benefits of high microbial diversity are extensive [1, 2]. In the environment, diverse microbial communities contribute to the breakdown of organic matter and nutrient cycling, thereby maintaining ecosystem stability and productivity [3]. Soils with greater biodiversity levels are more resistant to environmental disturbances than those with reduced biodiversity, allowing ecosystems to better withstand and recover from disturbances such as pollution, climate change and habitat destruction [4]. Moreover, a complex soil microbial community increases competition for nutrients, inhibiting the development or persistence of pathogens in the soil. Thus, soil microbial diversity can directly benefit human health by suppressing disease-causing organisms, providing clean air, water, and food, and serving as a source of antibiotics [4]. In human health, it has been reported that 'a healthy microbiota community often demonstrates high taxonomic diversity, high microbial gene richness, and stable core microbiota' [5]. Such healthy microbiota is associated with improved immune function, reduced inflammation, improved energy and nutrient extraction, and a lower incidence of allergies and autoimmune diseases [5].

In built environments, such as homes, offices, and public spaces, microbial communities are influenced by multiple factors, including building design, ventilation systems, occupancy and human activity, types of construction materials, cleaning protocols, and external environmental conditions [6–8]. These elements interact to shape the microbial communities found in indoor spaces, affecting everything from air quality to human health. Understanding how these factors influence microbial diversity is crucial for designing healthier and more sustainable built environments [6]. The design and integration of landscape elements within built environments is one factor that can significantly influence microbial communities. Plants, in particular, have been shown to enhance microbial abundance and diversity within indoor environments [9, 10]. However, the extent to which plants influence taxonomic diversity, including genes associated with various metabolic functions or antimicrobial resistance, remains inadequately explored.

From April 2022 until August 2023, the National Museum of Emerging Science and Innovation ('Miraikan') in Tokyo, Japan, held an exhibition titled "Visionary Lab; Microbes Actually Are All Around".

The exhibition featured a full-scale futuristic living space with an adjoining planting area of purposefully

curated floral and shrubbery arrangements. It was designed to encourage visitors to contemplate future lifestyles

where humans and microbes exist plentifully in harmony. Miraikan's planting area integrates two key concepts:

multispecies sustainability and human augmentation of ecosystems. Multispecies sustainability emphasises the

interconnectedness of humans, plants, and microbes, advocating for practices that support all species' health and

well-being [11]. Human augmentation of ecosystems involves deliberately enhancing of natural environments to

improve ecosystem services and resilience [12].

With its landscape design, the Visionary Lab provided an opportunity to investigate the potential

effects of these concepts in a real-world setting. While acknowledging the limitations of our study design,

particularly the small sample size and single time point of sampling, this study aimed to provide preliminary

insights into the effects of the Visionary Lab's landscape design by comparing microbial samples from within

the exhibition space to those from other areas of the Miraikan museum, using 16S rRNA gene sequencing as

well as shallow shotgun sequencing to supplement our understanding of the different microbiomes. Our primary

objectives were to ascertain controlled landscape design's potential effects on the microbial ecosystem in this

unique setting. We hypothesized that the specific landscape design of the Visionary Lab, particularly through

the strategic integration of plant life and landscape elements, might cultivate a unique and rich microbial

ecosystem that could markedly contrast with other areas within the museum. While we recognize that our study

is limited in scope and cannot provide definitive conclusions, we believe it offers valuable preliminary insights

that can guide future, more comprehensive research in this area. This study aims to contribute to informing

urban design and public health strategies. The findings from this research may help guide future studies

exploring how built environments can be designed to enhance microbial diversity, potentially contributing to

improved air quality, reduced pathogen transmission, and overall public health. This approach aligns with the

broader goal of sustainable urban design, which seeks to create living spaces that are resilient, ecologically

balanced, and conducive to human well-being.

Methods

Study design, sample collection, and DNA extraction

This study employed a cross-sectional design, collecting samples from different locations within the

National Museum of Emerging Science and Innovation ('Miraikan') in Tokyo, Japan. The study included nine

samples (n=9) collected from various surfaces and locations within the museum. The study was conducted at the

National Museum of Emerging Science and Innovation ('Miraikan'), located in 2--3-6 Aomi, Koto-ku, Tokyo

135-0064. Microbial samples were collected via vacuum collection or surface swabbing from different surfaces

and locations, as listed in Table 1 and mapped in Figure 1. For vacuum collection, a Makita vacuum cleaner

(CL107FDSH) equipped with a DUSTREAM Collector (DU-ST-1) and DUSTREAM Filter (DU-FL-1) from

Indoor Biotechnologies was used. For surface swabbing, a sterile cotton-tipped swab (ESwab™; Copan

Diagnostics, Brescia, Italy) was used to swab surfaces for three minutes before they were stored in tubes

containing Liquid Amies Medium solution. Care was taken to ensure that collection methods did not

contaminate or alter the microbial communities present. The samples were promptly sealed, stored at low

temperatures, and refrigerated until DNA extraction. Extraction was performed on all nine samples using the

DNeasy® PowerSoil® Pro Kit (QIAGEN, Germany) according to the manufacturer's protocol.

Library preparation and sequencing

Sequencing libraries were generated for the 16S rRNA amplicon sequencing, and indexes were

incorporated to attribute sequences to individual samples. PCR amplification of the targeted V3-V4 regions was

performed using specific primers, CCTAYGGGRBGCASCAG and GGACTACNNGGGTATCTAAT. PCR

products with proper size were selected via 2% agarose gel electrophoresis. An equal amount of PCR products

from each sample was pooled, end-repaired, A-tailed, and then ligated with Illumina adapters. The libraries were

sequenced on a paired-end Illumina platform, producing 250 base pair paired-end raw reads. Library quality was

evaluated and quantified using qPCR. Quantified libraries were then pooled and sequenced on Illumina

platforms according to the effective library concentration and the data amount required. DNA samples, library

preparation, and amplicon sequencing were performed using 250 base paired-end sequencing on the Illumina

NovaSeq 6000 system at Novogene (Beijing, China).

Sequencing libraries were prepared for the shallow shotgun metagenomic sequencing, and quality

control (QC) was performed at each step to ensure data reliability. The DNA samples were randomly

fragmented into shorter sequences: end-repaired, A-tailed, and ligated with Illumina adapters. These adapter-

linked fragments were size-selected, PCR-amplified, and purified. The libraries were quantified using Qubit and

qPCR, and their size distribution assessed using a fragment analyzer. Quantified libraries were then pooled and

sequenced using the Illumina NovaSeq 6000 platform (Illumina Inc., San Diego, CA, USA) according to the

effective library concentration and the required data amount, producing a 150-bp paired-end indexed recipe at

Novogene (Beijing, China). Low-quality and adapter-containing reads were filtered out to ensure that only high-

quality clean reads were used for downstream analysis.

Microbiome analysis of 16S amplicon sequences

Microbiome analysis was performed based on previous studies [13, 14]. To summarise, raw FASTQ

data files were imported into the QIIME2 platform (2022.8) as qza files [15] for quality control and denoising

using DADA2. The sequences were organised into amplicon sequence variants (ASVs), which were then

classified against the SILVA SSU 138 database using the QIIME feature-classifier classification scikit-learn

package.[16, 17]. The subsequent analysis excluded ASVs classified as mitochondria, chloroplast, or unassigned.

82,274 reads were subsampled to reduce bias due to differences in read depth between samples. Shannon

diversity, the number of observed features were calculated, as well as the relative species abundance. The

weighted and unweighted UniFrac beta diversity indices were calculated, and the microbial community structure

differences were visualised with principal coordinate analysis plots (PCoA). Data were visualised using R

(version 4.0.4), ggplot2 (version 3.4.3), and ggprism (version 1.0.4).

Microbiome analysis of shallow shotgun metagenomic sequences

Upon completion of sequencing, the Trim Galore pipeline (V0.6.1) [18] was employed for adapter

trimming and quality filtering for the paired-end reads, with a quality threshold of 25 and a minimum read

length of 50bp. To remove the human genome from quality-controlled sequencing, hg38 was obtained from the

UCSC genome browser [19] and mapped in bowtie2 (version 2.5.4) [20]. Metagenomic assembly was carried

out using Megahit (v1.2.9) [21]. Taxonomic annotations were assigned to FASTQ reads using Kraken2 (v.2.1.3)

against the standard Kraken2 database, made up of NCBI taxonomic information, as well as the complete

genomes in RefSeq [22]. Sequences classified as Homo sapiens were removed from the resulting taxonomy

table. Metabolic functional trait profiling was performed on the assembled genomes using the METABOLIC

software (v. 4.0) [23], using the METABOLIC-G implementation. Prokka (v. 1.14.5) [24] was used to annotate

the prokaryotic genome, and the outputs were used by SHORTBRED (v. 0.9.4) [25] in conjunction with pre-

computed shortbread markers [26] to determine the presence and absence of virulence factors and antimicrobial

resistance genes across the samples against VFDB [27] and CARD [28]. Data were visualised using R (version

4.04), ggplot2 (version 3.4.2), and complexheatmap (version 3.19).

<u>Results</u>

<u>Museum and soil environments characterised by distinct microbial communities</u>

To understand of the taxonomic composition across different environments, we classified our samples

at the genus level (16S rRNA amplicon) and at the species level (shallow shotgun). This revealed that the

microbial compositions between the museum environment (MIRAI1-6, MIRAI9) and the soil environment

(MIRAI7-8) were notably different, as revealed by both 16s rRNA amplicon sequencing and shallow shotgun

sequencing. The amplicon sequencing results, displayed in Figure 2, suggest that the museum environment was

characterized by a prevalence of genera such as *Acinetobacter* (9.73%), *Enhydrobacter* (11.31%),

Staphylococcus (11.31%), *Bacillus* (4.41%), *Streptomyces* (1.36%), *Corynebacterium* (1.51%)*, Cutibacterium*

(1.48%), and *Brevibacterium* (0.71%), while the soil environment exhibited a different microbial profile with

Pseudonocardia (4.02%), *Streptomyces* (6.67%), *Mycobacterium* (2.81%), *Nocardioides* (1.82%), and

Steroidobacter (0.95%) being the most prevalent. Shallow shotgun sequencing provided a more granular view,

identifying species-level differences within the genera detected by amplicon sequencing (Table 2). In the

museum environment, species such as *Moraxella osloensis* (8.03%), *Cutibacterium acnes* (7.81%),

Staphylococcus aureus (3.68%), *Micrococcus luteus* (3.60%), *Brevibacterium casei* (2.616%), and

Dermacoccus nishinomiyaensis (1.98%) were particularly abundant. Conversely, in the soil environment,

species like *Pseudonocardia sp. DSM 110487* (2.14%), *Nocardioides sp. NBC_00368* (1.01%), *Streptomyces*

bathyalis (0.796%), *Variovorax paradoxus* (0.70%), *Pseudonocardia sp. HH130630-07* (0.70%), and

Mycolicibacterium smegmatis (0.68%) were observed to be dominant.

Planting areas contributed to increasing microbial diversity within the exhibition

In order to generate further insights into the effectiveness of landscape design on microbial diversity,

we analysed the alpha diversity metrics. The alpha diversity indices, as shown in Table 3, indicate notable

differences between the Visionary Lab samples (MIRAI2, MIRAI3, MIRAI4, MIRAI5, MIRAI7, and MIRAI8)

and the non-Visionary-Lab samples (MIRAI1, MIRAI6, and MIRAI9). The Shannon diversity indices for the

Visionary Lab samples tended to be higher, ranging from 9.28 to 10.48, compared to the non-Visionary-Lab

samples, which ranged from 4.87 to 7.89. This suggests a potentially greater diversity of microbial species in the

Visionary Lab environment. Similarly, the number of observed features was markedly higher in the Visionary

Lab samples, with counts between 1,556 and 2,217, while the non-Visionary-Lab samples showed lower counts

ranging from 384 to 1,092. The phylogenetic diversity (Faith PD) also appeared to reflect this trend, with the

Visionary Lab samples exhibiting values from 85.99 to 150.13, higher than the non-Visionary-Lab samples,

which ranged from 38.27 to 69.42. Additionally, the Pielou evenness values are generally higher for the

Visionary Lab samples, mostly above 0.87, potentially indicating a more even distribution of species within

these microbial communities, whereas the non-Visionary-Lab samples displayed lower evenness values ranging

from 0.56 to 0.78. Within all samples and across all metrics, MIRAI8 (a soil sample from the visionary lab)

shows the highest overall diversity.

Location-Based Microbial Clustering in Museum Samples

To further understand the differences in microbial community structures, we performed beta diversity

analysis using weighted UniFrac Principal Coordinate Analysis on samples analysed by 16s rRNA amplicon

sequencing. The results, displayed in Figure 3, reveal clear clustering of microbial communities that correspond

to their specific sampling locations within the museum, as mapped in Figure 1.

The PCoA plot can be broadly separated into three main clusters. The first cluster includes samples

MIRAI6 and 9, taken from the "Future Backward Thinking" and "Calculator and Nature" exhibits, respectively.

These exhibits are located close to each other, as shown on the map, which may explain to their similar

microbial community structures. The second cluster consists of samples MIRAI2 and MIRAI1, which again are

located close to one another (whilst MIRAI2 was taken from within the Visionary Lab exhibition, it was

towards the entrance/exit). The third cluster, comprising the remaining samples, were all taken from various

locations within the main area of the visionary lab exhibit, and this proximity is reflected in their similar

microbial community compositions.

<u>No consistent patterns observed in the presence of virulence factors or antimicrobial resistance genes</u>

Having understood the structure of the microbiome of our samples; in order to gain further insights into

the implications, we performed analysis of virulence factors and antimicrobial resistance. Table 4 presents a

comparative analysis of virulence factors and antimicrobial resistance between samples taken from within the

Visionary Lab (MIRAI2-8), where landscape design was implemented, and an external sample (MIRAI1).

Sample MIRAI1, collected outside the premises, exhibited a moderate presence of virulence factors with notable

counts in elements such as fumarate hydratase (VFDBID: VFG038025) and transposase (virulence|961).

However, no antibiotic resistance genes were detected in this sample. Among the Visionary Lab samples,

sample MIRAI3 showed a significant presence of multiple virulence factors with higher counts and hits, such as

biopolymer transporter ExbD (gi 406029600), an uncharacterised protein previously found in *Penicillium*

rubens (gi 15839882) and M28 family peptidase (gi 120401443). Sample M3 also contained antibiotic

resistance genes *RbpA* and *Mycobacterium*. Interestingly, sample M7, collected from the soil around the

planting area, showed a presence of virulence factors like phosphoribosyl-AMP cyclohydrolase (VFDB ID:

VFG028663), UV excision repair protein RAD23 homolog B isoform X1 (VFDB ID: VFG009731), and the

same uncharacterised protein identified in MIRAI3 (gi 15839882). It also shared the same antibiotic resistance

gene as MIRAI3- *RbpA* (ADV91011). Sample MIRAI8, also from the soil around the planting area, shows no

detectable virulence factors or antibiotic resistance genes.

217
<u>Planting areas contribute to microbiomes with a larger array of metabolic functions</u>

Having not been able to elucidate any concrete patterns in virulence or antimicrobial resistance, we

next strived to further understand the functional landscape of the microbiomes. The heatmap in Figure 4

illustrates the presence or absence of genes coding for various metabolic functions across different samples

using a shallow shotgun sequencing approach. Overall, the samples exhibited some common patterns but also

displayed distinct differences. Categories like "Metal reduction" and "Chlorite reduction" appeared consistently

across all samples, indicating these metabolic functions are widely shared. However, other categories, such as

"Complex carbon degradation" and "Nitrogen cycling," show significant variation between samples, suggesting

differing metabolic capabilities. Sample MIRAI3, with a high presence of blue, indicates many metabolic

functions, suggesting it might be the most diverse in terms of metabolic activity. Conversely, samples MIRAI1

and MIRAI2 have the least amount of blue, indicating fewer metabolic functions and suggesting limited

metabolic diversity. Samples MIRAI7 and MIRAI8, collected from the soil around the planting area, showed

moderate metabolic diversity, with MIRAI7 exhibiting a broad range of metabolic capabilities, including

complex carbon degradation and nitrogen cycling, pointing to a rich microbial community.

Discussion

Microbial composition and diversity

The observed distinct microbial profiles between the museum and soil samples can be attributed to the

differential ecological niches and associated activities within these settings. These profiles appear to reflect the

influence of different environmental contexts. The museum environment, characterised by genera such as

Acinetobacter, *Enhydrobacter*, *Staphylococcus*, *Bacillus*, *Streptomyces*, *Corynebacterium*, *Cutibacterium*, and

Brevibacterium, predominantly appears to harbour human-associated microbes (Figure 2). These taxa are

commonly found on human skin, surfaces frequently touched by humans, and in indoor air, potentially

reflecting the high level of human activity and interaction within the museum environment [29]. For example,

Cutibacterium acnes, which was notably abundant in the museum samples, is a well-known skin commensal-

the most prevalent bacteria residing in human, healthy, sebum-rich skin areas such as the face and back [30].

Similarly, *Staphylococcus aureus and Staphylococcus hominis*, are also human-associated bacteria that thrive in

indoor environments, and their presence in the museum samples aligns with previous studies reporting that

staphylococci, usual colonisers of the human skin and upper respiratory tract, are commonly spread from

humans to built environment surface and air [31–33]. *Moraxella osloensis* was found to be the most abundant

species across the exhibition samples, and is a gram-negative aerobic bacterium that is considered to be an

opportunistic human-associated pathogen, previously isolated from the upper respiratory tract, blood,

genitourethral speciments, and other sites in humans [34–36]. Again, *Moraxella osloensis* has been found in

several locations within the indoor environment where humans frequent, including sinks and laundry rooms [35,

37]. *Corynebacterium* is another predominant genus in the skin microbiome, with *Corynebacterium*

tuberculostearicum being a ubiquitous skin-colonising bacterium which has been found to play a role in skin

health and disease [38, 39]. These findings are consistent with previous studies that have shown a high

prevalence of human-associated microbes in indoor environments with significant human traffic [29, 40, 41].

In contrast, the soil samples from the Visionary Lab's planting area exhibited a different microbial

profile characterized by environment-associated genera such as *Pseudonocardia, Streptomyces, Mycobacterium,*

Nocardioides, and *Steroidobacter* (Figure 2). These taxa are typically associated with soil and plant materials,

indicating the influence of natural environmental conditions in shaping microbial community composition. For

example, members of the genus *Pseudonocardia* are commonly found in soil, plants and the wider environment

[42]. In particular, *Pseudonocardia sp.* DSM 110487 and *Pseudonocardia sp. HH 130630-07,* two species

identified by shallow shotgun sequencing to be abundant in the soil samples, are known for their capabilities in

organic matter degradation and their symbiotic relationships with insects, respectively, contributing to the soil's

health and ecological balance [43, 44]. Regarding the genera *Streptomyces,* of which *Streptomyces bathyalis*

was identified as an abundant species within our soil microbiome, studies support our findings, showing that

that the bacteria belonging to this genus are mainly found in soil but are also occasionally isolated from manure

and other environmental sources [45]. The presence of *Mycobacterium* and *Nocardioides* further supports the

environmental origin of our samples, as species from these genera are again frequently isolated from soil and are

involved in nutrient cycling and biodegradation processes [46, 47]. By understanding the distinct microbial

compositions in different contexts, this study begins to shed light on the importance of considering both human

and environmental factors in designing and managing indoor spaces to enhance microbial diversity and promote

ecological balance.

The small discrepancies observed between the results obtained from 16s rRNA amplicon sequencing

and shallow shotgun sequencing regarding the most abundant genera and species can likely be attributed to the

limitations and biases inherent in each method. While amplicon sequencing provided a broad overview of the

dominant genera, it was subject to PCR biases that can skew the representation of microbial communities [48].

Additionally, the lack of comprehensive reference databases for soil microbes likely resulted in the

underrepresentation of many species under both sequencing methods [49]. Whilst shallow shotgun sequencing

offered higher resolution at the species level, uncovering specific species within the detected genera, this

method may also have been subjected to potential biases in DNA extraction and library preparation steps. These

findings underscore the importance of using complementary sequencing methods to achieve a comprehensive

characterization of microbial communities. By integrating data from both 16S rRNA amplicon and shallow

shotgun sequencing, we can obtain a more accurate and nuanced picture of the microbial ecosystems present in

different environments [50, 51]. Furthermore, understanding the distinct microbial profiles associated with

human-dominated indoor settings versus natural soil environments can inform strategies for managing microbial

diversity and maintaining healthy indoor microbiomes.

Our findings suggest a potential impact of landscape design on microbial diversity within the Visionary

Lab compared to other areas of the Miraikan museum. While limited by our small sample size, consistently

suggest higher microbial diversity in the Visionary Lab samples compared to the non-Visionary-Lab samples

(Table 3). Higher Shannon diversity indices in the Visionary Lab compared to the non-Visionary-Lab suggest a

more complex microbial community [52]. This higher diversity is further supported by the increased number of

observed features and higher phylogenetic diversity values, indicating a broader range of microbial species and

greater evolutionary variety in the Visionary Lab, which could potentially enhance ecosystem multifunctionality

[53]. The higher Pielou evenness values in the Visionary Lab samples indicate a more balanced distribution of

species, suggesting that the environment supports a wide range of microbial taxa without dominance by a few

species [54]. Higher evenness within microbial communities is often linked to enhanced ecosystem stability and

productivity [55]. This evenness may be attributed to the diverse plant life and landscape elements designed to

promote a harmonious coexistence of microbes. The notably high diversity observed in sample MIRAI8, a soil

sample from the Visionary Lab, further supports our hypothesis about the effectiveness of the landscape design

in enhancing microbial richness and evenness. This finding aligns with supports existing literature which

consistently suggests that soil, a critical component of landscape design, plays a crucial role in supporting a

diverse microbial ecosystem [56]. Similarly, a study by van der Heijden et al. (2008) found that plant diversity

enhances ecosystem productivity and stability, partly through its effects on soil microbial communities [57].

The samples outside the Visionary Lab exhibited lower microbial diversity, as evidenced by their alpha

diversity indices (Table 3). Several factors may contribute to this. Firstly, the lack of intentional landscape

design outside the exhibition means there are fewer plants and less environmental complexity to support a

diverse microbial community. As discussed, studies have shown that plant diversity and structural complexity

provide a range of niches and resources, which enhance microbial diversity. Without these elements, it is

expected that microbial communities will be less diverse. Secondly, human activity in these areas might

introduce human-associated microbes while reducing environmental microbes - something we can confirm

through our taxonomic assessment (Figure 2). These high-traffic areas can experience frequent cleaning and

disinfection, which can diminish microbial diversity by selectively eliminating certain taxa [58]. Additionally,

our taxonomic assessment revealed that there were few human-associated microbes which appeared to dominate

in the non-Visionary-Lab environments, leading to lower overall diversity (reflected in the observed features

metric). Thirdly, the environmental conditions outside the Visionary Lab might not be as conducive to microbial

diversity. Factors such as lower humidity, less organic matter, and fewer plant-derived resources can limit

microbial growth and diversity [59].

The beta diversity analysis provides significant insights into the spatial organisation and transfer

mechanisms of microbial communities within the museum. The observed clustering of microbial communities,

corresponding to their sampling locations, suggests the influence of spatial proximity on microbial diversity

(Figure 1 and Figure 3). These findings highlight the possible importance of environmental features and the

physical layout in shaping microbial ecosystems. Mechanisms of microbial transfer may play a crucial role in

explaining the formation of these distinct clusters. Human activity is a primary driver of microbial dispersal

within built environments [6]. As visitors move through the museum, they inadvertently carry and deposit

microbes through skin, clothing, and respiratory emissions, contributing to microbial transfer [7] This human-

mediated transfer is particularly evident in areas with high foot traffic, such as the entrances and exits of

exhibits. For example, the clustering of samples MIRAI2 and MIRAI1 could potentially be attributed to the high

visitor traffic near the Visionary Lab entrance, facilitating the homogenization of microbial communities in

these regions (Figures 1 & 3). Surface contact represents another critical mechanism [6]. Surfaces such as

exhibit materials, floors, and interactive displays serve as reservoirs for microbial communities. Visitors

touching these surfaces enhance microbial exchange, particularly in areas with interactive exhibits. Human

activity is a significant vector for microbial transfer, as people can carry microbes on their skin, clothing, and

personal items [6]. The interaction between visitors and the plant-rich environment of the Visionary Lab likely

facilitates the transfer and dissemination of diverse microbial species. Previous studies have shown that

environmental conditions, such as moisture levels, nutrient availability, and vegetation cover, can significantly

impact microbial diversity and composition [60, 61]. The clustering of samples within specific exhibits suggests

that the unique conditions and interactions within these microenvironments drive distinct microbial

assemblages. Although in this given study we did not collect data on airflow or ventilation, it is known that

airborne dispersal can also contribute to microbial transfer from different surfaces - for example transfer from

soil samples to non-soil samples [62]. Soil and plant surfaces themselves are reservoirs of microbial life, and

their interaction with the surrounding environment can lead to the spread of microbes, where air currents and

ventilation systems lift microbes from soil into the area and deposit them onto nearby surfaces [63]. This could

be the case in the Visionary Lab, where the strategic placement of environmental features, such as plants, create

microhabitats that support diverse microbial populations, contributing to the unique microbial ecosystems

observed within the Visionary Lab samples.

The clustering patterns observed in the beta diversity analysis are consistent with the theory that

proximity to curated environmental features significantly influences microbial composition [64]. This

correlation reinforces the idea that strategic environmental design can be leveraged to shape microbial

ecosystems in desired ways. By understanding the mechanisms of microbial transfer, such as human activity,

airflow, and surface contact, we can optimise the design of built environments to promote beneficial microbial

communities. In conclusion, the distinct community structures revealed by the beta diversity analysis highlight

the importance of spatial organisation and environmental design in shaping microbial ecosystems within the

museum.

Despite the valuable insights gained, it must be acknowledged that due to the small sample size (n=9),

of which only 5 were sequenced by shallow shotgun methods, it may not have been possible to fully capture the

variability and complexity of microbial communities within the Visionary Lab and the Miraikan museum. A

larger sample size would provide a more comprehensive understanding of the microbial diversity in these

environments and allow for further statistical tests to validate the significance of findings.

Virulence factors, antimicrobial resistance, and metabolic function profiles

The comparative analysis of virulence factors and antimicrobial resistance between samples revealed

no clear consistent patterns, highlighting the complexity of microbial community dynamics on human health

(Table 4). The prevalence of virulence factors in sample MIRAI1 suggests a baseline level of microbial

pathogenicity in the external environment, whilst the absence of antibiotic resistance genes in this sample

indicates a potentially lower risk of antibiotic-resistant infections in these settings. The significant presence of

virulence factors in MIRAI3 suggests that microbes in this controlled environment have adapted mechanisms to

enhance their survival and pathogenicity. ExbD, for instance, is involved in nutrient uptake and energy

transduction, which could enhance microbial fitness in nutrient variable conditions. It has been identified before

from *Verrucomicrobiota* bacterium**,** which in turn has been commonly isolated from environmental

environments such as marine and soil locations, as well as the human microbiome such as human faeces [65].

This makes sense in the context of our results, given it was found in sample MIRAI3 which is a sample taken

from a surface- so lots of contact with human environments, but close to the planting area- so transfer from the

soil samples. The detection of antibiotic resistance genes, such as *RbpA* and *Mycobacterium* in this sample also

raises concerns about the potential for these environments to harbour and disseminate resistant microbes. *RbpA*

is known to provide resistance to rifampicin, a critical antibiotic for treating tuberculosis [66]. This gene

indicates that even in curated environments, there is a risk of developing and spreading antimicrobial resistance,

likely due to selective pressures and microbial interactions. MIRAI 7, collected close to MIRAI 3 from the soil

around the planting area, shared several virulence factors and the RbpA resistance gene with MIRAI3. This

suggests that the landscape design, which includes soil and plant materials, creates microhabitats that support

diverse microbial communities with both pathogenic and resistant strains. Phosphoribosyl-AMP cyclohydrolase,

detected in MIRAI7, is crucial for nucleotide biosynthesis and microbial proliferation, indicating that soil

microbes are well-equipped for growth and survival in these enriched environments [67]. The absence of

detectable virulence factors and antibiotic resistance genes in MIRAI8 could be due to localised differences in

soil composition or microbial interactions that limit the growth of pathogenic and resistant strains.

A small number of studies [68–71] have concentrated on public settings in built-up environments, but

the majority of these studies have demonstrated that AMR resistance is highly abundant in public settings across

various species [72]. One study revealed that highly maintained environments (subject to intensive cleaning and

disinfection processes) exhibit a higher diversity of antimicrobial resistance genes and reduced microbial

diversity [73]. Based on this, we hypothesised that the soil samples taken from the planting area within the

visionary lab would show the least AMR genes and virulence factors (due to its high microbial diversity) and

those with more human contact (samples taken from surfaces) would have more AMR genes and virulence

factors. However, overall, these results suggest that the influence of landscape design on virulence factors and

antimicrobial resistance is complex and does not follow a clear, consistent pattern. This may be due to our

study's limited sample size and scope, which does not fully capture the broader patterns and interactions within

the microbial communities. PROKKA and SHORTBRED are tools which depend on the quality and

completeness of the reference databases and the contigs used in the analysis. Incomplete or fragmented contigs

may lead to underestimation or misidentification of genes, affecting the results. The variability observed

emphasised the need for further research to understand the multifaceted interactions between environmental

conditions, microbial communities, and human activity in shaping the presence and distribution of virulence

factors and antimicrobial resistance genes in built environments.

The distinct metabolic profiles highlight how controlled landscape design can shape microbial

communities, fostering unique ecological interactions and functions (Figure 4). Samples which share genes that

code for common functions can be attributed to these functions' broad applications in various environmental and

human settings. Within the group of functions which exhibit notable variation within the samples (function

present in 30-70% of samples), the most common pattern is for the function to be present in MIRAI3, MIRAI7,

and MIRAI8 and not present within MIRAI1 and MIRAI2. For example, the complex carbon degradation

category follows this pattern. This distribution pattern of these genes across the samples can be explained by the

functions within the category being central in an environment with plant material (e.g., cellulose-degrading,

hemicellulose-debranching, etc…) [74]. As a result, genes which code for these functions are more likely to be

found within the soil/plant samples (MIRAI7 and MIRAI8). The reason these genes are also found to be present

in sample MIRAI3, could be attributed to microbial transfer over short distances- as MIRAI3 was taken from a

location close to the planting area where MIRAI7 and MIRAI8 were taken. This same explanation can be

applied to virtually all the remaining functions within this variation category within samples. Fermentation is

another function that demonstrates significant variation across samples. Genes associated with fermentation are

predominantly found in plant/soil samples (MIRAI7 and MIRAI8) and are largely absent in samples from

human-associated environments (MIRAI1 and MIRAI2).

This analysis focuses solely on the presence or absence of genes coding for specific functions, which

presents certain limitations. For example, in the context of 'nitrogen cycling', the pattern of presence or absence

of genes varies across the five different functions, leading to the question of why one sample contains one type

of gene while lacking genes for closely related functions in the same nitrogen cycle. These ambiguities may

arise from biological factors or contamination, complicating interpretation. Thus, more comprehensive analyses

considering specific gene abundance are needed better to understand the genetic landscape and its functional

implications. Nonetheless, the broad application of certain metabolic functions across various settings

underscores their ecological importance, while other variability points to localised environmental influences.

<u>Conclusion</u>

This study comprehensively outlines the impact of green infrastructure on microbial diversity in the

context of the Visionary Lab compared to other areas of the Miraikan museum. The curated landscape design,

incorporating diverse plant life, supports a more complex and balanced microbial community, contributing to

enhanced microbial richness, evenness, and ecosystem multifunctionality. While the Visionary Lab samples

exhibited higher microbial diversity, the study did not reveal consistent patterns in virulence factors or

antimicrobial resistance genes. This variability underscores the complexity of microbial community dynamics

and the influence of localised environmental conditions. Future studies should increase the sample size and

variety of sampling locations within and outside the Visionary Lab and employ further advanced sequencing

techniques (such as deep metagenomic sequencing) to enhance the robustness and generalizability of results.

Additionally, exploring the impact of different environmental variables, such as humidity, temperature, and light

exposure, on microbial communities, and conducting longitudinal studies to track microbial changes over time

would further elucidate the dynamics of microbial ecosystems. Using this comprehensive approach to build

upon the findings from this study would contribute to developing evidence-based guidelines for creating

microbial-friendly and health-promoting indoor environments, aligning with the broader goal of sustainable

urban design.

<u>Declarations</u>

<u>Ethics approval and consent to participate:</u>

Not Applicable.

<u>Consent for publication</u>

Not Applicable

<u>Competing interests</u>

K.I. is a board member at BIOTA Inc., Tokyo, Japan. M.M. is employed by BIOTA Inc. as a research and

development intern.

<u>Funding</u>

<u>Authors' contributions</u>

K.I. designed the study and analysed the 16s rRNA Amplicon data. M.M. interpreted the 16s rRNA Amplicon

data and analysed and interpreted the shallow shotgun data. M.M was the major contributor in writing the

manuscript. All authors read and approved the final manuscript.

<u>Acknowledgements</u>

We would like to extend our sincere gratitude to Marin Yamaguchi and Keisuke Otsuka for their diligent work

in collecting the samples. Our thanks go to the National Museum of Emerging Science and Innovation

(Miraikan) for hosting the "Visionary Lab; Microbes Actually Are All Around" exhibition between April 2022

and   August   2023   and   for   allowing   us   to   collect   samples   within   their   facility

(https://www.miraikan.jst.go.jp/resources/visionarieslab/visionaries-archive-003.html). We also thank the Japan

Science and Technology Agency (JST) for their collaboration and support in the Visionary Lab exhibition. This

study would not have been possible without their collaboration and support.

Availability of Data and Materials

The datasets generated through 16S rRNA amplicon and metagenomic sequencing are available and deposited

in the DDBJ Sequence Read Archive (DRA) database under accession numbers DRR582054-DRR582069 and

BioProject PRJDB18223.

474
475 Figures and Tables

Figure 1: Illustration of the spatial layout of the Visionary Lab and the specific locations where microbial

samples were collected. Footprints and coloured paths indicate the movement flow within the exhibit.

Figure 2: Relative abundance of most abundant microbial genera across the samples from A) the museum environment in comparison to those from B) the soil samples (within the Visionary Lab).

Figure 3: Weighted unifrac Principal Coordinate Analysis (PCoA) for samples analysed by 16S rRNA amplicon sequencing. Samples are colour-coded by location and shaped according to surface type.

Figure 4: Heatmap illustrating the presence or absence of genes coding for various metabolic functions across samples sequenced by a shallow shotgun approach. The legend on the right categorises each function by its metabolic category, aiding in the interpretation of metabolic diversity and specialisation among the samples.

Table 1: Summary table regarding samples collected for genomic analysis, and data regarding the number of sequences processed for each sample using the two different sequencing methods. Some samples only have data for 16s rRNA amplicon sequencing as indicated by a dash in the corresponding shallow shotgun column.

Sample	Collection protocol	Surface type	Place	Reads processed	
				16s rRNA amplicon	Shallow shotgun
MIRAI1	Vacuum	Floor	Near the 'Earth'	114,111	21,916,203
MIRAI2	Vacuum	Floor	Visionary Lab	107,616	30,083,584

MIRAI3	Vacuum	Floor	Visionary Lab near the planting area	109,227	39,829,873
MIRAI4	Swab	Wall	Visionary Lab	121,281	-
MIRAI5	Swab	Desk	Visionary Lab	110,168	-
MIRAI6	Swab	Wall	Calculator and nature	117,977	-
MIRAI7	Swab	Soil	Visionary Lab planting area	104,243	21,768,309
MIRAI8	Swab	Soil	Visionary Lab planting area	82,274	27,126,330
MIRAI9	Swab	Desk	Future backward thinking	115,917	-

498 **Table 2:** Table displaying the relative abundance (%) of the 25 most abundant microbial species identified in
499 samples from museum surfaces and soil samples. The 'remainder' row represents the cumulative relative
500 abundance of all other species not listed in the top 25 for each sample.

Museum microbiome					Soil microbiome		
25 most abundant species	Relative abundance				25 most abundant species	Relative abundance	
	MIRAI1	MIRAI2	MIRAI3			MIRAI7	MIRAI8
Moraxella osloensis	16.88	7.19	0.02		*Pseudonocardia sp. DSM 110487*	3.94	0.33
Cutibacterium acnes	14.03	9.35	0.04		*Nocardioides sp. NBC_00368*	1.95	0.07
Staphylococcus aureus	0.26	10.79	0.00		*Streptomyces bathyalis*	1.56	0.03
Micrococcus luteus	9.35	1.44	0.02		*Variovorax paradoxus*	0.48	1.02
Nocardioides sp. NBC_00368	0.26	6.47	2.64		*Pseudonocardia sp. HH130630-07*	0.15	1.25
Brevibacterium casei	7.79	0.00	0.06		*Mycolicibacterium smegmatis*	1.30	0.07
Dermacoccus nishinomiyaensis	5.19	0.72	0.02		*Kribbella sp. NBC_00482*	1.19	0.03
Janibacter sp. CX7	0.78	4.32	0.50		*Pseudonocardia sp. AL041005-10*	0.15	1.05
Corynebacterium tuberculostearicum	4.68	0.72	0.00		*Pseudoxanthomonas suwonensis*	0.93	0.26
Janibacter limosus	0.52	4.32	0.32		*Stackebrandtia nassauensis*	1.10	0.03
Staphylococcus hominis	4.16	0.72	0.00		*Mycolicibacterium moriokaense*	1.02	0.07
Agrococcus carbonis	0.26	4.32	0.11		*Streptomyces xanthii*	1.04	0.03
Sphingomonas sp. FARSPH	3.12	1.44	0.06		*Ramlibacter tataouinensis*	0.13	0.89

Janibacter terrae	0.52	3.60	0.26		*Enhydrobacter sp.*	0.26	0.69
Pseudonocardia sp. DSM 110487	0.00	1.44	2.26		*Pseudonocardia sp. HH130629-09*	0.11	0.79
Brachybacterium sp. SGAir0954	0.26	2.16	1.18		*Sorangium cellulosum*	0.37	0.46
Janibacter melonis	0.52	2.16	0.88		*Streptomyces nigra*	0.78	0.03
Nocardioides sp. JQ2195	0.00	2.16	1.38		*Usitatibacter rugosus*	0.09	0.72
Corynebacterium afermentans	3.38	0.00	0.00		*Variovorax sp. HW608*	0.35	0.46
Dermacoccus sp. PAMC28757	2.60	0.72	0.02		*Luteitalea sp. TBR-22*	0.17	0.62
Janibacter sp. A1S7	0.26	2.88	0.07		*Croceibacterium sp. D39*	0.37	0.43
Brachybacterium sp. NBEC-018	0.00	2.16	0.97		*Lysobacter soli*	0.56	0.23
Janibacter indicus	1.30	1.44	0.19		*Luteitalea pratensis*	0.15	0.56
Mycolicibacterium smegmatis	0.00	0.72	1.79		*Kribbella sp. NBC_00709*	0.67	0.03
Nocardioides sp. zg-1228	0.00	1.44	0.95		*Lacunisphaera limnophila*	0.26	0.43
Remainder	23.90	27.34	86.26		Remainder	80.92	89.43

504 **Table 3:** Alpha diversity metrics for all samples. Metrics include Shannon diversity (measure of species

505 richness and evenness), observed features (number of distinct features observed), Faith's Phylogenetic Diversity

506 (measure of biodiversity that incorporates phylogenetic differences between species), and Pielou's Evenness

507 (measure of how evenly the species are distributed).

	Alpha diversity metrics			
Sample	Shannon diversity	Observed features	Faith PD	Pielou evenness
MIRAI1	7.893	1,092	69.428	0.782
MIRAI2	9.281	1,597	112.933	0.872
MIRAI3	9.736	1,694	105.363	0.908
MIRAI4	9.955	1,616	88.141	0.934
MIRAI5	9.368	1,556	85.999	0.883
MIRAI6	4.877	384	38.279	0.568

MIRAI7	9.991	1,702	86.597	0.568
MIRAI8	10.488	2,217	150.135	0.944
MIRAI9	5.702	524	44.321	0.631

510 **Table 4:** Virulence factors and antibiotic resistance genes identified in samples sequenced by shallow shotgun

511 sequencing. 'Family' represents the database family and gene identifier, 'count' is the total abundance count of

512 each gene, 'hits' is the number of times the gene was detected, and 'marker length' is the length of the gene

513 marker.

	Virulence Factor Database					Complex Antibiotic Resistance Database			
Sample	Family	Count	Hits	Marker length		Family	Count	Hits	Marker length
MIRAI1	VFDB|VFG038025(gi_384141332)	350.086	1	42		-			
	virulence|961|vfid|962|vsiid|961|ssid|transposase	231.307	1	50					
MIRAI2	Victors|gi|15926930|ref|NP_374463_1|	468.345	1	27		-			
	virulence|18940|vfid|42469|vsiid|63306|ssid|hypothetical	20.314	1	168					
MIRAI3	VFDB|VFG030641(gi_406029600)	140.943	3	30		gb|AAK44936_1|ARO_3003395|Mycobacterium	368.619	9	102
	VFDB|VFG031723(gi_15839882)	84.071	2	23		gb|ADV91011_1|ARO_3000245|RbpA	199.669	1	48
	VFDB|VFG022109(gi_120401443)	72.607	2	22					
	VFDB|VFG022671(gi_333990322)	44.371	1	28					
MIRAI7	VFDB|VFG028663(gi_433629379)	98.4040	1	24		gb|ADV91011_1|ARO_3000245|RbpA	418.201	1	48
	VFDB|VFG009731(gi_118618107)	176.0	1	14					

		85							
	VFDB|VFG031723(gi_158 39882)	17 6.0 85	2	23					
MIRAI 8	-					-			

References

1. Hu J, Amor DR, Barbier M, Bunin G, Gore J. Emergent phases of ecological diversity and dynamics mapped in microcosms. Science. 2022;378:85–9.

2. Huelsmann M, Ackermann M. Community instability in the microbial world. Science. 2022;378:29–30.

3. Nath A. The Significant Role of Microbial Diversity on Earth's Ecosystems. Applied Microbiology: Open Access. 2023;9:0–1.

4. Bertola M, Ferrarini A, Visioli G. Improvement of Soil Microbial Diversity through Sustainable Agricultural Practices and Its Evaluation by -Omics Approaches: A Perspective for the Environment, Food Quality and Human Safety. Microorganisms. 2021;9.

5. Hou K, Wu Z-X, Chen X-Y, Wang J-Q, Zhang D, Xiao C, et al. Microbiota in health and diseases. Signal Transduct Target Ther. 2022;7:135.

6. Gilbert JA, Stephens B. Microbiology of the built environment. Nat Rev Microbiol. 2018;16:661–70.

7. National Academies of Sciences, Engineering, and Medicine, National Academy of Engineering, Division on Engineering and Physical Sciences, Health and Medicine Division, Division on Earth and Life Studies, Board on Infrastructure and the Constructed Environment, et al. The Built Environment and Microbial Communities. In: Microbiomes of the Built Environment: A Research Agenda for Indoor Microbiology, Human Health, and Buildings. National Academies Press (US); 2017.

8. Ito K. Understanding the characteristics of the microbiomes of the built environment and their effects on the human body. Indoor Environment. 2023;**26**(1):29–42

9. Bruno A, Fumagalli S, Ghisleni G, Labra M. The Microbiome of the Built Environment: The Nexus for Urban Regeneration for the Cities of Tomorrow. Microorganisms. 2022;10:2311.

10. Mahnert A, Moissl-Eichinger C, Berg G. Microbiome interplay: plants alter microbial abundance and diversity within the built environment. Front Microbiol. 2015;6:153475.

11. Rupprecht CDD, Vervoort J, Berthelsen C, Mangnus A, Osborne N, Thompson K, et al. Multispecies sustainability. Global Sustainability. 2020;3:e34.

12. Funabashi M. Human augmentation of ecosystems: objectives for food production and science by 2045. NPJ Sci Food. 2018;2:16.

13. Niwa R, Chen D, Seong Y ah, Jo K, Ito K. Direct contact of fermented rice bran beds promotes food-to-hand transmission of lactic acid bacteria. FEMS Microbiol Lett. 2023;370:fnad120.

14. Ito K, Niwa R, Kobayashi K, Nakagawa T, Hoshino G, Tsuchida Y. A dark matter in sake brewing: Origin of microbes producing a Kimoto-style fermentation starter. Front Microbiol.

2023;14:1112638.

15. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol.
2019;37:852–7.

16. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA
gene database project: improved data processing and web-based tools. Nucleic Acids Res.
2012;41:D590–6.

17. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing
taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier
plugin. Microbiome. 2018;6:1–17.

18. FelixKrueger/TrimGalore: v0.6.10 - add default decompression path.
https://doi.org/10.5281/zenodo.7598955.

19. Raney BJ, Barber GP, Benet-Pagès A, Casper J, Clawson H, Cline MS, et al. The UCSC Genome
Browser database: 2024 update. Nucleic Acids Res. 2024;52:D1082–8.

20. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–
9.

21. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for
large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.
2015;31:1674–6.

22. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol.
2019;20:1–13.

23. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, et al. METABOLIC: high-throughput
profiling of microbial genomes for functional traits, metabolism, biogeochemistry, and community-
scale functional networks. Microbiome. 2022;10:1–22.

24. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30:2068–9.

25. Kaminski J, Gibson MK, Franzosa EA, Segata N, Dantas G, Huttenhower C. High-Specificity
Targeted Functional Profiling in Microbial Communities with ShortBRED. PLoS Comput Biol.
2015;11:e1004557.

26. Website. https://huttenhower.sph.harvard.edu/shortbred/.

27. VFDB: Virulence Factors of Bacterial Pathogens. http://www.mgc.ac.cn/VFs/main.htm. Accessed
27 May 2024.

28. McArthur AG, Wright GD. The Comprehensive Antibiotic Resistance Database.
https://card.mcmaster.ca/. Accessed 27 May 2024.

29. Adams RI, Bateman AC, Bik HM, Meadow JF. Microbiota of the indoor environment: a meta-
analysis. Microbiome. 2015;3:1–18.

30. Rozas M, Hart de Ruijter A, Fabrega MJ, Zorgani A, Guell M, Paetzold B, et al. From Dysbiosis
to Healthy Skin: Major Contributions of to Skin Homeostasis. Microorganisms. 2021;9.

31. D'Accolti M, Soffritti I, Bini F, Mazziga E, Mazzacane S, Caselli E. Pathogen Control in the Built
Environment: A Probiotic-Based System as a Remedy for the Spread of Antibiotic Resistance.
Microorganisms. 2022;10.

32. Cave R, Misra R, Chen J, Wang S, Mkrtchyan HV. Comparative Genomics Analysis
Demonstrated a Link Between Staphylococci Isolated From Different Sources: A Possible Public
Health Risk. Front Microbiol. 2021;12:576696.

33. Madsen AM, Moslehi-Jenabian S, Islam MZ, Frankel M, Spilak M, Frederiksen MW.
Concentrations of Staphylococcus species in indoor air as associated with other bacteria, season,
relative humidity, air change rate, and S. aureus-positive occupants. Environ Res. 2018;160:282–91.

34. Tan L, Grewal PS. Pathogenicity of Moraxella osloensis, a bacterium associated with the
nematode Phasmarhabditis hermaphrodita, to the slug Deroceras reticulatum. Appl Environ
Microbiol. 2001;67:5010–6.

35. Kubota H, Mitani A, Niwano Y, Takeuchi K, Tanaka A, Yamaguchi N, et al. Moraxella species
are primarily responsible for generating malodor in laundry. Appl Environ Microbiol. 2012;78:3317–
24.

36. Tan L, Grewal PS. Endotoxin activity of Moraxella osloensis against the grey garden slug,
Deroceras reticulatum. Appl Environ Microbiol. 2002;68:3943–7.

37. Zhang T, Liu M, Zhou D, Ma Z, Chen L, Wu D, et al. Environmental factors and particle size
shape the community structure of airborne total and pathogenic bacteria in a university campus. Front
Public Health. 2024;12:1371656.

38. Ahmed NM, Joglekar P, Deming C, NISC Comparative Sequencing Program, Lemon KP, Kong
HH, et al. Genomic characterization of the species complex, a ubiquitous member of the human skin
microbiome. bioRxiv. 2023. https://doi.org/10.1101/2023.06.16.545375.

39. Altonsy MO, Kurwa HA, Lauzon GJ, Amrein M, Gerber AN, Almishri W, et al. Corynebacterium
tuberculostearicum, a human skin colonizer, induces the canonical nuclear factor-κB inflammatory
signaling pathway in human skin cells. Immun Inflamm Dis. 2020;8:62–79.

40. Gupta M, Lee S, Bisesi M, Lee J. Indoor Microbiome and Antibiotic Resistance on Floor
Surfaces: An Exploratory Study in Three Different Building Types. Int J Environ Res Public Health.
2019;16.

41. National Academies of Sciences, Engineering, and Medicine, National Academy of Engineering,
Division on Engineering and Physical Sciences, Health and Medicine Division, Division on Earth and
Life Studies, Board on Infrastructure and the Constructed Environment, et al. Microorganisms in Built
Environments: Impacts on Human Health. In: Microbiomes of the Built Environment: A Research
Agenda for Indoor Microbiology, Human Health, and Buildings. National Academies Press (US);
2017.

42. Riahi HS, Heidarieh P, Fatahi☐Bafghi M. Genus Pseudonocardia: What we know about its
biological properties, abilities and current application in biotechnology. J Appl Microbiol.
2022;132:890–906.

43. Safaei N, Bunk B, Spröer C, Wink J, Mast Y. Complete Genome Sequences of Two Novel
Species from the Family Isolated from the Persian Gulf. Microbiol Resour Announc.
2021;10:e0091821.

44. Bruner-Montero G, Wood M, Horn HA, Gemperline E, Li L, Currie CR. Symbiont-Mediated
Protection of Leaf-Cutter Ants from the Entomopathogenic Fungus Metarhizium anisopliae. MBio.
2021;12:e0188521.

45. STREPTOMYCES. 1999;:2134–8.

46. Ma Y, Wang J, Liu Y, Wang X, Zhang B, Zhang W, et al. : "Specialists" for Hard-to-Degrade
Pollutants in the Environment. Molecules. 2023;28.

47. Walsh CM, Gebert MJ, Delgado-Baquerizo M, Maestre FT, Fierer N. A Global Survey of
Mycobacterial Diversity in Soil. Appl Environ Microbiol. 2019;85.

48. Straub D, Blackwell N, Fuentes AL, Peltzer A, Nahnsen S, Kleindienst S. Interpretations of
microbial community studies are biased by the selected 16S rRNA gene amplicon sequencing
pipeline. bioRxiv. 2019;:2019.12.17.880468.

49. Choi J, Yang F, Stepanauskas R, Cardenas E, Garoutte A, Williams R, et al. Strategies to improve
reference databases for soil microbiomes. ISME J. 2017;11:829–34.

50. La Reau AJ, Strom NB, Filvaroff E, Mavrommatis K, Ward TL, Knights D. Shallow shotgun
sequencing reduces technical variation in microbiome analysis. Sci Rep. 2023;13:1–8.

51. Hillmann B, Al-Ghalith GA, Shields-Cutler RR, Zhu Q, Gohl DM, Beckman KB, et al. Evaluating
the Information Content of Shallow Shotgun Metagenomics. mSystems. 2018;3.

52. Kitikidou K, Milios E, Stampoulidis A, Pipinis E, Radoglou K. Using Biodiversity Indices
Effectively: Considerations for Forest Management. Ecologies. 2024;5:42–51.

53. Maherali H, Klironomos JN. Influence of phylogeny on fungal community assembly and
ecosystem functioning. Science. 2007;316:1746–8.

54. Jost L. The Relation between Evenness and Diversity. Diversity . 2010;2:207–32.

55. Wittebolle L, Marzorati M, Clement L, Balloi A, Daffonchio D, Heylen K, et al. Initial
community evenness favours functionality under selective stress. Nature. 2009;458:623–6.

56. Delgado-Baquerizo M, Eldridge DJ, Ochoa V, Gozalo B, Singh BK, Maestre FT. Soil microbial
communities drive the resistance of ecosystem multifunctionality to global change in drylands across
the globe. Ecol Lett. 2017;20:1295–305.

57. van der Heijden MGA, Bardgett RD, van Straalen NM. The unseen majority: soil microbes as
drivers of plant diversity and productivity in terrestrial ecosystems. Ecol Lett. 2008;11:296–310.

58. Chen C-H, Tu C-C, Kuo H-Y, Zeng R-F, Yu C-S, Lu HH-S, et al. Dynamic change of surface
microbiota with different environmental cleaning methods between two wards in a hospital. Appl
Microbiol Biotechnol. 2017;101:771–81.

59. Gonzalez JM, Aranda B. Microbial Growth under Limiting Conditions-Future Perspectives.
Microorganisms. 2023;11.

60. Martiny JBH. Dispersal and the Microbiome. Microbe Wash DC. 2015;10:191–6.

61. Fierer N, Jackson RB. The diversity and biogeography of soil bacterial communities. Proc Natl
Acad Sci U S A. 2006;103:626–31.

62. Tong X, Leung MHY, Wilkins D, Lee PKH. City-scale distribution and dispersal routes of
mycobiome in residences. Microbiome. 2017;5:1–13.

63. Gupta A, Gupta R, Singh RL. Microbes and Environment. In: Principles and Applications of
Environmental Biotechnology for a Sustainable Future. Singapore: Springer Singapore; 2017. p. 43–
84.

64. Junkins EN, McWhirter JB, McCall L-I, Stevenson BS. Environmental structure impacts

microbial composition and secondary metabolism. ISME Commun. 2022;2:15.

65. Taxonomy browser (Verrucomicrobiota bacterium).
https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=2026799. Accessed
27 May 2024.

66. Hu Y, Morichaud Z, Chen S, Leonetti J-P, Brodolin K. Mycobacterium tuberculosis RbpA protein
is a new type of transcriptional activator that stabilizes the σ A-containing RNA polymerase
holoenzyme. Nucleic Acids Res. 2012;40:6547–57.

67. Fisher G, Pečaver E, Read BJ, Leese SK, Laing E, Dickson AL, et al. Catalytic Cycle of the
Bifunctional Enzyme Phosphoribosyl-ATP Pyrophosphohydrolase/Phosphoribosyl-AMP
Cyclohydrolase. ACS Catal. 2023. https://doi.org/10.1021/acscatal.3c01111.

68. Cave R, Misra R, Chen J, Wang S, Mkrtchyan HV. Whole genome sequencing revealed new
molecular characteristics in multidrug resistant staphylococci recovered from high frequency touched
surfaces in London. Sci Rep. 2019;9:9637.

69. Kang K, Ni Y, Li J, Imamovic L, Sarkar C, Kobler MD, et al. The Environmental Exposures and
Inner- and Intercity Traffic Flows of the Metro System May Contribute to the Skin Microbiome and
Resistome. Cell Rep. 2018;24:1190-1202.e5.

70. Roberts MC, Soge OO, No D. Comparison of Multi-Drug Resistant Environmental Methicillin-
Resistant Staphylococcus aureus Isolated from Recreational Beaches and High Touch Surfaces in
Built Environments. Front Microbiol. 2013;4:74.

71. Shen C, Feng S, Chen H, Dai M, Paterson DL, Zheng X, et al. Transmission of mcr-1-Producing
Multidrug-resistant Enterobacteriaceae in Public Transportation in Guangzhou, China. Clin Infect Dis.
2018;67 suppl_2:S217–24.

72. Surveillance and prevalence of antimicrobial resistant bacteria from public settings within urban
built environments: Challenges and opportunities for hygiene and infection control. Environ Int.
2021;157:106836.

73. Mahnert A, Moissl-Eichinger C, Zojer M, Bogumil D, Mizrahi I, Rattei T, et al. Man-made
microbial resistances in built environments. Nat Commun. 2019;10:1–12.

74. Yu T, Wu W, Liang W, Wang Y, Hou J, Chen Y, et al. Anaerobic degradation of organic carbon
supports uncultured microbial populations in estuarine sediments. Microbiome. 2023;11:81.

75. Vassileva M, Malusà E, Sas-Paszt L, Trzcinski P, Galvez A, Flor-Peregrin E, et al. Fermentation
Strategies to Improve Soil Bio-Inoculant Production and Quality. Microorganisms. 2021;9.

76. Fischbach MA, Sonnenburg JL. Eating for two: how metabolism establishes interspecies
interactions in the gut. Cell Host Microbe. 2011;10:336–47.

703