1   **Title Page**
2
3   A trait-based metric to prioritise the conservation of functionally
4   irreplaceable species at risk of extinction
5

6   Ceri Webster[1,2+], Joanna Barker[3], David Curnick[2], Matthew Gollock[3], James Hansford[2,4],
7   Michael Hoffmann[3], Thomas E. Martin[5,6], Thomas J. Matthews[7,8], Nathalie Pettorelli[2], Joseph
8   A. Tobias[1], Samuel T. Turvey[2], Patrick A. Walkden[1], Jiaqi Wang[9], Joseph P. Wayman[7], James
9   Rosindell[*1], Rikki Gumbs[*2]
10
11   [*]joint senior authors
12   [+]corresponding author (ceri.webster@ioz.ac.uk)
13
[1]Department of Life Sciences, Silwood Park Campus, Imperial College London, Buckhurst
Road, Ascot, Berkshire, SL5 7PY, UK
[2]Institute of Zoology, Zoological Society of London, London, NW8 7LS, UK
[3]Conservation and Policy, Zoological Society of London, London, NW1 4RY, UK
[4]Department of Earth Sciences, University College London, London, WC1E 6BS, UK
[5]Operation Wallacea, Wallace House, Old Bolingbroke, Lincolnshire, PE23 4EX, UK
[6] Department of Biological Sciences, Simon Fraser University, Burnaby, BC, Canada
[7]School of Geography, Earth and Environmental Sciences and Birmingham Institute of Forest
Research, University of Birmingham, Birmingham B15 2TT, UK
[8]CE3C – Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity
Group / CHANGE – Global Change and Sustainability Institute and Universidade dos Açores –
Faculty of Agricultural Sciences and Environment; PT-9700-042, Angra do Heroísmo, Açores,
Portugal.
[9]School of Biological Sciences, The University of Hong Kong, Hong Kong SAR

28
29
30

31   **Abstract**
32
Robust species-level methods for quantifying ecological differences have yet to be
incorporated into conservation strategies. Here, we present a conservation prioritisation
approach that integrates species trait data and extinction risk to quantify the contribution of
individual species to overall functional diversity. The Functionally Irreplaceable with Risk of
Extinction (FIRE) metric directs conservation action to species whose extinction is expected
to result in significant losses of functional diversity. We applied our framework to sets of
species at the global scale. First we assessed the world's birds, highlighting congruent and
divergent priorities identified by trait-based and phylogenetic approaches. Second, we
applied FIRE to the world's sharks, exploring the impact of imputed traits on prioritisation
robustness. For birds and sharks, we show that prioritising by functional irreplaceability is an
effective strategy to conserve exploited species. The FIRE metric provides a robust tool to
facilitate the incorporation of functional diversity into conservation policy and practice,
revealing species that may be overlooked by existing approaches.
46
47
48

1. Introduction

The biodiversity crisis is driving species declines and losses across all biomes, threatening the functioning of healthy ecosystems and the services they provide[1]. Efforts to mitigate these declines require strategic decision-making and often utilise established metrics of risk or priority, such as conservation status[2], taxonomic diversity[3], endemism[4], climate vulnerability[5,6] and evolutionary history[7]. Despite increasing interest in the variation of functional traits between species (i.e. functional diversity[8–10]), relatively little progress has been made at incorporating this into conservation decision-making[11]. For example, Goal A of the United Nations Convention on Biological Diversity (CBD) Kunming-Montreal Global Biodiversity Framework (GBF) focuses on maintaining, restoring and enhancing biodiversity to improve functions and services[12], yet the accompanying monitoring framework contains no means for tracking progress towards recovery of ecological integrity and functional diversity[13].

Measuring the functional diversity of species assemblages can help to reveal how extinction will impact ecosystem functions and services[14], providing an estimate of the ecological role or functionality of different species within a given assemblage, and the uniqueness or redundancy of that role in comparison to related or co-occurring taxa. Recent studies suggest that loss of species with functionally unique traits will drive functional homogenisation and erosion of ecosystem processes[15,16]. However, current prioritisation metrics that consider taxonomic or phylogenetic information alone do not adequately capture functional diversity[17,18].

Advances in metrics that prioritise species for conservation based on phylogenetic diversity (the sum of phylogenetic branch lengths connecting a set of taxa[19]) have produced approaches that are based on the established conceptual framework of averting expected losses of the most imperilled biodiversity[20–24]. For example, the updated Evolutionarily Distinct and Globally Endangered (EDGE) metric, based on the EDGE2 protocol[20], prioritises species based on the amount of threatened phylogenetic diversity their conservation is expected to secure. In contrast, existing approaches to integrate functional traits into conservation prioritisation metrics have largely been based on the original EDGE metric[7], which weighted a measure of distinctiveness by an index of extinction risk in an *ad hoc* manner[25] (though see Pavione[26]). The 'EcoDGE'[27] metric is an adaptation of the original EDGE metric that calculates the 'distinctiveness' component of the metric using dendrograms representing functional diversity rather than phylogenetic trees. The 'FUSE'[28] metric is another adaptation of the original EDGE calculation, whereby an index of priority is created by summing two measures; functional uniqueness[29] and functional specialisation[29], both of which are independently weighted by an ordinal ranking of extinction risk. However, growing focus on the maintenance of functionally diverse systems for conservation[8,9], the increasing availability of trait data across the tree of life[18,30], and established methods to quantify functional diversity[31] provide the opportunity to develop a more clearly mechanistic prioritisation approach to guide the conservation of functionally irreplaceable species.

Here, we detail a method that prioritises functionally irreplaceable and threatened species for conservation based on their expected contributions to trait space, given the landscape of

extinction risk across the entirety of trait space. We present the Functionally Irreplaceable
with Risk of Extinction (FIRE) metric, which builds on previous approaches[26,28,32] to quantify
the expected loss of functional diversity that can be averted through species-based
conservation action. We then provide two case studies illustrating the utility of our new
approach. The first applies FIRE to the world's birds, using a comprehensive species-level
trait dataset for over 10,000 species[33] to highlight unique and convergent conservation
priorities for maintaining avian functional and phylogenetic diversity. For the second, we
apply FIRE to 557 of the world's shark species, a smaller taxonomic group with more patchy
data, allowing us to explore how imputation of missing data influences the setting of
conservation priorities using the FIRE approach. We then use functional trait data for birds
and sharks to examine whether species threatened by hunting, collecting, and fishing (i.e.
exploitation) are particularly functionally or evolutionarily irreplaceable, and explore how
conservation strategies to prioritise irreplaceable species perform at safeguarding diversity
at risk from this threat. FIRE builds on existing prioritisation frameworks to provide
functional indices to inform conservation and policy.

111
112

2. Results

2.1 Functionally Irreplaceable with Risk of Extinction (FIRE) metric

The FIRE metric aims to identify and prioritise threatened species whose extinction is
expected to represent disproportionately large losses of unique functional trait space. FIRE
estimates the functional irreplaceability of a species by calculating its expected future
unique contribution to overall trait space given the extinction risk and functionality of all
other species in the same trait space. The trait space from which FIRE is calculated requires
independent trait axes to quantify functional distances between species (Fig. 1, see
Methods). In the case of continuous traits, including morphometric data, these independent
axes can be generated using ordination methods (PCA or PCoA, depending on data type).
Trait probability densities (TPDs) are then used to estimate a probabilistic trait space based
on the results of the ordination (PCA or PCoA)[34]. The expected loss of overall trait space is
then calculated using a large number (n ≥ 1000) of extinction iterations where, for each
iteration, species are removed from trait space with probabilities derived from their IUCN
Red List categories (Fig. 1, see Methods). For each species, a distribution of functional
irreplaceability scores is then calculated by quantifying the amount of trait space expected
to be uniquely occupied by the species if it were to avoid extinction in each of the *n*
extinction iterations. The FIRE of a species is then calculated as the product of its functional
irreplaceability and probability of extinction across the distribution of *n* iterations, and
represents the expected loss of trait space that could be averted through the conservation
of each species (i.e., if the species does not go extinct) (Fig. 1). To facilitate comparability
between functional irreplaceability scores calculated from different trait spaces (for a fixed
taxonomic group), we present the scores as the percentage of total trait space for which we
expect a species to be responsible in the future.

140
Fig. 1| Calculating the Functionally Irreplaceable with Risk of Extinction (FIRE) metric. Top panel:
Constructing a multivariate trait space requires trait data to quantify functional distances between species.
Probabilistic trait space is estimated from the functional distances between species and an estimated
variability using the TPD package (v1.1.0)[34]. Colours in trait space indicate high (red) or low (blue) probability of
occupancy. Middle panel: Post-extinction trait space is calculated by removing species predicted to go extinct
in a given iteration following an extinction scenario, whereby species become extinct or survive based on their
probability of extinction. White circles represent areas of trait space no longer occupied by species following
extinction. This step is repeated across 1000 iterations to derive a distribution of uncertainty in extinction
scenarios. IUCN Red List categories are converted to a continuous distribution of extinction probabilities,
following Gumbs et al. 2023[20] where the median of each Red List category was defined as; Critically
Endangered = 0.97, Endangered = 0.485, Vulnerable =0.2425, Near Threatened = 0.12125, and Least Concern =
0.060625. Species probability of extinction is randomly selected from the distribution of extinction
probabilities defined by their IUCN Red List categories. Species that were not assessed or had IUCN Red List
category listed as Data Deficient, were randomly sampled from the entire distribution of extinction
probabilities. Bottom panel: The functional irreplaceability of a species is calculated as the distinct contribution
of that species (which is immune to extinction during the calculation of its own functional irreplaceability
score) to post-extinction trait space. In the example shown there are five pixels in discretised trait space that
will in future only be covered by the focal species, if the focal species is protected from extinction. The FIRE
value of a species is then calculated as the product of its functional irreplaceability and probability of

extinction. The calculation is repeated 1000 times to capture the distribution of outcomes from the probability
of extinction. Image credits given in supplementary material.
162
163 ### 2.2 Case study 1: FIRE for the world's birds
164
165 Birds represent an ideal taxonomic group for testing the FIRE metric, given their long-
166 standing role in conservation research[35], the availability of comprehensive, high-resolution
167 trait data[33,36], and a robust literature base linking traits to ecological functions[37].
168
2.2.1 Trait space summary
To quantify the trait space of the world's birds (11,005 species), we selected eight
morphological traits that have been shown to provide accurate information on the
functional role and trophic status of birds at the global scale[33]: total beak length (from the
tip to the skull), beak length to the nares, beak width and depth, wing length, secondary
length (length from the carpal joint to the first of the wings secondary feathers), tail length
and tarsus length. We found consistently weak correlations between morphological traits
and extinction risk. Tarsus length ($\rho = 0.160$, $p < 0.001$), beak depth ($\rho = 0.157$, $p < 0.001$),
and secondary length ($\rho = 0.154$, $p < 0.001$) showed the strongest associations, though
effect sizes remained low. Other traits such as beak length (culmen: $\rho = 0.155$; nares: $\rho =$
0.147), wing length ($\rho = 0.149$), and beak width ($\rho = 0.133$) also showed significant but weak
positive associations (all $p < 0.001$). Tail length had the weakest relationship with extinction
risk ($\rho = 0.110$, $p < 0.001$). Overall, the low ρ values across traits suggest limited predictive
value of individual morphological variables for extinction vulnerability (Fig. S1).
183
We undertook a Principal Components Analysis (PCA) using these eight traits and extracted
the first three axes, which together represent 92% of the total variation. PCA axis 1 (PC1;
76% of total variation) represents an overall size axis, PC2 (10%) represents a trade-off
between beak length vs. tarsus, secondary and tail length, and PC3 (6%) represents a trade-
off between beak length vs. beak width and depth. We re-ran the analyses including body
mass as an additional trait. The functional irreplaceability ($r = 0.96$, $p < 0.001$) and FIRE ($r =$
0.97, $p < 0.001$) values were highly correlated between the two approaches and we only
discuss the results excluding body mass from herein. We used TPDs to account for the
probabilistic nature of species traits[38] (see Methods).
193
194 To estimate the predicted loss of trait space in the future, we projected species risk of
195 extinction onto global trait space (see Methods). Mapping extinction risk onto trait space
196 revealed marked variation in vulnerability (Fig. 2). We show that extinction in trait space is
197 likely to occur as a loss of distinct clusters rather than as a uniform contraction around the
198 periphery.
199

200
Fig. 2| Extinction risk of avian trait space. a-c, Trait spaces defined by three principal component axes (PC1,
PC2 and PC3) to show; the direction and weighting of the considered traits (black arrows), and projections of
the predicted erosion of trait space based on IUCN Red List assessments[2]. Light blue indicates low probability
of extinction and red indicates high probability of extinction. Dashed lines show examples of species position in
trait space, including California condor (*Gymnogyps californianus*), Brown pelican (*Pelecanus occidentalis*) and
Sword-billed hummingbird (*Ensifera enifera*). Traits include; total beak length (from the tip to the skull), beak
length to the nares, beak width and depth, wing length, secondary length (length from the carpal joint to the
first of the wings secondary feathers), tail length and tarsus length. Image credits given in supplementary
material.
210
2.2.2 Avian functional irreplaceability
The world's most functionally irreplaceable birds include several species widely considered
to be morphologically and ecologically unique, such as the Shoebill (*Balaeniceps rex*), Great
Hornbill (*Buceros bicornis*), Sunbittern (*Eurypyga helias*), kiwis (*Apteryx* spp.), and
Secretarybird (*Sagittarius serpentarius*; Fig. 3a). Other unique species are more obscure,
including the Coppery Thorntail (*Discosura letitiae*; DD on Red List) which is only known by a
single museum specimen of unknown provenance. Overall, there was a trend for
hummingbird species to be relatively highly ranked in the list of bird functional
irreplaceability values (discussed in more detail in Text S1). Conversely, the least functionally
irreplaceable species tend to be small passerines (e.g. warblers, tits; see full list in Data S1).
Eleven bird species do not overlap with any other species in trait space and thus receive
maximum irreplaceability scores (Data S1).
223
Threatened birds (i.e. Vulnerable [VU], Endangered [EN], and Critically Endangered [CR] on
the Red List) are significantly more functionally irreplaceable than non-threatened (i.e. Least
Concern [LC] and Near Threatened [NT]) birds (mean = 0.065 vs. 0.035, Welch's t-test: df =
1486.7, p < 0.001), corroborating earlier work highlighting this relationship[15,39,40]. Of the top
5% most functionally irreplaceable species (N = 551), 25.4% are threatened with extinction,
compared with 11.7% of all birds[2]. The highest priority FIRE species (i.e. highly functionally
irreplaceable species threatened with extinction) include the Extinct in the Wild Alagoas
Curassow (*Mitu mitu*), and the Critically Endangered Helmeted Hornbill (*Rhinoplax vigil*) and
Tristan Albatross (*Diomedea dabbenena*), alongside species such as the Secretarybird
(Endangered), Shoebill (Vulnerable), and Kākāpō (*Strigops habroptila*; Critically Endangered;
Fig. 3d; see full list in Data S1).
235

Fig. 3| Functionally irreplaceable birds. Correlations between (a) evolutionary distinctiveness and functional irreplaceability and (b) Evolutionarily Distinct and Globally Endangered (EDGE) and Functionally Irreplaceable with Risk of Extinction (FIRE) scores. Dotted lines represent the upper 5% of scores for each metric. Illustrations represent examples of species that fall within the top 5% of (c) evolutionary distinctiveness and functional irreplaceability (d) EDGE and FIRE. Bird species: 1. Sunbittern (*Eurypyga helias*), 2. Secretarybird (*Sagittarius serpentarius*), 3. Shoebill (*Balaeniceps rex*), 4. California Condor (*Gymnogyps californianus*). The top 25 highest ranked (c) functionally irreplaceable and (d) FIRE bird species and numbers represent the (c) functional irreplaceability and (d) FIRE rank of each species, where species with the same median value have the same rank. Boxplots represent the range, interquartile range and median of (c) functional irreplaceability and (d) FIRE scores calculated over 1000 iterations. IUCN Red List population trends (Popn) are denoted with arrows to represent increasing or decreasing population trends, a dash to represent stable population trends, and a question mark to represent unknown population trends. Species in bold text fall in the top 5% of both (c) functional irreplaceability and evolutionary distinctiveness scores and (d) FIRE and EDGE scores. Colours represent IUCN Red List categories (red = Critically Endangered, orange = Endangered, yellow = Vulnerable, light green = Near Threatened, green = Least Concern, grey = Data Deficient). Image credits given in supplementary material.

2.2.3 Unique and convergent FIRE priorities

To compare convergent and unique conservation priorities between FIRE and the phylogenetically-informed EDGE approach, we matched the bird species in our FIRE dataset with those with available EDGE data (N = 10971)[41]. There is a weak but significant positive correlation between functional irreplaceability and evolutionary distinctiveness for the

world's birds (ρ = 0.1, df = 10684, p < 0.001; Fig. 3a), though species that comprise
monotypic families have significantly higher functional irreplaceability than birds in general
(mean = 0.15 vs. 0.04, Welch's t-test: df = 33.021, p = 0.03). We identified species within the
highest 5% of FIRE scores, in a threatened or Extinct in the Wild Red List category, and with
available EDGE scores, as our set of priority birds (N = 316). Of these priority 5% species, 165
are also in the top 5% of EDGE scores, and include species such as the Helmeted Hornbill,
Secretarybird, Shoebill, Andean Condor (*Vultur gryphus*), and New Caledonian Owlet-
nightjar (*Aegotheles savesi*). We highlight these species as amongst the most important for
conservation, given their particularly high contribution to irreplaceable functional and
phylogenetic diversity globally (Fig. 3b, full list in Data S1).

Conversely, 102 (32.3%) of the top 5% FIRE species do not qualify as priority EDGE species
(i.e. above median EDGE with 95% confidence and threatened with extinction[20]), and thus
are not currently captured by phylogenetically-informed prioritisation. Further, 71 of the
top 5% FIRE species do not trigger any Key Biodiversity Areas (KBAs), and 45 (14.2%) are
underrepresented in the global KBA network (i.e. <8% of species area of habitat (AOH)
covered by KBAs; Lansley et al. 2025). While many of these species are nevertheless the
subject of targeted conservation actions (e.g. the Okarito Kiwi, *Apteryx rowi*[42]), additional
conservation efforts are likely required to promote and conserve some of the world's most
functionally irreplaceable birds. For example, FIRE scores tend to shift the focus of
conservation attention from less distinctive lineages towards maintaining interconnected
populations of widespread species with important ecological roles, such as Kori Bustard
Ardeotis kori and Greater Rhea *Rhea americana*.

2.3 Case study 2: FIRE prioritisation with incomplete data

Given the scale and urgency of the biodiversity crisis, it is impractical to limit conservation
research and efforts to only a select few taxonomic groups or regions with complete data [43–
45]. Major advances have been made to facilitate the inclusion of species lacking data into
assessments of extinction risk[46,47], conservation priorities[20,48], and functional diversity[49]. We
designed the FIRE metric to permit the inclusion of species lacking extinction risk data (Fig.
1). In addition, the metric can be calculated using imputed trait data[84]. To illustrate the
calculation of FIRE for a group with good – but incomplete – trait data coverage, we applied
the metric to the world's sharks (Selachimorpha). A substantial amount of trait and
extinction risk data are now available for the world's sharks, a group of high ecological
importance and significant conservation concern[28,50]. Given their elevated extinction risk
and functional roles in marine ecosystems, sharks have been the subject of previous trait-
based prioritisation efforts based on imputed data[32].

2.3.1 Imputing trait space
To quantify the trait space occupied by the world's sharks, we selected eight traits that
characterise a variety of functional guilds (maximum body length, maximum depth in the
water, trophic level, reproductive guild, habitat preference, growth ratio, and the first two
principal component axes from an analysis of body shape; see Methods) for 557 species
(>99% of all sharks). Taxonomic coverage of these eight traits was incomplete but high
overall (maximum body length: 96.1%, maximum depth: 87.6%, trophic level: 49.9%, body

shape variation PC1: 91.9%, body shape variation PC2: 91.9%, growth ratio: 61.4%,
reproductive mode: 84.2%, habitat preference: 100%; Table S1).

We used random forest models (missForest)[51] to impute missing trait values and attain
100% coverage for all traits. This method uses a random forest trained on the observed
values of the trait matrix to impute missing values. MissForest makes no prior assumptions
about the distributions of variables, does not incorporate phylogenetic relationships and can
be used to predict both continuous and categorical data[51]. Additionally, numerous studies
have found a high degree of accuracy when imputing traits using random forest methods
such as missForest[49,52,53]. Following imputation of all traits, we found generally weak or
inconsistent associations between traits and extinction risk. Shape PC1 and body length
showed the strongest relationships ($\rho = 0.41$ and $\rho = 0.44$, respectively; both $p < 0.001$).
Maximum depth ($\rho = -0.20$, $p < 0.001$) and growth ratio ($\rho = 0.16$, $p < 0.001$) showed modest
effects. Traits such as trophic level ($\rho = 0.06$, $p = 0.15$) and shape PC2 ($\rho = -0.09$, $p = 0.05$)
were weak predictors, with minimal explanatory power (Fig. S2). Categorical traits showed
significant differences in extinction score among groups for both habitat preference
(Kruskal-Wallis $\chi^2 = 53.54$, df = 7, $p < 0.001$) and reproductive guild (Kruskal-Wallis $\chi^2 =$
56.23, df = 3, $p < 0.001$). However, these differences were not clearly directional and did not
indicate a consistent trait gradient with increasing extinction risk (Fig. S2).

We defined trait space occupied by extant sharks by condensing the dissimilarity distance
matrix of compiled traits using a principal coordinate analysis (PCoA), and extracting the first
four axes, which together represent 45.28% of total variation. To assess how imputation
was distributed across trait space, we calculated the proportion of imputed data for each
species and projected this onto the first two PCoA axes (Fig. S3). We then summarised
imputation density across trait space by calculating the average proportion of imputed data
along both axes. This analysis revealed that imputation was relatively evenly distributed,
with no strong clustering of highly imputed species in any particular region of trait space.
Average imputation values along PC1 ranged from 0 to 0.26 (median: 0.18), and along PC2
from 0.05 to 0.32 (median: 0.15; Fig. S3). Species with a high proportion of imputed data
(more than 50% of data imputed) did not have high functionally irreplaceability scores (Fig.
4a).

To test the validity of our trait space derived from real and imputed data, we calculated
functional irreplaceability scores from a trait space constructed only with species with
complete trait data ('gapless dataset'; N = 176). We then generated artificially incomplete
data by randomly removing trait values at the observed proportion of incompleteness, then
imputing them back. We found a strong correlation between the median functional
irreplaceability scores from the gapless dataset and those from the imputed datasets ($\rho =$
0.78, $p < 0.001$; see Methods and Fig. 4b). The change in FIRE rankings of the 176 species in
the gapless dataset, when compared with their rankings in the full (imputed) dataset, was
consistent irrespective of rank ($\rho = 0.08$, df = 175, $p = 0.3$; Fig. 4c), indicating that the
inclusion of species with incomplete trait data did not affect priority rankings in a biased
manner. Indeed, 27 of 34 species (79%) remained in the highest 20% of rankings, and 13 of
18 (72%) remained in the top 10% across both datasets. However, the robustness of these
results is likely to be sensitive to the traits selected and the imputation method used,

particularly if trait gaps are non-randomly distributed or if certain traits disproportionately
influence ordination and irreplaceability scores[49,53].
354
355

356
357 **Fig. 4| Performance of functional irreplaceability estimations using imputed data for shark species.** (a) The
358 proportion of trait data imputed for species in our complete shark dataset (N = 557) and their functional
359 irreplaceability scores. (b) The correlation between functional irreplaceability scores of the gapless dataset (no
360 missing trait data; and a simulated imputed dataset (N = 176). (c) The change in FIRE rankings between the full
361 FIRE dataset and the gapless dataset (Accuracy) for species with full trait data availability (N = 176).
362

2.3.3 Priority FIRE species
The species with the highest FIRE scores are the Sand Tiger Shark (*Carcharias taurus*;
representing 0.44% avertable expected loss of trait space), Scalloped Hammerhead (*Sphyrna
lewini*; representing 0.33% avertable expected loss) and Basking Shark (0.31%; Fig. 5). Shark
orders Echinorhiniformes and Lamniformes have the highest average FIRE scores of 0.08%
(sd = 0.02) and 0.07% (sd = 0.12), respectively (more information on functional
irreplaceability and extinction risk in Text S2). We identified 76 shark species as priority FIRE
species, defined as threatened sharks for which we have 95% certainty that they are above
median FIRE (mirroring the approach to identifying priority EDGE species[20]; full list in Data
S2). All of the top 76 priority FIRE sharks and 67.9% of the 215 priority functionally
irreplaceable sharks have declining or unknown population trends, based on Red List data[2].
One Critically Endangered species in the top 28 FIRE sharks, the Lost Shark (*Carcharhinus
obsoletus*), is Possibly Extinct, and has not been recorded since 1934[54].

Fig. 5| Functionally irreplaceable sharks. Correlations between (a) evolutionary distinctiveness and functional irreplaceability and (b) Evolutionarily Distinct and Globally Endangered (EDGE) and Functionally Irreplaceable with Risk of Extinction (FIRE) scores. Dotted lines represent the upper 5% of scores for each metric. Silhouettes represent examples of species that fall within the top 5% of (c) evolutionary distinctiveness and functional irreplaceability (d) EDGE and FIRE. Shark species: 1. Broadnose Sevengill Shark (*Notorynchus cepedianus*), 2. Zebra Shark (*Stegostoma tigrinum*), 3. Basking Shark (*Cetorhinus maximus*), 4. Scalloped Hammerhead (*Sphyrna lewini*). The top 25 highest ranked (c) functionally irreplaceable and (d) FIRE shark species. Numbers represent the (c) functional irreplaceability and (d) FIRE rank of each species, where species with the same median value have the same rank. Boxplots represent the range, interquartile range and median of (c) functional irreplaceability and (d) FIRE scores calculated over 1000 iterations. IUCN Red List population trends (Popn) are denoted with arrows to represent increasing or decreasing population trends, a dash to represent stable population trends, and a question mark to represent unknown population trends. Species in bold text fall in the top 25 of both (c) functional irreplaceability and evolutionary distinctiveness scores and (d) FIRE and EDGE scores. Colours represent IUCN Red List categories (red = Critically Endangered, orange = Endangered, yellow = Vulnerable, light green = Near Threatened, green = Least Concern, grey = Data Deficient). Silhouette credit available in supplementary material.

2.3.4 Relationship between FIRE and EDGE

We applied the EDGE2 protocol[20] to calculate evolutionary distinctiveness and EDGE scores for sharks, and compared these with functional irreplaceability and FIRE. Functional

irreplaceability and evolutionary distinctiveness showed a weak but significant positive
correlation (Pearson's r = 0.18, p < 0.001; Fig. 5a). Six species ranked in the top 5% for both
metrics, while three of the top 5% of functionally irreplaceable species, including the
Southern Sleeper Shark (*Somniosus antarcticus*) and Gummy Shark (*Mustelus antarcticus*),
were among the bottom 5% for evolutionary distinctiveness. Ten species ranked in the top
5% for both FIRE and EDGE scores (Fig. 5b), with the Sand Tiger Shark (rank 1 for both),
Basking Shark (FIRE rank 3, EDGE rank 2), and Scalloped Hammerhead (FIRE rank 2, EDGE
rank 9) ranking highest across both frameworks.
405
406 **2.4 Case study 3: Functional irreplaceability and exploitation**
407
To determine whether there is a relationship between the different distinctiveness
measures (evolutionary distinctiveness and functional irreplaceability) and species targeted
for human exploitation, we selected sets of species of increasing richness based on
functional irreplaceability and based on evolutionary distinctiveness. Specifically, species
with higher scores in a given metric had a proportionally greater probability of being
selected in a set, we also used randomly selected sets for comparison (see Methods). There
are 468 shark species (84.0% of total) with biological resource use (BRU; i.e. intentional and
unintentional fishing) listed as a threat on the IUCN Red List[2]. For 168 species (30.2% of
total), BRU is listed as intentional, and for 300 (53.9% of total), BRU is listed as
unintentional. There are 1256 bird species with intentional BRU (i.e. hunting and collecting)
listed as a threat, with an additional 125 species threatened by unintentional BRU[2] (see
Methods).
420
For birds, selecting species based on functional irreplaceability highlighted more
intentionally exploited species than evolutionary distinctiveness at all sample sizes (Fig. 6a;
Table S2). For sharks, functional irreplaceability again highlights the greatest number of
intentionally exploited species (Fig. 6b; Table S2). Selecting shark species based on
evolutionary distinctiveness performs significantly better than random at highlighting
intentionally exploited species (Fig. 6b; ANOVA with Tukey's test all p < 0.001 compared
with random; all pairwise comparison results available in Table S2).
428

429
Fig. 6| The relationship between intentional exploitation of species and their Functional Irreplaceability and
Evolutionary Distinctiveness. The number of intentionally exploited a) bird and b) shark species highlighted
when 10%-50% of species are selected for conservation based on weighted random draws from different
distributions corresponding to different conservation strategies: uniform (fully random; orange), weighted by
functional irreplaceability (blue) or weighted by evolutionary distinctiveness (green). Boxplots represent
median value (solid line), 25th and 75th percentiles (box edges), and 5th and 95th percentiles (whiskers) from
1000 iterations. Silhouette credit available in supplementary material.

437
438 For both birds and sharks, species with intentional BRU listed as a threat had significantly
439 higher functional irreplaceability compared to species where intentional BRU is not listed as
440 a threat (sharks: mean = 0.10 vs. 0.06, df = 249.1, p < 0.001; birds: mean = 0.09 vs. 0.03, df =
441 1227.5, p < 0.001) and evolutionary distinctiveness (sharks: mean = 18.9 vs. 12.4, df = 225.1,
442 p < 0.001; birds: mean = 5.3 vs. 4.7, df = 1227.5, p = 0.001).

443
444   **3. Discussion**
445
The FIRE metric offers a novel approach to identify priority species whose conservation is
expected to avert the greatest losses of irreplaceable functional diversity. Given the wealth
of evidence linking functional diversity to the provision of ecosystem processes and services,
it is vital to prevent further erosion of functional trait space[55–57]. By applying FIRE to global-
scale datasets of birds and sharks, we illustrate the metric's ability to identify priorities for
conservation for different sets of taxa with contrasting levels of species richness and data
coverage. Our results highlight numerous species of both birds and sharks that are not
currently identified as priorities by phylogenetically-informed approaches, alongside
threatened species that are both phylogenetically and functionally irreplaceable. Further,
our results indicate that functionally irreplaceable species are disproportionately threatened
by exploitation (i.e. hunting, collecting, fishing, according to IUCN Red List data).

457
The FIRE metric, which measures the expected contribution of species to overall trait space,
builds on approaches such as EcoDGE[27] and FUSE[28] to integrate functional diversity into an
expected loss framework. As FIRE incorporates complementarity to quantify the risk to each

point in trait space due to the extinction risk of the species present, the approach also
presents further opportunities beyond species prioritisation. Our methods can be used to
measure the expected erosion of trait space due to extinctions, and to quantify variation in
the probability of occupancy in regions of trait space as extinction risk changes over time.
Through these advances, FIRE can potentially underpin the development of a coherent suite
of metrics to monitor risk to functional diversity, and guide conservation efforts, as has been
achieved with the EDGE and Phylogenetic Diversity indicators adopted by the GBF[58]. We
calculated FIRE for birds and sharks at the global scale. However, FIRE can also be calculated
at regional, national or community levels. These complementary levels of analysis would
allow us to identify species whose function within specific ecosystems or localities is
expected to be distinct and at risk of being lost[59,60]. However, the highlighted species in this
context may not be irreplaceable as they are with global scale analyses, because they could
be (re)introduced from other localities outside the focal area. Such assessments
nevertheless have strong potential to inform local, regional and global conservation
planning and policy.
476
The trait space underpinning FIRE calculations is generated using TPDs, as we wanted to
account for the probabilistic nature of species traits and consider intraspecific trait
variability when describing functional space[38]. TPDs measure the probability of occupancy
across trait space, and FIRE is designed to be calculated with either a binary or probabilistic
TPD trait space (see Methods). We used the probabilistic approach to calculate FIRE for
birds, given the large number of species and thus high potential for many species to occupy
areas of trait space fully occupied by multiple other species, leading to such species having a
functional irreplaceability and FIRE score of zero. For our shark case study, which comprised
~6% of the richness of birds, we illustrated the binary approach to the occupation of trait
space by a given species; a species occupies a region of trait space or it does not. As TPDs
use kernels to represent the probabilistic distribution of species in trait space, additional
work to explore the sensitivity of these approaches to kernel parameters is needed. Future
calculations of FIRE may incorporate intraspecific variation to quantitatively parameterise
kernel size and occupancy probability across trait space.
491
Though currently formulated based on the use of TPDs, FIRE in its broadest terms – i.e the
amount of expected functional diversity loss that can be averted through conservation
action on a single species – can in principle utilise any approach to measuring functional
richness (e.g., dendrograms). We simply apply extinction to the set of species and compare
total diversity (according to our chosen measure) of a case where our focal species is
permitted to go extinct with a case where our focal species is protected. As with all
functional diversity analyses, FIRE is sensitive to methodological choices, including the
metric used to quantify functional diversity, the traits selected to represent ecological
functions, and the number of PCA / PCoA axes retained. Consequently, any given FIRE
priority list should be viewed as contingent on these choices, rather than as a single fixed or
definitive ranking.
503
The growing interest in quantifying functional diversity for conservation purposes[61] has led
to an increase in studies examining the patterns and processes of functional diversity and
the impacts of biodiversity loss[62]. However, persistent knowledge gaps and limited trait data
continue to constrain functional diversity research, often restricting analyses to specific

taxonomic groups and geographic regions[17]. As a result, most functional diversity studies
have focused on well-documented groups such as plants and birds[33,63], potentially limiting
the generality of their conclusions. This taxonomic and geographic bias underscores the
need to broaden the scope of functional diversity assessments. Recent advances in the
collection and availability of trait data across a wider range of taxa[64–66] and the development
of methods for imputing missing data[51,67,68] present new opportunities to do so. In this
study, we demonstrate that incomplete datasets can still yield meaningful insights into
functional diversity. By calculating FIRE from an incomplete shark trait dataset using
imputation, we show that it is possible to assess functional diversity more broadly, even in
the face of substantial data gaps. However, reliance on single imputation may overlook the
uncertainty in missing data[69]. Multiple-imputation approaches offer a promising avenue for
future work, as they better capture this uncertainty[69].
520
As trait data availability increases across taxonomic groups[33,64,65,70–72], the number and types
of traits used to inform trait space will vary and will require careful consideration. For
example, an important aspect of shark life history that we were unable to include due to a
lack of trait availability is changes through different life stages[73]. Many species will display
changes in functional and behavioural features, such as shifts in trophic position and
foraging location[74] throughout their lifetime, and future work should aim to incorporate
these and other trait dynamics to understand the relationship between ecological function
and extinction risk more comprehensively.
529
The moderate correlation between FIRE and EDGE for both birds and sharks is to be
expected due to the incorporation of extinction risk in both metrics. However, there is only
a weak correlation between functional irreplaceability and evolutionary distinctiveness for
both birds and sharks - a finding that echoes recent studies that challenge the longstanding
assumption that phylogenetic diversity should be employed as a surrogate for functional
diversity[19,75]. This disparity is expected because of widespread convergent and divergent
evolution, which weakens the link between phylogeny and function[37]. Indeed, there is
growing evidence that species traits, rather than phylogeny, provide the most accurate
index of ecological function in a range of taxonomic groups[37]. Our results thus support the
argument that phylogenetic diversity and functional diversity should be considered as two
interlinked yet distinct dimensions of biodiversity, both of which are vital components of
conservation strategies to preserve ecosystems[56,76,77] and promote human wellbeing[78,79].
542
Indeed, our work further highlights the complex and non-linear relationship between major
adaptive change and evolutionary time[80–82]. For sharks, three species were in the top 5% of
functional irreplaceability but the bottom 5% of evolutionary distinct species. Exploring the
distribution of these species in trait space highlights that there are multiple clusters that are
occupied by many species. However, though these three species were clustered with their
congeners, they occupy areas on the periphery of their respective genus clusters. Their
unique trait combinations cause these species to occupy low density areas, in at least one
dimension of trait space, which is not occupied by many other species (Fig. S4). It is well
established that certain traits, such as body size and generation length[83], are associated with
increased extinction risk, and it is to be expected that regions of trait space dominated by
such traits will be inherently vulnerable[15,16,84]. Further exploration is needed to understand
how key traits are distributed across the multiple axes of trait space produced by TPDs.

Where intrinsically sensitive traits cluster in sparsely occupied regions of trait space, there is
the potential for cascading losses due to functionally clumped extinction risk[15,16].

Many of the highest-ranking FIRE species have been identified as priorities by earlier
approaches using different trait data[32], calculations of trait space[18,78], and treatment of
extinction risk[84], supporting the robustness of our identified priorities. While there is
moderate congruence in the priority species identified by FIRE and EDGE, there are high-
ranking FIRE species that could be overlooked if only focusing on high-ranking EDGE species,
such as the Slender-billed Vulture (*Gyps tenuirostris*), Sira Curassow (*Pauxi koepckeae*),
Angular Roughshark (*Oxynotus centrina*) and the Pacific Nurse Shark (*Ginglymostoma*
unami).

Indeed, prioritising any single biodiversity facet alone risks overlooking species that may
significantly contribute to other ecological or phylogenetic aspects of biodiversity. We
therefore highlighted species that are in the top 5% of both FIRE and EDGE scores as species
of particular interest whose conservation would represent large conservation gains across
multiple dimensions of biodiversity (Fig. 3 and 5). These include many iconic threatened bird
species, such as the Helmeted Hornbill, Shoebill and Kākāpō, along with high-profile shark
species, including Scalloped Hammerhead, Pelagic Thresher and Basking Shark.

Previous studies on plants[79,85] and birds[78] indicate that functionally and evolutionarily
distinct species are disproportionately utilised by humans. Our work corroborates these
earlier findings for birds and further shows that the same is true for sharks. Sets of bird and
shark species selected based on functional irreplaceability scores highlight more species
threatened by exploitation than selecting species at random. These findings support earlier
work that suggests that functionally irreplaceable species are targeted by hunting[86,87],
fishing[86,87] and the pet trade[88]. Beyond targeted fishing and exploitation, incidental catch is
also a problem for many marine species[50,89]. Transitioning to the sustainable use and
protection of species that are currently overexploited is essential to maintain functionally
irreplaceable species and the benefits provided to people through ecosystem services.

Over the last two decades, the EDGE metric has galvanised conservation action by
prioritising highly evolutionarily distinct and threatened species for practical
conservation[7,20]. Our results build on this foundation by demonstrating how the new FIRE
metric can effectively prioritise species to avert the greatest impending losses of functional
diversity. The erosion and reorganisation of ecological assemblages is already widespread[90–
92], driven by rapid declines of species populations, particularly those with unique and vital
ecological roles [90–93]. Using a trait-based approach to prioritise species with distinct
phenotypes and associated functional roles, the FIRE metric can help to bring a much-
discussed but perennially overlooked dimension of biodiversity to the forefront of
conservation planning, policy and action. Targeting this functional dimension is critical to
maintain the complex and resilient ecosystems needed to support human wellbeing now
and into the future[13].

4. Methods

Functionally Irreplaceable with Risk of Extinction (FIRE) calculations

The Functionally Irreplaceable with Risk of Extinction (FIRE) metric identifies species whose
effective conservation will avert the greatest expected losses of functional diversity. To do
this, for each species, FIRE combines two components: (i) functional irreplaceability, the
expected unique contribution of a given species to overall trait space into the future, based
on the extinction risk of all other species and (ii) the extinction risk of the species.

Species coverage of trait space

One way to measure functional irreplaceability, the property we are trying to maximise
through our prioritisation scheme, is to quantify the proportion of TPD trait space uniquely
covered by a species. As a first step it is necessary to find the hypervolume of trait space
where a given TPD summed across all species is non-zero. More generally, however, we can
calculate functional richness as;

$$\sum_{v \in V} \left(1 - \prod_{i \in S} (1 - x_{i,v}) \right)$$

where $x_{i,v}$ is the probability that species i is present in a section v of trait space, S is the set
of all species and V is the set of all possible parts of trait space (these could be voxels in n-
dimensional space). Our metric can be calculated as a binary or non-binary measure. In the
binary measure we constrain $x_{i,v}$ to be either 1 or 0. In the non-binary measure, we allow
$x_{i,v}$ to take any value between 0 and 1. These intermediate values may be based on
uncertainty and/or intraspecific variability, which may in turn be connected to the total
abundance of the species where such data are available. In cases where we assume some
degree of intraspecific variation in species traits, each species occupies multiple voxels in
trait space and the sum of occupancy across all voxels will be greater than 1. The equation
can be adapted to incorporate extinction risk E_i as a probability of extinction for each
individual species $i \in S$ to give an expression for the expected future hypervolume of trait
space.

$$\sum_{v \in V} \left(1 - \prod_{i \in S} \left(1 - x_{i,v}(1 - E_i) \right) \right)$$

Irreplaceable species-specific trait space (functional irreplaceability)

To measure the irreplaceability of a species based on how much functional diversity it
represents, we considered the extinction risk of all other species overlapping with the trait
space of the focal species to reflect the expected unique occupancy of trait space in the
future. If the focal species overlaps in trait space with few other species that are at high risk
of extinction, there will be a moderate likelihood that this focal species will be the sole
inhabitant of that region of trait space in the future. In contrast, if the focal species overlaps
in trait space with many other species or with species that have low risk of extinction, then
it is very unlikely that the focal species will be the sole inhabitant of that region of trait

space in the future. The functional irreplaceability can be expressed mathematically as
follows for species $j \in S$

$$FI(j) = 100 \cdot \frac{\sum_{v \in V} \left(x_{j,v} \left(1 - \prod_{i \in S \setminus \{j\}} \left(1 - x_{i,v}(1 - E_i) \right) \right) \right)}{\sum_{v \in V} \left(1 - \prod_{i \in S} \left(1 - x_{i,v} \right) \right)}$$

648
The numerator captures the amount of trait space that in the future, given extinction risk of
all species other than the focal species, will rely on the survival of the focal species to
remain covered. The denominator and factor of 100 expresses this as a percentage of total
trait space. Because of the numerator in particular, this equation is computationally
expensive to evaluate for all species. A naive implementation would have computational
time complexity proportional to the size of the trait space multiplied by the square of the
species richness. Furthermore, the whole calculation would still ideally need to be repeated
to capture an uncertainty distribution propagating the error in all the Ei values. Our solution
for calculating functional irreplaceability is to use the repeats required to capture
uncertainty as part of a monte carlo process to simulate the result that the above equation
captures in precise probabilistic terms. This requires three steps and is performed on every
species (summarised in Fig. 1 of the main text).

661
662 Step 1: Calculate the volume of trait space occupied by all species in the analysis.

663
Step 2: Stochastically simulate the removal of species based on their extinction risk but with
the constraint that the focal species does not go extinct (i.e., ignore the extinction risk of the
focal species for this step). We followed the Gumbs et al.[20] weighting of IUCN Red List
categories[94] to estimate probability of extinction for each species; IUCN Red List categories
were converted to represent probability of extinction based on a continuous distribution of
extinction probabilities, where the median of each Red List category was defined as, CR =
0.97; EN = 0.485; VU = 0.2425; NT = 0.12125; LC = 0.060625. Species that were not assessed
or had IUCN Red List category listed as Data Deficient, were randomly sampled from the
entire distribution of extinction probabilities. Using these probabilities, we ran simulations
to determine whether species were removed from the pool (i.e., went extinct).

674
675 Step 3: Calculate a post-extinction trait space with the focal species present, and again with
676 the focal species removed. Calculate the difference between these two trait space volumes,
677 to give the species of interest's distinct contribution to trait space. Divide the result by total
678 trait space volume (step 1) and multiply this by 100 to get a percentage of the unique trait
679 space occupied by the focal species resulting in a functional irreplaceability value.

680
681 *Avertable loss (FIRE)*

682
Step 4: To capture the avertable loss of irreplaceable functional diversity we multiplied the
functional irreplaceability of each species (from step 3) by its probability of extinction to
quantify the expected loss of functional diversity given the extinction risk of the species (i.e.,
the species' FIRE value). We calculated risk of extinction by following the Gumbs et al.[20]
weighting of IUCN Red List categories[94] to estimate the probability of extinction for each
species. The full equation for FIRE of species $j \in S$ is thus

689
$$FIRE(j) = E_j \cdot 100 \cdot \frac{\sum_{v \in V} \left(x_{j,v} \left(1 - \prod_{i \in S \setminus \{j\}} \left(1 - x_{i,v}(1 - E_i) \right) \right) \right)}{\sum_{v \in V} \left(1 - \prod_{i \in S} (1 - x_{i,v}) \right)}$$

Final functional irreplaceability and FIRE values. We first perform step 1, then calculate
functional irreplaceability and FIRE (perform steps 2-4 from above) over 1000 repeat
iterations to capture a distribution of uncertainty in the extinction scenarios. The final
functional irreplaceability and FIRE values are given by the median value of the sampled
distribution.

Case study 1: FIRE for the world's birds

Data collection

We collated published data for extinction risk, traits and evolutionary history of birds (Aves).
To taxonomically match species information from different datasets, we used nomenclature
from the latest Birdlife checklist[95] (accessed 10/2024), which included 11005 species. All
database manipulation and analyses were performed using R statistical software (v. 4.3.1)[96].

Extinction risk data. We use the IUCN Red List of Threatened Species (v. 2025.1)[2] to obtain
information of species extinction risk. This includes assessment data for 10975 species,
including 10936 data sufficient species (Least Concern, LC; Near Threatened, NT; Vulnerable,
VU; Endangered, EN; or Critically Endangered, CR) and 39 categorised as Data Deficient
(DD). The remaining 30 species in our dataset are Not Evaluated (NE) on the Red List. To
calculate a species probability of extinction we generate a continuous distribution of
extinction probabilities following methods outlined by Gumbs et al.[20], where the median of
each Red List category was defined as, CR = 0.97; EN = 0.485; VU = 0.2425; NT = 0.12125; LC
= 0.060625. Species probability of extinction was randomly selected from the distribution of
extinction probability values within their Red List categorisation. Where species were
categorised as DD or NE their extinction probability was randomly selected from the whole
distribution of extinction risk values.

Evolutionary history data. We used the EDGE2 protocol[20] to calculate evolutionary
distinctiveness (ED; the expected contribution of a species to overall evolutionary history
given the extinction risk to all other species) and Evolutionarily Distinct and Globally
Endangered (EDGE; the amount of expected loss of evolutionary history that can be
conserved by averting the extinction of a given species) scores for all birds. Evolutionary
distinctiveness and EDGE are the phylogenetic equivalents of functional irreplaceability and
FIRE, and we used the same weightings of extinction risk for EDGE and FIRE calculations. ED
and EDGE scores were calculated as the median scores from a sample of 1000 phylogenetic
trees generated from Jetz et al.[97] phylogeny, where species absent from the phylogenies
were inserted based on taxonomy using the same approach as Gumbs et al.[41].

Trait data. As trait data for birds, we used eight continuous morphological measurements
that have been demonstrated to reliably indicate the functional roles and trophic statuses of
birds on a global scale[37]: (1) total beak length (from the tip to the skull), (2) beak length to
the nares, (3) beak width and (4) depth (at the nares), (5) wing length, (6) secondary length

(length from the carpal joint to the first of the wings secondary feathers), (7) tail length and
(8) tarsus length. Data for all eight traits were sourced from the AVONET database[33]. We
also sourced information on species body mass from AVONET for use in a sensitivity test.
We investigated the relationship between traits and extinction risk using Spearman's rank
correlation.
739
740 *Quantifying trait space*
741
Principle component analysis. The eight morphological traits were all log-transformed and
then centred and scaled to a mean of zero and unit variance. A Principle Components
Analysis (PCA) was undertaken using the eight log-transformed and scaled traits and the
first three PCA axes taken to build the trait space. We used three axes as they explained
over 90% of the total variance and resulted in a more manageable number of species
(compared to when using four PCA axes) with the maximum functional irreplaceability
value. The latter is because, as trait space dimensionality increases, there is more trait space
available in which to measure differentiation and thus in most cases, and keeping all other
parameters equal, species become more functionally distinct from each other.
751
Multidimensional trait space. We use TPDs to account for intraspecific trait variability when
describing functional space[34]. TPDs represent each species in functional trait space as a
multivariate kernel density estimation. We approximated TPD values using the measured
traits available in the analysis and an estimated variability using the TPD package (v. 1.1.0)[34].
Each species occupancy is estimated in the same field defined by the three axes of the PCA,
with each grid divided into 20 equal parts. Trait space is held constant following removal of
species in subsequent analyses.
759
Functional irreplaceability/FIRE. We apply the methods stated above to calculate functional
irreplaceability and FIRE values for all birds using the non-binary method to calculate species
occupancy of trait space (where the species proportional probabilistic occupancy of each
voxel is calculated). Following the EDGE2 protocol[20] and applying it to our FIRE metric, we
defined sets of priority FIRE birds as being those that are both assessed as threatened on
the IUCN Red List and above the median, 75th percentile, and 95th percentile of functional
irreplaceability with 95% confidence (i.e. in 95% or more of the iterations of FIRE
calculation; Data S1 and S2).
768
Accounting for outliers in trait space. The five kiwi species (Apterygiformes) represent
extreme outliers in the global bird morphological trait space[37] and can have substantial
impact on functional diversity analyses if not accounted for[61]. Here, to mitigate this effect
we first built the trait space and calculated functional irreplaceability for all species with the
kiwis included, storing the functional irreplaceability values for the five kiwi species. We
then removed the five kiwi species and re-built the trait space (first re-scaling the traits and
re-running the PCA). This second trait space was used to calculate functional irreplaceability
for all other bird species (i.e., all species other than the kiwis). The functional irreplaceability
values for the kiwis were then added to these values prior to FIRE values being calculated
for all birds. This process means that, when the second trait space is analysed, the kiwis are
not included in the species removed based on their extinction risk (i.e., Step 2 above).

However, this is unlikely to be an issue given they occupy an entirely distinct area of trait
space and thus will not influence the functional irreplaceability of any other species.
782
Trait space extinction projections. To understand the future impact of species extinctions on
global trait space, we estimated the proportional loss of each voxel in trait space. Future
occupancy of voxels in trait space is calculated as a function of current occupancy of species
in trait space and extinction probabilities. To estimate this, we calculate the probability that
at least one species remains in a given voxel, after accounting for extinction probabilities,
and divide this by the probability that at least one species currently occupies that voxel:
789

790
$$\frac{1 - \prod_{i \in S}\left(1 - x_{i,v}(1 - E_i)\right)}{1 - \prod_{i \in S}\left(1 - x_{i,v}\right)}$$

791
Where $x_{i,v}$ is the probability that species i is present in voxel v of trait space, E_i is the
probability of extinction for species i and S is the set of all species. Each future projection is
calculated 1000 times to reflect the distribution of species extinction predictions. These
predictions were mapped onto the three principal component axes of trait space to visualise
the impact of extinction on different areas of trait space and identify certain trait
combinations that may be vulnerable to extinction.
798
799   ***Avian functional irreplaceability***
800
We used functional irreplaceability scores to identify bird species with little or no functional
redundancy, including those with no overlap with other species in our trait space. To
explore the relationship between functional irreplaceability and extinction risk, we used
Welch's t-test to compare the mean functional irreplaceability of threatened (i.e. VU, EN, CR
on the Red List) and non-threatened (LC and NT on the Red List) species. We selected the
species in the top 5% highest functional irreplaceability scores (following Safi et al. 2013[98]),
and calculated the proportion that are threatened, and compared this qualitatively with the
overall proportion of threatened birds, derived from the 2025.1 version of the IUCN Red
List[2].
810
811   ***Unique and convergent FIRE priorities***
812
To compare convergent and unique conservation priorities between FIRE and the
phylogenetically-informed EDGE approach, we matched the bird species in our FIRE dataset
with those with available EDGE data. To achieve this, we used existing EDGE2 scores for
birds[41,99], updated for the 2025.1 Red List extinction risk data[100]. We limited our
comparisons of FIRE and EDGE data to species for which both data were available (N =
10971). To quantify the relationship between avian functional irreplaceability and its
phylogenetic equivalent, ED, we used Spearman's rank correlation. We also used Welch's t-
test to test whether species in monotypic families had higher functional irreplaceability than
birds in general. We again followed Safi et al.[98] in highlighting species with top 5% highest
FIRE scores as top priorities, and restricted this set to species in a threatened or Extinct in
the Wild Red List category. Using this set of priority species, we identified the number of
species also in the top 5% threatened species based on EDGE scores. We also calculated

what proportion of these priority species qualify as priority EDGE species (i.e. above median
EDGE with 95% confidence and threatened with extinction[20]), to identify species not
currently captured by a primary phylogenetically-informed prioritisation. To explore how
well the priority FIRE species are represented by other conservation mechanisms, we
calculated how many species trigger Key Biodiversity Areas[101] (KBAs), and used published
estimates of range overlap with KBAs[102] to identify species potentially underrepresented in
the global KBA network. Lansley et al.[102] suggested that species with <8% of their area of
habitat[103] (AOH; IUCN range maps trimmed to suitable habitat and elevation) covered by
KBAs may not be effectively represented in priority areas for maintaining biodiversity.

Case study 2: Calculating FIRE for sharks

Data collection

We collated published data for extinction risk, traits and evolutionary history of sharks
(Selachimorpha). We limited our study to sharks, omitting other clades of Chondrichthyes
(Batoidea and Holocephali), due to limited availability of comparable data for several key
traits, including body shape. To taxonomically match species information from different
datasets, we used nomenclature from FishBase[64] (accessed 11/2023), which included 557
species.

Extinction risk data. We use the IUCN Red List of Threatened Species (v. 2023.1)[104] to obtain
information of species extinction risk. This includes assessment data for 540 species,
including 467 data sufficient species (Least Concern, LC; Near Threatened, NT; Vulnerable,
VU; Endangered, EN; or Critically Endangered, CR) and 73 categorised as Data Deficient
(DD). The remaining 17 species in our dataset are Not Evaluated (NE) on the Red List. To
calculate a species probability of extinction we generate a continuous distribution of
extinction probabilities following methods outlined by Gumbs et al.[20], where the median of
each Red List category was defined as, CR = 0.97; EN = 0.485; VU = 0.2425; NT = 0.12125; LC
= 0.060625. Species probability of extinction was randomly selected from the distribution of
extinction probability values within their Red List categorisation. Where species were
categorised as DD or NE their extinction probability was randomly selected from the whole
distribution of extinction risk values.

Evolutionary history data. We used the EDGE2 protocol[20] to calculate evolutionary
distinctiveness and EDGE scores for all sharks. ED and EDGE scores were calculated as the
median scores from a sample of 1000 phylogenetic trees generated from the Stein et al.[105]
tree distribution of 10,000 Chondrichthyes trees, where species absent from the
phylogenies were inserted based on taxonomy using the same approach as Gumbs et al.[41].

Trait data. We collated eight traits for our analysis (Data S4), obtained from FishBase[64] and
Sharks of the World: A Complete Guide[106]. We selected traits based on data availability and
representation of functional guilds[107] (Table S1). Our traits were selected to capture a broad
suite of traits that are important to describing species' roles within an ecosystem reflecting
life history strategies, habitat use, trophic interactions and morphological variation (Table
S1)[108,109]. To measure body shape, we modify methods shown in Siders et al.[108], which uses
Elliptical Fourier analysis (EFA) to represent variation in shark body shape. We collected 515

illustrated images from Sharks of the World: A Complete Guide to measure the variation in
shark body shape using EFAs. These images were first converted into outlined silhouettes
using photoshop (v. 23.5.5)[110]. Using the R package Momocs (v. 1.4.1)[111] each outline was
scaled and centred on a matrix (Fig. S5) and EFA was performed on the outlines, using 16
harmonics (Fig. S6). Results of the EFA were condensed using Principal Component Analysis
(PCA; Data S5). PC1 and 2 collectively explain 62% of total variation and were used to
represent body shape for our analysis (Fig. S7), PC1 represents 43% variation and PC2
represents 19% variation.
880
881 *Imputing missing data*
882
883 Our compiled database contained 557 species. To complete missing values in the trait
884 database, we imputed missing data using the missForest function in the missForest R
885 package (v. 1.5)[51]. This method uses a random forest trained on the observed values of the
886 trait matrix to impute missing values. It can be used to predict both continuous and
887 categorical data.
888
To test the performance of imputation we limited the dataset to only include species with
complete trait data and simulated incomplete trait data by artificially removing trait data.
We randomly removed data from each trait, with the percentage of the trait removal equal
to the percentage of the trait data missing from our full database. Simulated removal of
data was repeated 100 times to get a variable combination of missing values. We then
imputed the missing data using missForest and calculated each species functional
irreplaceability 100 times, and calculated this 100 times from the unimputed dataset. We
compared the median functional irreplaceability of each species from the complete dataset
and the simulated imputed dataset using Spearman's rank correlation to test the robustness
of imputation using missForest for the observed proportions of missing data.
899
To examine the robustness of FIRE rankings to the inclusion of species with missing data, we
calculated FIRE rankings for the set of species with complete trait data and compared these
to the rankings of the species when FIRE was calculated for all species (including imputed
data). To examine the robustness of priority species identified, we calculated the number
and proportion of species that were in the top 10% and 20% of both rankings. We then used
Spearman's rank correlation to examine the relationship between the absolute change in
rankings for each species between the two datasets, and the priority ranking of each species
in the FIRE rankings restricted to species with complete trait data. If there is no relationship,
this indicates that high priority rankings are similarly unstable as low-priority rankings,
whereas a positive correlation would indicate greater stability in the ranks of high priority
species.
911
912 To assess relationships between species traits and extinction risk, we used Spearman's rank
913 correlation to quantify relationships with extinction risk and continuous trait (body length,
914 trophic level, maximum depth, growth ratio, body shape variation PC1 and body shape
915 variation PC2). For categorical traits (habitat preference and reproductive guild), we used
916 Kruskal-Wallis tests to test for differences in extinction score among trait categories.
917
918 *Quantifying trait space*

Distance matrix. Prior to quantifying a distance matrix, we log transformed, scaled and centred continuous traits to ensure normal distribution and prevent overweighting. We calculated a dissimilarity distance matrix (Gower's distance) using the function gawdis (v. 0.1.5)[14]. This approach prevents disproportionate weighting of categorical traits to produce multi-trait dissimilarity with more uniform contributions of different traits[14].

Principal coordinate analyses. Using the dudi.pco function in the ade4 package (v. 1.7.22)[112] we summarised the functional dissimilarity matrix with PCoA to quantify distances in trait space between species. Multivariate trait space was built by extracting the first four PCoA axes. To assess how imputation was distributed across trait space, we calculated the proportion of imputed data for each species and summarised imputation density across trait space. We calculated the average proportion of imputed data along the first two PCoA axes by summarising the mean of each point along both PCoA axes.

Multidimensional trait space. We approximated TPD values using the measured traits available in the analysis and an estimated variability using the TPD package (v. 1.1.0)[34]. Each species occupancy is estimated in the same field defined by the four axes of the PCoA, with each grid divided into 25 equal parts. Trait space is held constant following removal of species in subsequent analyses.

Functional irreplaceability/FIRE. We apply the methods stated above to calculate functional irreplaceability and FIRE values for all sharks using the binary method to calculate species occupancy of trait space (either a species occupies a voxel (1) or it does not (0)). Following the EDGE2 protocol[20] and applying it to our FIRE metric, we defined sets of priority FIRE sharks as being those that are both assessed as threatened on the IUCN Red List and above the median, 75th percentile, and 95th percentile of functional irreplaceability with 95% confidence (i.e. in 95% or more of the iterations of FIRE calculation; Data S2).

Relationship between FIRE and EDGE

We used Pearson's correlations to examine the relationship between functional irreplaceability and evolutionary distinctiveness scores and between FIRE and EDGE scores. To determine species that are of particularly high priority for conserving irreplaceable functional and phylogenetic diversity we identified species that are in the top 5% of both functional irreplaceability and evolutionary distinctiveness as well as FIRE and EDGE.

To better understand patterns of high functional irreplaceability in highly speciose groups (i.e. those with many close evolutionary relatives) we investigated the distribution of three species— the Gummy Shark (*Mustelus antarcticus*), Rig (*Mustelus lenticulatus*) and Southern Sleeper Shark (*Somniosus antarcticus*)—and their congeners in trait space. We plot each dimension of trait space and highlight the three species and their congeners in trait space (Fig. S4).

Functional irreplaceability and exploitation

We identified exploited bird and shark species using the IUCN Red List threats classification
scheme (https://www.iucnredlist.org/resources/threat-classification-scheme)[2], where
threat type was listed as Biological resource use. We consider a species intentionally
exploited when the species being assessed is the target, as codified under Intentional use,
Intentional use: subsistence/small scale and Intentional use: large scale. Unintentionally
exploited species were those indicated as being impacted by Unintentional effects,
Unintentional effects: subsistence/small scale and Unintentional effects: large scale. Species
that were listed as both intentional use and unintentional effects were not included in the
unintentionally exploited categorisation.
975

To explore the potential of different prioritisation strategies at safeguarding intentionally
exploited species, we selected 10%, 20%, 30%, 40% and 50% of all bird and shark species,
weighted by different prioritisation measures: evolutionary distinctiveness score; functional
irreplaceability score and randomly selected species. We repeated this 1000 times for each
approach and compared the number of intentionally exploited species captured by each
approach for each sample size. To test if prioritisation strategies differed significantly at
safeguarding intentionally exploited species we used ANOVA with Tukey's Honestly
Significant Difference post hoc test to test the pairwise difference between strategies.
984

To consider species that are unintentionally exploited we compared functional
irreplaceability and evolutionary distinctiveness scores of species using t-tests. To
investigate the potential impact of human exploitation on shark functional diversity and
phylogenetic diversity, we removed species categorised as i) intentionally exploited and ii)
both intentionally and unintentionally exploited to identify the loss of trait space and
cumulative years of evolutionary history. We compared the loss of trait space and
evolutionary history in exploited species to a randomly selected sample of species of the
same number.
993

References

1. Mace, G. M., Norris, K. & Fitter, A. H. Biodiversity and ecosystem services: a multilayered relationship. *Trends in Ecology & Evolution* **27**, 19–26 (2012).

2. IUCN Red List of Threatened Species. The IUCN Red List of Threatened Species. *IUCN Red List of Threatened Species* https://www.iucnredlist.org/en (2025).

3. Hoekstra, J. M., Boucher, T. M., Ricketts, T. H. & Roberts, C. Confronting a biome crisis: global disparities of habitat loss and protection. *Ecology Letters* **8**, 23–29 (2005).

4. Mittermeier, R. A. *et al.* Wilderness and biodiversity conservation. *Proceedings of the National Academy of Sciences* **100**, 10309–10313 (2003).

5. IPCC. *Global Warming of 1.5°C: IPCC Special Report on Impacts of Global Warming of 1.5°C above Pre-Industrial Levels in Context of Strengthening Response to Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*. (Cambridge University Press, Cambridge, 2022).

6. Foden, W. B. *et al.* Climate change vulnerability assessment of species. *WIREs Climate Change* **10**, e551 (2019).

7. Isaac, N. J. B., Turvey, S. T., Collen, B., Waterman, C. & Baillie, J. E. M. Mammals on the EDGE: Conservation Priorities Based on Threat and Phylogeny. *PLOS ONE* **2**, e296 (2007).

8. Cadotte, M. W., Carscadden, K. & Mirotchnick, N. Beyond species: functional diversity and the maintenance of ecological processes and services. *Journal of Applied Ecology* **48**, 1079–1087 (2011).

9. Violle, C. *et al.* Let the concept of trait be functional! *Oikos* **116**, 882–892 (2007).

10. Palacio, F. X. *et al.* A protocol for reproducible functional diversity analyses. *Ecography* **2022**, e06287 (2022).

11. Miatta, M., Bates, A. E. & Snelgrove, P. V. R. Incorporating biological traits into

conservation strategies. *Annu. Rev. Mar. Sci.* **13**, 421–443 (2021).

12. Convention on Biological Diversity. Kunming-Montreal Global Biodiversity Framework.

Tech. rep. Montreal, Canada (2022).

13. Tobias, J. A., Bullock, J. M., Dicks, L. V., Forester, B. R. & Razgour, O. Biodiversity

conservation requires integration of species-centric and process-based strategies.

Proceedings of the National Academy of Sciences **122**, e2410936122 (2025).

14. de Bello, F., Botta-Dukát, Z., Lepš, J. & Fibich, P. Towards a more balanced combination

of multiple traits when computing functional differences between species. *Methods in*

Ecology and Evolution **12**, 443–448 (2021).

15. Ali, J. R., Blonder, B. W., Pigot, A. L. & Tobias, J. A. Bird extinctions threaten to cause

disproportionate reductions of functional diversity and uniqueness. *Functional Ecology*

37, 162–175 (2023).

16. Carmona, C. P. *et al.* Erosion of global functional diversity across the tree of life. *Science*

Advances **7**, eabf2675 (2021).

17. Oliveira, B. F., Scheffers, B. R. & Costa, G. C. Decoupled erosion of amphibians'

phylogenetic and functional diversity due to extinction. *Global Ecology and*

Biogeography **29**, 309–319 (2020).

18. Pollock, L. J., Thuiller, W. & Jetz, W. Large conservation gains possible for global

biodiversity facets. *Nature* **546**, 141–144 (2017).

19. Faith, D. P. Conservation evaluation and phylogenetic diversity. *Biological Conservation*

61, 1–10 (1992).

20. Gumbs, R. *et al.* The EDGE2 protocol: Advancing the prioritisation of Evolutionarily

Distinct and Globally Endangered species for practical conservation action. *PLOS Biology*

21, e3001991 (2023).

21. Faith, D. P. Threatened Species and the Potential Loss of Phylogenetic Diversity:

Conservation Scenarios Based on Estimated Extinction Probabilities and Phylogenetic

Risk Analysis. *Conservation Biology* **22**, 1461–1470 (2008).

22. Jensen, E. L., Mooers, A. Ø., Caccone, A. & Russello, M. A. I-HEDGE: determining the

optimum complementary sets of taxa for conservation using evolutionary isolation.

PeerJ **4**, e2350 (2016).

23. Robuchon, M. *et al.* Revisiting species and areas of interest for conserving global

mammalian phylogenetic diversity. *Nat Commun* **12**, 3694 (2021).

24. Steel, M., Mimoto, A. & Mooers, A. Ø. Hedging Our Bets: The Expected Contribution of

Species to Future Phylogenetic Diversity. *Evol Bioinform Online* **3**, 117693430700300024

(2007).

25. Gouhier, T. C. & Pillai, P. Avoiding Conceptual and Mathematical Pitfalls When

Developing Indices to Inform Conservation. *Frontiers in Ecology and Evolution* **8**, (2020).

26. Pavoine, S. & Ricotta, C. Identifying functionally distinctive and threatened species.

Biological Conservation **284**, 110170 (2023).

27. Hidasi-Neto, J., Loyola, R. & Cianciaruso, M. V. Global and local evolutionary and

ecological distinctiveness of terrestrial mammals: identifying priorities across scales.

Diversity and Distributions **21**, 548–559 (2015).

28. Pimiento, C. *et al.* Functional diversity of marine megafauna in the Anthropocene.

Science Advances **6**, eaay7650 (2020).

29. Mouillot, D. *et al.* Rare Species Support Vulnerable Functions in High-Diversity

Ecosystems. *PLOS Biology* **11**, e1001569 (2013).

30. Mammola, S., Carmona, C. P., Guillerme, T. & Cardoso, P. Concepts and applications in

functional diversity. *Functional Ecology* **35**, 1869–1885 (2021).

31. Tucker, C. M. *et al.* A guide to phylogenetic metrics for conservation, community ecology

and macroecology. *Biological Reviews* **92**, 698–715 (2017).

32. Pimiento, C. *et al.* Functional diversity of sharks and rays is highly vulnerable and

supported by unique species and locations worldwide. *Nat Commun* **14**, 7691 (2023).

33. Tobias, J. A. *et al.* AVONET: morphological, ecological and geographical data for all birds.

Ecology Letters **25**, 581–597 (2022).

34. Carmona, C. P., de Bello, F., Mason, N. W. H. & Lepš, J. Trait probability density (TPD):

measuring functional diversity across scales based on TPD with R. *Ecology* **100**, e02876

(2019).

35. Fraixedas, S. *et al.* A state-of-the-art review on birds as indicators of biodiversity:

Advances, challenges, and future directions. *Ecological Indicators* **118**, 106728 (2020).

36. Hughes, A. C. *et al.* Sampling biases shape our view of the natural world. *Ecography* **44**,

1259–1269 (2021).

37. Pigot, A. L. *et al.* Macroevolutionary convergence connects morphological form to

ecological function in birds. *Nat Ecol Evol* **4**, 230–239 (2020).

38. Carmona, C. P., de Bello, F., Mason, N. W. H. & Lepš, J. Traits Without Borders:

Integrating Functional Diversity Across Scales. *Trends in Ecology & Evolution* **31**, 382–

394 (2016).

39. Hughes, E. C., Edwards, D. P. & Thomas, G. H. The homogenization of avian

morphological and phylogenetic diversity under the global extinction crisis. *Current*

Biology **32**, 3830-3837.e3 (2022).

40. Matthews, T. J. *et al.* Threatened and extinct island endemic birds of the world:

Distribution, threats and functional diversity. *Journal of Biogeography* **49**, 1920–1940

(2022).

41. Gumbs, R. *et al.* Global conservation status of the jawed vertebrate Tree of Life. *Nat*

Commun **15**, 1101 (2024).

42. BirdLife International. Apteryx rowi. The IUCN Red List of Threatened Species 2021:

e.T22732871A180757495. https://dx.doi.org/10.2305/IUCN.UK.2021-

3.RLTS.T22732871A180757495.en (2021).

43. Díaz, S. *et al.* Set ambitious goals for biodiversity and sustainability. *Science* **370**, 411–

413 (2020).

44. Mace, G. M. *et al.* Approaches to defining a planetary boundary for biodiversity. *Global*

Environmental Change **28**, 289–297 (2014).

45. McKinney, M. L. High Rates of Extinction and Threat in Poorly Studied Taxa.

Conservation Biology **13**, 1273–1281 (1999).

46. Bland, L. M., Collen, B., Orme, C. D. L. & Bielby, J. Predicting the conservation status of

data-deficient species. *Conserv Biol* **29**, 250–259 (2015).

47. Bachman, S. P., Brown, M. J. M., Leão, T. C. C., Nic Lughadha, E. & Walker, B. E.

Extinction risk predictions for the world's flowering plants to support their conservation.

New Phytologist **242**, 797–808 (2024).

48. Weedop, K. B., Mooers, A. Ø., Tucker, C. M. & Pearse, W. D. The effect of phylogenetic

uncertainty and imputation on EDGE Scores. *Animal Conservation* **22**, 527–536 (2019).

49. Stewart, K. *et al.* Functional diversity metrics can perform well with highly incomplete

data sets. *Methods in Ecology and Evolution* **14**, 2856–2872 (2023).

50. Dulvy, N. K. *et al.* Overfishing drives over one-third of all sharks and rays toward a global

extinction crisis. *Current Biology* **31**, 4773-4787.e8 (2021).

51. Stekhoven, D. J. & Bühlmann, P. MissForest—non-parametric missing value imputation

for mixed-type data. *Bioinformatics* **28**, 112–118 (2012).

52. Gendre, M., Hauffe, T., Pimiento, C. & Silvestro, D. Benchmarking imputation methods

for categorical biological data. *Methods in Ecology and Evolution* **15**, 1624–1638 (2024).

53. May, J. A., Feng, Z. & Adamowicz, S. J. A real data-driven simulation strategy to select an

imputation method for mixed-type trait data. *PLOS Computational Biology* **19**, e1010154

(2023).

54. White, W. T., Kyne, P. M. & Harris, M. Lost before found: A new species of whaler shark

Carcharhinus obsolerus from the Western Central Pacific known only from historic

records. *PLOS ONE* **14**, e0209387 (2019).

55. Violle, C. *et al.* Functional Rarity: The Ecology of Outliers. *Trends in Ecology & Evolution*

32, 356–367 (2017).

56. Cadotte, M. W. Functional traits explain ecosystem function through opposing

mechanisms. *Ecology Letters* **20**, 989–996 (2017).

57. Díaz, S. *et al.* Functional traits, the phylogeny of function, and ecosystem service

vulnerability. *Ecology and Evolution* **3**, 2958–2975 (2013).

58. CBD. Monitoring framework for the Kunming-Montreal global biodiversity framework. in

Conference of the parties to the convention on biological diversity fifteenth meeting. p.

CBD/COP/DEC/15/5 (2022).

59. Mazel, F. *et al.* Prioritizing phylogenetic diversity captures functional diversity unreliably.

Nat Commun **9**, 2888 (2018).

60. Cadotte, M., Albert, C. H. & Walker, S. C. The ecology of differences: assessing

community assembly with trait and evolutionary distances. *Ecology Letters* **16**, 1234–

1244 (2013).

61. Cardoso, P. *et al.* Calculating functional diversity metrics using neighbor-joining trees.

Ecography **2024**, e07156 (2024).

62. Matthews, T. J. *et al.* The global loss of avian functional and phylogenetic diversity from

anthropogenic extinctions. *Science* **386**, 55–60 (2024).

63. Kattge, J. *et al.* TRY plant trait database–enhanced coverage and open access. *Global*

change biology **26**, 119–188 (2020).

64. Froese, R. & Pauly, D. FishBase. www.fishbase.org (2023).

65. Meiri, S. SquamBase—A database of squamate (Reptilia: Squamata) traits. *Global*

Ecology and Biogeography **33**, e13812 (2024).

66. Jones, K. E. *et al.* PanTHERIA: a species-level database of life history, ecology, and

geography of extant and recently extinct mammals: Ecological Archives E090-184.

Ecology **90**, 2648–2648 (2009).

67. Buuren, S. V. & Groothuis-Oudshoorn, K. **mice** : Multivariate Imputation by Chained

Equations in *R*. *J. Stat. Soft.* **45**, (2011).

68. Goolsby, E. W., Bruggeman, J. & Ané, C. Rphylopars: fast multivariate phylogenetic

comparative methods for missing data and within-species variation. *Methods in Ecology*

and Evolution **8**, 22–27 (2017).

69. Blomberg, S. P. & Todorov, O. S. The fallacy of single imputation for trait databases: Use

multiple imputation instead. *Methods in Ecology and Evolution* **16**, 658–667 (2025).

70. Oskyrko, O., Mi, C., Meiri, S. & Du, W. ReptTraits: a comprehensive dataset of ecological

traits in reptiles. *Sci Data* **11**, 243 (2024).

71. Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C. & Costa, G. C. AmphiBIO,

a global database for amphibian ecological traits. *Sci Data* **4**, 170123 (2017).

72. Mull, C. G. *et al.* Sharkipedia: a curated open access database of shark and ray life

history traits and abundance time-series. *Sci Data* **9**, 559 (2022).

73. Ahnelt, H., Sauberer, M., Ramler, D., Koch, L. & Pogoreutz, C. Negative allometric growth

during ontogeny in the large pelagic filter-feeding basking shark. *Zoomorphology* **139**,

71–83 (2020).

74. Curnick, D. J., Carlisle, A. B., Gollock, M. J., Schallert, R. J. & Hussey, N. E. Evidence for

dynamic resource partitioning between two sympatric reef shark species within the

British Indian Ocean Territory. *Journal of Fish Biology* **94**, 680–685 (2019).

75. Vane-Wright, R. I., Humphries, C. J. & Williams, P. H. What to protect?—Systematics and

the agony of choice. *Biological Conservation* **55**, 235–254 (1991).

76. Collen, B. *et al.* Investing in evolutionary history: implementing a phylogenetic approach

for mammal conservation. *Philosophical Transactions of the Royal Society B: Biological*

Sciences **366**, 2611–2622 (2011).

77. Forest, F. *et al.* Preserving the evolutionary potential of floras in biodiversity hotspots.

Nature **445**, 757–760 (2007).

78. Gumbs, R. *et al.* Conserving avian evolutionary history can effectively safeguard future

benefits for people. *Science Advances* **9**, eadh4686 (2023).

79. Molina-Venegas, R. Conserving evolutionarily distinct species is critical to safeguard

human well-being. *Sci Rep* **11**, 24187 (2021).

80. Rosindell, J., Manson, K., Gumbs, R., Pearse, W. D. & Steel, M. Phylogenetic Biodiversity

Metrics Should Account for Both Accumulation and Attrition of Evolutionary Heritage.

Systematic Biology **73**, 158–182 (2024).

81. Kemp, T. S. *The Origin of Higher Taxa: Palaeobiological, Developmental, and Ecological*

Perspectives. (University of Chicago Press, Chicago, IL, 2015).

82. Tucker, C. M., Davies, T. J., Cadotte, M. W. & Pearse, W. D. On the relationship between

phylogenetic diversity and trait diversity. *Ecology* **99**, 1473–1479 (2018).

83. Chichorro, F. *et al.* Trait-based prediction of extinction risk across terrestrial taxa.

Biological Conservation **274**, 109738 (2022).

84. Stewart, K. *et al.* Threat reduction must be coupled with targeted recovery programmes

to conserve global bird diversity. *Nat Ecol Evol* 1–13 (2025) doi:10.1038/s41559-025-

02746-z.

85. Molina-Venegas, R., Rodríguez, M. Á., Pardo-de-Santayana, M., Ronquillo, C. &

Mabberley, D. J. Maximum levels of global phylogenetic diversity efficiently capture

plant services for humankind. *Nat Ecol Evol* **5**, 583–588 (2021).

86. Ripple, W. J. *et al.* Extinction risk is most acute for the world's largest and smallest

vertebrates. *Proceedings of the National Academy of Sciences* **114**, 10678–10683 (2017).

87. Ripple, W. J. *et al.* Are we eating the world's megafauna to extinction? *Conservation*

Letters **12**, e12627 (2019).

88. Toomes, A. *et al.* Drivers of the Australian native pet trade: The role of species traits,

socioeconomic attributes and regulatory systems. *Journal of Applied Ecology* **59**, 1268–

1278 (2022).

89. Branch, T. A., Lobo, A. S. & Purcell, S. W. Opportunistic exploitation: an overlooked

pathway to extinction. *Trends in Ecology & Evolution* **28**, 409–413 (2013).

90. Blowes, S. A. *et al.* The geography of biodiversity change in marine and terrestrial

assemblages. *Science* **366**, 339–345 (2019).

91. Hillebrand, H. *et al.* Biodiversity change is uncoupled from species richness trends:

Consequences for conservation and monitoring. *Journal of Applied Ecology* **55**, 169–184

(2018).

92. Cooke, R. S. C., Eigenbrod, F. & Bates, A. E. Projected losses of global mammal and bird

ecological strategies. *Nat Commun* **10**, 2279 (2019).

93. IPBES. *Global Assessment Report on Biodiversity and Ecosystem Services of the*

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

https://zenodo.org/records/6417333 (2019) doi:10.5281/zenodo.6417333.

94. IUCN. *IUCN Red List Categories and Criteria, Version 3.1*. (IUCN Species Survival

Commission, 2001).

95. BirdLife International. BirdLife Taxonomic Checklist v6. (2021).

96. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing (2023).

97. Jetz, W. *et al.* Global Distribution and Conservation of Evolutionary Distinctness in Birds.

Current Biology **24**, 919–930 (2014).

98. Safi, K., Armour-Marshall, K., Baillie, J. E. M. & Isaac, N. J. B. Global Patterns of

Evolutionary Distinct and Globally Endangered Amphibians and Mammals. *PLOS ONE* **8**,

e63582 (2013).

99. McClure, C. J. W. *et al.* Conserving the evolutionary history of birds. *Conservation*

Biology **37**, e14141 (2023).

100. BirdLife International. BirdLife Taxonomic Checklist. (2024).

101. BirdLife International. The World Database of Key Biodiversity Areas. (2025).

102. Lansley, T. P., Crowe, O., Butchart, S. H. M., Edwards, D. P. & Thomas, G. H.

Effectiveness of Key Biodiversity Areas in representing global avian diversity.

Conservation Biology **n/a**, e70000 (2025).

103. Brooks, T. M. *et al.* Measuring Terrestrial Area of Habitat (AOH) and Its Utility for the

IUCN Red List. *Trends in Ecology & Evolution* **34**, 977–986 (2019).

104. IUCN Red List of Threatened Species. The IUCN Red List of Threatened Species. *IUCN*

Red List of Threatened Species https://www.iucnredlist.org/en (2023).

105. Stein, R. W. *et al.* Global priorities for conserving the evolutionary history of sharks,

rays and chimaeras. *Nat Ecol Evol* **2**, 288–298 (2018).

106. Ebert, D. A., Dando, M. & Fowler, S. *Sharks of the World: A Complete Guide*. (Princeton

University Press, 2021).

107. Villéger, S., Brosse, S., Mouchet, M., Mouillot, D. & Vanni, M. J. Functional ecology of

fish: current approaches and future challenges. *Aquat Sci* **79**, 783–801 (2017).

108. Siders, Z. A. *et al.* Functional and phylogenetic diversity of sharks in the Northeastern

Pacific. *Journal of Biogeography* **n/a**, (2022).

109. Henderson, C. J. *et al.* Long term declines in the functional diversity of sharks in the

coastal oceans of eastern Australia. *Commun Biol* **7**, 1–8 (2024).

110. Adobe Inc. Adobe Photoshop. (2023).

111. Bonhomme, V., Picq, S., Gaucherel, C. & Claude, J. Momocs: Outline Analysis Using R.

Journal of Statistical Software **56**, 1–24 (2014).

112. Dray, S. & Dufour, A.-B. The ade4 Package: Implementing the Duality Diagram for

Ecologists. *Journal of Statistical Software* **22**, 1–20 (2007).

1249
1250

1251 Supplementary material
1252
1253 **Supplementary text**
1254
1255 **S1 Text | Bird functional irreplaceability; hummingbirds as a case study.**
1256
Inspecting the list of the most functionally irreplaceable birds indicates that there are
numerous hummingbirds included: 12 of the top 50 functionally irreplaceable birds are
hummingbirds. Hummingbirds as a group are known to be particularly functionally
distinctive[1,2]. However, within hummingbirds, while some of these species are widely
recognised to be functionally unique species (e.g., sword-billed hummingbird *Ensifera*
ensifera – the only species with a beak longer than its body, excluding the tail), the inclusion
of some of the other species in the list of most functionally irreplaceable birds is arguably
less intuitive. To check that our analytical approach was working as intended, we inspected
the location of the most irreplaceable hummingbird species in the 3-dimensional global bird
trait space (the trait space excluding the five kiwi species; Fig. A). In Figure A hummingbirds
(Trochilidae) are shown as red points with the six most functionally irreplaceable
hummingbird species shown as blue points, and all other bird species as black points. Two
things are apparent when inspecting Fig. A: First, as expected, hummingbirds as a group are
relatively distinct relative to all other birds, at least in this 3-dimensional trait space. Second,
the hummingbirds assessed to be the most functionally irreplaceable in our analyses are
clearly located on the edge of trait space, thus indicating that our analyses are working as
expected. It is likely that, while the individual traits of some species may not make some
species appear particularly functionally distinct to the naked eye (i.e., they may not have
extreme individual trait values such as a very long bill), the use of PCA better highlights
species that have rare combinations of trait values (i.e., species that are relatively distinct in
PCA space). A similar process is observed with certain palm swift species, where swifts in
general are located on the edge of trait space, and then certain palm swift species – those
that have rare combinations of trait values – are located on the edge of this group and thus
occupying the extremities of trait space. As a final sense check, we highlighted the location
of the species with the top 100 functional irreplaceability values in the 3-dimensional global
bird trait space (Fig. B); this plot clearly shows that the species are, as expected, located
towards the edges of trait space.

1284

Fig. A | The location of hummingbirds (red points; n = 360) in the global bird trait space. The six
hummingbird species with the highest functional irreplaceability scores (*Coeligena orina*, *Discosura letitiae*,
Phaethornis yaruqui, *Pterophanes cyanopterus*, *Ensifera ensifera*, *Chaetocercus berlepschi*) are shown as blue
points and all other species (n = 10,634) as black points. The trait space is that used in the main analyses, built
after excluding the five kiwi species. Both plots show the same trait space at different rotations.

Fig. B | The location of the 100 most functionally irreplaceable birds (red points) in the global bird trait
space. All other species (n = 10,900) are shown as black points. The trait space is that used in the main
analyses, built after excluding the five kiwi species. Both plots show the same trait space at different rotations.
1293

1294   **S2 Text | Shark functional irreplaceability and extinction risk**

1295

Current unique irreplaceability and FIRE values.
Incorporating functional complementarity [i.e. considering the extinction risk of species
closely associated in trait space] into the functional irreplaceability calculations will change

the consequences of each species' functional irreplaceability score, depending on their
relation in trait space with other species, and the extinction probability of those species. To
test the correlation between current unique irreplaceability and our measure of functional
irreplaceability based on expected loss, we also calculate current irreplaceability as the area
of current trait space uniquely occupied by a species. We calculated the proportion of each
species' functional irreplaceability score that was due to their unique contribution to
current trait space compared with the proportion gained through extinction scenarios (Fig.
A).
1307

1308
1309 **Fig. A | Proportion of unique functional distinctiveness of each species represented by current trait space**
1310 **occupancy.** The proportion of each species functional distinctiveness score due to current trait space
1311 occupancy compared to future trait space occupancy. The median amount of functional distinctiveness
1312 represented by current trait space is 75% of the species functional distinctiveness score.
1313

1314 To ensure there is no overweighting of extinction probabilities, we calculated FIRE scores
1315 from these current unique irreplaceability scores rather than those from an extinction-
1316 based approach, and correlated those with the original FIRE scores. There was a strong
1317 correlation between functional irreplaceability and a measure of unique current
1318 contributions to trait space for each species ($\rho = 0.63$, $p < 0.001$; Fig. B).
1319

1320
1321 **Fig. B | Correlation between future and current functional distinctiveness.** A scatterplot to compare the
1322 relationship between future functional distinctiveness (where functional complementarity is accounted for)
1323 and current functional distinctiveness (where all species are included in trait space when calculating functional
1324 distinctiveness). Correlations between future and current functional distinctiveness were significantly positive
1325 (Spearman's correlation: r = 0.96, p < 0.001).

1326
Comparing functional metrics.
To investigate the variations between different metrics, we use Pearson's correlation to
compare functional irreplaceability against other approaches to quantifying species-specific
contributions to overall functional diversity. We investigate correlations between functional
irreplaceability, functional uniqueness (FUn), functional distinctiveness (FDis) and functional
uniqueness measured in probabilistic trait space (FU). FU is calculated as the functional
distance to the nearest neighbour of the species of interest, using the uniqueness function
in the funrar package (v. 1.5.0)[3]. FDis is calculated as the average functional distance from a
species to all the others in the given community, using the distinctiveness function in the
funrar package (v. 1.5.0)[3]. FU is calculated as the mean overlap of species in probabilistic
trait space, using the uniqueness function in the TPD package (v. 1.1.0)[4]. We compared our
functional irreplaceability measure against other approaches to quantifying species-specific
contributions to overall functional diversity, and found significant but only moderate
positive relationships between them (Fig. C).
1341

1342
1343
Fig. C | Correlation between functional indices. Scatterplots to compare species-specific functional diversity
indices between; a) Functional Uniqueness (FUn) and Functional Distinctiveness (FDis; Pearson's correlation: r
= 0.56, p < 0.001), b) Functional Uniqueness and Trait Probability Density Functional Uniqueness (TPD FU;
Pearson's correlation: r = 0.57, p < 0.001), c) FDis and TPD FU (Pearson's correlation: r = 0.47, p < 0.001), d)
TPD functional distinctiveness and TPD FU (Pearson's correlation: r = 0.66, p < 0.001), e) TPD functional
distinctiveness and FUn (Pearson's correlation: r = 0.55, p < 0.001) and f) TPD functional distinctiveness and
FDis (Pearson's correlation: r = 0.43, p < 0.001).
1351
1352 *Global shark functional irreplaceability.*

The top 5% of functionally irreplaceable shark species (28 spp.) together occupy 10.85% of unique trait space. Overall, 46% of the top 5% functionally irreplaceable species are threatened with extinction, compared with 30% of all sharks[5]. To estimate the projected impact of extinction on shark trait space, we calculated the proportion of trait space that would be lost with the extinction of all 167 currently threatened species (Vulnerable [VU], Endangered [EN], Critically Endangered [CR] on the IUCN Red List). We found that 17.18% (compared to an average random sample of 11.53%; sd = 1.24) of global shark trait space is at risk of extinction, potentially increasing to 23.58% (compared to an average random sample of 20.73%; sd = 1.58) if we were to also consider all Data Deficient [DD] and Not Evaluated [NE] species as at risk of extinction. The orders Hexanchiformes, Echinorhiniformes and Lamniformes have the highest average functional irreplaceability scores of 0.20% (sd = 0.021), 0.18% (sd = 0.001) and 0.18% (sd = 0.018) respectively (Fig. D). The orders with the lowest average functional irreplaceability score are Squatiniformes and Carcharhiniformes, which both have an average functional irreplaceability score of 0.06% (sd = 0.007 and 0.009, respectively) (Fig. D).

Fig. D | Distribution of functional irreplaceability and FIRE across the tree of life. A phylogenetic tree of all sharks with branches coloured to represent the median (a) functional irreplaceability and (b) FIRE of all descendant species across the shark tree of life. Numbers indicate shark orders. Density plots represent the distribution of (a) functional irreplaceability and (b) FIRE scores in each order, both on log scale. Illustrations represent the shark species with the highest (a) functional irreplaceability and (b) FIRE score in their respective order. Shark species: (a) 1. Barred Bullhead Shark (*Heterodontus zebra*), 2. Warren's Sixgill Sawshark (*Pliotrema warreni*), 3. Bramble Shark (*Echinorhinus brucus*), 4. Smoothback Angelshark (*Squatina oculata*), 5. Angular Roughshark (*Oxynotus centrina*), 6. Bluntnose Sixgill Shark (*Hexanchus griseus*), 7. Pacific Nurse Shark (*Ginglymostoma unami*), 8. Basking Shark (*Cetorhinus maximus*), 9. Japanese Swellshark (*Cephaloscyllium umbratile*), (b) 1. Horn Shark (*Heterodontus francisci*), 2. Anna's Sixgill Sawshark (*Pliotrema annae*), 3. Bramble Shark (*Echinorhinus brucus*), 4. Smoothback Angelshark (*Squatina oculata*), 5. Angular Roughshark (*Oxynotus centrina*), 6. Broadnose Sevengill Shark (*Notorynchus cepedianus),* 7. Pacific Nurse Shark (*Ginglymostoma unami*), 8. Sand Tiger Shark (*Carcharias taurus*), 9. Scalloped Hammerhead (*Sphyrna lewini*). The phylogenetic tree was selected at random from the set of 1,000 phylogenetic trees used in this study to calculate EDGE scores. Illustrations: Marc Dando.

Supplementary figures and tables

Fig. S1 | Relationships between bird morphological traits and extinction risk. Variation in morphological trait values across IUCN Red List categories (EX = Extinct, EW = Extinct in the Wild, CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least Concern, DD = Data Deficient). Traits include: (a) beak length (culmen), (b) beak length (nares), (c) beak width, (d) beak depth, (e) tarsus length, (f) secondary length, (g) tail length, and (h) wing length. Boxplots represent the range, interquartile range and median with points denoting outliers.

1396

1397
1398 **Fig. S2 | Relationships between shark traits and extinction risk.** Variation in trait values across IUCN Red List
1399 categories (CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least
1400 Concern, DD = Data Deficient). Traits include: (a) body length, (b) trophic level, (c)maximum depth, (d) growth
1401 ratio, (e) body shape variation PC1, (f) body shape variation PC2, (g) habitat preference, and (h) reproductive
1402 guild. Boxplots (a-f) represent the range, interquartile range and median with points denoting outliers.

1403
Fig. S3 | The distribution of imputed values across the shark trait space. (a) Distribution of the proportion of
imputed data across species, projected onto the first two principal coordinate (PCoA) axes. Points are coloured
according to the proportion of imputed data (blue = low, red = high). Grey bars show marginal distributions of
imputation density across species. Panels (b–i) show locations in trait space where imputation occurred for
individual traits: (b) body length, (c) maximum depth, (d) trophic level, (e) body shape variation PC1, (f) body
shape variation PC2, (g) growth ratio, (h) reproductive guild, and (i) habitat preference. Each red point
indicates a species with imputed values for the given trait, while blue points represent species with observed
values.

Full trait space

1412
1413 **Fig. S4 | Occupation of trait space for selected species and their congeners.** Probability of occupancy of full
1414 trait space (a). Area of trait space occupied by all species in Mustelus genus (orange) and space occupied by
1415 Mustelus antarcticus (b) and Mustelus lenticulatus (c). Area of trait space occupied by all species in Somniosus
1416 genus (orange) and space occupied by Somniosus antarcticus (red; d).

1417
1418 **Fig. S5 | Stacked shark outlines.** Scaled and centred outlines to show the body shape silhouettes of 515
1419 images of shark species collected from Sharks of the World: A Complete Guide[11].
1420

1421
1422 **Fig. S6 | Harmonic power required to fit shark outline.** a) Number of harmonics required to gather cumulative
1423 harmonic power from the outlines of sharks following elliptical fourier analysis. b) Boxplots to show the
1424 harmonic power for the sum of harmonics (h1-h29) for the body shape elliptical fourier analysis. 16 harmonics
1425 were selected to gather 99% of cumulated harmonic power.
1426

1427

1428 **Fig. S7 | Principal coordinate analysis of shark body shape elliptical fourier analysis.** The first and second
1429 principal component from a principal component analysis of the body shape elliptical fourier analysis.
1430 Coordinates quantify variation in shape of shark species, with outlines representing mean body shape for areas
1431 shown on the graph.

1432

1433 **Table S1 | Trait data.** Traits used to build trait space and the functional guild each trait represents for both
1434 birds and sharks.

1435

Clade	Trait	Definition	Source	Unit	Data Type	Number of species (% coverage)	Functional Guild
Sharks	Maximum body length	The maximum total length of an individual species (where the mean maximum values is taken if sex maximum length differs)[6]	Fishbase[7]	cm	Continuous	535 (96.1%)	Related to body size, metabolism and impact on the food web[8]
	Maximum depth	The maximum reported depth for juveniles and adults of the species[6]	Fishbase[7]	m	Continuous	488 (87.6%)	Can reflect vertical mobility, swim

							performance and nutrient cycling[9]
	Trophic level	Where the species operate in their respective foodwebs[6]	Fishbase[7]	1 + mean trophic level of species food items	Continuous	278 (49.9%)	Can reflect trophic interactions and the postion occupied on a food web[8]
	Body shape variation PC1	Principal component 1 of the PCA describing the elliptial fourier analysis of each species lateral sillouette[10]	Sharks of the world[11]		Continuous	512 (91.9%)	Can reflect morphological variation, swim performance and predator-prey interactions[12]
	Body shape variation PC2	Principal component 2 of the PCA describing the elliptial fourier analysis of each species lateral sillouette[10]	Sharks of the world[11]		Continuous	512 (91.9%)	Can reflect morphological variation, swim performance and predator-prey interactions[12]
	Growth ratio	The relative increase in length of the species based on the length at birth and maximum length	Sharks of the world[11], Fishbase[7]	length (cm) / length at birth (cm)	Continuous	342 (61.4%)	Can reflect life history strategies[13,14]
	Reproductive mode	Whether the species lays eggs (oviparous) or gives birth to live young (viviparous)	Sharks of the world[11], Fishbase[7]	oviparous, viviparous, aplacental viviparous, viviparous (with a yolk sac placenta)	Categorical	557 (100%)	Can reflect life history strategies[13,14]
	Habitat preference	The particular environment preferred by the species[6]	Fishbase[7]	pelagic, benthopelagic, demersal, reef-associated, bathypelagic, bathydemersal	Categorical	469 (84.2%)	Can reflect hunting or foraging behaviour, nutrient cycling and microhabitat formation[15]
Birds	**Beak culmen length**	Length from the tip of the beak to the base of the skull[16]	AVONET[16]	mm	Continuous	11005 (100%)	Diet; feeding action (e.g. probing, cracking, hammering); food type[17,18]
	Beak nares length	Length from the anterior edge of the nostrils to the tip of the beak[16]	AVONET[16]	mm	Continuous	11005 (100%)	Diet; feeding action (e.g. probing, cracking, hammering); food type[17,18]
	Beak width	Width of the beak at the anterior edge of the nostrils[16]	AVONET[16]	mm	Continuous	11005 (100%)	Diet; feeding action (e.g. probing, cracking, hammering); food type[17,18]
	Beak depth	Depth of the beak at the anterior edge of the nostrils[16]	AVONET[16]	mm	Continuous	11005 (100%)	Diet; feeding action (e.g. probing, cracking,

						hammering); food type[17,18]
Tarsus length	Length of the tarsus from the posterior notch between tibia and tarsus, to the end of the last scale of acrotarsium (at the bend of the foot) [16]	AVONET[16]	mm	Continuous	11005 (100%)	Foraging niche; foraging strategy (e.g. walking, sallying, gleaning)[19–22]
Wing length	Length from the carpal joint (bend of the wing) to the tip of the longest primary on the unflattened wing[16]	AVONET[16]	mm	Continuous	11005 (100%)	Foraging niche; foraging strategy (e.g. walking, sallying, gleaning)[19–22]
Length to first secondary	Length from the carpal joint (bend of the wing) to the tip of the first secondary, i.e. the outermost secondary adjacent to the innermost primary feather. Secondary1 is roughly equivalent to Wing length minus Kipp's distance (measured in a fully folded and flat wing)[16]	AVONET[16]	mm	Continuous	11005 (100%)	Foraging niche; foraging strategy (e.g. walking, sallying, gleaning)[19–22]
Tail length	Distance between the tip of the longest rectrix and the point at which the two central rectrices protrude from the skin, typically measured using a ruler inserted between the two central rectrices[16]	AVONET[16]	mm	Continuous	11005 (100%)	Foraging niche; foraging strategy (e.g. walking, sallying, gleaning)[19–22]

1436

1437

1438 **Table S2 |** ANOVA pairwise comparisons between prioritisation strategies at capturing intentionally exploited
1439 bird and shark species with Tukey's Honestly Significant Difference post hoc test.

Clade	Percentage of total species	Prioritisation	Difference in means	Lower bounds	Upper bounds	p
Birds	10%	Functional distinctiveness-Evolutionary distincitiveness	120.07	119.05	121.09	< 0.001
		Random-Functional distinctiveness	-128.783	-129.80126	-127.76475	< 0.001
		Random-Evolutionary distinctiveness	-8.71	-9.73	-7.69	< 0.001
	20%	Functional distinctiveness-Evolutionary distincitiveness	191.93	190.67	193.19	< 0.001
		Random-Functional distinctiveness	-203.393	-204.65105	-202.13495	< 0.001
		Random-Evolutionary distinctiveness	-11.47	-12.72	-10.21	< 0.001
	30%	Functional distinctiveness-Evolutionary distincitiveness	234.27	232.86	235.67	< 0.001
		Random-Functional distinctiveness	-248.35	-249.76	-246.95	< 0.001
		Random-Evolutionary distinctiveness	-14.09	-15.49	-12.68	< 0.001
	40%	Functional distinctiveness-Evolutionary distincitiveness	254.14	252.66	255.61	< 0.001
		Random-Functional distinctiveness	-267.59	-269.06	-266.11	< 0.001
		Random-Evolutionary distinctiveness	-13.45	-14.93	-11.98	< 0.001
	50%	Functional distinctiveness-Evolutionary distincitiveness	255.90	254.43	257.37	< 0.001
		Random-Functional distinctiveness	-266.729	-268.2035	-265.2545	< 0.001
		Random-Evolutionary distinctiveness	-10.83	-12.30	-9.35	< 0.001
Sharks	10%	Functional distinctiveness-Evolutionary distincitiveness	0.98	0.59	1.36	< 0.001
		Random-Functional distinctiveness	-5.95	-6.33	-5.57	< 0.001
		Random-Evolutionary distinctiveness	-4.97	-5.36	-4.59	< 0.001
	20%	Functional distinctiveness-Evolutionary distincitiveness	2.33	1.85	2.81	< 0.001
		Random-Functional distinctiveness	-10.90	-11.38	-10.42	< 0.001
		Random-Evolutionary distinctiveness	-8.57	-9.05	-8.09	< 0.001
	30%	Functional distinctiveness-Evolutionary distincitiveness	4.46	3.94	4.99	< 0.001
		Random-Functional distinctiveness	-14.83	-15.35	-14.30	< 0.001
		Random-Evolutionary distinctiveness	-10.36	-10.89	-9.84	< 0.001
	40%	Functional distinctiveness-Evolutionary distincitiveness	7.15	6.62	7.68	< 0.001
		Random-Functional distinctiveness	-18.44	-18.97	-17.91	< 0.001
		Random-Evolutionary distinctiveness	-11.28	-11.81	-10.75	< 0.001
	50%	Functional distinctiveness-Evolutionary distincitiveness	10.82	10.29	11.34	< 0.001
		Random-Functional distinctiveness	-20.97	-21.49	-20.44	< 0.001
		Random-Evolutionary distinctiveness	-10.15	-10.67	-9.62	< 0.001

1440

1441

1442 **List of Supplementary data**

1443

1444 Data S1 – Bird indices

1445

1446 Data S2 – Shark indices

1447

1448 Data S3 – Trait database for birds

1449

1450 Data S4 – Trait database for sharks

1451

1452 Data S5 – PCA shape data

1453

References:

1. Pigot, A. L. *et al.* Macroevolutionary convergence connects morphological form to ecological function in birds. *Nat Ecol Evol* **4**, 230–239 (2020).

2. Williams, P. J. & Curley, S. R. Functional and phylogenetic convergence of winter and breeding bird communities in the northeastern US. *Ecography* e07717 (2025) doi:10.1002/ecog.07717.

3. Grenié, M., Denelle, P., Tucker, C. M., Munoz, F. & Violle, C. funrar: An R package to characterize functional rarity. *Diversity and Distributions* **23**, 1365–1371 (2017).

4. Carmona, C. P., de Bello, F., Mason, N. W. H. & Lepš, J. Trait probability density (TPD): measuring functional diversity across scales based on TPD with R. *Ecology* **100**, e02876 (2019).

5. IUCN Red List of Threatened Species. The IUCN Red List of Threatened Species. *IUCN Red List of Threatened Species* https://www.iucnredlist.org/en (2025).

6. Froese, R. & Pauly, D. FishBase 2000: Concepts, design and data sources. *ICLARM* (2000).

7. Froese, R. & Pauly, D. FishBase. www.fishbase.org (2023).

8. Henderson, C. J. *et al.* Long term declines in the functional diversity of sharks in the coastal oceans of eastern Australia. *Commun Biol* **7**, 1–8 (2024).

9. Andrzejaczek, S. *et al.* Diving into the vertical dimension of elasmobranch movement ecology. *Science Advances* **8**, eabo1754 (2022).

10. Siders, Z. A. *et al.* Functional and phylogenetic diversity of sharks in the Northeastern Pacific. *Journal of Biogeography* **n/a**, (2022).

11. Ebert, D. A., Dando, M. & Fowler, S. *Sharks of the World: A Complete Guide*. (Princeton University Press, 2021).

12. Sternes, P. C. & Shimada, K. Body forms in sharks (Chondrichthyes: Elasmobranchii) and

their functional, ecological, and evolutionary implications. *Zoology* **140**, 125779 (2020).

13. Cortés, E. Life history patterns, demography, and population dynamics. *Biology of sharks*

and their relatives 449–469 (2004).

14. Cortés, E. Life History Patterns and Correlations in Sharks. *Reviews in Fisheries Science* **8**,

299–344 (2000).

15. Sorenson, L., Santini, F. & Alfaro, M. E. The effect of habitat on modern shark

diversification. *Journal of Evolutionary Biology* **27**, 1536–1548 (2014).

16. Tobias, J. A. *et al.* AVONET: morphological, ecological and geographical data for all birds.

Ecology Letters **25**, 581–597 (2022).

17. Felice, R. N., Tobias, J. A., Pigot, A. L. & Goswami, A. Dietary niche and the evolution of

cranial morphology in birds. *Proceedings of the Royal Society B: Biological Sciences* **286**,

20182677 (2019).

18. Sayol, F., Reijenga, B. R., Tobias, J. A. & Pigot, A. L. Ecophysical constraints on avian

adaptation and diversification. *Current Biology* **35**, 1326-1336.e6 (2025).

19. Fitzpatrick. Tail length in birds in relation to tail shape, general flight ecology and sexual

selection. *Journal of Evolutionary Biology* **12**, 49–60 (1999).

20. Einoder, L. D. & Richardson, A. M. M. Aspects of the Hindlimb Morphology of Some

Australian Birds of Prey: A Comparative and Quantitative Study. *Auk* **124**, 773–788

(2007).

21. Miles, D. B. & Ricklefs, R. E. The Correlation Between Ecology and Morphology in

Deciduous Forest Passerine Birds. *Ecology* **65**, 1629–1640 (1984).

22. Miles, D. B., Ricklefs, R. E. & Travis, J. Concordance of Ecomorphological Relationships in

Three Assemblages of Passerine Birds. *The American Naturalist* **129**, 347–364 (1987).

1502
1503