Sensory-Behavioral Deficits in Parkinson's Disease: Insights from a 6-OHDA Mouse Model

Savannah R. Linen[1], Nelson H. Chang[2], Ellen J. Hess[3], Garrett B. Stanley[2], Christian Waiblinger[2#]

[1]Program in Bioinformatics, Georgia Institute of Technology, Atlanta, GA, USA
[2]Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA, USA
[3]Departments of Pharmacology and Chemical Biology and Neurology, Emory University, Atlanta, GA USA

#Correspondence: Dr. Christian Waiblinger
                    Email: christian.waiblinger@gatech.edu

Running Title: Sensory Deficits in a 6-OHDA Mouse Model

With 24 text pages and 5 figures

Word counts: Abstract 238, Intro 488, Discussion 1159.

Acknowledgements: S.R.L. was supported by a McCamish Parkinson's Disease Innovation Program Blue Sky grant. E.J.H was supported by a McCamish Parkinson's Disease Innovation Program Blue Sky grant and NIH R01NS124764. N.H.C., G.B.S and C.W. were supported by NIH Brain Grants RF1NS128896 and R01NS104928. This study was supported by the Emory University Emory Integrated Cellular Imaging Core Facility (RRID:SCR_023534), with special thanks to April Reedy for her advice on experimental design and image acquisition. We thank the Emory HPLC Bioanalytical Core (EHBC), which was supported by the Emory University School of Medicine and the Georgia Clinical & Translational Science Alliance of the National Institutes of Health under Award Number UL1TR002378. Additionally, we thank Aqua Asberry from the Georgia Institute of Technology's Research Histology Core for her assistance in specimen embedding and cryostat sectioning.

Author contributions: C.W. and G.B.S. conceived the project. S.R.L. and C.W. designed the study. S.R.L., N.H.C. and C.W. carried out all experiments. S.R.L. and C.W. analyzed the data. S.R.L., C.W., E.J.H, and G.B.S. wrote the manuscript.

Conflict of Interest: The authors declare no competing interests

Abstract

Parkinson's disease (PD) is characterized by the degeneration of dopaminergic neurons in the
striatum, predominantly associated with motor symptoms. However, non-motor deficits,
particularly sensory symptoms, often precede motor manifestations, offering a potential early
diagnostic window. The impact of non-motor deficits on sensation behavior and the underlying
mechanisms remains poorly understood. In this study, we examined changes in tactile sensation
within a Parkinsonian state by employing a mouse model of PD induced by 6-hydroxydopamine
(6-OHDA) to deplete striatal dopamine (DA). Leveraging the conserved mouse whisker system as
a model for tactile-sensory stimulation, we conducted psychophysical experiments to assess
sensory-driven behavioral performance during a tactile detection task in both the healthy and
Parkinson-like states. Our findings reveal that DA depletion induces pronounced alterations in
tactile sensation behavior, extending beyond expected motor impairments. We observed diverse
behavioral deficits, spanning detection performance, task engagement, and reward
accumulation, among lesioned individuals. While subjects with extreme DA depletion
consistently showed severe sensory behavioral deficits, others with substantial DA depletion
displayed minimal changes in sensory behavior performance. Moreover, some exhibited
moderate degradation of behavioral performance, likely stemming from sensory signaling loss
rather than motor impairment. The implementation of a sensory detection task is a promising
approach to quantify the extent of impairments associated with DA depletion in the animal
model. This facilitates the exploration of early non-motor deficits in PD, emphasizing the
importance of incorporating sensory assessments in understanding the diverse spectrum of PD
symptoms.

SIGNIFICANCE STATEMENT

This study explores sensory-motor aspects of Parkinson's disease using a 6-OHDA mouse model.
Leveraging the mouse whisker system, we reveal diverse deficits in tactile sensation behavior
due to dopamine depletion. Our findings emphasize the importance of sensory assessments in
understanding the diverse spectrum of PD symptoms.

Introduction

Parkinson's disease (PD) is a prevalent neurodegenerative disorder, affecting approximately 1 million individuals in the United States annually, with 90,000 new diagnoses reported each year. Diagnosis is based on identifying common motor symptoms such as tremor, bradykinesia (slowed movement), and muscular rigidity. However, PD patients also experience non-motor impairments, including alterations in olfactory, auditory, tactile, nociceptive, thermal, and proprioceptive perception (Oppo et al. 2020; Jafari, Kolb, and Mohajerani 2020; Kesayan et al. 2015; José Luvizutto et al. 2020; Brim and Struhal 2021). Notably, these non-motor symptoms, often referred to collectively as sensory symptoms, precede the manifestation of motor symptoms by two or more years (Pont-Sunyer et al. 2015), offering a potential window for early diagnostic methods.

In the realm of PD research, animal models have become instrumental in understanding the cellular and circuit mechanisms underlying both motor and non-motor deficits, as well as in the development of therapeutic strategies. The 6-hydroxydopamine (6-OHDA) rodent model, involving the injection of the neurotoxin 6-OHDA into the medial forebrain bundle, is a widely used model that induces dopamine (DA) depletion in the striatum, mimicking Parkinson-like symptoms. Originally designed to study motor deficits (Brooks and Dunnett 2009; Francardo et al. 2011; Lundblad et al. 2004), this model has been extended to investigate non-motor deficits in sensory circuits (Ketzef et al. 2017; Reig and Silberberg 2014). However, questions persist regarding how animals behave and interact with their sensory environment, specifically how the perception of sensory stimuli and behavioral response are altered in the DA-depleted state.

Here, we investigate the behavioral aspects of mice trained on a sensory-based detection task in both healthy and Parkinson-like states induced by 6-OHDA lesioning. Leveraging the rodent whisker system for tactile-sensory stimulation, we anticipate reduced task performance in the Parkinson-like state. In addition to the standard behavioral readout of motor impairments, we designed a psychophysical experiment to systematically evaluate task performance in both states. As a complement to conventional measures (Aeed et al. 2021; Branchi et al. 2008), here we highlight the sensory dimension of the task in behaving animals. We hypothesized that a detailed analysis of behavioral readouts and lesion level assessment would reveal specific alterations in sensory processing and subsequent behavioral responses in the DA-depleted state.

Our findings unveil a spectrum of deficits following DA depletion, encompassing rotational asymmetry, weight loss, and varying degrees of altered tactile sensation evident in our behavioral detection task. Despite observing substantial DA depletion in the majority of mice, those displaying severe sensory behavioral deficits consistently exhibited pronounced impairments across all measures. However, some mice with substantial DA depletion, rotational bias, and weight loss displayed only minor sensory behavioral deficits. Others strikingly demonstrated diminished sensory behavioral performance without evident motor function loss, suggesting the emergence of a sensory deficit. Taken together, these results highlight the complex interactions between these factors and underscore the importance of developing a more comprehensive understanding of the effects of DA depletion across sensory, motor, and cognitive functions.

Methods

Experimental model and subject details

All experiments and surgical procedures were performed in accordance with the guidelines approved by the Georgia Institute of Technology Institutional Animal Care and Use Committee and conformed to guidelines established by the NIH. Subjects were 12 male mice (C57/BL/6J, Jackson Laboratories), aged 16-20 weeks at the age of experimentation. Only males were used in this study because the incidence rate of PD is 1.5 times higher in men than in women (Wooten 2004). Animals were housed together in pairs of two per cage (following surgical recovery when applicable) and housed in an inverted light cycle (light: 7pm-7am: dark: 7am-7pm) to ensure experiments occurred during animals' wakeful periods. Housing temperatures remained at 65-75°F (18-23°C) with 40-60% humidity.

Head-plate implantation

At least 1 week before experimentation, animals were anesthetized with isoflurane, 5% in an induction chamber, and maintained at 1-3% in a stereotaxic frame. Following anesthetization and analgesia, an incision was made across the length of the skull and the surrounding connective tissues and muscles were removed with a scalpel blade. A ring-shaped (inner radius 5 mm) titanium headplate was then attached to the skull to allow for head fixation during experimentation using MetaBond, a three-stage dental acrylic (Parkell, Inc) (Waiblinger et al. 2022). MetaBond was made on ice, allowed to thicken, applied to the skull, and allowed to set for five minutes to attach the headplate. The skull was then covered with a thin layer of MetaBond followed by a silicone elastomer (Kwik-Cast Sealant, World Precision Instruments) to protect future injection sites. During the procedure animals were placed on a heating pad to keep body temperature stable. Sterile techniques were employed throughout the procedure to minimize chances of infection, and no antibiotics were given. Opioid and non-steroidal anti-inflammatory analgesics were administered (SR-Buprenorphine, SC, post operatively and Ketoprofen, IP, post-operatively). Habituation to head-fixation and training began once animals had fully recovered from head-plate implantation.

Sensory detection task and training

Behavioral training and testing were conducted using a sensory-driven detection task, utilizing the vibrissa/whisker pathway of the mouse. During behavioral training and testing, water intake was restricted to experimental sessions where animals could earn water to satiety. For two days every week, testing was paused, and animals were given access to water *ad libitum*. Bodyweight was monitored daily and remained relatively stable during constant training. If weight dropped more than ~5 g, supplementary water was given outside of training sessions to maintain the animal's body weight. We conducted 1-2 training sessions each day comprising 50-300 trials. All experiments were performed in the dark to ensure no visual identification of the galvo-motor whisker actuator. A constant auditory white noise background noise (70 dB) was produced by an arbitrary waveform generator to mask any sound emission from the galvo-motor whisker actuator. All mice were trained on a Go/No-Go detection task employing a protocol as described previously (Ollerenshaw et al. 2012; Stüttgen, Rüter, and Schwarz 2006; Waiblinger et al. 2018, 2019, 2022). In this task, the whisker is deflected at intervals of 6-8 s (flat probability distribution)

with a single pulse (detection target). Trials were sorted into four categories, consisting of a "hit",
"miss", "correct rejection", or "false alarm". A trial was considered a "hit" if the animal generated
the "Go" response (a lick at a waterspout within 1000 ms of target onset). If the animal did not
lick in response to a whisker deflection the trial was considered a "miss". Additionally, catch trials
in which no deflection of the whisker occurred (A = 0°) were considered a "correct rejection" if
the animal did not lick (No-Go). If random licks did occur within 1000 ms of catch onset however,
the trial was classified as a "false alarm". Licking that occurred within a 2 s window prior to
stimulus onset was punished by resetting time (time-out) and beginning a new inter-trial interval
of 6-8 s, drawn at random from a flat probability distribution. These trial types were excluded
from the main data analysis.
153
Whisker Stimulation
Whisker deflections were carried out using a calibrated galvo-motor (galvanometer optical
scanner model 621OH, Cambridge Technology) as described previously (Chagas et al. 2013).
Dental cement was used to narrow the opening of the rotating arm to prevent whisker motion
at the insertion site. A single whisker was inserted into the rotating arm of the motor on the right
side of the mouse's face at a 5 mm (±1 mm tolerance) distance from the skin, directly stimulating
the whisker shaft and largely overriding bio elastic whisker properties (Waiblinger et al. 2022).
All whiskers were trimmed to avoid artifacts generated by the arm touching other whiskers.
Across mice, the whisker chosen differed (C1, C2, D1, or D2) but the same was used throughout
the experimentation within each mouse. Custom written code in Matlab and Simulink was
designed to elicit voltage commands for the actuator (Ver. 2015b; The MathWorks, Natick,
Massachusetts, USA). Stimuli were defined as a single event, a sinusoidal pulse (half period of a
100 Hz sine wave, starting at one minimum and ending at the next maximum). The pulse
amplitudes used A = [0, 2, 4, 8, 16]°, correspond to maximal velocities: Vmax = [0 628 1256 2512
5023]°/s or mean velocities: Vmean = [0 408 816 1631 3262]°/s and were well within the range
reported for frictional slips observed in natural whisker movement (Ritt, Andermann, and Moore
2008; Waiblinger et al. 2022; Wolfe et al. 2008). Figure 2A summarizes the behavior setup,
stimulus delivery and logic of the task.
172
6-OHDA lesions
After the collection of behavioral data in the healthy state (pre-lesion, at least 8 sessions of
detection testing and one session of rotation), unilateral lesions with 6-OHDA were performed in
a subset of animals (6-OHDA, n=10; sham, n=2). The procedures of 6-OHDA injection were
adapted from recent studies (Bagga, Dunnett, and Fricker 2015; Ketzef et al. 2017). Animals were
anesthetized with isoflurane and mounted on a stereotaxic frame as described in the head-plate
implantation section above. Kwik cast was removed, and a small craniotomy was created AP: -
1.2 mm, ML: +1.2 mm, relative to bregma according to the Mouse Brain Atlas in Stereotaxic
Coordinates (Frankin and Paxinos 2008). A unilateral injection of 0.2ul of 6-OHDA (15 mg/ml
solution 6-OHDA.HBr) was then administered into the left median forebrain bundle at a rate of
0.1ul/min (depth 4.8-5 mm according to the Mouse Brain Atlas in Stereotaxic Coordinates) via a
Hamilton syringe (Hamilton Neuros Syringe 700/1700) (Frankin and Paxinos 2008; Thiele, Warre,
and Nash 2012). Five-minute waiting periods following both injector insertion and removal were
implemented to allow for tissue relaxation. Sham lesioned mice received the correspondent

injection protocol with the same volume of vehicle (0.9% saline and 0.02% ascorbic acid).
Craniotomies were then sealed with a thin layer of MetaBond and the exposed skull was covered
with a silicone elastomer. Opioid and non-steroidal anti-inflammatory analgesics were
administered (SR-Buprenorphine, SC, post operatively and Ketoprofen, IP, post-operatively).
Animals were allowed to recover for 7–21 days until activity and body weight were normal and
then behavioral testing continued.
193
Post-lesion care
6-OHDA lesion mice can suffer from a variety of post-operative complications requiring special
post-operative care and a continuous health assessment (Masini et al. 2021). After the surgery,
animals were placed in single cage and monitored until fully awake. Additional warmth was
provided as an infrared light source. In addition to the standard post-operative monitoring and
free access to food and water, a maximum of 3ml sterile saline per day was administered SC to
encourage rehydration in lesioned mice. Furthermore, lesioned mice received nutritional
supplement in form of DietGel Boost (ClearH$_2$O) as well as sweetened condensed milk (ClearH$_2$O)
in food containers on the floor of the cage for 3-10 days (or until body weight stabilized) to
increase survival and maintain healthy body weight. The health status of individual mice was
assessed daily for a period of 7-21 days post-surgery until activity and body weight were normal.
Animals that reached the humane endpoint (defined by the ethical permit) were euthanized.
206
Rotation
Unilateral 6-OHDA lesions typically cause highly specific, reproducible rotational measurements
and the rotation test is a standard measurement of 6-OHDA lesion efficacy (Brooks and Dunnett
2009). After unilateral median forebrain bundle DA depletion, a postural bias towards the side of
the lesion is exhibited. Ipsilateral rotation is driven by an imbalance of DA between hemispheres
generating decreased movement on the side of lesion. Mice were placed in a clear plexiglass
cylinder (40 cm diameter, 20 cm height) and attached via a flexible harness to an automated
counter (Rotation Sensors LE902SR, Container & Sensor Support LE902RP, and Individual Counter
LE902CC, Panlab). Ipsilateral and contralateral turns relative to the site of the lesion were
recorded for 45 minutes. Turns were counted at 45 degrees and findings were presented as
percent left turns, where one turn is equivalent to 360 degrees (8 x 45 degrees). The test was
performed for all subjects both prior to and following 6-OHDA (n=10) or sham injection (n=2).
219
TH assessment via immunofluorescence
Following experimentation, animals were transcardially perfused using a 4% paraformaldehyde
(PFA) solution in phosphate-buffered saline (PBS) (Gage, Kipke, and Shain 2012). Brains were then
extracted and post-fixed in PFA for 24 hr before being embedded in OCT for long-term storage at
-80°C until sectioning and immunofluorescence. Immunofluorescence staining for tyrosine
hydroxylase (TH) was performed as described previously (Ketzef et al. 2017). Sections of 15 uL
were collected using a cryostat (CryoStar NX70), rinsed in phosphate buffered saline (PBS) and
subsequently incubated for 1 hr in blocking solution consisting of 5% normal goat serum, 0.3%
Triton X-100, and 1% BSA in PBS. Following permeabilization, sections were again rinsed in PBS,
then incubated with primary antibody, anti-TH rabbit monoclonal antibody (Millipore Sigma,
Billerica, MA) diluted 1:500 in BSA-PBS (1%) at 4°C for 24 hr. Sections were then rinsed with PBS

and incubated with Cy3-conjugated goat anti-rabbit secondary antibody (Millipore Sigma,
Billerica, MA; Jackson Immunoresearch, Philadelphia, PA) diluted 1:800 in BSA-PBS (1%).
Following staining, sections were imaged using either a Zeiss 900 confocal microscope (Carl Zeiss,
Jena, Germany) at 20x magnification or Leica Stellaris 8 (Leica, Wetzlar, Germany). Images were
quantified using ImageJ to assess percent fluorescence change between lesioned and non-
lesioned hemispheres.
237
<u>TH assessment via high performance liquid chromatography</u>
Sample Preparation & Protein Assay. The efficacy of lesioning was assessed using high
performance liquid chromatography (HPLC). Mice were sacrificed in the same way as described
previously, but no fixation was performed prior to tissue extraction (Gage, Kipke, and Shain
2012). Brain-hemispheres were then separated into non-lesioned (right) and lesioned (left) and
placed immediately on dry ice for flash freezing to preserve monoamine concentrations,
specifically DA, in the tissue. The tissue was resuspended in 600 uL 0.1 PCA just above freezing
temperature and probe sonicated (Branson 450 Digital Sonifier with microtip, Marshall Scientific)
on ice with setting 3 and 30% duty cycle. The homogenates were subsequently centrifuged at 4°C
at 10,000 x g for 15 min. Supernatants were transferred to 0.22 uM polyvinylidene fluoride
polymer (PVDF) microcentrifuge filter tubes and any remaining matter was removed via filtration
through spin filter at 800 rpm for 5 minutes. Concentrations of monoamines were acquired
through reverse phase HPLC with electrochemical detection. 1000 uL 2% Sodium dodecyl sulfate
(SDS) was used to dissolve protein pellets. Quantification of protein was carried out in triplicate
96-well microplates with SpectraMax M5e spectrophotometer (Molecular Devices, Sunnyvale,
CA) using the BCA method (Pierce BCA Protein Assay Kit, Thermo Scientific).
254
HPLC Conditions. An ESA 5600A CoulArray detection system equipped with an ESA Model 584
pump and an ESA 542 refrigerated autosampler was utilized to perform HPLC. Separations were
performed at 28°C via a Hypersil 150 x 3 mm (3uM particle size) C18 column. The mobile phase
consisted of 1.6 mM 1-octanesulfonic acid sodium, 75 mM NaH_2PO_4, 0.025% triethylamine, and
8% acetonitrile at pH 3. A 25 uL sample was injected and eluted isocratically at 0.4 mL/min. A
6210 electrochemical cell (ESA, Bedford, MA) equipped with 5020 guard cell with potential set at
475 mV was used for sample detection. Analytical cell potentials were –175, 200, 350, and 425
mV. Analytes were identified by matching retention time to known standards (Sigma Chemical
Co., St. Louis MO.) and compounds were quantified by comparing peak areas to those of
standards on the dominant sensor.
265
266 <u>Experimental timeline</u>
267 Figure 2B outlines the experimental timeline. Over the course of approximately eight weeks,
268 animals completed an entire set of experiments as described below.
269
270 *Habituation.* Habituation to head-fixation began once animals had fully recovered from head-
271 plate implantation for approximately 1 week. Mice were trained to tolerate head-fixation and
272 got accustomed to the experimental chamber for approximately 1 week.
273

Detection Training. Naïve mice received uncued single whisker stimulations in form of a single
pulse (A = 16°, P_{stim} = 0.8) interspersed by catch trials or no stimulation (A = 0°, P_{catch} = 0.2). In the
early training stage, a water droplet became available immediately after stimulus offset
regardless of the animal's action to condition the animal's lick response and shape the stimulus
reward association. Once animals displayed stable consumption behavior (usually 1-2 sessions),
water was only delivered after an indicator lick of the spout within 1000 ms, transitioning the
task into an operant conditioning paradigm where the response is only reinforced by reward if it
is correctly emitted following stimulus. Learning was measured by calculating the hit p(hit) and
false alarm rate p(fa) of successive daily training sessions. A criterion of p(hit) - p(fa) ≥ 0.75 was
used to determine successful acquisition of the task. Once an animal reached the criterion, they
were considered experts at the task and proceeded into the "Detection Testing" phase.
285
Detection Testing. In this phase, the psychometric curve was measured in repeated daily sessions
(1-2 sessions per day) for a minimum of six days. The psychometric curve was measured for all
animals using a constant stimuli method entailing the presentation of repeated stimulus blocks
containing multiple stimulus amplitudes (A = [0, 2, 4, 8, 16]°). In a single trial, one of multiple
stimuli was presented after a variable time interval (6-8 s), each with equal probability (uniform
distribution, P = 0.2). A stimulus block consisted of a trial sequence containing all stimuli and a
catch trial in a pseudorandom order (e.g. each stimulus is presented once per block). One
behavioral session consisted of repeated stimulus blocks until the animal disengaged from the
task. Animals had to complete a minimum of n=50 trials and a session was considered complete
once an entire block of stimuli (A = [0, 2, 4, 8, 16]°) was missed. Therefore, trial numbers varied
across animals and sessions (n=50-300 trials). The flexible session length ensures that the data is
not affected by the animal's potential impulsivity or satiation on any given day. Following the six-
day testing period, animals were unilaterally injected with 6-OHDA and allowed 7-14 days of
recovery. Following recovery animals were again subjected to a "Detection Testing" period which
was identical to the one before the lesion to determine the extent of sensory impairments
following 6-OHDA lesioning.
302
Length of testing. Over about eight weeks, animals completed a whole set of experiments (Fig.
2b). Note, sessions do not directly correlate with days as two sessions per day were performed
in some instances and testing was paused during recovery periods. To resolve the actual time
frame of testing, we tracked task performance across days, considering different recovery
periods (Fig. 4-1 and table 1). In cases where testing was repeated after longer periods (up to 50
days post-lesion), no differences in behavior was observed. N=2 out of 12 animals reached the
humane endpoint likely due to 6-OHDA lesioning and were euthanized before the end of the
normal testing period.

Animal ID	Pre-lesion		Recovery	Post-lesion	
	Days	Sessions	Days	Days	Sessions
21	6	6	13	2*	3
22	6	6	13	14	7
25	6	8	14	15	8
26	6	10	14	13	10
29	5	8	23	7	8
30	7	9	33	3	5
31	8	8	7	6	13
32	6	8	15	1*	2
33 (sham)	5	8	19	5	8
34 (sham)	5	8	19	5	8
35	6	6	8	7	8
36	6	6	8	7	8

311   Table 1: Number of sessions and days for each animal pre-lesion, in recovery, and post-lesion.
312   *Animals euthanized due to humane endpoint.
313
<u>Experimental Design and Statistical Analysis</u>
Body weight. The weight of each subject was measured daily during detection training, detection
testing and recovery phases post surgeries. Percent body weight was calculated by identifying
the minimum weight within the recovery period (after 6-OHDA and sham injection). Numbers
were normalized to each animals' weight on the day before the lesion. A two-sided Wilcoxon
rank-sum test was used to compare the significance of the effects of lesions on body weight in
lesioned versus healthy animals.
321
322   *Rotation test.* The rotation test was performed within subjects, prior to and following 6-OHDA
323   (n=10) or sham injection (n=2). Each animal performed the test twice, once before and once after
324   injection. A two-sided Wilcoxon rank-sum test was used to compare the significance of the effects
325   of lesions on rotation in lesioned versus healthy animals.
326
327   *Detection Training.* Learning was measured for all subjects by calculating the hit p(hit) and false
328   alarm rate p(fa) of successive daily training sessions. The learning curve was measured by
329   calculating a dprime, d'_{behav}, which quantifies the effect size from the observed hit rate and false
330   alarm rate of each training session
331
332    $$d'behav = Z(hit) - Z(fa) \qquad (1)$$
333
334   where the function Z(p), p ∈ [0,1], is the inverse of the cumulative distribution function of the
335   Gaussian distribution. A criterion of d' = 2.3 (calculated with p(hit) = 0.95 and p(fa) = 0.25) was
336   used to determine the end of the detection training period.
337
Detection Testing. The psychometric experiment was performed within subjects, prior to and
following 6-OHDA (n=10) or sham injection (n=2). The psychometric curve was measured in
repeated daily sessions (1-2 sessions per day) for at least six days.

For average psychometric curves across mice (Fig. 2), response probabilities were averaged from
all animals that performed the task pre- and post-lesion. For the analysis of individual subjects
(Fig. 3), psychometric data was assessed as response-probabilities averaged across sessions
within a given stimulus condition.
345
346   Psychometric curves were fitted using Psignifit (Frund, Haenel, and Wichmann 2011; Wichmann
347   and Hill 2001). Briefly, a constrained maximum likelihood method was used to fit a modified
348   logistic function with 4 parameters: α (the displacement of the curve), β (related to the inverse
349   of slope of the curve), γ (the lower asymptote or guess rate), and λ (the higher asymptote or lapse
350   rate) as follows:
351

352 $$P(GO|s_i) = y + (1 - y - \lambda)\frac{1}{1+\exp(-z(s_i))} \qquad\qquad (2a)$$

354 $$z(s_i) = \frac{s_i - a}{\beta} \qquad\qquad (2b)$$

355
356   where s_i is the stimulus on the i^{th} trial. Response thresholds were calculated from the average
357   psychometric function for a given experimental condition using Psignifit. The term "response
358   threshold" refers to the inverse of the psychometric function at some performance level with
359   respect to the stimulus dimension. Throughout this study, we use a performance level of 50%
360   (probability of detection of 0.5).
361
362   To assess the effects of the lesion on detection behavior, the psychometric curves and response
363   thresholds were compared in lesion and sham lesion animals, from before and after the injection.
364   Statistical differences between psychophysical curves were assessed using bootstrapped
365   estimates of 95% confidence limits for the response thresholds provided by the Psignifit toolbox.
366
Impairment severity. Based on the difference in response threshold (at P=0.5) before and after
6-OHDA lesion, animals were classified into three severity groups: "minor", "moderate" and
"severe" deficit. Animals with a minor deficit show no or very small changes in performance
(thresh$_{post-pre}$ <1, n=3). Animals with a moderate deficit show a clear decrease in performance
(thresh$_{post-pre}$>1 and <10, n=4), yet they are still performing the task. Animals with a severe deficit
display a substantial drop in performance (thresh$_{post-pre}$>10, n=3) and even stop performing the
task.
374
375

Results

The current study investigates sensory behavioral deficits in the 6-hydroxydopamine (6-OHDA) mouse model of PD. To assess the impact of dopamine (DA) depletion on sensory signaling, we utilized a trained detection behavior paradigm in the vibrissa system of the mouse. Unilateral 6-OHDA injections into the mouse medial forebrain bundle (MFB) were performed, and striatal DA depletion was confirmed through histology and standard behavioral assays. Behavioral performance in the sensory-driven detection task was then evaluated over time in both healthy and PD-like states to examine non-motor deficits. Finally, all metrics were compared to provide a comprehensive understanding of the effects of the DA depletion.

Validation of the 6-OHDA mouse model

Figure 1 illustrates the validation of the 6-OHDA mouse model through unilateral injections into the left MFB. DA innervation in the striatum was verified by assessing tyrosine hydroxylase (TH) immunofluorescence. In a control mouse, both right and left striatum are intact with no visible difference in fluorescence between hemispheres (Fig. 1A). In contrast, an animal injected with 6-OHDA shows an extensive reduction (81%) in DA innervation in the lesioned versus the non-lesioned hemisphere (Fig. 1B). The amount of DA loss was consistent among all mice that underwent histological validation (median loss: 77.25%, quantified via immunofluorescence, n=8; and 92.7% quantified via high performance liquid chromatography, n=1).

To further validate DA depletion, all mice underwent a standard rotameter test before and after 6-OHDA injection (Fig. 1C). This test assesses postural bias, reflecting an imbalance of DA between hemispheres and impairing motor abilities on the injection side (Brooks and Dunnett 2009). Prior to 6-OHDA injection, mice exhibited fairly symmetric rotation behavior (median left turns: 52%, n=10). Following 6-OHDA injection, the majority of mice showed a rotational bias consistent with the injection side (median left turns: 87.3%, n=10). Notably, seven mice displayed a clear bias (<80%), while the remaining three showed a minor deficit comparable to sham controls (~60%). To ensure uniformity of lesioning effects, conventional methods often exclude animals with minimal rotation bias (Przedborski and Jackson-Lewis 1995; Ungerstedt and Arbuthnott 1970). In contrast, we conducted a thorough analysis across all animals in this study, including all data to account for potential variability in the effects of 6-OHDA injections among subjects. Our approach, as demonstrated in the results, incorporates a wide range of metrics beyond standard tests, necessitating the comprehensive reporting of all data.

In addition to rotational behavior, we observed fluctuations in body weight following DA depletion (Fig. 1D), a common phenomenon in rodents treated with 6-OHDA (Barata-Antunes et al. 2020; Masini et al. 2021). Despite receiving extensive post-operative care, all subjects experienced weight loss within 7-14 days post-6-OHDA injection. Importantly, we noted a significant, positive correlation between the observed weight changes and rotational behavior,

with subjects displaying higher degrees of weight loss also exhibiting more pronounced rotational
biases post-6-OHDA injection (Pearson correlation coefficient r=.81, p<.01, n=10).

6-OHDA lesion affects detection behavior

Twelve mice were trained on a tactile Go/No-Go detection task (Ollerenshaw et al. 2012;
Stüttgen, Rüter, and Schwarz 2006; Waiblinger et al. 2018, 2019, 2022), and their performance
was evaluated over time in the healthy and PD-like state. Figure 2A depicts the task setup and
logic, where mice responded to whisker deflections by either licking a waterspout (Go) or
refraining from licking (No-Go) in the absence of a stimulus. Reward delivery depended on correct
responses and "time-outs" were used to penalize/discourage impulsive licking. Further details
regarding the task are provided in the Methods.

Figure 2B outlines the experimental timeline, with procedures detailed in the Methods section.
Briefly, over eight weeks, animals underwent a series of experiments. This included "Detection
Training," where animals received single whisker stimulations or no stimulation. Learning
progress was assessed daily, and upon meeting criteria, animals proceeded to "Detection
Testing." During this phase, the psychometric curve was measured in repeated daily sessions for
a minimum of 6 days. Following this testing period, animals underwent unilateral 6-OHDA
injection and allowed 7-14 days of recovery. After the recovery period, animals underwent
another round of "Detection Testing" post-lesion, mirroring the pre-lesion phase, to assess the
extent of behavioral impairments following 6-OHDA lesioning.

Figure 2C illustrates averaged psychometric curves from a subset of mice (n=10) before (gray)
and after 6-OHDA injection (red) across multiple sessions. The marked shift in the psychometric
curve post-lesion signifies a decline in performance, with varying effects noted among individual
mice. This decline in average psychometric performance implies compromised tactile sensation
following DA depletion. While evident degradation in performance was observed, complete
abolishment was not consistently present. This variability could stem from the degree of DA
depletion, inter-subject variability, or a combination of both. The high variability observed
precludes significant differences in standard statistical testing. Further investigation is warranted
given the variability across several measures.

6-OHDA lesion causes various deficits across individuals

To explore the behavioral response variability in the 6-OHDA lesioned mice in more detail, we
examined multiple behavioral metrics for each animal (Figure 3), including sensory detection
thresholds, motor lick responses, as well as overall task engagement and motivation.

Figure 3A displays psychometric curves and response thresholds for nine representative mice
that performed the detection task in both the healthy (gray) and DA depleted (colored) states.
When analyzing the data individually, a pattern of varying task performance emerges allowing us
to order subjects according to their impairment severity. Based on the difference in response
threshold (at P=0.5) before and after 6-OHDA lesion, animals were classified into three severity
groups; "minor", "moderate" and "severe" deficit (Fig. 3A, from left to right). Animals with minor
deficits show minimal performance changes (thresh$_{post-pre}$<1, green compared to gray) with a
slight potential improvement. Animals with a moderate deficit show a clear decrease in
performance (thresh$_{post-pre}$>1 and <10, blue compared to gray), yet they are still performing the
task. Animals with a severe deficit display a substantial drop in performance (thresh$_{post-pre}$>10,
red compared to gray) and even stop performing the task.

To assess whether the decline in performance correlated with the inability of animals to respond
following DA depletion, indicating a motor aspect to the behavioral deficit, we analyzed indicator
lick response rates (Fig. 3B, top) and first lick latencies (Fig. 3B, bottom). Animals with minor
deficits have similar response rates and latencies before and after lesioning (green compared to
gray). Intriguingly, in the moderate case, although licking behavior diminishes after lesioning, it
persists without a noticeable increase in latency (blue compared to gray). This suggests that these
animals retain the ability to generate timed motor responses, indicating intact motor capabilities.
However, the observed compromise in sensory-motor capabilities underscores the likely
presence of a sensory deficit. In severe cases, overall task activity drastically decreases, with
subjects exhibiting minimal or no directed lick responses (red compared to gray), indicating
potential motor impairments in addition to any underlying sensory deficits. For this group of
animals, increased response latency is not a reliable metric due to the absence of lick responses
for latency estimation. Categorizing severe cases as purely motor-related may be an
oversimplification, as there could be involvement of both motor and sensory aspects.
Importantly, these animals demonstrate eating, drinking, and grooming behaviors outside the
setup, suggesting some degree in resilience in motor control.

Associated with task performance is the accumulation of rewards, representing the total
instances of an animal successfully receiving a reward (Fig. 3C, vertical axis). To quantify this, we
computed the average total number of rewards per session both before (gray dashed lines) and
after the lesion (colored dashed lines). This metric serves as an approximation of the animal's
motivation, given that rewards act as a primary motivator for task engagement. The number of
trials completed (Fig. 3C, horizontal axis) serves as a proxy for the overall engagement in the task.
It is important to note that animals were required to complete a minimum of 50 trials, and a
session was considered complete once an entire block of stimuli (A = [0, 2, 4, 8, 16]°) was missed.
Consistent with their stable performance, animals with minor deficits accumulate a comparable
reward count before and after the lesion (left column). In some cases, animals even gain a few
more rewards by working more trials. Animals with moderate deficits (middle column), however,
show a clear decrease in reward accumulation and reduced task engagement post lesioning. As
severity of impairment increases, the number of rewards gained and total number of trials
worked decreases drastically compared to the healthy state. Severely impaired animals (right
column) are held in the setup for the minimum number of trials with very limited activity.

Our examination of multiple behavioral metrics for individual subjects revealed a spectrum of
impairments, reflecting variability in task performance among lesioned animals. This diversity
manifested in changes to response probabilities, response rates, latency, and associated reward
accumulation. Notably, our analysis enabled the classification of severity and identified a
sensory-only component in the moderate cases, alongside motivational factors.

6-OHDA lesion causes persistent deficits over time

Unlike the gradual progression of PD symptoms observed clinically, 6-OHDA induces a swift and
complete lesion in the nigrostriatal pathway, commonly injected into the substantia nigra or
medial forebrain bundle (Agid et al. 1973; Przedborski and Jackson-Lewis 1995; Tieu 2011).
Studies on the motor symptom progression indicate that 6-OHDA-induced lesions can be
established with high stability over time (Antony, Diederich, and Balling 2011; Iancu et al. 2005;
Quiroga-Varela et al. 2017). However, the progression of non-motor behavioral deficits in this
model remains poorly understood. To assess potential changes over time, we systematically
analyzed the data from the detection experiments by evaluating behavioral performance across
sessions and days in both healthy and PD-like states (Figure 4). Animals were once again
categorized into three severity groups (minor, moderate, and severe deficit) as explained in the
previous section.

Figure 4A illustrates the detection performance over multiple sessions for three animals with
minor deficits, both before and after the 6-OHDA lesion. Performance is represented as the
probability of a correct lick with a salient stimulus (8-degree whisker deflection) across 6 pre-
lesion sessions and up to 12 post-lesion sessions for each animal. Before the lesion, animals
exhibit slight variability in performance. After the lesion, performance remains unchanged, with
consistent levels observed across sessions, suggesting that the perceptual capabilities linked to
the task likely remained intact in these mice.

Figure 4B illustrates the detection performance over time for four animals exhibiting moderate
deficits. While some individuals show an initial decline in performance shortly after the lesion,
noticeable variability emerges across sessions, occasionally within a single day. The session-to-
session variability underscores the complexity of the animals' responses, potentially elucidating
the coexistence of sensory and motivational deficits alongside preserved motor capabilities
within the detection task, as previously observed for this group of mice (Figure 3, blue panels).

Figure 4C illustrates a substantial decline in detection performance in three severely affected
animals shortly after the 6-OHDA injection, evident within the initial two sessions. Following the
lesion, variability in performance quickly diminishes in the severe group, reaching nearly zero
levels. This pronounced and persistent change in performance confirms the previous observation
that severely impaired animals exhibit very limited activity in response to the DA depletion.

Notably, recovery periods varied among the three groups of mice, resulting in differences in the
data presented regarding the lesion timeframe. Additionally, due to occasional repetition of
sessions within a day and pauses on weekends, sessions do not directly correspond to days. To
address this, performance was tracked across days in a separate analysis, considering these
variations (Figure. 4-1 and Table 1). The day-based analysis yields results consistent with the
session-based analysis, confirming that task performance does not improve or degrade over time
within the measured timeframe.

Utilizing multiple metrics for comprehensive lesion assessment

Our findings underscore that inducing unilateral DA depletion leads to rotational defects, weight
loss, and diverse behavioral impairments, as evidenced in a sensory-driven detection task. While
these metrics have been discussed separately thus far, we now compare them to gain a more
comprehensive understanding of the lesion effects. Figure 5 presents an overview of all metrics
assessed in this study before and after the 6-OHDA lesion, comparing histological findings,
rotation data, weight loss, and behavioral data from the detection task. These metrics are
compared for each of the three severity groups – minor (green), moderate (blue), and severe
deficit (red) – to elucidate the distinct effects observed within each group.

Figure 5A summarizes all metrics for animals with minor deficits. By construction, animals in this
group demonstrate consistent detection thresholds before and after the lesion, indicating stable
sensory behavioral performance. In some cases, performance may have even slightly improved.
Strikingly, these animals exhibit a bias in rotation behavior, predominantly favoring left turns (60-
84%), and experience a modest loss of body weight (10-12%) during the recovery period.
Histological analysis reveals a substantial reduction (48-81%) in DA innervation in the lesioned
hemisphere, as quantified via immunofluorescence. This discrepancy between stable detection
performance and DA depletion metrics strongly suggests that the lesion may have impacted
certain physiological functions unrelated to the detection task or that the animals were able to
compensate in some manner.

Figure 5B summarizes all metrics for animals with moderate deficits. By construction, animals
with moderate deficits demonstrate a clear increase in detection threshold (despite lack of
apparent motor deficit – see Fig. 3), consistent with a sensory deficit post-6-OHDA lesion. Animals
in this group exhibit varying left turn biases (55-98%), suggesting inconsistent motor impairment.
This variability is also reflected in weight loss during the recovery period (9-21%). Histological
analysis confirms a substantial reduction (47-86%) in DA innervation, indicating the efficacy of
the lesion. The observed decrease in detection performance alongside inconsistent motor
impairments further underscores the likelihood of isolated sensory or motivational deficits in this
group of mice, emphasizing the inherent variability in lesion effects.

Figure 5C summarizes all metrics for animals with severe deficits. By design, animals with severe
deficits exhibit a substantial increase in detection threshold following 6-OHDA injection. These
mice display a pronounced rotational bias (90-95%) with minimal variability and experience
substantial weight loss during the extended recovery period (23-29% within 21 days). Histological
analysis for this group reveals the most profound reduction (85-90%) in DA innervation among
the three levels of behavioral deficit, emphasizing the severity of the 6-OHDA lesion within this
group. This pronounced change in all metrics confirms the previous observation that severely
impaired animals exhibit very limited activity in response to the lesion.

In summary, our study demonstrates the diverse effects of unilateral DA depletion induced by 6-
OHDA injection. Mice with severe deficits in sensory-driven behavior consistently exhibited
pronounced impairments across all measures, indicating robust and consistent effects. However,
while standard measures like rotation bias and histological assessment are crucial for evaluating
DA depletion, their correlation with sensory task performance is sometimes limited. For instance,
mice with minor sensory deficits still showed substantial DA depletion alongside rotational bias
and weight loss. Interestingly, mice with moderate declines in sensory performance displayed
the greatest variability across metrics, suggesting complex interactions between these factors.
These findings emphasize the necessity of comprehensive assessment using multiple behavioral
parameters to fully capture the effects of DA depletion.

Discussion

In this study, we investigate the relationship between DA depletion and behavior within the 6-OHDA mouse model, conducting a comprehensive exploration of impairments. Our findings illustrate that inducing unilateral DA depletion leads to rotational defects, weight loss, and a wide range of diverse sensory- and motor deficits, particularly evident in a behavioral detection task. The variations in task performance observed among animals underscore the inherent complexity of the model. By employing multiple metrics for comprehensive lesion assessment and dissecting sensory and motor components, our study offers detailed insights into the contributing factors underlying the observed diversity of impairments.

While Parkinson's Disease diagnosis traditionally relies on motor symptoms, emerging evidence of sensory deficits suggests a potential window for early diagnostic methods (Juri, Rodriguez-Oroz, and Obeso 2010; Konczak et al. 2012; Prätorius, Kimmeskamp, and Milani 2003; Richardson and Sussman 2019). Despite not replicating PD's natural progression, the 6-OHDA mouse model provides a stable platform for characterizing deficits in PD-like states, induced by DA depletion in the striatum (Bagga, Dunnett, and Fricker 2015; Francardo et al. 2011; Masini et al. 2021). This understanding of the model's fidelity extends to investigating the question: How does DA depletion impact sensory processing? Our findings reveal a wide spectrum of behavioral deficits, highlighting the complexity of the relationship between DA depletion and sensory processing. Within this spectrum, we observe both increased and decreased sensitivity in sensory detection performance. Such impaired tactile acuity has been reported in animal models of PD (Romero-Sánchez et al. 2020) as well as in clinical settings (Kesayan et al. 2015). While our study primarily focuses on the behavioral outcomes of DA depletion, the underlying mechanisms by which this neurotransmitter deficit affects sensory function are complex and multifaceted. One possible avenue for exploration is through the anatomy of the basal ganglia and its connections with sensory processing regions in the cortex. DA plays a crucial modulatory role in these circuits, and its depletion may disrupt the transmission of sensory information or alter the processing of sensory signals, leading to changes in detection performance. Studies reporting altered striatal sensory responses (Ketzef et al. 2017; Reig and Silberberg 2014) underscore the potential impact of DA depletion on sensory processing. In contrast to studies involving electrophysiology on non-behaving or anesthetized animals, our study deliberately focuses on understanding behavioral outcomes in the natural, behaving state, allowing speculation on the precise impact of striatal changes on sensory functions. Further research utilizing electrophysiology in behaving animals is necessary to elucidate the specific neural pathways and mechanisms underlying these effects.

PD patients exhibit a wide range of non-motor impairments, including alterations in olfactory, auditory, tactile, nociceptive, thermal, and proprioceptive perception (Oppo et al. 2020; Jafari, Kolb, and Mohajerani 2020; Kesayan et al. 2015; José Luvizutto et al. 2020; Brim and Struhal 2021). In our investigation, we focus on tactile sensation mediated through whisker deflection in mice, using the conserved rodent "whisker to barrel" pathway as a reliable model for tactile-sensory stimulation and processing. How do findings in the vibrissa system relate to other sensory pathways? While our investigation primarily centers on this model, our behavioral paradigm and findings may have broader implications across sensory modalities. The principles of sensory

integration and DA-mediated modulation observed in our study could extend to other sensory
pathways with similar organization, including vision, hearing, taste, and smell. Given dopamine's
known influence on neural circuits involved in diverse sensory modalities, alterations in DA levels
could impact sensory perception and processing across various modalities. Therefore, our study
may provide valuable insights into the broader mechanisms underlying sensory deficits in PD and
other neurodegenerative disorders.

Our findings unveil a large spectrum of impairments following DA depletion, revealing high
variability among individuals. Where does this variability come from? It may arise from multiple
factors. Differences in the precise placement and depth of the injection within the medial
forebrain bundle can lead to variations in the extent of DA depletion, impacting different
subregions of the striatum (Masini et al. 2021). Despite controlling factors such as age, genetic
background, and environmental conditions, variables like overall health status, social rank among
cage mates, or overall activity level may influence vulnerability to neurotoxic insults. These
factors can impact the magnitude of DA depletion induced by 6-OHDA injections, further
contributing to the observed variability in behavioral outcomes. Furthermore, intrinsic variability
in behavior is evident even among healthy subjects. Overall, the variability following 6-OHDA
injections likely arises from a combination of factors, including lesion accuracy, neuronal
susceptibility, pharmacokinetic factors, and general variability in behavioral responses.

Our study challenges the presumed direct correlation between behavioral deficits and the extent
of DA depletion, which is typically evaluated using rotation bias (Przedborski and Jackson-Lewis
1995; Ungerstedt and Arbuthnott 1970). While such tests are essential for assessing DA depletion
in vivo, they have limitations in capturing the full behavioral spectrum. Excluding subjects with
limited rotation bias from analysis narrows the perspective within this animal model, potentially
overlooking important aspects in the relationship between DA depletion and behavior. This is
particularly notable in our moderate cases, which show variable rotation bias but clear deficits in
stimulus detection without altered response latencies, indicating isolated sensory impairments.
Additionally, these moderate cases often show strong session-to-session fluctuations and
achieve fewer rewards, indicating a possible motivational deficit. These findings underscore the
complexity of lesion-induced behavioral changes and warrant a reevaluation of traditional
interpretations of PD models, emphasizing the need to consider multiple behavioral metrics and
individual variability in lesion effects.

The histological assessment, although limited to a subset of mice, reveals substantial DA
depletion, regardless of the assessed sensory behavioral deficit. The small increments in DA
depletion between severity groups underscore the impact of even slight changes in DA levels on
behavior and physiology. While our model does not mirror PD's natural progression, it predicts
an intriguing pattern: Minor deficits indicate substantial DA loss but stable sensory detection,
suggesting compensatory mechanisms. Moderate deficits involve compromised sensory and
motivational aspects, eventually leading to severe deficits akin to classic motor symptoms. This
pattern provides implications for understanding the progression of PD. Further exploration with
genetic mouse models of PD progression (Dovonou et al. 2023) and rigorous histological or
pathophysiological assessments could validate these predictions. Such a comprehensive

approach would elucidate the underlying mechanisms contributing to the observed behavioral
changes across disease progression.

In summary, our study sheds light on sensory deficits in the 6-OHDA mouse model, providing
insights into early PD symptoms. The proposed behavioral task proves valuable for uncovering
these symptoms and other hidden variables. Additionally, our severity-based classification
system suggests the ability to capture different disease stages depending on lesion extent,
hinting at the potential for early PD detection. This underscores the versatility of the 6-OHDA
model in simulating various disease stages. To fully leverage the potential of the behavioral task
and reveal circuitry alterations, further research with disease progression models, longitudinal
measurements, and physiological assessments is warranted. This approach holds promise for
deepening our understanding of PD and advancing intervention and treatment development.

References

Aeed, Fadi, Nathan Cermak, Jackie Schiller, and Yitzhak Schiller. 2021. "Intrinsic Disruption of the M1 Cortical Network in a Mouse Model of Parkinson's Disease." *Movement Disorders: Official Journal of the Movement Disorder Society* 36(7): 1565–77. doi:10.1002/mds.28538.

Agid, Yves, France Javoy, Jacques Glowinski, Dominique Bouvet, and Constantino Sotelo. 1973. "Injection of 6-Hydroxydopamine into the Substantia Nigra of the Rat. II. Diffusion and Specificity." *Brain Research* 58(2): 291–301. doi:10.1016/0006-8993(73)90002-4.

Antony, Paul M. A., Nico J. Diederich, and Rudi Balling. 2011. "Parkinson's Disease Mouse Models in Translational Research." *Mammalian Genome* 22(7–8): 401–19. doi:10.1007/s00335-011-9330-x.

Bagga, V., S.B. Dunnett, and R.A. Fricker. 2015. "The 6-OHDA Mouse Model of Parkinson's Disease – Terminal Striatal Lesions Provide a Superior Measure of Neuronal Loss and Replacement than Median Forebrain Bundle Lesions." *Behavioural Brain Research* 288: 107–17. doi:10.1016/j.bbr.2015.03.058.

Barata-Antunes, Sandra, Fábio G. Teixeira, Bárbara Mendes-Pinheiro, Ana V. Domingues, Helena Vilaça-Faria, Ana Marote, Deolinda Silva, Rui A. Sousa, and António J. Salgado. 2020. "Impact of Aging on the 6-OHDA-Induced Rat Model of Parkinson's Disease." *International Journal of Molecular Sciences* 21(10): 3459. doi:10.3390/ijms21103459.

Branchi, Igor, Ivana D'Andrea, Monica Armida, Tommaso Cassano, Antonella Pèzzola, Rosa Luisa Potenza, Maria Grazia Morgese, Patrizia Popoli, and Enrico Alleva. 2008. "Nonmotor Symptoms in Parkinson's Disease: Investigating Early-phase Onset of Behavioral Dysfunction in the 6-hydroxydopamine-lesioned Rat Model." *Journal of Neuroscience Research* 86(9): 2050–61. doi:10.1002/jnr.21642.

Brim, Bianca, and Walter Struhal. 2021. "Thermoregulatory Dysfunctions in Idiopathic Parkinson's Disease." In *International Review of Movement Disorders*, Elsevier, 285–98. doi:10.1016/bs.irmvd.2021.08.009.

Brooks, Simon P., and Stephen B. Dunnett. 2009. "Tests to Assess Motor Phenotype in Mice: A User's Guide." *Nature Reviews Neuroscience* 10(7): 519–29. doi:10.1038/nrn2652.

Chagas, André M., Lucas Theis, Biswa Sengupta, Maik C. Stüttgen, Matthias Bethge, and Cornelius Schwarz. 2013. "Functional Analysis of Ultra High Information Rates Conveyed by Rat Vibrissal Primary Afferents." *Frontiers in Neural Circuits* 7. doi:10.3389/fncir.2013.00190.

Dovonou, Axelle, Cyril Bolduc, Victoria Soto Linan, Charles Gora, Modesto R. Peralta Iii, and
Martin Lévesque. 2023. "Animal Models of Parkinson's Disease: Bridging the Gap
between Disease Hallmarks and Research Questions." *Translational Neurodegeneration*
12(1): 36. doi:10.1186/s40035-023-00368-8.

Francardo, Veronica, Alessandra Recchia, Nataljia Popovic, Daniel Andersson, Hans Nissbrandt,
and M. Angela Cenci. 2011. "Impact of the Lesion Procedure on the Profiles of Motor
Impairment and Molecular Responsiveness to L-DOPA in the 6-Hydroxydopamine
Mouse Model of Parkinson's Disease." *Neurobiology of Disease* 42(3): 327–40.
doi:10.1016/j.nbd.2011.01.024.

Frankin, Keith B.J., and George Paxinos. 2008. *The Mouse Brain in Stereotaxic Coordinates*
(2008). 3rd ed.

Frund, I., N. V. Haenel, and F. A. Wichmann. 2011. "Inference for Psychometric Functions in the
Presence of Nonstationary Behavior." *Journal of Vision* 11(6): 16–16.
doi:10.1167/11.6.16.

Gage, Gregory J., Daryl R. Kipke, and William Shain. 2012. "Whole Animal Perfusion Fixation for
Rodents." *Journal of Visualized Experiments* (65): 3564. doi:10.3791/3564.

Iancu, Ruxandra, Paul Mohapel, Patrik Brundin, and Gesine Paul. 2005. "Behavioral
Characterization of a Unilateral 6-OHDA-Lesion Model of Parkinson's Disease in Mice."
Behavioural Brain Research 162(1): 1–10. doi:10.1016/j.bbr.2005.02.023.

Jafari, Zahra, Bryan E. Kolb, and Majid H. Mohajerani. 2020. "Auditory Dysfunction in
Parkinson's Disease." *Movement Disorders* 35(4): 537–50. doi:10.1002/mds.28000.

José Luvizutto, Gustavo, Thanielle Souza Silva Brito, Eduardo De Moura Neto, and Luciane
Aparecida Pascucci Sande De Souza. 2020. "Altered Visual and Proprioceptive Spatial
Perception in Individuals with Parkinson's Disease." *Perceptual and Motor Skills* 127(1):
98–112. doi:10.1177/0031512519880421.

Juri, Carlos, MariCruz Rodriguez-Oroz, and Jose A. Obeso. 2010. "The Pathophysiological Basis
of Sensory Disturbances in Parkinson's Disease." *Journal of the Neurological Sciences*
289(1–2): 60–65. doi:10.1016/j.jns.2009.08.018.

Kesayan, Tigran, Damon G. Lamb, Adam D. Falchook, John B. Williamson, Liliana Salazar, Irene
A. Malaty, Nikolaus R. McFarland, et al. 2015. "Abnormal Tactile Pressure Perception in
Parkinson's Disease." *Journal of Clinical and Experimental Neuropsychology* 37(8): 808–
15. doi:10.1080/13803395.2015.1060951.

Ketzef, Maya, Giada Spigolon, Yvonne Johansson, Alessandra Bonito-Oliva, Gilberto Fisone, and
Gilad Silberberg. 2017. "Dopamine Depletion Impairs Bilateral Sensory Processing in the

Striatum in a Pathway-Dependent Manner." *Neuron* 94(4): 855-865.e5.
doi:10.1016/j.neuron.2017.05.004.

Konczak, Jürgen, Alessandra Sciutti, Laura Avanzino, Valentina Squeri, Monica Gori, Lorenzo
Masia, Giovanni Abbruzzese, and Giulio Sandini. 2012. "Parkinson's Disease Accelerates
Age-Related Decline in Haptic Perception by Altering Somatosensory Integration." *Brain*
135(11): 3371–79. doi:10.1093/brain/aws265.

Lundblad, M., B. Picconi, H. Lindgren, and M.A. Cenci. 2004. "A Model of L-DOPA-Induced
Dyskinesia in 6-Hydroxydopamine Lesioned Mice: Relation to Motor and Cellular
Parameters of Nigrostriatal Function." *Neurobiology of Disease* 16(1): 110–23.
doi:10.1016/j.nbd.2004.01.007.

Masini, Débora, Carina Plewnia, Maëlle Bertho, Nicolas Scalbert, Vittorio Caggiano, and Gilberto
Fisone. 2021. "A Guide to the Generation of a 6-Hydroxydopamine Mouse Model of
Parkinson's Disease for the Study of Non-Motor Symptoms." *Biomedicines* 9(6): 598.
doi:10.3390/biomedicines9060598.

Ollerenshaw, Douglas R., Bilal A. Bari, Daniel C. Millard, Lauren E. Orr, Qi Wang, and Garrett B.
Stanley. 2012. "Detection of Tactile Inputs in the Rat Vibrissa Pathway." *Journal of*
Neurophysiology 108(2): 479–90. doi:10.1152/jn.00004.2012.

Oppo, Valentina, Marta Melis, Melania Melis, Iole Tomassini Barbarossa, and Giovanni Cossu.
2020. "'Smelling and Tasting' Parkinson's Disease: Using Senses to Improve the
Knowledge of the Disease." *Frontiers in Aging Neuroscience* 12: 43.
doi:10.3389/fnagi.2020.00043.

Pont-Sunyer, Claustre, Anna Hotter, Carles Gaig, Klaus Seppi, Yaroslau Compta, Regina
Katzenschlager, Natalia Mas, et al. 2015. "The Onset of Nonmotor Symptomsin
Parkinson's Disease (the ONSET PD Study)." *Movement Disorders* 30(2): 229–37.
doi:10.1002/mds.26077.

Prätorius, B, S Kimmeskamp, and T.L Milani. 2003. "The Sensitivity of the Sole of the Foot in
Patients with Morbus Parkinson." *Neuroscience Letters* 346(3): 173–76.
doi:10.1016/S0304-3940(03)00582-2.

Przedborski, S, and V Jackson-Lewis. 1995. "DOSE-DEPENDENT LESIONS OF THE DOPAMINERGIC
NIGROSTRIATAL PATHWAY INDUCED BY INTRASTRIATAL INJECTION OF 6-
HYDROXYDOPAMINE." *Neuroscience* 67(3): 631–47.

Quiroga-Varela, A., E. Aguilar, E. Iglesias, J.A. Obeso, and C. Marin. 2017. "Short- and Long-Term
Effects Induced by Repeated 6-OHDA Intraventricular Administration: A New Progressive
and Bilateral Rodent Model of Parkinson's Disease." *Neuroscience* 361: 144–56.
doi:10.1016/j.neuroscience.2017.08.017.

Reig, Ramon, and Gilad Silberberg. 2014. "Multisensory Integration in the Mouse Striatum."
Neuron 83(5): 1200–1212. doi:10.1016/j.neuron.2014.07.033.

Richardson, Kelly C., and Joan E. Sussman. 2019. "Intensity Resolution in Individuals With
Parkinson's Disease: Sensory and Auditory Memory Limitations." *Journal of Speech,*
Language, and Hearing Research 62(9): 3564–81. doi:10.1044/2019_JSLHR-H-18-0424.

Ritt, Jason T., Mark L. Andermann, and Christopher I. Moore. 2008. "Embodied Information
Processing: Vibrissa Mechanics and Texture Features Shape Micromotions in Actively
Sensing Rats." *Neuron* 57(4): 599–613. doi:10.1016/j.neuron.2007.12.024.

Romero-Sánchez, Héctor Alonso, Liliana Mendieta, Amaya Montserat Austrich-Olivares,
Gabriela Garza-Mouriño, Marcela Benitez-Diaz Mirón, Arrigo Coen, and Beatriz Godínez-
Chaparro. 2020. "Unilateral Lesion of the Nigroestriatal Pathway with 6-OHDA Induced
Allodynia and Hyperalgesia Reverted by Pramipexol in Rats." *European Journal of*
Pharmacology 869: 172814. doi:10.1016/j.ejphar.2019.172814.

Stüttgen, Maik C., Johannes Rüter, and Cornelius Schwarz. 2006. "Two Psychophysical Channels
of Whisker Deflection in Rats Align with Two Neuronal Classes of Primary Afferents." *The*
Journal of Neuroscience 26(30): 7933–41. doi:10.1523/JNEUROSCI.1864-06.2006.

Thiele, Sherri L., Ruth Warre, and Joanne E. Nash. 2012. "Development of a Unilaterally-
Lesioned 6-OHDA Mouse Model of Parkinson's Disease." *Journal of Visualized*
Experiments (60): 3234. doi:10.3791/3234.

Tieu, K. 2011. "A Guide to Neurotoxic Animal Models of Parkinson's Disease." *Cold Spring*
Harbor Perspectives in Medicine 1(1): a009316–a009316.
doi:10.1101/cshperspect.a009316.

Ungerstedt, Urban, and Gordon W. Arbuthnott. 1970. "Quantitative Recording of Rotational
Behavior in Rats after 6-Hydroxy-Dopamine Lesions of the Nigrostriatal Dopamine
System." *Brain Research* 24(3): 485–93. doi:10.1016/0006-8993(70)90187-3.

Waiblinger, Christian et al. 2022. "Emerging Experience-Dependent Dynamics in Primary
Somatosensory Cortex Reflect Behavioral Adaptation." *Nature Communications* 13(1):
534.

Waiblinger, Christian, Clarissa J. Whitmire, Audrey Sederberg, Garrett B. Stanley, and Cornelius
Schwarz. 2018. "Primary Tactile Thalamus Spiking Reflects Cognitive Signals." *The*
Journal of Neuroscience 38(21): 4870–85. doi:10.1523/JNEUROSCI.2403-17.2018.

Waiblinger, Christian et al. 2019. "Stimulus Context and Reward Contingency Induce Behavioral
Adaptation in a Rodent Tactile Detection Task." *The Journal of Neuroscience* 39(6):
1088–99.

Wichmann, Felix A., and N. Jeremy Hill. 2001. "The Psychometric Function: I. Fitting, Sampling,
and Goodness of Fit." *Perception & Psychophysics* 63(8): 1293–1313.
doi:10.3758/BF03194544.

Wolfe, Jason, Dan N Hill, Sohrab Pahlavan, Patrick J Drew, David Kleinfeld, and Daniel E
Feldman. 2008. "Texture Coding in the Rat Whisker System: Slip-Stick Versus Differential
Resonance" ed. Garrett B Stanley. *PLoS Biology* 6(8): e215.
doi:10.1371/journal.pbio.0060215.

Figure 1. Validation of 6-OHDA mouse model. A unilateral injection of 0.2ul of 6-OHDA (15 mg/ml solution 6-OHDA.HBr) was administered into the left median forebrain bundle via Hamilton syringe. Mice were allowed to recover for up to 2 weeks before testing continued. **A.** Coronal sections showing the striatal hemispheres of a control mouse without lesion stained for TH expression. **B.** Coronal sections of a mouse after 6-OHDA-lesion, stained for TH expression. Over 80% reduction in fluorescence was found between DA-depleted and control hemisphere. Scale bar: 1 mm. **C.** Percent left turns performed by each mouse in the rotameter test before and after 6-OHDA injection. Bars represent medians across mice with percentiles (n=10). ***P < 0.001, two-sided Wilcoxon rank-sum test. **D.** Body weight before and after 6-OHDA injection. Percent body weight is calculated by identifying the minimum weight within the recovery period. Numbers are normalized to each animals' weight on the day before the lesion (n=10). ***P < 0.001, two-sided Wilcoxon rank-sum test. Gray areas represent median range across sham operated mice (n=2).

A **Tactile detection task**

B **Experimental Timeline**

C **Psychometric performance**

Figure 2. Assessment of sensory capabilities in the 6-OHDA model.
A. Schematic of the behavior setup, stimulus delivery and behavioral detection task. A punctuate whisker deflection (10 ms) has to be detected by the mouse with an indicator lick to receive reward ("Go"). Licking has to be withheld in catch trials ("No-go"). Reward is only delivered upon correct licks. **B.** Timeline of experiments. Mice were first trained on the detection task. Detection testing was done at least 6 days prior to 6-OHDA injection and repeated after the recovery period. **C.** Left: Psychometric curves and response thresholds before (gray) and after 6-OHDA injection (red). "Stim amplitude" represents the degree of whisker deflection. Filled circles correspond to the average response probabilities from all animals tested (n=10). Solid curves are logistic fits to the data. Response thresholds are shown as vertical dotted lines with 95 % confidence limits. Right: Response thresholds separately shown for all mice before and after the lesion (gray and red symbols).

Figure 3. Assessment of sensory capabilities in different individuals. A. Psychometric curves and response thresholds for each animal performing the detection task before (gray) and after 6-OHDA injection (colored). Mice are ordered by the performance drop (from left to right). Solid curves are logistic fits to response probabilities with different stimulus amplitudes from an individual, averaged across sessions. Response thresholds at P=0.5 are shown as vertical lines. The difference in response thresholds is used to define the 3 categories: minor deficit (thresh$_{post-pre}$ <1, green, n=3), moderate deficit (thresh$_{post-pre}$ >1 and <10, blue, n=3) and severe deficit (thresh$_{post-pre}$ >10, red, n=3). **B.** Top: Response (1st lick) latencies of individual mice before and after 6-OHDA injection shown as histograms. Bottom: Median 1st lick latencies with percentiles. **C.** Number of rewards (correct trials) accumulated as a function of trials worked by each animal before and after 6-OHDA injection. Each line corresponds to one session. Dashed horizontal lines correspond to the total rewards per session on average. Figure conventions and order are the same as in A.

Figure 4. Persistence of sensory deficits in the 6-OHDA mouse model. The 3 categories, minor, moderate and severe, are based on the difference in thresholds from the psychometric curves. **A.** Detection performance (P correct lick with 8 degree whisker deflection) over time (sessions) for animals with minor deficits, before and after the 6-OHDA lesion. **B.** Performance for animals with moderate deficits. **C.** Performance for animals with severe deficits. Symbols represent individual mice. Thick lines represent medians across mice with percentiles (n=3-4).

A Minor deficit

B Moderate deficit

C Severe deficit

Figure 5. Comparison of multiple behavioral metrics and DA depletion in the 6-OHDA mouse model. A. Minor deficit group (n=3). Detection: Perceptual thresholds (at P=0.5) from the detection task for all mice before and after the 6-OHDA lesion. Rotation: Percent left turns performed by animals in the rotameter test before and after 6-OHDA lesion. Weight loss: Maximum weight loss during the recovery period for each animal. DA depletion: Left. The percentage of DA preservation is quantified from histological analysis for all mice of this group. Right. Coronal sections of an example mouse after 6-OHDA-lesion, stained for tyrosine hydroxylase (TH) expression. **B.** Same metrics for the moderate deficit group (n=4). **C.** Same metrics for the severe deficit group (n=3). Symbols represent individual mice. Solid filled circles represent medians across mice with percentiles. Gray areas represent median range across sham operated mice (n=2). Note, histological assessment was performed for a subset of mice (minor deficit n=3; moderate deficit n=3.; severe deficit n=2). Scale bars: 1 mm.