1

Genome-wide analysis of Smad and Schnurri transcription factors in *C. elegans*

demonstrates widespread interaction and a function in collagen secretion

Mehul Vora[3,4,*], Jonathan Dietz[3,*], Zachary Wing[1], Jun Liu[5], Christopher Rongo[3,†], Cathy

Savage-Dunn[1,2,†]

[1]Department of Biology, Queens College, CUNY, NY, USA

[2]PhD Program in Biology, the Graduate Center, CUNY, NY, USA

[3]Waksman Institute, Dept. of Genetics, Rutgers University, NJ, USA

[4]ModOmics Ltd, Southampton, UK

[5]Department of Molecular Biology and Genetics, Cornell University, NY, USA

*These two authors contributed equally

[†]Co-corresponding authors

ABSTRACT

Smads and their transcription factor partners mediate the transcriptional responses of target cells to secreted ligands of the Transforming Growth Factor-β (TGF-β) family, including those of the conserved bone morphogenetic protein (BMP) family, yet only a small number of direct target genes have been well characterized. In *C. elegans,* the BMP2/4 ortholog DBL-1 regulates multiple biological functions, including body size, via a canonical receptor-Smad signaling cascade. Here, we identify functional binding sites for SMA-3/Smad and its transcriptional partner SMA-9/Schnurri based on ChIP-seq peaks (identified by modEncode) and expression differences of nearby genes identified from RNA-seq analysis of corresponding mutants. We found that SMA-3 and SMA-9 have both overlapping and unique target genes. At a genome-wide scale, SMA-3/Smad acts as a transcriptional activator, whereas SMA-9/Schnurri direct targets include both activated and repressed genes. Mutations in *sma-9* partially suppress the small body size phenotype of *sma-3,* suggesting some level of antagonism between these factors and challenging the prevailing model for Schnurri function. A functional analysis of target genes revealed a novel role in body size for genes involved in one-carbon metabolism and in the endoplasmic reticulum (ER) secretory pathway, including the disulfide reductase *dpy-11.* Our findings indicate that Smads and SMA-9/Schnurri have previously unappreciated complex genetic and genomic regulatory interactions that in turn regulate the secretion of extracellular components like collagen into the cuticle to mediate body size regulation.

INTRODUCTION

Members of the TGF-β family of secreted ligands play numerous roles in development and disease. In humans, there are 33 ligand genes that can be broadly separated into two subfamilies: the TGF-β/Activin subfamily and the BMP subfamily [1]. Due to the conservation of these ligands and their signaling pathways across metazoans, genetic studies in invertebrate systems have been instrumental in identifying signaling mechanisms [2, 3]. Canonical signaling occurs when ligand dimers bind to transmembrane receptors generating a heterotetrameric complex consisting of two type I and two type II serine/threonine kinase receptors. Following ligand binding and complex assembly, the constitutively active type II receptor phosphorylates the type I receptor on the GS domain and thereby activates its kinase domain [4]. The activated type I receptor phosphorylates the C-terminus of intracellular receptor-regulated Smads (R-Smads), promoting their heterotrimeric complex formation with co-Smads. The heterotrimeric Smad complex accumulates in the nucleus and binds DNA directly to elicit changes in gene expression [5-15]. Co-Smads for all ligands and R-Smads for TGF-β/Activin ligands bind a 4 bp GTCT Smad Binding Element (SBE); furthermore, R-Smads for BMP ligands associate with GC-rich sequences (GC-SBE) [16-18]. The SBE is considered too degenerate and low affinity to account fully for binding specificity, so transcription factor partners likely contribute to target gene selection [19]. To date, only a few direct target genes of Smads have been extensively studied, including Drosophila *brinker* [20]; Xenopus *mixer* [21] and *Xvent2* [22]; and the mammalian *ATF3* and *Id* genes [23]. Genome-wide studies have the potential to expand these examples and elucidate general principles of target gene selection [24-26]. More of these studies are needed to understand how Smad transcriptional partners influence target gene selection and contribute to the execution of specific biological functions.

In the nematode *Caenorhabditis elegans,* a BMP signaling cascade initiated by the ligand DBL-1 plays a major role in body size regulation [27]. In nematodes, body size is constrained by

a collagen-rich cuticle, which is secreted by an epidermal layer (the hypodermis) and remodeled

over four successive molts during larval growth and then continuously during adulthood [28, 29].

DBL-1 signals through type I receptor SMA-6, type II receptor DAF-4, and Smads SMA-2,

SMA-3, and SMA-4 (founding members of the Smad family), which act together in the

hypodermis to promote body size growth during the earliest larval growth stages [30-32]. The

exact mechanism by which the DBL-1 pathway regulates body size is not fully understood, but

is known to involve the regulated synthesis of cuticular collagen, of which there are over 170

genes [33-35]. A complete understanding of how DBL-1 regulates body size will require the

identification of all direct transcriptional targets of the pathway during larval growth.

In addition to body size, the DBL-1 pathway also regulates male tail patterning, mesodermal

lineage specification, innate immunity, and lipid metabolism. A transcription factor partner for

this pathway, SMA-9, has been identified that plays a role in each of these biological functions

[36, 37]. SMA-9 is the homolog of Drosophila Schnurri, which was identified for its roles in

Dpp/BMP signaling [38-40]. Three vertebrate Schnurri homologs regulate immunity,

adipogenesis, and skeletogenesis, acting through both BMP-dependent and BMP-independent

mechanisms [41-44]. Schnurri proteins are very large transcription factors with multiple Zn-

finger domains. At the *brinker* locus in Drosophila and the *Xvent2* locus in Xenopus, binding of

an R-Smad and a Co-Smad with a precise 5 bp spacing between binding sites has been shown

to recruit Schnurri, which controls the direction of transcriptional regulation [22]. This model for

Smad-Schnurri interaction has not been tested at a genomic scale.

In this study, we use BETA software to combine RNA-seq and ChIP-seq datasets for

SMA-3/Smad and SMA-9/Schnurri to identify direct versus indirect target genes of these factors,

as well as to identify common versus unique targets [45, 46]. Analysis of *sma-3; sma-9* double

mutants further extends our understanding of how these factors interact to produce locus-

specific effects on target genes. We use GO term analysis and loss-of-function studies that

shed light on the downstream effectors for body size regulation, lipid metabolism, and innate

immunity. Finally, we use a ROL-6::wrmScarlet reporter for collagen synthesis and secretion to

show that SMA-3, SMA-9, and the transcriptional target gene DPY-11 regulate body size growth

by promoting the secretion and delivery of collagen into the cuticle.

METHODS

C. elegans strains

C. elegans strains were grown at 20°C using standard methods unless otherwise indicated. N2 is the wild-type strain; some strains were provided by the *Caenorhabditis Genetics Center* (CGC), which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440) or generated in previous work. Strong loss-of-function or null alleles were used. The following genotypes were used: *sma-3(wk30), sma-9(ok1628), qcIs6[sma-3p::gfp::sma-3, rol-6(d)], jjIs1253[sma-9p::sma-9C2::gfp+unc-119(+)] (*generated via bombardment)*,* [47]*, rrf-3(pk1426), fat-6(tm331), fat-7(wa36), nhr-114(gk849), acs-22(gk373989), lbp-8(gk5151), clec-60(tm2319), sysm-1(ok3236), ins-7(ok1573), sma-2(e502), rol-6(syb2235[rol-6::wrmScarlet])* [48]*, and pwSi82[hyp-7p::VIT2ss::oxGFP::KDEL, HygR]*. The double mutants *sma-3(wk30); sma-9(wk97), and rrf-3(pk1426); lon-2(e678)* were generated in this study.

RNA-seq

Developmentally synchronized animals were obtained by hypochlorite treatment of gravid adults to isolate embryos. Animals were grown on NGM plates at 20°C until the late L2 stage. Total RNA was isolated from animals using Trizol (Invitrogen) combined with Bead Beater lysis in three biological replicates for each genotype [45]. Libraries were generated using polyA selection in a paired-end fashion and sequenced on an Illumina HiSeq (2x150 bp configuration, single index, per lane) by Azenta (formerly Genewiz). Reads were mapped to the *C. elegans* genome (WS245) and gene counts generated with STAR 2.5.1a. Normalization and statistical analysis on gene counts were performed with EdgeR using generalized linear model functionality and tagwise dispersion estimates. Principal Component Analysis showed tight clustering within four biological replicates, with a clear separation between genotypes in which SMA-3 and SMA-9 active versus inactive. Likelihood ratio tests were conducted in a pairwise fashion between genotypes with a Benjamini and Hochberg correction. All RNA-seq raw

sequence files as well as normalized counts after EdgeR can be accessed at GEO (Accession

Number: GSE266398)

Chromatin Immunoprecipitation Sequencing

Chromatin immunoprecipitation sequencing (ChIP-seq) was performed by Michelle Kudron

(Valerie Reinke Model Organism ENCyclopedia Of DNA Elements and model organism

Encyclopedia of Regulatory Networks group) on the *sma-3(wk30);qcIs6[GFP::SMA-3]* and

LW1253*: jjIs1253[sma-9p::sma-9C2::gfp+unc-119(+)]* strains at the late L2 stage [49]. Data are

available at encodeproject.org. To calculate distances between SMA-3 and SMA-9 ChIP-seq

peaks, each peak was reduced to a centroid position (midpoint between the two border

coordinates along the chromosome). For each chromosome, a matrix of SMA-3 and SMA-9

peak centroids was created, allowing the measurement of distance (in bps) between every

SMA-3 and SMA-9 centroid along that chromosome. The shortest distance in the matrix was

chosen to define each SMA-3/SMA-9 nearest neighboring pair. The resulting inter-centroid

distances were analyzed from all 6 chromosomes. To mimic a random distribution of SMA-3 and

SMA-9 peaks, each peak on a given chromosome was reassigned a location on that

chromosome using randomized values (generated by the Microsoft Excel randomization

function) within the size range for that chromosome. A matrix of SMA-3 and SMA-9 peak

centroids was then analyzed from this randomized dataset as described above for the actual

dataset.

Identification of Direct Targets Using BETA and LOA

To identify SMA-3 direct targets, differentially expressed genes (DEGs) from the RNA-seq

comparison of wild type vs. *sma-3(wk30)* using an FDR≤0.05 were compared against the

genomic coordinates of SMA-3 peaks from the ChIP-seq analysis using BETA basic and the

WS245 annotation of the *C. elegans* genome [46]. The following parameters were used: 3 kb

from TSS, FDR cutoff of 0.05 and one-tail KS test cutoff of 0.05. The input files consisted of

.bed files of IDR thresholded peaks and differential expression Log_2FC and FDR values from the

RNA-seq. An identical approach was used to identify SMA-9 direct targets using DEGs from the

RNA-seq comparison of wild type vs. *sma-9(ok1628).*

To identify direct targets co-regulated by both SMA-3 and SMA-9, the two pairwise RNA-seq

comparisons (wild type vs. *sma-3* and wild type vs. *sma-9*) were analyzed, measuring DEGs for

the same genes in both comparisons. Taking a conditional approach, the information from the

first comparison (wild type vs. *sma-3*) was examined to see if it affected interpretation in the

second (wild type vs. *sma-9*). Using the approach of Luperchio et al. [50], the genes in the

second comparison were split into two groups, conditional on the results in the first comparison,

with one group comprising genes found to show differential expression in the first comparison,

and the second group comprising genes found not to show differential expression. To estimate

which genes were differentially regulated, an FDR of 0.01 was used to generate an overlapping

list between the two comparisons. BETA basic was then used to identify potential direct targets

of the SMA-3/SMA-9 combination using just the ChIP-seq peaks that overlapped between the

two transcription factors. The following parameters were used: 3 kb from TSS, FDR cutoff of

0.05 and one-tail KS test cutoff of 0.05. Analysis tools can be obtained at GitHub:

https://github.com/shahlab/hypoxia-multiomics as per [45]. The WormBase database was used

to obtain information about candidate target genes, including sequence, genetic map position,

expression pattern, and available mutant alleles [51].

Quantitative RT-PCR analysis

Worms were synchronized using overnight egg lay followed by 4-h synchronization. When

animals reached L4 stage, they were collected and washed, and then RNA was extracted using

previously published protocol [52] followed by Qiagen RNeasy miniprep kit (Catalogue. No.

74104). Invitrogen SuperScript IV VILO Master Mix (Catalogue. No.11756050) was used to

generate cDNA, and qRT-PCR analysis was done using Applied Biosystems *Power* SYBR

Green PCR Master Mix (Catalogue. No. 4367659). Delta delta Ct analysis was done using

Applied Biosystems and StepOne software. All qRT-PCR analysis was repeated on separate

biological replicates. The following primer pairs were used: 5'-ATGTGTGACGACGAGGTTGCC-

3' and 5'-GTCTCCGACGTACGAGTCCTT-3' to detect *act-1*, 5'-GTGGATTCTTCTTCGCTCAT-

3' and 5'-CACAAGATGACAAGTGGGAA-3' to detect *fat-6*, 5'-CATTCGATGTTTTTGAGGCG-3'

and 5'-GATCGAAGTAGGCACCATCT-3' to detect *nhr-114*, and 5'-

GGCAGGTCTAATCCACGACTTG-3' and 5'-CTAATGTCCGGGTTCCCATCG-3' to detect

C54E4.5. All graphs were made using GraphPad Prism software and statistical analysis was

performed using One-way ANOVA with Multiple Comparison Test, as calculated using the

GraphPad software.

RNAi Analysis of Body Size

RNAi knockdown of individual target genes was performed in the RNAi-sensitive *C. elegans*

mutants *rrf-3(pk1426)* and *rrf-3(pk1426); lon-2(e678),* which were fed HT115 bacteria containing

dsRNA expression plasmid L4440, with or without gene targeting sequences between flanking

T7 promoters. NGM growth plates were used containing ampicillin for L4440 RNAi plasmid

selection and IPTG (isopropyl β-D-1-thiogalactopyranoside) to induce dsRNA expression. Both

rrf-3(pk1426) and *rrf-3(pk1426); lon-2(e678)* were exposed to the RNAi food during the L4 stage

and allowed to lay eggs for 3 hours and then removed. Following hatching and development to

adulthood exposed to the RNAi food, 2 adult hermaphrodites were transferred to fresh RNAi

plates, allowed to lay eggs, and removed from the plate. Upon hatching and development to the

L4 stage, hermaphrodites were imaged using an AxioImager M1m (Carl Zeiss, Thornwood, NY)

with a 5X (NA 0.15) objective. The RNAi feeding constructs were obtained from the Open

Biosystems library (Invitrogen). To analyze the body length of the RNAi exposed animals, 3

independent measurements were made per worm using the segmented line tool on Fiji/ImageJ

[53]. 3-5 biological replicates were completed for each RNAi construct. The data were analyzed

using ANOVA with Dunnett's post hoc test correction for multiple comparisons.

Hypodermal Imaging of ROL-6::wrmScarlet

Animals expressing *rol-6::wrmScarlet* in different genetic backgrounds were imaged using a

Chroma/89 North CrestOptics X-Light V2 spinning disk, a Chroma/89North Laser Diode

Illuminator, and a Photometrics PRIME95BRM16C CMOS camera via MetaMorph software.

Day 1 adults (unless otherwise noted) were used to ensure molting was completed. A 63X oil

objective (NA 1.4) was used to detect fluorescence. In order to visualize the cuticle and

hypodermis layers of each animal, a z-series was completed using a 0.5 micron step size

across 6.5 microns. Each image was analyzed using Fiji/ImageJ for fluorescence quantification

in the hypodermis of the animals. Background was subtracted using a rolling ball filter. An

outline was drawn around each nematode and the mean fluorescence intensity was calculated

within the outline. At least 10 animals were analyzed and pooled across 3-4 biological

replicates. Using GraphPad Prism, the individual mean fluorescence intensity values were

normalized to the mean for control animals in each experiment and analyzed using ANOVA with

Dunnett's post hoc test correction for multiple comparisons. Images were then deconvolved

using DeconvolutionLab2 [54].

Animals expressing *rol-6::wrmScarlet* together with the ER marker *VIT2ss::oxGFP::KDEL* were

imaged using a Zeiss Elyra 7 Lattice SIM. A 60X water objective (NA 1.2) was used to detect

fluorescence, and a z-series was completed as described above.

RNAi treatment of *rol-6::wrmScarlet* animals was performed similarly to the RNAi treatment in

the body length analysis with these exceptions: nematodes exposed to *dpy-11* RNAi grew for

one generation until day 1 adulthood before imaging, nematodes exposed to C54H2.5 and

F41C3.4-containing RNAi plasmids were introduced to the animals at the L1 development stage

and allowed to develop until day 1 adulthood, and nematodes exposed to Y25C1A.5 and

Y113G7A.3-containing RNAi plasmids were introduced to animals at the L4 development stage

and grown for 24 hours before imaging. Tunicamycin treatment of *rol-6::wrmScarlet* animals

was completed by allowing animals to develop from eggs to L4 stage in the presence of 5

µg/mL in NGM plates. Experiments were conducted over 3-4 biological replicates. The data

were analyzed using an unpaired two-tailed t test ANOVA with Dunnett's post hoc test

correction for multiple comparisons, where appropriate.

RESULTS

Transcription factors SMA-3 and SMA-9 bind overlapping and distinct genomic sites

Genetic evidence identifies SMA-3/Smad as a core component of the DBL-1/BMP signaling

cascade [2]. For many identified functions, *sma-3* mutant phenotypes are indistinguishable from

those of *dbl-1* ligand mutants, including body size, male tail patterning, mesodermal lineage

specification, and fat storage [2, 36, 55]. In contrast, SMA-9 contributes to each of these

functions, but in each case has a different effect, either a more limited role, a weaker effect, or

an antagonistic phenotype [36, 37, 55]. We sought to determine the transcriptional targets of

SMA-3 and SMA-9 on a genome-wide scale by analyzing ChIP-seq data from genome binding

of GFP-tagged SMA-3 and SMA-9 expressed from transgenes [56]. These constructs are

functional, as evidenced by their ability to rescue the mutant phenotypes of respective loss-of-

function *sma-3* and *sma-9* mutants [36, 55]. Genome binding was analyzed at the second larval

(L2) stage, a developmental stage at which a Smad reporter is highly active, and both SMA-3

and SMA-9 are first observed to affect body size [57]. ChIP sequencing reads identify 4205

peaks for GFP::SMA-3 and 7065 peaks for SMA-9::GFP (Supplementary File 1).

We considered the possibility that SMA-3 and SMA-9 bind as a protein complex in some

contexts. In *Drosophila*, the SMA-9 homolog Schnurri interacts physically with the Smads Mad

and Medea [58, 59]. We therefore analyzed the distances between the centroids of SMA-3 and

SMA-9 ChIP-seq peaks. If SMA-3 and SMA-9 bind independently, then we would expect a

Gaussian distribution of inter-centroid distances, whereas if they bind in a complex, we should

see a non-Gaussian distribution with increased representation of distances less than or equal to

the average peak size. This analysis demonstrated an increased representation (approaching

45%) of inter-centroid distances of 100 bp or less (Figure 1a,b), smaller than the average peak

size for SMA-3 (400 bp) or SMA-9 (250 bp) (Figure 1c), consistent with the interpretation that

SMA-3 and SMA-9 frequently bind as subunits in a complex. The midpoint of the cumulative

probability distribution of inter-centroid distances was 788 bp; by contrast, a randomization of

the positions of SMA-3 and SMA-9 ChIP peaks expanded the midpoint of the cumulative

probability distribution of inter-centroid distances to 8,211 bp (Figure 1a,b). From this analysis, a

substantial subset (3101 peaks) of the SMA-3 (73.7%) and SMA-9 (43.9%) peaks overlap

(Figure 1d). We therefore considered these instances of overlapping peaks to be complexes,

whereas adjacent but non-overlapping peaks likely represent independent binding, leading us to

conclude that (1) SMA-3 typically binds to sites in complex with SMA-9, and that (2) over half of

all SMA-9 sites are independent from these SMA-3/SMA-9 complex sites.

Identification of direct transcriptional targets of SMA-3 and SMA-9

To determine how these binding sites correlate with changes in gene expression of neighboring

genes, we performed RNA-seq on L2 stage samples of *sma-3* and *sma-9* mutants compared

with wild-type controls. Principal component analysis (PCA) demonstrated that all three

biologically independent replicates of each genotype clustered together (Figure 2a) and that

each genotype is transcriptionally distinct from the others. Using a false discovery rate (FDR) ≤

0.05, we identified 1093 differentially expressed genes (DEGs) downregulated and 774

upregulated DEGs in *sma-3* mutants (Figure 2b, Supplementary File 2). In *sma-9* mutants, we

identified 412 downregulated DEGs and 371 upregulated DEGs (Figure 2d, Supplementary File

2).

RNA-seq identifies both direct and indirect transcriptional targets. To identify direct functional

targets of each of these transcription factors, we employed BETA software (Figure 2c,e), which

infers direct target genes by integrating ChIP-seq and RNA-seq data [46]. BETA analysis

identified 367 direct targets for SMA-3 and 332 direct targets for SMA-9 (Supplementary File 3).

Every identified direct target of SMA-3 was downregulated in the *sma-3* mutant (Figure 2b),

indicating that SMA-3/Smad functions primarily as a transcriptional activator. In contrast, 46% of

direct targets of SMA-9 were upregulated and 53% were downregulated in the *sma-9* mutant

(Figure 2d, Supplementary File 3). Thus, SMA-9 likely acts as either a transcriptional activator

or repressor depending on the genomic context. This conclusion is consistent with our previous

analyses of SMA-9 function *in vivo* and in a heterologous system [34].

Significant overlap in directly regulated DEGs of SMA-3 and SMA-9

Because analysis of ChIP-seq peaks suggests that SMA-3 and SMA-9 can bind DNA as a

complex, we sought to identify a core subset of DEGs co-regulated by these transcription

factors. Rather than relying on individual RNA-seq analyses, in which arbitrary cut-offs for

significance may lead to an underestimation of the overlap, we performed Luperchio Overlap

Analysis (LOA) on *sma-3* and *sma-9* RNA-seq datasets to identify 882 shared DEGs

(Supplementary File 4) [45, 50]. From ChIP-seq data, we identified 3101 peaks that are

overlapping between SMA-3 and SMA-9. We used LOA to identify DEGs shared between the

pairwise comparison of *sma-3* vs wild type, and between the pairwise comparison of *sma-9* vs

wild type, using evidence of potential DEGs in one comparison to inform the state of those

potential DEGs in the other comparison. Processing common occupancy sites with the common

DEGs through BETA software (Figure 3a), we identified 129 co-regulated direct target genes

with overlapping SMA-3 and SMA-9 binding peaks (Supplementary File 5). These results are

consistent with SMA-3 and SMA-9 acting broadly in a protein complex. Most (114) of these

direct targets are activated by both SMA-3 and SMA-9 (Figure 3c), but 15 of them have

reversed regulation in *sma-9* mutants compared with *sma-3* (Figure 3d), suggesting an

antagonistic function that we analyze further below. For shared activated targets, loss of SMA-3

caused a greater fold change than loss of SMA-9 (Supplementary File 2).

Our BETA/LOA analysis next allowed us to deduce SMA-3 and SMA-9 exclusive targets,

revealing 238 SMA-3-exclusive direct targets (Figure 3b) and 279 SMA-9-exclusive direct

targets (Figure 3e,f). Surprisingly, many of these target genes contained overlapping SMA-3

and SMA-9 binding peaks (Figure 3b,e,f), although loss of one of the two factors did not result in

changes in gene expression, perhaps suggesting that the presence of the other factor at these

targets was sufficient to regulate gene expression to physiological levels. Interpretation is further

complicated for target genes surrounded by a mixture of distinct and overlapping SMA-3 and

SMA-9 peaks. Our results suggest that (1) SMA-3 and SMA-9 can act independent of one

another, (2) a SMA-3/SMA-9 complex usually (although not always) acts as a transcriptional

activator of shared target genes, and that (3) SMA-9 can act as either a transcriptional activator

or repressor of its own SMA-3-independent targets.

Mechanisms of integration of SMA-3 and SMA-9 function

SMA-3 and SMA-9 both regulate body size. We used double mutant analysis to determine

whether they do so independently or together. For two gene products that act together in the

same pathway, we expect the double mutants to resemble one of the single mutants. If they

function independently, then we expect the double mutant to be more severe than the single

mutants (additive phenotypes). We constructed a *sma-3; sma-9* double mutant and measured

its body length at the L4 stage in comparison with control, *sma-3* mutants, and *sma-9* mutants.

Contrary to expectations, the double mutant was neither the same as nor more severe than the

single mutants; instead, it showed an intermediate phenotype (Figure 4a). This result suggests

that there may be some antagonistic action between these two transcription factors, consistent

with our observation that some direct targets are regulated in opposite directions by SMA-3 and

SMA-9.

We hypothesized that the interaction between SMA-3 and SMA-9 may be context dependent,

with different target genes showing independent, coordinated, or antagonistic interactions

between the transcription factors. We tested this hypothesis by performing qRT-PCR on select

target genes in wild type, *sma-3* mutants, *sma-9* mutants, and *sma-3; sma-9* double mutants at

the L2 stage. We considered three target genes: *fat-6*, which has overlapping SMA-3 and SMA-

9 peaks and is downregulated in both *sma-3* and *sma-9* mutants; *nhr-114*, which has distinct

non-overlapping SMA-3 and SMA-9 peaks and is downregulated in both *sma-3* and *sma-9*

mutants; and C54E4.5, which has overlapping SMA-3 and SMA-9 peaks yet show opposite

direction of regulation in *sma-3* versus *sma-9* mutants. For two of the tested target genes, *fat-6*

and *nhr-114,* there was no significant difference in expression levels between the single and

double mutants (Figure 4b), consistent with the transcription factors acting together, regardless

of overlapping binding. The third target gene, C54E4.5, was selected because it is a co-direct

target yet the RNA-seq data shows changes in its expression in opposite directions in *sma-3*

versus *sma-9* mutants, downregulated in *sma-3* yet upregulated in *sma-9* relative to wild type. In

the double mutant, C54E4.5 is upregulated and indistinguishable from the expression in *sma-9*

single mutants (Figure 4b). Thus, for this target gene, the *sma-9* loss-of-function phenotype is

epistatic to that of *sma-3.* This relationship is possible if SMA-9 is required for SMA-3 binding at

this site, or if SMA-9 is required for SMA-3 to engage with the transcriptional machinery.

Biological functions of SMA-3 and SMA-9 target genes

Gene Ontology analysis showed that SMA-3 and SMA-9 shared some annotation clusters,

including for fatty acid metabolism and one-carbon metabolism, but also showed that each

regulated its own annotation clusters, including collagen genes, ribosome biogenesis,

mitochondrial proteins, and ER chaperones for SMA-3, as well as innate immunity factors and

cytochrome P450s for SMA-9 (Figure 3b-f; Figure 3 – figure supplement 1). A Smad-

independent role for SMA-9 in immunity is consistent with the pronounced role of vertebrate

Schnurri homologs in immunity [42], which have not been reported to overlap with TGF-β-

regulated functions. We expect that clusters of genes involved in fatty acid metabolism and

innate immunity mediate the physiological functions of BMP signaling in fat storage and

pathogen resistance, respectively. Using a candidate gene approach, we previously identified

several cuticle collagen genes that mediate regulation of body size downstream of BMP

signaling. Here, we sought to validate and extend this analysis in an unbiased manner by

screening target genes for a function in body size regulation. For nine genes for which mutants

were available, we measured body length at the L4 stage. As expected, *sma-2/Smad* mutants

were smaller. As we previously showed, *fat-6* and *fat-7* mutants were not significantly different

from wild type in larval stages [60]. Of the remaining six genes for which mutants were available,

only *ins-7* showed significantly reduced body size with a statistical effect size (Glass' effect size:

the difference between the mean of the mutant and the mean for wild type, divided by the

standard deviation of wild type) greater than one (Figure 5a,b). To test the functions of genes for

which mutants were not available, we used RNAi depletion in the *rrf-3* RNAi hypersensitive

mutant. If target genes truly act downstream of the DBL-1 signaling pathway to regulate body

size, then we expect them to also act downstream of the negative regulator LON-2, which

antagonizes the DBL-1 pathway with respect to body size at the level of ligand-receptor

interactions. Thus, we also performed RNAi depletion in a *lon-2; rrf-3* double mutant, which

demonstrates exaggerated DBL-1 signaling and elongated body size; we reasoned that

candidate transcriptional effectors of the DBL-1 pathway might be more limiting and hence show

a suppression phenotype in this genetic background. As expected, RNAi of *sma-3* or of *sma-9*

reduced the body length in both genetic backgrounds (Figure 5b,c,d; Figure 5 – figure

supplement 1). RNAi of one of the target genes, *mttr-1,* prevented the development of *rrf-3;*

lon-2 animals to the L4 stage, so body length could not be quantified at this stage. Nearly two

thirds of the 36 genes tested caused a statistically significant reduction in body length (with a

statistical effect size greater than one) upon RNAi treatment in at least one of the two genetic

backgrounds. Half of the genes tested caused a significant reduction in body length in both

genetic backgrounds (Figure 5 – figure supplement 1), making them strong candidates for direct

transcriptional effectors of body size regulation. These genes have GO terms associated with

either one-carbon metabolism or chaperone/ER secretion, suggesting that the upregulation of

these activities is a key aspect of how DBL-1 signaling promotes growth.

DBL-1 signaling promotes body size through collagen secretion

We observed that *dpy-11* depletion via RNAi resulted in the most severe reduction in body

length among the tested target genes. The *dpy-11* gene encodes a protein-disulfide reductase

involved in cuticle development. As we previously established that BMP signaling regulates

cuticle collagen gene expression, we posited whether BMP signaling also impacts cuticle

collagen secretion. We tested this hypothesis by monitoring the expression and localization of a

cuticle collagen, choosing to monitor ROL-6, a cuticle collagen gene with a demonstrated role in

body size [35].

The hypodermis synthesizes and secretes collagen into the cuticle in a pattern of circular annuli

that surround the animal along its length (Figure 6a). These collagen-rich annuli and the newly

synthesized collagen inside the hypodermal cells underneath the cuticle can be distinguished

using confocal microscopy (Figure 6b). We generated a functional endogenously tagged allele

of *rol-6* that expresses a ROL-6::wrmScarlet fusion protein, taking care to preserve the

proteolytic processing sites such that wrmScarlet remains attached to the final protein product.

As previously shown [48], this fusion protein localizes to the cuticle (Figure 6c). In L4 animals,

total cuticular fluorescence was reduced in *sma-3* and *sma-9* mutants, with clear patches of

decreased ROL-6::wrmScarlet (Figure 6d,e). Interestingly, the hypodermal subcellular

distribution of ROL-6::wrmScarlet was altered in *sma-3* and *sma-9* mutants, with

ROL-6::wrmScarlet protein accumulating intracellularly in these mutants compared to wild type,

often underneath the clear patches observed in the cuticular layer (Figure 6f). This phenomenon

is consistent with changes in collagen secretion in impaired BMP signaling. We used structured

illumination super-resolution microscopy to compare an endoplasmic reticulum (ER) marker,

KDEL::oxGFP [61], with subcellular ROL-6::wrmScarlet, and we found that ROL-6 becomes

trapped in and accumulates at the hypodermal ER in *sma-3* mutants (Figure 7a-l). We also

noted that ER structures in *sma-3* mutants were thinner and more skeletonized relative to wild

type (Figure 7j,l), consistent with a defect in secretion.

To further assess BMP signaling targets impact on collagen release, we depleted *dpy-11* via

RNAi in nematodes expressing ROL-6::wrmScarlet. Consistent with a role for DPY-11 in the

secretion of collagens, including ROL-6, we observed a severe disruption in both intracellular

and cuticular deposition of ROL-6::wrmScarlet (Figure 6g; Figure 6 – figure supplement 1). To

validate that the perturbed ROL-6::wrmScarlet distribution and localization exhibited in *sma-3*

and *sma-9* mutants and *dpy-11* RNAi was caused by disturbances in ER-specific processes, we

targeted the ER chemically and genetically. We observed similar ROL-6::wrmScarlet subcellular

distribution in tunicamycin treated ROL-6::wrmScarlet expressing animals as seen in *sma-3* and

sma-9 mutants (Figure 7m; Figure 7 – figure supplement 1). Tunicamycin inhibits the glycan

biosynthesis pathway, disrupts ER-mediated secretion, and induces ER stress. Moreover,

RNAi-mediated depletion of genes involved in various steps of ER-derived vesicle production

and transport, including *C54H2.5* (SURF4 ortholog/ER cargo release), *F41C3.4* (GOLT1A/b

ortholog/ER to Golgi apparatus transport), *Y25C1A.5* (COPB-1/COPI coat complex subunit),

and *Y113G7A.3* (COPII coat complex subunit) increased ROL-6::wrmScarlet hypodermal

intracellular accumulation and left empty patches in the cuticle, similar to BMP signaling mutants

(Figure 6g; Figure 6 – figure supplement 1). To test whether impaired secretion through the ER

alone could compromise body size, we examined nematodes treated with tunicamycin and

found that they showed a strong decrease in body size (Figure 7n). Taken together, our results

suggest that signaling by DBL-1/BMP signaling promotes body size growth by promoting ER-

specific processes involved in collagen maturation, transport, and secretion into the cuticular

ECM.

DISCUSSION

TGF-β signaling pathways regulate sets of target genes to execute biological functions in a

context-dependent manner. Canonical signaling is mediated by the Smad transcription factor

complex, but Smad binding sites are too degenerate and low affinity to account for the specific

context-dependent effects, so transcription factor partners must also be involved. Some of the

characterized partners are cell type-specific transcription factors. In contrast, Schnurri proteins

are transcriptional partners that co-regulate target genes across multiple cell types. Here we

used genome-wide RNA-seq and ChIP-seq and a novel software analysis pipeline to untangle

the roles of Smad and Schnurri transcription factors in the developing *C. elegans* larva. We

chose the second larval (L2) stage because Smad activity is elevated at this stage as

determined by the RAD-SMAD activity reporter [57], and because this stage is the earliest point

at which one can observe a clear difference for one of the best studied Smad mutant

phenotypes: body size growth. Our analysis provides a complete picture of the direct targets of

the DBL-1 signaling pathway during larval growth, and it identifies a novel set of effectors that

mediate changes in body size in response to DBL-1 signaling, including factors involved in

collagen secretion into the cuticle.

By using an analysis pipeline that combines BETA, which integrates ChIP-seq and RNA-seq

data to identify targets, with LOA, which integrates two separate RNA-seq pairwise comparisons

to identify shared DEGs, we demonstrated that SMA-3/Smad acts primarily as a transcriptional

activator, whereas SMA-9/Schnurri can function as either an activator or a repressor depending

on the locus. These dual functions for SMA-9/Schnurri are consistent with previous studies that

demonstrated that different domains of SMA-9 can act as activators or repressors in a

heterologous system [34]. Furthermore, SMA-9 DNA-binding domain fusions with known

transcriptional activation or repression domains could each rescue a subset of mutant defects in

sma-9 mutants [34].

Our analysis revealed that SMA-3/Smad and SMA-9/Schnurri have both co-regulated and

independently regulated target genes. Interestingly, the GO terms for both shared and

independent target genes partially overlap, suggesting broad similarity in biological functions. In

vertebrates, Schnurri homologs are shown to be direct DNA binding proteins with diverse

biological functions that include TGF-β-responsive and TGF-β-independent roles. In TGF-β-

independent roles they bind NFκB-like sequences [62], and can interact with other transcription

factors including TRAF2 and c-Jun [63, 64]. It will be interesting to determine whether the SMA-

9/Schnurri-specific target genes are responsive to TGF-β signals and/or to other signaling

ligands. Within the subset of shared target genes, our analysis indicated that a significant

number have overlapping binding peaks, consistent with a model in which SMA-3 and SMA-9

bind as a complex. Complex formation is consistent with the model for Smad-Schnurri

interaction in Drosophila and Xenopus. We have now generated new insights into how a Smad-

Schnurri complex may regulate target gene expression in a locus-specific manner. Analysis of

sma-3; sma-9 double mutants suggests that these factors may have antagonistic effects in

addition to cooperative effects. Antagonism would be consistent with the observed interaction

between SMA-9 and the BMP pathway in mesodermal lineage specification [36, 65]. For co-

activated target genes, genetic analysis is consistent with the proteins acting together rather

than additively, consistent with complex formation.

To identify a mechanism for antagonism between SMA-3 and SMA-9, we analyzed the co-

regulated target gene, C54E4.5, which is downregulated in *sma-3* and upregulated in *sma-9*

mutants. In the *sma-3; sma-9* double mutant, C54E4.5 is upregulated as it is in *sma-9* single

mutants. Thus, SMA-3 activation of C54E4.5 is dependent on the presence of SMA-9. This

dependence could occur if SMA-9 is needed to recruit SMA-3 to the DNA. This possibility would

be a novel mode of interaction, because for the previously analyzed *brinker* and *Xvent2* target

genes, Schnurri was shown to be recruited by the Smad complex rather than vice versa. A

second possibility is that SMA-3 can bind in the absence of SMA-9 but cannot engage with the

transcriptional machinery; that is, both proteins are required to form a transcriptional activation

complex. With either possibility, SMA-9 represses C54E4.5 expression in the absence of

SMA-3.

Finally, GO term analysis readily identified target genes involved in lipid metabolism and

pathogen response, but target genes required for body size regulation remain more difficult to

predict. To circumvent this barrier, we performed body size measurements on mutants. We

also performed body size measurements on RNAi knockdowns for identified target genes in an

RNAi sensitive strain, examining their effect in both a wild-type background and a *lon-2*

background in which DBL-1 signaling is exaggerated, resulting in an elongated body size.

These analyses confirmed previous work focusing on the role of the cuticle in mediating body

size regulation by DBL-1/BMP [35]. Although RNAi knockdowns of *clec-1, fah-1,* C52D10.3,

dre-1, hsp-12.3, wrt-1 reduced body size in the *rrf-3* mutant background, they failed to reduce

the body size in *lon-2*; *rrf-3* mutants, suggesting that they regulate body size upstream or

independently of the DBL-1 pathway. By contrast, RNAi knockdowns of *haf-9*, *his-32*, *zip-10*,

emb-8, *F25B5.6*, *nath-10*, and *hsp-3* reduced body size in *lon-2; rrf-3* double mutants but not in

rrf-3 single mutants, suggesting that they might be factors whose effects are only detectable in

the context of an overactive pathway; these warrant future study. Seventeen genes showed a

body reduction with a statistically significant effect size equal to or greater than one in both

genetic backgrounds, making them of particular interest. These genes had GO terms

associated with either one-carbon metabolism or chaperone/ER secretion, suggesting that the

upregulation of these activities is a key aspect of how DBL-1 signaling promotes growth.

How could one-carbon metabolism play a role in body size growth? This complex set of

interlinked metabolic cycles is critical for methionine and folate homeostasis. It also provides the

methyl groups needed to synthesize nucleotides, amino acids, the antioxidant glutathione,

creatine, and phospholipids like phosphatidylcholine, a fundamental component of membranes

[66]. One-carbon metabolism also provides the methyl groups needed to make epigenetic

marks on DNA and chromatin. The specific role of this metabolic pathway in body size growth

will be an important topic of future study.

An enrichment of ER secretion and chaperone factors in the list of direct targets involved in

body size growth was unexpected, but reasonable given that one of the key functions of the

hypodermis is to secrete cuticular collagen. The role of collagens in body size and morphology

is well documented [29, 67], and the secreted ADAMTS metalloprotease ADT-2 modifies cuticle

collagen organization and regulates body size [68]. To test whether DBL-1 regulates ER

secretion, we turned to endogenously tagged cuticle collagen ROL-6::wrmScarlet to analyze

subcellular localization. Using this reporter for collagen synthesis and secretion, we

demonstrated that the DBL-1 pathway influences the secretion of this cuticle collagen. In

particular, ROL-6::wrmScarlet accumulates in a perinuclear ER compartment in DBL-1 pathway

mutants. Interestingly, treatment with tunicamycin, which impairs ER secretion, was sufficient to

reduce body size, which is consistent with a model in which BMP signaling promotes collagen

secretion to foster growth. We remain cautious in our interpretation of this result, as blocking ER

secretion with tunicamycin could affect the secretion of the BMP receptors or other proteins that

function together with the receptors, which could also lead to a body size defect.

The collagenous cuticle is a major target of DBL-1/BMP signaling in body size regulation. For

example, others have also demonstrated that DBL-1 signaling regulates cuticle collagen LON-3

post-transcriptionally [69]. In addition to direct transcriptional regulation of cuticle components,

here we highlight the importance of regulating other target genes such as *dpy-11* that are

needed for collagen post-transcriptional processing. TGF-β family members are well known

regulators of collagen deposition and extracellular matrix composition, suggesting that this class

of transcriptional targets is conserved over evolutionary time [70, 71]. Thus, it is likely that the

multi-level interactions identified in *C. elegans* are also relevant to the functions of these factors

in vertebrates.

  FIGURE LEGENDS

Figure 1. Transcription factors SMA-3 and SMA-9 bind both overlapping and distinct

genomic sites

(a) Cumulative probability distribution graph of the distances between the centroids (base pair

position located centrally within each peak) of nearest neighbor SMA-3 and SMA-9 ChIP-seq

peaks. The black line represents actual inter-peak distances, whereas the green line represents

a hypothetical randomized dataset. The horizontal dotted line indicates the point in the curve at

which half of the peak pairs fall. (b) Same cumulative probability distribution as in (a), but

focused on distances less than 500 bps. The centroids of nearly half of all SMA-3/SMA-9

neighboring pairs fall within 500 bps of each other. (c) Histogram of the interpeak distances

(actual data in black, randomized data in green), as well as the ChIP-seq peak widths (SMA-3 in

red, SMA-9 in blue). Most peaks are larger in size than most interpeak centroid distances,

indicating substantial peak overlap. (d) Size proportional Venn diagram showing the number of

SMA-3 and SMA-9 peaks that either overlap with one another or remain independent. Although

most SMA-3 peaks overlap with SMA-9 peaks, more than half of SMA-9 peaks are located

independent from SMA-3 peaks.

Figure 2. Identification of direct transcriptional targets of SMA-3 and SMA-9

(a) Principal Component Analysis (PCA) over two dimensions (PC1 and PC2) for RNA-seq

datasets for wild type (in black), *sma-3(wk30)* (in red), and *sma-9(ok1628)* (in blue). The

percent of variance for each component is indicated. The three biological replicates for each

genotype are well clustered. (b) Volcano plot of RNA-seq FDR values versus log2 fold change

(FC) expression for individual genes (squares) in *sma-3* mutants relative to wild type. The direct

targets identified by BETA are indicated with red squares; the negative log2 FC values

demonstrate that SMA-3 promotes the expression of these genes. Non-target genes

nevertheless showing differential expression are indicated with green squares. (c) Strategy for

integrating SMA-3 ChIP-seq and mutant RNA-seq data to identify directly regulated targets. (d)

Volcano plot of RNA-seq FDR values versus log2 fold change expression for individual genes

(squares) in *sma-9* mutants relative to wild type. The direct targets identified by BETA are

indicated with blue squares; the combination of positive and negative log2FC values

demonstrates that SMA-9 promotes the expression of some of these genes and inhibits the

expression of others. Non-target genes nevertheless showing differential expression are

indicated with green squares. (e) Strategy for integrating SMA-9 ChIP-seq and mutant RNA-seq

data to identify directly regulated targets.

Figure 3. Significant overlap in directly regulated DEGs of SMA-3 and SMA-9

(a) Strategy for integrating SMA-3 and SMA-9 ChIP-seq and mutant RNA-seq data to identify

common versus unique directly regulated targets. (b-f) Cartoon representations of the different

types of direct target genes, their neighboring SMA-3 and/or SMA-9 binding sites, and the effect

of those sites on that gene's expression. The red circle labeled "3" represents SMA-3 binding

sites, whereas the blue circle labeled "9" represents SMA-9 binding sites. Arrows represent that

the wild type transcription factor promotes the expression of the neighboring DEG (gray),

whereas T-bars indicate that it inhibits the expression of the DEG. Types of regulation include

(b) SMA-3 alone promoting DEG expression, (c) SMA-3 and SMA-9 combined promoting

expression, (d) SMA-3 and SMA-9 showing antagonistic regulation of expression, (e) SMA-9

alone promoting DEG expression, and (f) SMA-9 alone inhibiting DEG expression. Example

DEGs and tables of annotation clusters for gene ontology terms for those DEGs (via DAVID,

with accompanying statistical EASE score) are shown under the cartoon demonstrating each

type of regulation.

Figure 3 – figure supplement 1. DAVID annotation clusters of gene ontology terms for

identified SMA-3 and SMA-9 targets.

Tables of annotation clusters for gene ontology terms for direct target DEGs (via DAVID, with

accompanying statistical EASE score) are shown for (a) SMA-3 and (b) SMA-9.

Figure 4. Genetic interactions between SMA-3 and SMA-9

(a) Mean body length of L4 animals (measured head to tail in microns). Dots indicate the size of

individual animals. ****$P<0.0001$, **$P<0.01$ One way ANOVA with Tukey's multiple comparison

test. (b) Mean mRNA levels for the indicated target gene (X-axis) for the indicated genotypes

(*sma-3*, *sma-9*, or the double mutant combination) relative to the level in wild type. Expression

values are in log2 fold change (FC). Individual genotype mRNA levels for each gene were first

normalized to actin mRNA levels in that genotype. *$P<0.05$ Two way ANOVA with Tukey's

multiple comparison test.

Figure 5. Multiple SMA-3 and SMA-9 target genes regulate body size

(a) Mean body length of L4 animals (normalized to wild type) for the indicated mutants. Dots

indicate the size of individual animals. Asterisks above each column indicate one way ANOVA

with Dunnett's multiple comparison test against wild type (****$P<0.0001$, ***$P<0.001$, **$P<0.01$,

*$P<0.05$). Asterisks over pairwise comparison bars indicate one way ANOVA with Sídák's

multiple comparison test. The dotted line indicates the size value falling two standard deviations

below the mean of wild type. (b) Glass' effect size (the difference between the mean of the

mutant and the mean for wild type, divided by the standard deviation of wild type) for the

indicated mutants. The dotted line indicates an effect size of one. (c,d) Mean body size and

Glass' effect size for RNAi knockdowns of the indicated gene in the *rrf-3* RNAi sensitive strain.

"EV" indicates the empty RNAi vector as a negative control. Asterisks above each column

indicate one way ANOVA with Dunnett's multiple comparison test against EV (****$P<0.0001$,

***P<0.001, **P<0.01, *P<0.05). For all graphs, red, blue, and purple columns indicate values

for *sma-3* (mutant or RNAi knockdown), *sma-9* (mutant or RNAi knockdown), and their double

mutant combination, respectively. Green bars indicate genotypes for which the mutant or RNAi

knockdown resulted in a statistically significant reduction (P<0.05) with a Glass' effect size of

one or greater.

Figure 5 – figure supplement 1. Target genes required for exaggerated growth in *lon-2*

mutants

(a) Mean body length of L4 animals (normalized to EV) for the indicated RNAi knockdowns in

the *rrf-3; lon-2* double mutant. "EV" indicates the empty RNAi vector as a negative control. Dots

indicate the size of individual animals. Asterisks above each column indicate one way ANOVA

with Dunnett's multiple comparison test against EV (****P<0.0001, ***P<0.001, **P<0.01,

*P<0.05). The dotted line indicates the size value falling two standard deviations below the

mean of EV (b) Glass' effect size (the difference between the mean of the RNAi knockdown and

the mean for EV, divided by the standard deviation for EV) for the indicated gene undergoing

knockdown. The dotted line indicates an effect size of one. Green bars indicate genotypes for

which the RNAi knockdown resulted in a statistically significant reduction (P<0.05) with a Glass'

effect size of one or greater. (c) The average of the Glass' effect size between the *rrf-3* wild-type

background and the *rrf-3 lon-2* mutant background (individual dots for each are shown). For (c),

yellow bars indicate genotypes for which the RNAi knockdown resulted in a statistically

significant reduction (P<0.05) with a Glass' effect size of one or greater in both the wild-type and

the *lon-2* mutant background. For all graphs, red and blue columns indicate values for *sma-3*

and *sma-9* RNAi knockdown, respectively.

Figure 6. DBL-1 signaling promotes body size through collagen secretion

(a) Cartoon illustrating a cross section through the nematode body (dorsal oriented up). Rings of cuticular collagen annuli (magenta), secreted by the underlying hypodermal cell layer (tan) surround the body. Lateral alae containing collagen, secreted by the underlying seam cells (gray), run orthogonal along the length of the body. (b) Cartoon illustrating a cross section through a portion of the nematode body (lateral oriented up) underneath a microscope coverglass. Horizontal green bars indicate the cuticular versus hypodermal plates captured by confocal microscopy in panels c-e. (c) ROL-6::wrmScarlet fluorescence detected in annuli and alae in the cuticular layer, as well as in nuclear envelopes in the underlying hypodermal layer in wild-type animals. The horizontal red line indicates the specific xz cross section shown below the cuticular and hypodermal plane panels. The bar indicates 5 microns. (d) ROL-6::wrmScarlet in *sma-3* mutants, visualized as per wild type. Patches of cuticular surface show diminished levels of ROL-6:wr:mScarlet, whereas the protein is detected in the hypodermal layer just under these patches (yellow arrowheads), suggesting a failure to deliver collagen to the surface cuticle (easily visualized in the xz panel). (e) Mutants for *sma-9* show the same phenotype as *sma-3*. (f) Quantification of ROL-6::wrmScarlet fluorescence in the hypodermal layer of indicated mutants. Dots indicate the fluorescence of individual animals. Asterisks over pairwise comparison bars indicate one way ANOVA with Sídák's multiple comparison test (***$P<0.001$, **$P<0.01$). (g) A similar analysis for RNAi knockdowns of the SMA-3 target gene *dpy-11*, as well as four known ER secretion factors as comparative controls. Asterisks above each column indicate one way ANOVA with Dunnett's multiple comparison test against wild type (***$P<0.001$, **$P<0.01$, *$P<0.05$).

Figure 6 – figure supplement 1. DPY-11 promotes body size through collagen secretion

(a) ROL-6::wrmScarlet fluorescence detected in annuli and alae in the cuticular layer, as well as in nuclear envelopes in the underlying hypodermal layer in wild-type animals exposed to an

empty vector for RNAi knockdown. (b-f) ROL-6::wrmScarlet in animals exposed to RNAi

knockdown for the indicated gene. (b) In animals knocked down for *dpy-11*, little

ROL-6::wrmScarlet makes it to the cuticle, instead accumulating intracellularly in the

hypodermis. (c-f) In animals knocked down for known ER secretory factors, patches of cuticular

surface show diminished levels of ROL-6::wrmScarlet, whereas the protein is detected in the

hypodermal layer just under these patches (yellow arrowheads), similar to what is observed in

sma-3 and *sma-9* mutants, and consistent with a failure to deliver collagen to the surface

cuticle. The bar indicates 5 microns.

Figure 7. ROL-6::wrmScarlet accumulates in the ER of *sma-3* mutants

ROL-6::wrmScarlet (magenta) and ssGFP::KDEL (an ER marker, shown here in yellow) shown

separately (a-d, g-j) or merged (e-f, k-l) in either (a-f) wild type or (g-l) a *sma-3* mutant.

(a,c,e,g,i,k) shows the interface between the cuticle and the hypodermis, as visualized by SIM

super resolution microscopy. (b,d,f,h,j,l) shows the hypodermal layer at a focal plane centered

around the nuclear envelope. In wild type, most ROL-6::wrmScarlet is delivered into the cuticle.

In sma-3 mutants, lower levels of ROL-6::wrmScarlet are present in the cuticle and rapidly

bleached under the SIM laser even under low power, whereas abundant ROL-6::wrmScarlet

colocalized with the ER ssGFP::KDEL marker (yellow). We noted that ER reticulation in *sma-3*

was thinner and skeletonized compared to wild type, perhaps suggesting reduced secretory

throughput. (m) Quantification of ROL-6::wrmScarlet fluorescence in the hypodermal layer of

tunicamycin treated versus untreated nematodes. (n) Mean body length of L4 animals

(normalized to untreated) for tunicamycin treated versus untreated nematodes. (m,n) Dots

indicate the values for individual animals. Asterisks over pairwise comparison bars indicate a

student t test (****$P<0.0001$).

Figure 7 – figure supplement 1. Tunicamycin treatment mimics the effect of *sma-3* **and**

sma-9 **mutations on collagen secretion**

(a) ROL-6::wrmScarlet fluorescence detected in annuli and alae in the cuticular layer, as well as

in nuclear envelopes in the underlying hypodermal layer in untreated wild-type animals. (b)

ROL-6::wrmScarlet in animals exposed to tunicamycin. Patches of cuticular surface show

diminished levels of ROL-6::wrmScarlet, whereas the protein is detected in the hypodermal

layer just under these patches (yellow arrowheads), similar to what is observed in *sma-3* and

sma-9 mutants, and consistent with a failure to deliver collagen to the surface cuticle. In

addition, collagen in the lateral alae is disorganized, suggesting secretion is impaired in the

seam cells. The bar indicates 5 microns.

SUPPLEMENTAL FILES

Supplemental File 1. SMA-3 and SMA-9 ChIP-seq Sites.

This file contains the chromosomal location of 4205 ChIP-seq peaks for SMA-3 and 7065 ChIP-seq peaks for SMA-9 in separate tabs labeled "SMA-3" and "SMA-9," respectively. SMA-3 sites that overlap with a SMA-9 site are listed on the "Overlapping Sites_S3" tab. SMA-9 sites that overlap with a SMA-3 site are listed on the "Overlapping Sites_S9" tab. Non-overlapping SMA-3 and SMA-9 sites are listed on the "Non-overlapping_S3" and "Non-overlapping_S9" tabs, respectively. For all tabs, column A indicates chromosome location, column B indicates the start of the peak sequence, and column C indicates the end of the peak sequence. Column labels are in row 1.

Supplemental File 2. Differential Gene Expression From *sma-3* and *sma-9* Mutants.

This file lists the differential gene expression from RNA-seq of *sma-3* vs wild type (the tab labeled "SMA3 vs N2"), as well as *sma-9* versus wild type (the tab labeled "SMA9 vs N2"). For each gene, WormBase GeneID, public gene name, log fold change, p-value, and FDR are listed. Column labels are in row 1.

Supplemental File 3. SMA-3 and SMA-9 Direct Targets.

This file lists the direct target genes identified by BETA analysis of RNA-seq and ChIP-seq data. Direct targets of SMA-3 are in the tab labeled "SMA3 Direct Targets." Direct targets of SMA-9 are in the tab labeled "SMA9 Direct Targets." For each gene, chromosomal location, transcriptional start site, transcriptional end site, public gene name, rank product from BETA analysis, RNA-seq log fold change from corresponding mutant vs wild type, and WormBase GeneID are listed. Column labels are in row 1.

Supplemental File 4. Differential Gene Expression Shared Between *sma-3* and *sma-9*

Mutants Analyzed Using LOA.

This file lists the differentially expressed genes identified by LOA analysis as being common to

both the RNA-seq of *sma-3* vs wild type as well as the RNA-seq of *sma-9* versus wild type. For

each gene, WormBase GeneID and public gene name are listed, followed by the log fold

change, p-value, and FDR from the *sma-3* vs wild type RNA-seq, followed by the log fold

change, p-value, and FDR from the *sma-9* vs wild type RNA-seq. Column labels are in row 1.

Supplemental File 5. Classes of Direct Targets for SMA-3 and SMA-9.

This file lists the direct target genes identified by combined LOA/BETA analysis of RNA-seq and

ChIP-seq data, as described in Figure 3. Direct targets of SMA-3 alone are in the tab labeled

"Figure 3b." Direct targets of SMA-3 and SMA-9 in which both factors promote the target's

expression are in the tab labeled "Figure 3c." Direct targets of SMA-3 and SMA-9 in which the

two factors have opposite effects on the target's expression are in the tab labeled "Figure 3d."

Direct targets of SMA-9 alone in which the factor either promotes or inhibits the target's

expression are in the tabs labeled "Figure 3e" and "Figure 3f," respectively. For each gene,

WormBase GeneID and public gene name are listed, followed by the log fold change and FDR

from the *sma-3* vs wild type RNA-seq, followed by the log fold change and FDR from the *sma-9*

vs wild type RNA-seq. Column labels are in row 1.

ACKNOWLEDGEMENTS

We thank Michelle Kudron with the Model Organism ENCyclopedia of DNA Elements and model

organism Encyclopedia of Regulatory Networks projects for performing chromatin

immunoprecipitation sequencing. Some strains were provided by the *Caenorhabditis* Genetics

Center, which is funded by the National Institutes of Health (NIH) Office of Research

Infrastructure Programs (P40 OD-101440). We thank Barth Grant for the *pwSi82* strain*,* and

Nanci Kane for her assistance with the Zeiss Elyra 7 Lattice SIM within the Waksman Institute

Shared Imaging Facility (Rutgers, The State University of New Jersey). We thank Derek Gordon

for guidance and assistance with the analysis of the ChIP-seq distance matrix. This work was

supported by NIH grants R15GM112147 to CSD, R21AG075315 to CSD and CR,

R35GM130351 to JL, and R01GM101972 to CR.

REFERENCES

1.   Massague J. TGF-beta signal transduction. Annu Rev Biochem. 1998;67:753-91. doi: 10.1146/annurev.biochem.67.1.753. PubMed PMID: 9759503.

2.   Savage C, Das P, Finelli AL, Townsend SR, Sun CY, Baird SE, et al. Caenorhabditis elegans genes sma-2, sma-3, and sma-4 define a conserved family of transforming growth factor beta pathway components. Proc Natl Acad Sci U S A. 1996;93(2):790-4. Epub 1996/01/23. doi: 10.1073/pnas.93.2.790. PubMed PMID: 8570636; PubMed Central PMCID: PMCPMC40134.

3.   Sekelsky JJ, Newfeld SJ, Raftery LA, Chartoff EH, Gelbart WM. Genetic characterization and cloning of mothers against dpp, a gene required for decapentaplegic function in Drosophila melanogaster. Genetics. 1995;139(3):1347-58. Epub 1995/03/01. PubMed PMID: 7768443; PubMed Central PMCID: PMCPMC1206461.

4.   Wrana JL, Attisano L, Wieser R, Ventura F, Massague J. Mechanism of activation of the TGF-beta receptor. Nature. 1994;370(6488):341-7. doi: 10.1038/370341a0. PubMed PMID: 8047140.

5.   Liu F, Ventura F, Doody J, Massague J. Human type II receptor for bone morphogenic proteins (BMPs): extension of the two-kinase receptor model to the BMPs. Mol Cell Biol. 1995;15(7):3479 - 86.

6.   Wrana J, Attisano L, Carcamo J, Zentella A, Doody J, Laiho M, et al. TGF-β signals through a heteromeric protein kinase receptor complex. Cell. 1992;71(6):1003 - 14. PubMed PMID: doi:10.1016/0092-8674(92)90395-S.

7.   Estevez M, Attisano L, Wrana JL, Albert PS, Massague J, Riddle DL. The daf-4 gene encodes a bone morphogenetic protein receptor controlling C. elegans dauer larva development. Nature. 1993;365(6447):644-9. Epub 1993/10/14. doi: 10.1038/365644a0. PubMed PMID: 8413626.

8.   Wu JW, Hu M, Chai J, Seoane J, Huse M, Li C, et al. Crystal structure of a phosphorylated Smad2. Recognition of phosphoserine by the MH2 domain and insights on Smad function in TGF-beta signaling. Mol Cell. 2001;8(6):1277-89. Epub 2002/01/10. PubMed PMID: 11779503.

9.   Chacko BM, Qin BY, Tiwari A, Shi G, Lam S, Hayward LJ, et al. Structural basis of heteromeric smad protein assembly in TGF-beta signaling. Molecular cell. 2004;15(5):813-23. doi: 10.1016/j.molcel.2004.07.016. PubMed PMID: 15350224.

10.   Souchelnytskyi S, Tamaki K, Engstrom U, Wernstedt C, ten Dijke P, Heldin CH. Phosphorylation of Ser465 and Ser467 in the C terminus of Smad2 mediates interaction with Smad4 and is required for TGF-β signaling. J Biol Chem. 1997;272(44):28107-15. Epub 1997/11/05. PubMed PMID: 9346966.

11.   Qin BY, Chacko BM, Lam SS, de Caestecker MP, Correia JJ, Lin K. Structural basis of Smad1 activation by receptor kinase phosphorylation. Mol Cell. 2001;8(6):1303-12. Epub 2002/01/10. PubMed PMID: 11779505.

12.   Zhang Y, Feng X, We R, Derynck R. Receptor-associated Mad homologues synergize as effectors of the TGF-beta response. Nature. 1996;383(6596):168 - 72. PubMed PMID: doi:10.1038/383168a0.

13.   Abdollah S, Macias-Silva M, Tsukazaki T, Hayashi H, Attisano L, Wrana JL. TGF-β RI phosphorylation of Smad2 on Ser465 and Ser467 is required for Smad2-Smad4 complex formation and signaling. J Biol Chem. 1997;272(44):27678-85. Epub 1997/11/05. PubMed PMID: 9346908.

14.   Lagna G, Hata A, Hemmati-Brivanlou A, Massague J. Partnership between DPC4 and SMAD proteins in TGF-beta signalling pathways. Nature. 1996;383(6603):832 - 6. PubMed PMID: doi:10.1038/383832a0.

15.   Nicolas FJ, De Bosscher K, Schmierer B, Hill CS. Analysis of Smad nucleocytoplasmic
shuttling in living cells. Journal of Cell Science. 2004;117(18):4113-25. doi: 10.1242/jcs.01289.
16.   Kim J, Johnson K, Chen HJ, Carroll S, Laughon A. Drosophila Mad binds to DNA and
directly mediates activation of vestigial by Decapentaplegic. Nature. 1997;388(6639):304-8. doi:
10.1038/40906. PubMed PMID: 9230443.
17.   Gao S, Steffen J, Laughon A. Dpp-responsive silencers are bound by a trimeric Mad-
Medea complex. The Journal of biological chemistry. 2005;280(43):36158-64. Epub 2005/08/20.
doi: 10.1074/jbc.M506882200. PubMed PMID: 16109720.
18.   Rushlow C, Colosimo PF, Lin M-c, Xu M, Kirov N. Transcriptional regulation of the
Drosophila gene zen by competing Smad and Brinker inputs. Genes & Development.
2001;15(3):340-51. doi: 10.1101/gad.861401.
19.   Hill CS. Transcriptional Control by the SMADs. Cold Spring Harb Perspect Biol.
2016;8(10). Epub 20161003. doi: 10.1101/cshperspect.a022079. PubMed PMID: 27449814;
PubMed Central PMCID: PMCPMC5046698.
20.   Pyrowolakis G, Hartmann B, Muller B, Basler K, Affolter M. A simple molecular complex
mediates widespread BMP-induced repression during Drosophila development. Dev Cell.
2004;7(2):229-40. doi: 10.1016/j.devcel.2004.07.008. PubMed PMID: 15296719.
21.   Germain S, Howell M, Esslemont GM, Hill CS. Homeodomain and winged-helix
transcription factors recruit activated Smads to distinct promoter elements via a common Smad
interaction motif. Genes Dev. 2000;14(4):435-51. PubMed PMID: 10691736; PubMed Central
PMCID: PMCPMC316385.
22.   Yao LC, Blitz IL, Peiffer DA, Phin S, Wang Y, Ogata S, et al. Schnurri transcription
factors from Drosophila and vertebrates can mediate Bmp signaling through a phylogenetically
conserved mechanism. Development. 2006;133(20):4025-34. doi: 10.1242/dev.02561. PubMed
PMID: 17008448.
23.   Kang Y, Chen CR, Massague J. A self-enabling TGFbeta response coupled to stress
signaling: Smad engages stress response factor ATF3 for Id1 repression in epithelial cells. Mol
Cell. 2003;11(4):915-26. doi: 10.1016/s1097-2765(03)00109-6. PubMed PMID: 12718878.
24.   Deignan L, Pinheiro MT, Sutcliffe C, Saunders A, Wilcockson SG, Zeef LA, et al.
Regulation of the BMP Signaling-Responsive Transcriptional Network in the Drosophila Embryo.
PLoS Genet. 2016;12(7):e1006164. Epub 20160705. doi: 10.1371/journal.pgen.1006164.
PubMed PMID: 27379389; PubMed Central PMCID: PMCPMC4933369.
25.   Chiu WT, Charney Le R, Blitz IL, Fish MB, Li Y, Biesinger J, et al. Genome-wide view of
TGFbeta/Foxh1 regulation of the early mesendoderm program. Development.
2014;141(23):4537-47. Epub 20141030. doi: 10.1242/dev.107227. PubMed PMID: 25359723;
PubMed Central PMCID: PMCPMC4302925.
26.   Morikawa M, Koinuma D, Tsutsumi S, Vasilaki E, Kanki Y, Heldin CH, et al. ChIP-seq
reveals cell type-specific binding patterns of BMP-specific Smads and a novel binding motif.
Nucleic Acids Res. 2011;39(20):8712-27. Epub 20110715. doi: 10.1093/nar/gkr572. PubMed
PMID: 21764776; PubMed Central PMCID: PMCPMC3203580.
27.   Suzuki Y, Yandell MD, Roy PJ, Krishna S, Savage-Dunn C, Ross RM, et al. A BMP
homolog acts as a dose-dependent regulator of body size and male tail patterning in
Caenorhabditis elegans. Development. 1999;126(2):241-50.
28.   Lazetic V, Fay DS. Molting in C. elegans. Worm. 2017;6(1):e1330246. Epub 2017/07/14.
doi: 10.1080/21624054.2017.1330246. PubMed PMID: 28702275; PubMed Central PMCID:
PMCPMC5501215.
29.   Page AP, Johnstone IL. The cuticle. WormBook. 2007:1-15. Epub 2007/12/01. doi:
10.1895/wormbook.1.138.1. PubMed PMID: 18050497; PubMed Central PMCID:
PMCPMC4781593.

30.   Gumienny TL, Savage-Dunn C. TGF-beta signaling in C. elegans. WormBook. 2013:1-
34. Epub 2013/08/03. doi: 10.1895/wormbook.1.22.2. PubMed PMID: 23908056; PubMed
Central PMCID: PMCPMC5081272.
31.   Yoshida S, Morita K, Mochii M, Ueno N. Hypodermal expression of Caenorhabditis
elegans TGF-beta type I receptor SMA-6 is essential for the growth and maintenance of body
length. Dev Biol. 2001;240(1):32-45. Epub 2002/01/11. doi: 10.1006/dbio.2001.0443. PubMed
PMID: 11784045.
32.   Wang J, Tokarz R, Savage-Dunn C. The expression of TGFbeta signal transducers in
the hypodermis regulates body size in C. elegans. Development. 2002;129(21):4989-98. Epub
2002/10/25. PubMed PMID: 12397107.
33.   Roberts AF, Gumienny TL, Gleason RJ, Wang H, Padgett RW. Regulation of genes
affecting body size and innate immunity by the DBL-1/BMP-like pathway in Caenorhabditis
elegans. BMC Dev Biol. 2010;10:61. Epub 2010/06/10. doi: 10.1186/1471-213X-10-61. PubMed
PMID: 20529267; PubMed Central PMCID: PMCPMC2894779.
34.   Liang J, Yu L, Yin J, Savage-Dunn C. Transcriptional repressor and activator activities of
SMA-9 contribute differentially to BMP-related signaling outputs. Dev Biol. 2007;305(2):714-25.
Epub 2007/04/03. doi: 10.1016/j.ydbio.2007.02.038. PubMed PMID: 17397820.
35.   Madaan U, Yzeiraj E, Meade M, Clark JF, Rushlow CA, Savage-Dunn C. BMP Signaling
Determines Body Size via Transcriptional Regulation of Collagen Genes in Caenorhabditis
elegans. Genetics. 2018;210(4):1355-67. Epub 2018/10/03. doi: 10.1534/genetics.118.301631.
PubMed PMID: 30274988; PubMed Central PMCID: PMCPMC6283163.
36.   Foehr ML, Lindy AS, Fairbank RC, Amin NM, Xu M, Yanowitz J, et al. An antagonistic
role for the C. elegans Schnurri homolog SMA-9 in modulating TGFbeta signaling during
mesodermal patterning. Development. 2006;133(15):2887-96. Epub 2006/06/23. doi:
10.1242/dev.02476. PubMed PMID: 16790477.
37.   Liang J, Lints R, Foehr ML, Tokarz R, Yu L, Emmons SW, et al. The Caenorhabditis
elegans schnurri homolog sma-9 mediates stage- and cell type-specific responses to DBL-1
BMP-related signaling. Development. 2003;130(26):6453-64. Epub 2003/11/25. doi:
10.1242/dev.00863. PubMed PMID: 14627718.
38.   Staehling-Hampton K, Laughon AS, Hoffmann FM. A Drosophila protein related to the
human zinc finger transcription factor PRDII/MBPI/HIV-EP1 is required for dpp signaling.
Development. 1995;121(10):3393-403. doi: 10.1242/dev.121.10.3393. PubMed PMID: 7588072.
39.   Grieder NC, Nellen D, Burke R, Basler K, Affolter M. Schnurri is required for Drosophila
Dpp signaling and encodes a zinc finger protein similar to the mammalian transcription factor
PRDII-BF1. Cell. 1995;81(5):791-800. doi: 10.1016/0092-8674(95)90540-5. PubMed PMID:
7774018.
40.   Arora K, Dai H, Kazuko SG, Jamal J, O'Connor MB, Letsou A, et al. The Drosophila
schnurri gene acts in the Dpp/TGF beta signaling pathway and encodes a transcription factor
homologous to the human MBP family. Cell. 1995;81(5):781-90. doi: 10.1016/0092-
8674(95)90539-1. PubMed PMID: 7774017.
41.   Jin W, Takagi T, Kanesashi SN, Kurahashi T, Nomura T, Harada J, et al. Schnurri-2
controls BMP-dependent adipogenesis via interaction with Smad proteins. Dev Cell.
2006;10(4):461-71. Epub 2006/04/04. doi: 10.1016/j.devcel.2006.02.016. PubMed PMID:
16580992.
42.   Jones DC, Glimcher LH. Regulation of bone formation and immune cell development by
Schnurri proteins. Adv Exp Med Biol. 2010;658:117-22. doi: 10.1007/978-1-4419-1050-9_13.
PubMed PMID: 19950022.
43.   Jones DC, Wein MN, Oukka M, Hofstaetter JG, Glimcher MJ, Glimcher LH. Regulation
of adult bone mass by the zinc finger adapter protein Schnurri-3. Science.
2006;312(5777):1223-7. doi: 10.1126/science.1126313. PubMed PMID: 16728642.

44.   Steinfeld H, Cho MT, Retterer K, Person R, Schaefer GB, Danylchuk N, et al. Mutations
in HIVEP2 are associated with developmental delay, intellectual disability, and dysmorphic
features. Neurogenetics. 2016;17(3):159-64. Epub 20160322. doi: 10.1007/s10048-016-0479-z.
PubMed PMID: 27003583; PubMed Central PMCID: PMCPMC4907844.
45.   Vora M, Pyonteck SM, Popovitchenko T, Matlack TL, Prashar A, Kane NS, et al. The
hypoxia response pathway promotes PEP carboxykinase and gluconeogenesis in C. elegans.
Nat Commun. 2022;13(1):6168. Epub 20221018. doi: 10.1038/s41467-022-33849-x. PubMed
PMID: 36257965; PubMed Central PMCID: PMCPMC9579151.
46.   Wang S, Sun H, Ma J, Zang C, Wang C, Wang J, et al. Target analysis by integration of
transcriptome and ChIP-seq data with BETA. Nat Protoc. 2013;8(12):2502-15. Epub 20131121.
doi: 10.1038/nprot.2013.150. PubMed PMID: 24263090; PubMed Central PMCID:
PMCPMC4135175.
47.   Praitis V, Casey E, Collar D, Austin J. Creation of low-copy integrated transgenic lines in
Caenorhabditis elegans. Genetics. 2001;157(3):1217-26. doi: 10.1093/genetics/157.3.1217.
PubMed PMID: 11238406; PubMed Central PMCID: PMCPMC1461581.
48.   Aggad D, Brouilly N, Omi S, Essmann CL, Dehapiot B, Savage-Dunn C, et al.
Meisosomes, folded membrane microdomains between the apical extracellular matrix and
epidermis. Elife. 2023;12. Epub 20230313. doi: 10.7554/eLife.75906. PubMed PMID: 36913486;
PubMed Central PMCID: PMCPMC10010689.
49.   Gerstein MB, Lu ZJ, Van Nostrand EL, Cheng C, Arshinoff BI, Liu T, et al. Integrative
analysis of the Caenorhabditis elegans genome by the modENCODE project. Science.
2010;330(6012):1775-87. Epub 2010/12/24. doi: 10.1126/science.1196914. PubMed PMID:
21177976; PubMed Central PMCID: PMCPMC3142569.
50.   Luperchio TR, Boukas L, Zhang L, Pilarowski G, Jiang J, Kalinousky A, et al. Leveraging
the Mendelian disorders of the epigenetic machinery to systematically map functional epigenetic
variation. Elife. 2021;10. Epub 20210831. doi: 10.7554/eLife.65884. PubMed PMID: 34463256;
PubMed Central PMCID: PMCPMC8443249.
51.   Davis P, Zarowiecki M, Arnaboldi V, Becerra A, Cain S, Chan J, et al. WormBase in
2022-data, processes, and tools for analyzing Caenorhabditis elegans. Genetics. 2022;220(4).
doi: 10.1093/genetics/iyac003. PubMed PMID: 35134929; PubMed Central PMCID:
PMCPMC8982018.
52.   Yin J, Madaan U, Park A, Aftab N, Savage-Dunn C. Multiple cis elements and GATA
factors regulate a cuticle collagen gene in Caenorhabditis elegans. Genesis. 2015;53(3-4):278-
84. Epub 2015/02/26. doi: 10.1002/dvg.22847. PubMed PMID: 25711168; PubMed Central
PMCID: PMCPMC4412796.
53.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub
20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID:
PMCPMC3855844.
54.   Sage D, Donati L, Soulez F, Fortun D, Schmit G, Seitz A, et al. DeconvolutionLab2: An
open-source software for deconvolution microscopy. Methods. 2017;115:28-41. Epub
20170103. doi: 10.1016/j.ymeth.2016.12.015. PubMed PMID: 28057586.
55.   Clark JF, Meade M, Ranepura G, Hall DH, Savage-Dunn C. Caenorhabditis elegans
DBL-1/BMP Regulates Lipid Accumulation via Interaction with Insulin Signaling. G3 (Bethesda).
2018;8(1):343-51. Epub 2017/11/23. doi: 10.1534/g3.117.300416. PubMed PMID: 29162682;
PubMed Central PMCID: PMCPMC5765361.
56.   Kudron MM, Victorsen A, Gevirtzman L, Hillier LW, Fisher WW, Vafeados D, et al. The
ModERN Resource: Genome-Wide Binding Profiles for Hundreds of Drosophila and
Caenorhabditis elegans Transcription Factors. Genetics. 2018;208(3):937-49. Epub 2017/12/30.
doi: 10.1534/genetics.117.300657. PubMed PMID: 29284660; PubMed Central PMCID:
PMCPMC5844342.

57.   Tian C, Sen D, Shi H, Foehr ML, Plavskin Y, Vatamaniuk OK, et al. The RGM protein DRAG-1 positively regulates a BMP-like signaling pathway in Caenorhabditis elegans. Development. 2010;137(14):2375-84. Epub 2010/06/11. doi: 10.1242/dev.051615. PubMed PMID: 20534671; PubMed Central PMCID: PMCPMC2889605.

58.   Dai H, Hogan C, Gopalakrishnan B, Torres-Vazquez J, Nguyen M, Park S, et al. The zinc finger protein schnurri acts as a Smad partner in mediating the transcriptional response to decapentaplegic. Dev Biol. 2000;227(2):373-87. doi: 10.1006/dbio.2000.9901. PubMed PMID: 11071761.

59.   Muller B, Hartmann B, Pyrowolakis G, Affolter M, Basler K. Conversion of an extracellular Dpp/BMP morphogen gradient into an inverse transcriptional gradient. Cell. 2003;113(2):221-33. doi: 10.1016/s0092-8674(03)00241-1. PubMed PMID: 12705870.

60.   Clark JF, Savage-Dunn C. Delta-9 fatty acid desaturase mutants display increased body size. MicroPubl Biol. 2018;2018. Epub 2018/09/18. doi: 10.17912/SS8E-6587. PubMed PMID: 32550399; PubMed Central PMCID: PMCPMC7282526.

61.   Tang LTH, Trivedi M, Freund J, Salazar CJ, Rahman M, Ramirez-Suarez NJ, et al. The CATP-8/P5A-type ATPase functions in multiple pathways during neuronal patterning. PLoS Genet. 2021;17(7):e1009475. Epub 20210701. doi: 10.1371/journal.pgen.1009475. PubMed PMID: 34197450; PubMed Central PMCID: PMCPMC8279360.

62.   Gaynor RB, Muchardt C, Diep A, Mohandas TK, Sparkes RS, Lusis AJ. Localization of the zinc finger DNA-binding protein HIV-EP1/MBP-1/PRDII-BF1 to human chromosome 6p22.3-p24. Genomics. 1991;9(4):758-61. doi: 10.1016/0888-7543(91)90371-k. PubMed PMID: 2037300.

63.   Oukka M, Kim ST, Lugo G, Sun J, Wu LC, Glimcher LH. A mammalian homolog of Drosophila schnurri, KRC, regulates TNF receptor-driven responses and interacts with TRAF2. Mol Cell. 2002;9(1):121-31. doi: 10.1016/s1097-2765(01)00434-8. PubMed PMID: 11804591.

64.   Oukka M, Wein MN, Glimcher LH. Schnurri-3 (KRC) interacts with c-Jun to regulate the IL-2 gene in T cells. J Exp Med. 2004;199(1):15-24. doi: 10.1084/jem.20030421. PubMed PMID: 14707112; PubMed Central PMCID: PMCPMC1887724.

65.   Liu Z, Shi H, Szymczak LC, Aydin T, Yun S, Constas K, et al. Promotion of bone morphogenetic protein signaling by tetraspanins and glycosphingolipids. PLoS Genet. 2015;11(5):e1005221. Epub 20150515. doi: 10.1371/journal.pgen.1005221. PubMed PMID: 25978409; PubMed Central PMCID: PMCPMC4433240.

66.   Clare CE, Brassington AH, Kwong WY, Sinclair KD. One-Carbon Metabolism: Linking Nutritional Biochemistry to Epigenetic Programming of Long-Term Development. Annu Rev Anim Biosci. 2019;7:263-87. Epub 20181109. doi: 10.1146/annurev-animal-020518-115206. PubMed PMID: 30412672.

67.   McMahon L, Muriel JM, Roberts B, Quinn M, Johnstone IL. Two sets of interacting collagens form functionally distinct substructures within a Caenorhabditis elegans extracellular matrix. Mol Biol Cell. 2003;14(4):1366-78. Epub 2003/04/11. doi: 10.1091/mbc.e02-08-0479. PubMed PMID: 12686594; PubMed Central PMCID: PMCPMC153107.

68.   Fernando T, Flibotte S, Xiong S, Yin J, Yzeiraj E, Moerman DG, et al. C. elegans ADAMTS ADT-2 regulates body size by modulating TGFbeta signaling and cuticle collagen organization. Dev Biol. 2011;352(1):92-103. Epub 2011/01/25. doi: 10.1016/j.ydbio.2011.01.016. PubMed PMID: 21256840; PubMed Central PMCID: PMCPMC3049821.

69.   Suzuki Y, Morris GA, Han M, Wood WB. A cuticle collagen encoded by the lon-3 gene may be a target of TGF-beta signaling in determining Caenorhabditis elegans body shape. Genetics. 2002;162(4):1631-9. Epub 2003/01/14. PubMed PMID: 12524338; PubMed Central PMCID: PMCPMC1462364.

70.   Kim KK, Sheppard D, Chapman HA. TGF-beta1 Signaling and Tissue Fibrosis. Cold
Spring Harb Perspect Biol. 2018;10(4). Epub 2017/04/23. doi: 10.1101/cshperspect.a022293.
PubMed PMID: 28432134; PubMed Central PMCID: PMCPMC5880172.
71.   MacFarlane EG, Haupt J, Dietz HC, Shore EM. TGF-beta Family Signaling in
Connective Tissue and Skeletal Diseases. Cold Spring Harb Perspect Biol. 2017;9(11). Epub
2017/03/02. doi: 10.1101/cshperspect.a022269. PubMed PMID: 28246187; PubMed Central
PMCID: PMCPMC5666637.

1036

Figure 1

Figure 2

a

b SMA-3 Targets

- Direct DEGs
- Non-Direct DEGs
- Non-DEGs

c

RNA-seq
1867 DEGs by edgeR
FDR≤0.05
Wild Type vs. *sma-3(wk30)*

ChIP-seq
Genomic Coordinates of 4205 SMA-3 Peaks

BETA
3 kb Window
FDR ≤ 0.05

367 Directly Regulated Target Genes *Down in Mutant*

d SMA-9 Targets

- Direct DEGs
- Non-Direct DEGs
- Non-DEGs

e

RNA-seq
783 DEGs by edgeR
FDR ≤ 0.05
Wild Type vs. *sma-9(ok1628)*

ChIP-seq
Genomic Coordinates of 7065 SMA-9 Peaks

BETA
3 kb Window
FDR ≤ 0.05

176 Directly Regulated Target Genes *Down in Mutant*

156 Directly Regulated Target Genes *Up in Mutant*

Figure 3

a

129 Directly Co-regulated Target Genes:
114 DEGs Activated By SMA-3 and SMA-9
15 DEGs Activated By SMA-3 But Repressed By SMA-9

b

238 DEGs
dpy-17, dpy-10, nhr-114, dpy-9, dpy-2, sma-6, sma-2, acdh-1, hsp-3, dpy-11

Annotation Cluster	EASE Score
Nucleus	12.08
Ribosome Biogenesis	3.97
Zinc-finger	3.76
mRNA Splicing	3.38
RNA Helicases	3.08
Chromosome Organization	2.84
ER Chaperones	2.61
DNAJ Domain	1.88

c

114 DEGs
lbp-8, col-94, col-153, ins-37, fat-6, hsp-60, sams-1, cysl-2

Annotation Cluster	EASE Score
Ribosome Biogenesis	3.46
Flavoproteins	3.01
One-carbon Metabolism	2.63
AAA-type ATPases	1.94
Mitochondria	1.92
Glutathione S-Transferase	1.38

d

15 DEGs
mpc-1, gdh-1 C54E4.5

e

129 DEGs
cyp-13A9, lys-4, clec-186, nhr-114

Annotation Cluster	EASE Score
Lipid Metabolism	3.80
Fatty Acid Oxidation	2.99
Peptidases	2.85
Crontonases	2.52
Long FA Metabolism	2.42
Oxido-reductases	2.34
Cytochrome P450s	2.19

f

150 DEGs
zip-10, ins-7, lon-1, sma-3

Annotation Cluster	EASE Score
Innate Immunity	5.23

Figure 3 – figure supplement 1

a

SMA-3 Direct Targets Annotation Clusters

Annotation Cluster	EASE Score
Ribosome Biogenesis	6.18
Oxido-reductases	3.31
Mitochondria	3.10
One-carbon Metabolism	3.03
RNA Helicases	2.95
ER Chaperones	2.89
Chromosome Organization	2.66
mRNA Splicing	2.64
Fatty Acid Metabolism	2.58
Amino Acid Biosynthesis	2.25
RNA Binding	2.08
Methionine Biosynthesis	2.00
Apple Domain	1.86
Acyl-CoA Dehydrogen	1.74
Isomerases	1.67
DNAJ Domain	1.46

b

SMA-9 Direct Targets Annotation Clusters

Annotation Cluster	EASE Score
Innate Immunity	7.44
Fatty Acid Metabolism	4.03
One-carbon Metabolism	3.23
Amino Acid Metabolism	2.73
Anti-Bacterial	2.67
Fatty Acid Transporters	2.62
Cytochrome P450	2.58
Lysosomes & Proteolysis	2.56
Cofactor Biosynthesis	2.33
Crontonases	2.14
Long FA Metabolism	2.05
Glutathione Metabolism	1.61
C-type Lectins	1.60

Figure 4

Figure 5

a

b

c

d

a

b

c

Figure 6

Figure 6 – figure supplement 1

Figure 7

Figure 7 – figure supplement 1

a Untreated

b Tunicamycin (5 μg/mL)