Preeclampsia-specific immune cell network in placenta revealed by
Cytometry by time of flight and single-cell RNA-seq

Haiyi Fei[#1,2,3], Xiaowen Lu [#1,2,3], Zhan Shi[#1,2,3], Xiu Liu[1,2,3], Cuiyu Yang[1,2,3], Xiaohong Zhu[4],
Yuhan Lin[1,2,3], Ziqun Jiang[1,2,3], Jianmin Wang[1,2,3], Dong Huang[1,2,3], Liu Liu[1,2,3], Songying
Zhang*[1,2,3], Lingling Jiang*[1,2,3]

[1]Assisted Reproduction Unit, Department of Obstetrics and Gynecology, Sir Run Run Shaw
Hospital, School of Medicine, Zhejiang University, Hangzhou, 310016, China.

[2]Zhejiang Provincial Clinical Research Center for Obstetrics and Gynecology, Hangzhou,
310016, China.

[3]Zhejiang Key Laboratory of Precise Protection and Promotion of Fertility, Hangzhou, 310016,
China.

[4]Department of Obstetrics and Gynecology, Zhejiang Xiaoshan Hospital, 310016, Hangzhou,
China

*Co-corresponding authors
Lingling Jiang
Assisted Reproduction Unit, Department of Obstetrics and Gynecology,
Sir Run Run Shaw Hospital, Zhejiang University School of Medicine,
310000, Hangzhou, China
E-mail: linglingjiang@zju.edu.cn
20
21 Songying Zhang
22 Assisted Reproduction Unit, Department of Obstetrics and Gynecology,
23 Sir Run Run Shaw Hospital, Zhejiang University School of Medicine,
24 310000, Hangzhou, China
25 E-mail: zhangsongying@zju.edu.cn
26

Abstract

Preeclampsia (PE), a major cause of maternal and perinatal mortality with highly
heterogeneous causes and symptoms, is usually complicated by gestational diabetes mellitus
(GDM). However, a comprehensive understanding of the immune microenvironment in the
placenta of PE and the differences between PE and GDM is still lacking. In this study,
Cytometry by time of flight (CyTOF) indicated that the frequencies of memory-like Th17 cells
(CD45RA$^-$CCR7$^+$IL-17A$^+$CD4$^+$), memory-like CD8$^+$ T cells
(CD45RA$^-$CCR7$^+$CD38$^+$pAKT^{mid}CD127^{low}) and pro-inflam Macs
(CD206$^-$CD163$^-$CD38^{mid}CD107a^{low}CD86^{mid}HLA-DR^{mid}CD14$^+$) were increased, while the
frequencies of CD69^{hi}Helios^{mid}CD127^{mid} γδT cells, anti-inflam Macs
(CD206$^+$CD163$^-$CD86^{mid}CD33$^+$HLA-DR$^+$) and granulocyte myeloid-derived suppressor cells
(gMDSCs, CD11b$^+$CD15^{hi}HLA-DR^{low}) were decreased in the placenta of PE compared with
that of NP, but not in that of GDM or GDM&PE. The pro-inflam Macs were positively correlated

with memory-like Th17 cells and memory-like CD8$^+$ T cells but negatively correlated with
gMDSCs. Single-cell RNA sequencing revealed that transferring the F480$^+$CD206$^-$ pro-inflam
Macs with a Folr2$^+$Ccl7$^+$Ccl8$^+$C1qa$^+$C1qb$^+$C1qc$^+$ phenotype from the uterus of PE mice to
normal pregnant mice induced the production of memory-like
IL-17a$^+$Rora$^+$Il1r1$^+$TNF$^+$Cxcr6$^+$S100a4$^+$CD44$^+$ Th17 cells via IGF1-IGF1R, which contributed to
the development and recurrence of PE. Pro-inflam Macs also induced the production of
memory-like CD8$^+$ T cells but inhibited the production of
Ly6g$^+$S100a8$^+$S100a9$^+$Retnlg$^+$Wfdc21$^+$ gMDSCs at the maternal-fetal interface, leading to
PE-like symptoms in mice. In conclusion, this study revealed the PE-specific immune cell
network, which was regulated by pro-inflam Macs, providing new ideas about the
pathogenesis of PE.
51
Introduction
In recent years, low fertility is a global problem (Agbaglo et al., 2022), exacerbated by
pregnancy complications. Preeclampsia (PE) is a progressive systemic disease during
pregnancy, with pregnancy-induced hypertension, proteinuria, and liver and kidney injury as
the main diagnostic indicators (Chappell et al., 2021). With a global prevalence of 7–10%, PE
is a major cause of maternal and perinatal mortality and morbidity, especially in low-income
and middle-income countries. The only specific treatment is delivery (Chappell et al., 2021).
However, due to the variety of symptoms and the heterogeneity of the disease, the
pathogenesis of PE is still unclear (Burton et al., 2019; Grotegut, 2016).
PE is often complicated by gestational diabetes mellitus (GDM), which is characterized by
insulin resistance and associated with aberrant maternal immune cell adaption (Corrêa-Silva
et al., 2018; McElwain et al., 2021; McIntyre et al., 2019). Studies have shown that GDM is an
independent risk factor for preeclampsia (Nerenberg et al., 2013; Ostlund et al., 2004;
Schneider et al., 2012). However, whether a different pathogenesis underlies between PE and
GDM is still unclear.
The fetus is equivalent to a semi-homogenous graft for the mother. Therefore,
maternal-fetal immune tolerance plays an important role in the maintenance of pregnancy.
Elevated levels of maternal inflammation are now considered to be an important mechanism in
the pathogenesis of preeclampsia (Aneman et al., 2020; Deer et al., 2023; Jung et al., 2022),
including macrophages (Faas et al., 2014; Yao et al., 2019), granulocytes (Lampé et al., 2015;
Lampé et al., 2011; D. Liu et al., 2021), natural killer (NK) cells (Bachmayer et al., 2006; Fukui
et al., 2011; Travis et al., 2020), innate B1 cells (Jensen et al., 2012; LaMarca et al., 2011;
Zhong et al., 2007) and γδ T cells (Chatterjee et al., 2017). Moreover, adaptive immune
response is also critical for the pathogenesis of PE (Deer et al., 2023). In PE, CD4$^+$ T cells
prefer to differentiate into pro-inflammatory Th1 and Th17 phenotypes (Eghbal-Fard et al.,
2019; Fu et al., 2014; Lang et al., 2021; Lu et al., 2020); Saito et al. (2007) instead of
non-inflammatory Th2 and Treg phenotypes (Care et al., 2018; Cornelius et al., 2015a; Saito
et al., 2007; Santner-Nanan et al., 2009; Sasaki et al., 2007). However, a comprehensive and
in-depth understanding of the maternal-fetal interface of PE is still lacking.
The immune cells crosstalk with each other at the maternal-fetal interface elaborately. The
interaction between B2 cells and Th cells depends on CD40 and its ligand CD40L, which helps
B2 cells provide AT1-AAs and differentiate into memory B cells (Cornelius et al., 2015b). The

cytokines derived from NK cells can promote the differentiation of Th17 to co-work in recurrent
miscarriage (Fu et al., 2013). Nevertheless, studies about the regulation of the immune
network at the maternal-fetal interface of PE still need to be completed.
In this study, we combined cytometry by time of flight (CyTOF), single-cell RNA
sequencing (scRNA-seq), and rodent experiment to identify the overall immune cell
composition and their interactions at the maternal-fetal interface in PE, GDM, and GDM
complicated with PE (GDM&PE). This study revealed the PE-specific immune cell network,
which was regulated by pro-inflammatory macrophages, providing new ideas about the
pathogenesis of PE.
93
Results
Overall immune cell profile in the placenta of PE, GDM, and GDM&PE
To fully characterize the immune microenvironment at the maternal-fetal interface of PE,
we collected full-term placentas of PE and normal pregnancy (NP), and did a CyTOF test
(**Table 1**), which panel was shown in **Table 2**. Since PE is usually complicated by GDM, the
subjects of GDM and GDM complicated with PE (GDM&PE) were also included. The
experimental workflow for CyTOF is shown in the schematic diagram (**Figure 1A**). We found
that in addition to the well-known macrophages and T cells, there are also γδ T cells, B cells,
NK cells, granulocytes, dendritic cells (DCs), and myeloid-derived suppressor cells (MDSCs).
An overall distribution of CD45$^+$ cell subsets in placentas was shown in the t-SNE maps
(**Figure 1B**). There were no significant differences in the proportion of these large
subpopulations comparing the PE, GDM, GDM&PE group to the NP group (**Figure 1C**). To
identify the ten cell subpopulations, CD4, CD8, and γδTCR were used as markers to
distinguish CD4$^+$ T cells, CD8$^+$ T cells, and γδ T cells. CD11b is a marker that distinguishes
myeloid cells, including macrophages and granulocytes (**Figure 1D**). The ten cell subsets
were further annotated by 41 markers in the heatmap (**Figure 1E, Table 3**).
Specific altered T cell profile in the placenta of PE
To fully understand the distribution in each cell subset, we first analyzed 15 clusters of
CD4$^+$ T cells from the placentas of NP, PE, GDM, and GDM&PE (**Figure 2A**). CD4$^+$ T cell
clusters were defined by canonical marker set signatures (**Figure 2-figure supplement 1 A**).
The frequencies of memory-like CD45RA$^-$CCR7^{mid}CD28$^+$CD127$^+$CD4$^+$ T cells (cluster 8) were
significantly increased in individuals with PE compared with NP controls, while. At the same
time, there was no significance in the GDM and GDM&PE groups compared with the NP group
(**Figure 2B, Table 4**). We analyzed the expression of common intracellular molecules in CD4$^+$
memory-like T cells by sorting CD45RO$^+$CCR7$^+$CD4$^+$ T cells from placental samples by flow
cytometry. Higher levels of IL-17A and lower levels of Foxp3 were found in memory-like
CD45RO$^+$CCR7$^+$CD4$^+$ T cells in the placentas of individuals with PE (**Figure 2C**). Consistent
with the results of CyTOF, the higher fluorescence intensity of CD45RO$^+$CD4$^+$ T cells was
found in the placentas of individuals with PE (**Figure 2D**). Moreover, lower expression of
immune checkpoint molecules including T-cell immunoglobulin mucin-3 (Tim-3) and
programmed cell death 1 (PD-1) was found in CD45RA$^-$CCR7$^+$CD4$^+$ memory T cells in the PE
group, suggesting that these cells have a lower immunosuppressive capacity (Chen et al.,
2022; Fanelli et al., 2021; Rasmussen et al., 2022; Wang et al., 2016), though there is no
significant difference in the GDM or GDM&PE groups comparing with the NP group (**Figure**

2E). Then, 18 clusters of CD8$^+$ T cells were analyzed (**Figure 2F**). CD8$^+$ T cell clusters
annotation was based on the expression of canonical marker signatures (**Figure 2-figure**
supplement 1 B). CD45RA$^-$CCR7$^+$CD38$^+$pAKT^{mid}CD127^{low} (cluster 2) memory-like CD8$^+$ T
cells were were significantly increased in the PE group (**Figure 2G, Table 4**). Cluster 2
expressed lower expression of PD-1 and TIGIT, suggesting the activation of cytotoxicity of
these cells (**Figure 2H**) (Morita et al., 2020). Then, 22 clusters of γδT cells were analyzed and
annotated (**Figure 2I, Figure 2-figure supplement 1 C**). Cluster 15, which expressed lower
expression of Tim3, was significantly decreased in the PE group (**Figure 2J, 2K**). However,
regardless of CD4$^+$ T cells or CD8$^+$ T cells, there is no significant change in the memory-like
clusters in the GDM or GDM&PE group compared to the NP group.
In conclusion, significant changes in placental T cell profile were found in the placentas of
PE but not in those of GDM or GDM&PE, suggesting that abnormal activation of T cells in the
placenta is associated with the pathogenesis of PE.
Abnormal polarization of macrophages was correlated with specific immune cell
subsets in individuals with PE
Except for T cells, we also analyzed 29 clusters of CD45$^+$CD3$^-$CD11b$^+$ cells from
placentas of NP, PE, GDM, and GDM&PE, including 9 clusters of macrophages, 11 clusters of
granulocytes, and 5 clusters of NK/NK-like cells (**Figure 3A**). The clusters of
CD45$^+$CD3$^-$CD11b$^+$ cells were defined by canonical marker set signatures (**Figure 3 B**).
Moreover, significantly decreased frequencies of CD11b$^+$CD15^{hi}HLA-DR^{low} granulocytes
(cluster 12), which are identified as granulocyte myeloid-derived suppressor cells (gMDSCs),
were found in the PE group (**Figure 3C, Table 4**). In addition, the frequency of
anti-inflammatory macrophages (anti-inflam Macs) (CD206$^+$CD163$^-$CD86^{mid}CD33$^+$HLA-DR$^+$,
cluster 25) was also significantly decreased, whereas the frequency of pro-inflammatory
macrophages (pro-inflam Macs) (CD206$^-$CD163$^-$CD38^{mid}CD107a^{low}CD86^{mid}HLA-DR^{mid}CD14$^+$,
cluster 23) was significantly increased in the PE group (**Figure 3C, Table 4**). However, the
frequencies of macrophages and gMDSCs were unchanged significantly in the GDM or
GDM&PE groups.
Pearson correlation analysis indicated positive correlations between pro-inflam Macs
(cluster 23 in CD11b$^+$ cells) and CD4$^+$ memory-like T cells (cluster 8 of CD4$^+$ T cells), as well as
CD8$^+$ memory-like T cells (cluster 2 in CD8$^+$ T cells). CD206$^-$ pro-inflam Macs were negatively
correlated with gMDSCs (cluster 12 in CD11b$^+$ cells), but were not statistically significant.
Conversely, CD206$^+$ anti-inflam Macs (cluster 25 in CD11b$^+$ cells) were positively correlated
with gMDSCs and negatively correlated with CD8$^+$ memory-like T cells (**Figure 3D, 3E**).
These results suggested that abnormally polarized CD206$^-$pro-inflam Macs are positively
correlated with memory-like CD4$^+$ and CD8$^+$ T cells in the placentas of individuals with PE and
negatively correlated with gMDSCs.
F480$^+$CD206$^-$ pro-inflam Macs induced immune imbalance at the maternal-fetal interface
and PE-like symptoms
Though increased pro-inflam Macs have been reported in the placenta from individuals
with PE (Faas et al., 2014), few studies have reported the interaction between macrophages
and other immune cells in the placenta. CD45$^+$F4/80$^+$CD206$^-$ pro-inflam Macs or
CD45$^+$F4/80$^+$CD206$^+$ anti-inflam Macs were isolated from the uterus and placentas of mice
with RUPP and injected into normal pregnant mice (**Figure 4A**). RNA-seq was used to analyse

the difference between CD45$^+$F4/80$^+$CD206$^-$ pro-inflam Macs and CD45$^+$F4/80$^+$CD206$^+$

anti-inflam Macs. Significant differences were found between the two groups of macrophages

(**Figure 4B**). We found that the expression of 2123 genes was increased, and 2699 genes

decreased significantly in the pro-inflam Macs compared to the anti-inflam Macs (**Figure 4C**).

The expression levels of genes associated with inflammatory response (Ccl7, Ccl8, Ccl2, IL6)

(He et al., 2019; Wu et al., 2023), complement system activation (C1qa, C1qb) (Chen et al.,

2021), and lipid metabolism (Pltp, Apoe) (Desrumaux et al., 2016) were significantly increased

in the pro-inflam Macs; whereas the expression levels of genes associated with the regulation

of angiogenesis and vascular endothelial growth factor production (VEGFα, Macro) and tissue

development (TGF-β1, Thbs1, Fn1, Slpi) (Jin et al., 2023; Nugteren et al., 2021; Li et al., 2022)

were significantly decreased (**Figure 4C, 4D**). Moreover, Gene Set Enrichment Analysis

(GSEA) was performed to further explore the most significantly enriched functional terms

between the two groups of macrophages. We found that NF-κB signaling and interspecies

interaction between organisms were enriched in the pro-inflam Macs, while tissue

development and epithelial cell morphogenesis were enriched in the anti-inflam Macs (**Figure**

4E).

To further investigate the effect of the pro-inflam Macs polarization in immune imbalance

at the maternal-fetal interface, PLX3397, the inhibitor of CSF1R, which is needed for

macrophage development, was used to deplete the macrophages of pregnant mice (X. Chen

et al., 2023;J. Chen et al., 2022). As expected, an increased embryo resorption rate,

decreased fetal top-rump length and fetal weight were found in mice injected with pro-inflam

Macs (**Figure 4F**). Increased SBP and UACR were also found in mice injected with pro-inflam

Macs (**Figure 4G**). Moreover, it is shown that CD44$^+$ memory-like Th17 cells and memory-like

CD8$^+$ T cells increased while CD11b$^+$Ly6G$^+$ gMDSCs decreased in mice injected pro-inflam

Macs compared with mice injected anti-inflam Macs, which validated our findings in CyTOF

(**Figure 4H**). Clodronate liposomes were also used to deplete the macrophages of pregnant

mice (X. Liu et al., 2022) before pro-inflam Macs or anti-inflam Macs were injected into the

mice. The same experimental results were obtained (**Figure 4-figure supplement 1**). In

conclusion, the F480$^+$CD206$^-$ pro-inflam Macs induced immune imbalance at the

maternal-fetal interface and PE-like symptoms.

Pro-inflam Macs and anti-inflam Macs subsets are functionally heterogeneous

To further explore the role that macrophages play in the immune imbalance at the

maternal-fetal interface in PE, scRNA-seq was performed to analyze the uterine CD45$^+$

immune cells from mice that were transferred with pro-inflam Macs or anti-inflam Macs from

the the uterus and placentas of PE mice. An unsupervised cluster detection algorithm

(SEURAT) was applied and eight types of immune cells were detected by mostly

distinguishable cell type-specific genes, including macrophages, monocytes, granulocytes,

T/NK cells, B cells, plasma, mast cells, and basophils (**Figure 5A, 5B**).

Macrophages were further identified into 15 clusters defined by marker set signatures.

Single-cell differential expression analysis (SCDE) was performed for each population and

characteristic gene expression patterns were detected for cluster 0-cluster 14 to characterize

the phenotypes of these subsets in detail (**Figure 5C**). We found that the frequency of cluster

0 was significantly increased in mice injected with pro-inflam Macs, while the frequency of

cluster 1 was significantly decreased (**Figure 5D**). We also found that cluster 0 highly

expresses genes associated with foetal and tissue resident (Folr2) (Nalio et al., 2022;
Thomas et al., 2021), complement system activation (C1qa, C1qb, C1qc) (Chen et al., 2021),
and inflammatory response (Ccl7, Ccl8) (He et al., 2019; Wu et al., 2023); while cluster 1
highly expresses genes associated with tissue repair (Chil3, Slpi, Fn1) (Jin et al., 2023;
Nugteren et al., 2021; Li et al., 2022), blood vessel morphogenesis (Thbs1) (Che et al., 2021)
and preventing oxidative stress (Gsr, Mgst1) (Coppo et al., 2022) (**Figure 5E**). Changes in
gene expression patterns of cluster 0 (pro-inflam Macs) and cluster 1 (anti-inflam Macs) were
analyzed in mice injeced the pro-inflam Macs or anti-inflam Macs. We found that in the
pro-inflam Macs group, enriched GO terms in cluster 0 and cluster1 including 'antigen
processing and presentation of exogenous antigen' and 'inflammatory response', in which
high expression of genes such as CCL5, CCL8, CALR, IRF7, IL10, IFI44, IFI30 , OAS3, etc.
could be observed (**Figure 5F, 5G**). The control group showed elevated expression of
MMP14, Chil3, EGR1, ATF3 and VASP; and these genes were enriched in 'vascular
endothelial growth factor production' and 'tissue development' (**Figure 5F, 5G**). These results
were consistent with the results of the transcriptome RNA sequencing results of
macrophages.
These data suggested that the CD45$^+$F480$^+$CD206$^-$ pro-inflam Macs with a
Folr2$^+$Ccl7$^+$Ccl8$^+$C1qa$^+$C1qb$^+$C1qc$^+$ phenotype play an improtant role in the development of
PE.

Pro-inflam Macs induced the memory-like Th17 cells, which was associatied with the
development and recurrence of PE
We also analyzed the scRNA-seq data of uterine T/NK cells from mice that were injected
with or without pro-inflam Macs. T/NK cells were further identified into 12 clusters defined by
marker set signatures (**Figure 6A, 6B**). We found that the frequency of cluster 0 was increased
in mice injected with pro-inflam Macs, while the frequency of cluster 1 and cluster 2 were
significantly decreased (**Figure 6A**). In addition to genes associated with Th17 cells (IL-17a,
IL17f, Rora, Il1r1, TNF) (Wu et al., 2019; Leite et al., 2023), cluster 0 also exhibits high
expression of genes associated with memory phenotype (Cxcr6, S100a4, CD44) (Schroeder
et al., 2023; Bieberich et al., 2021), suggesting that cluster 0 was the memory-like Th17 cells
(**Figure 6A, 6C**). Cluster 1 exhibits high expression of genes associated with
immunoregulation (Lef1, Tcf7, Ccr7) (Qiu et al., 2022; Sekine et al., 2020); Cluster 2 exhibits
high expression of genes associated with immunosuppression (Gata3, GITR, CD28),
suggesting that cluster 1 and 2 were T cells with immunosuppressive function (Esensten et al.,
2016; Pai et al., 2023) (**Figure 6A, 6C**).
Changes in gene expression patterns of cluster 0 (memory-like Th17 cells) were analyzed
in the pro-inflam Macs group and Control group. We found that in the pro-inflam Macs group,
enriched GO terms including 'positive regulation of response to external stimulus' and 'tumor
necrosis factor production', in which highly express genes such as TNFSF11, TNF, IL27RA,
IGF1R, CD226, LAMP1 (**Figure 6C, 6D**). The control group showed elevated expression of
ASS1, EIF5, S100A9, CTLA4, S100A8 and CXCR4; these genes were enriched in GO terms
including 'regulation of programmed cell death' and 'cellular biosynthetic process' (**Figure 6D,**
6E).
Combining the CyTOF and sc-RNA seq data, we found the frequency of memory-like
IL-17A$^+$CD4$^+$ T cells significantly increased at the maternal-fetal interface of individuals with

PE. To confirm the effect of memory-like Th17 cells in PE, we constructed an animal model of
PE by reducing uterine perfusion pressure (RUPP), and mice with sham operation were
considered as controls (**Figure 6-figure supplement 1A**). We found that mice of PE showed
an increased embryo absorption rate, decreased fetal weight, and pup crown-rump length
compared with the sham operation group (**Figure 6-figure supplement 1B**). Increased
systolic blood pressure (SBP) and urine albumin creatine ratio (UACR) were also observed in
mice with RUPP, which indicated a successful PE mice mode was built (**Figure 6-figure**
supplement 1C). Our studies revealed that the frequency of memory-like Th17 cells
(CD4$^+$CD44$^+$IL-17A$^+$ T cells) was significantly increased in the PE mice (**Figure 6F**).
To confirm the importance of memory-like CD4$^+$ T cells in the pathogenesis of PE, the
CD4$^+$CD44$^+$ T cells in the uterus and placentas from PE or NP mice were sorted and
intravenously injected the cells into normal pregnant mice on day12.5 of gestation (**Figure 6F**).
Increased embryo absorption rate, decreased fetal weight and pup crown-rump length were
found in mice injected with PE mouse-derived CD4$^+$CD44$^+$ T cells (**Figure 6G**). The typical
PE-like symptoms, including increased SBP and UACR, were also observed in those mice
(**Figure 6H**).
It has been reported that sustained expansion of immunosuppressive memory Tregs
during prior pregnancy is beneficial for maintaining a second pregnancy (Rowe et al., 2012). To
verify whether memory-like Th17 cells promote the recurrence of PE, we established a second
pregnant mouse model with a history of PE or NP in the first pregnancy. An increased embryo
resorption rate, decreased crown-rump length, and fetal weight were found in mice with a
history of PE pregnancy compared with those with a history of NP pregnancy (**Figure**
6—figure supplement 1D). Moreover, mice with a history of PE in the first pregnancy showed
increased SBP and UACR during the second pregnancy (**Figure 6—figure supplement 1E**).
Consistently, higher levels of IL-17A were also found in memory-like CD4$^+$ T cells in mice with
a history of PE in the first pregnancy (**Figure 6—figure supplement 1F**).
Therefore, pro-inflam Macs induced the IL-17a$^+$IL17f$^+$Rora$^+$Il1r1$^+$TNF$^+$Cxcr6$^+$S100a4$^+$
CD44$^+$ memory-like Th17 cells, which was associatied with the development and recurrence of
PE.
Pro-inflam Macs lead to a reduced proportion of gDMSCs
The uterine granulocytes from mice that were transferred with pro-inflam Macs from the
uterus and placentas of PE mice were further analyzed using sc RNA-seq. Granulocytes were
identified into 12 clusters defined by marker set signatures (**Figure 7A, 7B**).We found that
cluster 0-cluster 3 were different between the two groups: the frequency of cluster 0 and
cluster 2 were increased in mice injected with pro-inflam Macs, while the frequency of cluster 1
and cluster 3 were decreased (**Figure 7A**). Cluster 0 has the signature of a mature
granulocyte (FOS, FOSB) (Montaldo et al., 2022), and also highly expresses IL-1R2 and
CSF3R; Cluster 1 highly express genes associated with immunoregulation genes (Hmox1,
CCL4, CCL3, Lcn2) (Wenzel et al., 2022; Montaldo et al., 2022); Cluster 2 highly expressed
pro-inflammatory genes (IFITM1, IL1B, Ptgs2, Cxcl3) (Ulff-Møller et al., 2018; Drummond al.,
2022); while cluster 3 highly expressed gMDSCs associated genes (Ly6g, S100a8, S100a9,
Retnlg, Wfdc21) (von Wulffen et al., 2023; Kao al., 2023) (**Figure 7C**). Our animal experiments
above (Fig 4) have identified a reduced proportion of CD11b$^+$Ly6G$^+$ gMDSCs in the mice
injected with pro-inflam Macs. Changes in gene expression patterns of cluster 3 (gMDSCs)

were analyzed in the pro-inflam Macs group and the control group. Enriched GO terms
including 'tumor necrosis factor production' and 'leukocyte mediated immunity' were found in
the pro-inflam Macs group, which highly express the genes such as IRF7, EGR1, C1QA,
C1QB, CCL2, CSF3R, TNFRSF1B (Figure 7D, 7E). The control group showed elevated
expression of CCL4, CXCL10, CD177, MMP8, S100A13, LCN2 and SPP1; these genes were
enriched in 'granulocyte migration' and 'localization' ((Figure 7D, 7E).

In conclusion, from the rodent experiments and the data of scRNA-seq, we demonstrated
that pro-inflam Macs suppressed the production of $Ly6g^+S100a8^+S100a9^+Retnlg^+Wfdc21^+$
gMDSCs.

Pro-inflam Macs induced the production of memory-like Th17 cells via IGF1-IGF1R

Our results above showed that pro-inflam Macs induced memory-like Th17 cells in PE,
however, the underlying molecular mechanisms were still unknown. CellPhoneDB analysis
indicated that increased communication counts and signaling pathways numbers between
macrophages and T/NK cells were observed in mice injected with pro-inflam Macs (**Figure
8—figure supplement 1A, 1B**). Then we identified the interacting ligand-receptor pairs
between different types of immune cells and macrophages. Insulin-like growth factor 1 (IGF1)
-IGF1 receptor (IGF1R) ligand-receptor pair was significantly enhanced between
macrophages and T/NK cells in mice injected with pro-inflam Macs (**Figure 8A**). IGF1 has
significant effects on immune function maintenance, and signaling through IGF1R could cause
increased aerobic glycolysis, favoring Th17 cell differentiation over that of Treg cells
(Bekkering et al., 2018; DiToro et al., 2020). For further demonstration, we analyzed the
frequencies of $IGF1^+CD14^+$ cells and $IGF1R^+CD4^+$ cells in placentas of individuals with PE and
NP and found them significantly increased in the PE group (**Figure 8B**).

Trophoblast-derived extracellular vesicles from the placenta of PE (PE-EVs), which carry
the fetal antigen, could induce M1 macrophage polarization to participate in the development
of PE according to our previous study (X. Liu et al., 2022). To confirm whether
pro-inflammatory macrophages induced the production of memory-like Th17 cells via
IGF1-IGF1R, macrophages from human peripheral blood, after incubating with PBS,
trophoblast-derived extracellular vesicles from NP (NP-EVs) or PE-EVs in vitro, were
co-cultured with $CD4^+$ naïve T cells (**Figure 8C**). The frequencies of memory-like $CD45RO^+$
$CCR7^+$ $IL-17A^+$ $CD4^+$ cells significantly increased in the PE-EVs-induced macrophages group
rather than in the NP-EVs-induced macrophages group. However, the frequencies of
memory-like Th17 cells significantly decreased after $CD4^+$ naïve T cells were treated with the
IGF1R inhibitor BMS-754807, an inhibitor of IGF1R (**Figure 8D**).

To investigate the role of IGF1-IGF1R in the development of PE in vivo, $CD45^+$ $CD4^+$ T
cells from NP mice were isolated and treated with BMS-754807 or PBS, then injected into NP
mice at day 11.5 of gestation (**Figure 8E**). Lipopolysaccharide (LPS)-induced PE mice model
was constructed by intraperitoneal injection of LPS on day12.5 and day15.5 of gestation (Han
et al., 2021), and anti-CD4 antibody was used to deplete the $CD4^+$ T cells in pregnant mice.
The mice injected with $CD4^+$ T cells treated by BMS-754807 presented decreased embryo
resorption rate and increased pup crown-rump length and fetal weight (**Figure 8F**), decreased
SBP and UACR (**Figure 8G**), and decreased frequency of memory-like Th17 cells at the
maternal-fetal interface (**Figure 8H**). These data suggested that pro-inflam Macs could induce
the production of memory-like Th17 via the IGF1-IGF1R, leading to the development of PE.

Discussion

Preeclampsia, a progressive systemic disease during pregnancy, is closely related to the alterations in the immune environment at the maternal-fetal interface. In this study, the overall immune cell profile in the placenta of individuals with NP, PE, GDM and GDM&PE were detected by CyTOF and a PE-specific change in immune microenvironment of maternal-fetal interface were described for the first time. Further, we provided a novel insight that pro-inflam Macs induced memory-like Th17 cells, memory-like CD8$^+$ T cells, and suppressing the production of gMDSCs in mice maternal-fetal interface by scRNA-seq. In addition, we first validated that IGF1-IGF1R was involved in producing memory-like Th17 cells induced by pro-inflammatory macrophages in vitro and in vivo, thus leading to the development of PE.

Various studies represented the effect of a particular subset of immune cells in the pathogenesis of PE or GDM (Bachmayer et al., 2006; Care et al., 2018; Cornelius et al., 2015a; Eghbal-Fard et al., 2019; Faas et al., 2014; Fu et al., 2014; Fukui et al., 2011; Lampé et al., 2015; Lampé et al., 2011; Lang et al., 2021; D. Liu et al., 2021; Lu et al., 2020; Santner-Nanan et al., 2009; Sasaki et al., 2007; Travis et al., 2020; Yao et al., 2019). However, only one report has provided an overall analysis of immune cells in the human placental villi in the presence and absence of spontaneous labor at term by scRNA-seq (Miller et al., 2022). Here we used CyTOF and scRNA-seq to comprehensively analyze the immune cell profile and explore the interaction of immune cells in PE. Additionally, most previous studies have studied immune cells from maternal systemic circulation rather than directly from the tissue of maternal-fetal interface (Cornelius et al., 2015b; Deer et al., 2021; Santner-Nanan et al., 2009; Shields et al., 2018; Wallace et al., 2012). In this study, cells were directly isolated from the tissue of maternal-fetal interface, which is considered as a more accurate approach to mimic the real immune environment of the maternal-fetal interface.

Abnormal inflammatory responses were reported to play an important role in the pathogenesis of both PE and GDM (Aneman et al., 2020; Corrêa-Silva et al., 2018; Deer et al., 2023; Jung et al., 2022; McElwain et al., 2021). However, It is unclear how the maternal-fetal interface immune microenvironment differs between the two diseases. We analyzed the overall placental immune cell profile in NP, PE, GDM and GDM&PE by CyTOF and revealed a PE-specific immune cell profile: The frequencies of memory-like Th17 cells (CD45RA$^-$CCR7$^+$IL-17A$^+$CD4$^+$), memory-like CD8$^+$ T cells (CD45RA$^-$CCR7$^+$CD38$^+$pAKT^{mid}CD127^{low}) and pro-inflam Macs (CD206$^-$CD163$^-$CD38^{mid}CD107a^{low}CD86^{mid}HLA-DR^{mid}CD14$^+$) were increased, while the decreased frequencies of CD69^{hi}Helios^{mid}CD127^{mid} γδT cells, anti-inflam Macs (CD206$^+$CD163$^-$CD86^{mid}CD33$^+$HLA-DR$^+$) and granulocyte myeloid-derived suppressor cells (gMDSCs, CD11b$^+$CD15^{hi}HLA-DR^{low}) were observed in the placenta of PE, but not in that of GDM or GDM&PE compared with that of NP.

PE is often divided into two subtypes based on the time of onset. Late-onset PE, occurring at 34 weeks of gestation or later, accounts for most preeclampsia cases, while early-onset PE, occurring before or at 33 weeks of gestation(Lisonkova and Joseph, 2013). However, our previous study observed no significant difference in the M1/M2 macrophage polarization in placentas with early-onset or late-onset PE (X. Liu et al., 2022). Therefore, our research was conducted regardless of the preeclampsia-onset type to ensure consistency in macrophage polarization of the two subtypes.

There have been studies indicating that macrophages, Th17 cells, Treg cells, and CD8[+] T
cells play an essential role in PE development (Care et al., 2018; Eghbal-Fard et al., 2019;
Lager et al., 2020; Yao et al., 2019). Macrophages was indicated with a pro-inflammatory
phenotype in the placenta with PE (Faas et al., 2014; Yao et al., 2019). Care et al. reported
that placental Tregs helped suppress inflammation at the maternal-fetal interface (Care et al.,
2018). Similarly, Lu et al. found an abnormal increase of IL-17A in the placenta in the
development of PE (Lu et al., 2020). Meanwhile, the abnormal infiltration of CD8[+] T cells in the
placenta may contribute to PE (Lager et al., 2020). Consistent with the previous findings, we
found that memory-like CD4[+] and memory-like CD8[+] T cells significantly increased in the PE
group and that IL-17A was significantly higher expressed in CD4[+] memory-like T cells from the
PE group, which is vital in vascular inflammation, leading to the development of PE (Amador et
al., 2014; Madhur et al., 2021). Though memory T cells play a key role in fetal-maternal
tolerance in normal pregnancy (Kieffer et al., 2019), few studies have reported the
characteristics of memory-like T cells in the development of PE.
We constructed an adoptive transferred mouse model and confirmed that pregnant mice
developed PE symptoms after transferring PE mouse-derived memory-like CD4[+] T cells. It is
worth noting that we isolated CD4[+] memory-like T cells directly from the maternal-fetal
interface, which is different from previous studies that the CD4[+] T cells for adoptive transfer
were isolated from the spleen and induced in vitro (Cornelius et al., 2015b; Deer et al., 2021;
Shields et al., 2018; Wallace et al., 2012). Recurrence is an important concern in individuals
with PE (Bernardes et al., 2019). We found that a prior PE pregnancy increased the probability
of PE in the next pregnancy in mice, accompanied by an increased frequency of memory-like
Th17 cells at the maternal-fetal interface, suggesting that increased memory-like Th17 cells
are positively related to the recurrence of PE. However, additional animal experiments are
needed to confirm the function of the fetal-specific memory Th17 cells in the recurrence of PE.
Activation of CD8[+] T cells and Th1 cells can be induced by abnormal polarized
macrophages in recurrent miscarriage (Li et al., 2022). gMDSCs have been reported to have
immune suppressive activity and are necessary for maintaining maternal-fetal tolerance
(Köstlin-Gille et al., 2019; Zhou et al., 2018). However, the correlation between macrophages
and other immune cells in the development of PE remained unclear. We verified that
pro-inflam Macs induced memory-like Th17 cells and memory-like CD8[+] T cells and inhibited
the production of gMDSCs in PE by correlation analysis and adoptive transfer in animal
experiments for the first time, indicating that the immune cells orchestrate a network in the
maternal-fetal interface elaborately.
The function of IGR1 and IGF1R in PE is controversial. IGF1 plays a significant role in
maintaining immune function, and IGF1R facilitates the differentiation of naïve CD4[+] T cells
into Th17 cells (DiToro et al., 2020). A study showed that IGF-1R was downregulated in the
serum of individuals with PE (Liao et al., 2021). Decreased amounts of IGF1R were also found
in the placentas of individuals with PE (Robajac et al., 2015). However, another study found no
significance in the affinity and the number of IGF-1R between placentas from individuals with
PE or not (Díaz et al., 2005). We found that mice exhibiting PE symptoms had significantly
enhanced IGF1-IGF1R interaction between inflammatory macrophages and memory-like Th17
cells by analyzing the data of scRNA-seq. Then, this finding was further confirmed in human
placentas from individuals with PE and NP, for the frequencies of IGF1[+] CD14[+] cells and

IGF1R$^+$ CD4$^+$ cells were found to significantly increase in the PE group. In vitro, we found that PE-EVs-treated macrophages secreted more IGF1 than NP-EV-treated macrophages and induced a higher frequency of memory-like Th17 cells. The inhibition of IGF1R on CD4$^+$ naïve T cells resulted in a decreased frequency of memory-like Th17 cells, implying that the IGF1-IGF1R is critical for the production of memory-like Th17 cells induced by inflammatory macrophages. Animal experiments confirmed that IGF1R inhibition on CD4$^+$ T cells at the maternal-fetal interface reduced the frequency of memory-like Th17 cells, improving PE.

In addition, placentas from different groups for CyTOF analysis and flow cytometry were matched by age, preterm pregnancies, and previous abortions; however, there are differences in body mass index (BMI), gestational age, term pregnancies, and the number of living children. Obesity can increase inflammatory and oxidative stress markers in the placental environment (Spradley et al., 2015). Also, it has been reported that the placental immune state shifts with gestational age (Lewis et al., 2018). However, as PE is often accompanied by obesity and early termination of pregnancy, it is difficult to exclude these factors in sample collection. Moreover, limited placental samples in the GDM&PE group are the shortage of this study, for it is hard to collect enough clean samples that exclude interference factors because the number of pregnant women exposed to COVID-19 has increased sharply since December 2022 in China.

In summary, this study demonstrated the statistically distinguished placental immune microenvironment in individuals with PE, but not in those with GDM or GDM&PE. More importantly, macrophages orchestrate a network in the maternal-fetal interface elaborately. These findings provide novel insights into PE pathogenesis and a potential immune target for the clinical prevention and treatment of PE.

Materials and Methods

Clinical sample collection

The samples used in this study were collected from Sir Run Run Shaw Hospital between October 2020 and August 2023. Informed consent was obtained from all volunteers, and the Ethics Committee of Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, approved the study. Human placentas were obtained from women with NP, PE, GDM, or GDM&PE who underwent elective cesarean delivery. The diagnostic criteria for PE included new-onset hypertension after 20 weeks of gestation with SBP ≥140 mmHg and/or diastolic blood pressure ≥90 mmHg and proteinuria (≥300 mg) on at least two occasions. A positive glucose tolerance test diagnoses gestational diabetes mellitus. Women with normal blood pressure, full-term pregnancies, and no complications were designated as controls. The detailed clinical characteristics of the pregnant women in this study are presented in Table 5 and Table 6.

Mice

Eight-week-old female C57 mice and ten-week-old male BALB/c mice were purchased from Hangzhou Ziyuan Laboratory Animal Technology Co., Ltd (Hangzhou, China) and Shanghai Jihui Experimental Animal Breeding Co., Ltd (Shanghai, China), respectively. All animals were maintained under pathogen-free conditions. The Guide for the Care and Use of Laboratory Animals (China) conducted all experimental procedures involving animals, and the

Animal Research Ethics Committee of the Sir Run Run Shaw Hospital of Zhejiang University
approved the protocols.

Female C57 mice were mated with male BALB/c mice to establish an allogeneic
pregnancy model (Rowe et al., 2012). The day of the vaginal plug detection was considered
day 0.5 of pregnancy. SBP was measured using a noninvasive mouse tailcuff BP analyzer
(BP-2010A, Softron, Japan) at 12.5, and 16.5 days of gestation. Three random urine samples
were collected after day 16.5 of gestation, and UACR was measured by urinary microalbumin
(CH0101060, Maccura, China). Mice were euthanized on day 18.5 of gestation.

488 Reduction in uterine perfusion pressure mouse model

489   To construct a mouse model of PE, we ligate uterine arteries in pregnant mice on day 12.5
490 of gestation. Briefly, after 4% chloral hydrate was intraperitoneal injected for anesthesia,
491 bilateral incisions were made on the back of the pregnant mice, and surgical sutures were
492 used to reduce the blood flow of the bilateral uterine arcades.

493 Isolation of single cells from the mouse uterus and human placenta

Uterus from pregnant mice and placentas from volunteers were washed twice with
ice-cold PBS and cut into small pieces. The tissues were digested with collagenase type IV (1
mg/ml, Sigma-Aldrich, USA) and DNase I (0.01 mg/ml, Sigma-Aldrich, U.S.A) in RPMI 1640
medium (Thermo Fisher Scientific) for 40 min at 200 rpm and 37 °C. The suspensions were
strained through 70-µm nylon mesh and centrifuged at 500×g for 5 minutes. Leaving the
supernatants, the cell pellets from human placentas need an extra purification by Ficoll (P4350,
Solarbio, China) according to the manufacturer's instructions. Human CD4$^+$ memory T cells
were isolated using a Human Central and Effector Memory CD4$^+$ T Cell Isolation Kit (17865,
STEMCELL, Canada).

503 Adoptive transferred mouse model

Endogenous macrophages or CD4$^+$ T cells were depleted by injecting PLX3397 (S7818,
Selleck, USA), clodronate liposomes (40337ES08, YEASEN, China) or anti-CD4 antibody
(BE0003-1, BioXcell, USA) intraperitoneally every three days starting from day 10.5 of
gestation.

For the adoptive transferred mouse model, phycoerythrin-conjugated anti-mouse CD4
(12-0041-82, eBioscience, USA) and Pc5.5-conjugated anti-mouse CD44 (45-0441-82,
eBioscience, USA) were used to label CD4$^+$ memory-like T cells from uterus and placentas.
PE-conjugated anti-mouse CD45 (E-AB-F1136D, Elabscience, China), FITC-conjugated
anti-mouse F480 (11-4801-82, eBioscience, USA), and PE-Cyanine7-conjugated anti-mouse
CD206 (141719, Biolegend, USA) were used to label anti-inflam Macs from uterus and
placentas, and the remaining CD45$^+$F480$^+$CD206$^-$ cells were considered pro-inflam Macs. The
MoFlo Astrios EQ FACS Cell Sorter at Core Facilities, ZheJiang University School of Medicine
(China), was used to obtain specific cells with a purity of > 96%. Briefly, 2–5 × 10^5 CD4$^+$ CD44$^+$
T cells, 5×10^5 pro-inflam Macs, or 5×10^5 anti-inflam Macs were injected to pregnant C57 mice
intravenously at day 12.5 of gestation.

2–5 × 10^5 CD45$^+$CD4$^+$ T cells from uterus and placentas of pregnant mice were sorted
and transferred to pregnant C57 mice at day 11.5 of gestation after treating with 100nM
BMS-754809 (HY10200; MedChemExpress; USA) or PBS for three days in vitro.

522 Mass cytometry by time of flight

7 × 10^5 cells from NP (n=9), PE (n=8), GDM (n=8), GDM&PE (n=7) presented in Table 1

were stained for cell surface markers for 30 min at 4°C in staining medium (PBS containing 1%

BSA and 0.02% NaN_3). Cells were washed with protein-free PBS, stained with 2.5 µM cisplatin

for 5 min at 25°C and fixed using transcription factor buffer set (BD Biosciences), followed by

intracellular staining for 60 min at 4 °C. Then the cells were washed and stored at 4 °C until

acquisition. Finally, cells were analyzed on a CyTOF mass spectrometer (Fluidigm, South San

Francisco, CA, USA). The data were normalized for each experiment using EQ four-element

calibration beads (Fluidigm) and the reference EQ passport P13H2302.

RNA sequencing

5×10^5 CD45$^+$F4/80$^+$CD206$^-$ pro-inflam Macs (n=3) and 5×10^5 CD45$^+$F4/80$^+$CD206$^+$

anti-inflam Macs (n=3) were isolated from the uterus and placentas of mice with RUPP

and high throughput sequencing and bioinformatics analyses were conducted at Shenzhen

Huada Gene Technology Service Co. Ltd.(Shenzhen, China). Only differential genes with

more than two-fold change and a corrected P value less than 0.05 were considered statistically

significant.

Single-cell RNA sequencing (scRNA-seq)

Mice were injected with 5×10^5 CD45$^+$F4/80$^+$CD206$^-$ pro-inflam Macs or 5×10^5

CD45$^+$F4/80$^+$CD206$^+$ anti-inflam Macs at 12.5 days of gestation and euthanized on day 18.5 of

gestation. CD45$^+$ immune cells from the uterus and placenta were isolated from mice

transferred pro-inflam Macs and anti-inflam Macs. scRNA-seq was conducted at PLTTECH

Service Co. Ltd. (Hangzhou, China). Differential genes with a corrected P < 0.05 were

considered statistically significant.

Isolation of NP-EVs and PE-EVs

NP-EVs and PE-EVs were isolated and identified using our published protocols (Jiang et

al., 2021; X. Liu et al., 2022). Briefly, after digesting the placental tissues from NP or PE, the

suspensions were filtered through 100-µm nylon mesh and centrifugated at 3,000×g for 15

minutes. Then the supernatants were filtered with a 0.22-µm filter and centrifuged at

100,000×g for 1h at 4 °C. Then the pellets of EVs were resuspended and centrifuged at

100,000×g once again. A BCA assay kit conducted Protein quantitation of EVs (23235,

Thermo Fisher Scientific). To isolate the T-EVs, 500 µg of the EVs were incubated with 1 µg

placental alkaline phosphatase antibody (SC-47691, Santa Cruz Biotechnology) at 4 °C

overnight, then washed in the recommended buffer (PBS containing 2% exosome-free FBS

and 1 mmol/L ethylenediaminetetraacetic acid), and centrifuged at 100 000×g for 1 hour at

4 °C. After resuspended, the total EVs were sorted by EasySep Mouse PE Positive Selection

Kit II (17666, STEMCELL) to collect the final NP-EVs or PE-EVs.

Induction of CD4$^+$ memory-like T cells

Macrophages were obtained after five days of culture of mononuclear cells isolated from

human peripheral blood following previous protocols (X. Liu et al., 2022). NP-EVs or PE-EVs in

a 50 µg/mL concentration were added to macrophages and co-cultured at 37 °C for 8 h. The

human CD4$^+$ naïve T cells Isolation Kit (19555, STEMCELL, Canada) was used to isolate

purified CD4$^+$ naïve T cells from human peripheral blood cells according to the manufacturer's

instructions. Then, CD4$^+$ naïve T cells were co-cultured with EV-treated macrophages for six

days. Flow cytometry was performed to measure the frequency of memory-like Th17 cells. For

animal experiments, CD4$^+$ naïve T cells sorted from the uterus and placentas of pregnant mice

were cultured with the IGF1R inhibitor BMS-754807 at a concentration of 10 µM for three days

before co-cultured with macrophages.

Flow cytometry

Single immune cells from the mice uterus were obtained following the method described
above. After incubation with Cell Stimulation Cocktail (00-4975-93, Invitrogen, USA) for five
hours at 37 °C, cells were collected and surface staining for either phycoerythrin-conjugated
anti-mouse CD4 (12-0041-82, eBioscience, USA), APC-conjugated anti-mouse CD8 (100711,
BioLegend, USA), phycoerythrin-Cy5.5-conjugated anti-mouse CD44 (45-0441-82,
eBioscience, USA), APC-conjugated anti-mouse CD11b (101212, BioLegend, USA),
FITC-conjugated anti-mouse Gr-1 (108406, BioLegend, USA), PE-Cyanine7-conjugated
anti-mouse Ly6G (E-AB-F1108H, elabscience, China) or intracellular staining for
488-conjugated anti-mouse IL-17A (506910, BioLegend, USA) was performed according to the
manufacturer's instructions.

Single placental and peripheral lymphocytes and were obtained following the method
described above. The Human Central and Effector Memory CD4$^+$ T cell Isolation Kit (17865,
STEMCELL, Canada) was used to obtain purified memory CD4$^+$ T cells, according to the
manufacturer's instructions. After incubation with Cell Stimulation Cocktail (00-4975-93,
Invitrogen, USA) for five hours at 37 °C, cells were collected and surface staining for
phycoerythrin-conjugated anti-human CD4 (2384240, eBioscience, USA), FITC-conjugated
anti-human CD45RO (304204, BioLegend, USA), and phycoerythrin-Cy7-conjugated
anti-human CCR7 (353227, BioLegend, USA) or intracellular staining for APC-conjugated
anti-human IL-17A (17-7179-42, eBioscience, USA) and Foxp3 (320014, BioLegend, USA)
was performed according to the manufacturer's instructions.

Immunofluorescence

Frozen sections of the placentas were permeabilized with PBS containing 0.5% Triton
X-100 (PBST) for 20 min and incubated for 1 h with a blocking buffer in PBST. The sections
were then incubated with anti-CD4 (sc-1176, Santa Cruz Biotechnology, China) and
FITC-conjugated anti-human CD45RO (304204, BioLegend, USA) at 4 °C overnight, followed
by Alexa Fluor 568-conjugated secondary antibodies (dilution: 1:200, Yeasen, China) for 1 h.
The slides were counterstained with 4,6-diamidino-2-phenylindole (DAPI, 1 µg/ml; Roche,
Switzerland) for 20 min. Digital images were obtained using confocal fluorescence microscopy
(ZEISS LSM 800, Germany). ImageJ software was used to quantify the fluorescence intensity
from immunofluorescence (IF) images.

Statistical analysis

601   Data were analysed using SPSS version 20. After the Shapiro-wilk test, the data were
602 confirmed to be non-normal distribution. Kruskal-wallis test was used to compare the results of
603 experiments with multiple groups. All data are presented as mean ± SEM (*$P < 0.05$, ** $P <$
604 0.01, *** $P < 0.001$; NS, not significant).

Acknowledgments

606   The authors would like to express their heartfelt gratitude to the participants for their
607 contributions. This work was financially supported by the National Natural Science Foundation
608 of China (82271694), the Zhejiang Province Natural Science Foundation key project fund
609 (LZ24H040002), the Zhejiang Medicine and Health Science and Technology Plan Project
610 (2023KY800).

References

Agbaglo, E., Agbadi, P., Tetteh, J. K., Ameyaw, E. K., Adu, C., & Nutor, J. J. 2022. Trends in total fertility rate in Ghana by different inequality dimensions from 1993 to 2014. *BMC Womens Health* **22**: 49. DOI: https://doi.org/10.1186/s12905-022-01629-w, PMID: 35197037

Amador, C. A., Barrientos, V., Peña, J., Herrada, A. A., González, M., Valdés, S., Carrasco, L., Alzamora, R., Figueroa, F., Kalergis, A. M., & Michea, L. 2014. Spironolactone decreases DOCA-salt-induced organ damage by blocking the activation of T helper 17 and the downregulation of regulatory T lymphocytes. *Hypertension* **63**: 797-803. DOI: https://doi.org/10.1161/hypertensionaha.113.02883, PMID: 24420551

Aneman, I., Pienaar, D., Suvakov, S., Simic, T. P., Garovic, V. D., & McClements, L. 2020. Mechanisms of Key Innate Immune Cells in Early- and Late-Onset Preeclampsia. *Front Immunol* **11**: 1864. DOI: https://doi.org/10.3389/fimmu.2020.01864, PMID: 33013837

Bachmayer, N., Rafik Hamad, R., Liszka, L., Bremme, K., & Sverremark-Ekström, E. 2006. Aberrant uterine natural killer (NK)-cell expression and altered placental and serum levels of the NK-cell promoting cytokine interleukin-12 in pre-eclampsia. *Am J Reprod Immunol* **56**: 292-301. DOI: https://doi.org/10.1111/j.1600-0897.2006.00429.x, PMID: 17076673

Bekkering, S., Arts, R. J. W., Novakovic, B., Kourtzelis, I., van der Heijden, C., Li, Y., Popa, C. D., Ter Horst, R., van Tuijl, J., Netea-Maier, R. T., van de Veerdonk, F. L., Chavakis, T., Joosten, L. A. B., van der Meer, J. W. M., Stunnenberg, H., Riksen, N. P., & Netea, M. G. 2018. Metabolic Induction of Trained Immunity through the Mevalonate Pathway. *Cell* **172**: 135-146.e139. DOI: https://doi.org/10.1016/j.cell.2017.11.025, PMID: 29328908

Bernardes, T. P., Mol, B. W., Ravelli, A. C. J., van den Berg, P. P., Boezen, H. M., & Groen, H. 2019. Recurrence risk of preeclampsia in a linked population-based cohort: Effects of first pregnancy maximum diastolic blood pressure and gestational age. *Pregnancy Hypertens* **15**: 32-36. DOI: https://doi.org/10.1016/j.preghy.2018.10.010, PMID: 30825924

Burton, G. J., Redman, C. W., Roberts, J. M., & Moffett, A. 2019. Pre-eclampsia: pathophysiology and clinical implications. *Bmj* **366**: l2381. DOI: https://doi.org/10.1136/bmj.l2381, PMID: 31307997

Care, A. S., Bourque, S. L., Morton, J. S., Hjartarson, E. P., Robertson, S. A., & Davidge, S. T. 2018. Reduction in Regulatory T Cells in Early Pregnancy Causes Uterine Artery Dysfunction in Mice. *Hypertension* **72**: 177-187. DOI: https://doi.org/10.1161/hypertensionaha.118.10858, PMID: 29785960

Chappell, L. C., Cluver, C. A., Kingdom, J., & Tong, S. 2021. Pre-eclampsia. *Lancet* **398**: 341-354. DOI: https://doi.org/10.1016/s0140-6736(20)32335-7, PMID: 34051884

Chatterjee, P., Chiasson, V. L., Seerangan, G., De Guzman, E., Milad, M., Bounds, K. R., Gasheva, O., Tobin, R. P., Hatahet, M., Kopriva, S., Jones, K. A., Newell-Rogers, M. K., & Mitchell, B. M. 2017. Depletion of MHC class II invariant chain peptide or γ-δ T-cells ameliorates experimental preeclampsia. *Clin Sci (Lond)* **131**: 2047-2058. DOI: https://doi.org/10.1042/cs20171008, PMID: 28642294

Chen, B., Ye, B., Li, M., Wang, S., Li, J., Lai, Y., Yang, N., Ke, Z., & Zhang, H. 2022. TIGIT Deficiency Protects Mice From DSS-Induced Colitis by Regulating IL-17A-Producing CD4(+) Tissue-Resident Memory T Cells. *Front Immunol* **13**: 931761. DOI: https://doi.org/10.3389/fimmu.2022.931761, PMID: 35844584

Cornelius, D. C., Amaral, L. M., Harmon, A., Wallace, K., Thomas, A. J., Campbell, N., Scott, J., Herse, F., Haase, N., Moseley, J., Wallukat, G., Dechend, R., & LaMarca, B. 2015a. An increased population of regulatory T cells improves the pathophysiology of placental ischemia in a rat model of preeclampsia. *Am J Physiol Regul Integr Comp Physiol* **309**: R884-891. DOI: https://doi.org/10.1152/ajpregu.00154.2015, PMID: 26290102

Cornelius, D. C., Castillo, J., Porter, J., Amaral, L. M., Campbell, N., Paige, A., Thomas, A. J., Harmon, A., Cunningham, M. W., Jr., Wallace, K., Herse, F., Wallukat, G., Dechend, R., & LaMarca, B. 2015b. Blockade of CD40 ligand for intercellular communication reduces hypertension, placental oxidative stress, and AT1-AA in response to adoptive transfer of CD4+ T lymphocytes from RUPP rats. *Am J Physiol Regul Integr Comp Physiol* **309**: R1243-1250. DOI: https://doi.org/10.1152/ajpregu.00273.2015, PMID: 26310940

Corrêa-Silva, S., Alencar, A. P., Moreli, J. B., Borbely, A. U., de, S. L. L., Scavone, C., Damasceno, D. C., Rudge, M. V. C., Bevilacqua, E., & Calderon, I. M. P. 2018. Hyperglycemia induces inflammatory mediators in the human chorionic villous. *Cytokine* **111**: 41-48. DOI: https://doi.org/10.1016/j.cyto.2018.07.020, PMID: 30114628

Deer, E., Herrock, O., Campbell, N., Cornelius, D., Fitzgerald, S., Amaral, L. M., & LaMarca, B. 2023. The role of immune cells and mediators in preeclampsia. *Nat Rev Nephrol* **19**: 257-270. DOI: https://doi.org/10.1038/s41581-022-00670-0, PMID: 36635411

Deer, E., Reeve, K. E., Amaral, L., Vaka, V. R., Franks, M., Campbell, N., Fitzgerald, S., Herrock, O., Ibrahim, T., Cornelius, D., & LaMarca, B. 2021. CD4+ T cells cause renal and placental mitochondrial oxidative stress as mechanisms of hypertension in response to placental ischemia. *Am J Physiol Renal Physiol* **320**: F47-f54. DOI: https://doi.org/10.1152/ajprenal.00398.2020, PMID: 33196321

Díaz, E., Cárdenas, M., Ariza, A. C., Larrea, F., & Halhali, A. 2005. Placental insulin and insulin-like growth factor I receptors in normal and preeclamptic pregnancies. *Clin Biochem* **38**: 243-247. DOI: https://doi.org/10.1016/j.clinbiochem.2004.10.013, PMID: 15708546

DiToro, D., Harbour, S. N., Bando, J. K., Benavides, G., Witte, S., Laufer, V. A., Moseley, C., Singer, J. R., Frey, B., Turner, H., Bruning, J., Darley-Usmar, V., Gao, M., Conover, C., Hatton, R. D., Frank, S., Colonna, M., & Weaver, C. T. 2020. Insulin-Like Growth Factors Are Key Regulators of T Helper 17 Regulatory T Cell Balance in Autoimmunity. *Immunity* **52**: 650-667.e610. DOI: https://doi.org/10.1016/j.immuni.2020.03.013, PMID: 32294406

Eghbal-Fard, S., Yousefi, M., Heydarlou, H., Ahmadi, M., Taghavi, S., Movasaghpour, A., Jadidi-Niaragh, F., Yousefi, B., Dolati, S., Hojjat-Farsangi, M., Rikhtegar, R., Nouri, M., & Aghebati-Maleki, L. 2019. The imbalance of Th17/Treg axis involved in the pathogenesis of preeclampsia. *J Cell Physiol* **234**: 5106-5116. DOI: https://doi.org/10.1002/jcp.27315, PMID: 30277561

Faas, M. M., Spaans, F., & De Vos, P. 2014. Monocytes and macrophages in pregnancy and pre-eclampsia. *Front Immunol* **5**: 298. DOI: https://doi.org/10.3389/fimmu.2014.00298, PMID: 25071761

Fanelli, G., Romano, M., Nova-Lamperti, E., Werner Sunderland, M., Nerviani, A., Scottà, C., Bombardieri, M., Quezada, S. A., Sacks, S. H., Noelle, R. J., Pitzalis, C., Lechler, R. I., Lombardi, G., & Becker, P. D. 2021. PD-L1 signaling on human memory CD4+ T cells induces a regulatory phenotype. *PLoS Biol* **19**: e3001199. DOI:

https://doi.org/10.1371/journal.pbio.3001199, PMID: 33901179

Fu, B., Li, X., Sun, R., Tong, X., Ling, B., Tian, Z., & Wei, H. 2013. Natural killer cells promote
immune tolerance by regulating inflammatory TH17 cells at the human maternal-fetal
interface. *Proc Natl Acad Sci U S A* **110**: E231-240. DOI:
https://doi.org/10.1073/pnas.1206322110, PMID: 23271808

Fu, B., Tian, Z., & Wei, H. 2014. TH17 cells in human recurrent pregnancy loss and
pre-eclampsia. *Cell Mol Immunol* **11**: 564-570. DOI: https://doi.org/10.1038/cmi.2014.54,
PMID: 25027967

Fukui, A., Funamizu, A., Yokota, M., Yamada, K., Nakamua, R., Fukuhara, R., Kimura, H., &
Mizunuma, H. 2011. Uterine and circulating natural killer cells and their roles in women
with recurrent pregnancy loss, implantation failure and preeclampsia. *J Reprod Immunol*
90: 105-110. DOI: https://doi.org/10.1016/j.jri.2011.04.006, PMID: 21632120

Grotegut, C. A. 2016. Prevention of preeclampsia. *J Clin Invest* **126**: 4396-4398. DOI:
https://doi.org/10.1172/jci91300, PMID: 27841760

Han, X., Li, W., Li, P., Zheng, Z., Lin, B., Zhou, B., Guo, K., He, P., & Yang, J. 2021. Stimulation
of α7 Nicotinic Acetylcholine Receptor by Nicotine Suppresses Decidual M1 Macrophage
Polarization Against Inflammation in Lipopolysaccharide-Induced Preeclampsia-Like
Mouse Model. *Front Immunol* **12**: 642071. DOI:
https://doi.org/10.3389/fimmu.2021.642071, PMID: 33995360

Jensen, F., Wallukat, G., Herse, F., Budner, O., El-Mousleh, T., Costa, S. D., Dechend, R., &
Zenclussen, A. C. 2012. CD19+CD5+ cells as indicators of preeclampsia. *Hypertension*
59: 861-868. DOI: https://doi.org/10.1161/hypertensionaha.111.188276, PMID: 22353610

Jiang, L., Fei, H., Jin, X., Liu, X., Yang, C., Li, C., Chen, J., Yang, A., Zhu, J., Wang, H., Fei, X.,
& Zhang, S. 2021. Extracellular Vesicle-Mediated Secretion of HLA-E by Trophoblasts
Maintains Pregnancy by Regulating the Metabolism of Decidual NK Cells. *Int J Biol Sci*
17: 4377-4395. DOI: https://doi.org/10.7150/ijbs.63390, PMID: 34803505

Jung, E., Romero, R., Yeo, L., Gomez-Lopez, N., Chaemsaithong, P., Jaovisidha, A., Gotsch,
F., & Erez, O. 2022. The etiology of preeclampsia. *Am J Obstet Gynecol* **226**: S844-s866.
DOI: https://doi.org/10.1016/j.ajog.2021.11.1356, PMID: 35177222

Kieffer, T. E. C., Laskewitz, A., Scherjon, S. A., Faas, M. M., & Prins, J. R. 2019. Memory T
Cells in Pregnancy. *Front Immunol* **10**: 625. DOI:
https://doi.org/10.3389/fimmu.2019.00625, PMID: 31001255

Köstlin-Gille, N., Dietz, S., Schwarz, J., Spring, B., Pauluschke-Fröhlich, J., Poets, C. F., &
Gille, C. 2019. HIF-1α-Deficiency in Myeloid Cells Leads to a Disturbed Accumulation of
Myeloid Derived Suppressor Cells (MDSC) During Pregnancy and to an Increased
Abortion Rate in Mice. *Front Immunol* **10**: 161. DOI:
https://doi.org/10.3389/fimmu.2019.00161, PMID: 30804946

Lager, S., Sovio, U., Eddershaw, E., van der Linden, M. W., Yazar, C., Cook, E., Happerfield, L.,
Jessop, F. A., Sebire, N. J., Charnock-Jones, D. S., & Smith, G. C. S. 2020. Abnormal
placental CD8(+) T-cell infiltration is a feature of fetal growth restriction and pre-eclampsia.
J Physiol **598**: 5555-5571. DOI: https://doi.org/10.1113/jp279532, PMID: 32886802

LaMarca, B., Wallace, K., Herse, F., Wallukat, G., Martin, J. N., Jr., Weimer, A., & Dechend, R.
2011. Hypertension in response to placental ischemia during pregnancy: role of B
lymphocytes. *Hypertension* **57**: 865-871. DOI:

744   https://doi.org/10.1161/hypertensionaha.110.167569, PMID: 21357287

745 Lampé, R., Kövér, Á., Szűcs, S., Pál, L., Árnyas, E., Ádány, R., & Póka, R. 2015. Phagocytic
746   index of neutrophil granulocytes and monocytes in healthy and preeclamptic pregnancy. *J*
747   *Reprod Immunol* **107**: 26-30. DOI: https://doi.org/10.1016/j.jri.2014.11.001, PMID:
748   25534923

749 Lampé, R., Szucs, S., Adány, R., & Póka, R. 2011. Granulocyte superoxide anion production
750   and regulation by plasma factors in normal and preeclamptic pregnancy. *J Reprod*
751   *Immunol* **89**: 199-206. DOI: https://doi.org/10.1016/j.jri.2011.01.019, PMID: 21529965

752 Lang, X., Liu, W., Hou, Y., Zhao, W., Yang, X., Chen, L., Yan, Q., & Cheng, W. 2021. IL-17A
753   polymorphism (rs2275913) and levels are associated with preeclampsia pathogenesis in
754   Chinese patients. *BMC Med Genomics* **14**: 5. DOI:
755   https://doi.org/10.1186/s12920-020-00840-8, PMID: 33407460

756 Lewis, E. L., Sierra, L. J., Barila, G. O., Brown, A. G., Porrett, P. M., & Elovitz, M. A. 2018.
757   Placental immune state shifts with gestational age. *Am J Reprod Immunol* **79**: e12848.
758   DOI: https://doi.org/10.1111/aji.12848, PMID: 29577513

759 Li, M., Sun, F., Xu, Y., Chen, L., Chen, C., Cui, L., Qian, J., Li, D., Wang, S., & Du, M. 2022.
760   Tim-3(+) decidual Mφs induced Th2 and Treg bias in decidual CD4(+)T cells and
761   promoted pregnancy maintenance via CD132. *Cell Death Dis* **13**: 454. DOI:
762   https://doi.org/10.1038/s41419-022-04899-2, PMID: 35550500

763 Liao, G., Cheng, D., Li, J., & Hu, S. 2021. Clinical significance of microRNA-320a and
764   insulin-like growth factor-1 receptor in early-onset preeclampsia patients. *Eur J Obstet*
765   *Gynecol Reprod Biol* **263**: 164-170. DOI: https://doi.org/10.1016/j.ejogrb.2021.06.032,
766   PMID: 34218203

767 Lisonkova, S., & Joseph, K. S. 2013. Incidence of preeclampsia: risk factors and outcomes
768   associated with early- versus late-onset disease. *Am J Obstet Gynecol* **209**:
769   544.e541-544.e512. DOI: https://doi.org/10.1016/j.ajog.2013.08.019, PMID: 23973398

770 Liu, D., Li, Q., Ding, H., Zhao, G., Wang, Z., Cao, C., Dai, Y., Zheng, M., Zhu, X., Wu, Q., Wang,
771   Y., Duan, H., Tang, H., Lu, X., Hou, Y., & Hu, Y. 2021. Placenta-derived IL-32β activates
772   neutrophils to promote preeclampsia development. *Cell Mol Immunol* **18**: 979-991. DOI:
773   https://doi.org/10.1038/s41423-021-00636-5, PMID: 33707686

774 Liu, X., Fei, H., Yang, C., Wang, J., Zhu, X., Yang, A., Shi, Z., Jin, X., Yang, F., Wu, D., Jiang, L.,
775   & Zhang, S. 2022. Trophoblast-Derived Extracellular Vesicles Promote Preeclampsia by
776   Regulating Macrophage Polarization. *Hypertension* **79**: 2274-2287. DOI:
777   https://doi.org/10.1161/hypertensionaha.122.19244, PMID: 35993233

778 Lu, D., Peng, Q., Chen, D., Chen, X., & Jiang, M. 2020. Expression imbalance of IL-17/IL-35 in
779   peripheral blood and placental tissue of pregnant women in preeclampsia. *Taiwan J*
780   *Obstet Gynecol* **59**: 409-414. DOI: https://doi.org/10.1016/j.tjog.2020.03.013, PMID:
781   32416889

782 Madhur, M. S., Elijovich, F., Alexander, M. R., Pitzer, A., Ishimwe, J., Van Beusecum, J. P.,
783   Patrick, D. M., Smart, C. D., Kleyman, T. R., Kingery, J., Peck, R. N., Laffer, C. L., & Kirabo,
784   A. 2021. Hypertension: Do Inflammation and Immunity Hold the Key to Solving this
785   Epidemic? *Circ Res* **128**: 908-933. DOI: https://doi.org/10.1161/circresaha.121.318052,
786   PMID: 33793336

787 McElwain, C. J., McCarthy, F. P., & McCarthy, C. M. 2021. Gestational Diabetes Mellitus and

Maternal Immune Dysregulation: What We Know So Far. *Int J Mol Sci* **22**. DOI:
https://doi.org/10.3390/ijms22084261, PMID: 33923959

McIntyre, H. D., Catalano, P., Zhang, C., Desoye, G., Mathiesen, E. R., & Damm, P. 2019.
Gestational diabetes mellitus. *Nat Rev Dis Primers* **5**: 47. DOI:
https://doi.org/10.1038/s41572-019-0098-8, PMID: 31296866

Miller, D., Garcia-Flores, V., Romero, R., Galaz, J., Pique-Regi, R., & Gomez-Lopez, N. 2022.
Single-Cell Immunobiology of the Maternal-Fetal Interface. *J Immunol* **209**: 1450-1464.
DOI: https://doi.org/10.4049/jimmunol.2200433, PMID: 36192116

Morita, K., Tsuda, S., Kobayashi, E., Hamana, H., Tsuda, K., Shima, T., Nakashima, A.,
Ushijima, A., Kishi, H., & Saito, S. 2020. Analysis of TCR Repertoire and PD-1 Expression
in Decidual and Peripheral CD8(+) T Cells Reveals Distinct Immune Mechanisms in
Miscarriage and Preeclampsia. *Front Immunol* **11**: 1082. DOI:
https://doi.org/10.3389/fimmu.2020.01082, PMID: 32582176

Nerenberg, K. A., Johnson, J. A., Leung, B., Savu, A., Ryan, E. A., Chik, C. L., & Kaul, P. 2013.
Risks of gestational diabetes and preeclampsia over the last decade in a cohort of Alberta
women. *J Obstet Gynaecol Can* **35**: 986-994. DOI:
https://doi.org/10.1016/s1701-2163(15)30786-6, PMID: 24246398

Novotny, S. R., Wallace, K., Heath, J., Moseley, J., Dhillon, P., Weimer, A., Wallukat, G., Herse,
F., Wenzel, K., Martin, J. N., Jr., Dechend, R., & Lamarca, B. 2012. Activating
autoantibodies to the angiotensin II type I receptor play an important role in mediating
hypertension in response to adoptive transfer of CD4+ T lymphocytes from placental
ischemic rats. *Am J Physiol Regul Integr Comp Physiol* **302**: R1197-1201. DOI:
https://doi.org/10.1152/ajpregu.00623.2011, PMID: 22461177

Ostlund, I., Haglund, B., & Hanson, U. 2004. Gestational diabetes and preeclampsia. *Eur J
Obstet Gynecol Reprod Biol* **113**: 12-16. DOI:
https://doi.org/10.1016/j.ejogrb.2003.07.001, PMID: 15036703

Rasmussen, T. A., Zerbato, J. M., Rhodes, A., Tumpach, C., Dantanarayana, A., McMahon, J.
H., Lau, J. S. Y., Chang, J. J., Gubser, C., Brown, W., Hoh, R., Krone, M., Pascoe, R.,
Chiu, C. Y., Bramhall, M., Lee, H. J., Haque, A., Fromentin, R., Chomont, N., Milush, J., et
al. 2022. Memory CD4(+) T cells that co-express PD1 and CTLA4 have reduced response
to activating stimuli facilitating HIV latency. *Cell Rep Med* **3**: 100766. DOI:
https://doi.org/10.1016/j.xcrm.2022.100766, PMID: 36198308

Robajac, D., Masnikosa, R., Miković, Ž., Mandić, V., & Nedić, O. 2015. Oxidation of placental
insulin and insulin-like growth factor receptors in mothers with diabetes mellitus or
preeclampsia complicated with intrauterine growth restriction. *Free Radic Res* **49**:
984-989. DOI: https://doi.org/10.3109/10715762.2015.1020798, PMID: 25812587

Rowe, J. H., Ertelt, J. M., Xin, L., & Way, S. S. 2012. Pregnancy imprints regulatory memory
that sustains anergy to fetal antigen. *Nature* **490**: 102-106. DOI:
https://doi.org/10.1038/nature11462, PMID: 23023128

Saito, S., Shiozaki, A., Nakashima, A., Sakai, M., & Sasaki, Y. 2007. The role of the immune
system in preeclampsia. *Mol Aspects Med* **28**: 192-209. DOI:
https://doi.org/10.1016/j.mam.2007.02.006, PMID: 17433431

Santner-Nanan, B., Peek, M. J., Khanam, R., Richarts, L., Zhu, E., Fazekas de St Groth, B., &
Nanan, R. 2009. Systemic increase in the ratio between Foxp3+ and IL-17-producing

CD4+ T cells in healthy pregnancy but not in preeclampsia. *J Immunol* **183**: 7023-7030. DOI: https://doi.org/10.4049/jimmunol.0901154, PMID: 19915051

Sasaki, Y., Darmochwal-Kolarz, D., Suzuki, D., Sakai, M., Ito, M., Shima, T., Shiozaki, A., Rolinski, J., & Saito, S. 2007. Proportion of peripheral blood and decidual CD4(+) CD25(bright) regulatory T cells in pre-eclampsia. *Clin Exp Immunol* **149**: 139-145. DOI: https://doi.org/10.1111/j.1365-2249.2007.03397.x, PMID: 17459078

Schneider, S., Freerksen, N., Röhrig, S., Hoeft, B., & Maul, H. 2012. Gestational diabetes and preeclampsia--similar risk factor profiles? *Early Hum Dev* **88**: 179-184. DOI: https://doi.org/10.1016/j.earlhumdev.2011.08.004, PMID: 21890288

Schonkeren, D., van der Hoorn, M. L., Khedoe, P., Swings, G., van Beelen, E., Claas, F., van Kooten, C., de Heer, E., & Scherjon, S. 2011. Differential distribution and phenotype of decidual macrophages in preeclamptic versus control pregnancies. *Am J Pathol* **178**: 709-717. DOI: https://doi.org/10.1016/j.ajpath.2010.10.011, PMID: 21281803

Shields, C. A., McCalmon, M., Ibrahim, T., White, D. L., Williams, J. M., LaMarca, B., & Cornelius, D. C. 2018. Placental ischemia-stimulated T-helper 17 cells induce preeclampsia-associated cytolytic natural killer cells during pregnancy. *Am J Physiol Regul Integr Comp Physiol* **315**: R336-r343. DOI: https://doi.org/10.1152/ajpregu.00061.2018, PMID: 29718698

Spradley, F. T., Palei, A. C., & Granger, J. P. 2015. Immune Mechanisms Linking Obesity and Preeclampsia. *Biomolecules* **5**: 3142-3176. DOI: https://doi.org/10.3390/biom5043142, PMID: 26569331

Travis, O. K., White, D., Baik, C., Giachelli, C., Thompson, W., Stubbs, C., Greer, M., Lemon, J. P., Williams, J. M., & Cornelius, D. C. 2020. Interleukin-17 signaling mediates cytolytic natural killer cell activation in response to placental ischemia. *Am J Physiol Regul Integr Comp Physiol* **318**: R1036-r1046. DOI: https://doi.org/10.1152/ajpregu.00285.2019, PMID: 32320265

Wallace, K., Novotny, S., Heath, J., Moseley, J., Martin, J. N., Jr., Owens, M. Y., & LaMarca, B. 2012. Hypertension in response to CD4(+) T cells from reduced uterine perfusion pregnant rats is associated with activation of the endothelin-1 system. *Am J Physiol Regul Integr Comp Physiol* **303**: R144-149. DOI: https://doi.org/10.1152/ajpregu.00049.2012, PMID: 22647295

Wang, S., Zhu, X., Xu, Y., Zhang, D., Li, Y., Tao, Y., Piao, H., Li, D., & Du, M. 2016. Programmed cell death-1 (PD-1) and T-cell immunoglobulin mucin-3 (Tim-3) regulate CD4+ T cells to induce Type 2 helper T cell (Th2) bias at the maternal-fetal interface. *Hum Reprod* **31**: 700-711. DOI: https://doi.org/10.1093/humrep/dew019, PMID: 26908841

Weissgerber, T. L., & Mudd, L. M. 2015. Preeclampsia and diabetes. *Curr Diab Rep* **15**: 9. DOI: https://doi.org/10.1007/s11892-015-0579-4, PMID: 25644816

Yao, Y., Xu, X. H., & Jin, L. 2019. Macrophage Polarization in Physiological and Pathological Pregnancy. *Front Immunol* **10**: 792. DOI: https://doi.org/10.3389/fimmu.2019.00792, PMID: 31037072

Zhong, X., Gao, W., Degauque, N., Bai, C., Lu, Y., Kenny, J., Oukka, M., Strom, T. B., & Rothstein, T. L. 2007. Reciprocal generation of Th1/Th17 and T(reg) cells by B1 and B2 B cells. *Eur J Immunol* **37**: 2400-2404. DOI: https://doi.org/10.1002/eji.200737296, PMID: 17683116

Zhou, J., Nefedova, Y., Lei, A., & Gabrilovich, D. 2018. Neutrophils and PMN-MDSC: Their
biological role and interaction with stromal cells. *Semin Immunol* **35**: 19-28. DOI:
https://doi.org/10.1016/j.smim.2017.12.004, PMID: 29254756

Chen,X. , Maria ,F., Melissa,M., Jasmin,H., Igor,S., Ekaterina,A., Chanung,W., Xin,B.,
Mary.B.F., Hu,H., Irina,S., Kim,M.W. , Carla, M. , Jonathan,K., Maxim,N., Jason, D.,
David,M. 2023. Microglia-mediated T cell infiltration drives neurodegeneration in
tauopathy.*Nature*.615(7953):668-677.doi:10.1038/s41586-023-05788-0,PMID: 36890231.

Chen,J., Yang,J., Wang, W., Guo, D., Zhang, C., Wang,S., Lu, X., Huang,X., Wang,P.,
Zhang,G., Zhang,J., Wang,J., Cai,Z. 2022. Tumor extracellular vesicles mediate
anti-PD-L1 therapy resistance by decoying anti-PD-L1. Cell Mol Immunol.
19(11):1290-1301. doi: 10.1038/s41423-022-00926-6, PMID: 36220994.

Nalio Ramos R, Missolo-Koussou Y, Gerber-Ferder Y, Bromley CP, Bugatti M, Núñez NG,
Tosello Boari J, Richer W, Menger L, Denizeau J, Sedlik C, Caudana P, Kotsias F, Niborski
LL, Viel S, Bohec M, Lameiras S, Baulande S, Lesage L, Nicolas A, Meseure D,
Vincent-Salomon A, Reyal F, Dutertre CA, Ginhoux F, Vimeux L, Donnadieu E, Buttard B,
Galon J, Zelenay S, Vermi W, Guermonprez P, Piaggio E, Helft J. 2022. Tissue-resident
FOLR2+ macrophages associate with CD8+ T cell infiltration in human breast cancer. Cell.
2022 Mar 31;185(7):1189-1207.e25. doi: 10.1016/j.cell.2022.02.021, PMID: 35325594.

Thomas JR, Appios A, Zhao X, Dutkiewicz R, Donde M, Lee CYC, Naidu P, Lee C, Cerveira J,
Liu B, Ginhoux F, Burton G, Hamilton RS, Moffett A, Sharkey A, McGovern N. 2021.
Phenotypic and functional characterization of first-trimester human placental
macrophages, Hofbauer cells. J Exp Med. 2021 Jan 4;218(1):e20200891. doi:
10.1084/jem.20200891.PMID: 33075123.

Che LH, Liu JW, Huo JP, Luo R, Xu RM, He C, Li YQ, Zhou AJ, Huang P, Chen YY, Ni W, Zhou
YX, Liu YY, Li HY, Zhou R, Mo H, Li JM. A single-cell atlas of liver metastases of colorectal
cancer reveals reprogramming of the tumor microenvironment in response to preoperative
chemotherapy.Cell Discov. 2021 Sep 7;7(1):80. doi: 10.1038/s41421-021-00312-y. PMID:
34489408.

Desrumaux C, Lemaire-Ewing S, Ogier N, Yessoufou A, Hammann A, Sequeira-Le Grand A,
Deckert V, Pais de Barros JP, Le Guern N, Guy J, Khan NA, Lagrost L. Plasma
phospholipid transfer protein (PLTP) modulates adaptive immune functions through
alternation of T helper cell polarization. Cell Mol Immunol. 2016 Nov;13(6):795-804. doi:
10.1038/cmi.2015.75. Epub 2015 Aug 31. PMID: 26320740.

He J, Song Y, Li G, Xiao P, Liu Y, Xue Y, Cao Q, Tu X, Pan T, Jiang Z, Cao X, Lai L, Wang Q.
Fbxw7 increases CCL2/7 in CX3CR1hi macrophages to promote intestinal inflammation.
J Clin Invest. 2019 Jun 27;129(9):3877-3893. doi: 10.1172/JCI123374.PMID: 31246581.

Wu Y, Wu C, Shi T, Cai Q, Wang T, Xiong Y, Zhang Y, Jiang W, Lu M, Chen Z, Chen J, Wang J,
He R. FAP expression in adipose tissue macrophages promotes obesity and metabolic
inflammation. Proc Natl Acad Sci U S A. 2023 Dec 19;120(51):e2303075120. doi:
10.1073/pnas.2303075120. Epub 2023 Dec 15. PMID: 38100414.

Jin G, Guo N, Liu Y, Zhang L, Chen L, Dong T, Liu W, Zhang X, Jiang Y, Lv G, Zhao F, Liu W,
Hei Z, Yang Y, Ou J. 5-aminolevulinate and CHIL3/CHI3L1 treatment amid ischemia aids
liver metabolism and reduces ischemia-reperfusion injury. Theranostics. 2023 Aug
28;13(14):4802-4820. doi: 10.7150/thno.83163. eCollection 2023. PMID: 37771779.

Nugteren S, Samsom JN. Secretory Leukocyte Protease Inhibitor (SLPI) in mucosal tissues:
Protects against inflammation, but promotes cancer. Cytokine Growth Factor Rev. 2021
Jun;59:22-35. doi: 10.1016/j.cytogfr.2021.01.005. Epub 2021 Feb 10. PMID: 33602652.
Li R, Ferdinand JR, Loudon KW, Bowyer GS, Laidlaw S, Muyas F, Mamanova L, Neves JB,
Bolt L, Fasouli ES, Lawson ARJ, Young MD, Hooks Y, Oliver TRW, Butler TM, Armitage
JN, Aho T, Riddick ACP, Gnanapragasam V, Welsh SJ, Meyer KB, Warren AY, Tran MGB,
Stewart GD, Cortés-Ciriano I, Behjati S, Clatworthy MR, Campbell PJ, Teichmann SA,
Mitchell TJ. Mapping single-cell transcriptomes in the intra-tumoral and associated
territories of kidney cancer. Cancer Cell. 2022 Dec 12;40(12):1583-1599.e10. doi:
10.1016/j.ccell.2022.11.001. Epub 2022 Nov 23.PMID: 36423636.
Che LH, Liu JW, Huo JP, Luo R, Xu RM, He C, Li YQ, Zhou AJ, Huang P, Chen YY, Ni W,
Zhou YX, Liu YY, Li HY, Zhou R, Mo H, Li JM. A single-cell atlas of liver metastases of
colorectal cancer reveals reprogramming of the tumor microenvironment in response to
preoperative chemotherapy. Cell Discov. 2021 Sep 7;7(1):80. doi:
10.1038/s41421-021-00312-y.PMID: 34489408.
Coppo L, Mishra P, Siefert N, Holmgren A, Pääbo S, Zeberg H. A substitution in the
glutathione reductase lowers electron leakage and inflammation in modern humans. Sci
Adv. 2022 Jan 7;8(1):eabm1148. doi: 10.1126/sciadv.abm1148. Epub 2022 Jan 5. PMID:
34985944.
Hang S, Paik D, Yao L, Kim E, Trinath J, Lu J, Ha S, Nelson BN, Kelly SP, Wu L, Zheng Y,
Longman RS, Rastinejad F, Devlin AS, Krout MR, Fischbach MA, Littman DR, Huh JR.
Bile acid metabolites control TH17 and Treg cell differentiation.Nature. 2019
Dec;576(7785):143-148. doi: 10.1038/s41586-019-1785-z. Epub 2019 Nov 27. PMID:
31776512.
Leite JA, Menezes L, Martins E, Rodrigues TS, Tavares L, Ebering A, Schelmbauer C,
Martelossi Cebinelli GC, Zinina V, Golden A, Soshnikova N, Zamboni DS, Cunha FQ,
Huber M, Silva JS, Waisman A, Carlos D, Saraiva Câmara NO. AIM2 promotes TH17
cells differentiation by regulating RORγt transcription activity. iScience. 2023 Oct
4;26(11):108134. doi: 10.1016/j.isci.2023.108134. eCollection 2023 Nov 17. PMID:
37867943.
Evrard M, Becht E, Fonseca R, Obers A, Park SL, Ghabdan-Zanluqui N, Schroeder J, Christo
SN, Schienstock D, Lai J, Burn TN, Clatch A, House IG, Beavis P, Kallies A, Ginhoux F,
Mueller SN, Gottardo R, Newell EW, Mackay LK. Single-cell protein expression profiling
resolves circulating and resident memory T cell diversity across tissues and infection
contexts. Immunity. 2023 Jul 11;56(7):1664-1680.e9. doi: 10.1016/j.immuni.2023.06.005.
Epub 2023 Jun 30. PMID: 37392736.
Bieberich F, Vazquez-Lombardi R, Yermanos A, Ehling RA, Mason DM, Wagner B,
Kapetanovic E, Di Roberto RB, Weber CR, Savic M, Rudolf F, Reddy ST. A Single-Cell
Atlas of Lymphocyte Adaptive Immune Repertoires and Transcriptomes Reveals
Age-Related Differences in Convalescent COVID-19 Patients. Front Immunol. 2021 Jul
12;12:701085. doi: 10.3389/fimmu.2021.701085. eCollection 2021. PMID: 34322127.
Qiu C, Du G. Loss of LEF-1 expression as a diagnostic indicator for extranodal NK/T-cell
lymphoma: An immunohistochemical study of 88 cases. Eur J Haematol. 2022
Nov;109(5):513-518. doi: 10.1111/ejh.13836. Epub 2022 Aug 2.PMID: 35871391.

Sekine T, Perez-Potti A, Nguyen S, Gorin JB, Wu VH, Gostick E, Llewellyn-Lacey S, Hammer
Q, Falck-Jones S, Vangeti S, Yu M, Smed-Sörensen A, Gaballa A, Uhlin M, Sandberg JK,
Brander C, Nowak P, Goepfert PA, Price DA, Betts MR, Buggert M. TOX is expressed by
exhausted and polyfunctional human effector memory CD8+ T cells. Sci Immunol. 2020
Jul 3;5(49):eaba7918. doi: 10.1126/sciimmunol.aba7918. PMID: 32620560.
Esensten JH, Helou YA, Chopra G, Weiss A, Bluestone JA. CD28 Costimulation: From
Mechanism to Therapy. Immunity. 2016 May 17;44(5):973-88.
doi:10.1016/j.immuni.2016.04.020. PMID: 27192564.
Pai JA, Hellmann MD, Sauter JL, Mattar M, Rizvi H, Woo HJ, Shah N, Nguyen EM, Uddin FZ,
Quintanal-Villalonga A, Chan JM, Manoj P, Allaj V, Baine MK, Bhanot UK, Jain M, Linkov I,
Meng F, Brown D, Chaft JE, Plodkowski AJ, Gigoux M, Won HH, Sen T, Wells DK,
Donoghue MTA, de Stanchina E, Wolchok JD, Loomis B, Merghoub T, Rudin CM, Chow
A, Satpathy AT. Lineage tracing reveals clonal progenitors and long-term persistence of
tumor-specific T cells during immune checkpoint blockade. Cancer Cell. 2023 Apr
10;41(4):776-790.e7. doi: 10.1016/j.ccell.2023.03.009. Epub 2023 Mar 30. PMID:
37001526.
Montaldo E, Lusito E, Bianchessi V, Caronni N, Scala S, Basso-Ricci L, Cantaffa C,
Masserdotti A, Barilaro M, Barresi S, Genua M, Vittoria FM, Barbiera G, Lazarevic D,
Messina C, Xue E, Marktel S, Tresoldi C, Milani R, Ronchi P, Gattillo S, Santoleri L, Di
Micco R, Ditadi A, Belfiori G, Aleotti F, Naldini MM, Gentner B, Gardiman E, Tamassia N,
Cassatella MA, Hidalgo A, Kwok I, Ng LG, Crippa S, Falconi M, Pettinella F, Scapini P,
Naldini L, Ciceri F, Aiuti A, Ostuni R. Cellular and transcriptional dynamics of human
neutrophils at steady state and upon stress. Nat Immunol. 2022 Oct;23(10):1470-1483.
doi: 10.1038/s41590-022-01311-1. Epub 2022 Sep 22.PMID: 36138183.
Wenzel P, Rossmann H, Müller C, Kossmann S, Oelze M, Schulz A, Arnold N, Simsek C,
Lagrange J, Klemz R, Schönfelder T, Brandt M, Karbach SH, Knorr M, Finger S, Neukirch
C, Häuser F, Beutel ME, Kröller-Schön S, Schulz E, Schnabel RB, Lackner K, Wild PS,
Zeller T, Daiber A, Blankenberg S, Münzel T. Heme oxygenase-1 suppresses a
pro-inflammatory phenotype in monocytes and determines endothelial function and
arterial hypertension in mice and humans. Eur Heart J. 2015 Dec 21;36(48):3437-46. doi:
10.1093/eurheartj/ehv544. Epub 2015 Oct 29. PMID: 26516175.
Ulff-Møller CJ, Asmar F, Liu Y, Svendsen AJ, Busato F, Grønbaek K, Tost J, Jacobsen S. Twin
DNA Methylation Profiling Reveals Flare-Dependent Interferon Signature and B Cell
Promoter Hypermethylation in Systemic Lupus Erythematosus. Arthritis Rheumatol. 2018
Jun;70(6):878-890. doi: 10.1002/art.40422. Epub 2018 May 9. PMID: 29361205.
Drummond RA, Swamydas M, Oikonomou V, Zhai B, Dambuza IM, Schaefer BC, Bohrer AC,
Mayer-Barber KD, Lira SA, Iwakura Y, Filler SG, Brown GD, Hube B, Naglik JR, Hohl TM,
Lionakis MS. CARD9+ microglia promote antifungal immunity via IL-1β- and
CXCL1-mediated neutrophil recruitment. Nat Immunol. 2019 May;20(5):559-570. doi:
10.1038/s41590-019-0377-2. Epub 2019 Apr 17. PMID: 30996332.
von Wulffen M, Luehrmann V, Robeck S, Russo A, Fischer-Riepe L, van den Bosch M, van
Lent P, Loser K, Gabrilovich DI, Hermann S, Roth J, Vogl T. S100A8/A9-alarmin promotes
local myeloid-derived suppressor cell activation restricting severe autoimmune arthritis.

Cell Rep. 2023 Aug 29;42(8):113006. doi: 10.1016/j.celrep.2023.113006. Epub 2023 Aug
22. PMID: 37610870.
Kao KD, Grasberger H, El-Zaatari M. The Cxcr2+ subset of the S100a8+ gastric granylocytic
myeloid-derived suppressor cell population (G-MDSC) regulates gastric pathology. Front
Immunol. 2023 Sep 8;14:1147695. doi: 10.3389/fimmu.2023.1147695. eCollection 2023.
PMID: 37744359.

1013

Figures with figure Legends

Figure 1 Identification and characterization of placental immune cells using CyTOF

(A) Schematic of the experimental workflow in CyTOF experiment. The placentas were
obtained from individuals with normal pregnancy (NP, n=9), preeclampsia (PE, n=8),
gestational diabetes mellitus (GDM, n=8) or GDM&PE (n=7).

(B) t-SNE maps showing 7×10^5 CD45$^+$ cells from the placenta overlaid with color-coded
clusters and the distributions of B cells, CD4$^+$ T cells, CD8$^+$ T cells, DC, γδT cells, monocytes,
granulocytes, MDSC, and NK cells.

(C) Percentages of each cell type of CD45$^+$ cells in placentas.

(D) t-SNE maps showing the expression of CD3, CD8, CD4, γδTCR, CD14, CD15, CD56.

(E) Heatmap showing the expression levels of markers in CD45$^+$ cell subsets.

Data were compared between NP and PE, NP and GDM, NP and GDM&PE using the
Shapiro-wilk test and represented as mean±SEM (*P < 0.05, ** P < 0.01, *** P < 0.001; NS, not
significant).

1028

Figure 2 Specific altered T cell profile in the placentas of individuals with PE

(A) Distribution of the CD4$^+$ T cells analyzed using t-SNE.

(B) Scatter dot plots showing the frequencies of cluster 8 of CD4$^+$ T cells in the placentas of
individuals with NP, PE, GDM and GDM&PE.

(C) Expression of IL-17A and Foxp3 in CD45RO$^+$CCR7$^+$CD4$^+$ T cells in placentas of
individuals with NP and PE using flow cytometry. (IL-17A: n=10 in NP group, n=15 in PE group;
Foxp3: n=7 in NP group, n=9 in PE group).

(D) Immunofluorescence co-staining of CD4 (red), CD45RO (green) and DAPI (blue) in frozen
placental sections. The right panels show the enlargement of the dotted box in the left panel.
Scale bar, 20 µm.

(E) Scatter dot plots showing significantly altered markers of in cluster 8 of CD4$^+$ T cells.

(F) Distribution of the CD8$^+$ T cells analyzed using t-SNE.

(G) Scatter dot plots showing the frequencies of cluster 2 of CD8$^+$ T cells in the placentas of
individuals with NP, PE, GDM and GDM&PE.

(H) Scatter dot plots showing significantly altered markers in cluster 2 of CD8$^+$ T cells.

(I) Distribution of the γδT cells analyzed using t-SNE.

(J) Scatter dot plots showing the frequencies of cluster 15 of γδT cells in the placentas of
individuals with NP, PE, GDM and GDM&PE.

(K) Scatter dot plots showing significantly altered markers in cluster 15 of γδT cells.

Data were compared between NP and PE, NP and GDM, NP and GDM&PE using the
Shapiro-wilk test and represented as mean±SEM (*P < 0.05, ** P < 0.01, *** P < 0.001; NS, not
significant).

Figure 2—figure supplement 1 Identification of the placental T cell subsets.

(A) Heatmap showing the expression levels of markers in the $CD4^+$ T subsets.

(B) Heatmap showing the expression levels of markers in the $CD8^+$ T cells.

(C) Heatmap showing the expression levels of markers in the γδT cells.

1055

Figure 3 Identification of the placental $CD11b^+$ cell subsets and the Interaction between
placental immune cells.

(A) Distribution of the $CD11b^+$ immune cells analyzed using t-SNE.

(B) Heatmap showing the expression levels of markers in the $CD11b^+$ cells.

(C) Scatter dot plots showing the frequencies of cluster 12, cluster 23, and cluster 25 of
$CD11b^+$ cells in the placentas.

(D) Interaction between placental immune cells showed in heatmap.

(E) Scatter plots of Pearson's correlation analysis between placental immune cells.

Data were compared between NP and PE, NP and GDM, NP and GDM&PE using the
Shapiro-wilk test and represented as mean±SEM (*$P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

1066

Figure 4 The immune imbalance at the maternal-fetal interface induced by $F480^+CD206^-$
pro-inflam Macs.

(A) Schematic of the mice model adoptively transferred $CD45^+F4/80^+CD206^-$ pro-inflam Macs
or $CD45^+F4/80^+CD206^+$ anti-inflam Macs.

(B) Principal component analysis (PCA) reflected the differences between the two groups of
macrophages (n=3).

(C) The volcano map shows a comparison of the content and P value of gene expression
between pro-inflam Macs and anti-inflam Macs. Differential expression genes were screened
out when $P < 0.05$. Red dots indicate genes with increased expression in pro-inflam Macs.
Blue dots indicate genes with decreased expression.

(D) The volcano map shows differential expression genes between pro-inflam Macs and
anti-inflam Macs.

(E) Representative pathways enriched in the identified genes as determined by GSEA (p
value<0.05).

(F) Embryo abortion rate of the pregnant mice, body weight and crown-rump length of pups
measured on day 18.5 of gestation. Black represents mice treated with DMSO (n=8); gray
represents mice treated with PLX3397 (n=8); blue represents mice injected with
$CD45^+F4/80^+CD206^+$ anti-inflammatory macrophages (n=8); red represents mice injected with
$CD45^+F4/80^+CD206^-$ pro-inflammatory macrophages (n=8).

(G) SBP and UACR of pregnant mice in the four groups.

(H) Frequencies of $CD44^+CD4^+IL-17A^+$ cells, $CD44^+CD8^+$ T cells and $CD11b^+Ly6G^+$
granulocytes analyzed by flow cytometry.

Figure 4—figure supplement 1 Clodronate liposomes were used to used to deplete the
macrophages of pregnant mice to demonstrate that pro-inflam Macs lead to immune
imbalance.

(A) Embryo abortion rate of the pregnant mice, body weight and crown-rump length of pups
measured on day 18.5 of gestation. Black represents mice treated with control liposomes
(n=6); gray represents mice treated with clodronate liposomes (n=6); blue represents mice

injected with CD45$^+$F4/80$^+$CD206$^+$ anti-inflam Macs (n=6); red represents mice injected with
CD45$^+$F4/80$^+$CD206$^-$ pro-inflam Macs (n=6).
(B) SBP and UACR of pregnant mice in the four groups.
(C) Frequencies of CD44$^+$CD4$^+$IL-17A$^+$ cells, CD44$^+$CD8$^+$ T cells and CD11b$^+$Gr1$^+$
granulocytes analyzed by flow cytometry.
1100
Figure 5 Identification of the macrophage subsets in mice injected pro-inflam Macs and
anti-inflam Macs use sc RNA-seq.
(A) UMAP maps showing the 8 types of mouse immune cells at the maternal-fetal interface.
(B) Heatmap showing clustering analysis for markers distinguished different type of immune
cells.
(C) Heatmap showing clustering analysis for markers distinguished different clusters of
macrophages.
(D) UMAP maps showing the 15 clusters of mouse macrophages was listed in the left panel.
Bar graph showing the frequencies of clusters of macrophages in the two groups of mice was
listed in the right panel.
(E) UMAP maps showing the distribution of specific markers of cluster 0 and cluster 1.
(F) Dot plot depicting GO enrichment terms that were significantly enriched in the
differentially expressed genes in cluster 0 and cluster 1 from the pro-inflam Macs group
and the control group.
(G) Violin plot of specific differential gene expression in cluster 0 and cluster 1 between the
pro-inflam Macs group and the control group.
1117
1118
Figure 6 Identification of the T/NK cells subsets in mice injected pro-inflam Macs and
anti-inflam Macs use sc RNA-seq.
(A) UMAP maps showing the 12 clusters of mouse T/NK cells was listed in the up panel. Bar
graph showing the frequencies of clusters of T/NK cells in the two groups of mice was listed in
the down panel.
(B) Heatmap showing clustering analysis for markers distinguished 12 different clusters of
T/NK cells.
(C) UMAP maps showing the distribution of specific markers of cluster 0, 1 and 2.
(D) Dot plot depicting GO enrichment terms that were significantly enriched in the
differentially expressed genes in cluster 0 from the pro-inflam Macs group and the control
group.
(E) Violin plot of specific differential gene expression in cluster 0 between the pro-inflam
Macs group and the control group.
(F) Frequencies of CD4$^+$CD44$^+$ T cells and the percentages of IL-17A$^+$ cells in
CD4$^+$CD44$^+$ T cells at the maternal-fetal interface in Sham and RUPP group analyzed by
flow cytometry.
(G) The embryo abortion rate of pregnant mice, body weight, and crown-rump length of pups
measured on day 18.5 of gestation in mice injected PBS, Sham mouse-derived or RUPP
mouse-derived CD4$^+$CD44$^+$ T cells. Black represents mice injected with PBS (n=6); blue
represents mice injected with Sham mouse-derived CD4$^+$CD44$^+$ T cells (n=6); red represents

mice injected with RUPP mouse-derived CD4$^+$CD44$^+$ T cells (n=6).

(H) SBP and UACR of pregnant mice injected with PBS, Sham mouse-derived or RUPP

mouse-derived CD4$^+$CD44$^+$ T cells.

Figure 6—figure supplement 1 Memory-like Th17 cells may be associated with the

recurrence of PE.

(A) Experimental design of mice model with PE by reducing uterine perfusion pressure (RUPP)

on day 12.5 of gestation. Mice with sham operation were considered as controls. Systolic

blood pressure (SBP) and urine albumin creatine ratio (UACR) were measured on day 12.5

and 16.5 of gestation respectively. Mice were sacrificed on day 18.5 of gestation.

(B) The embryo abortion rate of pregnant mice, body weight, and crown-rump length of pups

measured on day 18.5 of gestation in the Sham and RUPP group. Embryo abortion rate =

number of absorbed embryos/total number of embryos. Blue represents mice in the Sham

group (n=9); Red represents mice in the RUPP group (n=9).

(C) SBP and UACR of pregnant mice in Sham and RUPP group.

(D) Embryo abortion rate of pregnant mice, body weight and crown-rump length of pups

measured on day 18.5 of gestation. Black represents mice with previous normal pregnancy

(n=9); gray represents mice with previous pregnancy with PE (n=9).

(E) SBP and UACR of second pregnant mice with a history of PE or NP in the first pregnancy

measured on day 16.5 of gestation.

(F) Frequencies of CD4$^+$ CD44$^+$ T cells and the levels of IL-17A in CD4$^+$ CD44$^+$ T cells in mice

with a pregnancy history with NP and PE analyzed by flow cytometry.

Data were compared using the Student's t-test and represented as mean±SEM (*P < 0.05, ** P

< 0.01, *** P < 0.001; NS, not significant).

1162

Figure 7 Identification of the granulocytes subsets in mice injected pro-inflam Macs and

anti-inflam Macs use sc RNA-seq.

(A) UMAP maps showing the 9 clusters of mouse granulocytes was listed in the up panel. Bar

graph showing the frequencies of clusters of granulocytes in the two groups of mice was listed

in the down panel.

(B) Heatmap showing clustering analysis for markers distinguished 9 different clusters of

granulocytes cells.

(C) UMAP maps showing the distribution of specific markers of cluster 0, 1, 2 and 3.

(D) Dot plot depicting GO enrichment terms that were significantly enriched in the

differentially expressed genes in cluster 3 from the pro-inflam Macs group and the control

group.

(H) Violin plot of specific differential gene expression in cluster 3 between the pro-inflam

Macs group and the control group.

1176

Figure 8 Pro-inflam Macs induce the generation of memory-like Th17 cells via

IGF1-IGF1R

(A) Signaling modules indicated by ligand-receptor pairing between macrophages and other

types of immune cells at the maternal-fetal interface using CellPhoneDB.

(B) Frequencies of IGF1$^+$CD14$^+$ cells and IGFIR$^+$CD4$^+$ cells in placentas of individuals with NP

and PE (n=12 in NP group, n=4 in PE group).

(C) Schematic of the experimental workflow to induce memory-like T cells in vitro.
Macrophages, after incubating with PBS, NP-EVs or PE-EVs, were co-cultured with CD4[+]
naïve T cells treated with DMSO or BMS-754807. Cells were isolated from human peripheral
blood.
(D) Frequencies of CD45RO[+]CCR7[+]Th17 cells. Black represents CD4[+] naïve T cells treated
with DMSO; gray represents CD4[+] naïve T cells cocultured with NP-EVs-treated macrophages;
blue represents CD4[+] naïve T cells co-cultured with PE-EVs-treated macrophages; red
represents CD4[+] naïve T cells treated with BMS-754807 before co-cultured with
PE-EVs-treated macrophages (n=10 in each group).
(E) Schematic of mice transferred CD4[+] T cells treated with BMS-754807 or PBS. Anti-CD4
antibody was used to deplete CD4[+] T cells in mice on day 10.5 of gestation. CD4[+] T cells were
transferred into mice on day 11.5 of gestation. 20 µg/kg lipopolysaccharide (LPS) was
intraperitoneally injected on day 12.5 and 15.5 of gestation to induce a PE-like pregnant mice
model. Mice were sacrificed on day 18.5 of gestation.
(F) Embryo abortion rate of pregnant mice, body weight and crown-rump length of pups were
measured on day 18.5 of gestation. Black represents the control group mice (n=6); gray
represents mice treated with LPS (20µg/kg) to construct an animal model of PE (n=6); blue
represents anti-CD4 antibody treated PE mice (n=6); red represents anti-CD4 antibody treated
PE mice injected with CD4[+] T cells with DMSO treatment (n=7); orange represents anti-CD4
antibody treated PE mice injected with CD4[+] T cells with BMS754807 treatment (n=7).
(G) SBP and UACR of pregnant mice in the five groups.
(H) The frequencies of CD4[+] CD44[+] IL-17A[+] cells analyzed by flow cytometry.
The results were compared using one-way ANOVA and represented as mean±SEM (*P < 0.05,
** P < 0.01, *** P < 0.001; NS, not significant).
Figure 8—figure supplement 1 Cell–cell communications in immune cells when
pro-inflam Macs accumulated at the maternal–fetal interface.
(A) Abundance of connections between different cell types at the maternal-fetal interface
analyzed by CellPhoneDB.
(B) Capacities for interactions between immune cells are showed. Each line indicates the
ligands expressed by the cell population represented by the same color (labeled). The lines
connect to cell types that express cognate receptors. Line thickness is proportional to the
number of ligands when cognate receptors are present in the recipient cell type.
1215
1216 **Tables**
1217 **Table 1 Details of the individuals included in the CyTOF**

Parameters	NP (n=9)	PE (n=8)	GDM (n=8)	GDM&PE (n=7)	p Value
Age (years)	31.33 ± 2.108	31.75 ± 4.206	35.25 ±3.419	32.43 ± 2.321	0.1008
BMI (kg/m^2)	27.35 ± 1.740	30.84 ±2.888	29.56 ± 2.689	30.16 ± 2.280	0.0518
Gestational age (weeks)	38.89 ± 0.558	37.39 ± 1.253	38.73 ± 0.419	36.22 ± 1.372	<0.001
Number of living children	0.4444 ± 0.497	0.500 ± 0.500	1.125 ± 0.331	0.429 ± 0.495	0.0236
Previous abortions	0.7778 ± 1.030	1.000 ± 0.866	1.000 ± 1.000	1.714 ± 2.050	0.5848
Mean Systolic Blood Pressure	109.7 ± 5.400	151 ± 7.225	106.9 ± 2.848	150.7 ± 9.161	<0.0001

1218

1219 **Table 2 CyTOF antibody panel used for analyzing placentas from individuals with**
1220 **normal pregnancy (NP), preeclampsia (PE), gestational diabetes mellitus (GDM) and**
1221 **GDM complicated with PE (GDM&PE)**

Target	Metal tag
CD45	HI30
CD3	UCHT1
CD68	Y1/82A
CD56	NCAM16.2
gd TCR	5A6.E9
CD19	HIB19
CCR6	G034E3
CD38	HIT2
CD103	B-Ly7
CD39	A1
CXCR3	G025H7
PD-L1	29E.2A3
PD-1	EH12.2H7
CD11C	BU15
CD107a	H4A3
pAKT	D9E
CCR4	L291H4
TIGIT	A15153G
CD206	15-2
Helios	22F6
CD28	CD28.2
pSTAT3	4/p-Stat3
GITR	110416
CD33	WM53
CTLA-4	BN13
FOXP3	PCH101
CD163	GHI/61
CD45RA	HI100
ICOS	C398.4A
CD69	FN50
CCR7	G043H7
NKG2A	131411
CD15	W6D3
TIM-3	F38-2E2
CD86	Fun-1
HLA-DR	L243
Granzyme B	QA16A02
CD14	M5E2
pS6	A17020B
CD127	A019D5
CD4	RPA-T4
CD8	RPA-T8
CD11b	M1/70

1222

1223

1224

    Table 3 Expression levels of markers identified in each immune subset in the placentas

	B cells	CD4[+] T	CD8[+] T	gdTCR	NK cells	Macrophage	Granulocyte	MDSC	Dendritic cells
CD45	++	++	++	++	++	++	++	++	++
CD3	-	++	++	-	-	-	-	-	-
CD68	+	+	+	+	+	++	+	++	++
CD56	-	-	-	-	+	-	-	-	-
gd TCR	++	-	-	++	-	-	-	-	-
CD19	++	-	-	++	-	-	-	-	-
CCR6	+	-	-	-	-	-	-	-	-
CD38	++	++	+	+	++	++	-	-	+
CD103	-	-	-	-	-	-	-	-	-
CD39	++	-	-	-	+	++	+	+	+
CXCR3	+	+	+	+	+	+	-	+	++
PD-L1	-	-	-	-	-	-	-	+	-
PD-1	-	-	-	-	-	-	-	-	-
CD11C	+	-	-	-	+	++	++	+	++
CD107a	+	-	-	-	-	+	-	+	+
pAKT	+	+	+	+	+	++	+	+	+
CCR4	-	-	-	-	-	-	-	-	-
TIGIT	-	-	-	-	-	-	-	-	-
CD206	-	-	-	-	-	+	-	+	-
Helios	+	+	+	+	+	+	+	+	+
CD28	-	+	+	+	-	-	-	-	-
pSTAT3	+	+	-	+	-	+	+	-	-
GITR	-	-	-	-	-	-	-	-	-
CD33	-	-	-	-	-	+	-	+	-
CTLA-4	-	-	-	-	-	-	-	-	-
FOXP3	-	-	-	-	-	-	-	-	-
CD163	-	-	-	-	-	-	-	-	-
CD45RA	++	+	+	+	++	-	-	-	-
ICOS	-	+	+	-	-	-	-	-	-
CD69	+	+	+	+	++	+	-	++	-
CCR7	+	++	+	+	-	-	-	+	-
NKG2A	-	-	-	-	-	-	-	-	-
CD15	-	-	-	-	-	-	++	-	-
TIM-3	-	-	-	-	-	-	-	-	-
CD86	-	-	-	-	-	++	-	-	+
HLA-DR	++	-	+	+	-	++	+	-	++
Granzyme	+	-	-	+	++	+	+	-	-
CD14	-	-	-	-	-	++	-	-	-
pS6	-	-	-	-	-	-	-	-	-
CD127	-	++	+	-	-	-	-	-	-
CD4	-	++	-	-	-	-	-	-	-
CD8	-	-	++	-	-	-	-	-	-
CD11b	-	-	-	+	+	++	++	-	+

Table 4 The marker profile of PE-specific immune subsets

	memory-like CD4+ T cells Cluster 08 of CD4+ T cells	memory-like CD8+ T cells Cluster 02 of CD8+ T cells	gMDSCs Cluster 12 of CD3-CD11b+ cells	pro-inflammatory macrophages Cluster 23 of CD3-CD11b+ cells	anti-inflammatory macrophages Cluster 25 of CD3-CD11b+ cells
CD45	++	++	++	++	++
CD3	++	++	-	-	-
CD68	-	-	+	++	++
CD56	-	-	-	-	-
gd TCR	-	-	-	-	-
CD19	-	-	-	-	-
CCR6	-	-	-	-	-
CD38	++	++	-	+	+
CD103	-	-	-	+	+
CD39	+	-	+	-	-
CXCR3	+	+	-	-	-
PD-L1	-	-	-	-	-
PD-1	-	-	-	-	-
CD11C	-	-	++	++	++
CD107a	-	-	-	+	+
pAKT	+	+	-	-	-
CCR4	-	-	-	-	-
TIGIT	-	-	-	-	-
CD206	-	-	+	-	+
Helios	+	+	-	-	-
CD28	+	+	-	-	-
pSTAT3	-	-	-	-	-
GITR	-	-	-	-	-
CD33	-	-	-	-	+
CTLA-4	-	-	-	-	-
FOXP3	-	-	-	-	-
CD163	-	-	-	-	-
CD45RA	-	-	-	-	-
ICOS	+	+	-	-	-
CD69	+	-	-	-	-
CCR7	++	++	-	-	-
NKG2A	-	-	-	-	-
CD15	-	-	++	-	-
TIM-3	-	-	-	-	-
CD86	-	-	-	+	+
HLA-DR	-	-	+	+	++
Granzyme B	-	-	+	-	-
CD14	-	-	-	+	+
pS6	-	-	-	-	-
CD127	+	-	-	-	-
CD4	++	-	-	-	-
CD8	-	++	-	-	-
CD11b	-	-	++	++	++

1229   **Table 5 Details of the individual with NP or PE included in the study**

Parameters	NP (n=30)	PE (n=30)	p Value
Age (years)	31.13 ± 2.9747	31.50 ± 4.1693	0.0741
BMI (kg/m^2)	27.24 ± 2.8048	29.48±3.9295	0.0155
Gestational age (weeks)	38.68± 0.6038	35.53 ± 3.29	<0.0001
Number of living children	0.6000 ± 0.4900	0.4667 ± 0.6182	0.3665

1230

1231

1232   **Table 6 Details of the individual with early-onset preeclampsia or late-onset**
1233   **preeclampsia included in the study**

Parameters	Early-onset preeclampsia(n=11)	Late-onset preeclampsia (n=19)	p Value
The gestational week when blood pressure begins to rise	28.82± 2.8220	36.05± 1.129	< 0.0001
Age (years)	32.45± 4.6768	32.36± 3.8472	0.9568
BMI (kg/m^2)	29.64± 2.6840	28.71±4.4506	0.5388
Gestational age (weeks)	33.45± 3.7887	36.77± 1.2171	0.0014
Number of living children	0.36± 0.5045	0.36 ± 0.4956	0.98

1234

1235

1236

1237

1238

A CD4+ T cells

NP   PE   GDM   GMD&PE

C01 C09
C02 C10
C03 C11
C04 C12
C05 C13
C06 C14
C07 C15
C08

B Cluster 8 of CD4+ T cells

NS
NS
*

NP
PE
GDM
GDM&PE

C

CD45RO / CCR7

NP   PE

IL-17A   10.87%   23.88%

Foxp3   23.47%   8.39%

CD4

Th17A   Treg

D CD45RO / CD4 / DAPI

NP

PE

E Cluster 08

NS
NS
*

NS
NS
*

PD-1   TIM3

F CD8+ T cells

NP   PE   GDM   GMD&PE

C01 C10
C02 C11
C03 C12
C04 C13
C05 C14
C06 C15
C07 C16
C08 C17
C09 C18

G Cluster 2 of CD8+ T cells

NS
NS
*

I γδT cells

NP   PE   GDM   GMD&PE

C01 C12
C02 C13
C03 C14
C04 C15
C05 C16
C06 C17
C07 C18
C08 C19
C09 C20
C10 C21
C11 C22

H Cluster 2 of CD8+ T cells

NS
NS
*

NS
NS
*

PD-1   TIGIT

J Cluster 15 of γδT cells

NS
NS
*

K Cluster 15 of γδT cells

NS
NS
*

Tim3

CD4⁺ T cells

CD8⁺ T cells

γδT cells

A

B