UNC-6/Netrin promotes both adhesion and directed growth within a single axon

Ev L. Nichols[1], Joo Lee[1,4], and Kang Shen[1,2,3,5]*

1. Department of Biology, Stanford University, Stanford, CA 94305, USA
2. Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA
3. Wu Tsai Neurosciences Institute, Stanford University, Stanford, CA 94305, USA
4. Present address: Department of Developmental Biology, Stanford, CA 94305, USA
5. Lead Contact

*Correspondence: kangshen@stanford.edu

Summary

During development axons undergo long-distance migrations as instructed by guidance molecules and their receptors, such as UNC-6/Netrin and UNC-40/DCC. Guidance cues act through long-range diffusive gradients (chemotaxis) or local adhesion (haptotaxis). However, how these discrete modes of action guide axons *in vivo* is poorly understood. Using time-lapse imaging of axon guidance in *C. elegans*, we demonstrate that UNC-6 and UNC-40 are required for local adhesion to an intermediate target and subsequent directional growth. Exogenous membrane-tethered UNC-6 is sufficient to mediate adhesion but not directional growth, demonstrating the separability of haptotaxis and chemotaxis. This conclusion is further supported by the endogenous UNC-6 distribution along the axon's route. The intermediate and final targets are enriched in UNC-6 and separated by a ventrodorsal UNC-6 gradient. Continuous growth through the gradient requires UNC-40, which recruitsUNC-6 to the growth cone tip. Overall, these data suggest that UNC-6 stimulates stepwise haptotaxis and chemotaxis *in vivo*.

Keywords

Axon guidance, Netrin, DCC, chemotaxis, haptotaxis, Trim9, nervous system

development, cellular migration, time-lapse imaging

Introduction

Axon guidance is critical for establishing functional neural circuits. Given the

importance of axon growth in laying the morphological foundations for neural circuitry,

axon guidance has been of interest to neuroscientists since Ramón y Cajal first observed

growing axons (termed growth cones) in histological examinations of the embryonic

nervous system (Ramón y Cajal, S., 1911). Early cell ablation studies established that

growing axons rely on anatomical landmarks, or intermediate targets, to properly navigate

through their environment. A classic example of intermediate targets is in the grasshopper

hindlimb, where pioneering Ti1 sensory axons navigate from the tip of the limb to the

animal's central nervous system. Instead of following the shortest route, Ti1 axons grow

toward a series of guidepost cells whereby they sequentially readjust their trajectory upon

reaching a guidepost (Keshishian and Bentley, 1983). In fact, these intermediate cells are

necessary for proper sensory axon guidance as their ablation results in Ti1 axon

misrouting (Bentley and Caudy, 1983a, 1983b). Studies in other model organisms have

identified additional examples of intermediate targets across phylogeny: the ventral nerve

cord (VNC) in *C. elegans* (Hedgecock et al., 1990), the *Drosophila* midline (Klämbt et al.,

1991), and the floor plate (Bernhardt et al., 1992; Kennedy et al., 1994; Serafini et al.,

1994) and optic chiasm in vertebrates (Chung et al., 2000). However, given the transient

nature of a growth cone's interactions with its intermediate targets, detailed *in vivo*, cell

biological studies of these interactions remain sparse.

The stereotyped axonal behaviors at the intermediate target require molecular

cues which confer directional information to growing axons. A critical advance of the field

was the identification of numerous ligands (Netrins, Slits, Semaphorins, and Ephrins)

responsible for axon guidance and their cognate receptors (Kolodkin and Tessier-

Lavigne, 2011). These signaling pathways are generally characterized as either attractive

or repulsive, depending on the receptor-ligand combination (Comer et al., 2019). For

example, Netrin attracts axons when signaling through a receptor homodimer of Deleted

in Colorectal Cancer (DCC) or Neogenin1 (Neo1) whereas it repels axons when bound

to a receptor dimer that includes UNC-5 (Kolodkin and Tessier-Lavigne, 2011).

Netrin was initially proposed to attract axons as a chemotactic cue mediated by a

long-range action of diffusible Netrin present in a gradient(Comer et al., 2019). Initial

characterizations of Netrin in the vertebrate spinal cord identified it as a diffusive cue

produced by the floor plate through spinal explants (Kennedy et al., 1994; Serafini et al.,

1994). More recently, *in vivo* examinations have visualized a Netrin-1 gradient emanating

from the floor plate (Dominici et al., 2017; Kennedy et al., 2006). Together, these data

support a role for Netrin in stimulating chemotaxis. Together, these data support a role for

Netrin in stimulating chemotaxis.

Alternatively, Netrin can stimulate haptotactic migration through local, contact-

dependent interactions between the growth cone and an anchored Netrin ligand (Comer

et al., 2019). Haptotactic mechanisms of Netrin guidance are supported by experiments

in the *Drosophila* nervous system where Netrin is required for commissural axons to cross

the midline. Artificial tethering of Netrin to the cell membrane at the midline is sufficient to

drive commissural axon orientation and midline crossing (Brankatschk and Dickson,

2006). Similarly, in the *Drosophila* visual system, R8 photoreceptors utilize local Netrin

and Frazzled (Fra)/DCC to stabilize their axons at the neuropil (Timofeev et al., 2012).

Mutants for *net*, *fra,* or *trim9* (a downstream regulator of Netrin signaling) navigate

normally to the neuropil but fail to stabilize, eventually retracting (Akin and Zipursky,

2016). Notably, evidence of Netrin-induced haptotaxis is not limited to *Drosophila.*

Investigations in the vertebrate spinal cord and rhombic lip have identified local roles for

Netrin in promoting axon growth and confinement within the central nervous system

(Varadarajan et al., 2017; Yung et al., 2018).

In *C. elegans*, current understandings posit that the VNC produces the Netrin

homolog UNC-6 which forms a ventrodorsal gradient (Wadsworth et al., 1996). Axons

navigating ventrally sense this UNC-6/Netrin gradient and chemotax toward the UNC-6

source at the VNC. Here, we utilize time-lapse imaging in *C. elegans* to visualize growth

cone behaviors during ventral axon guidance in an intact nervous system. These

experiments identify the ventral sublateral region of the worm as a common intermediate

target where ventral growth cones stabilize before further migration. Both stabilization and

growth depend on UNC-6/Netrin and UNC-40/DCC. Non-diffusible UNC-6 is sufficient to

rescue axon stabilization but not growth. Last, imaging of endogenous UNC-6 localization

during axon outgrowth is consistent with a model where the extending growth cone

enriches UNC-6 around the cell membrane. Together, these experiments support a model

where haptotaxis and chemotaxis act successively in UNC-6-mediated axonal migration.

Results

The ventral sublateral region is an intermediate target for ventral axons

C. elegans is a canonical model for axon guidance given its conservation of axon

guidance genes and stereotyped neuronal identities and morphologies (Chisholm et al.,

2016). While the majority of *C. elegans* neurons are born embryonically, a subset is born

during larval development which enables direct observations of axon guidance by time-

lapse imaging experiments. Of these postembryonic neurons, 5 (AVM, HSN, PVM, PDE,

and PQR) initiate pioneering axons toward the ventral region of the animal where they

join the ventral nerve cord (VNC) (Figure 1A). We sought to leverage this common

guidance trajectory and accessible visualization to provide new insights into mechanisms

of axon guidance. First, we developed a paradigm that permits complete visualization of

PDE axon's ventral growth in wildtype animals. This strategy used a *Plin-32::mCherry-*

PH integrated transgene to label the PDE cell membrane and imaged PDE axon

outgrowth every 3 min for 4 hr (Figure 1B, Video S1). The mature axonal morphology of

PDE is unbranched along the dorsoventral axis before bifurcating at the VNC (Figure 1B).

The continuous, unbranched morphology of the mature PDE axon suggests continuous

ventral growth after axon initiation. Surprisingly, however, we observed that the PDE axon

consistently halts its ventral growth upon reaching the region near the ventral sublateral

nerve cord (vSLNC), a small axon tract that runs along the anteroposterior body axis of

the animal. During its pause, the axon establishes transient collateral branch from which

additional dorsal and ventral branches emerge. Eventually, one ventral branch stabilizes

at the VNC, bifurcates, and initiates fasciculated growth along VNC axons. Following VNC

stabilization, the transient collateral branch at the vSLNC is pruned, resulting in the

unbranched PDE axonal morphology observed in adulthood. The stereotyped PDE

axonal pause and consistent transient branching at vSLNC indicate that this is an

intermediate guidance target.

The vSLNC sits atop the ventral body wall muscles of the worm and adjacent to

the hypodermis, thereby demarcating an anatomical region that separates two distinct

environments through which the axon grows (Figure 1C,D). In adult animals, the vSLNC

is ensheathed by the hypodermis, but the ensheathment process begins during larval

stages and is not yet completed by PDE axonogenesis (Chalfie and Sulston, 1981). Prior

to reaching the vSLNC, the PDE axon grows along the hypodermal basal lamina, but

additional ventral growth past the sublateral region requires that axons wedge themselves

between fibrous adhesive junctions between the body wall muscles and the hypodermis

(Figure 1C,D) (Cox and Hardin, 2004). This stark contrast in growth environments

correlates with alternate forms of morphological axon growth. Prior to reaching the

vSLNC, the growth cone of PDE is wide and reminiscent of a lamellipodium. However,

ventral axon branching from the transient collateral branch are consistent with filopodia-

based motility.

Given that the body wall muscles tile the anteroposterior axis of the entire animal,

other ventrally directed axons also must undergo a migration through these distinct

regions. We next hypothesized that the other ventrally growing axons would similarly

utilize the ventral sublateral region as an intermediate target. To test this, we imaged axon

emergence in AVM, HSN, PVM, and PQR neurons. While these neurons position their

somas more ventrally, we observe similar axonal trajectories as the PDE axon: axons

initially emerge ventrally as lamellipodia, then adhere upon reaching the vSLNC. Multiple

ventrally directed filopodia form and a ventral branch eventually stabilizes at the axons'

ultimate ventral target, the VNC (Figure 1E). Taken together, these data reveal that ventral

axon guidance in *C. elegans* is a stepwise process facilitated by an intermediate target

near the vSLNC. Notably, dorsally migrating axons in *C. elegans* also stall their growth

cones at the borders of longitudinal nerve tracks and muscle borders, forming anvil-

shaped structures (Knobel et al., 1999). These results argue that the stepwise migration

and pausing of growth cone at specific tissue structures is a general phenomenon in *C.*

elegans, prompting us to understand the molecular bases of this cellular behavior.

UNC-6/Netrin signaling is required for multiple steps of ventral axon guidance

Circumferential axon migration in *C. elegans* is primarily driven by signaling

through UNC-6/Netrin and its receptors (Hedgecock et al., 1990; Wadsworth et al., 1996).

Dorsal guidance requires UNC-6 to signal through both its receptors, UNC-5 and UNC-

40/DCC. Ventral guidance does not require UNC-5 signaling but is dependent on UNC-

40 (Hedgecock et al., 1990). Most evidence supporting this notion are based on analyses

of post-developmental axonal morphologies in mutants instead of visualizing the

developmental process. We used time-lapse imaging of PDE axon outgrowth in null

mutants of *unc-6* (*ev400)* and *unc-40* (*e1430)* to understand whether UNC-6/UNC-40

signaling is required for one or all of the events: 1. initial axon outgrowth direction, 2.

stabilization at the intermediate target, 3. directional branching from the sublateral region

or 4. ventral growth from intermediate target to VNC. Importantly, the guidance of the

vSLNC axons do not rely on UNC-6 or UNC-40, so the putative intermediate target is

intact in these mutants(Li et al., 2008). The resulting movies demonstrate that each step

of ventral growth is disrupted in *unc-6* and *unc-40* mutants (Figure 2A, Video S1). First,

in most wildtype animals, the axon emerges from the ventral side of PDE, while axons

emerge from non-ventral directions of PDE in *unc-6* and *unc-40* mutants (Figure S2A).

Second, over 80% of wildtype axons form a stable collateral branch immediately upon

reaching the vSLNC. In contrast, nearly half of the *unc-6* and *unc-40* mutant axons form

collateral branches at vSLNC but subsequently retract (Figure 2A,B). Typically, these

axons "probe" the vSLNC again after retraction and establish contacts with vSLNC again

during the imaging window. However, retraction upon initial sublateral contact is rarely

observed in wildtype animals, indicating that UNC-6 signaling stabilizes the growth cone

at the sublateral intermediate target.

To further understand how UNC-6 and UNC-40 regulate axon development, we

inserted a GFP tag at the C-terminus of the endogenous *unc-40* locus. The resulting

animals show UNC-40::GFP fluorescence in the nerve ring and major nerve cords of the

worm (VNC and dorsal nerve cord), consistent with a published endogenous UNC-

40::mNeonGreen strain (Figure S1A-D) (Jayadev et al., 2022). We next analyzed movies

of UNC-40::GFP during PDE axon outgrowth. These movies demonstrate that the PDE

growth cone contains low levels of UNC-40::GFP prior to reaching the sublateral region.

However almost immediately upon contacting the sublateral, UNC-40::GFP fills the

growth cone along the transient collateral branch (Figure 2C,D). However, *unc-6(ev400)*

mutants fail to increase growth cone UNC-40::GFP upon reaching the sublateral region

and instead demonstrate seemingly random fluctuations in growth cone UNC-40::GFP

intensity (Figure 2C,D). These results suggest that UNC-6 regulates UNC-40 levels in the

growth cone.

Given similar morphological polarization PDE and HSN growth cones, we sought

to generalize these findings in HSN which develops more slowly than PDE and is

therefore more amenable to imaging axon engagement with the ventral sublateral

intermediate target. Indeed, endogenous UNC-40::GFP is enriched at the ventral surface

of HSN cell body where its axon emerges. However, this enrichment is absent in *unc-6*

mutants, consistent with previous studies using overexpressed UNC-40::GFP constructs

(Figure 2E) (Adler et al., 2006). To quantify UNC-40 polarity, we measured the total GFP

intensity along the periphery of the dorsal soma and sublateral contact site of HSN. The

overall UNC-40::GFP polarity was calculated by dividing the difference between the two

subcellular locations by the total GFP intensity along the cell periphery (Figure 2F). The

resulting polarity index ranges from -1 to 1 with negative numbers indicating a ventral

UNC-40::GFP bias and positive numbers indicating a dorsal bias. A polarity value of 0

denotes an equal dorsoventral distribution of UNC-40::GFP. We also calculated the

polarity of UNC-40 in AVM, PVM, and PQR in wildtype and *unc-6(ev400)* animals.

Wildtype animals display negative polarity calculations, but in *unc*-6 mutant animals, the

polarity index is close to 0 indicating that UNC-40 is evenly distributed between the dorsal

and ventral neuronal domains (Figures 2E,G; S1E-J). Together, these data indicate that

upon vSLNC contact, UNC-6 signaling enriches UNC-40 at the contact site and facilitates

adhesion with the surrounding environment.

Once the axon establishes a lateral branch at the vSLNC, it generates dorsally and

ventrally-oriented filopodia (Fig. 2A). These filopodia are dynamic and undergo constant

extension and retraction. Using time-lapse movies to capture the dynamics, we quantified

the directionality, length, growth speed and stability of these branches (Fig.2H-K).

Wildtype animals generate slightly more ventral branches than dorsal ones (55.6%

ventrally directed branches vs. 44.4% dorsal). However, *unc-6* and *unc-40* mutants have

a strong reversal of this bias where 61.4% and 71.9% of branching, respectively, is

directed dorsally (Figure 2H). These data suggest that UNC-6/UNC-40 signaling

promotes ventral branching. To further test this, we analyzed the growth trajectories of

individual ventral branches. This analysis demonstrates that ventral growth stalls in *unc-*

6 and *unc-40* mutants, manifesting as a decreased average growth speed (Figure 2I,J).

While *unc-6* and *unc-40* mutant PDE axonal filopodia reach the VNC, very few stabilize

within the imaging window. Instead, most branches retract, indicating that UNC-6 and

UNC-40 are also require for stabilizing filopodia once it reaches the VNC (Figure 2K).

Taken together, these data indicate that UNC-6 and UNC-40 are required for

several steps of axon guidance including the direction of axonal initiation, axonal

adhesion at the intermediate target, directional growth toward the VNC and axonal

stabilization at the VNC. While the developmental process is dynamic with numerous

growth and retraction events, UNC-6 and UNC-40's impacts on the stepwise events

compound to ultimately disrupt the adult morphology of ventrally directed axons. In *unc-*

6(ev400) mutants, 67.9% $\pm$ 5.1% of PDE axons and 66.9% $\pm$ 4.3% of HSN axons fail to

reach the VNC by adulthood, compared with only 0% of PDE axons and 0.91% $\pm$ 0.9%

of HSN axons in wildtype animals (Tables 1,2; Figure S1L).

MADD-2/Trim9/Trim67 promotes growth cone adhesion

Our data thus far demonstrate that PDE axon outgrowth is an excellent *in vivo*

model of Netrin-mediated guidance where precise axonal behaviors can be probed

across distinct change-of-function genotypes. Therefore, we next asked whether

regulators of UNC-6/Netrin signaling might only affect growth or adhesion. For this

analysis, we chose the E3 ubiquitin ligase MADD-2/Trim9/Trim67 because of its

conserved role in Netrin-dependent guidance and morphological polarization of the HSN

emerging axon (Akin and Zipursky, 2016; Hao et al., 2010; Plooster et al., 2017). We

captured movies of PDE axon guidance in *madd-2* null (*ok2226)* animals (Figure 3A).

These analyses revealed that *madd-2* mutants showed normal growth cone orientation

at emergence. However, the majority of the *madd-2* mutant axons retracted upon vSLNC

contact similar to *unc-6* mutants, suggesting that MADD-2 contributes to UNC-6 mediated

adhesion to the intermediate target (Figures 3B).

Given UNC-40's requirement for adhesion and ventral polarization, we next asked

whether MADD-2 is required for the polarized subcellular localization of UNC-40. To test

this, we observed endogenous UNC-40::GFP polarization within HSN, AVM, PVM, and

PQR in *madd-*2 mutants. These images reveal that UNC-40 loses its ventral polarity in

madd-2 mutants (Figure 3C,D; S2A-F). We confirmed these results in PDE by analyzing

growth cone UNC-40 levels during axon guidance. Consistent with the results in HSN,

each of AVM, PVM, and PQR, the PDE growth cone showed reduced UNC-40

polarization at the vSLNC (Figure S2B-H). Together, these data reveal a requirement for

MADD-2 to polarize UNC-40 at endogenous levels. Further, we analyzed *madd-2; unc-6*

double mutants for UNC-40::GFP polarity in HSN. These animals demonstrate enhanced

mispolarization of UNC-40 (Figure 3C,D), suggesting that UNC-6 and MADD-2 might

promote the polarity of the UNC-40 receptor in parallel pathways.

Existing literature suggests that Trim9 ubiquitinates DCC and that Netrin

antagonizes this activity (Plooster et al., 2017). To test whether UNC-40 ubiquitination is

required for its polarization at the vSLNC, we generated *unc-40(4KR)* animals with lysine

to arginine mutations at the 4 intracellular lysine residues adjacent to the UNC-40

transmembrane domain to hamper potential ubiquitination. These mutations were made

in the UNC-40::GFP::AID background to allow visualization of endogenous levels of

mutant UNC-40. These four amino acid substitutions are sufficient to mispolarize UNC-

40 in HSN, comparable to polarity measurements in *madd-2(ok2226)* animals (Figure

3C,E,F). Given that MADD-2 directly regulates UNC-40 localization through

ubiquitination, these data suggest that MADD-2 ensures that UNC-40 is polarized and

enriched at the growth cone at the sublateral region to promote adhesion.

In the next stage of growth, the movies of *madd-2* mutant demonstrate that PDE

axons primarily initiate ventrally directed branches similar to wildtype animals, indicating

that the directional growth of PDE axon is independent of MADD-2 (Figure 3G). However,

these ventral branches also display decreased stabilization at the VNC similar to *unc-6*

mutants (Figure 3H). Together, these experiments reveal two important insights: axonal

adhesion and stabilization are dependent on MADD-2 but directional growth is not.

Diffusible UNC-6 is dispensable for axon stabilization

UNC-6/Netrin has been proposed to function via either chemotaxis or haptotaxis

(Comer et al., 2019; Varadarajan et al., 2017; Wu et al., 2019). Chemotactic growth occurs

in response to a gradient where axons sense the Netrin concentration in their immediate

environment to determine their direction of travel toward or away from the Netrin source.

In contrast, haptotactic growth does not rely on a long-distance gradient and instead uses local, minimally diffusive Netrin to anchor growth along a laminar surface. In the vertebrate spinal cord, commissural axons utilize both mechanisms of Netrin-mediated guidance to properly establish ventral tracts that avoid the developing motor column (Varadarajan et al., 2017; Wu et al., 2019).

Therefore, we sought to distinguish between these two UNC-6 growth mechanisms during stepwise PDE axon guidance by expressing a non-diffusible form of UNC-6 in the intermediate target region in the *unc-6* mutants. Specifically, we fused the *unc-6* coding sequence to the transmembrane domain sequence of *nlg-1/neuroligin* so that the molecule is tethered to the plasma membrane (Teichmann and Shen, 2011). We generated two membrane-tethered UNC-6 expression constructs: one under the control of *mec-7* promoter elements for expression along the vSLNC and one under the control of the *unc-4* promoter for expression at the VNC (Figure 4A). Importantly, UNC-4 expressing cholinergic motor neurons produce endogenous UNC-6 in larval stages and MEC-7 expressing mechanosensory neurons do not. However, membrane-tethered UNC-6 would be present on the axons in VNC but unable to diffuse towards the PDE cell body. Using these constructs, we generated three different transgenic strains in the *unc-6* mutant background, a strain expressing *Pmec-7::unc-6^{TM}*, a strain expression *Punc-4::unc-6^{TM}* and a strain carrying both. With these transgenes, we aimed to determine whether the distinct axon behaviors at the vSLNC intermediate target and at VNC are mediated by UNC-6's chemotactic or haptotactic activity.

These transgenes are unable to restore ventral directionality of axon emergence (Figure S3). Interestingly, stabilization at vSLNC is restored when non-diffusible UNC-6

is present at vSLNC (*Pmec-7::unc-6^{TM}*) but not when expressed only at the VNC (*Punc-*

4::unc-6^{TM}), consistent with stabilization being contact-mediated (Figure 4B,C). Given the

rapid and spatially specific accumulation of growth cone UNC-40 observed in the movies

of PDE axon outgrowth, we next tested whether membrane-tethered UNC-6 could rescue

UNC-40::GFP polarization in HSN. Indeed, non-diffusible UNC-6 along the vSLNC

restored UNC-40::GFP polarity in HSN (Figure 4D,E). On the contrary, non-diffusible

UNC-6 expressed along the VNC could not restore ventral UNC-40 polarization (Figure

4D,E), presumably due to the inability of HSN to contact this UNC-6 population. These

data indicate that axonal stabilization at the vSLNC intermediate target requires local

UNC-6, and membrane tethered UNC-6 is sufficient for this function. Therefore,

haptotaxis is sufficient to account for adhesion and UNC-40 polarization.

Next, we analyzed the directional filopodia growth in animals expressing

membrane tethered UNC-6. Interestingly, none of the transgenes were sufficient to

restore biased ventral branching (Figure 4F). This is consistent with a model where the

biased direction of axon growth after adhesion at the vSLNC intermediate target requires

diffusible UNC-6. Lastly, the frequency of axon stabilization at the VNC is partially rescued

only in animals with membrane tethered UNC-6 at both the vSLNC and VNC (Figure 4G).

This result suggests that VNC stabilization is contact-mediated, but its efficiency requires

UNC-6 at the intermediate target along earlier portions of the axon's path of growth. Taken

together, these data suggest that contact-mediated haptotaxis drives axon stabilization at

the vSLNC intermediate target and VNC, and that polarized UNC-40 towards the ventral

cell body and growth cone is required for adhesion.

Given the ability of non-diffusible UNC-6 constructs to restore contact-mediated

axon guidance behaviors, we considered their impact on the ultimate morphology of HSN

and PDE neurons. By assaying adult neuronal morphology, we found that only the double

transgene of membrane tethered UNC-6 was sufficient to partially restore adult axonal

morphology with 68.6% ± 5.0% of PDE axons and 47.4% ± 4.6% of HSN axons reaching

the VNC by adulthood (Tables 1,2). This is in comparison to 100% of PDE axons and

99.1% ± 0.9% of HSN axons doing so in wildtype animals and 32.1% ± 5.1% of PDE

axons and 33.1% ± 4.3% of HSN axons in *unc-6* mutants. These data indicate that

membrane-tethered UNC-6 can partially rescue axon guidance when present at both

vSLNC and VNC. This implies that diffusible UNC-6, possibly in the form of a gradient, is

also required for axon guidance of PDE and HSN.

UNC-6 is present in a shallow gradient

To further understand UNC-6's distribution and activity, we used an endogenous

C-terminus mNeonGreen (mNG) tag of UNC-6 (UNC-6::mNG) (Naegeli et al., 2017) to

visualize its localization *in vivo*. Our genetic evidence suggests that UNC-6 functions not

only locally at vSLNC and VNC but also likely over the entire distance between VNC and

vSLNC. We closely examined the localization of UNC-6::mNG in this region. Within the

zone between VNC and vSLNC, numerous faint UNC-6 puncta are present. Interestingly,

bright UNC-6 puncta are localized along the axonal filopodia of HSN. A similar localization

along the axonal filopodia is also observed for PDE and AVM (Figure 5A). Such signal is

completely absent in strains without the endogenous UNC-6::mNG tag, validating that the

signal represents endogenous UNC-6 (Figure S4). We measured the intensity of UNC-6

through this region along the ventrodorsal axis. The average intensity over this region is

the highest at the VNC. Moving dorsally, the average intensity steadily decreases but

increases again near the vSLNC (Figure 5B). These data are consistent with a model

where UNC-6 is enriched in both the vSLNC and VNC regions separated by a shallow

gradient.

Given the striking enrichment of UNC-6 along axonal filopodia, we hypothesized

that UNC-40 is required for UNC-6 clustering along the filopodia. Animals carrying *unc-*

40 null mutations display a dorsoventral gradient similar to wildtype animals (Figure

5C,D). However, axonal filopodia that extend toward the VNC are not associated with

UNC-6 puncta (Figure 5C). Taken together, these data suggest that UNC-6 is present in

a gradient and decorates the periphery of the growth cone during growth.

UNC-6 dynamically assemble around the growth cone during extension

The filopodia between vSLNC and VNC during axon outgrowth are dynamic, with

constant growth and retraction. This implies that the UNC-6 puncta associated with the

filopodia membrane are also likely not static. To directly test this, we simultaneously

imaged HSN growth cone and UNC-6::mNG puncta during its outgrowth in the region

between the VSLNC and VNC. Indeed, we observed dynamic assembly and disassembly

of mNG puncta near the growth cone and filopodia on ~1 min time scales (Figure 6A). In

movies of *unc-40* null animals, these foci assemble but quickly disperse and do not

associate with the growth cone (Figure 6A). To quantify this, we calculated the intensity

of each UNC-6 cluster adjacent to the cell membrane. These measurements reveal

decreased UNC-6 labeling around the cell in *unc-40* mutants compared to wildtype

animals (Figure 6B). Axonal growth was rarely observed in *unc-40* mutants within the

120s imaging window (Figure 6A), consistent with the reduced growth speed (Figure 2I).

Together, these data suggest that there is a diffusive population of UNC-6 in the axonal

path and that the extending growth cone remodels the UNC-6 landscape by assembling

transient, highly concentrated foci around the cell to stimulate growth via UNC-40.

Discussion

Here, we use time-lapse imaging of *in vivo* growth cone behaviors to demonstrate

that UNC-6/Netrin and UNC-40/DCC mediate both axon adhesion and directed growth.

UNC-6 and UNC-40/DCC are required for stepwise axon adhesion and growth. Axon

stabilization requires contact with UNC-6 which cooperates with the E3 ubiquitin ligase

MADD-2 to polarize UNC-40, providing adhesive traction upon reaching an intermediate

target. Biased growth requires UNC-6 and UNC-40 but cannot be explained by contact-

dependent UNC-6 signaling. Indeed, we observe a gradient of UNC-6 in the region of

filopodial extension. Growth through the gradient requires capturing of UNC-6 clusters by

UNC-40. These insights underscore the dynamic process of Netrin guidance as growth

cones sequentially utilize haptotaxis and chemotaxis (Figure 6C).

Complementary roles for haptotaxis and chemotaxis in UNC-6/Netrin guidance

Netrin has conserved functions in nervous system development across phylogeny.

Initially, Netrin was proposed to function as a chemotactic cue that stimulates directional

axon growth (Kennedy et al., 1994; Serafini et al., 1994). Indeed, secreted Netrin from

the vertebrate floor plate functions to attract axons toward the ventral midline, avoiding

the developing motor column (Wu et al., 2019). However, additional investigations in the

vertebrate spinal cord as well as in the *Drosophila* visual system and midline have

identified haptotactic roles for Netrin in axon guidance (Akin and Zipursky, 2016;

Brankatschk and Dickson, 2006; Varadarajan et al., 2017). Our data clearly demonstrate

roles for both chemotaxis and haptotaxis in Netrin-mediated attraction. Specifically,

haptotaxis (mediated by membrane-tethered UNC-6) is sufficient to explain growth cone

stabilization by providing adhesive force to maintain axonal contact with an intermediate

target. Conversely, chemotaxis likely provides the basis of directional growth toward the

VNC.

Importantly, the endogenous distribution of UNC-6/Netrin supports dual migration

mechanisms of haptotaxis and chemotaxis. In *C. elegans*, the ventral sublateral

intermediate target is separated from the VNC by a gradient of UNC-6 with the highest

concentration along the VNC. Yet, we also observe that this gradient is modified by an

accumulation of UNC-6 in the sublateral region, consistent with the location of the axon's

intermediate target. The precise mechanisms which produce elevated levels of UNC-6 in

the sublateral region remains unclear. As a laminin-related protein, Netrin is likely

interacting with extracellular matrix (ECM) proteins, and the enrichment of an ECM-

binding partner in the sublateral could restrain UNC-6 in such a way as to promote

haptotaxis. One candidate class of molecules is heparin sulfate proteoglycans which have

high affinity to UNC-6 (Serafini et al., 1994). In fact, structural analysis of the Netrin/DCC

interaction has identified a putative binding site dependent on extracellular sulfate (Finci

et al., 2014).

A recent study in *C. elegans* supports these conclusions using a transmembrane-

tethered allele of *unc-6* (Hooper and Lundquist, 2024). This allele displays a partial loss-

of-function phenotype that can be restored through UNC-6 produced by an ectopic

source. These data suggest that the diffusibility of UNC-6 is more critical than the location

of its transcription in *C. elegans*. Therefore, the gradient that we observe is likely

established using local interactions between UNC-6 and the surrounding ECM. Additional

studies combining cell biological and structural information may provide further insight on

UNC-6 interactions with the ECM and the dichotomous signaling mechanisms that

promote haptotactic and chemotactic responses.

Regulation and localization of UNC-40/DCC is crucial to haptotaxis

Haptotactic growth cone adhesion serves a central function in growth cone

migrations (Comer et al., 2019). Investigations of the molecular signaling mechanisms

that promote Netrin-driven haptotaxis have implicated an "adhesion-clutch" model" where

DCC generates force through connections to the actin cytoskeleton (Qiu et al., 2024). Our

data identify another molecular regulatory feature of haptotaxis: guidance receptor

localization. In ventrally migrating axons, endogenous levels of UNC-40 receptor are

polarized toward the ventral side of the cell upon contact with the sublateral intermediate

target. The E3 ubiquitin ligase MADD-2/Trim9/Trim67 cooperates with UNC-6 in this

polarization. These results are consistent with cell invasion mechanisms in the non-

neuronal anchor cell in *C. elegans* (Wang et al., 2014)*.* In our neuronal system, genetic

paradigms that disrupt UNC-40 polarization also exhibit decreased adherence to the

sublateral region. Therefore, we posit that polarization of UNC-40 likely amplifies the cell's

capacity to adhere to its intermediate target.

Notably, *Drosophila* photoreceptors depend on Trim9 for axon stabilization but not

elongation in Netrin guidance, consistent with our movies of *madd-2* mutant axon

guidance (Akin and Zipursky, 2016). Additionally, studies of vertebrate neurons

demonstrate that Trim9 ubiquitinates DCC when the concentration of Netrin is low to

regulate DCC exocytic balance (Plooster et al., 2017). Given the requirement of four

juxtamembrane lysine residues in UNC-40 polarization, our data suggest that this

regulatory function of MADD-2/Trim9 is conserved and crucial for UNC-40 cell polarity

and growth cone adhesion. While the data here do not directly examine cytoskeletal

regulation in relation to haptotaxis, Trim9 and Trim 67 are known to regulate members of

the Ena/VASP family through ubiquitination to control filopodia density in response to

Netrin-1 (Boyer et al., 2019; McCormick et al., 2024; Menon et al., 2015). Additionally, the

lone Ena/VASP family member in *C. elegans* (UNC-34) is required for ventral axon

guidance downstream of UNC-40 (Gitai et al., 2003; Norris et al., 2009). Systematic

investigations of the interplay between ubiquitination, receptor localization, and

cytoskeletal dynamics are required to fully elaborate haptotactic signaling mechanisms in

growth cones.

Collectively, our data provide spatiotemporal access to Netrin-mediated attraction. During

growth, the axon alternates between haptotactic and chemotactic migration, and we

delineate the precise axonal behaviors controlled by each migration mechanism. Overall,

this study predicates future investigations of the precise molecular cascades that govern

discrete chemotactic and haptotactic responses to Netrin within the same growth cone.

Acknowledgements

We thank the members of the Shen Lab for their feedback on the study and on the

manuscript. We particularly thank Richard Fetter, Callista Yee, Kelsie Eichel, and Dane

Kawano for their detailed input on the study and feedback on the manuscript. We also

thank Richard Fetter for Fig. 1D. This work was funded by the Howard Hughes Medical

Institute (where K.S. is an investigator) and the NSF Graduate Research Fellowship

Program (E.L.N.).

Author contributions

E.L.N. and K.S. conceived of the project. E.L.N. and J.L. generated all reagents and

carried out all experiments. E.L.N. performed all the analyses. E.L.N. wrote the

manuscript with editing from K.S. and J.L.

Declaration of interests

The authors have no competing interests to declare.

Figure legends

Figure 1. The ventral sublateral nerve cord (vSLNC) is an intermediate target for

ventrally growing axons. A. Diagram of *C. elegans* illustrating the anteroposterior

locations of the neurons which extend ventral axons in the larval stages: AVM, HSN, PDE,

PVM, and PQR. **B.** Confocal time-lapse images of PDE axon guidance. Images were taken every 3 min for 4 hr in the L2 stage. The final image is a confocal still image visualizing the mature PDE morphology in the L4 stage. Blue arrow denotes PDE soma. White text denotes PVD, PDE sheath glia (PDEsh), PDE socket glia (PDEso). Green arrowhead denotes the axonal pause in the sublateral region. Orange arrowhead indicates a dorsal branch, and blue arrowhead denotes a ventral branch. **C.** Cartoons showing lateral and cross-sectional views of ventral axon navigation in *C. elegans.* Ventral axons travel along the hypodermal edge until they reach the sublateral region. Further ventral growth requires navigation between the hypodermis and ventral body wall muscles. **D.** Electron microscopic image of PDE axon navigation path (blue). The dashed yellow box denotes the sublateral region where the axonal trajectory changes. This region is enlarged as an inset on the top right. Orange area denotes body wall muscle. **E.** Confocal images of AVM, HSN, PVM, and PQR axon emergence. The top row visualizes axonal contact with the vSLNC. The bottom row visualizes branching toward the VNC. Dashed green line denotes the vSLNC and the dashed gray line denotes the VNC. White arrow denotes PQR dendrite. Scale bar is 5 μm in **A, E** and 1 μm in **D.**

Figure 2. UNC-6 and UNC-40 promote stabilization and directed growth at the sublateral region. A. Confocal time-lapse images of PDE axons guidance in wildtype, *unc-6(ev400),* and *unc-40(e1430)* animals. Images were taken every 3m for 4hr in the L2 stage. Orange arrowhead indicates a dorsal branch, and blue arrowhead denotes a ventral branch. Dashed green line denotes the vSLNC and the dashed gray line denotes the VNC. **B.** Percentage of axons that stabilize in the sublateral region upon initial contact

in wildtype (n=23), *unc-6(ev400)* (n=18)*,* and *unc-40(e1430)* (n=21) animals. SE is shown. **C.** Airyscan images of PDE axon navigation and endogenous UNC-40::GFP growth cone localization in wildtype and *unc-6(ev400)* animals. Dashed white box denotes growth cone. **D.** Normalized intensity traces of growth cone UNC-40::GFP during axon navigation normalized to sublateral contact (t=0). Black line denotes wildtype animals (n=8), and red line denotes *unc-6(ev400)* animals (n=8). SEM is shown. **E.** Airyscan images of HSN axon emergence and endogenous UNC-40::GFP localization in wildtype and *unc-6(ev400)* animals. **F.** Cartoon schematic of polarity measurements and calculation in **G. G.** UNC-40::GFP polarity in HSN in wildtype (n=20) and *unc-6(ev400)* (n=21) animals. Positive values denote dorsal polarization, and negative values denote ventral polarization. SEM is shown **H.** Proportion of branches formed dorsally (orange) and ventrally (cyan) from the sublateral region in wildtype (n=90 branches, 24 animals), *unc-6(ev400)* (n=114 branches, 25 animals), and *unc-40(e1430)* (n=114 branches, 28 animals) animals. SE is shown. **I.** Representative traces of ventral branch extension in wildtype (black), *unc-6(ev400)* (magenta), and *unc-40(e1430)* (green) animals. **J.** Average ventral branch extension speed in wildtype (n=50 branches, 24 animals), *unc-6(ev400)* (n=43 branches, 25 animals), and *unc-40(e1430)* (n=32 branches, 28 animals) animals. SEM is shown. **K.** Percent of axons that stabilize that the VNC within the 4h imaging window in wildtype (n=24), *unc-6(ev400)* (n=25), and *unc-40(e1430)* (n=28) animals. SE is shown. Scale bars denote 5 μm in **A, C, E.** Fisher's exact test is used in **B, H, K.** Unpaired Student's t-test is used in **G.** Ordinary one-way ANOVA with multiple comparisons is used in **J.** * denotes $p<0.05$, ** denotes $p<0.01$, **** denotes $p<0.0001$.

Figure 3. MADD-2 promotes axon stabilization but not growth. A. Confocal time-lapse images of PDE axons guidance in wildtype and *madd-2(ok2226)* animals. Images were taken every 3m for 4hr in the L2 stage. Dashed green line denotes the vSLNC and the dashed gray line denotes the VNC. **B.** Percentage of axons that stabilize in the sublateral region upon initial contact in wildtype (n=23), *unc-6(ev400)* (n=18)*,* and *madd-2(ok2226)* (n=11) animals. SE is shown. **C.** Airyscan images of HSN axon emergence and endogenous UNC-40::GFP localization in wildtype, *madd-2(ok2226), madd-2(ok2226); unc-6(ev400),* and *unc-40(4KR)* animals. **D.** UNC-40::GFP polarity in HSN in wildtype (n=20), *unc-6(ev400)* (n=21), *madd-2(ok2226)* (n=23), and *unc-6(ev400); madd-2(ok2226)* (n=19) animals. Positive values denote dorsal polarization, and negative values denote ventral polarization. SEM is shown. **E.** Cartoon schematic of the UNC-40 protein sequence. The dashed red box denotes the intracellular lysine residues that are mutated in *unc-40(4KR)* animals. **F.** UNC-40::GFP polarity in HSN in wildtype (n=20), *madd-2(ok2226)* (n=23), and *unc-40(4KR)* (n=21) animals. Positive values denote dorsal polarization, and negative values denote ventral polarization. SEM is shown. **G.** Proportion of branches formed dorsally (orange) and ventrally (cyan) from the sublateral region in wildtype (n=90 branches, 24 animals), *unc-6(ev400)* (n=114 branches, 25 animals), and *madd-2(ok2226)* (n=52 branches, 12 animals) animals. SE is shown. **H.** Percent of axons that stabilize that the VNC within the 4h imaging window in wildtype (n=24), *unc-6(ev400)* (n=25), and *madd-2(ok2226)* (n=12) animals. SE is shown. Scale bars denote 5 μm in **A,C.** Fisher's exact test is used in **B,G,H.** Ordinary one-way ANOVA with multiple comparisons is used in **D,F.** * denotes $p < 0.05$, *** denotes $p < 0.001$, **** denotes $p < 0.0001$.

Figure 4. Diffusible UNC-6 is dispensable for axon stabilization. A. Cartoon

schematic of the non-diffusible rescue paradigms at the vSLNC (green, using *Pmec-7)*

and at the VNC (magenta, using *Punc-4*). **B.** Confocal time-lapse images of PDE axons

guidance in wildtype, *unc-6(ev400); Pmec-7::unc-6^{TM}, unc-6(ev400); Punc-4::unc-6^{TM}*

and *unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}* animals. Images were taken every

3m for 4hr in the L2 stage. Green lines denote the vSLNC and gray lines denote the VNC.

Solid lines denote the location of membrane-tethered UNC-6. **C.** Percentage of axons

that stabilize in the sublateral region upon initial contact in wildtype (n=23), *unc-6(ev400)*

(n=18)*, unc-6(ev400); Pmec-7::unc-6^{TM}* (n=10)*, unc-6(ev400); Punc-4::unc-6^{TM}* (n=8) and

unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM} (n=7) animals. SE is shown. **D.**

Airyscan images of HSN axon emergence and endogenous UNC-40::GFP localization in

wildtype, *unc-6(ev400), unc-6(ev400); Pmec-7::unc-6^{TM},* and *unc-6(ev400); Punc-4::unc-*

6^{TM} animals. **E.** UNC-40::GFP polarity in HSN in wildtype (n=20), *unc-6(ev400)* (n=21)*,*

unc-6(ev400); Pmec-7::unc-6^{TM} (n=25)*,* and *unc-6(ev400); Punc-4::unc-6^{TM}* (n=18)

animals. Positive values denote dorsal polarization, and negative values denote ventral

polarization. SEM is shown. **F.** Proportion of branches formed dorsally (orange) and

ventrally (cyan) from the sublateral region in wildtype (n=90 branches, 24 animals), *unc-*

6(ev400) (n=114 branches, 25 animals), *unc-6(ev400); Pmec-7::unc-6^{TM}* (n=105

branches, 14 animals)*, unc-6(ev400); Punc-4::unc-6^{TM}* (n=83 branches, 10 animals), and

unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM} (n=108 branches, 15 animals) animals.

SE is shown. **G.** Percent of axons that stabilize that the VNC within the 4h imaging window

in wildtype (n=24), *unc-6(ev400)* (n=25), *unc-6(ev400); Pmec-7::unc-6^{TM}* (n=14)*, unc-*

6(ev400); Punc-4::unc-6^{TM} (n=10), and *unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}*

(n=15) animals. SE is shown. Scale bars denote 5 μm in **B,D.** Fisher's exact test is used

in **C,F, G.** Ordinary one-way ANOVA with multiple comparisons is used in **E.** * denotes

$p<0.05$, ** denotes $p<0.01$, **** denotes $p<0.0001$.

Figure 5. UNC-6 is present in a gradient. A. Airyscan images of axon extension in HSN,

PDE, and AVM and endogenous UNC-6::mNG distribution in wildtype animals. White

arrowheads denote UNC-6 clusters around axonal filopodia. Dashed green line denotes

vSLNC and dashed gray line denotes VNC. **B.** Average intensity of UNC-6::mNG between

the VNC (left) and vSLNC (right) in wildtype animals (n=49 animals). **C** Airyscan images

of axon extension in HSN, PDE, and AVM and endogenous UNC-6::mNG distribution in

unc-40(e1430) animals. Dashed green line denotes vSLNC and dashed gray line denotes

VNC. **D.** Average intensity of UNC-6::mNG between the VNC (left) and vSLNC (right) in

unc-40(e1430) animals (n=37 animals). SEM is shown. Scale bars denote 5 μm in **A,C.**

Figure 6. UNC-6 clusters are dynamically assembled at the growth cone tip during

axon extension. A. Airyscan time-lapse images of HSN axon extension and endogenous

UNC-6::mNG in wildtype and *unc-40(e1430)* animals. Images were taken every 30s for

30min. White arrowhead denotes the growth cone tip. Dashed white circle denotes UNC-

6 clusters on the lateral edges of the growth cone. **B.** Intensity of UNC-6 foci adjacent to

the growth cone membrane in wildtype (n=861 foci, 5 animals) and *unc-40(e1430)*

(n=1306 foci, 5 animals) animals. **C.** Model of ventral axon migration via alternating

haptotaxis and chemotaxis. Scale bar denotes 1 μm in **A.** Unpaired Student's t-test is

used in **B.** **** denotes $p<0.0001$.

Tables

Table 1. Genetic analysis of PDE axon morphology in adulthood. "% of axons at

VNC" denotes percentage of PDE neurons with axons that extend to the ventral nerve

cord (VNC). "% of axons not at VNC" denotes percentage of PDE neurons with axons

that do not extend to the ventral nerve cord (VNC). N indicates number of animals scored.

P-values generated from Fisher's exact test.

Genotype	% of Axons at VNC	% of Axons not at VNC	Standard Error	n	Comparison with wildtype	Comparison with *unc-6(ev400)*
wildtype	100.0%	0.0%	0.0%	118	■■■■	p<0.0001
unc-6(ev400)	32.1%	67.9%	5.1%	84	p<0.0001	■■■■
unc-40(e1430)	33.0%	67.0%	4.8%	97	p<0.0001	p>0.9999
unc-6(ev400); Pmec-7::unc-6^{TM}	31.6%	68.4%	4.8%	95	p<0.0001	p>0.9999
unc-6(ev400); Punc-4::unc-6^{TM}	35.2%	64.8%	5.1%	88	p<0.0001	p=0.7475
unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}	68.6%	31.4%	5.0%	86	p<0.0001	p<0.0001
madd-2(ok2226)	94.7%	5.3%	2.3%	95	p=0.0166	p<0.0001
unc-6(ev400); madd-2(ok2226)	43.6%	56.4%	4.9%	101	p<0.0001	p=0.1299
unc-40(4KR)	94.9%	5.1%	2.2%	99	p=0.0187	p<0.0001

11   **Table 2. Genetic analysis of HSN axon morphology in adulthood.** "% of axons at

12   VNC" denotes percentage of HSN neurons with axons that extend to the ventral nerve

13   cord (VNC). "% of axons not at VNC" denotes percentage of HSN neurons with axons

14   that do not extend to the ventral nerve cord (VNC). N indicates number of animals scored.

15   P-values generated from Fisher's exact test.

Genotype	% of Axons at VNC	% of Axons not at VNC	Standard Error	n	Comparison with wildtype	Comparison with *unc-6(ev400)*
wildtype	99.1%	0.91%	0.90%	110		p<0.0001
unc-6(ev400)	33.1%	66.9%	4.3%	121	p<0.0001	
unc-40(e1430)	30.6%	69.4%	4.1%	124	p<0.0001	p=0.7840
unc-6(ev400); Pmec-7::unc-6^{TM}	33.3%	66.7%	4.5%	108	p<0.0001	p>0.9999
unc-6(ev400); Punc-4::unc-6^{TM}	29.5%	70.5%	4.5%	105	p<0.0001	p=0.6666
unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}	47.4%	52.6%	4.6%	116	p<0.0001	p=0.0252
madd-2(ok2226)	92.2%	7.8%	2.5%	115	p=0.0191	p<0.0001
unc-6(ev400); madd-2(ok2226)	44.4%	55.6%	4.5%	124	p<0.0001	p=0.0881
unc-40(4KR)	91.9%	8.1%	2.4%	124	p=0.0114	p<0.0001

STAR Methods

Resource Availability

Lead contact. Further information and requests for resources and reagents should be

directed to and will be fulfilled by the lead contact, Kang Shen (kangshen@stanford.edu).

Materials availability. All materials used and generated in this study, including plasmids

and worm strains, are available for sharing by contacting Kang Shen

(kangshen@stanford.edu).

Data and code availability. The published article includes all generated or analyzed during

this study. Raw data generated for this study are available from Kang Shen

(kangshen@stanford.edu) upon request. Any additional information required to reanalyze

the data reported in this paper are available upon request. This paper does not report

original code.

Experimental model and subject details

<u>C. elegans *methods.*</u> All *C. elegans* animals used in this study were handled at 20°C on

NGM plates with OP50 *E. coli* as food, according to standard protocols (Brenner, 1974).

N2 Bristol was used as a wildtype reference strain. Transgenic strains were generated

via microinjection of 1-5 ng/μl of plasmid DNA and of 1 ng/μl of selection markers plasmids

(*Pmyo-2::mCherry*). Pharyngeal mCherry signal was used to identify transgenic worms.

All analyses were performed on *C. elegans* hermaphrodites. Imaging of axon outgrowth

used animals aged to the L2 stage. Imaging of mature neuronal morphologies used

animals aged to the L4 and young adult stages. Detailed description of the strains used

in this study are available in the Key Resources Table.

Method details

<u>*Confocal imaging.*</u> All images were taken at room temperature using live *C. elegans*. L2

animals were anesthetized using 6mM levamisole in M9 buffer and mounted on 4%

agarose pads. For still images, worms were imaged on an inverted Zeiss Axio Observer

Z1 microscope with a Yokogawa CSU-X1 spinning-disk unit, a Hamamatsu EM-CCD

digital camera controlled by Metamorph (version 7.8.12.0). Images were acquired using

a Plan-Apochromat 100x 1.4 NA objective. For dynamic imaging of axon outgrowth,

worms were imaged on an inverted Zeiss Axio Observer Z1 microscope with a Yokogawa

CSU-W1 spinning-disk unit, a Prime 95B Scientific CMOS camera controlled by 3i

Slidebook (v6). Images were acquired using a C-Apochromat 63x 1.2 NA objective.

Images were acquired every 3min for 4hr. For long-term dynamic imaging, the coverslip

was sealed on all sides using valap to prevent sample dehydration.

Airyscan imaging. All images were taken at room temperature using live _C. elegans_. L2

animals were anesthetized using 6mM levamisole in M9 buffer and mounted on 4%

agarose pads. Worms were imaged on a Zeiss LSM 980 Airyscan 2 system controlled by

Zen (version 3.5 blue edition). Images were acquired using a Plan-Apochromat 63x 1.4

NA objective. For dynamic imaging of UNC-6::mNG, images were acquired every 30sec

for 30min.

Constructs and cloning. All constructs used in this study are listed in the Key Resources

Table. Plasmid constructs generated for this study were created using isothermal

assembly with overlapping oligonucleotides (Gibson et al., 2009). Expression constructs

were generated in a pSM delta vector. Non-diffusible UNC-6 constructs were made by

fusing the _nlg-1_ transmembrane domain sequence to the C-terminus of the _unc-6_ coding

sequence. A SNAP tag was fused to the intracellular side of the _nlg-1_ transmembrane

domain.

Germline editing using CRISPR/Cas9. Endogenous fluorophore insertions and base pair

changes were generated by gonadal microinjection of CRISPR/Cas9 protein complexes.

Genome editing was performed using standard protocols (Dokshin et al., 2018). Cas9

protein, tracRNA, crRNAs were injected at a concentration of 1.525 μM. crRNAs used in

this study are available in the Key Resources Table. DNA repair templates were created

by PCR or ordered as ultramers and injected at 2 μM. pRF4 (_rol-6(su1006)_) was used as

a co-injection marker and selected against in the F_2 generation. F_2 animals were screened

for the desired edits using PCR and confirmed by sequencing. Newly generated strains

with the proper edits were outcrossed 5 times before use in experiments.

Electron microscopy. N2 Bristol worms staged to L4 were prepared for conventional EM

by high pressure freezing/freeze-substitution. Worms were suspended in M9 containing

20% BSA and _E. coli_ and frozen in 100 μm well carriers (Type A) across from a

hexadecane-coated flat carrier (Type B) with a BalTec HPM 01 high-pressure freezer. A

Leica AFS2 unit was used for freeze-substitution in 1% OsO_4, 0.1% uranyl acetate, 1%

methanol in acetone, and 3% water. After substitution, the samples were rinsed in

acetone, infiltrated and polymerized in Eponate 12 resin. A Leica UCT ultramicrotome with

a Diatome diamond knife was used to cut serial 50 nm sections. The sections were picked

up on Pioloform film coated slot grids and stained with uranyl acetate and Sato's lead. An

FEI Tecnai T12 TEM at 120 kV with a Gatan 4k x 4k U895 camera was used to image the

sections. Serial sections were aligned using TrakEM2 in Fiji.

UNC-40::GFP visualization. Images of UNC-40::GFP in single cells were generated by

masking using Fiji. In short, the image channel of the cell membrane marker was used to

generate a binary mask. This binary mask was applied to the UNC-40::GFP channel,

removing fluorescence not associated with the cell membrane mask and retaining the

original intensity values of UNC-40::GFP. This protocol was used to generate all of the

UNC-40::GFP images in this manuscript.

Quantification and statistical analysis

UNC-40::GFP polarity measurements. Fiji was used to measure the integrated

fluorescent density along the cell dorsal and ventral periphery in single z-planes as

visualized in Figure 2F. The ventral integrated density was subtracted from the dorsal

integrated density, and this value was divided by the total integrated density in the ventral

and dorsal regions. This calculated value is reported as the polarity of the fluorescent

marker. All measurements were taken in raw images prior to processing of UNC-40::GFP

for visualization as described above.

Analysis of dynamic imaging of axon growth. Fiji was used to analyze movies of axon

outgrowth. Growth cone and UNC-40::GFP tracings were completed using the MTrackJ

plugin on Fiji. Intensity values were normalized to the traced intensity in the growth cone

upon contact with the sublateral region. All other analyses were reported by manual

counting or length measurements. Dorsal and ventral branching were determined by

protrusions longer than 0.6 μm. Growth speed was calculated as the total displacement

of the branch divided by its duration in the movies.

Analysis of dynamic imaging of UNC-6::mNG. Fiji was used to analyze movies of UNC-

6::mNG dynamics. First, UNC-6::mNG foci overlapping with the cell membrane marker

were isolated via the masking method described above for UNC-40::GFP, retaining the

UNC-6::mNG intensity values. Next, z-projections were made of the movies showing

UNC-6::mNG contacts with the cell membrane. Last, the Analyze Particles analysis on

Fiji was run to generate the average intensity of each identified UNC-6::mNG cluster.

Data visualization and statistical analysis. Plotly (via Python) was used to generate the

roseplots. GraphPad Prism 10 was used for all other data visualization and statistical

analyses. SEM and SE are shown as indicated in figure legends. Statistical analyses used

are also available in figure legends.

Supplemental information

Document S1. Figures S1-S4.

Table S1. Summary of *C. elegans* strains used in this study, related to STAR Methods.

Video S1. Comparison of PDE axon guidance in wildtype and *unc-6(ev400)* mutants,

related to Figure 2.

References

Adler, C.E., Fetter, R.D., Bargmann, C.I., 2006. UNC-6/Netrin induces neuronal
asymmetry and defines the site of axon formation. Nat. Neurosci. 9, 511–518.
https://doi.org/10.1038/nn1666
Akin, O., Zipursky, S.L., 2016. Frazzled promotes growth cone attachment at the source
of a Netrin gradient in the Drosophila visual system. eLife 5, e20762.
https://doi.org/10.7554/eLife.20762
Bentley, D., Caudy, M., 1983a. Pioneer axons lose directed growth after selective killing
of guidepost cells. Nature 304, 62–65. https://doi.org/10.1038/304062a0
Bentley, D., Caudy, M., 1983b. Navigational Substrates for Peripheral Pioneer Growth
Cones: Limb-axis Polarity Cues, Limb-segment Boundaries, and Guidepost
Neurons. Cold Spring Harb. Symp. Quant. Biol. 48, 573–585.
https://doi.org/10.1101/SQB.1983.048.01.062
Bernhardt, R.R., Nguyen, N., Kuwada, J.Y., 1992. Growth cone guidance by floor plate
cells in the spinal cord of zebrafish embryos. Neuron 8, 869–882.
https://doi.org/10.1016/0896-6273(92)90201-N
Boyer, N.P., McCormick, L.E., Menon, S., Urbina, F.L., Gupton, S.L., 2019. A pair of E3
ubiquitin ligases compete to regulate filopodial dynamics and axon guidance. J.
Cell Biol. 219, e201902088. https://doi.org/10.1083/jcb.201902088
Brankatschk, M., Dickson, B.J., 2006. Netrins guide Drosophila commissural axons at
short range. Nat. Neurosci. 9, 188–194. https://doi.org/10.1038/nn1625
Brenner, S., 1974. THE GENETICS OF CAENORHABDITIS ELEGANS. Genetics 77,
71–94. https://doi.org/10.1093/genetics/77.1.71

Chalfie, M., Sulston, J., 1981. Developmental genetics of the mechanosensory neurons of *Caenorhabditis elegans*. Dev. Biol. 82, 358–370. https://doi.org/10.1016/0012-1606(81)90459-0

Chisholm, A.D., Hutter, H., Jin, Y., Wadsworth, W.G., 2016. The Genetics of Axon Guidance and Axon Regeneration in Caenorhabditis elegans. Genetics 204, 849–882. https://doi.org/10.1534/genetics.115.186262

Chung, K.Y., Taylor, J.S.H., Shum, D.K.Y., Chan, S.O., 2000. Axon routing at the optic chiasm after enzymatic removal of chondroitin sulfate in mouse embryos. Development 127, 2673–2683. https://doi.org/10.1242/dev.127.12.2673

Comer, J.D., Alvarez, S., Butler, S.J., Kaltschmidt, J.A., 2019. Commissural axon guidance in the developing spinal cord: from Cajal to the present day. Neural Develop. 14, 9. https://doi.org/10.1186/s13064-019-0133-1

Cox, E.A., Hardin, J., 2004. Sticky worms: adhesion complexes in C. elegans. J. Cell Sci. 117, 1885–1897. https://doi.org/10.1242/jcs.01176

Dokshin, G.A., Ghanta, K.S., Piscopo, K.M., Mello, C.C., 2018. Robust Genome Editing with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in Caenorhabditis elegans. Genetics 210, 781–787. https://doi.org/10.1534/genetics.118.301532

Dominici, C., Moreno-Bravo, J.A., Puiggros, S.R., Rappeneau, Q., Rama, N., Vieugue, P., Bernet, A., Mehlen, P., Chédotal, A., 2017. Floor-plate-derived netrin-1 is dispensable for commissural axon guidance. Nature 545, 350–354. https://doi.org/10.1038/nature22331

Finci, L.I., Krüger, N., Sun, X., Zhang, J., Chegkazi, M., Wu, Y., Schenk, G., Mertens, H.D.T., Svergun, D.I., Zhang, Y., Wang, J., Meijers, R., 2014. The Crystal Structure of Netrin-1 in Complex with DCC Reveals the Bifunctionality of Netrin-1 As a Guidance Cue. Neuron 83, 839–849. https://doi.org/10.1016/j.neuron.2014.07.010

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., Smith, H.O., 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods 6, 343–345. https://doi.org/10.1038/nmeth.1318

Gitai, Z., Yu, T.W., Lundquist, E.A., Tessier-Lavigne, M., Bargmann, C.I., 2003. The Netrin Receptor UNC-40/DCC Stimulates Axon Attraction and Outgrowth through Enabled and, in Parallel, Rac and UNC-115/AbLIM. Neuron 37, 53–65. https://doi.org/10.1016/S0896-6273(02)01149-2

Hao, J.C., Adler, C.E., Mebane, L., Gertler, F.B., Bargmann, C.I., Tessier-Lavigne, M., 2010. The Tripartite Motif Protein MADD-2 Functions with the Receptor UNC-40 (DCC) in Netrin-Mediated Axon Attraction and Branching. Dev. Cell 18, 950–960. https://doi.org/10.1016/j.devcel.2010.02.019

Hedgecock, E.M., Culotti, J.G., Hall, D.H., 1990. The unc-5, unc-6, and unc-40 genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in C. elegans. Neuron 4, 61–85. https://doi.org/10.1016/0896-6273(90)90444-K

Hooper, K.M., Lundquist, E.A., 2024. Short- and long-range roles of UNC-6/Netrin in dorsal-ventral axon guidance in vivo in Caenorhabditis elegans. https://doi.org/10.1101/2024.04.23.590737

Jayadev, R., Morais, M.R.P.T., Ellingford, J.M., Srinivasan, S., Naylor, R.W., Lawless, C., Li, A.S., Ingham, J.F., Hastie, E., Chi, Q., Fresquet, M., Koudis, N.-M., Thomas, H.B., O'Keefe, R.T., Williams, E., Adamson, A., Stuart, H.M., Banka, S., Smedley, D., Genomics England Research Consortium, Sherwood, D.R., Lennon, R., 2022. A basement membrane discovery pipeline uncovers network complexity, regulators, and human disease associations. Sci. Adv. 8, eabn2265. https://doi.org/10.1126/sciadv.abn2265

Kennedy, T.E., Serafini, T., de la Torre, JoséR., Tessier-Lavigne, M., 1994. Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. Cell 78, 425–435. https://doi.org/10.1016/0092-8674(94)90421-9

Kennedy, T.E., Wang, H., Marshall, W., Tessier-Lavigne, M., 2006. Axon Guidance by Diffusible Chemoattractants: A Gradient of Netrin Protein in the Developing Spinal Cord. J. Neurosci. 26, 8866–8874. https://doi.org/10.1523/JNEUROSCI.5191-05.2006

Keshishian, H., Bentley, D., 1983. Embryogenesis of peripheral nerve pathways in grasshopper legs: II. The major nerve routes. Dev. Biol. 96, 103–115. https://doi.org/10.1016/0012-1606(83)90315-9

Klämbt, C., Jacobs, J.R., Goodman, C.S., 1991. The midline of the drosophila central nervous system: A model for the genetic analysis of cell fate, cell migration, and growth cone guidance. Cell 64, 801–815. https://doi.org/10.1016/0092-8674(91)90509-W

Knobel, K.M., Jorgensen, E.M., Bastiani, M.J., 1999. Growth cones stall and collapse during axon outgrowth in Caenorhabditis elegans. Development 126, 4489–4498. https://doi.org/10.1242/dev.126.20.4489

Kolodkin, A.L., Tessier-Lavigne, M., 2011. Mechanisms and Molecules of Neuronal Wiring: A Primer. Cold Spring Harb. Perspect. Biol. 3, a001727. https://doi.org/10.1101/cshperspect.a001727

Lee, J., Magescas, J., Fetter, R.D., Feldman, J.L., Shen, K., 2021. Inherited apicobasal polarity defines the key features of axon-dendrite polarity in a sensory neuron. Curr. Biol. 31, 3768-3783.e3. https://doi.org/10.1016/j.cub.2021.06.039

Li, H., Kulkarni, G., Wadsworth, W.G., 2008. RPM-1, a Caenorhabditis elegans Protein That Functions in Presynaptic Differentiation, Negatively Regulates Axon Outgrowth by Controlling SAX-3/robo and UNC-5/UNC5 Activity. J. Neurosci. 28, 3595–3603. https://doi.org/10.1523/JNEUROSCI.5536-07.2008

McCormick, L.E., Suarez, C., Herring, L.E., Cannon, K.S., Kovar, D.R., Brown, N.G., Gupton, S.L., 2024. Multi-monoubiquitylation controls VASP-mediated actin dynamics. J. Cell Sci. 137, jcs261527. https://doi.org/10.1242/jcs.261527

Menon, S., Boyer, N.P., Winkle, C.C., McClain, L.M., Hanlin, C.C., Pandey, D., Rothenfußer, S., Taylor, A.M., Gupton, S.L., 2015. The E3 ubiquitin ligase TRIM9 is a filopodia off switch required for netrin dependent axon guidance. Dev. Cell 35, 698–712. https://doi.org/10.1016/j.devcel.2015.11.022

Naegeli, K.M., Hastie, E., Garde, A., Wang, Z., Keeley, D.P., Gordon, K.L., Pani, A.M., Kelley, L.C., Morrissey, M.A., Chi, Q., Goldstein, B., Sherwood, D.R., 2017. Cell invasion in vivo via rapid exocytosis of a transient lysosome-derived membrane domain. Dev. Cell 43, 403-417.e10. https://doi.org/10.1016/j.devcel.2017.10.024

Norris, A.D., Dyer, J.O., Lundquist, E.A., 2009. The Arp2/3 complex, UNC-115/abLIM, and UNC-34/Enabled regulate axon guidance and growth cone filopodia formation in Caenorhabditis elegans. Neural Develop. 4, 38. https://doi.org/10.1186/1749-8104-4-38

Plooster, M., Menon, S., Winkle, C.C., Urbina, F.L., Monkiewicz, C., Phend, K.D., Weinberg, R.J., Gupton, S.L., 2017. TRIM9-dependent ubiquitination of DCC constrains kinase signaling, exocytosis, and axon branching. Mol. Biol. Cell 28, 2374–2385. https://doi.org/10.1091/mbc.E16-08-0594

Qiu, Z., Minegishi, T., Aoki, D., Abe, K., Baba, K., Inagaki, N., 2024. Adhesion-clutch between DCC and netrin-1 mediates netrin-1–induced axonal haptotaxis. Front. Mol. Neurosci. 17, 1307755. https://doi.org/10.3389/fnmol.2024.1307755

Ramón y Cajal, S., 1911. Histology of the Nervous System of Man and Vertebrates. Oxford University Press, New York.

Serafini, T., Kennedy, T.E., Gaiko, M.J., Mirzayan, C., Jessell, T.M., Tessier-Lavigne, M., 1994. The netrins define a family of axon outgrowth-promoting proteins homologous to C. elegans UNC-6. Cell 78, 409–424. https://doi.org/10.1016/0092-8674(94)90420-0

Teichmann, H.M., Shen, K., 2011. UNC-6 and UNC-40 promote dendritic growth through PAR-4 in Caenorhabditis elegans neurons. Nat. Neurosci. 14, 165–172. https://doi.org/10.1038/nn.2717

Timofeev, K., Joly, W., Hadjieconomou, D., Salecker, I., 2012. Localized Netrins Act as Positional Cues to Control Layer-Specific Targeting of Photoreceptor Axons in *Drosophila*. Neuron 75, 80–93. https://doi.org/10.1016/j.neuron.2012.04.037

Varadarajan, S.G., Kong, J.H., Phan, K.D., Kao, T.-J., Panaitof, S.C., Cardin, J., Eltzschig, H., Kania, A., Novitch, B.G., Butler, S.J., 2017. Netrin1 Produced by Neural Progenitors, Not Floor Plate Cells, Is Required for Axon Guidance in the Spinal Cord. Neuron 94, 790-799.e3. https://doi.org/10.1016/j.neuron.2017.03.007

Wadsworth, W.G., Bhatt, H., Hedgecock, E.M., 1996. Neuroglia and Pioneer Neurons Express UNC-6 to Provide Global and Local Netrin Cues for Guiding Migrations in C. elegans. Neuron 16, 35–46. https://doi.org/10.1016/S0896-6273(00)80021-5

Wang, Z., Linden, L.M., Naegeli, K.M., Ziel, J.W., Chi, Q., Hagedorn, E.J., Savage, N.S., Sherwood, D.R., 2014. UNC-6 (netrin) stabilizes oscillatory clustering of the UNC-40 (DCC) receptor to orient polarity. J. Cell Biol. 206, 619–633. https://doi.org/10.1083/jcb.201405026

Wu, Z., Makihara, S., Yam, P.T., Teo, S., Renier, N., Balekoglu, N., Moreno-Bravo, J.A., Olsen, O., Chédotal, A., Charron, F., Tessier-Lavigne, M., 2019. Long-Range Guidance of Spinal Commissural Axons by Netrin1 and Sonic Hedgehog from Midline Floor Plate Cells. Neuron 101, 635-647.e4. https://doi.org/10.1016/j.neuron.2018.12.025

Yung, A.R., Druckenbrod, N.R., Cloutier, J.-F., Wu, Z., Tessier-Lavigne, M., Goodrich, L.V., 2018. Netrin-1 Confines Rhombic Lip-Derived Neurons to the CNS. Cell Rep. 22, 1666–1680. https://doi.org/10.1016/j.celrep.2018.01.068

Figure 1

A

AVM HSN

PDE
PVM
PQR

Dorsal
Anterior — Posterior
Ventral

B

PDEsh
PDEso
PVD
PDE
5μm
0 min

Axon emergence
18 min

Axon pause
near vSLNC
33 min

Branching
from vSLNC
72 min

Stabilization
at VNC
105 min

Adult morphology
5μm

C

Lateral view (shown in B)

emerging PDE axon

ventral sublateral
nerve cord (vSLNC)

body wall muscle

ventral axon
trajectory

ventral nerve cord (VNC)

Cross-section (shown in D)

epidermis

Dorsal
Left — Right
Ventral

emerging
PDE axon

ventral sublateral nerve
cords (vSLNCs)

body wall muscles

ventral axon trajectories

ventral nerve cords (VNCs)

D

Ventral axon trajectory

1μm

1μm

E

AVM   HSN   PVM   PQR

Pausing
at sublateral
5μm

Growth
to VNC

dendrite

Figure 2

A (wildtype) 0 min, 21 min, 60 min, 108 min; *unc-6(ev400)* 0 min, 21 min, 45 min, 72 min; *unc-40(e1430)* 0 min, 24 min, 66 min, 72 min. Scale bar 5μm.

B Percent of axons that stabilize after initial vsLNC contact: wildtype, *unc-6(ev400)* *, *unc-40(e1430)* **.

D Normalized UNC-40::GFP intensity (au) in PDE growth cone vs time (min) relative to vSLNC contact. wildtype (black), *unc-6(ev400)* (red).

C PDE UNC-40::GFP — pre-sublateral, at sublateral; wildtype, *unc-6(ev400)*. Scale bar 5μm.

E HSN UNC-40::GFP / UNC-40::GFP; wildtype, *unc-6(ev400)*. Scale bar 5μm.

F dorsal cell body / HSN / vSLNC contact site.

$$polarity = \frac{\left(\begin{array}{c}\text{sum intensity of} \\ \text{dorsal cell body}\end{array} - \begin{array}{c}\text{sum intensity of} \\ \text{vSLNC contact site}\end{array}\right)}{\text{sum intensity of dorsal cell body and vSLNC contact site}}$$

G UNC-40::GFP polarity in HSN: wildtype, *unc-6(ev400)* ****. more dorsal / more ventral.

H Percent of axon branching — Dorsal branch (orange), Ventral branch (cyan): wildtype, *unc-6(ev400)* *, *unc-40(e1430)* ***.

I branch length (μm) vs time (min), VNC to vSLNC: wildtype (black), *unc-6(ev400)* (magenta), *unc-40(e1430)* (green).

J Average ventral branch speed (μm/min): wildtype, *unc-6(ev400)* **, *unc-40(e1430)* **.

K Percent of axons that stabilize at the VNC within 4h: wildtype, *unc-6(ev400)* ****, *unc-40(e1430)* ****.

Figure 3

A wildtype
0 min 27 min 42 min 78 min
5µm

madd-2(ok2226)
0 min 24 min 48 min 69 min

B Percent of axons that stabilize after initial vsLNC contact
100%
80%
60%
40%
20%
0%
wildtype *unc-6(ev400)* *madd-2(ok2226)*

C HSN UNC-40::GFP // UNC-40::GFP
wildtype   *madd-2(ok2226); unc-6(ev400)*
5µm
madd-2(ok2226)   *unc-40(4KR)*

D UNC-40::GFP polarity in HSN
1.0
0.5
0.0
-0.5
-1.0
more dorsal / more ventral
wildtype *madd-2(ok2226)* *unc-6(ev400)* *madd-2(ok2226); unc-6(ev400)*
**** **** * *

E UNC-40
dashed red box denotes juxtamembrane lysine residues mutated in *unc-40(4KR)* animals
intracellular lysine residues
extracellular domain
transmembrane domain
intracellular domain

F UNC-40::GFP polarity in HSN
1.0
0.5
0.0
-0.5
-1.0
more dorsal / more ventral
wildtype *madd-2(ok2226)* *unc-40(4KR)*
*** ****

G Percent of axon branching
Dorsal branch Ventral branch
100%
80%
60%
40%
20%
0%
wildtype *unc-6(ev400)* *madd-2(ok2226)*
**

H Percent of axons that stabilize at the VNC within 4h
100%
80%
60%
40%
20%
0%
wildtype *unc-6(ev400)* *madd-2(ok2226)*
**** *

Figure 4

A
Dorsal | Ventral
Pmec-7 tethered UNC-6
Punc-4 tethered UNC-6

B
wildtype
0 min | 24 min | 45 min | 72 min
5µm

unc-6(ev400); Pmec-7::unc-6^{TM}
0 min | 27 min | 54 min | 87 min

unc-6(ev400); Punc-4::unc-6^{TM}
0 min | 30 min | 57 min | 87 min

unc-6(ev400); Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}
0 min | 30 min | 84 min | 114 min

C
Percent of axons that stabilize after initial vsLNC contact
wildtype, unc-6(ev400), Pmec-7::unc-6^{TM}, Punc-4::unc-6^{TM}, Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}
unc-6(ev400)

D
HSN
UNC-40::GFP | UNC-40::GFP
wildtype
5µm
unc-6(ev400); Pmec-7::unc-6^{TM}
unc-6(ev400); Punc-4::unc-6^{TM}

E
UNC-40::GFP polarity in HSN
**** | ns | ****
more dorsal | more ventral
wildtype, unc-6(ev400), Pmec-7::unc-6^{TM}, Punc-4::unc-6^{TM}
unc-6(ev400)

F
Dorsal branch | Ventral branch
Percent of axon branching
wildtype, unc-6(ev400), Pmec-7::unc-6^{TM}, Punc-4::unc-6^{TM}, Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}
unc-6(ev400)

G
Percent of axons that stabilize at the VNC within 4h
wildtype, unc-6(ev400), Pmec-7::unc-6^{TM}, Punc-4::unc-6^{TM}, Pmec-7::unc-6^{TM}; Punc-4::unc-6^{TM}
unc-6(ev400)

Figure 5

A wildtype

HSN   PDE   AVM

Endogenous UNC-6::mNG
High
Low

Fasciculation at sublateral

5μm

Branching to VNC

C *unc-40(e1430)*

HSN   PDE   AVM

B

UNC-6::mNG intensity (au)

0   1   2   3   4   5
VNC   vSLNC

Dorsoventral distance (μm)

D

UNC-6::mNG intensity (au)

0   1   2   3   4   5
VNC   vSLNC

Dorsoventral distance (μm)

Figure 6

A

wildtype   *unc-40(e1430)*

0 sec   60 sec   120 sec   0 sec   60 sec   120 sec

UNC-6::mNG / HSN

1μm

UNC-6::mNG

B

Intensity (au) of UNC-6 foci contacting HSN

****

wildtype   *unc-40(e1430)*

C

Directed emergence — Chemotaxis

Stabilization at vSLNC — Haptotaxis

Directed growth — Chemotaxis

Stabilization at VNC — Haptotaxis

UNC-40 enrichment

UNC-6 gradient

ventral sublateral nerve cord (vSLNC)

ventral nerve cord (VNC)