ERH elicits cell lineage restriction in mammalian preimplantation development and differentiation

from pluripotency via H3K9me3-heterochromatin

Andrew Katznelson*[1], Blake Hernandez[1], Kylea Tapia[2], Holly Fahning[2], Adam Burton[3], Jingchao

Zhang[1], Maria-Elena Torres-Padilla[3,4], Nicolas Plachta[1], Kenneth S. Zaret[1]**, Ryan L.

McCarthy[2,5,6]**

[1]Institute for Regenerative Medicine and Department of Cell and Developmental Biology,

Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[2]Center for Developmental Biology and Regenerative Medicine, Seattle Children's Research

Institute, Seattle, WA 98101, USA

[3]Institute of Epigenetics and Stem Cells (IES), Helmholtz Zentrum München D-81377 München,

Germany

[4]Faculty of Biology, Ludwig-Maximilians Universität, München, Germany

[5]Department of Pediatrics, University of Washington School of Medicine, Seattle, WA 98195, USA

[6]Lead contact

**co-corresponding authors

R. McCarthy, Ryan.McCarthy@seattlechildrens.org

K. Zaret, zaret@pennmedicine.upenn.edu

Key words: ERH, H3K9me3, heterochromatin, embryo, pre-implantation, differentiation,
pluripotency, naïve

Abstract

Enhancer of Rudimentary Homolog (ERH) is an evolutionarily conserved protein originally characterized as promoting fission yeast heterochromatin[1] and recently shown to maintain H3K9me3 heterochromatin in human fibroblasts[2]. Here, we find that ERH depletion in fibroblasts reverts the somatic cell H3K9me3 landscape of broad megabase size domains to an embryonic stem cell (ESC) state composed of mainly H3K9me3 peaks and enables activation of naïve and pluripotency genes and transposable elements during induced pluripotent stem cell (iPSC) reprogramming. Concordantly, we find that ERH represses totipotent and alternative lineage programs during mouse preimplantation development and is required for proper segregation of the inner cell mass and trophectoderm cell lineages. During human ESC differentiation into germ layer lineages, ERH silences naïve and pluripotency genes, transposable elements, and alternative lineage somatic genes. As in fission yeast, we find that mammalian ERH interacts with RNA-binding proteins to engage and repress its chromatin targets. Our findings reveal a conserved, fundamental role for ERH in cell fate specification via the initiation and maintenance of early developmental gene repression.

Main text

Heterochromatin marked by histone H3 lysine 9 trimethylation (H3K9me3) represses the

expression of many protein-coding genes and DNA repeat elements[3–5]. Yet despite its stability,

H3K9me3-heterochromatin is dynamically established, maintained, and rearranged during

embryogenesis[3,6,7]. Zygotic genome activation (ZGA), marked by the initial transcription of key

genes and transposable elements[8], occurs at the 1-2 cell stage in mice and the 4-8 cell stage in

humans[8] and requires demethylation of maternal H3K9me3 marks created in oocytes[9,10]. As the

embryo loses totipotency and differentiates into the first two developmentally-specified lineages,

the trophectoderm of the blastocyst and the inner cell mass (ICM), many ZGA-specific protein-

coding genes and repeats gain H3K9me3 and are subsequently silenced [6,11,12]. As development

proceeds, H3K9me3 is further established on lineage-inappropriate genes and repeat elements

[6,11,13], with the highest number of genes marked by H3K9me3 at the germ-layer stage[3].

Throughout germ layer specification to mature somatic lineages, H3K9me3 is removed from

genes that become active and is deposited onto early developmental genes that become

repressed[3]. The H3K9me3 landscape changes substantially from pluripotency to early and

terminally differentiated states[14,15]. The mechanisms operating in the initial targeting of histone

methyltransferases to deposit H3K9me3 and the maintenance of the initially established

H3K9me3 domains remain poorly understood. In this study, we compared how *enhancer of*

rudimentary homolog (ERH), a highly conserved heterochromatin protein distinct from histone

methyltransferases, contributes to H3K9me3 dynamics and gene repression in differentiated

somatic cells, pluripotent cell lines, and early mammalian development.

ERH is an evolutionarily conserved protein originally discovered in fission yeast to regulate

sporulation, the primary form of yeast cell differentiation, via H3K9 methylation [1,16]. In *S. pombe,*

Erh1 forms a complex with the RNA binding protein Mmi1, which is recruited by binding of nascent

meiotic transcripts through its YTH domain[1,15]. Erh1-Mmi1 then causes the degradation of its

bound RNA transcripts, the establishment of H3K9 methylation, and consequent repression of

meiotic and developmental genes, transposons, and rDNA[1,16]. We recently found that in

terminally-differentiated primary human fibroblasts, ERH is highly enriched in H3K9me3-

heterochromatin, where it represses meiotic and germ cell genes, similar to that observed in

fission yeast[1,2,5]. We further found that ERH in mammals has evolutionarily expanded its function

to include the majority of regions that exhibit H3K9me3 heterochromatin in somatic cells, including

alternative-lineage protein coding genes and repeat elements[2]. While ERH is highly conserved

among eukaryotes, its fission yeast partner Mmi1 lacks a metazoan homolog[17]. Thus, how cell-

type specific cofactors recruit ERH to chromatin to regulate cell identity throughout mammalian

development is unknown.

Here, we find that upon ERH loss in fibroblasts, there is reversion to a pluripotency-like

H3K9me3 state which is permissive to binding of pluripotency transcription factors, which in turn

activate pluripotency and alternative lineage genes and repeat elements. The derepression of

pluripotency and alternative lineage programs led us to investigate the role of ERH in the earliest

developmental stages, where repression by H3K9me3 has been implicated during trophectoderm

and inner cell mass segregation in the blastocyst[6,11,12] and the differentiation from pluripotency to

the germ layer lineage[3]. By genetic perturbation, chromatin and gene expression profiling, and

imaging, we identify a role for ERH across mouse and human systems in repressing early naïve

pluripotency and alternative lineage genes. We find that targeting of mammalian ERH to

chromatin occurs through cell-type specific interactions with RNA binding proteins, enabling

context-dependent functions. Together, our findings demonstrate a fundamental role for ERH

during developmentally acquired H3K9me3-mediated gene repression and maintenance of such

repression in somatic cells.

Results

ERH depletion in primary human fibroblasts causes H3K9me3 to revert to an ESC-like state

We assessed how H3K9me3 is disrupted after ERH loss by CUT&RUN for H3K9me3 after siRNA

knockdown in human primary fibroblasts (Fig. 1a). Consistent with our previous study[2], we

confirmed that H3K9me3 is globally reduced in human fibroblasts after ERH knockdown (Fig. 1b).

Yet while large megabase-scale domains were diminished (Fig. 1c top, purple domains in top two

tracks), as seen previously[2],10,526 H3K9me3 residual peaks were retained in the ERH

knockdown (Fig. 1c, lower, second track) and Supplementary Table 1). Remarkably, the residual

H3K9me3 peaks in siERH fibroblasts resemble the H3K9me3 profile in H1 ESCs (Fig. 1c lower,

third track). Across the genome, 85% of the residual fibroblast siERH H3K9me3 sites overlap with

H3K9me3 peaks in H1 hESCs (Extended Data Fig. 1a). Pearson correlation of H3K9me3 peaks

from fibroblasts after siControl or siERH, as well as in hESCs, supports a shift in the H3K9me3

heterochromatin landscape to an embryonic state (Fig. 1d,e). Thus, while loss of ERH in primary

fibroblasts causes a global loss of H3K9me3 domains, embryonic H3K9me3-marked targets are

retained.

We hypothesized that since ERH depleted fibroblasts retain hESC-like H3K9me3 peaks

(Fig. 1c-e), ERH may be dispensable for maintaining H3K9me3 in hESCs. To test this hypothesis

we depleted ERH in H1 hESCs cultured in pluripotency-maintaining conditions, which caused no

observable changes in colony morphology or loss of pluripotency (Fig. 1f), even after sustained

knockdown (Extended Data Fig. 1d). H3K9me3 level and nuclear organization by

immunofluorescence was not changed in siERH-treated hESCs (Fig. 1g; Extended Data Fig. 1e).

We also observed no change in H3K27me3 (Extended Data Fig. 1f). Despite no apparent

disruption of the pluripotent state or global H3K9me3 levels, ERH-depleted hESCs show an

increased nuclear area and nuclear blebbing (Extended Data Fig. 1g), which are changes

associated with chromatin decompaction[18]. We previously found that ERH depletion did not

change the expression or protein level of H3K9me3 HMTs[2], we similarly observe no significant

decrease in H3K9me3 HMT expression upon ERH depletion in pluripotent hESCs (Extended Data

Fig. 1h). To investigate potential locus-specific changes in hESC H3K9me3 upon ERH depletion,

we conducted H3K9me3 CUT&RUN in siControl and siERH treated hESCs. Consistent with our

immunofluorescence results, we observed minimal change in H3K9me3 (Extended Data Fig. 1i)

and high correlation between H3K9me3 peaks in siControl and siERH treated hESCs (Extended

Data Fig. 1j).

To identify chromatin-associated proteins and pathways regulating the residual H3K9me3

peak sites in human fibroblasts, we analyzed the regions using the cistromeDB Toolkit, which

uses a compendium of published ChIP-seq data sets[16,17]. We find that most proteins enriched at

the residual peak sites in ES cells (i.e., prior to ERH knockdown) are either KRAB-ZNFs or

involved in H3K9me3 repression via KAP1, which is recruited to chromatin through binding the

KRAB domain (Fig. 1h)[19]. Likewise, de novo motif analysis finds that enriched sequence motifs in

the residual peak sites are for KRAB-ZNFs factors (Extended Data Fig. 1k). Consistent with the

enrichment analysis, the residual siERH H3K9me3 peak sites are highly enriched for H3K9me3,

KAP1, and HP1 in ES cells (Fig. 1e,i; Extended Data Fig. 1l). As KRAB-ZNFs repress transposons

via H3K9me3, we hypothesized the residual peaks would be enriched over repeats. Repeat

enrichment analysis found that the residual peaks are enriched over canonical KRAB-ZNF

targets: LINEs, LTRs, and satellite repeats (Extended Data Fig. 1m). We conclude that ERH is

required to maintain the broad H3K9me3 domains that are established over the course of

development and are present in terminally differentiated cells, but does not control H3K9me3

peaks attributed to the DNA-dependent association of KAP1 that persists from pluripotency over

constitutively repressed repeats.

ERH-dependent H3K9me3 suppresses SOX2 binding and activation of pluripotency genes

and repeats during OSKM reprogramming

H3K9me3 impairs induced pluripotent stem cell (iPSC) reprogramming by blocking transcription

factor binding, and pluripotency genes marked by H3K9me3 show delayed activation compared

to genes in heterochromatin marked with H3K27me3 or neither mark (Extended Data Fig. 2a)[20].

We therefore assessed the impact of ERH depletion in the context of OSKM transcription factor

binding and gene activation during iPSC reprogramming. We focused on a 48 hour time point

after initiation of iPSC reprogramming (Fig. 1a and Supplementary Tables 2-4), as H3K9me3 is

inhibitory to transcription factor activity at this stage[20].

CUT&RUN of SOX2 after 48 hours of OSKM reprogramming showed that while most

binding sites were shared between siControl and siERH fibroblasts, 2,538 binding sites were

gained (Extended Data Fig. 3a and Supplementary Table 2). We observed that ERH knockdown

enabled SOX2 to target loci that lose H3K9me3, including within heterochromatin domains

previously annotated as resistant to reprogramming factor binding in fibroblasts[20] (Fig. 2a, and

Extended Data Fig. 3b-d). By assessing the genomic locations of siERH-gained SOX2 peaks in

the cistromeDB toolkit[21,22], we found that the newly gained SOX2 sites had the most similarity to

ChIP-seq peak sets of SOX2 and its pluripotency partners NANOG and OCT4 in embryonic stem

cells (Extended Data Fig. 3e). Inspection of genome browser tracks identified specific loci where

loss of ERH-controlled H3K9me3 enabled SOX2 binding over normal SOX2 and NANOG targets

in ES cells (Fig. 2b, Extended Data Fig. 3f). We next assessed if loss of ERH-dependent

H3K9me3 enables other transcription factors to bind previously refractory sites by performing

CUT&RUN footprinting of SOX2's primary binding cofactors during reprogramming, KLF4 and

OCT4, and find enhanced footprints for each factor at SOX2 sites gained by siERH (Extended

Data Fig. 3g). Thus, ERH loss during pluripotency reprogramming enables transcription factor

binding in what were heterochromatic H3K9me3 regions.

Protein coding genes embedded in H3K9me3 domains, normally refractory to transcription

factor activation, were activated after siERH (136 up, 13 down; Fig. 2c). Significantly, genes

associated with the human 8-cell like cell (8CLC) state[23] were activated, including *ZSCAN4,*

DUXA, DPPA4, LIN28A, TPRXL[24]*, KLF5,* and the histone variant *H3.Y*[25] (Fig. 2d and Extended

Data Fig. 4a). We compared the activation levels of these genes in siERH+OSKM with existing

RNA-seq datasets[26] and found that *H3.Y, DUXA* and *TPRXL* do not normally get activated at

similar levels to siERH+OSKM at any stage during iPSC reprogramming (Extended Data Fig. 4b).

However, activation of 8CLC genes in siERH+OSKM is incomplete, as many genes activated in

naïve ESCs[27] are not expressed or expressed at lower levels in siERH+OSKM (Extended Data

Fig. 4c), suggesting low activation of naïve programs but not full conversion to a naïve cell state.

While the activation of marker genes such as DUXA and ZSCAN4 is lower than in 8C embryos[28],

we conclude that ERH-dependent H3K9me3 safeguards early totipotent and pluripotent genes

and that ERH loss creates a permissive environment for transcription factor binding and

activation.

We observed that *LTR7* and *HERVH* repeats, which are expressed in and required to

maintain pluripotency[29,30], lose H3K9me3 after ERH depletion and are activated in siERH+OSKM

(Fig. 2e). *LTR7Y* repeats, which promote naïve pluripotency[31], exhibited similar ERH-dependent

H3K9me3 and repression (Fig. 2f,g and Extended Data Fig. 4d). Notably, residual H3K9me3

peaks retained at *LTR7Y* repeats were insufficient to maintain repression (Fig. 2f,g). Furthermore,

we observed siERH+OSKM activated repeat elements that function specifically during the human

8 cell-like cell stage, and are later marked by H3K9me3 and silenced in the human blastocyst[6]

(Extended Data Fig. 4e). These results demonstrate that ERH maintains H3K9me3 repression of

a subset of 8-cell stage and pluripotency repeats in terminally differentiated human cells.

Depletion of H3K9 HMTs has been shown to increase iPSC reprogramming[20]. However,

despite increased activation of pluripotency and 8-cell genes at 48 hours after initiation of OSKM-

based iPSC reprogramming in siERH treated fibroblasts, by day 2 we observed substantial cell

death that persisted 7 days into reprogramming, past the initial apoptotic wave (Extended Data

Fig. 4f,g) and resulting in no iPSC colonies forming in siERH+OSKM conditions, compared to the

abundant colonies in siControl+OSKM conditions (Extended Data Fig. 4h,i). The failure to

generate iPSC colonies was not rescued by simultaneous knockdown of p53 (Extended Data Fig.

4i). Lower ERH siRNA decreased cell death, but did not change the emerging iPSC colony

number (Extended Data Fig. 4g,h). Using XDeathDB[32] on our RNA-seq data identified increased

expression of genes activated in several cell death pathways, including RIPK2 signaling

(Extended Data Fig. 4j). We investigated if simultaneous inhibition of RIPK2 along with inhibition

of RIPK3 and Caspases 3 and 7, which has been shown to improve survival of cells activating 8

cell like genes[23], could enable survival of siERH+OSKM iPSCs. Notably, we found that adding

small molecule inhibitors GSK2983559, Ac-DEVD-CHO, and GSK872 to inhibit RIPK2, Caspases

3 and 7, and RIPK3, respectively, during reprogramming enabled the generation of iPSC colonies

with ERH depletion at a significantly higher efficiency compared to siControl (Extended Data Fig.

4k). RIPK2 inhibition alone was necessary and sufficient to successfully generate iPSC colonies

expressing pluripotency factors during ERH depletion, while inhibition of RIPK and Caspases 3

and 7 was not sufficient but did increase colony size (Extended Data Fig. 4l). These findings

demonstrate that the increased accessibility for transcription factor binding in H3K9me3 domains

(Fig. 2a,b) and improved activation of the pluripotency network (Fig. 2c,d) observed upon ERH

depletion can increase iPSC reprogramming efficiency.

ERH is necessary for trophectoderm development, the first embryonic differentiation

Our finding that ERH represses naïve genes that are normally expressed in 2-8 cell embryos, in

fully differentiated fibroblasts, led us to investigate if ERH establishes silencing of 2-8 cell genes

during pre-implantation development. We first analyzed existing RNA-seq data of mouse

embryogenesis timepoints from zygote to blastocyst[33] and found that ERH mRNA is present

across pre-implantation development and is significantly increased from the 8-cell stage through

the blastocyst stage (Extended Data Fig. 5a). Immunostaining mouse embryos for ERH protein

at the zygote, 2-cell, 8-cell and blastocyst stages revealed that, consistent with its mRNA levels,

ERH is expressed and nuclear-localized at all stages analyzed, with increasing abundance in the

early and late blastocyst stages (Fig. 3a,b). At the blastocyst stage, ERH levels are similar in ICM

and trophectoderm cells (Extended Data Fig. 5b).

To investigate if ERH is involved in pre-implantation lineage specification, we depleted

ERH levels by siRNA micro-injection at the 1-cell stage of mouse embryos, compared to a

nontargeting siRNA control, and assessed impacts on the blastocysts (Extended Data Fig. 5c).

ERH depletion did not cause major morphological changes in blastocyst structure, as both

siControl and siERH blastocysts form a cavity and segregate ICM and trophectoderm cells

(Extended Data Fig. 5c,d). Staining blastocysts for key transcription factors involved in lineage

specification revealed similar levels of the pluripotency marker OCT4 in ICM and TE cells between

siControl and siERH blastocysts (Extended Data Fig. 5e). However, the levels of the

trophectoderm-determining transcription factor CDX2[34] (Fig. 3b,c) and the trophectoderm marker

Keratin 8[35] (Fig. 3d,e) were significantly decreased in the ERH depleted blastocysts. Thus, ERH

is required for proper trophectoderm differentiation.

The role of ERH in H3K9me heterochromatin regulation in *S. pombe*[1,16] and human

fibroblasts[2] suggests that defects in trophectoderm differentiation may arise from alterations in

gene silencing mechanisms. Indeed, we observed a 32% decrease (p=0.0025) in H3K9me3 in

blastocysts of ERH-depleted embryos (Fig. 3f,g and Supplementary Table 5). We also observed

a significant increase in nuclear DNA shedding in the siERH blastocytes (Fig. 3h,i; Extended Data

Fig. 5f), similar to the observed nuclear DNA shedding in siERH hESCs (Extended Data Fig. 1g).

Decreased keratin levels as observed in siERH blastocytes (Fig. 3d,e) have been shown to make

trophectoderm cells of blastocyts more susceptible to DNA shedding[36], as has decreased

H3K9me3[18].

To characterize how these changes impact gene expression, we performed RNA-seq on

mouse blastocysts 4 days after injecting siControl or siERH at the 1 cell stage (Supplementary

Table 6), which confirmed an ~80% knockdown of ERH (Extended Data Fig. 5g). We classified

genes upregulated in ERH depleted blastocysts into 4 groups, based upon their expression during

normal mouse preimplantation development[33] (Fig. 3j; Supplementary Table 7). Group 1 consists

of transcripts present at the zygote stage, including maternally inherited transcripts, that become

silenced or degraded as development progresses. Group 2 consists of transcripts that are highly

expressed in the 2 to 8 cell stages and low or absent in the zygote and blastocyst stage. Group 3

consists of transcripts that are absent or lowly expressed at all stages measured. Group 4 consists

of transcripts with the highest expression in the blastocyst stage. Of the ERH-repressed genes

(i.e., genes whose expression goes up upon ERH knockdown), 25 normally have peak expression

during the 2 to 8 cell stage (group 2) and 54 are normally not expressed during preimplantation

development (group 3). ERH-repressed group 2 genes included the 2-cell specific *Zscan4d*[37] and

the 4-cell specific *D1Pas1*[38] (Fig. 3k). ERH-repressed group 3 alternative lineage genes included

genes specific to the brain[39], liver, and intestine [40] (Fig. 3k).

Comparing our RNA-seq data with existing mouse preimplantation H3K9me3 ChIP-seq

data[11], we found that the genes repressed by ERH at the 2-cell stage and later, alternative lineage

genes were initially marked by H3K9me3 (Fig. 3k). We further found that upon ERH depletion,

key markers of primitive endoderm, including *Gata4* and *Sox17*[41], were significantly repressed

(Extended Data Fig. 5h), while epiblast genes were unchanged (Extended Data Fig. 5i). We

conclude that ERH is necessary to silence 2-8 cell stage genes and prevent the spontaneous

activation of later developmental genes during the blastocyst stage, and that failure to silence

results in improper lineage specification in the blastocyst.

ERH is necessary for proper lineage commitment and restriction during human ESC

differentiation

Our finding that ERH is critical for blastocyst development led us to ask if ERH depletion impacted

the ability of hESCs to differentiate. When we used Activin A and the GSK-3 Inhibitor CHIR99021

together to differentiate hESCs to an endoderm identity[42], as expected within 24 hours we

observed repression of pluripotency factors and initial expression of endodermal markers in a

subset of cells (Extended Data Fig. 6a). Yet depletion of ERH in H1 hESCs followed by 24 hour

endoderm differentiation showed a failure to decrease NANOG protein levels (Fig. 4a and

Extended Data Fig. 6b) and a reduced emergence of SOX17+ cells (Fig. 4a). To assess if this

observation was specific to endoderm, we also performed differentiation to ectoderm, with control

and ERH depleted cells, for 2 days with retinoic acid treatment. We observed sustained

expression of the pluripotency factor OCT4 and decreased abundance of the ectodermal

transcription factor PAX6 (Fig. 4b).

To investigate transcriptional changes caused by ERH depletion, we performed RNA-seq

of siRNA control and ERH depleted H1 hESCs (Extended Data Fig. 6c and Supplementary Table

8) during pluripotent culture, after 1 day of endoderm or 2 days of ectoderm differentiation. While

ERH depletion in fibroblasts+OSKM resulted in more gene activation than repression (Fig. 2c;

Extended Table 3), depletion in pluripotent hESCs exhibited less derepression (Extended Data

Fig. 4d; Extended Table 8), consistent with the difference in the importance of ERH in H3K9me3

maintenance between fibroblasts and hESC state (Fig. 1). Genes normally repressed in the

siControl endoderm and ectoderm differentiation were preferentially upregulated by ERH

depletion (endoderm, Extended Data Fig. 6f; ectoderm Extended Data Fig. 6g).

During endoderm differentiation, we observed that ERH depletion preferentially

upregulated genes identified as higher in naïve vs primed hESCs[43] (Extended Data Fig. 6h),

beyond what was observed in the pluripotent state (Extended Data Fig. 6e). For example, the

gene for the pluripotency factor NANOG, previously demonstrated to be repressed by H3K9me3

during pluripotency exit[44], was significantly upregulated by ERH depletion (Fig. 4c), consistent

with our immunostaining (Fig. 4a and Extended Data Fig. 6b). Additionally, genes found to be

expressed in human pluripotent stem cells were upregulated during endoderm and ectoderm

differentiation after ERH depletion, including *DUSP5/6, NODAL, LEFTY1/2*, and *DACT1*

(Extended Data Fig. 6i and Supplementary Table 8). Genes associated with the 2 cell stage and

naïve pluripotency, including *ZSCAN4*[37]*, KLF5*[31]*, ESRRB*[45]*, NR5A2*[46] and *NANOGP1*[47]*,* were

upregulated during siERH-treated hESC differentiation (Fig. 4d and Extended Data Fig. 6j). The

expression of naïve genes after ERH depletion was at a much lower level than in true naïve

hESCs (Extended Data Fig. 6k), indicating permissive expression but not spontaneous primed to

naïve conversion. We conclude that ERH is required to repress pluripotency genes during hESC

pluripotency exit.

During differentiation, a key function of H3K9me3 heterochromatin is to repress alternative

lineage genes[3,11]. Our finding that ERH depletion in blastocysts upregulates lineage-incompatible

genes (Fig. 3j,i) led us to investigate if a similar consequence occurs during hESC differentiation.

GO analysis of genes upregulated by ERH depletion during endoderm and ectoderm

differentiation confirmed activation of transcriptional networks typically not induced by the specific

differentiation treatment (Fig. 4e, Extended Data Fig. 7a and Supplementary Table 10). ERH

depletion upregulated meiotic and gametogenic genes in all tested developmental stages

(Extended Data Fig. 7b), consistent with previous findings in human fibroblasts[2] and the role of

ERH in S. pombe sporulation[1]. To further assess the cell lineages in which the upregulated genes

function, we generated composite expression profiles of the genes upregulated by ERH depletion

during endoderm (Fig. 4f) and ectoderm differentiation (Fig. 4g) from expression data downloaded

from the BBI Descartes human single cell atlas[48], and found that heart, muscle and liver genes

are preferentially activated. Directed differentiation of hESCs to endoderm[42], instead induced

both early and late mesodermal genes[49] when ERH was depleted (Fig. 4h,i). Notable

inappropriate gene activation in siERH definitive endoderm included *NXK2-5, NPPB* and *FLNC*,

typically expressed in heart muscle, as well as *TAGLN,* a canonical smooth muscle marker (Fig.

4i). Increased levels of BRACHYURY protein, a key determinant of early mesoderm[50], was also

confirmed by immunofluorescence (Fig. 4j). Inappropriate activation during ectoderm

differentiation with ERH knockdown included genes typically expressed in liver development

(Extended Data Fig. 7c). Thus, while ERH appears dispensable to maintain hESC pluripotency,

it is essential to restrict cell identity for proper germ layer differentiation.

ERH represses transposable elements and proximal genes

Having observed that ERH represses pluripotency and alternative lineages during hESC

differentiation, we proceeded to investigate how this repression was elicited. During pluripotency-

to-endoderm differentiation at protein-coding genes that are upregulated in siERH DE vs siControl

DE, H3K9me3 levels only slightly increase in and around those genes (i.e., without siERH) during

endoderm differentiation (Extended Data Fig. 8a). However, repeat elements including L1 LINEs

do normally exhibit a substantial H3K9me3 gain during endoderm differentiation (Fig. 5a).

Repression of transposable elements is a canonical function of H3K9me3 heterochromatin[51] and

transposable elements can regulate developmental gene networks by acting as enhancers for

transcription factor binding[31,52–55]. We therefore assessed transposable elements in our RNA-seq

data from siControl and siERH-treated H1 hESCs during self-renewal and differentiation

(Supplementary Table 9), with the hypothesis that dysregulation of specific transposable elements

might underlie the observed differentiation defects in siERH treated hESCs.

We found that transposable elements were upregulated by ERH knockdown in H1 hESCs,

definitive endoderm, and ectoderm, including classes of L1 elements and the hominid-specific[56]

SINE VNTR and Alu (SVA) elements (Fig. 5b; Extended Data Fig. 8b). Satellite repeats were also

derepressed (Fig. 5b), as were morula-blastocyst specific *LTR7Ys*[31] and human 8 cell stage

expressed *HERVKs*[57] (Fig. 5b; Extended Data Fig. 8c,d), which act as transcription factor binding

sites for pluripotency[52,58] and lineage specific genes[55]. After stratifying L1 elements by their

evolutionary age[59], we found that ERH preferentially represses evolutionarily young L1 repeats in

pluripotent and differentiating hESCs, but broadly represses L1 elements in terminally

differentiated fibroblasts (Fig. 5c). By filtering our data for genes expressed during pluripotency

that exhibit ERH-dependent repression during differentiation, we uncovered evolutionarily young

transposable elements (Fig. 5d and Extended Data Fig. 8e,f) and L1 repeats within 25 kb

(Extended Data Fig. 8g) in regions normally marked by H3K9me3 in hESC-derived endoderm

(Fig. 5e note different scales on subset to right; Extended Data Fig. 9a).

To assess if repeat expression upon ERH knockdown during endoderm differentiation was

due to H3K9me3 loss, we performed H3K9me3 CUT&RUN in siControl and siERH hESCs

differentiated to endoderm. We found the increased L1 element RNA expression upon ERH

depletion (Fig. 5c) did not globally correspond to a loss of H3K9me3 (Extended Data Fig. 9b).

However, when we restricted the analysis to L1 TEs that gained H3K9me3 with at least a 2-fold

increase in RNA in siERH endoderm, the L1 loci showed decreased H3K9me3 upon ERH

depletion (Fig. 5f). We found repeat element loci with increased RNA expression with (Fig. 5g left)

and without H3K9me3 loss (Fig. 5g right) indicating that RNA expression from TEs upon ERH

depletion during endoderm differentiation was not necessarily dependent upon H3K9me3 loss.

We observe instances of endoderm specific upregulation of genes proximal to evolutionarily

young repeats, such as L1 and LTR7Ys, with reduced H3K9me3 in ERH depleted endoderm

(Extended Data Fig. 9c). ERH therefore represses evolutionarily young and lineage-specific

repeats, thus preventing improper activation of pluripotency and alternative lineage genes during

differentiation to maintain lineage specification integrity. The role of H3K9me3 and ERH at this

cell fate change likely occurs predominantly at transposable element regions and not gene bodies.

ERH is recruited to chromatin by cell-specific cofactors

Our observation that ERH is required for maintenance of H3K9me3 in fibroblasts (Fig. 1b) but

dispensable for hESC H3K9me3 (Fig. 4b) led us to investigate potential ERH protein interactions

that may drive the difference in regulatory mechanism. We focus the experiments on the

pluripotent and terminally differentiated states as they exhibit contrasting requirements for ERH

in H3K9me3 maintenance (Fig. 1b-g) and very different H3K9me3 distribution patterns (Fig. 1d,

i; Extended Data Fig 1g). In *S. pombe* Erh is recruited by the RNA binding factor Mmi1[1]. Mmi1

itself is not seen in mammalian cells[60,61], but since it is a YTH domain-containing RNA binding

factor, we investigated potential interactions of ERH with human YTH domain proteins YTHDC1

and YTHDC2. YTHDC1 regulates H3K9me3 through SETDB1 recruitment and repression of

repeat elements in mouse ESCs[62,63] and YTHDC2 has been shown to positively regulate

transposable elements and the non-coding RNA LINC-ROR in human hESCs[64]. To specifically

interrogate on chromatin ERH interacting proteins, we isolated soluble nucleoplasm (NPL) and

insoluble chromatin-associated fractions (CAF) and then performed immunoprecipitation of ERH

(Fig. 6a). In fibroblasts but not hESCs, we observed that ERH interacts with YTHDC1 in the

chromatin fraction (Fig. 6b,c). ERH did not interact with YTHDC2 in the chromatin fraction in either

hESCs or fibroblasts, despite being present in both cell types (Fig. 6b,c). We also probed for

SAFB1, a known ERH interacting protein[65], and observed consistent ERH-SAFB1 interactions in

both fibroblasts and pluripotent hESCs (Fig. 6b,c), demonstrating that YTHDC1 but not SAFB1

interactions with ERH are cell type-dependent. To investigate the cause of this differential

interaction we first analyzed ERH levels, finding that while ERH RNA is not significantly different

between pluripotency and fibroblasts (Extended Data Fig. 10a) or between pluripotent and

differentiated hESCs (Extended Data Fig. 6c), fibroblasts stain more strongly for ERH protein

(Extended Data Fig. 10b). We observe that ERH protein levels are higher in cells on the edge of

hESC colonies (Extended Data Fig. 10c) and become higher in all cells after 1 day of endoderm

differentiation (Extended Data Fig. 10c). These results indicate that ERH involvement in

H3K9me3 regulation and RNA binding protein regulation may be regulated by ERH protein level.

As S. pombe Erh1 chromatin recruitment is RNA and Mmi1 dependent, we next tested if

chromatin association of ERH was RNA dependent and if either YTHDC1 or YTHDC2 facilitated

the recruitment. We employed an immunostaining approach[66] in which permeabilized nuclei were

treated with RNase A and then stained for ERH, and found that ERH nuclear retention was

significantly decreased (Fig. 6d). In contrast, RNase H treatment, which digests DNA:RNA

hybrids, did not release ERH from the nucleus (Fig. 6d). In an orthogonal approach, RNase A

treatment of fibroblasts during soluble nuclear protein isolation released a portion of chromatin-

bound ERH into the nucleoplasm, indicating that its chromatin retention is at least partially RNA-

dependent (Extended Data Fig. 10d). To determine if YTHDC1 or YTHDC2 facilitate ERH

chromatin recruitment, we targeted each separately with siRNAs in human fibroblasts and

observed that YTHDC1 but not YTHDC2 depletion led to decreased levels of ERH in the

chromatin fraction and increased ERH in the soluble nuclear fraction (Fig. 7e). We confirmed the

role of YTHDC1 in ERH retention utilizing our cell permeabilization and immunostaining assay

(Extended Data Fig. 10e). From these experiments we concluded that ERH associates with

chromatin with YTHDC1 in fibroblasts, but not in hESCs, and that a subset of ERH protein requires

RNA and YTHDC1 for chromatin localization.

We next assessed for co-regulation of common targets by ERH, YTHDC1, and YTHDC2,

first by evaluating expression changes of published YTHDC1 and YTHDC2 targets, ERVK

elements repressed by YTHDC1[67], and HERVH elements activated by YTHDC2[64]. ERH

represses HERVK and HERVH elements in fibroblasts treated with OSKM factors but not in

pluripotent hESCs (Fig. 6f), matching the interaction pattern of ERH with YTHDC1 (Fig. 6b,c).

Major satellite repeats, which are regulated by SAFB1[68], are repressed by ERH in both fibroblasts

and pluripotent hESCs (Fig. 6f).

As we found that ERH interacts with YTHDC1 in differentiated cells but not pluripotent

cells, we investigated if YTHDC1 depletion recapitulates aspects of ERH depletion during hESC

differentiation. Depletion of YTHDC1 but not YTHDC2 during endoderm differentiation of hESCs

resulted in an increase of Brachyury staining indicative of a mesodermal fate (Fig. 6g), matching

our findings from ERH depletion during endoderm differentiation (Fig. 5f). We found that ERH

represses many genes not regulated by its interacting partners YTHDC1 and SAFB1, which

predominately repress different subsets of ERH targets (Fig. 6h). Taken together, the results

indicate that various RNA binding proteins function non-redundantly to recruit ERH to different

targets. Based upon these findings, we propose that ERH interactions with RNA binding proteins,

including YTHDC1 and SAFB1, facilitate cell-specific repression of genes and repeat elements

(Fig. 6i).

Discussion

We discovered that depletion of ERH reverts the H3K9me3 landscape in terminally differentiated

cells to an hESC-like state, permissive to activation of early embryonic genes. This surprising

result led us to investigate the role of ERH in establishing gene repression during the earliest

stages of mammalian lineage segregation, in maintaining repression of pluripotency, and enabling

germ layer specification. We found that ERH controls H3K9me3 for repression of naïve and

alternative lineage genes during the cell fate commitment of totipotent cells to ICM and

trophectoderm in mouse blastocysts as well as of hESCs to the endoderm and ectoderm cell

lineages, to enforce cell fate and prevent anomalous gene activation. Between the three systems

we employed, mouse preimplantation development, hESC differentiation, and fibroblast iPSC

reprogramming, we observe that ERH commonly repressed a shared set of naïve genes such as

ZSCAN4 across all systems, indicating that ERH is involved in both the initial repression of this

early transcriptional program and in maintaining repression in terminally differentiated cells.

Remarkably, ERH is the sole apparent mechanism for repressing meiotic genes in *S. pombe,* is

conserved in repressing germ cell genes in mammals[2], and has expanded to foundationally

control diverse other types of developmental genes and stages.

In a developmental context, the failure to repress 2-cell, 8-cell and pluripotency genes

after ERH knockdown results in delays of specific-lineage gene expression, including primitive

endoderm genes in mouse blastocysts and various endoderm and ectoderm genes during hESC

differentiation. Furthermore, we demonstrated that ERH is necessary to maintain repression of

alternative lineage genes, as ERH depletion elicited precocious expression of genes typical of

later cell differentiation in mouse blastocysts and human hESCs, as well as heterochromatinized

genes during initial iPSC reprogramming. We observed that evolutionarily young transposable

elements, including the human-specific *LTR7Ys*[31], are especially upregulated by ERH depletion

during hESC differentiation and iPSC reprogramming. Much of the lineage specific expression

changes upon ERH depletion during hESC differentiation may be attributed to increased

accessibility of transcription factor binding sites among derepressed repeats[31,52,55].

A key finding is that ERH depletion in human fibroblasts reverted the H3K9me3 chromatin

landscape to a hESC-like state. The reversion to a hESC-like H3K9me3 landscape permits

binding of SOX2 in fibroblasts to endogenous hESC SOX2 targets, activating embryonic genes.

That ERH is necessary for maintaining the broad H3K9me3 domains observed in somatic cells,

but is dispensable for maintaining hESC H3K9me3 peaks, is consistent with our observation that

ERH depletion in hESCs did not change global H3K9me3 levels. We suggest that the residual

H3K9me3 peaks in pluripotent cells, enriched for zinc finger motifs and KAP1 binding, may be

H3K9me3 nucleation sites and that, during differentiation, ERH is essential for the spread of

H3K9me3 from these sites. Importantly, when ERH is knocked down in fibroblasts, we observe

increased activation of many genes and transposable elements that harbor a residual H3K9me3

peak, yet lose broad H3K9me3, indicating that residual H3K9me3 peaks are not sufficient to block

gene activation. Our knowledge of H3K9 methylation nucleation and spread has been primarily

elucidated through studies in *S. pombe*[69–71], leaving further investigation needed into how

H3K9me3 nucleation and spread occurs in multicellular organisms.

We identify that ERH interacts with and is partially recruited to the chromatin by the RNA

binding protein YTHDC1 in somatic cells, but not in pluripotent hESCs. We find that ERH and

YTHDC1 repress similar targets in fibroblasts and differentiating hESCs, but not in homeostatic

pluripotent hESCs. In contrast, we observe that ERH interacts with the RNA binding protein

SAFB1 in both hESCs and human fibroblasts and likewise both repress the satellite repeat

expression in hESCs and fibroblasts. Our knockdowns of YTHDC1 and gene expression analysis

of YTHDC1 and SAFB1 fail to fully explain how ERH is recruited to chromatin to repress

transcription. We thus speculate that ERH is recruited to cell-type specific genes and repeats

through a diverse network of chromatin-associated RNA binding factors that are yet to be fully

identified.

We observe in mouse development and hESC systems that ERH is functionally required

in multiple developmental contexts to repress meiotic and gametogenic genes, as it is in *S.*

pombe[1], although the gene members targeted by ERH has greatly evolved and expanded in

mammals. Our findings on meiotic gene repression are supported by the recent identification of

ERH in a screen for proteins silencing germline genes in mESCs[72]. Other aspects of *Erh1* function

in *S. pombe,* such as the targeting of RNAs for degradation by the RNA exosome[1], appear to be

conserved as we observed upregulation of developmental genes typically downregulated by RNA

exosome degradation[73]. ERH was recently found by mass spectrometry to associate with the

Microprocessor and NEXT complexes in mouse ESCs[74]. Evolutionarily divergent functions of

ERH may be driven by evolutionarily young transposable elements that retain RNA expression,

similar to how transposable elements have influenced the evolution of KRAB Zinc finger proteins

DNA dependent targeting of H3K9me3[75,76].

We found that ERH enacts gene repression at the earliest stages of development and that

it is essential for maintaining this repression in terminally differentiated cells. Our findings advance

the understanding of how heterochromatin-based repression is established and maintained

during development. We propose that understanding mechanisms by which ERH is employed to

direct H3K9me3 to specific loci in a developmentally coordinated fashion will be crucial to

modulating developmental H3K9me3 rearrangements to achieve greater control of cell identity.

References

1. Sugiyama, T. *et al.* Enhancer of Rudimentary Cooperates with Conserved RNA-Processing Factors to Promote Meiotic mRNA Decay and Facultative Heterochromatin Assembly. *Mol Cell* **61**, 747–759 (2016).

2. McCarthy, R. L. *et al.* Diverse heterochromatin-associated proteins repress distinct classes of genes and repetitive elements. *Nat Cell Biol* **23**, 905–914 (2021).

3. Nicetto, D. *et al.* H3K9me3-heterochromatin loss at protein-coding genes enables developmental lineage specification. *Science* **363**, 294–297 (2019).

4. Bilodeau, S., Kagey, M. H., Frampton, G. M., Rahl, P. B. & Young, R. A. SetDB1 contributes to repression of genes encoding developmental regulators and maintenance of ES cell state. *Genes & development* **23**, 2484–9 (2009).

5. Becker, J. S. *et al.* Genomic and Proteomic Resolution of Heterochromatin and Its Restriction of Alternate Fate Genes. *Mol Cell* **68**, 1023-1037.e15 (2017).

6. Xu, R. *et al.* Stage-specific H3K9me3 occupancy ensures retrotransposon silencing in human pre-implantation embryos. *Cell Stem Cell* **29**, 1051-1066.e8 (2022).

7. Zhang, J. *et al.* Distinct H3K9me3 heterochromatin maintenance dynamics govern different gene programmes and repeats in pluripotent cells. *Nat Cell Biol* **26**, 2115–2128 (2024).

8. Schulz, K. N. & Harrison, M. M. Mechanisms regulating zygotic genome activation. *Nat Rev Genet* **20**, 221–234 (2019).

9. Xu, R. *et al.* Unreprogrammed H3K9me3 prevents minor zygotic genome activation and lineage commitment in SCNT embryos. *Nat Commun* **14**, 4807 (2023).

10. Sankar, A. *et al.* KDM4A regulates the maternal-to-zygotic transition by protecting broad H3K4me3 domains from H3K9me3 invasion in oocytes. *Nat Cell Biol* **22**, 380–388 (2020).

11. Wang, C. *et al.* Reprogramming of H3K9me3-dependent heterochromatin during mammalian

embryo development. *Nature cell biology* **20**, 620–631 (2018).

12. Yu, H. *et al.* Dynamic reprogramming of H3K9me3 at hominoid-specific retrotransposons during

human preimplantation development. *Cell Stem Cell* **29**, 1031-1050.e12 (2022).

13. Burton, A. *et al.* Heterochromatin establishment during early mammalian development is regulated

by pericentromeric RNA and characterized by non-repressive H3K9me3. *Nat Cell Biol* **22**, 767–778

(2020).

14. Bhanu, N. V., Sidoli, S. & Garcia, B. A. Histone modification profiling reveals differential signatures

associated with human embryonic stem cell self-renewal and differentiation. *Proteomics* **16**, 448–

458 (2016).

15. Ugarte, F. *et al.* Progressive Chromatin Condensation and H3K9 Methylation Regulate the

Differentiation of Embryonic and Hematopoietic Stem Cells. *Stem Cell Reports* **5**, 728–740 (2015).

16. Xie, G. *et al.* A conserved dimer interface connects ERH and YTH family proteins to promote gene

silencing. *Nat Commun* **10**, 251 (2019).

17. Kuznetsov, D. *et al.* OrthoDB v11: annotation of orthologs in the widest sampling of organismal

diversity. *Nucleic Acids Res* **51**, D445–D451 (2023).

18. Stephens, A. D. *et al.* Chromatin histone modifications and rigidity affect nuclear morphology

independent of lamins. *Mol Biol Cell* **29**, 220–233 (2018).

19. Stoll, G. A., Pandiloski, N., Douse, C. H. & Modis, Y. Structure and functional mapping of the KRAB-

KAP1 repressor complex. *EMBO J* **41**, e111179 (2022).

20. Soufi, A., Donahue, G. & Zaret, K. S. Facilitators and impediments of the pluripotency

reprogramming factors' initial engagement with the genome. *Cell* **151**, 994–1004 (2012).

21. Taing, L. *et al.* Cistrome Data Browser: integrated search, analysis and visualization of chromatin

data. *Nucleic Acids Res* **52**, D61–D66 (2024).

22. Zheng, R. *et al.* Cistrome Data Browser: expanded datasets and new tools for gene regulatory

analysis. *Nucleic Acids Res* **47**, D729–D735 (2019).

23. Yu, X. *et al.* Recapitulating early human development with 8C-like cells. *Cell Rep* **39**, 110994 (2022).

24. Zou, Z. *et al.* Translatome and transcriptome co-profiling reveals a role of TPRXs in human zygotic

genome activation. *Science* **378**, abo7923 (2022).

25. Taubenschmid-Stowers, J. *et al.* 8C-like cells capture the human zygotic genome activation program

in vitro. *Cell Stem Cell* **29**, 449-459.e6 (2022).

26. Cacchiarelli, D. *et al.* Integrative Analyses of Human Reprogramming Reveal Dynamic Nature of

Induced Pluripotency. *Cell* **162**, 412–424 (2015).

27. Bi, Y. *et al.* Cell fate roadmap of human primed-to-naive transition reveals preimplantation cell

lineage signatures. *Nat Commun* **13**, 3147 (2022).

28. Xue, Z. *et al.* Genetic programs in human and mouse early embryos revealed by single-cell RNA

sequencing. *Nature* **500**, 593–597 (2013).

29. Wang, J. *et al.* Primate-specific endogenous retrovirus-driven transcription defines naive-like stem

cells. *Nature* **516**, 405–409 (2014).

30. Lu, X. *et al.* The retrovirus HERVH is a long noncoding RNA required for human embryonic stem cell

identity. *Nat Struct Mol Biol* **21**, 423–425 (2014).

31. Ai, Z. *et al.* Krüppel-like factor 5 rewires NANOG regulatory network to activate human naive

pluripotency specific LTR7Ys and promote naive pluripotency. *Cell Rep* **40**, 111240 (2022).

32. Gadepalli, V. S., Kim, H., Liu, Y., Han, T. & Cheng, L. XDeathDB: a visualization platform for cell death

molecular interactions. *Cell Death Dis* **12**, 1156 (2021).

33. Gao, Y. *et al.* Protein Expression Landscape of Mouse Embryos during Pre-implantation

Development. *Cell reports* **21**, 3957–3969 (2017).

34. Ralston, A. & Rossant, J. Cdx2 acts downstream of cell polarization to cell-autonomously promote

trophectoderm fate in the early mouse embryo. *Dev Biol* **313**, 614–629 (2008).

35. Lim, H. Y. G. *et al.* Keratins are asymmetrically inherited fate determinants in the mammalian

embryo. *Nature* **585**, 404–409 (2020).

36. Domingo-Muelas, A. *et al.* Human embryo live imaging reveals nuclear DNA shedding during

blastocyst expansion and biopsy. *Cell* **186**, 3166-3181.e18 (2023).

37. Falco, G. *et al.* Zscan4: a novel gene expressed exclusively in late 2-cell embryos and embryonic

stem cells. *Dev Biol* **307**, 539–550 (2007).

38. Hamatani, T., Carter, M. G., Sharov, A. A. & Ko, M. S. H. Dynamics of global gene expression changes

during mouse preimplantation development. *Dev Cell* **6**, 117–131 (2004).

39. Boussicault, L. *et al.* CYP46A1, the rate-limiting enzyme for cholesterol degradation, is

neuroprotective in Huntington's disease. *Brain* **139**, 953–970 (2016).

40. Chalhoub, G. *et al.* Carboxylesterase 2 proteins are efficient diglyceride and monoglyceride lipases

possibly implicated in metabolic disease. *J Lipid Res* **62**, 100075 (2021).

41. Artus, J., Piliszek, A. & Hadjantonakis, A.-K. The primitive endoderm lineage of the mouse blastocyst:

sequential transcription factor activation and regulation of differentiation by Sox17. *Dev Biol* **350**,

393–404 (2011).

42. Teo, A. K. K., Valdez, I. A., Dirice, E. & Kulkarni, R. N. Comparable generation of activin-induced

definitive endoderm via additive Wnt or BMP signaling in absence of serum. *Stem Cell Reports* **3**, 5–

14 (2014).

43. Messmer, T. *et al.* Transcriptional Heterogeneity in Naive and Primed Human Pluripotent Stem Cells

at Single-Cell Resolution. *Cell Rep* **26**, 815-824.e4 (2019).

44. Dubois, A. *et al.* H3K9 tri-methylation at Nanog times differentiation commitment and enables the

acquisition of primitive endoderm fate. *Development* **149**, dev201074 (2022).

45. Martello, G. *et al.* Esrrb is a pivotal target of the Gsk3/Tcf3 axis regulating embryonic stem cell self-

renewal. *Cell Stem Cell* **11**, 491–504 (2012).

46. Gassler, J. *et al.* Zygotic genome activation by the totipotency pioneer factor Nr5a2. *Science* **378**,

1305–1315 (2022).

47. Maskalenka, K. *et al.* NANOGP1, a tandem duplicate of NANOG, exhibits partial functional

conservation in human naïve pluripotent stem cells. *Development* **150**, dev201155 (2023).

48. Domcke, S. *et al.* A human cell atlas of fetal chromatin accessibility. *Science* **370**, eaba7612 (2020).

49. Evseenko, D. *et al.* Mapping the first stages of mesoderm commitment during differentiation of

human embryonic stem cells. *Proc Natl Acad Sci U S A* **107**, 13742–13747 (2010).

50. Bernardo, A. S. *et al.* BRACHYURY and CDX2 mediate BMP-induced differentiation of human and

mouse pluripotent stem cells into embryonic and extraembryonic lineages. *Cell Stem Cell* **9**, 144–

155 (2011).

51. Allshire, R. C. & Madhani, H. D. Ten principles of heterochromatin formation and function. *Nat. Rev.*

Mol. Cell Biol. **19**, 229–244 (2018).

52. Pontis, J. *et al.* Hominoid-Specific Transposable Elements and KZFPs Facilitate Human Embryonic

Genome Activation and Control Transcription in Naive Human ESCs. *Cell Stem Cell* **24**, 724-735.e5

(2019).

53. Wu, F. *et al.* Species-specific rewiring of definitive endoderm developmental gene activation via

endogenous retroviruses through TET1-mediated demethylation. *Cell Rep* **41**, 111791 (2022).

54. Macfarlan, T. S. *et al.* Embryonic stem cell potency fluctuates with endogenous retrovirus activity.

Nature **487**, 57–63 (2012).

55. Pontis, J. *et al.* Primate-specific transposable elements shape transcriptional networks during

human development. *Nat Commun* **13**, 7178 (2022).

56. Wang, H. *et al.* SVA elements: a hominid-specific retroposon family. *J Mol Biol* **354**, 994–1007

(2005).

57. Grow, E. J. *et al.* Intrinsic retroviral reactivation in human preimplantation embryos and pluripotent

cells. *Nature* **522**, 221–5 (2015).

58. Theunissen, T. W. *et al.* Molecular Criteria for Defining the Naive Human Pluripotent State. *Cell Stem*

Cell **19**, 502–515 (2016).

59. Khan, H., Smit, A. & Boissinot, S. Molecular evolution and tempo of amplification of human LINE-1

retrotransposons since the origin of primates. *Genome Res* **16**, 78–87 (2006).

60. Hazra, D., Andrić, V., Palancade, B., Rougemaille, M. & Graille, M. Formation of S. pombe Erh1

homodimer mediates gametogenic gene silencing and meiosis progression. *Sci Rep* **10**, 1034 (2020).

61. Hazra, D., Chapat, C. & Graille, M. m6A mRNA Destiny: Chained to the rhYTHm by the YTH-

Containing Proteins. *Genes (Basel)* **10**, E49 (2019).

62. Liao, S., Sun, H. & Xu, C. YTH Domain: A Family of N6-methyladenosine (m6A) Readers. *Genomics*

Proteomics Bioinformatics **16**, 99–107 (2018).

63. Liu, J. *et al.* The RNA m(6)A reader YTHDC1 silences retrotransposons and guards ES cell identity.

Nature **591**, 322–326 (2021).

64. Sun, T. *et al.* Crosstalk between RNA m6A and DNA methylation regulates transposable element

chromatin activation and cell fate in human pluripotent stem cells. *Nat Genet* **55**, 1324–1335 (2023).

65. Drakouli, S., Lyberopoulou, A., Papathanassiou, M., Mylonis, I. & Georgatsou, E. Enhancer of

rudimentary homologue interacts with scaffold attachment factor B at the nuclear matrix to

regulate SR protein phosphorylation. *FEBS J* **284**, 2482–2500 (2017).

66. Maison, C. *et al.* Higher-order structure in pericentric heterochromatin involves a distinct pattern of

histone modification and an RNA component. *Nature genetics* **30**, 329–34 (2002).

67. Xu, W. *et al.* METTL3 regulates heterochromatin in mouse embryonic stem cells. *Nature* **591**, 317–

321 (2021).

68. Huo, X. *et al.* The Nuclear Matrix Protein SAFB Cooperates with Major Satellite RNAs to Stabilize

Heterochromatin Architecture Partially through Phase Separation. *Mol Cell* **77**, 368-383 e7 (2020).

69. Obersriebnig, M. J., Pallesen, E. M. H., Sneppen, K., Trusina, A. & Thon, G. Nucleation and spreading

of a heterochromatic domain in fission yeast. *Nat Commun* **7**, 11518 (2016).

70. Murawska, M. *et al.* The histone chaperone FACT facilitates heterochromatin spreading by

regulating histone turnover and H3K9 methylation states. *Cell Rep* **37**, 109944 (2021).

71. Greenstein, R. A. *et al.* Noncoding RNA-nucleated heterochromatin spreading is intrinsically labile

and requires accessory elements for epigenetic stability. *Elife* **7**, (2018).

72. Al Adhami, H. *et al.* Systematic identification of factors involved in the silencing of germline genes in

mouse embryonic stem cells. *Nucleic Acids Res* gkad071 (2023) doi:10.1093/nar/gkad071.

73. Wu, D. & Dean, J. RNA exosome ribonuclease DIS3 degrades Pou6f1 to promote mouse pre-

implantation cell differentiation. *Cell Rep* **42**, 112047 (2023).

74. Imamura, K. *et al.* A functional connection between the Microprocessor and a variant NEXT

complex. *Mol Cell* **84**, 4158-4174.e6 (2024).

75. Yang, P., Wang, Y. & Macfarlan, T. S. The Role of KRAB-ZFPs in Transposable Element Repression and

Mammalian Evolution. *Trends Genet* **33**, 871–881 (2017).

76. Imbeault, M., Helleboid, P.-Y. & Trono, D. KRAB zinc-finger proteins contribute to the evolution of

gene regulatory networks. *Nature* **543**, 550–554 (2017).

77. Cao, J. *et al.* A human cell atlas of fetal gene expression. *Science* **370**, eaba7721 (2020).

78. Jin, Y., Tam, O. H., Paniagua, E. & Hammell, M. TEtranscripts: a package for including transposable

elements in differential expression analysis of RNA-seq datasets. *Bioinformatics* **31**, 3593–3599

(2015).

79. Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution mapping of

DNA binding sites. *Elife* **6**, e21856 (2017).

80. Katznelson, A. & Zaret, K. A fluorescence-based protocol to quantitatively titrate CUT&RUN buffer

components. *STAR Protoc* **5**, 102866 (2024).

81. Liu, N. *et al.* Direct Promoter Repression by BCL11A Controls the Fetal to Adult Hemoglobin Switch.

Cell **173**, 430-442.e17 (2018).

82. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357–359

(2012).

83. Jin, H. *et al.* ChIPseqSpikeInFree: a ChIP-seq normalization approach to reveal global changes in

histone modifications without spike-in. *Bioinformatics* **36**, 1270–1272 (2020).

84. Meers, M. P., Tenenbaum, D. & Henikoff, S. Peak calling by Sparse Enrichment Analysis for

CUT&RUN chromatin profiling. *Epigenetics Chromatin* **12**, 42 (2019).

85. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Molecular cell* **38**, 576–89

(2010).

86. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.

Nucleic Acids Res **44**, W160-165 (2016).

87. Khan, A. & Mathelier, A. Intervene: a tool for intersection and visualization of multiple gene or

genomic region sets. *BMC Bioinformatics* **18**, 287 (2017).

88. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

89. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

90. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol* **15**, 550 (2014).

91. Filippi-Chiela, E. C. *et al.* Nuclear morphometric analysis (NMA): screening of senescence, apoptosis

and nuclear irregularities. *PLoS One* **7**, e42522 (2012).

92. Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci* **31**,

8–22 (2022).

93. Lau, K. X. *et al.* Unique properties of a subset of human pluripotent stem cells with high capacity for

self-renewal. *Nat Commun* **11**, 2420 (2020).

Figure legends

Figure 1. ERH depletion reverts fibroblast H3K9me3 patterns to an hESC-like state

a. Schematic of siRNA treatments and timing for H3K9me3 CUT&RUN experiments shown
in figure 1 and SOX2 CUT&RUN and RNA-seq experiments shown in figure 2.

b. Immunofluorescence of ERH (red), H3K9me3 (red) and DAPI (blue) in siControl and
siERH human fibroblasts. Scale bars, 50µm.

c. Browser track of H3K9me3 levels in siControl and siERH treated human fibroblasts and
H1 hESCs (GSM1003585) at the megabase-domain level and gene level. Example
H3K9me3 peaks common between siERH fibroblasts and H1 hESCs are indicated by red
triangles.

d. Heatmap of fibroblast siControl and siERH H3K9me3, H1 hESC H3K9me3
(GSM1003585) and H1 hESC HP1α (GSE123229) levels centered at siERH retained
H3K9me3 peaks located within fibroblast H3K9me3 domains +/-10kb.

e. Correlation of H3K9me3 from BJ fibroblasts after siControl or siERH with H1 hESC
H3K9me3.

f. Representative images showing hESC morphology and SOX2 (green) pluripotency
marker staining in siControl and siERH H1 hESCs under pluripotency maintaining
conditions. Scale bars, 50µm.

g. Representative images of H3K9me3 (red) and DAPI (blue) in siControl and siERH H1
hESCs under pluripotency maintaining conditions. Scale bars, 50µm.

h. CistromeDB Toolkit enrichment analysis of fibroblast siERH H3K9me3 peaks. The top 20
proteins scored by the analysis are visualized. Proteins colored by characterization as
H3K9me3-associated (orange), KRAB zinc-finger transcription factors (purple), and
euchromatin-associated (gray).

i. Browser track comparing siControl and siERH fibroblast H3K9me3 with H1 hESC
H3K9me3 (GSM1003585) and HP1a (GSM1701825) patterns. Peak locations called by
SEACR analysis on siERH H3K9me3 data are shown.

See also Extended Data Fig. 1 and Supplementary Table 1.

Figure 2. ERH depletion enhances gene activation of pluripotency and alternative lineage genes
during iPSC reprogramming

    a. Heatmap of H3K9me3 CUT&RUN from siControl and siERH human fibroblasts and SOX2
721
722   CUT&RUN from siControl and siERH fibroblasts + OSKM at siERH-gained SOX2 peaks.

    b. Browser track of example hESC SOX2 and NANOG peak coincident with fibroblast siERH
723
724   H3K9me3 loss and SOX2 peak gain.

    c. Volcano plot of expression changes (DESeq2) in siERH vs siControl fibroblasts+OSKM of
725
726   iPSC activated genes located in fibroblast H3K9me3 domains.

    d. Expression of ERH and example naïve pluripotency genes activated in 48 hours OSKM
727
728   by ERH depletion, transcripts per million (TPM) values from RNA-seq replicates shown
729   (DESeq2 adjusted p-value, n=2).

    e. Profile plots of H3K9me3 and RNA levels from LTR7 and HERVH repeats in siERH vs
730
731   siControl fibroblasts+OSKM.

    f. Profile plots of H3K9me3 and RNA levels from LTR7Y repeats in siERH vs siControl
732
733   fibroblasts+OSKM.

    g. Browser track showing H3K9me3 in siControl and siERH fibroblasts and RNA levels in
734
siControl+OSKM and siERH+OSKM for gene and adjacent LTR7Y naïve pluripotency
repeat.

See also Extended Data Fig. 2-4 and Supplementary Tables 2-4.

Figure 3. ERH is expressed during pre-implantation mouse development and is critical for normal
blastocyst development by repressing 2-cell and alternative lineage genes.

a. ERH (white) and DAPI (blue) staining during mouse pre-implantation development. Representative confocal z-sections are shown (zygote n=15, N=3, 2-cell stage n=18, N=3, 8-cell stage n=6, N=2, blastocyst n=10, N=3). Scale bars, 10μm.

b. Representative images of trophectoderm marker Cdx2 (green) in siControl and siERH blastocysts with Phalloidin (orange) marking cell boundaries. Scale bars, 10μm.

c. Quantification of CDX2 intensity in siControl vs siERH blastocysts (unpaired t-test, n=8 blastocysts for each siRNA).

d. Representative image of trophectoderm marker KRT8 (orange) in siControl and siERH blastocysts with DAPI (blue). Scale bars, 10μm.

e. Quantification of KRT8 intensity in siControl vs siERH blastocysts (unpaired t-test, n=9 blastocysts, n=8 blastocysts for each siRNA).

f. Representative images of H3K9me3 (magenta), DAPI (teal) in siControl vs siERH blastocysts. Scale bars, 10μm.

g. Quantification of H3K9me3 level in siControl vs siERH blastocysts. (unpaired t-test, n=8 for siControl and n=9 for siERH blastocysts).

h. Percentage of cells per blastocyst with abnormal nuclei. (unpaired t-tes, n=11 for siControl and n=9 for siERH blastocysts).

i. Representative image of DAPI staining siCTRL and siERH blastocysts with white arrows indicating DNA shedding.

j. ERH repressed blastocyst genes (adjusted p-value<=0.05, log2fc>=0.5) grouped based upon normal expression patterns exhibited during mouse pre-implantation development re-analyzed from Gao et al., 2017[33].

k. Example gene tracks of genes significantly upregulated in siERH vs siControl blastocysts (adjusted p-value<=0.05, log2fc>=0.5, n=3) with H3K9me3 tracks for 2 cell, trophectoderm and ICM cells from[11].

See also Extended Data 5 and Supplementary Tables 5,6.

Figure 4. ERH represses pluripotency and alternative lineage gene activation during hESC
differentiation

a. Representative images of NANOG (red), SOX17 (red) and DAPI (blue) during 1 day of
endoderm differentiation of siControl and siERH H1 hESCs. Scale bars, 50μm.

b. Representative images of OCT4 (red), PAX6 (green) and DAPI (blue) during 2 days of
ectoderm differentiation of siControl and siERH H1 hESCs. Scale bars, 50μm.

c. Plot of NANOG expression in transcripts per million (TPM) by RNA-seq in siControl and
siERH treated H1 hESCs cultured in pluripotency (H1), 1 day endoderm differentiation
(DE) and 2 day ectoderm differentiation (EC). (DESeq2 adjusted p-value reported).

d. Plot of KLF5 and ZSCAN4 expression in transcripts per million (TPM) by RNA-seq in
siControl and siERH treated H1 hESCs cultured in pluripotency (H1), 1 day endoderm
differentiation (DE) and 2 day ectoderm differentiation (EC). (DESeq2 adjusted p-value
reported).

e. GO   analysis   (PantherDB,   statistical   overrepresentation)   of   genes   upregulated
(padj<=0.05, log2 fold change >= 0.5) in siERH vs siControl H1 hESCs 1 day endoderm
differentiation.

f. Radar plot showing cumulative expression profile across indicated human tissues, data
downloaded from BBI Descartes database[77], for genes upregulated (padj<=0.05, log2 fold
change >= 0.5) in siERH vs siControl H1 hESCs 1 day endoderm differentiation.

g. Radar plot showing cumulative expression profile across indicated human tissues, data
downloaded from BBI Descartes database[77], for genes upregulated (padj<=0.05, log2 fold
change >= 0.5) in siERH vs siControl H1 hESCs 2 day ectoderm differentiation.

h. Example early mesodermal genes activated during endoderm differentiation by ERH
depletion, transcripts per million (TPM) values from RNA-seq replicates shown (DESeq2
adjusted p-value).

i. Example late mesodermal genes activated during endoderm differentiation by ERH
depletion, transcripts per million (TPM) values from RNA-seq replicates shown DESeq2
adjusted p-value).

j. Representative image of DAPI (blue), BRACHYURY (red) of siControl and siERH treated
H1 hESCs after 1 day of endoderm differentiation. Scale bars, 50μm.

For all RNA-seq TPM values n=3, except for siCTRL UT where n=2. See also Extended
Data Fig. 6 and Supplementary Tables 7,8.

Figure 5. ERH represses evolutionarily young transposable elements with regulatory function

a. H3K9me3 data from H1 hESCs (GSM1003585) and hESC derived endoderm
(GSM916057) at 65,656 L1 element loci +/- 1kb upregulated (adjusted pvalue<=0.05;
log2foldchange>=0.5) by ERH depletion during 1 day endoderm differentiation.

b. Venn diagram showing intersect of repeat classes upregulated (TETranscripts, padj<0.05,
log2fc>0.5) by ERH depletion in H1 hESCs untreated (UT), 1 day endoderm (DE) or 2 day
ectoderm (EC) conditions.

c. Expression change and significance, TEtranscripts[78] analysis, of L1 LTR transposable
elements in ERH depletion in H1 hESCs untreated (UT), 1 day endoderm (DE) or 2 day
ectoderm (EC) listed in order of evolutionary age as determined in[59].

d. Example browser track showing expression of L1 element and proximal gene in siControl
and siERH treated H1 hESCs in the pluripotent and 1 day DE conditions.

e. Example browser track of gene with promoter proximal repeat element gaining H3K9me3
during endoderm differentiation that is upregulated by ERH depletion.

f. H3K9me3 CUT&RUN signal at L1 elements that exhibited 2-fold RNA increase in siERH
vs siControl H1 hESC endoderm and switched from H3K9me3 unmarked in pluripotent to
marked in endoderm.

g. Browser track showing RNA-seq of example repeat element and nearby genes in siControl
and siERH treated H1 hESCs after 1 day of endoderm differentiation and H3K9me3 tracks
from H1 hESCs (GSM1003585) and hESC derived endoderm (GSM916057).

See also Extended Data Fig. 7,8 and Supplementary Table 10.

Figure 6. Cell type specific and independent recruitment of ERH to chromatin by RNA binding

proteins regulates gene and repeat repression

a. Strategy for identifying chromatin-bound ERH interaction partners by isolation of soluble

nucleoplasm and chromatin-associated protein fractions prior to immunoprecipitation.

b. Immunoprecipitation (IP) of ERH from the chromatin fraction of human fibroblasts.

Western blot analysis of YTHDC1, YTHDC2, SAFB1, and ERH in the input and IP

samples.

c. Immunoprecipitation (IP) of ERH from the chromatin fraction of human ESCs. Western

blot analysis of YTHDC1, YTHDC2, SAFB1, and ERH in the input and IP samples.

d. Immunofluorescence (IF) of ERH after RNase A and RNase H treatment.

e. Western blot of ERH, YTHDC1, and YTHDC2 in nucleoplasm and chromatin-associated

protein fractions of human fibroblasts after siYTHDC1 and siYTHDC2 treatment.

f. Expression change and significance, TEtranscripts[78] analysis, of HERVK HERVH and

satellite repeats in hESCs and human fibroblasts after siERH.

g. Immunofluorescence (IF) of BRACHYURY in siCTRL, siYTHDC1 and siYTHDC2 after 1

day endoderm differentiation.

h. Venn diagram of gene coregulation after siERH, siYTHDC1 and siSAFB1 in human

fibroblasts after addition of hepatic reprogramming factors, FOXA3, HNF1A, HNF4A.

i. Proposed model for cell-type specific, and cell-type agnostic gene and repeat regulation

by ERH, YTHDC1, and SAFB1.

See also Extended Data Fig. 9

Methods

Cell lines culture and condition

Human H1 ES cells were obtained from WiCell (WA01), grown on Matrigel coated wells and

cultured in mTeSR1 (Stem Cell Tech, 85850) medium at 37^{o}C and 5% CO_2. Medium was

changed daily. Human BJ foreskin fibroblasts were obtained from ATCC (CRL-2522) and

cultured in Eagle's Minimum Essential Medium (EMEM, Sigma-Aldrich M2279) supplemented

with 10% fetal bovine serum (FBS, Hyclone SH30071) and 2mM Glutamax (ThermoFisher

35050061) at 37^{o}C and 5% CO_2.

Mouse embryo work

Mouse embryo experimentation was approved by the Biological Resource Center Institutional

Animal Care and Use Committee (IACUC), Agency for Science, Technology and Research

(IACUC Protocol 806983). C57BL/6 wild-type 3–4-week-old female mice were superovulated

using 5 IU of pregnant mare serum (PMS, National Hormone and Peptide Program)

gonadotropin given intraperitoneally and 5 IU of recombinant chorionic gonadotrophin (CG,

National Hormone and Peptide Program) given 48 h after and immediately before mating,

according to animal ethics guidelines of the University of Pennsylvania. Embryos were flushed

from oviducts of plugged females using M2 medium (Merck) and cultured in KSOM+AA (Merck)

covered by mineral oil (Sigma, M-5310), at 37 °C and 5% CO2.

Mouse embryo immunofluorescence and microscopy

Embryos were fixed in 4% paraformaldehyde (Electron Microscopy Sciences, 15710) for 30 min

at room temperature washed twice in PBS with 0.1% Triton X-100 (Millipore Sigma, T8787),

permeabilized for 20 min in PBS with 0.5% Triton X-100, and blocked in 3% bovine serum

albumin (Gemini Bio, 700-100P) for 30 min. Embryos were then incubated at 4 °C overnight in

primary antibodies diluted in blocking solution at the following concentrations: ERH (Abcam,

ab166620) at 1:200, Keratin-8 (DSHB, THROMA-I) at 1:20, H3K9me3 (Abcam, ab8898) at

1:1000, and CDX2 (Invitrogen, ZC007) at 1:200. After primary antibody incubation, embryos

were washed 5 times for 20 min in PBS with 0.1% Triton X-100 and incubated for 30 min in

Alexa Fluor-conjugated secondary antibodies (Invitrogen, A32731, A32723, A32733, A32728

and A11006) diluted in blocking solution to 1:500. Phalloidin-Alexa Fluor 555 (Invitrogen,

A34055) diluted to 1:500 and DAPI (Sigma, 10236276001) diluted to 1:500 in blocking solution

were also used to label F-actin and DNA, respectively. Embryos were washed three times in

PBS with 0.1% Triton X-100 before mounting in PBS covered with mineral oil in an 8-well μ-

Slide (Ibidi, #80827). Embryos were imaged using a laser-scanning confocal microscope (Leica

SP8) with a water-immersion Apochromat 40× 1.1 NA objective and HyD detectors. Samples

were excited with standard laser lines (405-nm, 488-nm, 552-nm, 638-nm).

Mouse embryo siRNA injection experiments

Microinjections were performed using a FemtoJet (Eppendorf). For ERH knockdown

experiments, mouse embryos were injected with two siRNAs (Qiagen, S104403294 &

S104403287) against ERH at a total concentration of 400 nM. For knockdown experiments

AllStars negative control siRNA (Qiagen, 1027281) was injected in embryos at a concentration

of 400nM.

Differentiation of hESCs

Prior to differentiation, for samples to be analyzed by RNA-seq, cells grown to a density of 80%

were digested with Accutase (Stem Cell Technologies, 07920) for 7 min. Then, DMEM/F12 was

added and gently pipetted to generate a cell suspension. After centrifugation, cells were

resuspended in mTeSR1 medium with 5uM ROCK inhibitor Y-27632 (Cayman Chemical,

10005583) and seeded at $8X10^5$-$1X10^6$ cells per well in a 6 well plate coated with Matrigel

(Millipore Sigma, CLS354277) 24 hours prior to differentiation treatment. For definitive

endoderm differentiation of hESCs, mTeSR1 medium was removed and cells were cultured in

DMEM/F12 (Thermo Fisher, 11320033) supplemented with 10% (v/v) Fetal Bovine Serum

(Hyclone, SH30071) with 100ng/ml Activin A (Peprotech, 120-14P) and 3uM CHIR99021

(Cayman Chemical, 13122). For ectoderm differentiation of hESCs, mTeSR medium was

removed and cells were cultured in DMEM/F12 with 5uM Retinoic Acid (Cayman Chemical,

11017).

Cell culture siRNA transfection experiments

All ERH knockdown experiments were performed with two cycles of siRNA transfection

(Silencer Select siRNA, Thermo Fisher, s4815), three days apart. Transfections were performed

in fibroblasts using 5nM final concentration of siRNA and in hESCs using a 10nM final

concentration of siRNA and 2.5ul/ml final concentration of Lipofectamine RNAiMAX (Thermo

Fisher #31985070).

Reprogramming factor expression experiments

We utilized the CytoTune-iPS 2.0 Sendai reprogramming kit (ThermoFisher, A16517) to express

Oct4, SOX2, KLF4 and cMyc in BJ fibroblasts. Cells were treated according to the

manufacturer's instructions three days after the second of two cycles of siRNA transfection.

Cell culture immunofluorescence and microscopy

For immunofluorescence of human fibroblasts and hESCs cells were washed briefly with PBS,

and fixed in 4% paraformaldehyde in PBS for 10 minutes. Fixed cells were washed 3 times with

PBS, permeabilized with ice-cold 0.1% Triton-X in PBS for 10 minutes, and washed twice with

PBS-T (PBS, 0.05% Tween-20). Samples were blocked with 3% donkey serum (Millipore

Sigma, D9663) in PBS for 2 hours at room temperature and incubated overnight at 4°C using

the following concentrations: anti-H3K9me3 (1:300, Abcam ab8898), anti-H3K27me3 (1:300,

Active Motif, 39155), anti-SOX2 (1:200, Cell Signaling, 23064), anti-OCT4 (1:200, Abcam,

ab19857), anti-PAX6 (1:200, BD Biosciences, 561462), anti-Brachyury (1:200, Cell Signaling,

81694) in 3% donkey serum in PBS. Cells were washed 3 times with PBS-T and then incubated

with AlexaFluor 488- or 594-conjugated secondary antibodies raised in donkey (Thermo Fisher,

See Methods) at a 1:500 dilution in 3% donkey serum in PBS.

Total RNA isolation and sequencing

BJ fibroblasts grown in a 6-well dish were washed twice with cold PBS, followed by addition of 1

mL TRIzol reagent (ThermoFisher 15596026) and transferred to a chilled 1.5 mL tube using a

cell scraper. The cells were homogenized by pipetting through a 25G needle attached to a 2 mL

syringe 10 times. 200 µl of chloroform was added to the cells and vortexed for 15 seconds,

followed by a room temperature incubation of 3 minutes. Phase separation of nucleic acids and

protein was performed by centrifugation at 12,000 x g for 15 minutes at 4C. The upper aqueous

phase containing RNA was then transferred to a fresh chilled tube. RNA was precipitated from

the aqueous phase by adding an equal volume of isopropyl alcohol, followed by a 10 minute

incubation at room temperature. The RNA was pelleted by centrifugation at 12,000 x g for 10

minutes at 4C. After removal of supernatant, the pellet was washed twice with 1 mL 75%

ethanol, and resuspended in nuclease free water. RNA quality was verified using a high-

sensitivity RNA screentape (Agilent 5067-5576). 1ug of total RNA was sent to Novogene, who

assembled and sequenced bulk RNA-seq libraries. Two to three biological replicates were

prepared for each condition.

CUT&RUN experiments

CUT&RUN was performed as described in[79], with digitonin and NaCl in wash buffers optimized

for permeabilization of BJ fibroblasts and retainment of SOX2 on chromatin as performed in[80].

Next, BJ fibroblasts were detached from cell culture plates with Accutase (STEMCELL

Technologies 07920), washed, bound to magnetic concanavalin A beads, and permeabilized with

a dig-wash buffer (0.1% digitonin, 20 mM HEPES pH 7.5, 75 mM NaCl, 0.5 mM Spermidine,

EDTA-free protease inhibitor). Bead-bound cells were incubated with a SOX2 (CST, clone

D9B8N, 23064, 1:100) or H3K9me3 antibody (abcam ab8898, 1:100) at 4C overnight. The

following morning, the cells were washed twice with dig-wash buffer, incubated with pA/G-MNase

for an hour at 4C, and washed twice more. After chilling cells on an ice block for 5 minutes, MNase

was activated for 30 minutes by addition of 2mM $CaCl_2$. The reaction was stopped by adding an

equal volume of 2X STOP buffer (340 mM NaCl, 20 mM EDTA, 4 mM EGTA, 0.1% Digitonin, 0.05

mg/mL glycogen, 5 mg/mL RNase A). Digested chromatin was extracted from permeabilized cells

at 37C for 30 minutes, and DNA was purified by phenol-chloroform. Libraries were prepared as

previously described[81]. Two to three biological replicates were prepared for each condition.

CUT&RUN data processing

Paired-end CUT&RUN reads were mapped to the human genome build hg38 using bowtie2[82]

(v2.3.4.1) with run parameters --local --very-sensitive --no-unal --no-mixed --no-discordant -I 10 -

X 1000. A quality-filtered BAM file was generated with the command samtools (v1.16.1) view -q

5 -f 2 -bS. Global scaling factors (SF) for siERH and siControl H3K9me3 data sets were calculated

using the default parameters of ChIPSeqSpikeInFree[83] (R v3.6.3). Bigwig files for visualization

were generated from indexed BAM files with command bamCoverage --smoothLength 10 --

normalizeUsing CPM --scaleFactor 1/SF --ignoreForNormalization chrM (deeptools 3.5.0). BED

files were generated from BAMs using the bedtools command bamToBed (bedtools v2.27.1). BED

files were converted to bedgraph format using the bedtools command genomecov. Peaks were

called on the bedgraph files with SEACR[84] (v1.3), using the stringent setting and selecting the top

0.01% of regions by AUC. Peak lists for downstream analyses were derived from individual

replicates or the union of overlapping sites from biological triplicates, as indicated in the text and

figures. Enrichment analysis of known DNA binding motifs was performed using HOMER[85].

Heatmaps

973 Heatmaps were generated with deeptools[86] version 3.5.0. Counts matrices were generated with

974 command computeMatrix reference-point --missingDataAsZero --referencePoint center. Figures

975 were produced with the plotHeatmap command.

Correlation plots

To assess genome wide similarity between hBJ siERH H3K9me3, hBJ siControl H3K9me3, and

H1-hESC H3K9me3 data sets, the Pearson correlation of peak overlap was calculated using the

Intervene tool[87]. In brief, peak files for individual H3K9me3 replicates were processed with the

command intervene pairwise --I *H3K9me3.bed --compute fract --htype color. Visualization of

Pearson correlation of fractional overlap between peak sets were generated on

https://intervene.shinyapps.io/intervene.

RNA-seq data analysis

RNA-seq reads (paired-end) were mapped to the mm10 or hg38 genome for mouse and human

experiments respectively using STAR[88] (2.7.10b). The mapped reads to genes were counted

using featureCounts[89]. For counting repeat mapping reads the TEcount function from

TEtranscripts[78] was run on STAR aligned bam files using the hg38 repeatmasker gtf file

annotation. The output count files were analyzed by DESeq2[90] to detect differentially expressed

genes (Benjamini adjusted p-value < 0.05, log2 fold change > 1 or log2 fold change > 1.5).

Genes and repeats with fewer than 10 total mapped reads were excluded from DESeq2 and

TETranscripts analysis. TPM, FPKM and RPM were calculated using an in-house R script.

Mouse blastocyst image analysis

3D visualizations and analysis of embryos were performed using Imaris 8.2 or Imaris 9.7

software (Bitplane). The manual surface rendering module was used for cellular and nuclear

segmentation, using phalloidin and DAPI signals to define cellular and nuclear boundaries,

respectively. Quantifications of immunofluorescence intensity were performed by measuring the

mean fluorescence intensity of voxels within the segmented object, standardized DAPI to

account for light scattering.

1002

Cell culture image analysis

Quantification of nuclear area and signal intensity was performed using the Nuclear

Morphometric Analysis (NMA) plugin for ImageJ[91]. In ImageJ saturated pixels was set to 0.35%

prior to processing. NMA plugin was run using the following settings: show ellipse and use

watershed options selected, minimum nuclei area=150, maximum nuclei area=6000, nuclei

intensity segmentation threshold=95, nuclei insertion intensity=93%.

1009

1010 **Gene ontology enrichment analysis**

1011 Gene ontology enrichment analysis was performed using PANTHER[92] (v18.0) with settings,

1012 statistical overrepresentation test: GO biological process complete, all genes in the database

1013 were used as the background reference list, and statistical significance was determined using

1014 Fisher's Exact test with false discovery rate.

1015

Statistical analysis

Statistical analyses were performed in GraphPad Prism, the R software for statistical computing

and with DESeq2. Pairwise comparisons were analyzed using an unpaired, two-tailed Student's

t-test. Correlation was analyzed with a two-tailed Pearson's Correlation test. Significant overlap

between gene sets was analyzed with the hypergeometric test. Differential gene expression

analysis was performed using DESeq2 which calculates p-values by the Wald test and adjusted

p-values corrected for multiple testing using the Banjamini and Hochberg method. Reproducibility

was confirmed by independent experiments.

1024

1025   **Acknowledgements**

1026   R.L.M. was supported by an NIH NIDDK K01 postdoctoral fellowship DK117970-01. N.P. was

1027   supported by NIGMS (GM139970-01) and NICHD (HD102013-01A1). R35GM153180 supported

1028   work in the lab of K.S.Z.

1029

1030   **Author contributions**

1031   Conceptualization, R.L.M. and K.Z.; Formal analysis, A.K., B.H., H.F., K.T. and R.L.M.; Funding

1032   acquisition, K.Z; Investigation, A.K., B.H., H.F., K.T., J.Z., A.B. and R.L.M.; Methodology, A.K.,

1033   B.H., R.L.M. and K.Z.; Project administration, R.L.M and K.Z.; Resources, K.Z,; Supervision, M.T-

1034   P., N.P., R.L.M. and K.Z.; Writing – Original Draft, A.K., R.L.M. and K.Z.

1035

1036   **Declaration of Interests**

1037   The authors declare no competing interests.

1038

1039   **Corresponding author**

1040   Correspondence to Ryan McCarthy (ryan.mccarthy@seatttlechildrens.org) and Kenneth Zaret

1041   (zaret@pennmedicine.upenn.edu).

1042

1043   **Data availability**

1044   • RNA-seq and CUT&RUN datasets collected in this study have been deposited in the

1045   GEO of NCBI and are publicly available as of the date of publication. Accession

1046   numbers are listed in the Methods. GEO reviewer tokens for accessing the data

1047    while it remains in private status are for RNA-seq GSE268901 (wbsvkukqhzobpan),

1048    and for CUT&RUN GSE268961 (qngpsiywdvyfjyj).

1049

1050 **Code availability**

1051  •   This paper does not report original code.

1052  •   Any additional information required to reanalyze the data reported in this paper is

1053   available from the lead contact upon request.

1054

1055

1056 **Supplementary Tables**

1057 Supplementary Table 1. H3K9me3 peak coordinates in siERH human fibroblasts.

1058 Supplementary Table 2. Fibroblast +OSKM 48 hours, SOX2 CUT&RUN peaks in siControl and

1059 siERH.

1060 Supplementary Table 3. Fibroblast +OSKM 48 hours, siERH/siControl RNA-seq DESeq2

1061 differential gene expression analysis results.

1062 Supplementary Table 4. Fibroblast +OSKM 48 hours, siERH/siControl TETranscripts differential

1063 repeat expression analysis results.

1064 Supplementary Table 5. Blastocyst immunofluorescence quantification values.

1065 Supplementary Table 6. Blastocyst RNA-seq DESeq2 siERH/siControl differential gene

1066 expression analysis results.

1067 Supplementary Table 7. List of genes categorized as group 1-4 by preimplantation expression

1068 from Gao et al., 2017[33] and TPM values from this study's siControl and siERH mouse blastocysts

1069 RNA-seq.

1070 Supplementary Table 8. H1 hESC untreated, endoderm and ectoderm RNA-seq DESeq2

1071 siERH/siControl differential gene expression analysis results.

1072   Supplementary Table 9. H1 hESC untreated, endoderm, and ectoderm TETranscripts

1073   siERH/siControl differential repeat expression analysis

1074   Supplementary Table 10. H1 hESC untreated, endoderm and ectoderm GO analysis results.
1075

Extended data

Extended Data Fig. 1. ERH depletion reverts fibroblast H3K9me3 patterns to an hESC-like state

a. Browser tracks comparing H3K9me3 distribution patterns in human fibroblasts compared
to hESC.

b. Venn diagram of H3K9me3 peaks called using SEACR from fibroblast siControl and
siERH H3K9me3 CUT&RUN and H1 hESC ChIP-seq (GSM1003585).

c. Browser track showing SEACR peaks called in siControl H3K9me3 domain.

d. Nanog staining in hESCs treated for 5 weeks with siRNAs to sustain ERH depletion or
control.

e. Immunofluorescence for SOX2 (green) and H3K9me3 (red) in siControl and siERH H1
hESCs in pluripotent conditions at higher magnification to show nuclear organization.
Scale bars, 50μm

f. Immunofluorescence for DAPI (blue) and H3K27me3 (red) in siControl and siERH H1
hESCs in pluripotent conditions. Scale bars, 50μm.

g. Nuclear area, white arrows indicate nuclear blebs. $p=5.6*10^{-16}$, welch's two sample t-test.
Scale bars, 50μm.

h. Expression (TPM) of H3K9me3 HMTs in siControl and siERH H1 hESCs (n=2 for siControl
H1 hESCs and n=3 for siERH H1 hESCs).

i. Browser tracks showing H3K9me3 in human fibroblasts or hESCs treated with siControl
or siERH.

j. Correlation of H3K9me3 from hESCs after siControl or siERH.

k. Browser tracks showing H3K9me3 (GSM1003585) and HP1a (GSM1701825) in H1
hESCs and H3K9me3 in siControl and siERH human fibroblasts.

l. Top 10 statistically enriched transcription factor binding motifs at H3K9me3 peaks in
siERH treated human fibroblasts (homer findMotifs).

m. Occurrence of repeats from RepEnrich database and repeat occurrence at sites of siERH
fibroblasts H3K9me3 peaks.

Katznelson et al., Extended Data Figure 1

a H3K9me3 distribution in fibroblasts and hESCs

chr3:179,676,569-182,850,262   3,173 kb

Fibroblast H3K9me3   [0 - 0.10]

H3K9me3 domain

H1 hESC H3K9me3   [0 - 18]

H3K9me3 peak

b Venn diagram of overlap of H3K9me3 peaks called by SEACR in H1 hESCs, and siControl and siERH treated fibroblasts

Fib. siERH H3K9me3
1,057
5,748
475
H1 hESC H3K9me3
127,969
3,246
18,901
4,752
Fib. siControl H3K9me3

c Browser track showing limitations of peak calling to capture domain architecture of H3K9me3

megabase H3K9me3 domain   5 mB window

hFib. siControl H3K9me3

H3K9me3 peaks   ← Many SEACR-called peaks in single broad domain

genes   7633 5843 9645 9454 7620 13797 15712 14958 8623 14958 7203 21211

134620.1   MEAT6   C6orf118   LINC00473   PRR18

d Pluripotency maintenance in hESCs after repeated siRNA treatments

Weekly siRNA treatment schedule
siRNA siRNA
Mon Wed
repeated 5x

DAPI   NANOG
pluripotent conditions siControl
pluripotent conditions siERH

e Higher magnification showing nuclear organization of SOX2 and H3K9me3

SOX2   H3K9me3
pluripotent conditions siControl
pluripotent conditions siERH

f H3K27me3 immunofluorescence in siControl and siERH H1 hESCs

DAPI   H3K27me3
pluripotent conditions siControl
pluripotent conditions siERH

g Quantification of nuclear area in siControl and siERH H1 hESCs

pluripotent conditions siControl
pluripotent conditions siERH

count   p=5.6e-16
siControl
siERH
nuclear area

i chr13:102,905,680-103,325,317   419 kb

hFib.
siControl H3K9me3   [0 - 0.10]
siERH H3K9me3   [0 - 0.10]

H1 hESC
siControl H3K9me3   [0 - 10.00]
siERH H3K9me3   [0 - 10.00]

RefSeq genes

h H3K9me3 HMT expression in H1 hESCs

SETDB1   SUV39h1   SUV39h2
Expression (TPM)
siControl siERH   siControl siERH   siControl siERH

l Browser track of H3K9me3 in siControl and siERH compared to H3K9me3 and Hp1a in H1 hESC

chr2:195,052,698-195,952,928   900 kb

H3K9me3 Fibroblast siControl
                       siERH
HP1a H1 hESC

H3K9me3 peaks
siERH fibroblasts:

j
siControl rep1
siERH rep1
hESC
siControl rep2
siERH rep2

siControl rep1 siERH rep1 siControl rep2 siERH rep2
hESC

Correlation of peak overlap
-1 -0.5 0 0.5 1

k Top 10 statistically enriched transcription factor motifs at siERH fibroblast H3K9me3 peaks

	p-adj	Motif
ZNF136	<10⁻⁵	
SOX10	<10⁻⁵	
ZFX	<10⁻⁵	
SOX21	<10⁻⁵	
SOX3	<10⁻⁵	
SOX4	<10⁻⁵	
ZNF711	<10⁻⁵	
ZNF317	<10⁻⁵	
SOX15	<10⁻⁵	
SOX9	<10⁻⁵	

m Proportion of repeat class at siERH fibroblast H3K9me3 peaks vs background abundance

background repeat frequency   siERH H3K9me3 peaks

DNA
LINE
Simple
LTR
Satellite
SINE
snRNA

1103   Extended Data Fig. 2. Kinetics of iPSC gene activation is effected by heterochromatin marks.

1104   **a.** Expression of genes activated during fibroblasts to iPSC reprogramming timepoints[26] that

1105   are located in unmarked, H3K27me3 or H3K9me3 marked heterochromatin as annotated

1106   by Becker et al., 2017[5].

1107

a delayed activation iPSC genes in fibroblast H3K9me3 domains during reprogramming

1108

Extended Data Fig. 3. ERH depletion enhances SOX2 binding in H3K9me3 domains.

a. Venn diagram showing overlap between SOX2 peak calls in siControl and siERH
fibroblasts+OSKM.

b. Browser track of example hESC SOX2 and NANOG peak coincident with fibroblast siERH
H3K9me3 loss and SOX2 peak gain.

c. Characterization by fibroblast H3K9me3 status of SOX2 binding sites gained in siControl
and siERH fibroblasts+OSKM.

d. SOX2 CUT&RUN average profile in siControl and siERH fibroblasts+OSKM for genome
regions identified as being differentially bound by reprogramming transcription factors [20].

e. Rank ordered GIGGLE enrichment scores. Each circle indicates similarity between siERH
SOX2-gained sites and a ChIP-seq peak set from cistromeDB. Blue circles are peak sets
profiled in hESCs, white circles are peaks from a non-embryonic origin.

f. Browser track of H3K9me3 in siControl and siERH fibroblasts and RNA in siControl and
siERH fibroblasts+OSKM at DPPA4.

g. KLF4 and OCT4 footprint analysis at SOX2 siERH gained peaks.

1124

a overlap of SOX2 peaks in siERH and siControl

b gained SOX2 binding in H3K9me3 after siERH

c ERH siRNA enables SOX2 binding in H3K9me3

d SOX2 binding in differentially bound regions enhanced by siERH

e Similarity of siERH gained SOX2 peaks to hESC and non-hESC ChIP-seq

f chr3:109,308,070-109,329,272

g CUT&RUN footprints at Sox2 siERH gained peaks

Extended Data Fig. 4. ERH depletion enhances gene activation of pluripotency

a. Browser track of H3K9me3 loss and gene activation of DPPA4 in siERH treated human
fibroblasts+OSKM.

b. Example 8 cell genes activated in siERH+OSKM (n=2 per condition) but not activated
during eight time points of iPSC reprogramming time course[26].

c. Comparison of expression (FPKM) of naïve genes KLF5 and DNMT3L of siControl and
siERH fibroblasts+OSKM from this study (n=2 per condition) with pluripotent and naïve
human ESC data from Bi et al., 2022.

d. Browser tracks of H3K9me3 in siControl and siERH fibroblasts and RNA in siControl and
siERH fibroblasts+OSKM, at a locus of an LTR7Y-HERHV transposable element and
nearby gene.

e. Volcano plot showing differential repeat expression in siERH vs siControl
fibroblast+OSKM for repeats annotated in[6] as repressed by H3K9me3 in human
blastocysts. (TETranscripts, DESeq2, padj<=0.05, log2fc>0.5).

f. Representative phase image of siControl and siERH treated fibroblasts+OSKM 2 days
after infection with OSKM sendai virus.

g. Representative phase image showing cell confluency of siControl and siERH
fibroblasts+OSKM cells 2 weeks after initial OSKM infection with titrated siRNA
concentration. All treatments were plated at identical cell numbers 1 day prior to infection.

h. Plot of iPSC colony count from siControl and siERH fibroblasts+OSKM cells 3 weeks after
initial OSKM infection with titrated siRNA concentration (n=2 per condition). Each value
represents the number of colonies generated in 1 well of a 24 well plate.

i. Alkaline phosphatase staining of siControl and siERH +/- sip53 treated fibroblasts+OSKM
21 days after infection.

j. Expression (TPM) of RIPK2 in siControl and siERH fibroblasts+OSKM 48 hours after
infection (n=2 per condition).

k. Plot of iPSC colony count 3 weeks after initial OSKM infection for siControl and siERH
cells reprogrammed in media containing the inhibitors GSK2983559, Ac-DEVD-CHO and
GSK872 (n=4 per condition).

l. Plot of iPSC colony count (n=2 per condition) and imaging of siERH fibroblasts+OSKM 3
weeks after infection and treated with indicated combinations of inhibitors or DMSO
control.

1157

a Browser track of H3K9me3 and expression change at example naïve pluripotency gene

b Expression of naïve pluripotency genes during iPSC reprogramming and in siControl and siERH fibroblasts+OSKM

c Average expression in plutipotent (pESC) and naive (nESC) cells from Bi et al., 2022 compared to siCTRL and siERH +OSKM fibroblasts

d H3K9me3 and expression change at example gene and proximal repeat

e Expression changes of human blastocyst repressed repeats in siERH vs siControl fibroblasts+OSKM

f 2 days post infection

g 7 days post infection

h iPSC colony number 14 days post infection

i

j

k +3 inhibitors (GSK2983559 Ac-DEVD-CHO GSK872)

l

1158   Extended Data Fig. 5. Characterization of ERH and the effects of ERH depletion in mouse
1159   blastocysts

1160   **a.** ERH RNA expression in reads per kilobase million (RPKM) during stages of mouse pre-
1161   implantation development, RNA-seq data from[33]. unpaired two tail t-test; n=6.

1162   **b.** Fluorescent imaging of ERH (magenta) and quantification in ICM and trophectoderm cells
1163   (n=12 for ICM and n=13 for Trophectoderm, unpaired t-test, ns=not significant).

1164   **c.** ERH (magenta) and DAPI (teal) staining in mouse blastocysts injected at the 1 cell stage
1165   with siControl and siERH siRNAs showing protein depletion. Scale bars, 10µm.

1166   **d.** Representative DIC images of mouse blastocysts injected at the 1 cell stage with siControl
1167   and siERH siRNAs, two examples for each shown.

1168   **e.** Fluorescent imaging of trophectoderm marker CDX2 (green), pluripotency marker OCT4
1169   (magenta), with Phalloidin (orange) to define cell boundary and quantification of OCT4
1170   levels in siControl siERH blastocysts with DAPI (blue). (n=8 per condition, t-test, ns=not
1171   significant).

1172   **f.** Representative DAPI staining of siControl and siERH blastocysts. White arrows indicated
1173   DNA shedding.

1174   **g.** Expression of Erh in siControl and siERH treated mouse blastocysts by RNA-seq. (n=3
1175   per condition, t-test, p>0.0001).

1176   **h.** TPM expression values from RNA-seq for primitive endoderm genes. (n=3 per condition,
1177   t-test)

1178   **i.** TPM expression values from RNA-seq for epiblast genes. (n=3 per condition, t-test,
1179   ns=not significant)

1180

a *Erh* expression during mouse preimplantation

b ERH immunofluorescence in mouse blastocyst comparing protein levels in trophectoderm and ICM cells

c ERH protein staining in mouse blastocysts injected with siRNA at the 1-cell stage

e Comparison of OCT4 protein levels between siControl and siERH mouse blastocysts

d Example brightfield images of mouse blastocysts injected at the 1 cell stage with siControl or siERH siRNAs

f DAPI staining in siControl and siERH blastocysts showing increased DNA shedding

g *Erh* expression in siControl and siERH mouse blastocysts

h Expression of primitive endoderm genes in siControl and siERH mouse blastocysts

i Expression of epiblast genes in siControl and siERH mouse blastocysts

Extended Data Fig. 6. ERH is involved in silencing of pluripotency genes during differentiation

a. Imaging of H1 hESCs in pluripotent and 1 day endoderm differentiated showing changes
in colony morphology (phase), DAPI (blue), silencing of SOX2 (green) and induction of
endoderm marker SOX17 (red). Scale bars, 100µm.

b. Quantification of nuclear NANOG immunofluorescence intensity in siControl vs siERH 1
day endoderm differentiated H1 hESCs. (unpaired t-test).

c. ERH transcripts per million (TPM) values from RNA-seq in siControl and siERH H1 hESCs
in untreated, endoderm (DE) and ectoderm (EC) differentiation conditions.

d. Volcano plot showing differential gene expression in siERH vs siControl H1 in pluripotency
conditions. Inset pie chart indicates the number of genes upregulated and down regulated
in siERH vs siControl H1 (DESeq2, padj<=0.05, log2fc>0.5).

e. Volcano plot showing differential expression of genes upregulated in naïve vs primed
hESCs (List from Messmer et al., 2019[43]) in siERH vs siControl H1 in pluripotency
conditions. Inset pie chart indicates the number of genes upregulated and down regulated
in siERH vs siControl H1 (DESeq2, padj<=0.05, log2fc>0.5).

f. Volcano plot showing differential gene expression in siERH vs siControl 1 day endoderm
for genes down regulated in siControl DE vs siControl H1. Inset pie chart indicates the
number of genes upregulated and down regulated in siERH vs siControl DE (DESeq2,
padj<=0.05, log2fc>0.5).

g. Volcano plot showing differential gene expression in siERH vs siControl 2 day ectoderm
for genes down regulated in siControl EC vs siControl H1. Inset pie chart indicates the
number of genes upregulated and down regulated in siERH vs siControl EC (DESeq2,
padj<=0.05, log2fc>0.5).

h. Volcano plot showing differential expression of genes upregulated in naïve vs primed
hESCs (List from Messmer et al., 2019[43]) in siERH vs siControl H1 hESCs in 1 day
endoderm conditions. Inset pie chart indicates the number of genes upregulated and down
regulated in siERH vs siControl H1 (DESeq2, padj<=0.05, log2fc>0.5).

i. Intersection of genes enriched in hESCs with high self-renewal[93] with genes repressed by
ERH during endoderm and ectoderm differentiation. (hypergeometric test)

j. Graphs of naïve gene RNA-seq, transcripts per million (TPM) values from RNA-seq
replicates shown (DESeq2 adjusted p-value).

k. Graphs of key naïve gene expression in siControl and siERH H1 hESCs compared to
FPKM values for naïve and primed ESCs from Bi et al., 2022.
For all RNA-seq n=3, except for siCTRL UT where n=2.

a Imaging of endoderm differentiation of H1 hESCs

b NANOG abundance in siControl and siERH endoderm

c ERH expression in siRNA treated pluripotent and differentiated H1 hESCs

d H1 siERH vs siControl

e Genes upregulated in naïve vs primed hESCs

f Endoderm siERH vs siControl of genes down in siCTRL DE vs H1

g Ectoderm siERH vs siControl of genes down in siCTRL EC vs H1

h Genes upregulated in naïve vs primed hESCs

i Overlap of hESC self-renewal genes with ERH repressed genes

j Expression of naïve pluripotency genes in siRNA treated pluripotent and differentiated H1 hESCs

k Average expression in primed (pESC) and naive (nESC) cells from Bi et al., 2022 compared to siCTRL and siERH hESCs

1215 Extended Data Fig. 7. ERH represses alternative lineage gene activation during hESC
1216 differentiation

    a. GO analysis (PantherDB, statistical overrepresentation) of genes upregulated (padj<=0.05, log2 fold change >= 0.5) in siERH vs siControl H1 hESCs 2 day ectoderm differentiation.

    b. Example meiotic and gametogenic genes activated during endoderm and ectoderm differentiation by ERH depletion, transcripts per million (TPM) values from RNA-seq replicates shown (DESeq2 adjusted p-value).

    c. Example alternative lineage genes activated during ectoderm differentiation by ERH depletion, transcripts per million (TPM) values from RNA-seq replicates shown (DESeq2 adjusted p-value).

    For all RNA-seq n=3, except for siCTRL UT where n=2.

a

GO analysis for statisticaly enrichment of genes
upregulated in siERH ectoderm

b

Expression of example gametogenic genes

c

Expression of example liver genes in siControl and siERH H1
cells during pluripotency and ectoderm differentiation

Extended Data Fig. 8. ERH represses evolutionarily young transposable elements with regulatory
function

a. H3K9me3 data from H1 hESCs (GSM1003585) and hESC derived endoderm
(GSM916057) at gene bodies +/- 1kb for upregulated (adjusted pvalue<=0.05;
log2foldchange>=0.5) by ERH depletion during 1 day endoderm differentiation.

b. SVA class element expression plots in reads per million (RPM) from RNA-seq data
analyzed by TETranscripts. (DESeq2 adjusted p-value).

c. LTR7Y expression plot from RNA-seq data analyzed by TETranscripts in untreated (UT)
and endoderm (DE) differentiated H1 hESCs treated with siControl or siERH. (DESeq2
adjusted p-value).

d. LTR5_Hs and HERVK expression plots in reads per million (RPM) from RNA-seq data
analyzed by TETranscripts. (DESeq2 adjusted p-value).

e. Browser track showing RNA-seq of example repeat element and nearby gene that is
expressed in pluripotency, silenced in siControl DE and EC but expressed in siERH DE
and EC conditions.

f. Browser track showing RNA-seq of example repeat element and nearby gene that is
expressed in pluripotency, silenced in siControl DE and EC but expressed in siERH DE
condition.

g. Plot of expression change of specific L1 element loci vs expression change of
corresponding nearest gene within 25kb in 1 day endoderm. R, Pearson correlation.
For all RNA-seq n=3, except for siCTRL UT where n=2.

1250

Katznelson et al., Extended Data Figure 8

Extended Data Fig. 9. ERH dependent repression of H3K9me3 marked repeat elements

a. Browser track showing RNA-seq of example repeat element and nearby gene in siControl

and siERH treated H1 hESCs after 1 day of endoderm differentiation and H3K9me3 tracks

from H1 hESCs (GSM1003585) and hESC derived endoderm (GSM916057).

b. Plots of H3K9me3 CUT&RUN signal at L1 elements in endoderm differentiated H1 hESCs

treated with siControl or siERH. Data from loci at all L1 element loci with at least 2-fold

increased RNA expression upon ERH depletion.

c. Browser track showing example repeat element gaining H3K9me3 during endoderm

differentiation in siControl and gaining less H3K9me3 in siERH. RNA-seq track showing

proximal gene to repeat element being expressed in siERH endoderm.

a Browser track of example coordinated expression of gene
   and proximal repeat element with H3K9me3

b H3K9me3 CUT&RUN signal in siControl and siERH endoderm differentiated
   H1 hESCs at siERH upregulated L1 element loci at all or endoerm H3K9me3 gained loci

c Browser tracks of example repeats and proximal genes upregulated by ERH
   knockdown during endoderm differentiation with and without decreased H3K9me3

Extended Data Fig. 10. ERH protein levels are higher in differentiated cells and ERH is recruited
to the chromatin by RNA and RNA binding proteins.

a. ERH expression (TPM) from RNA-seq in siControl treated human fibroblasts, pluripotent
hESCs, 1 day endoderm differentiated hESCs (DE), and 2 day ectoderm differentiated
hESCs (EC). (n=2 for siControl fibroblasts and UT H1 hESCs, n=3 for siControl DE and
EC H1 hESCs).

b. ERH staining in human fibroblasts and pluripotent H1 hESCs, DAPI (blue), ERH (red).
Staining and imaging performed in parallel using same conditions and microscope settings
for both samples.

c. ERH staining in pluripotent H1 hESCs and 1 day endoderm differentiated H1 hESCs, DAPI
(blue), ERH (red). Scale bars, 50µm. Staining and imaging performed in parallel using
same conditions and microscope settings for both samples.

d. Western blot after RNase A treatment during soluble nuclear protein isolation. Western for
ERH and YTHDC1 in nuclear and chromatin fractions with and without RNase. RBMX and
H3 included as known RNA-sensitive and RNA-insensitive, chromatin-associated
proteins.

e. Quantification of immunofluorescence (IF) of nuclear ERH signal after siYTHDC1 followed
by washout, in human fibroblasts.

1279

1280

a

ERH expression by RNA-seq in
human fibroblasts and human hESCs

b

ERH in human fibroblasts and
H1 hESCs stained in parallel,
imaged using same settings

c

ERH in H1 hESCs in pluripotent and
1 day endoderm differentiaiton conditions,
stained in parallel, imaged using same settings

d

RNase A treatment of soluble
nuclear and chromatin fractions

e

Quantification of ERH intensity
by immunofluorescence
after permeabilization and
washout for indicated siRNA