Microbial interactions affect the tempo and mode of antibiotic resistance evolution

Laurens E. Zandbergen[1], Joost van den Heuvel[2], Andrew D. Farr[2,3], Bas J. Zwaan[2], J. Arjan G.

M. de Visser[2], Marjon G. J. de Vos[1,2]*

[1] Groningen Institute of Evolutionary Life Sciences (GELIFES), University of Groningen,

Groningen, The Netherlands

[2] Laboratory of Genetics, Wageningen University, Wageningen, The Netherlands

[3] Max Planck Institute for Evolutionary Biology, Plön, Germany

*corresponding author: m.g.j.de.vos@rug.nl

Competing interests

The authors declare no conflict of interest.

Abstract

The global rise of antibiotic resistance impedes the treatment of bacterial infections. To limit the

emergence and evolution of antibiotic resistance it is important to understand how bacterial

interactions in multispecies communities affect the course of evolution. We investigated how

ecological interactions between microbes derived from polymicrobial urinary tract infections

affect the tempo and mode of antibiotic resistance evolution. We show that for representative

strains of three uropathogens, *Escherichia coli, Klebsiella pneumoniae* and *Enterococcus*

faecium, the rate and evolutionary trajectories towards antibiotic resistance differ depending on

the conditioned medium mediated interactions with other microbes that alter their growth and

antibiotic tolerance. Replicate lineages of the same species evolved under similar ecological

conditions show parallel evolutionary trajectories, and resistance mutations and other functional

targets selected differed between these conditions. Our findings demonstrate that bacterial

interactions differentially affect the evolutionary potential of antibiotic resistance evolution.

Introduction

Bacterial pathogens often live in multispecies communities. In such microbial communities, bacteria interact, which affects their growth, survival and coexistence (1–5). Bacterial infections are often treated with antibiotics, whose efficacy may also depend on ecological interactions with community members (6). For instance, co-existing bacterial species in pathogenic communities can mediate the growth and survival of community members in the presence of antibiotics (7–9) through antibiotic degradation (10,11), change the metabolic landscape affecting pathogen fitness (12–14), lower antimicrobial efficacy or alter the transport of antibiotics (7,15), or change the induction of signal molecules that induce resistance mechanisms (16).

Ecological interactions can therefore affect the evolvability of antibiotic resistance in several ways. Firstly, they may alter the mutation supply, by affecting the population size or mutation rate (17–20). Additionally, ecological interactions may alter the fitness effect of mutations, mediating the selective accessibility of adaptive mutations, as genetic and environmental factors are known to modulate the effect of mutations (21–23) and therefore potentially morph the shape of the adaptive landscape (24,25).

In clinical practice, urinary tract infections (UTIs) are treated with antibiotics. Recurrence of infection after the discontinuation of the antimicrobial therapy in treated patients can be high (up to 50% in the elderly population) (26). This is not surprising, as infections are often polymicrobial and many UTI isolates are resistant to commonly prescribed antibiotics (1,27,28). To investigate whether different microbial interactions have an effect on the evolution of antibiotic resistance, we evolved three focal species (*Escherichia coli, Klebsiella pneumoniae* and *Enterococcus faecium*) in the conditioned medium of several species (*K. pneumoniae, E. coli, E. faecium, Staphylococcus haemolyticus* and *Pseudomonas aeruginosa*). The bacteria were isolated from elderly patients diagnosed with polymicrobial urinary tract infections (1). These pathogens are

part of the so-called ESKAPEE bacteria, which are known for their ability to evade antibiotics

due to their increasing multi-drug resistance (29). We hypothesize that ecological interactions

between uropathogens determine the evolutionary potential of antibiotic resistance evolution

through the modulation of population sizes and changing the shape of the adaptive landscape.

Materials and Methods

Bacterial isolates and growth medium

The five uropathogen isolates used in this study were previously isolated anonymously from

different elderly patients (>70 years old) suffering from urinary tract infection (1). These five

isolates (*E. coli*, *K. pneumonia*, *E. faecium*, *S. haemolyticus*¸ and *P. aeruginosa*) were selected

based on their consistent planktonic growth in Artificial Urine Medium (AUM) and for their

observed growth affecting interactions in a previous study (2,6). Isolates were cultured in

Artificial Urine Medium (AUM) (30).

Preparation of conditioned medium

Spent medium of *E. coli*, *K. pneumonia*, *E. faecium*, *S. haemolyticus*¸ and *P. aeruginosa* isolates

was prepared by culturing in AUM, followed by centrifugation and filtering. To prepare

conditioned medium, the AUM spent medium fraction was replenished with AUM constituents.

Serial dilution experimental evolution protocol

K. pneumoniae and *E. faecium* were evolved for 30 bi-daily transfers, *E. coli* was evolved for 35

bi-daily transfers. Focal species were cultured in 5 ml preheated conditioned medium, in 15 ml

screw cap falcon tubes, with the caps slightly opened. Each transfer *K. pneumoniae* and *E. coli*

were diluted 500 times, and *E. faecium* was diluted 100 times, due to its inherent smaller

population size in the artificial urine medium and urine. *K. pneumoniae* and *E. coli* were evolved

in the conditioned media of *K. pneumoniae*, *E. coli*, *E. faecium*, *Staphylococcus haemolyticus. E.*

faecium was additionally also evolved in *P. aeruginosa* conditioned medium. Each focal isolate-

conditioned medium combination was replicated five times. The Gram-negative isolates *K.*

pneumoniae and *E. coli* were evolved in trimethoprim-sulfamethoxazole (ratio 1:5, as in clinical

use (31)) and *E. faecium* in vancomycin (32). The level of antibiotics at the start of the

experiments was set at a 50% sub-MIC concentration, 0.05 and 0.0125 µg/mL trimethoprim for

K. pneumoniae, and *E. coli* respectively, and 0.25 µg/mL vancomycin for *E. faecium*. If the

cultures exceeded a population size threshold (for *K. pneumoniae* and *E. coli* 0.1 OD600 and

0.05 OD600 for *E. faecium)*, inferred through an optical density assessment, similar to

mcFarland standard assay (33), the antibiotic concentration was increased by 20%, otherwise

the antibiotic concentration was kept at the same level as the previous transfer.

Growth Measurements

Three clones of three of the five evolved replicate lineages of each focal species, evolved in

each conditioned medium, and the wild types of each focal species were tested in phenotypic

growth assays. Growth rates were obtained by incubation in a BMG Clariostar or Fluorostar

microplate reader at 37°C under aerobic conditions and shaken continuously at 500 rpm. The

growth rate was defined by the shortest obtained doubling time and quantified by the steepest

linear fit of the exponential phase of log(OD600). Population density was inferred from OD600

absorbance measurements as well, taking the average for all the measurements of the

stationary phase of the growth curve.

Antibiotic tolerance assays

Growth measurements were taken of wild type isolates of *K. pneumoniae*, *E. faecium* and *E. coli*

were inoculated with 2 µl of overnight cultures in triplicates in conditioned medium and AUM

reference medium in a concentration gradient with two-fold increases in antibiotic concentration

in 96-well plates. Plates were incubated for 24h at 37°C. *K. pneumoniae* and *E. coli* were

assayed in the presence of trimethoprim-sulfamethoxazole and *E. faecium* in the presence of

vancomycin. The fold change tolerance was calculated by dividing the concentration that limited

growth in a particular conditioned medium by the concentration of antibiotic that limited growth in

AUM.

MIC assays

Starting Minimum Inhibitory Concentration (MIC) of the wild-types were measured in a serial

dilution assay of the desired antibiotics in AUM. 200 µL cultures were created in a 96-well plate

including a two-fold serial dilution of the desired antibiotic (vancomycin for *E. faecium* and a 1:5

mixture of trimethoprim and sulfamethoxazole for both *E. coli* and *K. pneumoniae*) ranging from

0.125 to 1 µg/mL for the vancomycin and 0.00625 to 0.2 µg/mL for

trimethoprim:sulfamethoxazole, in triplicate. Cultures without antibiotics were used as control.

Cultures were incubated over 24h at 37°C at 200 rpm shaking conditions. Whether or not

bacteria had visibly grown was determined by measuring OD600 and subtracting AUM without

bacteria as blank. OD600 values <0.05 were inspected visually to determine complete inhibition.

MIC of final transfer mutants were determined on agar using a modified version of the TDtest

described previously by Gefen et al. (34), Supplementary Text 1.

DNA extraction

Cultures of evolved clones were grown overnight in 5ml LB, cells were pelleted, and DNA was

extracted using a Maxwell 16 Instrument in combination with the Maxwell® 16 Tissue LEV Total

RNA Purification Kit Custom (XAS 1220) (Promega, Madison, WI, USA) following manufacturer's

instructions. The DNA was eluted in 30 µL DNAse free water, and quantified using Nanodrop

2000c Spectrophotometer (Thermo Fisher Scientific, USA) and gel electrophoresis.

Sequencing

Three clones of three of the five evolved replicate lineages of each focal species, evolved in

each conditioned medium, were prepared for sequencing. DNA was extracted using the

Maxwell® 16 Tissue LE Total RNA purification kit (article number XAS1220, specifically tailored

towards DNA extraction). Sequencing was performed at Baseclear BV Leiden. Of the ancestor

and the evolved clones, paired-end 125 bp reads were generated using the Illumina HiSeq2500

or MiSeq system. Of the ancestors, long sequence reads were generated using the Oxford

Nanopore GridION X5 system to facility the mapping of Illumina reads of the evolved mutants,

and ancestor.

dN/dS

We calculated the dN/dS ratio of all mutations in the genomes of each sequenced end point

clone, by dividing the number of non-synonymous mutations by the number of synonymous

mutations, and controlling for mutation bias (35).

Calculation *H*-index

To quantify the repeatability of genomic evolution, the *H*-index previously described by Schenk

et al. (17) was used to estimate the pairwise fraction of shared mutations. The *H*-index was

calculated at the function (COG) and gene level. All mutants were compared pairwise, calculated

using (17):

$$H_{A,B} = \sum_{j=1}^{m} \sum_{k=1}^{n} \left(\frac{|A_j \cap B_k|}{|A_j|} \right) / \sum_{j=1}^{m} \sum_{k=1}^{m} \left(\frac{|A_j \cap A_l|}{|A_j|} \right) \qquad (1)$$

Which compares genotype *A* and *B* with *m* and *n* mutations (with index *j* and *k*), respectively.

Here, $A_j \cap A_l$ denotes the mutations that genotypes *A* and *B* have in common. The *H*-index is

calculated for both genotypes separately ($H_{A,B}$ and $H_{B,A}$) to take into account asymmetries in

overlap between mutations of different sequence size and compared using (17):

$H = \frac{H_{A,B} + H_{B,A}}{2}$ (2)

The final *H*-index value would fall between 0 (no repeated evolution) and 1 (complete

parallelism). When the *H*-index is 0.5, it indicates that the average of the two compared mutants

have half of their mutations shared between them.

Mutation rate assay

To estimate the intrinsic mutation rate of *E. faecium* and *K. pneumoniae*, we used a high-

throughput test which is similar to the fluctuation test developed by Luria and Delbrück (36).

Statistical analysis

For each genotype, measurements (of phenotypes such as on growth rate, population size,

resistance or tolerance levels) were performed in either triplicates or quadruplicates. The null

hypotheses were rejected if the *P*-value obtained from the test was lower than a significance

level of 0.05. For statistical tests with multiple pairwise comparisons, *P*-values were adjusted

using the Holm-Bonferroni method (37).

Additional information on all Materials and Methods can be found in Supplementary text 1.

Results

Conditioned media promote the adaptive radiation of antibiotic resistance evolution

To investigate the differential effect of microbial interactions on the evolution of antibiotic

resistance, we evolved single isolates of focal species *K. pneumoniae*, *E. coli* and *E. faecium* by

bi-daily serial transfers in conditioned artificial urine medium (AUM) supplemented with

antibiotics. Conditioned medium was used as a proxy for microbial interactions. It was prepared

by creating spent media of cultures grown in AUM followed by replenishment of the medium with

fresh nutrients (Materials and Methods) (6). When the evolving isolates had reached a

predetermined population size threshold at the moment of transfer, the antibiotic concentration

was increased by 20% (Materials and Methods). Each combination of focal isolate and

conditioned medium was replicated five times. The two Gram-negative species, *K. pneumoniae*

and *E. coli*, were evolved in the presence of trimethoprim-sulfamethoxazole (ratio 1:5, similar to

clinical use (31)), which inhibits DNA synthesis (38). *E. faecium* was evolved in the presence of

the Gram-positive specific antibiotic vancomycin, which inhibits cell wall synthesis (39); initial

antibiotic concentrations were 50% of the minimal inhibitory concentration (MIC) in all cases. *E.*

coli was evolved over 70 days (35 transfers) and *K. pneumoniae* and *E. faecium* for 60 days (30

transfers) (Materials and Methods).

We find that conditioned media differentially affect the rate of evolution and final level of

antibiotics in the environment (Figure 1). Notably, the evolutionary trajectories of the replicates

per conditioned medium are highly parallel. This is especially the case for *K. pneumoniae* and *E.*

faecium (Figure 1B,C). Comparing the rate of evolution defined as the slope of the graphs and

the final antibiotic concentration reached per focal species and conditioned medium, we find

significant differences for all conditioned medium for *E. faecium* (*P*<0.05 for all pairwise

comparisons, Welch's t-test with Holm-Bonferroni correction). For *K. pneumoniae* we find

significant differences (*P*<0.05, evolutionary rate and final antibiotic concentration) for all

conditioned media, except for the overlapping trajectories of *E. coli* and *S. haemolyticus*

conditioned media. The evolutionary trajectories of *E. coli* per conditioned medium environment

also display parallelism (Figure 1A), but are more variable compared to *K. pneumoniae* and *E.*

faecium (Figure 1B,C); we find significant differences for all conditioned media between

conditions (*P*<0.05, evolutionary rate and final antibiotic concentration), except for the partially

overlapping trajectories of *S. haemolyticus* and *E. faecium,* and *S. haemolyticus and E. coli*

conditioned medium (*P*=0.059 and *P*=0.10, respectively). This shows that the ecological

interactions affect the course of antibiotic resistance evolution.

Response to selection correlates with ecological effects on population size

Bacterial interactions, mediated by direct interactions or conditioned medium, are known to

affect the population sizes, for instance by competition for nutrients or cross-feeding. To

investigate whether population size effects due to bacterial interactions are related to the level of

evolution observed, we correlate the maximum population sizes, as inferred by OD600

measurements, of the ancestors in conditioned medium, with the final antibiotic levels tolerated

in the evolution experiments (Figure 2A-C). Indeed, the antibiotic concentration at the final

transfer correlates significantly with the population size in conditioned medium for all three focal

species. We find a strong positive correlation for *E. coli* and *E. faecium*, and a moderate

correlation for *K. pneumoniae* (R^2=0.86, 0.93 and 0.45, for *E. coli, E. faecium* and *K.*

pneumoniae resp., *P*<0.001 for *E. coli* (*N*=20) and *E. faecium* (*N*=25), and *P*=0.017 for *K.*

pneumoniae (*N*=20), Figure 2A-C).

Because phylogenetically similar species are expected to have the most niche overlap, we

hypothesize that the conditioned medium of phylogenetically similar species leads to the most

niche overlap, and consequently, the smallest population sizes, and hence the slowest evolution,

as established above. Indeed, we find that the rate and level of antibiotic resistance evolution is

slowest in the conditioned medium of the focal isolate for two of the three focal species (*K.*

pneumoniae in *K. pneumoniae* conditioned medium, and *E. faecium* in *E. faecium* conditioned

medium, Figure 1B,C). Moreover, for two of the three focal isolates the rate of evolution is

fastest in the conditioned medium of the isolates that are phylogenetically least related

(Supplementary Figure 1), *E. faecium* evolves fastest in the Gram-negative conditioned media

(*H*=16.39, *P*<0.005, Kruskal-Wallis), and *E. coli* evolves fastest in the Gram-positive conditioned

media (*H*=12.09, *P*<0.005). For *K. pneumoniae* we observe no such niche effects (*H*=0.006,

P=0.94), other than the slowest evolution in its own conditioned medium. Alternatively, the

conditioned medium of donor species may affect population sizes by other means than niche

overlap. For instance, we find that *K. pneumoniae* conditioned medium was associated with the

highest pH (pH 8), compared to the other conditioned media (pH 7.5 *S. haemolyticus and P.*

aeruginosa; pH 7.25 *E. faecium* and *E. coli* conditioned medium)*.* This increased alkaline

condition, was associated with the slower evolution of *E. coli* in *K. pneumoniae* conditioned

medium (40), but not the slower evolution of *E. faecium*, which is known to tolerate a wider

range of pH conditions (Figure 1A,C) (41,42). This suggests that niche specific effects such as

avoidance of competition, but also more general niche effectors, such as pH, could affect

bacterial interactions that promote the divergence of antibiotic resistance evolution, as also

observed in other studies (43–46).

We previously found that growth-mediating effects of bacterial interactions in the absence of

antibiotics correlate with the level of antibiotic tolerance (6). Here we define the tolerance as the

level of an antibiotic that a population can sustain under the influence of bacterial interactions

(6). The fold change in antibiotic tolerance was calculated based on the change in growth

inhibiting antibiotic concentration of conditioned media compared to unconditioned medium.

Here, we find that the concentration of antibiotics at the last transfer of the experiment correlates

positively with the antibiotic tolerance levels conferred by the conditioned media (R^2=0.50, 0.68

and 0.64 for *E. coli* (*N*=20), *K. pneumoniae* (*N*=20), and *E. faecium* (*N*=25) resp. for all *P*<0.001,

Figure 2D-F). Possibly, these effects of conditioned media on antibiotic tolerance are driven by

affecting the population size, similar to the inoculum effect in assays measuring the minimal

inhibitory concentration of certain antibiotics (11,47–49).

Ecological interactions do not affect the number of genetic changes

To investigate the consequences of these ecological interactions on evolutionary trajectories, we

determined the genetic changes in these evolved lineages through the sequencing of three

clones per evolved lineage (Materials and Methods). A fundamental determinant of evolutionary

trajectories is the population size, as it affects both the supply of mutations and the efficacy of

natural selection to select beneficial mutations (17,20,50). Therefore, effects on population size

may lead to different mutations being selected (51,52).

To quantify the impact of natural selection in our populations, we measured the dN/dS ratio

(35,53) (Materials and Methods). Generally, we find dN/dS ratios >> 1, resp. *E. coli* 4.5, *K.*

pneumoniae 7.2, *E. faecium* 17, indicating strong positive selection in all populations (chi-

squared test to test deviation from expected dN/dS in the absence of selection, *P*=0.024, 0.004,

0.001, for respectively *E. coli*, *K. pneumoniae*, *E. faecium*, Supplementary Table 1, Materials

and Methods).

SNPs outnumbered other mutations in the evolved isolates (Supplementary Figure 2A-C). Yet,

we find no positive correlation between the number of SNPs in the end-point mutants and

population size for *E. coli* and *E. faecium* (R^2=0.18 and 0.47, *P*=0.17, 0.20, resp.,

Supplementary Figure 3A,C), and a negative correlation for *K. pneumoniae* (R^2=0.70, *P*<0.001,

Supplementary Figure 3B). Other genetic differences were also observed, such as deletions,

inversions and copy-number differences. Moreover, in several of these *K. pneumoniae* lineages

one of the plasmids was entirely lost. Especially for the second plasmid (Supplementary Figure

4B, Supplementary Table 2), we observed that blocks of the plasmid were deleted during

evolution. These mutational patterns were not observed to be associated with specific

conditioned media.

We find no positive correlation between the different conditioned media associated with different

population sizes, and the total number of genetic changes (R^2=0.023 and 0.03, 0.15, P=0.68,

0.59, 0.19 for *E. coli, K. pneumoniae* and *E. faecium* lineages respectively, Supplementary

Figure 5A-C). Therefore, population size effects did not affect the number of genetic changes in

the end-point mutants.

Ecological interactions do affect the types of genetic changes

To test whether different conditioned media lead to the incorporation of different genetic changes

in the adaptive trajectories of the three focal species, we compared the genetic similarity of

isolates evolved in the same with those evolved in different conditioned media. We calculated

the *H*-index (17), as a measure for the fraction of mutated genes that two evolved isolates have

in common (Materials and Methods). If we calculate this index at the level of Clusters of

Orthologous Groups (COGs) (54), i.e. genes grouped by function, we find more parallelism

among replicate populations evolved within the same conditioned media compared to those

same lineages in different conditioned media (P=0.043, 0.043, 0.032 (Welch's t-test), excluding

self-comparisons, for *E. coli*, *K. pneumoniae* and *E. faecium,* respectively, Figure 3). This

indicates that the parallel evolution observed (Figure 1), compliments the parallelism observed at

the functional level in the same conditioned media.

For *E. coli*, particularly the lineages that evolved in the Gram-positive conditioned media of *S.*

haemolyticus and *E. faecium* display a high degree of parallelism at the COG-level (H>0.6,

Figure 3A, P<0.05 compared to the other *H*-values, Welch's t-test). This pattern is conserved at

the gene-level for lineages evolved in *S. haemolyticus* conditioned medium (Supplementary

Figure 6A). The high genetic parallelism corresponds to a similar phenotypic growth response of

E. coli replicates to these conditioned media (Figure 2D), and their partially overlapping

evolutionary trajectories (Figure 1A). Interestingly, the population size of *E. coli* in the

conditioned medium of itself is also similar to the population size in the conditioned medium of *S.*

haemolyticus (Figure 2A, *P*>0.05 when comparing the two set, Welch's t-test). Yet, the *H*-index

values that correspond to the *E. coli* lineages evolved in these Gram-positive (*S. haemolyticus)*

and Gram-negative (*E. coli)* conditioned media are relatively low (*H*<0.6), indicating lower

parallelism within these conditions. This shows that population size alone cannot explain the

level of parallelism, which suggests that the constituents of the conditioned medium, also play a

role in the genetic divergence of these evolutionary trajectories.

For *E. faecium*, we find a particularly strong parallelism at the COG-level for evolved lineages in

P. aeruginosa conditioned medium, but less so for other conditioned media (Figure 3C). At the

gene-level, this strong parallelism for evolved lineages in *P. aeruginosa* conditioned medium is

not present (Supplementary Figure 6C). The high level of parallelism among *E. faecium* lineages

evolved in *P. aeruginosa* conditioned medium coincides with the largest population sizes, and

the highest rate of evolution (Figure 1C). Yet, the level of parallelism of *E. faecium* evolved in the

conditioned media of *K. pneumoniae* and *S. haemolyticus* is lower, despite similar resistance

trajectories and population sizes (*H*-index means of 0.75 for *P. aeruginosa,* and 0.35 (*P*=0.022)

and 0.48 (*P*=0.013) for *K. pneumoniae* and *S. haemolyticus* resp., Welch's t-tests *H*-index

comparison of *E. faecium* in the respective conditioned media) (Figure 1B, 3B). This again

suggests that particular metabolic constituents of the conditioned medium also play a role in the

genetic divergence.

For *K. pneumoniae*, we find consistently high *H*-indices (*H*>0.6) at the COG-level across all

conditions, but a little lower index for the comparison of the isolates evolved in *E. faecium* and *E.*

coli conditioned media (Figure 3B). In addition, at the gene-level we also find relatively high

parallelism across all conditions (although slightly lower than at the COG-level), especially for

the isolates evolved within *S. haemolyticus* and *K. pneumoniae* conditioned medium (Figure 3B).

These findings highlight the rather limited effect of conditioned media on the antibiotic resistance

evolution of *K. pneumoniae* (Figure 2B).

As observed before (17,55), the *H*-indices show a higher parallelism at the COG-level than at

the gene-level (Figure 3, Supplementary Figure 6). This suggests that there is genetic

redundancy in the functionalities that are selected under these particular ecological

circumstances. Generally, the higher *H*-index of lineages evolved within the same conditioned

medium (along the diagonal axis in Figure 3) shows that the antibiotic resistance evolution in the

conditioned media leads to parallel genotypic (Figure 3) and phenotypic (Figure 1) adaptation

under the influence of species-specific bacterial metabolites.

Conditioned medium causes specific genetic changes

Zooming in on the genetic changes, we investigated whether the species-specific ecological

interactions also lead to the selection of different resistance mechanisms. We do find that in

particular conditioned media specific mutations occur repeatedly, whereas others do not (Figure

4A-C). In accordance with other research, we hypothesized that in larger populations, larger

effect-size mutations are selected, because these are not limited in their mutation supply and are

under strong selection, whereas in smaller populations, different types of smaller effect size

mutations would be selected (17,19). Indeed, we find that the lineages with the smallest

population size, which is *E. faecium* cultured in its own conditioned medium, repeatedly lacked

mutations in known antibiotic resistance targets (Figure 4C) (56–58). When evolved in other

conditioned media, with larger population sizes, they reached higher levels of resistance and

incorporated mutations in well-known vancomycin resistance targets, *VanR, VanS* (56,59,60), or

both. *VanS* is a sensor histidine kinase that detects vancomycin and activates *VanR* (56,59,61).

VanR is a transcriptional activator, regulating the transcription of the vancomycin-resistance

operon together with regulating transcription of the vanHAX operon, resulting in higher tolerance

due to changes in peptidoglycan formation (56,60). Indeed, lineages that had mutations in well-

known resistance targets were generally associated with larger population sizes and higher

resistance levels (Figure 4C, 2C).

The two Gram-negative lineages (*K. pneumoniae* and *E. coli*) were selected for resistance to

trimethoprim-sulfamethoxazole. All sequenced isolates had mutations in *folA*, dihydrofolate

reductase (Figure 4A,B, Supplementary table 3-4), which is involved in the folate pathway that is

essential for DNA synthesis and is inhibited by trimethoprim-sulfamethoxazole (62). *FolA* is the

direct target of trimethoprim. One lineage of *E. coli* evolved in *E. coli* conditioned medium even

had a 20-fold amplification of the copy-number of this gene.

Both *K. pneumoniae* and *E. coli* incorporated *folM* mutations. *FolM* encodes an isozyme of

dihydrofolate reductase and is also involved in DNA synthesis. Inhibition of this gene causes the

upregulation of *folA,* but usually also requires resistance mutations in *folA* to be effective (63). *E.*

coli incorporated *folM* mutations only in lineages that evolved slowly in the *K. pneumoniae*

conditioned medium (Figure 4A). Yet, in *K. pneumoniae*, *folM* mutations appeared in the slowest

evolving lineages in *K. pneumoniae* medium*,* but also in the fastest, namely the *E. coli* and *S.*

haemolyticus conditioned media.

In *K. pneumoniae* in *E. faecium* conditioned medium, *folX* was targeted instead of *folM* (Figure

4B). *E. coli* lineages evolved in *E. coli* and *S. haemolyticus* conditioned medium incorporated

mutations in yet another gene in the folate pathway, *folP* (64) (Figure 4A). All these mutations in

the folate pathway correspond to clinically relevant mutations (Supplementary text 2). The

observation that different genes within the folate pathway are targeted in the different

conditioned media suggests that ecological interactions may cause different trajectories of

antibiotic resistance evolution.

Antibiotic resistance and cost of resistance depend on the genetic background and the

ecological interactions

To assess whether the lineages evolved to the conditioned media or to the antibiotics, we

assessed the growth of the end-point mutants of evolved lineages in each conditioned medium,

and measured the antibiotic resistance level in the absence of conditioned media. The

relationship between the evolved the minimal inhibitory concentration (MIC) and the antibiotic

concentration at the final transfer shows a negative correlation between the MIC and the end-

point antibiotic concentrations for *K. pneumoniae*, and a positive correlation for *E. coli* and *E.*

faecium (Supplementary Figure 7). For these latter focal isolates the highest resistance evolved

in the most and fastest evolving populations (Supplementary Figure 7A,C), whereas for *K.*

pneumoniae resistance evolved fast in all conditioned media (Figure 1), but the highest

resistance evolved during the slowest adaptation (Supplementary figure 7). We also find a cost

of resistance of most mutants, as the growth is mostly below that of the ancestor

(Supplementary figure 8). We find no correlation between the MIC in the absence of conditioned

medium, and the growth of the mutants.

Adaptation to the conditioned medium is shown by the fact that the growth of mutants evolved in

a particular conditioned medium mostly (9 out of 13) lie in the upper right quadrant

(Supplementary Figure 9-11, for those 9 *P<0.05*, one-way ANOVA), showing improved growth in

those conditioned media, compared to the growth in other conditioned media. The fact that

resistance evolution is subject to ecological interactions, is further exemplified by the growth of

end-point clones of *E. coli* evolved in *E. faecium* conditioned medium (Supplementary Figure

9C,E), in the presence and absence of antibiotics. This shows that costs and benefits of accrued

mutations in these lineages are dependent on both environmental factors, the presence of

antibiotics and the conditioned medium. Evolution of antibiotic resistance in the presence of

microbial interactions is therefore a truly eco-evolutionary process.

Discussion and conclusion

We find that the bacterial interactions mediated by conditioned medium differentially altered the rate and mutational trajectories of antibiotic resistance evolution of uropathogenic isolates *E. coli*, *K. pneumoniae* and *E. faecium*. These results are in line with other work that showed that community composition can alter the selection for antibiotic resistance, for instance through modulating population densities via competitive or cooperative interactions (49,65–70). In the conditioned media environment, cultured bacteria were not in physical contact with the bacteria used to create conditioned media. Such contact-mediated interactions may affect eco-evolutionary outcomes, since two populations of bacteria can respond to the presence and the produced metabolites of the other when in the same environment (71,72). Moreover, higher-order interactions between more than two bacterial species could potentially affect growth (73,74). Lastly, the donor species were not cultured in the presence of antibiotics, these could therefore not 'soak' (75) or degrade the level of antibiotics (76). Future studies should address such additional factors that can affect the evolution of antibiotic resistance.

Overall, we found parallelism in lineages evolved under the same ecological conditions (Figure 1), and that resistance mutations and other functional targets selected were dependent on these conditions (Figure 3, 4). Initial standing variation cannot account for this pattern, since different mutations in the same genes were incorporated (Supplementary tables 3-5). Parallelism has been observed to depend on environmental conditions before, such as resource availability (77), ecological interactions (78) and population size of evolving lineages (17), but the phenotypic parallelism does not always correspond to molecular parallelism (79). This work shows that ecological interactions, which may be through avoidance of resource competition (utilizing different niches), cross-feeding, or pH changes, and potentially other antibiotic tolerance effects (6,45,66), can drive both phenotypic and genetic parallel antibiotic resistance evolution.

K. pneumoniae showed the least variation across conditioned media, in terms of growth,

tolerance as well as the rate and final level of adaptation. It also showed the highest parallelism

within each conditioned medium at the genetic level, which correlated with the overall

consistently large population sizes and therefore a smaller effect on mutation supply (19,20). In

contrast, *E. coli* and *E. faecium* showed more variation across conditioned media. Particularly, *E.*

faecium lineages that evolved in the smallest population sizes (in *E. faecium* conditioned

medium), did not incorporate such large effect resistance mutations. These lineages also

showed the highest conditioned medium dependence at the genetic level, which was not always

associated with the population size effects. This suggests that for these species different

conditioned media may have a differential effect on the functions under selection, such that the

underlying shape of the adaptive landscapes of resistance evolution would be more

environmentally dependent (80,81), or that the mechanisms that confer evolutionary benefits are

different depending on the antibiotic concentration (82). Following the mutational trajectories

during the emergence of antibiotic resistance evolution could lead to more insight in the

ecological impact on resistance by identifying genotypes that alter these trajectories (83–87).

Concluding, antimicrobial resistance is turning into a "silent pandemic" (58), as infections caused

by resistant microorganisms fail to respond to conventional treatment, resulting in prolonged

illness, greater risk of death and higher healthcare costs (88). Ecological interactions, mediating

population sizes and other ecological factors, can explain the deviating antibiotic resistance

levels as measured in isolation (e.g., in the clinic) (6,47,69). We find that such interactions

between bacteria, and conferred antibiotic tolerance, affect the mode, the genetic functions

selected, and the tempo, the rate of antibiotic resistance evolution. The identification of the

nature of such ecological interactions can aid in forecasting the evolutionary potential of

antibiotic resistance evolution. Further knowledge of such predictive parameters can improve

decision making, which will be crucial for developing interventions, and for curbing the

emergence and further evolution of antibiotic resistance (89).

Acknowledgements

We thank Ineke Heikamp-de Jong for help with the DNA extraction, and Bart Thomma for the

use of the MLII laboratory at Wageningen University. MGJDV was supported by an NWO VENI

Fellowship (863.14.015), ZonMw ETH grant (40-43500-98-4049/435004009) and L'Oreal

UNESCO LNVH NIAS-KNAW For Women in Science Fellowship, part of this work was

performed at NIAS-KNAW. LEZ was supported by the Adaptive Life program of the Faculty of

Science and Engineering, within GELIFES.

Contributions

Conceived and designed the study: LEZ JAGMDV BJZ MGJDV. Performed the experiments:

LEZ ADF MGJDV. Analyzed the data: LEZ JVDH MGJDV. Wrote the manuscript: LEZ JVDH

ADF JAGMDV MGJDV.

Data availability statement

Bacterial sequence data have been deposited with links to BioProject accession number

PRJNA1103400 in the NCBI BioProject database. The other datasets generated during and/or

analyzed during the current study are available from the corresponding author on reasonable

request.

References

1.   Croxall G, Hale J, Weston V, Manning G, Cheetham P, Achtman M, et al. Molecular
epidemiology of extraintestinal pathogenic *Escherichia coli* isolates from a regional cohort
of elderly patients highlights the prevalence of ST131 strains with increased antimicrobial
resistance in both community and hospital care settings. J Antimicrob Chemother.
66(11):2501–8.
2.   Zandbergen LE, Halverson T, Brons JK, Wolfe AJ, de Vos MGJ. The good and the bad:
Ecological interaction measurements between the urinary microbiota and uropathogens.
Front Microbiol. 2021;12(NA):659450.
3.   Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks, competition,
and stability. Science (1979). 2015;350(6261):663–6.
4.   Chang CY, Bajić D, Vila JCC, Estrela S, Sanchez A. Emergent coexistence in
multispecies microbial communities. Science (1979). 2023;381(6655):343–8.
5.   Fredrickson AG, Stephanopoulos G. Microbial competition. Science (1979).
1981;213(4511):972–9.
6.   De Vos MGJ, Zagorski M, McNally A, Bollenbach T. Interaction networks, ecological
stability, and collective antibiotic tolerance in polymicrobial infections. Proc Natl Acad Sci
U S A . 2017;114(40):10666–71.
7.   Lara EG, van der Windt I, Molenaar D, de Vos MGJ, Melkonian C. Using functional
annotations to study pairwise interactions in urinary tract infection communities. Genes
(Basel). 2021;12(8):1221.
8.   Ramsey MM, Rumbaugh KP, Whiteley M. Metabolite cross-feeding enhances virulence in
a model polymicrobial infection. PLoS Pathog. 2011;7(3):e1002012.
9.   Wintermute EH, Silver PA. Emergent cooperation in microbial metabolism. Mol Syst Biol.
2010;6:407.
10. Yurtsev EA, Chao HX, Datta MS, Artemova T, Gore J. Bacterial cheating drives the
population dynamics of cooperative antibiotic resistance plasmids. Mol Syst Biol.
2013;9(683):683.
11. Geyrhofer L, Ruelens P, Farr AD, Pesce D, de Visser JAGM, Brenner N. Minimal
surviving inoculum in collective antibiotic resistance. Regoes RR, Levin BR, editors. mBio.
2023 Apr 25;14(2).
12. Melkonian C, Seidl MF, van der Hooft JJJ, de Vos MGJ. Metabolic interactions shape a
community's phenotype. Trends Microbiol. 2022;30(7):609–11.
13. Yu JSL, Correia-Melo C, Zorrilla F, Herrera-Dominguez L, Wu MY, Hartl J, et al. Microbial
communities form rich extracellular metabolomes that foster metabolic interactions and
promote drug tolerance. Nat Microbiol. 2022;7(4):542–55.
14. Colegrave N, Buckling A. Microbial experiments on adaptive landscapes. Bioessays.
2005;27(11):1167–73.
15. Delcour AH. Outer membrane permeability and antibiotic resistance. Biochim Biophys
Acta Proteins Proteom. 2009;1794(5):808–16.
16. Li Y, Feng T, Wang Y. The role of bacterial signaling networks in antibiotics response and
resistance regulation. Mar Life Sci Technol. 2022;4(2):163–78.

17.   Schenk MF, Zwart MP, Hwang S, Ruelens P, Severing E, Krug J, et al. Population size
505   mediates the contribution of high-rate and large-benefit mutations to parallel evolution.
506   Vol. 6, Nature Ecology and Evolution. Dryad; 2022. p. 439–47.

507   18.   Mahrt N, Tietze A, Künzel S, Franzenburg S, Barbosa C, Jansen G, et al. Bottleneck size
508   and selection level reproducibly impact evolution of antibiotic resistance. Nat Ecol Evol.
509   2021 Jul 26;5(9):1233–42.

510   19.   Garoff L, Pietsch F, Huseby DL, Lilja T, Brandis G, Hughes D. Population bottlenecks
511   strongly influence the evolutionary trajectory to fluoroquinolone resistance in *Escherichia*
512   *coli*. Barlow M, editor. Mol Biol Evol. 2020 Jun 1;37(6):1637–46.

513   20.   De Visser JAGM, Zeyl CW, Gerrish PJ, Blanchard JL, Lenski RE. Diminishing returns
514   from mutation supply rate in asexual populations. Science (1979). 1999;283(5400):404–6.

515   21.   de Vos MGJ, Poelwijk FJ, Battich N, Ndika JD, Tans SJ. Environmental dependence of
516   genetic constraint. PLoS Genet. 2013;9(6):e1003580.

517   22.   Flynn KM, Cooper TF, Moore FB, Cooper VS. The environment affects epistatic
518   interactions to alter the topology of an empirical fitness landscape. PLoS Genet.
519   2013;9(4):e1003426.

520   23.   Lindsey HA, Gallie J, Taylor S, Kerr B. Evolutionary rescue from extinction is contingent
521   on a lower rate of environmental change. Nature. 2013;494(7438):463–7.

522   24.   De Vos MGJ, Schoustra SE, De Visser JAGM. Ecology dictates evolution? About the
523   importance of genetic and ecological constraints in adaptation. Epl. 2018;122(5).

524   25.   Wright S. The roles of mutation, inbreeding, crossbreeding and selection in evolution.
525   Proceedings of the Sixth International Congress of Genetics. 1932;1:356–366.

526   26.   Nicolle LE. Urinary tract infections in long-term–care facilities. Infect Control Hosp
527   Epidemiol. 2001;22(3):167–75.

528   27.   Siegman-Igra Y, Kulka T, Schwartz D, Konforti N. Polymicrobial and monomicrobial
529   bacteraemic urinary tract infection. Journal of Hospital Infection. 1994;28(1):49–56.

530   28.   Kline KA, Lewis AL. Gram-positive uropathogens, polymicrobial urinary tract infection, and
531   the emerging microbiota of the urinary tract. Urinary Tract Infections: Molecular
532   Pathogenesis and Clinical Management. 2016;459–502.

533   29.   Y. Mahmood H, Jamshidi S, Mark Sutton J, M. Rahman K. Current advances in
534   developing inhibitors of bacterial multidrug efflux pumps. Curr Med Chem. 2016 Apr
535   7;23(10):1062–81.

536   30.   Brooks T, Keevil CW. A simple artificial urine for the growth of urinary pathogens. Lett
537   Appl Microbiol. 1997;24(3):203–6.

538   31.   Hitchings GH. Mechanism of action of trimethoprim-sulfamethoxazole. I. J Infect Dis.
539   1973;128(Supplement 3):S433.

540   32.   Levine DP. Vancomycin: A history. Clinical Infectious Diseases. 2006 Jan 1;42(SUPPL.
541   1):S5–12.

542   33.   Leber AL. Clinical microbiology procedures handbook. Washington DC, USA: ASM Press;
543   2020.

544   34.   Gefen O, Chekol B, Strahilevitz J, Balaban NQ. TDtest: Easy detection of bacterial
545   tolerance and persistence in clinical isolates by a modified disk-diffusion assay. Sci Rep.
546   2017;7(December 2016):1–9.

35. Hershberg R, Petrov DA. Evidence that mutation is universally biased towards AT in
bacteria. PLoS Genet. 2010 Sep 9;6(9):e1001115.
36. Luria SE, Delbrück M. Mutations of bacteria from virus sensitivity to virus resistance.
Genetics. 1943 Nov 20;28(6):491–511.
37. Holm S. A simple sequentially rejective multiple test procedure. Scandinavian Journal of
Statistics. 1979;6(2):65–70.
38. Masters PA, O'Bryan TA, Zurlo J, Miller DQ, Joshi N. Trimethoprim-sulfamethoxazole
revisited. Arch Intern Med. 2003;163(4):402–10.
39. Bourgeois-Nicolaos N, Moubareck C, Mangeney N, Butel MJ, Doucet-Populaire F.
Comparative study of *vanA* gene transfer from *Enterococcus faecium* to *Enterococcus*
faecalis and to *Enterococcus faecium* in the intestine of mice. FEMS Microbiol Lett.
2006;254(1):27–33.
40. Hamdallah I, Torok N, Bischof KM, Majdalani N, Chadalavada S, Mdluli N, et al.
Experimental evolution of *Escherichia coli* K-12 at High pH and with *RpoS* Induction. Appl
Environ Microbiol. 2018 Aug;84(15).
41. Khanum N, Ansari T, Khan HMS, Khan Y. Influence of pH and temperature on stability of
sulfamethoxazole alone and in combination with trimethoprim (co trimoxazole). Asian
Journal of Chemistry. 2012;24(4):1851.
42. Giraffa G. *Enterococcus*. Encyclopedia of Food Microbiology (Second Edition). Oxford:
Academic Press; 2014. p. 674–9.
43. Zampieri M, Enke T, Chubukov V, Ricci V, Piddock L, Sauer U. Metabolic constraints on
the evolution of antibiotic resistance. Mol Syst Biol. 2017;13(3):917.
44. Laborda P, Martínez JL, Hernando-Amado S. Evolution of habitat-dependent antibiotic
resistance in *Pseudomonas aeruginosa*. Microbiol Spectr. 2022 Aug 31;10(4).
45. Windels EM, Cool L, Persy E, Swinnen J, Matthay P, Van B, et al. Antibiotic dose and
nutrient availability differentially drive the evolution of antibiotic resistance and
persistence. ISME J. 2024 Jan 8;18(1):1–30.
46. Ratzke C, Gore J. Modifying and reacting to the environmental pH can drive bacterial
interactions. PLoS Biol. 2018 Mar 14;16(3):e2004248.
47. Brook I. Inoculum effect. Rev Infect Dis. 1989;11(3):361–8.
48. Alvarez S, DeMaria A, Kulkarni R, Klein JO, McCabe WR. Interactions of rifampin and
trimethoprim in vitro. Rev Infect Dis. 1982 Mar 1;4(2):390–401.
49. Karslake J, Maltas J, Brumm P, Wood KB. Population density modulates drug inhibition
and gives rise to potential bistability of treatment outcomes for bacterial infections. PLoS
Comput Biol. 2016;12(10):e1005098.
50. Jiang X, Mu B, Huang Z, Zhang M, Wang X, Tao S. Impacts of mutation effects and
population size on mutation rate in asexual populations: A simulation study. BMC Evol
Biol. 2010 Dec 30;10(1):298.
51. Kauffman S, Levin S. Towards a general theory of adaptive walks on rugged landscapes.
J Theor Biol. 1987;128(1):11–45.
52. Izutsu M, Lake DM, Matson ZWD, Dodson JP, Lenski RE. Effects of periodic bottlenecks
on the dynamics of adaptive evolution in microbial populations. Microbiology (N Y). 2024
Sep 18;170(9).

53. Kryazhimskiy S, Plotkin JB. The population genetics of dN/dS. PLoS Genet. 2008 Dec
12;4(12):e1000304.
54. Tatusov RL, Koonin E V, Lipman DJ. A genomic perspective on protein families. Science.
1997 Oct 24;278(5338):631–7.
55. Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, et al.
The molecular diversity of adaptive convergence. Science (1979). 2012 Jan
27;335(6067):457–61.
56. Arthur M, Molinas C, Courvalin P. The VanS-VanR two-component regulatory system
controls synthesis of depsipeptide peptidoglycan precursors in *Enterococcus faecium*
BM4147. J Bacteriol. 1992;174(8):2582–91.
57. Hancock L, Perego M. Two-component signal transduction in *Enterococcus faecalis*. J
Bacteriol. 2002;184(21):5819–25.
58. Cars O, Chandy SJ, Mpundu M, Peralta AQ, Zorzet A, So AD. Resetting the agenda for
antibiotic resistance through a health systems perspective. Lancet Glob Health.
2021;9(7):e1022–7.
59. Evers S, Courvalin P. Regulation of *vanB*-type vancomycin resistance gene expression by
the VanSB-VanRB two-component regulatory system in *Enterococcus faecalis* V583. J
Bacteriol. 1996;178(5):1302–9.
60. Depardieu F, Courvalin P, Kolb A. Binding sites of VanRB and σ70 RNA polymerase in
the *vanB* vancomycin resistance operon of *Enterococcus faecium* BM4524. Mol Microbiol.
2005;57(2):550–64.
61. Guffey AA, Loll PJ. Regulation of resistance in vancomycin-resistant enterococci: The
vanrs two-component system. Microorganisms. 2021 Sep 25;9(10):2026.
62. Hitchings GH, Burchall JJ. Inhibition of folate biosynthesis and function as a basis for
chemotherapy. In: Advances in enzymology and related areas of molecular biology.
Advances in enzymology and related areas of molecular biology; vol. 27. 1965. p. 417–
68.
63. Giladi M, Altman-Price N, Levin I, Levy L, Mevarech M. *FolM*, A new chromosomally
encoded dihydrofolate reductase in *Escherichia coli*. J Bacteriol. 2003;185(23):7015–8.
64. Buwembo W, Aery S, Rwenyonyi CM, Swedberg G, Kironde F. Point mutations in the *folP*
gene partly explain sulfonamide resistance of *Streptococcus mutans*. Int J Microbiol.
2013;2013:367021.
65. Rezzoagli C, Granato ET, Kümmerli R. Harnessing bacterial interactions to manage
infections: A review on the opportunistic pathogen *Pseudomonas aeruginosa* as a case
example. J Med Microbiol. 2020;69(2):147–61.
66. Meredith HR, Srimani JK, Lee AJ, Lopatkin AJ, You L. Collective antibiotic tolerance:
Mechanisms, dynamics and intervention. Nat Chem Biol. 2015;11(3):182–8.
67. Vega NM, Gore J. Collective antibiotic resistance: Mechanisms and implications. Curr
Opin Microbiol. 2014;21:28–34.
68. Adamowicz EM, Muza M, Chacón JM, Harcombe WR. Cross-feeding modulates the rate
and mechanism of antibiotic resistance evolution in a model microbial community of
Escherichia coli and *Salmonella enterica*. Boshoff HI, editor. PLoS Pathog. 2020 Jul
20;16(7):e1008700.

69. Bottery MJ, Pitchford JW, Friman VP. Ecology and evolution of antimicrobial resistance in
bacterial communities. ISME Journal. 2021;15(4):939–48.

70. Klümper U, Recker M, Zhang L, Yin X, Zhang T, Buckling A, et al. Selection for
antimicrobial resistance is reduced when embedded in a natural microbial community.
ISME Journal. 2019;13(12):2927–37.

71. Brunet YR, Espinosa L, Harchouni S, Mignot T, Cascales E. Imaging type VI secretion-
mediated bacterial killing. Cell Rep. 2013;3(1):36–41.

72. Weiland-Bräuer N. Friends or foes—microbial interactions in nature. Biology (Basel). 2021
Jun 2;10(6):496.

73. Kelsic ED, Zhao J, Vetsigian K, Kishony R. Counteraction of antibiotic production and
degradation stabilizes microbial communities. Nature. 2015;521(7553):516–9.

74. Weinreich DM, Lan Y, Jaffe J, Heckendorn RB. The influence of higher-order epistasis on
biological fitness landscape topography. J Stat Phys. 2018;172(1):208–25.

75. Inglis RF, Hall AR, Buckling A. The role of "soaking" in spiteful toxin production in
Pseudomonas aeruginosa. Biol Lett. 2013;9(1):20120569.

76. Pathak A, Angst DC, León-Sampedro R, Hall AR. Antibiotic-degrading resistance
changes bacterial community structure via species-specific responses. ISME J. 2023 Sep
1;17(9):1495–503.

77. Turner CB, Marshall CW, Cooper VS. Parallel genetic adaptation across environments
differing in mode of growth or resource availability. Evol Lett. 2018 Aug 1;2(4):355–67.

78. Venkataram S, Kuo HY, Hom EFY, Kryazhimskiy S. Mutualism-enhancing mutations
dominate early adaptation in a two-species microbial community. Nat Ecol Evol. 2023 Jan
2;7(1):143–54.

79. Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. Tests of parallel molecular
evolution in a long-term experiment with *Escherichia coli*. Proc Natl Acad Sci U S A.
2006;103(24):9107–12.

80. AlRabiah H, Allwood JW, Correa E, Xu Y, Goodacre R. pH plays a role in the mode of
action of trimethoprim on *Escherichia coli*. PLoS One. 2018 Jul 13;13(7):e0200272.

81. Burian A, Erdogan Z, Jandrisits C, Zeitlinger M. Impact of pH on activity of trimethoprim,
fosfomycin, amikacin, colistin and ertapenem in human urine. Pharmacology. 2012;90(5–
6):281–7.

82. Ruelens P, de Visser JAGM. Choice of β-Lactam resistance pathway depends critically on
initial antibiotic concentration. Antimicrob Agents Chemother. 2021 Jul 16;65(8).

83. Mira PM, Meza JC, Nandipati A, Barlow M. Adaptive landscapes of resistance genes
change as antibiotic concentrations change. Mol Biol Evol. 2015 Oct;32(10):2707–15.

84. Das SG, Direito SO, Waclaw B, Allen RJ, Krug J. Predictable properties of fitness
landscapes induced by adaptational tradeoffs. Elife. 2020 May 19;9.

85. Hall BG. Predicting the evolution of antibiotic resistance genes. Nat Rev Microbiol. 2004
May;2(5):430–5.

86. Palmer AC, Toprak E, Baym M, Kim S, Veres A, Bershtein S, et al. Delayed commitment
to evolutionary fate in antibiotic resistance fitness landscapes. Nat Commun. 2015 Jun
10;6(1):7385.

87.   Ogbunugafor CB, Wylie CS, Diakite I, Weinreich DM, Hartl DL. Adaptive landscape by
676   environment interactions dictate evolutionary dynamics in models of drug resistance.
677   PLoS Comput Biol. 2016 Jan 25;12(1):e1004710.
678   88.   Organization WH. Antimicrobial resistance, Fact sheet N°194,. 2013.
679   89.   Rolff J, Bonhoeffer S, Kloft C, Leistner R, Regoes R, Hochberg ME. Forecasting
680   antimicrobial resistance evolution. Trends Microbiol. 2024 Jan;

Figure legends

Figure 1 | Evolutionary trajectories of uropathogenic bacteria in conditioned media. The

gray dotted graph in each figure represents the maximum antibiotic concentration possible in the

experimental setup per transfer. **A,** The phenotypic trajectories of *E. coli* evolution in the

presence of sulfamethoxazole-trimethoprim, with ratio sulfamethoxazole:trimethoprim 5:1. *E. coli*

was evolved in the presence of the conditioned media of *K. pneumoniae, E. coli, E. faecium* and

S. haemolyticus. The five replicates evolved along similar trajectories, but are more divergent

than those of *K. pneumoniae* and *E. faecium*. **B,** The phenotypic trajectories of *K. pneumoniae*

evolution in the presence of sulfamethoxazole-trimethoprim. Ratio

sulfamethoxazole:trimethoprim is 5:1. *K. pneumoniae* was evolved in the presence of the

conditioned media of *K. pneumoniae, E. coli, E. faecium* and *S. haemolyticus*. Five replicates

were evolved in each conditioned medium and all replicate trajectories overlapped. **C,** The

phenotypic trajectories of *E. faecium* evolution in the presence of vancomycin. *E. faecium* was

evolved in the presence of the same conditioned media as *K. pneumoniae* and *E. coli*, and also

in the conditioned medium of *P. aeruginosa*. The trajectories of all five replicates per conditioned

medium overlapped.

Figure 2 | Correlation of antibiotic concentration at the final transfer with ecological

interactions; WT population size in conditioned medium, and the fold change of antibiotic

tolerance in conditioned medium. Colors indicate the conditioned medium in which the focal

species were cultured, with blue indicating *K. pneumoniae* conditioned medium, red *E. coli*

conditioned medium, yellow *E. faecium* conditioned medium and green *S. haemolyticus*

conditioned medium. The *E. faecium* mutants were also evolved in *P. aeruginosa* conditioned

medium, represented by orange markers. **A**, Trimethoprim concentration (ratio

sulfamethoxazole:trimethoprim is 5:1) at final transfer of *E. coli* lineages in the serial propagation

experiment correlated with the population size interactions of the wild type in the absence of

antibiotics (R^2=0.86, *N*=20, *P*<0.001, means ± s.e.m.). **B**, Trimethoprim concentration (ratio

sulfamethoxazole:trimethoprim is 5:1) at final transfer of *K. pneumoniae* lineages in the serial

propagation experiment correlated with the population size interactions of the wild type in the

absence of antibiotics (R^2=0.45, *N*=20, *P*=0.017, means ± s.e.m.). **C**, Vancomycin concentration

at final transfer of *E. faecium* lineages in the serial propagation experiment correlated with the

population size of the wild type in the absence of antibiotics (R^2=0.93, *N*=25, *P*<0.001,

means ± s.e.m.). **D**, Correlation between antibiotic concentration at final transfer of *E. coli*

mutants of in the serial propagation experiment versus the fold change of the antibiotic tolerance

of the wild-type *E. coli* in the conditioned media compared to artificial urine medium (R^2=0.50,

N=20, *P*<0.001). **E**, Correlation between antibiotic concentration at final transfer of *K.*

pneumoniae mutants of in the serial propagation experiment versus the fold change of the

antibiotic tolerance of the wild-type *K. pneumoniae* in the conditioned media compared to

artificial urine medium (R^2=0.68, *N*=20, *P*<0.001). **F**, Correlation between antibiotic concentration

at final transfer of *E. faecium* mutants of in the serial propagation experiment versus the fold

change of the antibiotic tolerance of the wild-type *E. faecium* in the conditioned media compared

to artificial urine medium (R^2=0.64, *N*=25, *P*<0.001).

Figure 3 | Clustering and mutation repeatability of the final propagation step mutants

clustered on COG-level. Higher values indicate a higher parallelism of the two genotypes

compared. Three mutant lineages were sequenced per conditioned medium (four for *E. faecium*

mutants evolved in *P. aeruginosa* conditioned medium). Comparisons in the blocks around the

diagonal in each figure show the intra-conditioned medium comparisons, whereas the other

blocks show the inter-conditioned media comparisons. **A,** *H*-index of mutation repeatability at

COG-level of *E. coli* evolved in the four conditioned media. Mutator lineages are not taken into

account (Supplementary Figure 1A). **B,** *H*-index of mutation repeatability at COG-level of *K.*

pneumoniae evolved in the four conditioned media. Mutator lineages are not taken into account

(Supplementary Figure 1B). **C,** *H*-index of mutation repeatability at COG-level of *E. faecium*

evolved in the five conditioned media.

Figure 4 | Evolved genetic targets separated by type of mutation. Genetic targets shown

were hit multiple times (at least twice) in either replicate populations or environment, or both.

Size of the spheres indicate the number of unique mutations in the specific gene that were

observed. Genes are separated into their respective COG functional group as shown on the right

side of the figures, with each letter corresponding to a different COG-group. COG-groups: C –

Energy production and conversion, E – Amino acid transport and metabolism, F – Nucleotide

transport and metabolism, G – Carbohydrate transport and metabolism, H – Coenzyme transport

and metabolism, K – Transcription, L – Replication, recombination and repair, M – Cell

wall/membrane/envelope biogenesis, O – Posttranslational modification, protein turnover,

chaperones, P – Inorganic ion transport and metabolism, Q – Secondary metabolites

biosynthesis, transport and catabolism, R – General function prediction only, T Signal

transduction mechanisms, and V – Defense mechanisms. **A,** Genetic targets in *E. coli* evolved in

four different conditioned media. See Supplementary Table 3 for a list of the genes and proteins.

B, Genetic targets in *K. pneumoniae* evolved in four different conditioned media. See

Supplementary Table 4 for a list of the genes and proteins. **C,** Genetic targets in *E. faecium*

evolved in five different conditioned media. See Supplementary Table 5 for a list of the genes

and proteins.