Microbial interactions affect the tempo and mode of antibiotic resistance evolution

Laurens E. Zandbergen[1], Joost van den Heuvel[2], Andrew D. Farr[2,3], Bas J. Zwaan[2], J. Arjan G.

M. de Visser[2], Marjon G. J. de Vos[1,2]*

[1] Groningen Institute of Evolutionary Life Sciences (GELIFES), University of Groningen,

Groningen, The Netherlands

[2] Laboratory of Genetics, Wageningen University, Wageningen, The Netherlands

[3] Max Planck Institute for Evolutionary Biology, Plön, Germany

*corresponding author: m.g.j.de.vos@rug.nl

Competing interests

The authors declare no conflict of interest.

Abstract

The global rise of antibiotic resistance impedes the treatment of bacterial infections. To limit the

emergence and evolution of antibiotic resistance it is important to understand how bacterial

interactions in multispecies communities affect the course of evolution. We investigated how

ecological interactions between microbes derived from polymicrobial urinary tract infections affect

the tempo and mode of antibiotic resistance evolution. We show that for representative strains of

three uropathogens, *Escherichia coli, Klebsiella pneumoniae* and *Enterococcus faecium,* the rate

and evolutionary trajectories towards antibiotic resistance depend on interactions with community

members that mediate their growth and antibiotic tolerance. Replicate lineages of the same

species evolved under similar ecological conditions show parallel evolutionary trajectories. Our

findings demonstrate that bacterial interactions determine the evolutionary potential of antibiotic

resistance evolution.

Introduction

Bacterial pathogens often live in multispecies communities. In such microbial communities bacteria interact, which affects their growth, survival and coexistence (1–5). Bacterial infections are often treated with antibiotics, whose efficacy may also depend on ecological interactions with community members (6). For instance, co-existing bacterial species in pathogenic consortia can mediate the growth and survival of community members in the presence of antibiotics (7–9) through antibiotic degradation (10,11), change the metabolic landscape affecting pathogen fitness (12–14), lower antimicrobial efficacy or alter the transport of antibiotics (7,15), or change the induction of signal molecules that induce resistance mechanisms (16).

Ecological interactions can therefore affect the evolvability of antibiotic resistance in several ways. Firstly, they may alter the mutation supply, by affecting the population size or mutation rate (17–20). Additionally, ecological interactions may alter the fitness effect of mutations, mediating the selective accessibility of adaptive mutations, as genetic and environmental factors are known to modulate the effect of mutations (21–23) and therefore potentially morph the shape of the adaptive landscape (24,25).

In clinical practice, urinary tract infections (UTIs) are treated with antibiotics. Recurrence of infection after the discontinuation of the antimicrobial therapy in treated patients can be high (up to 50% in the elderly population) (26). This is not surprising, as infections are often polymicrobial and many UTI isolates are resistant to commonly prescribed antibiotics (1,27,28). It is therefore important to understand how antibiotic resistance evolves in a community context. To investigate the role of ecological interactions on the evolution of antibiotic resistance, we evolved three focal species (*Escherichia coli, Klebsiella pneumoniae* and *Enterococcus faecium*) in the conditioned medium of several species (*K. pneumoniae*, *E. coli*, *E. faecium*, *Staphylococcus haemolyticus* and *Pseudomonas aeruginosa*). The bacteria were isolated from elderly patients diagnosed with

polymicrobial urinary tract infections (1). These pathogens are part of the so-called ESKAPEE

bacteria, which are known for their ability to evade antibiotics due to their increasing multi-drug

resistance (29). We hypothesize that ecological interactions between uropathogens determine the

evolutionary potential of antibiotic resistance evolution.

Materials and Methods

Bacterial isolates and growth medium

The five uropathogen isolates used in this study were previously collected anonymously from

elderly patients (>70 years old) suffering from urinary tract infection (1). These five isolates (*E.*

coli, *K. pneumonia*, *E. faecium*, *S. haemolyticus¸* and *P. aeruginosa*) were selected based on their

consistent planktonic growth in Artificial Urine Medium (AUM) and for their observed interactions

in a previous study (2,6). Isolates were cultured in Artificial Urine Medium (AUM) (30).

Preparation of conditioned medium

Spent medium of *E. coli*, *K. pneumonia*, *E. faecium*, *S. haemolyticus¸* and *P. aeruginosa* isolates

was prepared by culturing in AUM, followed by centrifugation and filtering. To prepare conditioned

medium, the AUM spent medium fraction was mixed with AUM medium. The resulting conditioned

medium contained metabolite concentrations 0.75x AUM if all of a metabolite was consumed

during culturing and 1.25x AUM if nothing of a metabolite was consumed.

Serial dilution experimental evolution protocol

K. pneumoniae and *E. faecium* were evolved for 30 bi-daily transfers, *E. coli* was evolved for 35

bi-daily transfers. In 5 ml preheated conditioned medium, in 15 ml screw cap falcon tubes, with

the caps slightly opened. Each transfer *K. pneumoniae* and *E. coli* were diluted 500 times (max

~9 generations per bi-daily growth cycle, 2^9=512), and *E. faecium* was diluted 100 times (max

~6/7 generations per bi-daily growth cycle, 2^7=128), due to its smaller population size in the

artificial urine medium. The population sizes of *K. pneumoniae* and *E. coli* are ~10^8 CFU/ml, and

E. faecium ~10^4 in AUM. *K. pneumoniae* and *E. coli* were evolved in the conditioned media of *K.*

pneumoniae, *E. coli*, *E. faecium*, *Staphylococcus haemolyticus* and *E. faecium* also additionally

evolved in *P. aeruginosa* conditioned medium. Each focal isolate-conditioned medium

combination was replicated five times. The Gram-negative isolates *K. pneumoniae* and *E. coli*

were evolved in trimethoprim-sulfamethoxazole (ratio 1:5, as in clinical use (31)) and *E. faecium*

in vancomycin (32). The level of antibiotics at the start of the experiments was set at a 50% sub-

MIC concentration. If the cultures exceeded a predetermined population size threshold, the

antibiotic concentration was increased by 20%, otherwise the antibiotic concentration stayed at

the same level as the previous transfer.

Growth Measurements

Three clones of three of the five evolved replicate lineages of each focal species, evolved in each

conditioned medium, and the wild types of each focal species were tested in the phenotypic growth

assays. Due to a -80°C freezer incident, only one lineage of *E. faecium* evolved in *P. aeruginosa*

conditioned medium could be revived. Each isolate was incubated in three wells (triplicate

cultures) of a preheated 96-well plate (transparent non-treated flat bottom, Brand) containing 200

μL of conditioned medium for 24 h. The plates were inoculated using a 96 solid pin multi-blot

replicator tool, transferring ~1 μL of stock from a −80°C glycerol stock. The population size of wild

type cultures was obtained using the average OD600 at stationary growth after 24h. Growth rates

were obtained by incubation in a BMG Clariostar or Fluorostar microplate reader at 37°C under

aerobic conditions and shaken continuously at 500 rpm. The optical density at 600 nm (OD600)

was measured every 6–7 min in the microplate reader (three flashes, 10 ms settle time). Before

each measurement, the plate was shaken with a double orbital pattern. The growth rate was

defined by the shortest obtained doubling time and quantified by the steepest linear fit of the

exponential phase of log(OD600). Population density was inferred from OD600 absorbance

measurements as well, taking the average for all the measurements of the stationary phase of the

growth curve. Evaporation of media from the plates was not detected. 96-well plates also

contained non-inoculated wells as a control for contamination of both the conditioned medium and

the inoculation step. Contamination was found to be negligible.

Tolerance assays

Wild type isolates of *K. pneumoniae*, *E. faecium* and *E. coli* were inoculated with 2 ul of overnight

cultures in triplicates in conditioned medium and AUM reference medium in a concentration

gradient with two fold increases in antibiotic concentration in 96-well plates. Plates were incubated

for 24h at 37°C. *K. pneumoniae* and *E. coli* were assayed in the presence of trimethoprim-

sulfamethoxazole and *E. faecium* in the presence of vancomycin. The fold change tolerance was

calculated by dividing the concentration that limited growth in a particular conditioned medium by

the concentration of antibiotic that limited growth in AUM.

MIC assays

Starting Minimal Inhibitory Concentration (MIC) of the wild-types were measured in a serial dilution

assay of the desired antibiotics. Bacterial cultures in AUM with a volume of 200 µL were created

in a 96-well plate including a two-fold serial dilution of the desired antibiotic (vancomycin for *E.*

faecium and a 1:5 mixture of trimethoprim and sulfamethoxazole respectively for *E. coli* and *K.*

pneumoniae) ranging from 0.125 to 1 µg/mL for the vancomycin and 0.00625 to 0.2 µg/mL for

trimethoprim:sulfamethoxazole in triplicate. Cultures without antibiotics were used as control. The

serial dilution assay was grown over 24h at 37°C at 200 rpm. Whether or not bacteria had visibly

grown was determined by measuring OD600 and subtracting AUM without bacteria as blank.

OD600 values <0.05 were inspected visually to determine whether inhibition was complete or not.

MIC of final transfer mutants were determined on agar using a modified version of the TDtest

described previously by Gefen et al. (33).

DNA extraction

Cultures of evolved clones were grown overnight in 5ml LB, cells were pelleted, and DNA was

extracted using a Maxwell 16 Instrument in combination with the Maxwell® 16 Tissue LEV Total

RNA Purification Kit Custom (XAS 1220) (Promega, Madison, WI, USA) following manufacturer's

instructions. The DNA was eluted in 30 µL DNAse free water, and quantified using Nanodrop

2000c Spectrophotometer (Thermo Fisher Scientific, USA) and gel electrophoresis.

Sequencing

Three clones of three of the five evolved replicate lineages of each focal species, evolved in each

conditioned medium, were prepared for sequencing. Sequencing was performed at Baseclear BV

Leiden. Paired-end sequence reads were generated using the Illumina HiSeq2500 or MiSeq

system.

Calculation *H*-index

To quantify how repeatable the evolution of the mutants was, the *H*-index of genome

repeatability previously described by Schenk et al. (17) was used. The *H*-index was calculated

on the functional (COG) and at the gene level. All mutants were compared pairwise, calculated

using (17):

$$H_{A,B} = \sum_{j=1}^{m} \sum_{k=1}^{n} \left(\frac{|A_j \cap B_k|}{|A_j|} \right) / \sum_{j=1}^{m} \sum_{k=1}^{m} \left(\frac{|A_j \cap A_l|}{|A_j|} \right) \qquad (1)$$

Which compares genotype *A* and *B* with mutations *m* and *n* respectively. The *H*-index is

calculated for both genotypes separately ($H_{A,B}$ and $H_{B,A}$) and compared using (17):

$$H = \frac{H_{A,B} + H_{B,A}}{2} \qquad (2)$$

The final *H*-index value would fall between 0 (no repeated evolution) and 1 (complete

parallelism). When the *H*-index is 0.5, it indicates that the average of the two compared mutants

have half of their mutations shared between them.

Statistical analysis

For each genotype, measurements (of phenotypes such as on growth rate, population size,

resistance or tolerance levels) were performed in either triplicates or quadruplicates. Throughout

all of the analyses, whenever the values between two conditions of mutants were compared, a

two-tailed Welch's *t*-test was performed to test the null hypothesis that the two parameters do not

have significantly different values (hypothesis of neutrality). We used a Poisson-test to assess

whether the number of mutations in the mutators was significantly different in the hypermutators

and the non-hypermutators. We used a Pearson correlation coefficient and corresponding *p*-

values to determine the level of correlation. The null hypothesis was rejected if the *p*-value

obtained from the test was lower than a significant level of 0.05.

Additional information on Materials and Methods can be found in Supplementary text 1.

Results

Conditioned media promote the adaptive radiation of antibiotic resistance evolution

To investigate the effect of ecological interactions on the evolution of antibiotic resistance, we

evolved single isolates of focal species *K. pneumoniae*, *E. coli* and *E. faecium* by bi-daily serial

transfers in conditioned artificial urine medium (AUM) supplemented with increasing

concentrations of antibiotics. Conditioned medium was used as a proxy for microbial interactions,

which was prepared by culturing isolates in AUM followed by filtering out bacteria and

replenishment of the medium with fresh nutrients (Materials and Methods) (6). When the evolving

isolates had reached a predetermined population size threshold at the moment of transfer, the

antibiotic concentration was increased by 20% (Materials and Methods). Each combination of focal

isolate and conditioned medium was replicated five times. The two Gram-negative species, *K.*

pneumoniae and *E. coli*, were evolved in the presence of trimethoprim-sulfamethoxazole (ratio

1:5, similar to clinical use (31)), which inhibits DNA synthesis (34). *E. faecium* was evolved in the

presence of the Gram-positive specific antibiotic vancomycin, which inhibits cell wall synthesis

(35); initial antibiotic concentrations were 50% of the minimal inhibitory concentration (MIC) in all

cases. *E. coli* was evolved over 70 days (35 transfers) and *K. pneumoniae* and *E. faecium* for 60

days (30 transfers) (Materials and Methods).

We find that the ecological interactions alter the course of antibiotic resistance evolution.

Specifically, these interactions affect the rate and final level of antibiotic tolerance increase (Figure

1). Notably, the evolutionary trajectories of the replicates per conditioned medium are highly

parallel, and the parallel trajectories in each conditioned medium diverge from other conditioned

media. This is especially the case for *K. pneumoniae* and *E. faecium* (Figure 1B,C). The

evolutionary trajectories of *E. coli* per conditioned medium environment also display parallelism

(Figure 1A), but are more variable compared to *K. pneumoniae* and *E. faecium* (Figure 1B,C).

Generally, we find that the rate and level of antibiotic resistance evolution is slowest in the

conditioned medium of the focal isolate (e.g. *K. pneumoniae* in *K. pneumoniae* conditioned

medium, and *E. faecium* in *E. faecium* conditioned medium, Figure 1B,C). Moreover, the rate of

evolution is often fastest in the conditioned medium of the isolate that is phylogenetically least

related, e.g., *E. faecium* evolves fastest in the Gram-negative conditioned media, and the *E. coli*

evolves fastest in the Gram-positive conditioned media. This suggests that niche specific effects

such as avoidance of competition promote the divergence of antibiotic resistance evolution, as

also observed in other studies (36–38).

Response to selection correlates with ecological effects on population size

Bacterial interactions, due to for instance metabolic interactions, can induce niche specific effects that dictate the course of evolution. Such effects can be mediated by the specific chemical composition of the conditioned media affecting the population size and/or growth rate, and therefore the mutation supply rate (20). To investigate whether ecological interactions mediate the evolutionary potential by affecting the mutation supply rate of evolving populations, we correlate the maximum population sizes, as inferred by OD600 measurements, of the ancestors in conditioned medium, with the final antibiotic levels tolerated in the evolution experiments (Figure 2A-C). Indeed, the antibiotic concentration at the final transfer correlates significantly with the population size in conditioned medium for all three focal species. We find a strong positive correlation for *E. coli* and *E. faecium*, and a moderate correlation for *K. pneumoniae* (Pearson correlation coefficient 0.86, 0.93 and 0.45, for *E. coli*, *E. faecium* and *K. pneumoniae* resp., $p <$ 0.001 for *E. coli* ($N = 20$) and *E. faecium* ($N = 25$), and $p = 0.017$ for *K. pneumoniae* ($N = 20$)) (Figure 2A-C).

We previously found that growth-mediating effects of bacterial interactions in the absence of antibiotics correlate with antibiotic tolerance levels (6). The fold change in antibiotic tolerance was calculated based on the change in growth inhibiting antibiotic concentration of conditioned media compared to unconditioned medium. Here, we find that the concentration of antibiotics at the last transfer of the experiment correlates positively and significantly with the antibiotic tolerance levels conferred by the conditioned media (Pearson correlation coefficients, 0.50, 0.68 and 0.64 for *E. coli* ($N = 20$), *K. pneumoniae* ($N = 20$), and *E. faecium* ($N = 25$) resp., for all $p < 0.001$, Figure 2D-F). This suggests that ecological interactions may accelerate antibiotic resistance evolution via increased antibiotic tolerance, which enhances the mutation supply by allowing a larger standing population size.

Ecological interactions do not affect the number of genetic changes

To investigate the consequences of ecological interactions on mutational trajectories, we determined the genetic changes in these evolved lineages through Illumina sequencing of three clones per evolved lineage (Materials and Methods). A fundamental determinant of evolutionary trajectories is the population size, as it affects both the supply of mutations and the efficacy of natural selection to select beneficial mutations (17,20,39). Therefore, effects on population size may lead to different mutations being selected (40).

To quantify the impact of natural selection in our populations, we measured the dN/dS ratio (41). Generally, we find dN/dS ratios $\gg$ 1, resp. *K. pneumoniae* 7, *E. faecium* 17, and *E. coli* 5, indicating strong positive selection in all populations.

We did not find a strong positive correlation between the number of SNPs in the evolved lineages correlates and population size in conditioned media in the absence of antibiotics. We find a weak positive correlation for *E. coli* and *E. faecium* (Pearson correlation coefficient 0.18, 0.47, $p = 0.17$, 0.20, resp., Figure 3A,C), and a negative correlation for *K. pneumoniae* (Pearson correlation coefficient, -0.70, $p < 0.001$, Figure 3B). SNPs outnumbered other mutations in the evolved isolates (Figure 3D-F). Yet other genetic differences were also observed in the evolved lineages, such as deletions, inversions and copy-number differences. Moreover, in several of these *K. pneumoniae* lineages one of the plasmids was entirely lost. Especially for the second plasmid (Supplementary Figure 1B), we observed that blocks of the plasmid were deleted during evolution. In general, mutational patterns did not seem related to the specific conditioned media.

Additionally, hyper-mutator lineages appeared. These lineages accumulated many more mutations compared to the other evolved lineages of these species: around 60 mutations in comparison to the average of ≈10, for *E. coli* and *K. pneumoniae* (Figure 3D, E, Poisson-test, $p < 0.001$). For this reason, these lineages were removed from the SNP count analysis. The

sequences of the *E. coli* hyper-mutator lineages showed large deletions (>3KB) in the region which

includes *mutH*, a gene involved in mismatch repair (42), which points to a role for *mutH* in the

generation of these hyper mutator lineages. The *K. pneumoniae* hyper-mutator lineage appeared

in *E. faecium* conditioned medium, and was associated with a missense SNP mutation in *mutS*.

We do not find a strong relationship between the total number of genetic changes in evolved

lineages, and the population size in conditioned media (Pearson correlation coefficient, 0.023 and

0.03, 0.15, *p* = 0.68, 0.59, 0.19 for *E. coli, K. pneumoniae* and *E. faecium* lineages respectively,

Supplementary Figure 2A-C). This suggests an important role for clonal interference during

adaptation, which affects the type and not the number of mutations in populations with different

mutation supplies (17).

Ecological interactions do affect the types of genetic changes

To test whether different conditioned media caused different adaptive trajectories in the three focal

species, we compared the genetic similarity of isolates evolved in the same with those evolved in

different conditioned media. To this aim we calculated the *H*-index (17), i.e. a measure for the

fraction of mutated genes that two evolved isolates have in common (Materials and Methods). If

we calculate this index at the level of Clusters of Orthologous Groups (COGs) (43), i.e. genes

grouped by function, we find more parallelism among replicate populations evolved within the

same conditioned media compared to those same lineages in different conditioned media (*p* =

0.043, 0.043, 0.032 (*Welch's t-test*) for *E. coli*, *K. pneumoniae* and *E. faecium,* respectively, Figure

4). This indicates that the parallel evolution observed (Figure 1), compliments the parallelism

observed at the functional level in the same conditioned media.

For *E. coli*, particularly the lineages that evolved in the Gram-positive conditioned media of *S.*

haemolyticus and *E. faecium* display a high degree of parallelism at the COG-level (*H* > 0.6, Figure

4A). This pattern is conserved at the gene-level for lineages evolved in *S. haemolyticus*

conditioned medium (Supplementary Figure 3A). The high genetic parallelism corresponds to a

similar phenotypic growth response of *E. coli* replicates to these conditioned media (Figure 2D),

and to their similar evolutionary trajectories (Figure 1A). Interestingly, the population size of *E. coli*

in the conditioned medium of itself is also similar to the population size in the conditioned medium

of *S. haemolyticus* (Figure 2A). Yet, the *H*-index values that correspond to the *E. coli* lineages

evolved in these Gram-positive (*S. haemolyticus)* and Gram-negative (*E. coli)* conditioned media

are relatively low (*H* < 0.6), indicating lower parallelism within these conditions. This shows that

population size alone cannot explain the level of parallelism, which suggests that the particular

metabolic constituents of the conditioned medium, also play a role in the genetic divergence of

these evolutionary trajectories.

For *E. faecium*, we find a particularly strong parallelism at the COG-level for evolved lineages in

P. aeruginosa conditioned medium, but less so for other conditioned media (Figure 4C). At the

gene-level, this strong parallelism for evolved lineages in *P. aeruginosa* conditioned medium is

not present (Supplementary Figure 3C). The high level of parallelism among *E. faecium* lineages

evolved in *P. aeruginosa* conditioned medium coincides with their highest rate of evolution (Figure

1C). The level of parallelism of *E. faecium* evolved in the conditioned media of *K. pneumoniae* and

S. haemolyticus is lower, despite similar resistance trajectories and population sizes (*H*-index

means of 0.75 for *P. aeruginosa,* and 0.35 (p = 0.022) and 0.48 (p = 0.013) for *K. pneumoniae*

and *S. haemolyticus* resp., *Welch's t*-tests *H*-index comparison of *E. faecium* in the respective

conditioned media) (Figure 1B, 4B). This again suggests that particular metabolic constituents of

the conditioned medium play a role in the genetic divergence.

For *K. pneumoniae*, we find consistently high *H*-indices (*H* > 0.6) at the COG-level across

conditions, except for the comparison of the isolates evolved in *E. faecium* and *E. coli* conditioned

media (Figure 4B). In addition, at the gene-level we also find relatively high parallelism (although

slightly lower than at the COG-level), especially for the isolates evolved within *S. haemolyticus*

and *K. pneumoniae* conditioned medium (Supplementary Figure 3B). Interestingly, the generally

high parallelism at the COG-level for *K. pneumoniae* is associated with small population size

differences due to the exposure to these conditioned media (approximately two-fold). These

combined highlight the rather limited effect of conditioned media on the antibiotic resistance

evolution of *K. pneumoniae* (Figure 2B).

As observed before (17,44), the *H*-indices at the COG-level show a higher parallelism than at the

gene-level (Figure 4, Supplementary Figure 3). This indicates that there is genetic redundancy in

the functionalities that are selected for under these particular ecological circumstances. Generally,

the higher *H*-index of lineages evolved within the same conditioned medium (along the diagonal

axis in Figure 4) shows that the antibiotic resistance evolution in the conditioned media leads to

parallel genotypic (Figure 4) and phenotypic (Figure 1) adaptation under the influence of species-

specific bacterial metabolites.

Conditioned medium causes specific genetic changes

Next, we investigated whether the species-specific ecological interactions also lead to the

selection of different resistance mechanisms. We do find that in particular conditioned media

specific mutations occur repeatedly, whereas others do not (Figure 5A-C). We hypothesized that

in larger populations, larger effect-size mutations are selected (17). Indeed, if we look at the

resistance conferring mutations of *E. faecium*, we find that the lineages with the smallest

population size, which are those cultured in its own conditioned medium, repeatedly had

substitutions in *ompR*, a two-component response regulator of *ompF* involved in the transport of

antibiotics 41 42 (45,46) (Figure 5C). Efflux-enhancing mutations have previously been associated

with lower selective pressures (47). *E. faecium* lineages evolved in other conditioned media

reached higher levels of resistance and incorporated mutations in well-known vancomycin

resistance targets, *VanR, VanS* (45,48,49), or both. *VanS* is a sensor histidine kinase that detects

vancomycin and activates *VanR* (45,48,50). *VanR* is a transcriptional activator, regulating the

transcription of the vancomycin-resistance operon together with regulating transcription of the

vanHAX operon, resulting in higher tolerance due to changes in peptidoglycan formation (45,49).

Indeed, lineages that had mutations in these targets generally had higher resistance levels and

larger population sizes (Figure 5C, 2C).

The two Gram-negative lineages (*K. pneumoniae* and *E. coli*) were selected for resistance to

trimethoprim-sulfamethoxazole. All sequenced isolates had mutations in *folA*, dihydrofolate

reductase (Figure 5A,B), which is involved in the folate pathway that is essential for DNA synthesis

and is inhibited by trimethoprim-sulfamethoxazole (51). *FolA* is the direct target of trimethoprim.

One lineage of *E. coli* evolved in *E. coli* conditioned medium even had a 20-fold amplification of

the copy-number of this gene.

Both *K. pneumoniae* and *E. coli* incorporated *folM* mutations. *FolM* encodes an isozyme of

dihydrofolate reductase and is also involved in DNA synthesis. Inhibition of this gene causes the

upregulation of *folA,* but usually also requires resistance mutations in *folA* to be effective (52). *E.*

coli incorporated *folM* mutations only in lineages that evolved slowly in the *K. pneumoniae*

conditioned medium (Figure 5A). In *K. pneumoniae*, *folM* mutations appeared in lineages that

were evolved in *K. pneumoniae* medium*,* but also in the *E. coli* and *S. haemolyticus* conditioned

media, which allowed fast resistance evolution.

In *K. pneumoniae* in *E. faecium* conditioned medium, *folX* was targeted instead of *folM* (Figure

5B). *E. coli* lineages evolved in *E. coli* and *S. haemolyticus* conditioned medium incorporated

mutations in yet another gene in the folate pathway, *folP*, (53) (Figure 5A). All these mutations in

the folate pathway correspond to clinically relevant mutations (Supplementary text 2). The

observation that different genes within the folate pathway are targeted in the different conditioned

media suggests that ecological interactions may cause different trajectories of antibiotic resistance

evolution.

The observed mutations also confer resistance to antibiotics in the absence of conditioned media

(Supplementary text 2, Supplementary figure 5). Specifically, for *E. coli* and *E. faecium*, the

antibiotic level tolerated by the populations at the end of the serial transfer experiment correlated

positively with the measured MICs, while for *K. pneumoniae* we found this relationship to be

inverted. Moreover, the growth rate and population size of evolved lineages is mediated by

exposure to specific conditioned media (Supplementary text 3, Supplementary Figure 6-8) and

antibiotics interactions (Supplementary text 3, Supplementary Figure 6E). Concluding, specific

antibiotic resistance genes were targeted repeatedly within, but not across conditioned media

(Figure 5). This shows that the genetic basis of antibiotic resistance evolution is mediated by the

genetic background of the focal species, as well as by the ecological interactions.

Discussion and conclusion

We find that the bacterial interactions mediated by conditioned medium altered the rate and

mutational trajectories of antibiotic resistance evolution of uropathogenic isolates *E. coli*, *K.*

pneumoniae and *E. faecium*, derived from polymicrobial urinary tract infections. The conditioned

medium approach has some important caveats. In the conditioned media environment, cultured

bacteria were not in physical contact with the bacteria used to create conditioned media. Such

contact-mediated interactions may affect eco-evolutionary outcomes, since two populations of

bacteria can respond to the presence and the produced metabolites of the other when in the same

environment (54,55). Moreover, higher-order interactions between more than two bacterial

species could potentially affect growth (56,57). Also, cross-feeding metabolites may mediate

ecological interactions by altering gene expression in both interaction partners, in these

experiment cross-feeding was only limited to metabolites produced by the donor in conditioned

media. Future studies are needed to investigate such additional factors that can affect the

evolution of antibiotic resistance in the context of ecological interactions.

We find a positive relationship between the level of evolved antibiotic resistance and the ecological

interactions in the absence of the antibiotics: the larger the population size of the unevolved strains

in the conditioned medium and the higher the tolerance conferred by the conditioned medium, the

higher the final concentration of antibiotic that can be sustained in the serial dilution experiment.

Population size is known to affect the sensitivity to antibiotics known as the inoculum effect, and

although this is usually observed for β-lactamases (11), there is evidence that it plays a role for

antibiotics like trimethoprim as well, even though these are not degraded enzymatically (58,59).

This is also in line with other work that showed that community composition can alter the selection

for antibiotic resistance in a community context, for instance through modulating population

densities via competitive or cooperative interactions (58,60–65).

Overall, we found parallelism in lineages evolved under the same ecological conditions or in the

same conditioned medium (Figure 1), and that resistance mutations and other functional targets

selected were dependent on these conditions (Figure 4, 5). Parallelism has been observed to

depend on the environmental conditions before, such as resource availability (66), ecological

interactions (67) and population size of evolving lineages (17), but the phenotypic parallelism does

not always correspond to molecular parallelism (68). This work shows that ecological interactions,

which may be caused by avoidance of resource competition, cross-feeding and potentially other

antibiotic tolerance effects (6,38,61), can drive both phenotypic and genetic parallel antibiotic

resistance evolution.

It is important to recognize that only the final transfer replicates were sequenced, meaning that it

is unclear when specific mutations fixed in the population and how those affected the evolutionary

trajectories and by extent the adaptive landscape. Following mutational trajectories during the

emergence of antibiotic resistance evolution could lead to more insight in the environmental impact

on resistance by identifying genotypes that alter these trajectories (69–73).

For *E. coli* and *E. faecium*, the antibiotic level tolerated by the populations at the end of the serial

transfer experiment correlated with their respective MICs, while for *K. pneumoniae* we found this

relationship to be inverted. The *K. pneumoniae* lineages with the highest evolved MIC

corresponded to those cultured in the lowest concentration of antibiotics. Potentially, larger

population sizes corresponded to more 'greedy' adaptation, promoting faster initial adaptation, but

lower final adaptation, as measured by evolved MIC levels (Supplementary Figure 4B) (74). Such

balance between greedy and slower-and-steady evolution may have led to higher resistance

levels for the initially smallest population sizes, such as *K. pneumoniae* in *K. pneumoniae*

conditioned medium (Figure 1). Additionally, we showed that evolved resistance levels depend on

the environment. Ecological interactions, mediating population sizes and other ecological factors,

affect the antibiotic resistance levels as measured in isolation (e.g., in the clinic), which is in

accordance with other studies (6,64,75).

Concluding, antimicrobial resistance is turning into a 'silent pandemic' (47), as infections caused

by resistant microorganisms fail to respond to conventional treatment, resulting in prolonged

illness, greater risk of death and higher healthcare costs (76). We find that ecological interactions

between bacteria, by altering population size and conferred antibiotic tolerance, affect the mode,

the genetic functions selected, and the tempo, the rate of antibiotic resistance evolution. The

identification of the nature of such ecological interactions can aid in forecasting the evolutionary

potential of antibiotic resistance evolution. Further knowledge of such predictive parameters can

improve decision making, which will be crucial for developing interventions, and for curbing the

emergence and further evolution of antibiotic resistance (77).

413 **Acknowledgements**

414

415 We thank Ineke Heikamp-de Jong for help with the DNA extraction, and Bart Thomma for the use

416 of the MLII laboratory at Wageningen University. MGJDV was supported by an NWO VENI

417 Fellowship (863.14.015), ZonMw ETH grant (40-43500-98-4049/435004009) and L'Oreal Unesco

418 LNVH NIAS-KNAW For Women in Science Fellowship, part of this work was performed at NIAS-

419 KNAW. LEZ was supported by the Adaptive Life program of the Faculty of Science and

420 Engineering, within GELIFES.

421 **Contributions**

422

423 Conceived and designed the study: LEZ JAGMDV BJZ MGJDV. Performed the experiments: LEZ

424 ADF MGJDV. Analyzed the data: LEZ JVDH MGJDV. Wrote the manuscript: LEZ JVDH ADF

425 JAGMDV MGJDV.

426 **Data availability statement**

427

428 Bacterial sequence data have been deposited with links to BioProject accession number

429 PRJNA1103400 in the NCBI BioProject database. The other datasets generated during and/or

430 analysed during the current study are available from the corresponding author on reasonable

431 request.

432 **References**

433 1.   Croxall G, Hale J, Weston V, Manning G, Cheetham P, Achtman M, et al. Molecular

434   epidemiology of extraintestinal pathogenic *Escherichia coli* isolates from a regional cohort

435 of elderly patients highlights the prevalence of ST131 strains with increased antimicrobial

436 resistance in both community and hospital care settings. J Antimicrob Chemother.

437 66(11):2501–8.

438 2. Zandbergen LE, Halverson T, Brons JK, Wolfe AJ, de Vos MGJ. The Good and the Bad:

439 Ecological Interaction Measurements Between the Urinary Microbiota and Uropathogens.

440 Front Microbiol. 2021;12(NA):659450.

441 3. Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks, competition,

442 and stability. Science (80-). 2015;350(6261):663–6.

443 4. Chang CY, Bajić D, Vila JCC, Estrela S, Sanchez A. Emergent coexistence in

444 multispecies microbial communities. Science 2023;381(6655):343–8.

445 5. Fredrickson AG, Stephanopoulos G. Microbial competition. Science 1981;213(4511):972–

446 9.

447 6. De Vos MGJ, Zagorski M, McNally A, Bollenbach T. Interaction networks, ecological

448 stability, and collective antibiotic tolerance in polymicrobial infections. Proc Natl Acad Sci

449 U S A. 2017;114(40):10666–71.

450 7. Lara EG, van der Windt I, Molenaar D, de Vos MGJ, Melkonian C. Using functional

451 annotations to study pairwise interactions in urinary tract infection communities. Genes

452 2021;12(8):1221.

453 8. Ramsey MM, Rumbaugh KP, Whiteley M. Metabolite cross-feeding enhances virulence in

454 a model polymicrobial infection. PLoS Pathog. 2011;7(3):e1002012.

455 9. Wintermute EH, Silver PA. Emergent cooperation in microbial metabolism. Mol Syst Biol.

456 2010;6:407.

457 10. Yurtsev EA, Chao HX, Datta MS, Artemova T, Gore J. Bacterial cheating drives the

458 population dynamics of cooperative antibiotic resistance plasmids. Mol Syst Biol.

459 2013;9(683):683.

460 11. Geyrhofer L, Ruelens P, Farr AD, Pesce D, de Visser JAGM, Brenner N. Minimal

Surviving Inoculum in Collective Antibiotic Resistance. Regoes RR, Levin BR, editors.

MBio. 2023 Apr 25;14(2).

12.   Melkonian C, Seidl MF, van der Hooft JJJ, de Vos MGJ. Metabolic interactions shape a

community's phenotype. Trends Microbiol. 2022;30(7):609–11.

13.   Yu JSL, Correia-Melo C, Zorrilla F, Herrera-Dominguez L, Wu MY, Hartl J, et al. Microbial

communities form rich extracellular metabolomes that foster metabolic interactions and

promote drug tolerance. Nat Microbiol. 2022;7(4):542–55.

14.   Colegrave N, Buckling A. Microbial experiments on adaptive landscapes. Bioessays.

2005;27(11):1167–73.

15.   Delcour AH. Outer membrane permeability and antibiotic resistance. Biochim Biophys

Acta - Proteins Proteomics. 2009;1794(5):808–16.

16.   Li Y, Feng T, Wang Y. The role of bacterial signaling networks in antibiotics response and

resistance regulation. Mar Life Sci Technol. 2022;4(2):163–78.

17.   Schenk MF, Zwart MP, Hwang S, Ruelens P, Severing E, Krug J, et al. Population size

mediates the contribution of high-rate and large-benefit mutations to parallel evolution.

Vol. 6, Nature Ecology and Evolution. Dryad; 2022. p. 439–47.

18.   Garoff L, Pietsch F, Huseby DL, Lilja T, Brandis G, Hughes D. Population bottlenecks

strongly influence the evolutionary trajectory to fluoroquinolone resistance in *escherichia*

coli. Barlow M, editor. Mol Biol Evol. 2020 Jun 1;37(6):1637–46.

19.   Mahrt N, Tietze A, Künzel S, Franzenburg S, Barbosa C, Jansen G, et al. Bottleneck size

and selection level reproducibly impact evolution of antibiotic resistance. Nat Ecol Evol.

2021 Jul 26;5(9):1233–42.

20.   De Visser JAGM, Zeyl CW, Gerrish PJ, Blanchard JL, Lenski RE. Diminishing returns

from mutation supply rate in asexual populations. Science 1999;283(5400):404–6.

21.   de Vos MGJ, Poelwijk FJ, Battich N, Ndika JD, Tans SJ. Environmental dependence of

genetic constraint. PLoS Genet. 2013;9(6):e1003580.

22. Flynn KM, Cooper TF, Moore FB, Cooper VS. The environment affects epistatic

interactions to alter the topology of an empirical fitness landscape. PLoS Genet.

2013;9(4):e1003426.

23. Lindsey HA, Gallie J, Taylor S, Kerr B. Evolutionary rescue from extinction is contingent

on a lower rate of environmental change. Nature. 2013;494(7438):463–7.

24. De Vos MGJ, Schoustra SE, De Visser JAGM. Ecology dictates evolution? about the

importance of genetic and ecological constraints in adaptation. Epl. 2018;122(5).

25. Wright S. The roles of mutation, inbreeding, crossbreeding and selection in evolution.

Proc Sixth Int Congr Genet. 1932;1:356–366.

26. Nicolle LE. Urinary Tract Infections in Long-Term–Care Facilities. Infect Control Hosp

Epidemiol. 2001;22(3):167–75.

27. Siegman-Igra Y, Kulka T, Schwartz D, Konforti N. Polymicrobial and monomicrobial

bacteraemic urinary tract infection. J Hosp Infect. 1994;28(1):49–56.

28. Kline KA, Lewis AL. Gram-positive uropathogens, polymicrobial urinary tract infection, and

the emerging microbiota of the urinary tract. Urin Tract Infect Mol Pathog Clin Manag.

2016;459–502.

29. Y. Mahmood H, Jamshidi S, Mark Sutton J, M. Rahman K. Current Advances in

Developing Inhibitors of Bacterial Multidrug Efflux Pumps. Curr Med Chem. 2016 Apr

7;23(10):1062–81.

30. Brooks T, Keevil CW. A simple artificial urine for the growth of urinary pathogens. Lett

Appl Microbiol. 1997;24(3):203–6.

31. Hitchings GH. Mechanism of action of trimethoprim-sulfamethoxazole. J Infect Dis.

1973;128(Supplement 3):S433.

32. Levine DP. Vancomycin: A history. Clin Infect Dis. 2006 Jan 1;42(SUPPL. 1):S5–12.

33. Gefen O, Chekol B, Strahilevitz J, Balaban NQ. TDtest: Easy detection of bacterial

tolerance and persistence in clinical isolates by a modified disk-diffusion assay. Sci Rep.

2017;7:1–9.

34.   Masters PA, O'Bryan TA, Zurlo J, Miller DQ, Joshi N. Trimethoprim-sulfamethoxazole

revisited. Arch Intern Med. 2003;163(4):402–10.

35.   Bourgeois-Nicolaos N, Moubareck C, Mangeney N, Butel MJ, Doucet-Populaire F.

Comparative study of vanA gene transfer from *Enterococcus faecium* to *Enterococcus*

faecalis and to *Enterococcus faecium* in the intestine of mice. FEMS Microbiol Lett.

2006;254(1):27–33.

36.   Zampieri M, Enke T, Chubukov V, Ricci V, Piddock L, Sauer U. Metabolic constraints on

the evolution of antibiotic resistance. Mol Syst Biol. 2017;13(3):917.

37.   Laborda P, Martínez JL, Hernando-Amado S. Evolution of Habitat-Dependent Antibiotic

Resistance in *Pseudomonas aeruginosa*. Garcia-Solache MA, editor. Microbiol Spectr.

2022 Aug 31;10(4).

38.   Windels EM, Cool L, Persy E, Swinnen J, Matthay P, Van B, et al. Antibiotic dose and

nutrient availability differentially drive the evolution of antibiotic resistance and

persistence. ISME J. 2024 Jan 8;18(1):1–30.

39.   Jiang X, Mu B, Huang Z, Zhang M, Wang X, Tao S. Impacts of mutation effects and

population size on mutation rate in asexual populations: A simulation study. BMC Evol

Biol. 2010 Dec 30;10(1):298.

40.   Kauffman S, Levin S. Towards a general theory of adaptive walks on rugged landscapes.

J Theor Biol. 1987;128(1):11–45.

41.   Kryazhimskiy S, Plotkin JB. The population genetics of dN/dS. Gojobori T, editor. PLoS

Genet. 2008 Dec 12;4(12):e1000304.

42.   Acharya S, Foster PL, Brooks P, Fishel R. The coordinated functions of the *E. coli* MutS

and MutL proteins in mismatch repair. Mol Cell. 2003;12(1):233–46.

43.   Tatusov RL, Koonin E V, Lipman DJ. A genomic perspective on protein families. Science.

1997 Oct 24;278(5338):631–7.

44.   Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, et al.

The molecular diversity of adaptive convergence. Science 2012;335(6067):457–61.

45.   Arthur M, Molinas C, Courvalin P. The VanS-VanR two-component regulatory system

controls synthesis of depsipeptide peptidoglycan precursors in *Enterococcus faecium*

BM4147. J Bacteriol. 1992;174(8):2582–91.

46.   Hancock L, Perego M. Two-component signal transduction in *Enterococcus faecalis*. J

Bacteriol. 2002;184(21):5819–25.

47.   Cars O, Chandy SJ, Mpundu M, Peralta AQ, Zorzet A, So AD. Resetting the agenda for

antibiotic resistance through a health systems perspective. Lancet Glob Heal.

2021;9(7):e1022–7.

48.   Evers S, Courvalin P. Regulation of VanB-type vancomycin resistance gene expression

by the VanSB-VanRB two-component regulatory system in *Enterococcus faecalis* V583. J

Bacteriol. 1996;178(5):1302–9.

49.   Depardieu F, Courvalin P, Kolb A. Binding sites of VanRB and σ70 RNA polymerase in

the vanB vancomycin resistance operon of *Enterococcus faecium* BM4524. Mol Microbiol.

2005;57(2):550–64.

50.   Guffey AA, Loll PJ. Regulation of resistance in vancomycin-resistant *enterococci*: The

vanrs two-component system. Microorganisms. 2021 Sep 25;9(10):2026.

51.   Hitchings GH, Burchall JJ. Inhibition of folate biosynthesis and function as a basis for

chemotherapy. In: Advances in enzymology and related areas of molecular biology.

Hoboken, NJ, USA: John Wiley & Sons, Inc.; 1965. p. 417–68.

52.   Giladi M, Altman-Price N, Levin I, Levy L, Mevarech M. FolM, A New Chromosomally

Encoded Dihydrofolate Reductase in *Escherichia coli*. J Bacteriol. 2003;185(23):7015–8.

53.   Buwembo W, Aery S, Rwenyonyi CM, Swedberg G, Kironde F. Point mutations in the folP

gene partly explain sulfonamide resistance of *streptococcus* mutans. Int J Microbiol.

2013;2013:367021.

54. Brunet YR, Espinosa L, Harchouni S, Mignot T, Cascales E. Imaging Type VI Secretion-

Mediated Bacterial Killing. Cell Rep. 2013;3(1):36–41.

55. Weiland-Bräuer N. Friends or Foes: Microbial Interactions in Nature. 2021 Jun

2;10(6):496.

56. Kelsic ED, Zhao J, Vetsigian K, Kishony R. Counteraction of antibiotic production and

degradation stabilizes microbial communities. Nature. 2015;521(7553):516–9.

57. Weinreich DM, Lan Y, Jaffe J, Heckendorn RB. The Influence of Higher-Order Epistasis

on Biological Fitness Landscape Topography. J Stat Phys. 2018;172(1):208–25.

58. Karslake J, Maltas J, Brumm P, Wood KB. Population Density Modulates Drug Inhibition

and Gives Rise to Potential Bistability of Treatment Outcomes for Bacterial Infections.

PLoS Comput Biol. 2016;12(10):e1005098.

59. Alvarez S, DeMaria A, Kulkarni R, Klein JO, McCabe WR. Interactions of rifampin and

trimethoprim in vitro. Rev Infect Dis. 1982 Mar 1;4(2):390–401.

60. Rezzoagli C, Granato ET, Kümmerli R. Harnessing bacterial interactions to manage

infections: A review on the opportunistic pathogen *Pseudomonas aeruginosa* as a case

example. J Med Microbiol. 2020;69(2):147–61.

61. Meredith HR, Srimani JK, Lee AJ, Lopatkin AJ, You L. Collective antibiotic tolerance:

Mechanisms, dynamics and intervention. Nat Chem Biol. 2015;11(3):182–8.

62. Vega NM, Gore J. Collective antibiotic resistance: Mechanisms and implications. Curr

Opin Microbiol. 2014;21:28–34.

63. Adamowicz EM, Muza M, Chacón JM, Harcombe WR. Cross-feeding modulates the rate

and mechanism of antibiotic resistance evolution in a model microbial community of

Escherichia coli and *Salmonella enterica*. Boshoff HI, editor. PLoS Pathog. 2020 Jul

20;16(7):e1008700.

64. Bottery MJ, Pitchford JW, Friman VP. Ecology and evolution of antimicrobial resistance in

bacterial communities. ISME J. 2021;15(4):939–48.

65.   Klümper U, Recker M, Zhang L, Yin X, Zhang T, Buckling A, et al. Selection for

592   antimicrobial resistance is reduced when embedded in a natural microbial community.

593   ISME J. 2019;13(12):2927–37.

594   66.   Turner CB, Marshall CW, Cooper VS. Parallel genetic adaptation across environments

595   differing in mode of growth or resource availability. Evol Lett. 2018 Aug 1;2(4):355–67.

596   67.   Venkataram S, Kuo HY, Hom EFY, Kryazhimskiy S. Mutualism-enhancing mutations

597   dominate early adaptation in a two-species microbial community. Nat Ecol Evol. 2023 Jan

598   2;7(1):143–54.

599   68.   Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. Tests of parallel molecular

600   evolution in a long-term experiment with *Escherichia coli*. Proc Natl Acad Sci U S A.

601   2006;103(24):9107–12.

602   69.   Mira PM, Meza JC, Nandipati A, Barlow M. Adaptive Landscapes of Resistance Genes

603   Change as Antibiotic Concentrations Change. Mol Biol Evol. 2015 Oct;32(10):2707–15.

604   70.   Das SG, Direito SO, Waclaw B, Allen RJ, Krug J. Predictable properties of fitness

605   landscapes induced by adaptational tradeoffs. Elife. 2020 May 19;9.

606   71.   Hall BG. Predicting the evolution of antibiotic resistance genes. Nat Rev Microbiol. 2004

607   May;2(5):430–5.

608   72.   Palmer AC, Toprak E, Baym M, Kim S, Veres A, Bershtein S, et al. Delayed commitment

609   to evolutionary fate in antibiotic resistance fitness landscapes. Nat Commun. 2015 Jun

610   10;6(1):7385.

611   73.   Ogbunugafor CB, Wylie CS, Diakite I, Weinreich DM, Hartl DL. Adaptive Landscape by

612   Environment Interactions Dictate Evolutionary Dynamics in Models of Drug Resistance.

613   PLOS Comput Biol. 2016 Jan 25;12(1):e1004710.

614   74.   Salverda MLM, Koomen J, Koopmanschap B, Zwart MP, De Visser JAGM. Adaptive

615   benefits from small mutation supplies in an antibiotic resistance enzyme. Proc Natl Acad

616   Sci U S A . 2017;114(48):12773–8.

617   75.   Brook I. Inoculum Effect. Rev Infect Dis. 1989;11(3):361–8.

618   76.   Organization WH. Antimicrobial resistance, Fact sheet N°194,. 2013.

619   77.   Rolff J, Bonhoeffer S, Kloft C, Leistner R, Regoes R, Hochberg ME. Forecasting

620   antimicrobial resistance evolution. Trends Microbiol. S0966-842X (23) 00361-X.

 Figure legends

Figure 1 | Evolutionary trajectories of uropathogenic bacteria in conditioned media. A, The
phenotypic trajectories of *E. coli* evolution in the presence of sulfamethoxazole-trimethoprim, with
ratio sulfamethoxazole:trimethoprim 5:1. *E. coli* was evolved in the presence of the conditioned
media of *K. pneumoniae, E. coli, E. faecium* and *S. haemolyticus*. The five replicates evolved along
similar trajectories, but are more divergent than those of *K. pneumoniae* and *E. faecium*. **B,** The
phenotypic trajectories of *K. pneumoniae* evolution in the presence of sulfamethoxazole-
trimethoprim. Ratio sulfamethoxazole:trimethoprim is 5:1. *K. pneumoniae* was evolved in the
presence of the conditioned media of *K. pneumoniae, E. coli, E. faecium* and *S. haemolyticus*. Five
replicates were evolved in each conditioned medium and all replicate trajectories overlapped. **C,**
The phenotypic trajectories of *E. faecium* evolution in the presence of vancomycin. *E. faecium* was
evolved in the presence of the same conditioned media as *K. pneumoniae* and *E. coli*, and also in
the conditioned medium of *P. aeruginosa*. The trajectories of all five replicates per conditioned
medium overlapped.

Figure 2 | Correlation of antibiotic concentration at the final transfer with ecological
interactions; WT population size in conditioned medium, and the fold change of antibiotic
tolerance in conditioned medium. Colors indicate the conditioned medium in which the focal
species were cultured, with blue indicating *K. pneumoniae* conditioned medium, red *E. coli*
conditioned medium, yellow *E. faecium* conditioned medium and green *S. haemolyticus*
conditioned medium. The *E. faecium* mutants were also evolved in *P. aeruginosa* conditioned
medium, represented by orange markers. **A**, Trimethoprim concentration (ratio
sulfamethoxazole:trimethoprim is 5:1) at final transfer of *E. coli* lineages in the serial propagation
experiment correlated with the population size interactions of the wild type in the absence of
antibiotics (Pearson correlation coefficient of 0.8556, N = 20, p < 0.001, means ± s.e.m.). **B**,
Trimethoprim concentration (ratio sulfamethoxazole:trimethoprim is 5:1) at final transfer of *K.*
pneumoniae lineages in the serial propagation experiment correlated with the population size
interactions of the wild type in the absence of antibiotics (Pearson correlation coefficient of 0.4509,
N = 20, p = 0.017, means ± s.e.m.). **C**, Vancomycin concentration at final transfer of *E. faecium*
lineages in the serial propagation experiment correlated with the population size of the wild type
in the absence of antibiotics (Pearson correlation coefficient of 0.9259, N = 25, p < 0.001,
means ± s.e.m.). **D**, Correlation between antibiotic concentration at final transfer of *E. coli* mutants
of in the serial propagation experiment versus the fold change of the antibiotic tolerance of the
wild-type *E. coli* in the conditioned media compared to artificial urine medium (Pearson correlation
coefficient of 0.4986, N = 20, p < 0.001). **E**, Correlation between antibiotic concentration at final
transfer of *K. pneumoniae* mutants of in the serial propagation experiment versus the fold change
of the antibiotic tolerance of the wild-type *K. pneumoniae* in the conditioned media compared to
artificial urine medium (Pearson correlation coefficient of 0.6757, N = 20, p < 0.001). **F**, Correlation

between antibiotic concentration at final transfer of *E. faecium* mutants of in the serial propagation
experiment versus the fold change of the antibiotic tolerance of the wild-type *E. faecium* in the
conditioned media compared to artificial urine medium (Pearson correlation coefficient of 0.6418,
N = 25, p < 0.001).

Figure 3 | Mutation supply, number of SNPs and type of mutations in final propagation step
mutants. Colors indicate the conditioned medium in which the focal species were cultured, with
blue showing mutants evolved in *K. pneumoniae* conditioned medium, red showing mutants
evolved in *E. coli* conditioned medium, yellow showing mutants evolved in *E. faecium* conditioned
medium, and green showing mutants evolved in *S. haemolyticus* conditioned medium. For *E.*
faecium mutants were also evolved in *P. aeruginosa* conditioned medium, which is shown with the
orange marker. The error bars show the standard error of the mean for the number of SNPs in the
non-mutator trains. **A,** The average number of single nucleotide polymorphisms (SNPs) in the
clones of evolved *E. coli* in the conditioned media correlated with the population size of the wild
type in those conditioned media (Pearson correlation coefficient = 0.1792, N = 12, p = 0.17,
means ± s.e.m.). **B,** The average number of SNPs in the clones of *K. pneumoniae* in the conditioned
media correlated with the population size of the wild type in those conditioned media (Pearson
correlation coefficient = 0.7045, N = 12, p < 0.001, means ± s.e.m.). **C,** The average number of SNPs
in the clones of evolved *E. faecium* in the conditioned media correlated with the population size
of the wild type in those conditioned media (Pearson correlation coefficient =0.7616, N = 16, p =
0.1989, means ± s.e.m.). **D,** All mutations in the *E. coli* mutants evolved in the conditioned media
shown on the x-axis per replicate lineage. Two mutator strains were identified, shown here by the
broken y-axis/bars. **E,** All mutations in the *K. pneumoniae* mutants evolved in the conditioned
media shown on the x-axis per replicate lineage. One mutator strain was identified, shown here by
the broken y-axis/bars. **F,** All mutations in the *E. faecium* mutants evolved in the conditioned media
shown on the x-axis per replicate lineage.

Figure 4 | Clustering and mutation repeatability of the final propagation step mutants
clustered on COG-level. Higher values indicate a higher parallelism of the two genotypes
compared. Three mutant lineages were sequenced per conditioned medium (four for *E. faecium*
mutants evolved in *P. aeruginosa* conditioned medium). Comparisons in the blocks around the
diagonal in each figure show the intra-conditioned medium comparisons, while the other blocks
show the inter-conditioned media comparisons. **A,** *H*-index of mutation repeatability at COG-level
of *E. coli* evolved in the four conditioned media. Mutator lineages are not taken into account
(Figure 3D). **B,** *H*-index of mutation repeatability at COG-level of *K. pneumoniae* evolved in the
four conditioned media. Mutator lineages are not taken into account (Figure 3F). **C,** *H*-index of
mutation repeatability at COG-level of *E. faecium* evolved in the five conditioned media.

Figure 5 | Evolved genetic targets separated by type of mutation. Genetic targets shown were
hit multiple times (at least twice) in either replicate populations or environment, or both. Size of
the spheres indicate the number of unique mutations in the specific gene that were observed.
Genes are separated into their respective COG functional group as shown on the right side of the
figures, with each letter corresponding to a different COG-group. COG-groups: C – Energy
production and conversion, E – Amino acid transport and metabolism, F – Nucleotide transport
and metabolism, G – Carbohydrate transport and metabolism, H – Coenzyme transport and
metabolism, K – Transcription, L – Replication, recombination and repair, M – Cell
wall/membrane/envelope biogenesis, O – Posttranslational modification, protein turnover,
chaperones, P – Inorganic ion transport and metabolism, Q – Secondary metabolites biosynthesis,
transport and catabolism, R – General function prediction only, T Signal transduction mechanisms,
and V – Defense mechanisms. **A,** Genetic targets in *E. coli* evolved in four different conditioned
media. See Supplementary Table 2 for a list of the genes and proteins. **B,** Genetic targets in *K.*
pneumoniae evolved in four different conditioned media. See Supplementary Table 3 for a list of
the genes and proteins. **C,** Genetic targets in *E. faecium* evolved in five different conditioned media.
See Supplementary Table 4 for a list of the genes and proteins.

Gram-negative pathogens

Evolved in — *E. coli*, *K. pneumoniae*

Spent media: *E. coli*, *S. haemolyticus*, *K. pneumoniae*, *E. faecium*

+ Antibiotics — Trimethoprim - Sulfamethoxazole (1:5) Starting at 50% of MIC

Serial Transfer — Every 48 hours If OD600>0.1, antibiotic concentration increased 20%

Determine evolutionary trajectory — [antibiotic] vs Time

Gram-positive pathogen

Evolved in — *E. faecium*

Spent media: *E. coli*, *S. haemolyticus*, *K. pneumoniae*, *P. aeruginosa*, *E. faecium*

+ Antibiotic — Vancomycin Starting at 50% of MIC

Serial Transfer — Every 48 hours If OD600>0.05, antibiotic concentration increased 20%

Determine evolutionary trajectory — [antibiotic] vs Time