Location of NMDARs and co-agonist control the generation of

bursts in nigral dopamine neurons

Short title: Extrasynaptic NMDARs in nigral dopamine neurons

Sofian Ringlet[1,3], Zoraide Motta[2], Laura Vandries[3], Vincent Seutin[3], Kevin Jehasse[4],

Laura Caldinelli[2], Loredano Pollegioni[2], and Dominique Engel[1,*]

[1]GIGA-Neurosciences, Laboratory of Molecular Regulation of Neurogenesis,

University of Liege, Avenue Hippocrate 15, B-4000 Liege, Belgium

[2]The Protein Factory 2.0 Lab, Dipartimento di Biotecnologie e Scienze della Vita,

Università degli Studi dell'Insubria, via Dunant, 3 - 21100 - Varese (VA)

[3]GIGA-Neurosciences, Neurophysiology group, University of Liege, Avenue

Hippocrate 15, B-4000 Liege, Belgium

[4]Montefiore Institute of Electrical Engineering and Computer Science, Systems and

Modeling research unit at University of Liège, Quartier Polytech 1, allée de la

Découverte 10, 4000 Liège 1, Belgium

[*] corresponding author (dominique.engel@uliege.be)

Abstract

NMDA receptor activation in pars compacta substantia nigra dopamine neurons is

central to the generation of bursting activity, a key signal temporally associated to

movement initiation. The site of the NMDAR pool (synaptic and/or extrasynaptic) as

well as the identity of the co-agonist involved in the ignition of this phasic activity

remains unknown. Using *ex vivo* electrophysiological recordings, we demonstrate

that NMDARs located outside synapses are preponderant for this firing. This pool of

receptors is recruited during intense synaptic activity via spillover of glutamate and

require the binding of NMDAR co-agonist glycine for their full activation. Synaptic

NMDARs are not directly involved in bursting and are activated by D-serine, a distinct

co-agonist. Location dependency of NMDARs and co-agonist underlying burst

generation may serve as a guideline in understanding the physiological role of

dopamine neurons in health and disease.

Teaser

Extrasynaptic NMDARs recruited by spillover and glycine allow the generation of

bursts in dopamine neurons.

INTRODUCTION

Activity of dopamine (DA) neurons in the substantia nigra pars compacta (SNc) provides essential signal to the dorsal striatum for the control of movement (*1*, *2*). The fragility of DA neurons located specifically in this brain region is at the origin of the severe motor deficits expressed in Parkinson's disease (PD) (*3*). The fundamental function ensured by SNc DA neurons is encoded by transitions of their output signal from spontaneous low-frequency action potential (AP) firing to high-frequency bursts of APs. Bursting is a key electrophysiological signal producing an increase of DA release in postsynaptic target areas (*4*, *5*), which is temporally correlated to locomotion initiation (*1*, *6*). Induction of burst firing is operated by N-methyl-D-aspartate receptors (NMDARs) activated by glutamatergic afferents (*7*) originating from several brain areas but predominantly from the subthalamic nucleus (STN), pedunculopontine nucleus (PPN) and cortex (*8*). A substantial fraction of NMDARs in DA neurons are composed of GluN2B and GluN2D subunits (*9*) as in subthalamic (*10*) and hippocampal interneurons (*11*) generating slow deactivating excitatory postsynaptic currents (EPSCs). These subunits are also commonly expressed at extrasynaptic sites (*12–15*) and form triheteromeric GluN1/GluN2B/GluN2D receptors as in SNc DA neurons at both synaptic and extrasynaptic sites (*16*). While bursts critically depend on the presence of the GluN1 subunit (*17*, *18*), the link between the activation of NMDARs and the generation of a burst is not well understood. This lack of information is surprising since bursts mediate the increased dopamine release which is correlated to movement initiation, a sequence of events that is malfunctioning in the context of PD. Bursting activity is observed in vivo (*19*) but also in *ex vivo* tissue by exogenous applications of NMDA in combination to the SK

channel blocker apamin (*20–22*), by glutamate applications or synaptic stimulation (*23*).

In addition to glutamate, a co-agonist such as glycine (*24*) or D-serine (*25*) is an absolute requirement for NMDAR channel opening (*26*). While both amino acids have been shown to serve as endogenous agonists at the NMDAR glycine site, the contribution of the one or the other is dependent on neuronal development (*27*), expression of specific GluN2 subunits (*27*, *28*) and level of synaptic activity (*29*). D-serine is generally associated to the synaptic site where GluN2A-containing NMDARs are preferentially expressed, and glycine to the extrasynaptic site where GluN2B-containing NMDARs are present (*28*). The source of D-serine and glycine could potentially be neuronal (*30*, *31*) and astroglial, respectively, but this is still under debate. The amount of co-agonists in and around the synaptic cleft is tightly controlled by their corresponding synthesis and degrading enzymes and by their specific transporters (*32*). In SNc DA neurons, both EPCSs and bursts can be potentiated by exogenous co-agonists or by the blockade of glycine uptake (*21*) indicating that the co-agonist site of NMDARs is not saturated. In addition, blockade of the co-agonist binding-site on NMDARs abolished bursts, revealing the importance of the co-agonist in this firing pattern (*21*). However, the identity of the endogenous co-agonist as well as the population of NMDARs involved in burst generation remain unknown.

In this study, we determined the location of the NMDAR pool and the identity of the co-agonist implicated in the generation of bursting activity in SNc DA neurons. We found that distinct co-agonists activate synaptic and extrasynaptic NMDARs, although both of these NMDARs populations are composed of the same set of GluN2 subunits. Interestingly, the pool of NMDARs implicated in bursting and the pool of

82　extrasynaptic NMDARs require the same co-agonist, indicating that both pools of

83　NMDARs are identical. These findings shed light on the importance of the location of

84　both the NMDAR pool and the co-agonist implicated in the generation of bursts in

85　SNc DA neurons.

RESULTS

The co-agonist glycine activates NMDARs to regulate bursting activity

Bursting activity in SNc DA neurons relies on the tonic activation of NMDARs (*33*) through the binding of the neurotransmitter glutamate. In addition to this main agonist, a co-agonist such as glycine or D-serine participates to the activation of NMDARs as the blockade of the co-agonist site abolishes bursts (*21*). However, which endogenous co-agonist is predominant to activate NMDARs during this phasic activity remains unknown. To identify the co-agonist activating NMDARs implicated in burst generation, enzymatic depletion of the co-agonists was combined to patch-clamp recordings in acute brain slices. DA neurons were identified in current- and voltage-clamp and using post hoc immunohistochemistry for tyrosine hydroxylase (**fig. S1**). Bursting activity was induced by bath application of NMDA and apamin and hyperpolarizing current injection (*21*). In these conditions, 100% of neurons (22 out of 22 neurons) exhibited burst firing. Slices were incubated for 45 minutes with either D-amino acid oxidase (DAAO) (**Fig. 1A**) or glycine oxidase (GO) (**Fig. 1B**) to degrade D-serine or glycine, respectively. While the application of DAAO did not affect the profile of the interspike interval (ISI) distribution, burst duration, number of APs in a burst and burst frequency (**Fig. 1, C to F; Table S1**), the incubation with GO strongly modified the profile of the ISI distribution (**Fig. 1G**) and reduced burst frequency but without changing burst duration and the number of AP/burst (**Fig. 1, H to J; Table S1**). Adding D-serine (100 µM) to DAAO-treated samples modified slightly the ISI distribution profile, increased the depolarizing envelope duration and the number of APs riding on a burst without changing the burst frequency (**Fig. 1, C to F; Table S1).** Supplementing glycine (1 mM) to GO-treated samples modified the ISI

distribution profile, increased the depolarizing envelope duration and the number of

APs riding on a burst but did however not restore the frequency observed in control

(**Fig. 1, G to J; Table S1**). In the absence of the enzymes, the bursting activity

remained stable over 45 minutes (**fig. S2; Table S2**). These results indicate that

glycine regulate the occurrence of bursts via the activation of NMDARs.

To control for changes in the amount of glycine and D-serine in slices in the presence

of either GO or DAAO, concentrations of glycine and D-serine were evaluated using

high-performance liquid chromatography (HPLC) experiments. In slices incubated for

45 min with either DAAO (0.2 U mL^{-1}) or GO (0.2 U mL^{-1}), D-serine and glycine

concentration was significantly decreased (35.00 ± 12.97% of control; 42.40 ±

11.01% of control respectively). These results confirm that DAAO and GO specifically

degrade D-serine and glycine, respectively (**fig. S3**). The specificity of the enzymes

observed here is in line with previous observations in comparable experimental

conditions (*27–29*) and indicate that the reduction of burst frequency is due to a

selective reduction of the respective co-agonist amount.

Synaptic NMDARs are activated by the co-agonist D-serine

NMDARs are implicated in synaptic transmission in SNc DA neurons (*34, 35*) but the

nature of the endogenous co-agonist activating the synaptic pool of NMDARs has not

been identified for these neurons. In addition, whether the identity of the co-agonist

for synaptic NMDARs and for bursting activity is identical is not known. Slices were

treated with either DAAO to degrade D-serine (**Fig. 2A**) or with GO to reduce the

glycine level (**Fig. 2B**) for 45 min. Treatment with DAAO induced a significant

reduction in peak amplitude of the NMDAR-mediated component of sEPSCs at +40

mV. Addition of exogenous D-serine (100 μM) reversed the effects of DAAO on the

amplitude of sEPSCs which was similar to the EPSC amplitude in control (**Fig. 2C**). Treatment with GO did not change the amplitude of sEPSCs but supplementation of glycine to GO-treated samples increased the amplitude of sEPSCs which was significantly larger than in control conditions (**Fig. 2D**). Currents at -70 mV, representing principally the activation of AMPARs, were not modified, either in the presence of DAAO or GO (**Fig. 2, C and D**). These results provide evidence that D-serine, but not glycine, maintains synaptic activation of NMDARs in DA neurons under basal synaptic activity. Unexpectedly, the co-agonist activating synaptic NMDARs and NMDARs for bursting differed in their identity. This indicates that the pool of NMDARs implicated in the generation of bursts might be different from the pool of synaptic NMDARs and that those may not be directly implicated in the generation of bursts.

DA neurons in the SNc receive glutamatergic input principally from the STN, the PPN and motor cortex (*8*). Among these inputs, substantial innervation originates from the STN (*8*). To reveal the identity of the co-agonist in this specific synapse, electrically evoked EPSCs (eEPSCs) were recorded in DA neurons during stimulation of STN excitatory afferent fibers. Synaptic transmission originating from the STN was identified using paired-pulse ratio (see Materials and methods and **fig. S4**) and the stimulation intensity was adjusted to slightly above minimal stimulation to activate the smallest number of fibers. Incubation with DAAO reduced significantly the peak amplitude of the NMDAR-mediated component of eEPSCs at +40 mV and addition of exogenous D-serine (100 µM) reversed the effects of DAAO on the amplitude of eEPSCs (**Fig. 3, A and C**). Treatment with GO did not change the mean amplitude of eEPSCs but addition of glycine to GO-treated samples increased the mean amplitude of eEPSCs, which was significantly larger than in control conditions (**Fig.**

3, B and D). Currents at -70 mV were neither changed by the treatment with DAAO,

nor with GO. These results obtained for the STN-SNc synapse are similar to those

observed on sEPSCs, indicating that the same co-agonist, D-serine, is probably

shared by distinct excitatory inputs in DA neurons. In addition, these results further

reveal that synaptic NMDARs and NMDAR-mediating bursts are activated by distinct

co-agonists.

Synaptic NMDARs in nigral DA neurons are triheteromers composed of GluN2B

and GluN2D subunits

A strong association between the nature of the predominant co-agonist in excitatory

synapses and the expression of a specific GluN2 subunit has been previously

established (*27, 28*). D-serine is usually the main co-agonist found to activate

synaptic GluN2A-containing NMDARs. In juvenile DA neurons, a high proportion of

synaptic NMDARs are GluN1/GluN2B/GluN2D triheteromers and the presence of the

GluN2A subunit has not been detected (*9*). Since D-serine is often the co-agonist of

GluN2A-containing NMDARs, the subunit composition of NMDARs in DA neurons

has been reexamined in adolescent animals in which a potential switch from GluN2B

to GluN2A subunits occurring during the second postnatal week (*36*) already

happened. Ambient zinc (Zn^{2+}) levels exert a tonic inhibition of EPSCs mediated by

GluN2A-containing NMDARs due to a high-affinity binding site for Zn^{2+} specifically on

this subunit (*37*). To reveal the functional contribution of the GluN2A subunit, we

bath-applied the Zn^{2+} chelator tricine (10 mM) to buffer Zn^{2+} (*38*). This treatment did

not modify the amplitude of the NMDAR component of eEPSCs at +40 mV (**Fig. 4, A**

and B). The subsequent addition of Zn^{2+} (60 µM, corresponding to 300 nM free Zn^{2+})

was also unable to modify the amplitude of eEPSCs. The absence of effect of tricine

was also observed for applications of longer durations (**fig. S5**). In contrast,

application of the antagonist of NMDARs containing GluN2B (Ro25-6981, 1 µM)

reduced significantly the amplitude of eEPSCs to ~60% of control (**Fig. 4, C and D**).

Similarly application of the GluN2D-containing NMDAR antagonist DQP-1105 (20

µM) reduced the amplitude of sEPSCs to ~53% of control (**Fig. 4, E and F**).

Weighted decay time constant of EPSCs was not modified by the application of either

Ro25-6981 or DQP-1105. The absence of effect on the decay time constant by either

Ro25-6981 or DQP-1105 suggests that synaptic NMDARs are essentially composed

of a high proportion of GluN1/GluN2B/GluN2D triheteromers making the participation

of diheteromeric GluN1/GluN2B and GluN1/GluN2D receptors unlikely (*11*). Taken

together, these results indicate that a large fraction of synaptic NMDARs may be

GluN1/GluN2B/GluN2D triheteromers in adolescent SNc DA neurons as in juvenile

tissue (*9*) and seem to exclude a switch from Glu2B to GluN2A subunits in DA

neurons. In excitatory synapses of DA neurons, D-serine activates triheteromeric

NMDARs containing GluN2B and GluN2D but not GluN2A.

The expression of NMDAR subunits in DA neurons was further explored using

immunohistochemistry for GluN2A, GluN2B and GluN2D subunits. The distribution of

the subunits was determined in TH$^+$ DA neurons in the region in which DA neurons

were selected for patch-clamp recordings (**fig. S6**). Both GluN2B and GluN2D

subunits were present in a large majority (~80%) of TH$^+$ neurons but the GluN2A

subunit was only present in ~20% of TH$^+$ neurons (**fig. S6, B and C**). Within TH$^+$

neurons, there were ~3 times more fluorescent dots for GluN2B and GluN2D

subunits than for Glun2A subunits (**fig. S6D**). The distribution of the three distinct

NMDARs subunits was further examined at higher magnification at the cellular level

by   taking   into   account   membrane   and   intracellular   locations   (**fig.   S7**).

Immunolabeling for GluN2B and GluN2D subunits was found predominantly at the

membrane (GluN2B membrane, ~75% of the total dots; GluN2D membrane, ~77% of

the total dots) in comparison to the intracellular compartment (GluN2B intracellular,

~25% of the total dots; GluN2D intracellular, ~23%). In contrast, the distribution of

GluN2A was inverted and found at a high proportion in cell's intracellular milieu

(GluN2A intracellular, ~85%) and at a very low proportion at the membrane (GluN2A

membrane, ~15%) (**fig. S7, A and B**). This GluN2 subunits expression profile is

consistent with the pharmacological data obtained using subunit specific NMDAR

blockers (**Fig. 4**).

Glycine controls the activation of the NMDAR-mediated tonic current

NMDARs are generally concentrated in synapses but are also encountered at

extrasynaptic sites (*39*). Activation of extrasynaptic NMDARs by ambient glutamate

generates a whole-cell tonic current manifested by the application of the NMDAR

blocker D-AP5 (*40*). An NMDAR-mediated tonic current has also been described in

SNc DA neurons (*41*) but the identity of the co-agonist activating extrasynaptic

NMDARs in SNc DA neurons remains undetermined. At +40 mV, the shift in current

produced by D-AP5 had an averaged amplitude of ~30 pA which is comparable in

amplitude to previous reports for the same neurons (*41*) (**Fig. 5, A and B**). In slices

incubated with DAAO, the amplitude of the tonic current was not significantly different

from control after application of D-AP5. In contrast, in the presence of GO, the

amplitude of the tonic current was reduced to ~46% of control, indicating that

reducing the amount of glycine impaired the activation of NMDARs mediating this

current. Consistent with previous observations made in other neuronal types (*28*),

these results support the hypothesis that glycine is the predominant co-agonist acting

at extrasynaptic NMDARs. Interestingly, since glycine is the predominant co-agonist

for extrasynaptic NMDARs activated during tonic current and is also controlling the

occurrence of bursts, extrasynaptic NMDARs might be critical for the generation of

bursts.

NMDAR subunit composition of NMDAR-mediated tonic current

Since glycine is the co-agonist for the activation of extrasynaptic NMDARs, the

GluN2B subunit is expected to be a component of NMDARs at this location (*28*).

While NMDAR subtypes at synaptic sites generally contain GluN2A subunits , those

at extrasynaptic sites contain GluN2B (*42*, *43*) and GluN2D subunits (*12*, *13*). In

adolescent SNc DA neurons the subunit composition of extrasynaptic NMDARs is not

known. A large proportion could be GluN1/GluN2B/GluN2D triheteromers as

suggested for juvenile DA neurons using outside-out patch recordings (*16*). To

determine the subunit composition of extrasynaptic NMDARs in DA neurons, subunit

specific NMDAR blockers were used. In the presence of tricine, the magnitude of the

tonic current was similar as in control conditions (**Fig. 5, B and C**). Adding Zn^{2+} (60

nM) to tricine did not modify the current amplitude too, indicating that the GluN2A

subunit may not compose extrasynaptic NMDARs. The GluN2B blocker Ro25-6981

and the GluN2D blocker DQP-1105 reduced significantly the tonic current to ~31% of

control and ~26% of control, respectively (**Fig. 5, B and C**). These results indicate

that extrasynaptic NMDARs may be composed of GluN2B and GluN2D subunits, as

observed for juvenile tissue (*16*) and that extrasynaptic, like synaptic sites, may

express a high proportion of the same GluN1/GluN2B/GluN2D NMDAR subtype (*9*).

In addition, these observations confirm that GluN2B is not enriched in the

extrasynaptic membrane and is present both at synaptic and extrasynaptic sites (*43*)

at least in SNc DA neurons. Taken together, our results show that despite similar

NMDAR subtypes at synaptic and extrasynaptic sites, distinct co-agonists activate

NMDARs at synaptic and extrasynaptic sites. The preference of D-serine to act at

synaptic NMDARs and glycine at extrasynaptic NMDARs is therefore more probably

due to their availability rather than to the expression of a specific GluN2 subunit.

Activation of extrasynaptic NMDARs generates tonic current

It is generally accepted that the tonic NMDAR-mediated current is largely due to the

activation of extrasynaptic NMDARs (*40*). However, tonically activated synaptic

NMDARs could also contribute to the tonic current. To evaluate the potential

contribution of synaptic NMDARs to the tonic current, we used the irreversible open

channel blocker MK-801 to block this pool of receptors (*44–46*). After a 20 min

application of MK-801, low frequency stimulation elicited eEPSCs at +40 mV with

strongly reduced amplitude to ~30% of control due to the blockade of synaptic

NMDARs (**Fig. 6A**). The consecutive application of D-AP5 (50 µM) induced a tonic

current with an amplitude of ~22 pA that was similar to the amplitude measured

without the suppression of synaptic NMDARs (**Fig. 6, A and C**). In contrast, the

blockade of both synaptic and extrasynaptic NMDARs using low frequency repetitive

afferent fiber stimulation under MK801 incubation at +40 mV reduced strongly the

amplitude of the tonic current (**Fig. 6, B and C**). The absence of a change in the tonic

current when synaptic NMDARs are blocked indicates that the pool of extrasynaptic

NMDARs sustains the tonic NMDAR-mediated current in DA neurons.

Bursting activity relies on the contribution of extrasynaptic NMDARs

In vivo, bursting activity in SNc DA neurons results from the tonic activation of

NMDARs (*33*). In vitro, spontaneous bursts can be generated by exogenous

applications of NMDA and the SK channel blocker apamin but whether synaptic or

extrasynaptic NMDARs contribute to this firing activity is not known. To determine the

possible participation of synaptic NMDARs to the bursting activity, MK-801 (20 µM)

was applied to specifically block this pool of receptors. As shown in the previous

recordings, the irreversible open-channel inhibitor reduced strongly the amplitude of

eEPSCs at +40 mV to ~26% of control, indicating that a large fraction of synaptic

NMDARs were blocked (**Fig. 7A**). The recording configuration was then switched

from voltage-clamp at +40 mV to current-clamp ($V_{membrane}$ ~ -80 mV) to determine

whether the spontaneous bursting activity can be induced. After a short time period

(~10 min) to allow stabilization of the membrane potential, NMDA and apamin were

bath-applied in combination with small DC current injections. In these conditions, DA

neurons exhibited spontaneous bursting activity, similar to the activity recorded

without the blockade of synaptic NMDARs (**Fig. 1, A and 1B and fig. S2A**). These

recordings were performed using a K^+-based pipette internal solution and TEA and 4-

AP in aCSF to block a majority of the potassium conductances in order to stabilize

the current baseline at +40 mV in voltage-clamp. TEA and 4-AP were only present in

the bath for voltage-clamp recordings and were washed out while switching to

current-clamp. In these conditions, bursting could be generated in a majority of DA

neurons (**Fig. 7A**). Bursts generated in these conditions were similar to those in

control conditions with comparable burst duration, number of AP/burst and frequency

(**Fig. 7, B to E**). These results strongly suggest that bursts in DA neurons are elicited

mainly by the activation of extrasynaptic NMDARs.

Synaptic glutamate from STN axons allows burst generation via extrasynaptic

NMDARs

Glutamate released during synaptic transmission activates postsynaptic NMDARs in DA neurons to allow the generation of bursts (*23*, *47*). In vivo, the stimulation of the glutamatergic STN neurons favors bursting (*7*). To evaluate the contribution of glutamate released by STN neurons to bursting, fibers originating from this nucleus were stimulated using a focal theta glass electrode and voltage recorded in DA neurons in current-clamp configuration while AMPARs were blocked. Fibers from the STN were identified using paired-pulse facilitation (*48*). In control conditions a train of stimuli at low frequency (10 Hz) for 500 ms did not evoke associated APs (**Fig. 8, A and C to E**). At this stimulation frequency, extrasynaptic NMDARs are not recruited by spillover (*49*). Incubation with either DAAO or GO to reduce D-serine or glycine, respectively, did also not modify the firing. Addition of glycine (1 mM) to GO was not able to change the firing too. In contrast, stimulation at a higher frequency (50 Hz) for 500 ms induced in control conditions and in the presence of DAAO short bursts of APs (**Fig. 8B**) which were similar in both conditions. In the presence of GO, bursts had a reduced number of AP, a decreased intraburst frequency but an increased time to first AP. However, supplementation of glycine to GO induced burst at 50 Hz, increased the AP number, intraburst frequency and decreased time to first AP in comparison to the condition in which GO is alone (**Fig. 8, B and C to E**). At this stimulation frequency (50 Hz), extrasynaptic NMDARs are recruited by spillover of glutamate (*49*) and co-activated by the presence of glycine. These results obtained with synaptically generated bursting corroborate those with spontaneous bursting and further confirm the contribution of extrasynaptic NMDARs and glycine to the generation of bursts.

DISCUSSION

The contribution of NMDARs located at synaptic or extrasynaptic sites and the availability of co-agonists within and outside synapses strongly influence the excitability of DA neurons and therefore the downstream release of dopamine in the striatum. Here, by using electrophysiological and biochemical experiments, we determined the nature of the co-agonist and the pool of NMDARs implicated in bursts generated by SNc DA neurons. First, by combining enzymatic depletion for distinct co-agonists and whole-cell recordings, we identified the co-agonist controlling the initiation of bursts. Second, we uncovered that the co-agonists acting during glutamatergic synaptic transmission and bursting are distinct. Third, we found that the identity of the co-agonist implicated in the activation of extrasynaptic NMDARs and in bursting are identical. Fourth, we showed that the tonic NMDAR-mediated current is predominantly mediated by extrasynaptic NMDARs. Fifth, bursting activity is supported principally by the contribution of extrasynaptic but not synaptic NMDARs. Sixth, immunohistochemical labeling is consistent with electrophysiological recordings and confirms the presence of GluN2B and GluN2D but not GluN2A subunits at the membrane of DA neurons, indicating that the same NMDAR subtype is expressed both at synaptic and extrasynaptic sites. The same NMDAR subunits GluN2B and GluN2D at both locations might therefore invalidate an influence of a co-agonist on the expression of a specific GluN2 subunit or the latter on the distribution of a co-agonist at a given location. Finally, we concluded that bursting in DA neurons is mediated by the activation of glycine-gated extrasynaptic NMDARs.

Extrasynaptic NMDAR population mediate bursting in SNc DA neurons

NMDARs located at extrasynaptic versus synaptic sites have been shown to have

different subunit compositions and to deserve distinct cellular functions (*50*). In

adolescent SNc DA neurons, synaptic NMDARs are composed of GluN2B and

GluN2D subunits, an expression pattern already observed at postnatal ages (*9*).

NMDAR-EPSCs were reduced in amplitude to ~60% of control and ~50% of control

by the GluN2B-containing NMDAR blocker Ro25-6981 and the GluN2D-containing

NMDAR blocker DQP-1105, respectively. These percentages of inhibition are similar

to those reported by previous observations for midbrain neurons (*10*, *51*) and

hippocampal interneurons (*11*) where GluN1/GluN2B/GluN2D triheteromers have

been suggested to be present. A small proportion of diheteromeric GluN1/GluN2B or

GluN1/GluN2D NMDARs might participate to the synaptic current, but the absence of

change of the decay time course of NMDAR-eEPSCs in the presence of either Ro25-

6981 or DQP-1105 argues against the presence of these diheteromeric NMDARs (*9*,

52). Diheteromeric GluN1/GluN2B and GluN1/GluN2D NMDARs produce shorter and

longer deactivation time constants than triheteromeric receptors, respectively (*11*).

Tricine and Zn^{2+} were ineffective to modify NMDAR-EPSC amplitude, indicating the

absence of GluN2A in synaptic receptors. Interestingly, some GluN2A subunits were

present   in   a   small   proportion   of   SNc   DA   neurons   as   revealed   by

immunohistochemical labelling, but these were mainly located in the intracellular

space with only a minor proportion at the cell membrane. At extrasynaptic sites of DA

neurons NMDARs are composed of GluN2B and GluN2D subunits as revealed by

GluN2 specific NMDAR blockers on the tonic current, probably the same NMDAR

subtypes   as   those   at   synaptic   locations.   A   substantial   proportion   of   these

triheteromeric receptors is also present in juvenile tissue (*16*), ruling out a subunit

exchange   until   adolescence.   A   substantial   proportion   of   triheteromeric

GluN1/GluN2B/GluN2D NMDARs may therefore be present at both synaptic and

extrasynaptic sites. The presence of GluN2B and GluN2D subunits is generally

observed at extrasynaptic sites in distinct types of neurons at postnatal or juvenile

ages (*12*, *14*, *15*) but also in mature tissue (*13*, *53*). The presence of GluN2B in

synaptic and extrasynaptic NMDARs has been also observed in the hippocampus

(*43*) suggesting that in this region the subunit present at both sites is mainly in

triheteromeric   receptors   and   associated   to   either   GluN2A   in   synapses

(GluN1/GluN2A/GluN2B) NMDARs (*54*) or GluN2D at extrasynaptic locations

(GluN1/GluN2B/GluN2D) NMDARs.

The same NMDAR subtype is activated by distinct co-agonists depending on

the location

The identity of the co-agonist has been determined for several glutamatergic

synapses in distinct principal neurons and brain regions (*27–29*, *55–57*). D-serine is

generally engaged in the activation of synaptic NMDARs and our results confirms this

rule for SNc DA neurons. In a unique case, however, glycine has been shown to

activate synaptic NMDARs in mature dentate granule cells (*27*). The predominant

role of D-serine for synaptic NMDARs has been associated to the presence the

GluN2A subunit in postsynaptic NMDARs for instance in adolescent cortical layer 5

neurons (*57*) and hippocampal CA1 pyramidal neurons (*27*, *28*). GluN2A is indeed a

component of synaptic NMDARs after the maturational switch from GluN2B to

GluN2A (*36*). In the case of DA neurons, our results show that the preponderance of

D-serine to activate synaptic NMDARs is not accompanied by the postsynaptic

presence of GluN2A but either with the expression of GluN2B and GluN2D in SNc

DA neurons. In addition, the developmental switch from GluN2B subunits to GluN2A

reported for cortical and hippocampal excitatory synapses (Le Bail et al., 2015) does not seem to occur in SNc DA neurons. Enzymatic reduction of glycine revealed that this co-agonist predominates for the activation of extrasynaptic NMDARs in SNc DA neurons. A comparable correspondence has been reported for CA1 pyramidal neurons where glycine activates also preferentially extrasynaptic GluN2B-containing NMDARs (*28*). The expression of a specific NMDAR subtype (containing GluN2B and GluN2D but not GluN2A) at both the synaptic and extrasynaptic sites challenges the hypothesized association between a given co-agonist and a specific GluN2 subunit expressed in postsynaptic NMDARs (*27*, *28*, *58*). Despite the similar NMDAR subunit composition within and outside synapses, D-serine is active in the synaptic cleft and glycine at extrasynaptic sites, suggesting that a direct influence of the identity of the co-agonist on the presence of a specific GluN2 subtype is unlikely. Since both sites have the same NMDAR subtype composed of GluN2B and GluN2D and D-serine and glycine have similar affinities for a given GluN2 subunit (*58–60*), a discrimination between the two co-agonists cannot be operated by NMDARs. Rather, the organization of the microenvironment, the presence of transporters for the co-agonists and the proximity of astrocytic leaflets in the vicinity of NMDARs certainly influences the predominant role of co-agonists at synaptic and extrasynaptic locations. The glycine transporter GLYT1 expressed in the SNc (*61*) might maintain a high concentration ratio of D-serine/glycine within the synaptic cleft and a low concentration ratio at the extrasynaptic membrane (*62*). D-Serine reuptake by ASCT and EAAT transporters is not so efficient since they are not stereoselective and display low to moderate affinity for D-Serine. The mechanisms for the segregation of glycine and D-serine in the extrasynaptic and synaptic compartments, respectively, are not known despite substantial work dedicated to the distinct functions of the

424 tripartite synapse (*63*). More work is needed to shed light on the establishment and

425 role of the compartmentalization of co-agonists. Interestingly, manipulating the

426 availability of the co-agonists might represent a tool to target specifically synaptic

427 versus extrasynaptic NMDAR pools.

428 Contribution of glycine and extrasynaptic NMDARs to bursting activity in SNc

429 DA neurons

Bursting observed in vivo from SNc DA neurons (*19*) can be reproduced in slices by

bath application of NMDA and apamin (*20*). The firing in vivo and in vitro is

spontaneous and show comparable voltage time courses. The pivotal role of

NMDARs in the generation of bursts has been previously revealed in vivo using

pharmacological blockade of NMDARs (*33*) and in vitro using genetic inactivation of

the essential GluN1 subunit specifically in DA neuron (*17*, *18*). Both of these studies

showed that inactivated NMDARs induced a decrease in burst frequency as

observed in our work using depletion of glycine. Moreover, glycine is also

preponderant for the activation of extrasynaptic NMDARs underlying for a large part

the tonic NMDAR-mediated current. These results are in line with work showing that

in vivo bursting is generated by tonically activated NMDARs (*33*). Spontaneous

bursting activity could be induced after the pharmacological blockade of synaptic

NMDARs, further supporting the predominant contribution of extrasynaptic NMDARs

to bursting activity. Altogether, these observations strongly suggest that the pool of

NMDARs implicated in the generation of bursts and the pool of extrasynaptic

NMDARs, both depending on glycine, correspond to the same pool of NMDARs, in

line the observations made in vivo (*33*). Functionally, how can extrasynaptic

NMDARs be activated to generate bursts? This pool of receptors could be recruited

during temporal summation of excitatory synaptic inputs, mediating slow inactivating

NMDAR-EPSCs (*18*) due to their slow kinetics (*64*). Extrasynaptic NMDARs are most

probably recruited during high frequency synaptic activity by glutamate spillover (*15*)

when glutamate uptake is limited or overwhelmed (*41*, *65*). Plateau potentials in layer

5 cortical pyramidal neurons reminiscent of burst in DA neurons are similarly induced

by the activation of extrasynaptic NMDARs via spillover (*66*) and hippocampal

neurons (*67*, *68*). Extrasynaptic GluN2D-containing NMDARs might favor EPSP-spike

coupling, as recently shown for GABAergic neurons (*69*).

Exogenous application of glycine potentiates bursts in SNc DA neurons (*21*) and

enzymatic degradation of the co-agonist decreases their frequency as shown here.

Glycine is predominant for the activation of extrasynaptic NMDARs which support

bursting. These findings represent mechanistic elements that might be considered for

pharmacological treatment of PD. Interestingly, stimulating the activation of NMDARs

via its co-agonist site could be beneficial during the late phase of PD (*70*). In vivo

studies and clinical studies showed indeed that administration of sarcosine, a glycine

transporter 1 inhibitor, ameliorates some symptoms of PD (*71*, *72*). Increasing the

amount of glycine might act at extrasynaptic NMDARs and favor bursting activity by

SNc DA neurons in patients with PD.

MATERIALS AND METHODS

Animals

Experiments on Wistar rats were performed in strict accordance with institutional

(protocole 2203, Commission d'Éthique Animale, Université de Liège), national

(Comité déontologique), and European guidelines for animal experimentation.

Slice preparation

Adolescent 4- to 7-week-old Wistar rats of either sex were anaesthetized with

isoflurane (4% in O_2, Matrx VIP 3000, Midmark) and subsequently killed by

decapitation. Before anesthesia animals were kept in an oxygen-enriched chamber

for fifteen minutes in order to increase cell survival. Brain slices were prepared as

previously described (*21*). Briefly, after decapitation, the brain was rapidly dissected

out and immersed in ice-cold slicing solution containing: 87 mM NaCl, 25 mM

$NaHCO_3$, 2.5 mM KCl, 1.25 mM NaH_2PO_4, 10 mM D-glucose, 75 mM sucrose, 0.5

mM $CaCl_2$, and 7 mM $MgCl_2$ (pH 7.4 in 95% O_2/5% CO_2, ~325 mOsm). Parasagittal

300 μm-thick (for spontaneous EPSC recordings) or horizontal 250 μm-thick slices

(for electrically evoker EPSCS) containing the midbrain were cut using a vibratome

(Leica VT-1200, Nussloch, Germany) with a cutting velocity of 0.06 mm/s and a

horizontal vibration amplitude of 1.0 mm. Slices containing the SNc were transferred

to a storing chamber and allowed to recover at 34°C for 30–45 min. After recovery,

slices were kept at room temperature (20 – 25°C) until recording, for up to 4–5 hours.

Electrophysiology

Slices were transferred to a recording chamber and superfused with aCSF containing

(in mM): 125 NaCl, 25 $NaHCO_3$, 25 glucose, 2.5 KCl, 1.25 NaH_2PO_4, 2 $CaCl_2$, and 1

$MgCl_2$ (equilibrated with 95% O_2/5% CO_2, Osmolarity ~320 mOsm). DA neurons

were visualized using infrared-Dodt gradient contrast (IR-DGC) optics and a 40x

objective (numerical aperture 0.75, Zeiss, Oberkochen, Germany) on a Zeiss Axio

Examiner.A1 microscope equipped with a CCD camera (C7500-51, Hamamatsu,

Japan). The location of the medial terminal (MT) nucleus of the accessory optic tract

was used as a reference for the SNc area in horizontal slices. DA neurons were

identified in whole-cell using current-clamp and voltage-clamp protocols and post hoc

using immunohistochemistry. Selected DA neurons had a voltage rectification with hyperpolarizing current steps, action potentials with long half-width, slow autonomous pacemaking (1-4 Hz) and were positive for tyrosine hydroxylase (**fig. S1**).

Voltage-clamp recordings of DA neurons were conducted with patch pipettes from thick walled borosilicate glass tubing (outer diameter: 2 mm, inner diameter: 1 mm, Hilgenberg, Germany) with a horizontal puller (P-97, Sutter Instruments, Novato, USA). Pipettes were first filled with a potassium gluconate-based internal solution (~1 mm from the tip) and consecutively with a cesium gluconate solution (*73*) for whole-cell voltage-clamp recordings. For current-clamp recordings, pipettes were only filled with a potassium-based solution. The potassium gluconate solution consisted of, in mM: 120 K-gluconate, 20 KCl, 2 $MgCl_2$, 4 Na_2ATP, 0.5 NaGTP, 5 Na_2-phosphocreatine, 0.1 EGTA, 10 HEPES and 0.1% biocytin (pH = 7.3; osmolarity: ~302 mOsm) and the cesium solution: 120 Cs-gluconate, 20 CsCl, 2 $MgCl_2$, 4 Na_2ATP, 0.5 NaGTP, 5 Na_2-phosphocreatine, 0.1 EGTA, 10 HEPES, 3 QX-314 and 0.1% biocytin (pH = 7.3; osmolarity: ~306 mOsm). When filled with these solutions, tip resistances were between 3 and 6 MΩ.

Voltage- and current-clamp recordings were obtained using a Multiclamp 700B amplifier (Molecular devices, Palo Alto, CA) connected to a PC via a Digidata 1550 interface (Molecular devices). Bridge balance was used to compensate the series resistance (Rs) in current-clamp recordings (Rs <15 MΩ). The current-clamp signal was sampled at 10 kHz. Whole-cell recordings were discarded if the access resistance increased above 15 MΩ for voltage-clamp and above 20 MΩ for current-clamp or if the values varied over 30% during recording. All voltage-clamp and current-clamp recordings were performed at near-physiological temperature (32-34°C).

Bursting activity in SNc DA neurons was generated by bath-applications of NMDA

(30 µM) and apamin (300 nM). Hyperpolarizing current injection (from -80 to -140 pA)

was applied to maintain membrane potential in a range (~-60 to -80 mv) that

facilitated the occurrence of bursts. Spontaneous EPSCs were isolated

pharmacologically with picrotoxin (50 µM), CGP 55485 (1 µM) and strychnine (10

µM) to block $GABA_A$, $GABA_B$ and glycine receptor, respectively. Tonic NMDAR-

mediated currents were recorded in the presence of CNQX, picrotoxin, CGP 55485

and strychnine. The amplitude of the tonic current was measured by the baseline

shift after D-AP5 (50 µM) application.

Extracellular synaptic stimulation

Afferent fibers were stimulated electrically using theta-glass (borosilicate; Hilgenberg)

pipettes (*74*) filled with a HEPES-buffered sodium-rich solution containing (in mM):

135 NaCl, 5.4 KCl, 1.25 NaH_2PO_4, 1.8 $CaCl_2$, and 1 $MgCl_2$, 5 HEPES (pH adjusted

to 7.2 with NaOH). The stimulation pipette was placed rostrally close (within ~100

µm) to the recorded DA neuron and short current pulses were applied (200-600 µA,

200-400 ms) to be slightly above minimal stimulation. Excitatory afferent fibers

originating from the STN were selected using paired-pulse ratio. Paired-pulse

facilitation was used to identify fibers from the STN as reported previously (*48*).

Electrically evoked EPSCs were isolated pharmacologically with $GABA_A$, $GABA_B$ and

glycine receptor blockers (picrotoxin, CGP 55485 and strychnine, respectively).

Synaptic NMDARs were selectively blocked by using the irreversible NMDAR

antagonist MK-801 (20 µM) applied for 20 min. Electrically evoked EPSCs (eEPSCs)

were recorded at a holding potential of V_{hold} = +40 mV under 0.1 Hz stimulation until

stable baseline was obtained. Recording configuration was switched to current-clamp

with $V_{membrane}$ ~ -80 mV for 20 min during the bath-application of MK-801 and

stimulation was stopped. MK-801 was washed out and recording configuration was

switched back to voltage-clamp with V_{hold} = +40 mV. Stimulation was resumed and

delivered at 5 Hz for 16 s to block selectively synaptic NMDARs (*45, 46, 75*).

Stimulation <10 Hz does not induce measurable extrasynaptic NMDAR-mediated

current (*49*).

Drugs and reagents

Picrotoxin (50 µM, Alomone Lab), CGP 55845 hydrochloride (1 µM, Hellobio),

strychnine (10 µM, Sigma) and D-AP5 (50 µM, Alomone labs) were dissolved in

doubled distilled water. CNQX disodium (10 µM, Alomone labs) was dissolved in bi-

distilled water. Picrotoxin was dissolved in 90% ethanol. Biocytin and ProLong

Diamond (Life Technologies). Fluorescein Avidin DCS (Labconsult). The remaining

chemicals and all salts were purchased from Sigma-Aldrich (Belgium).

Immunohistochemistry

Immunohistochemical analysis for identification of DA neurons was done on 300 µm

horizontal or parasagittal slices taken after electrophysiological recordings. Biocytin-

filled cells were fixed in 4% paraformaldehyde overnight at 4°C. After washing in

phosphate buffer saline (PBS, 0.1 M) and then in PBS containing 10% normal goat

serum (NGS) and 0.3% triton X-100, slices were incubated with primary antibody

against TH (mouse anti-TH 1:1000, Immunostar 22941, Abcam, UK) in PBS

containing 5% goat-serum and 0.3% Triton X-100 for 22 hours at room temperature.

Subsequently, slices were rinsed with PBS and incubated with the secondary

antibody (goat-anti-mouse-Alexa Fluor 568 1:500, Invitrogen A-11004, Thermo

Fischer Scientitfic, Waltham, MA, USA) together with Fluorescein Avidin DCS (1

μl/ml, Vector Laboratories A-2001-5,) in PBS and 0.3% triton X-100 overnight at 4°C.

After the final rinsing, slices were mounted on a slide containing a 300 μm iSpacer

(SunjinLab #IS307, Hsinchu City, Taiwan) with ProLong Gold Antifade (Invitrogen

P36934), and imaged using an FV1000 confocal microscope (Olympus, Tokyo,

Japan). Images were analyzed with Fiji/ImageJ software (https://imagej.net) and

Imaris v9.5.1 (Oxford Instruments, Santa Barbara, CA, USA).

To determine the expression of GluN2 subunits in the SNc immunohistochemical

analysis was done on 5- to 6-week-old Wistar rats of either sex. Animals were

perfused using a peristaltic pump (Watson Marlow, 101U) with PBS (150 mL),

followed by PBS containing with 4% PFA (100 mL). Brains were carefully collected,

post-fixed in 4% PFA for 24 hours at 4 °C, subsequently stored in sucrose azide

0.01% for 48 hours at 4°C and sliced (40 μm) with a NX70 cryostat (Epredia Belgium,

Machelen, Belgium). Slices were exposed to 10% NGS and 0.1% Triton X-100

followed by primary antibody incubations in PBS containing 5% NGS and 0.1% Triton

X-100 overnight at 4 °C. Primary antibodies were against GluN2A subunit (rabbit

anti-GluN2A 1:200, Alomone AGC-002) combined to mouse anti-TH (1:1000,

Immunostar 22941, Abcam, UK), or GluN2B subunit (rabbit anti-GluN2B 1:200,

Alomone AGC-003) with mouse anti-TH or GluN2D subunit (rabbit anti-GluN2D

1:200, Alomone AGC-020) with mouse anti-TH in 5% NGS in PBS and 0.1% Triton

X-100 overnight at 4 °C. After rinsing, slices were incubated with secondary

antibodies (goat anti-mouse alexa Fluor 568 1:500, Invitrogen A-11004 and goat anti-

rabbit Alexa Fluor 488 1:400, Invitrogen A-11011) in PBS with 0.1% Triton X-100 for

2 hours at room temperature. Slices were rinsed with PBS and shortly exposed to

Dapi (0.5 μg/mL). After mounting with Aqua-poly/mount (Polysciences 18606,

Warrington, PA, USA) images were obtained with an FV1000 confocal microscope

(Olympus, Tokyo, Japan) using 5x, 40x and 60x oil immersion objectives (N.A. 0.16,

0.8 and 1.4, respectively).

The distribution of the NMDAR subunits within neurons and at their membranes was

analyzed with Imaris using movies (**Movie S1 to S6**). Membrane transparency of

reconstructed neurons was adjusted by varying the transparency setting

implemented in Imaris to differentiate NMDAR subunits expressed in the cytoplasm

(100% transparency - 0% opacity) from NMDAR subunits in the membrane (0%

transparency - 100% opacity). TH+ DA neurons were selected around the MT

randomly from both hemispheres.

Enzymes

Recombinant wild-type and M213R variants of *Rhodotorula gracilis* D-amino acid

oxidase (RgDAAO; EC 1.4.3.3) and recombinant H244K glycine oxidase from

Bacillus subtilis (BsGO; EC 1.4.3.19) were produced in *E. coli* cells and purified as

previously described (*76–78*). The RgDAAO and BsGO preparations had a specific

activity of 75 U/mg protein on D-serine as substrate and 2.5 U/mg protein on glycine

as substrate, respectively. Midbrain slices were perfused with RgDAAO (0.2 U/mL) or

BsGO (0.2 U/mL) for 45 min. For details about RgDAAO and BsGO substrate

specificity see (*77, 79*).

High Performance Liquid Chromatography (HPLC) analysis

Slices containing the substantia nigra were stored in the reserve chamber near-

physiological temperature (32-34 °C) using three experimental conditions. The first

condition was in physiological saline (aCSF), the second in aCSF + GO (0.2 U mL^{-1})

and the third in aCSF + DAAO (0.2 U mL^{-1}) for 45 minutes. The aCSF collected

during all experimental conditions were freeze-dried and then resuspended in 25 mM

HCl. Following a neutralization step adding 0.2 M NaOH, the samples were

precolumn derivatizated with o-phthaldialdehyde/N-acetyl-L-cysteine in a 1:4 volume

ratio of 0.2 M borate, pH 8.2, and 0.2 M sodium phosphate, pH 10. Diastereoisomer

derivatives were resolved on a Kinetex EVO C18 reverse-phase column (5 μm, 4.6 x

150 mm; Phenomenex, Bologna, MI, Italy) using a HPLC PU-2089 System (Jasco

Europe, Cremella, LC, Italy) equipped with a fluorescence detector (344/443 nm,

gain 1000X), in isocratic conditions (0.1 M sodium acetate buffer, pH 6.2, 3%

tetrahydrofuran, 0.7 mL/min flow rate), modified from (*80*). Identification and

quantification of the D- and L-serine and glycine was based on retention times and

peak areas, compared with those associated with external standards (calibration

curves were built by injecting 0.005-0.2 pmol of standards). To confirm the identity of

D-enantiomers, a selective degradation of D-amino acids in aCSF samples was

obtained by adding 10 μg of M213R RgDAAO for 4 h at 30 °C. Statistical analyses

were performed using GraphPad Prism 9.0.

Data acquisition and analysis

Pulse generation and data acquisition were performed using clampEx (clampEx 10.5,

Molecular Devices). Data analysis was performed using pClamp 10.5 (clampFit),

Mathematica 13 (Wolfram Research, Champaign, IL), Stimfit

(https://github.com/neurodroid/stimfit/wiki/Stimfit) and Prism 8 (GraphPad software,

La Jolla, CA). To detect sEPSCs, MiniAnalysis (v6.0.7, Synaptosoft Inc.) was used

and every single event was inspected and selected visually. To obtain an averaged

EPSC, events were superimposed to the initial phase of the current rise. To measure

the parameters of the averaged EPSC (amplitude, 20-80% rise time and decay), a

short baseline (5 ms) was used immediately preceding the rise of the event. The

averaged event was filtered at 2 kHz.

For the analysis of the NMDAR current component of sEPSCs at +40 mV, a dashed

vertical line was placed at 10 ms after the onset of the averaged EPSC to evaluate

the amplitude of the NMDAR-dependent synaptic current and for eEPSC, it was

placed 40 ms after the onset of the event (*29*, *35*). At this time point, the AMPAR-

current component is back to baseline. The peak amplitude of the AMPAR-current

component was measured at the peak of the averaged EPSC at -70 mV.

Bursts were defined as spontaneously occurring biphasic changes in the membrane

potential during which hyperpolarized periods between -60 and -80 mV alternated

with depolarized plateau potentials > -60 mV superimposed by APs. This firing

activity was stable and reproducible in the presence of NMDA with the SK channel

blocker apamin and hyperpolarizing current injection (*20*, *21*).

Statistical analysis

Experimental values are expressed as mean ± standard error of the mean (SEM).

Error bars in the figures also indicate SEM. Statistical analysis was performed using

Prism 8. Data were tested for normal distribution with Shapiro-wilk and d'Agostino

test. Accordingly, paired-data were either tested by ANOVA 1 test followed by a

Tukey post hoc test or Friedman test followed by Dunn's post hoc when more than 2

conditions. Unpaired-data were tested with Kruskal-Wallis test followed by Dunn's

test. In the case of 2 conditions, either T-test or Wilcoxon test (paired data) and

Mann-Withney test (unpaired data) were used. In the case of multiple conditions in 2

different stimulations protocol , ANOVA 2 test followed by a Tukey post hoc test were

applied. Significance level is given as * < 0.05, ** < 0.01, *** < 0.001, **** < 0.0001

and ns (not significant) corresponding to > 0.05 is represented in the histograms.

Acknowledgements

We thank Drs. Jean-Marc Goaillard and for critical reading of the manuscript and
helpful comments, Sandy El Sayed and Laurent Massotte for excellent technical
assistance and the collaboration with the GIGA-imaging platform for using the
confocal microscope and Imaris. **Funding:** This work was supported by grants from
the Belgian F.R.S. - F.N.R.S. (FRIA), the Fondation Leon Fredericq, Prix de l'espoir
from Marie-José De Ridder and S. R. Fonds facultaires de l'Université de Liège to
D.E. Z.M. is a PhD student of the Life Sciences and Biotechnology course at the
University of Insubria. L.P. thanks the support from FAR Fondo di Ateneo per la
Ricerca.

666
Author contributions:

Conceptualization: DE

Methodology: DE, LP, SR, KJ, LC, ZM and LV

Investigation: SR, LV, and LC

Writing – Original Draft: DE

Writing – Review & Editing: DE and SR

Funding Acquisition: SR, LP and DE

Resources: VS and LP

Supervision: DE

Competing Interests

The authors declare no competing interests.

Data and materials availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

 Figures and Tables

 Fig. 1. Glycine controls the occurrence of spontaneous bursting

(**A**), Voltage traces from a SNc DA neuron in the presence of NMDA (30 µM) and

apamin (300 nM, top trace black), following incubation with DAAO (45 min, 0.2

units/ml; middle trace red) and with DAAO supplemented by D-serine (100 µM,

bottom trace light red). Individual neurons were submitted sequentially to DAAO and

then to DAAO+D-serine.

(**B**), Voltage traces in the presence of NMDA (30 µM) and apamin (300 nM, top trace

black), following treatment with GO (45 min, 0.2 units/ml; middle trace blue) and with

GO supplemented by glycine (1 mM, bottom trace light blue). Individual neurons

were submitted sequentially to GO and then to GO+glycine.

(**C**), Histogram for the distribution of interspike interval (ISI) for a DA neuron in control

(light grey), in DAAO (red) and in DAAO+D-serine (light red). Note the similar

distributions for the three conditions.

(**D**), Summary plot showing mean burst duration. Control: 0.90 ± 0.15 s; DAAO: 0.81

± 0.12 s; DAAO+D-serine: 1.41 ± 0.23 s. Duration was not changed in DAAO

(Friedman test followed by post hoc Dunn's test, ns versus control). Burst duration

was increased in DAAO+D-serine (Friedman test followed by post hoc Dunn's test,

*$P < 0.05$ for DAAO+D-serine versus DAAO and for DAAO+D-serine versus control).

(**E**), Summary plot showing mean number of action potentials per burst. Control: 3.61

± 0.48; DAAO: 4.34 ± 0.72; DAAO+D-serine: 6.16 ± 1.05. AP number was not

changed in DAAO (Friedman test followed by post hoc Dunn's test, ns versus

control). AP number was increased in DAAO+D-serine (Friedman test followed by

post hoc Dunn's test, *$P < 0.05$ for DAAO+D-serine versus DAAO and **$P < 0.01$ for

DAAO+D-serine versus control).

(**F**), Summary plot showing mean burst frequency. Control: 0.36 ± 0.08 Hz; DAAO:

0.26 ± 0.04 Hz; DAAO+D-serine: 0.23 ± 0.03 Hz. Burst frequency was changed in

neither condition (Friedman test followed by post hoc Dunn's test, ns versus control,

ns for DAAO+D-serine versus DAAO and ns for DAAO+D-serine versus control). (**D-**

F), Bars represent mean ± SEM; points indicate data from individual experiments.

Data points are in control, n=8; DAAO, n=8; DAAO+D-serine, n=8.

(**G**), ISI histogram for the distribution of burst activity in control (grey), in GO (blue)

and in GO supplemented by glycine (light blue). Note the shift of the second large

peak in GO towards higher intervals in comparison to control.

(**H**), Summary plot showing burst duration. Control: 0.92 ± 0.18 s; GO: 1.11 ± 0.23 s;

GO+glycine: 1.90 ± 0.51 s. Duration was not changed in GO (Friedman test followed

by post hoc Dunn's test, ns versus control). Burst duration was increased in

GO+glycine (Friedman test followed by post hoc Dunn's test, *$P < 0.05$ for

GO+glycine versus GO and **$P < 0.01$ for GO+glycine versus control).

(**I**), Summary plot showing AP per burst. Control: 2.84 ± 0.22; GO: 2.87 ± 0.29; GO +

glycine: 4.71 ± 1.10. AP number was not changed in GO (Friedman test followed by

post hoc Dunn's test, ns versus control). AP number was increased in GO+glycine

(Friedman test followed by post hoc Dunn's test, *$P < 0.05$ for GO+glycine versus

GO and for GO+glycine versus control).

(**J**), Summary plot showing mean burst frequency. Control: 0.47 ± 0.06 Hz; GO: 0.17

± 0.05 Hz; GO + glycine: 0.19 ± 0.05 Hz. Burst frequency was decreased in GO

(Anova test followed by post hoc Tukey's test, **$P < 0.01$ versus control). Frequency

was not further changed in GO+glycine (Anova test followed by post hoc Tukey's

test, ns versus GO). Frequency was changed in GO+glycine in comparison to control

(Anova test followed by post hoc Tukey's test, **$P < 0.01$). (**H-J**), Bars represent

mean ± SEM; points indicate data from individual experiments. Data points are in

control, n=8; GO, n=8; GO+glycine, n=8.

 Fig. 2. D-serine is the main co-agonist of synaptic NMDA receptors in SNc DA

neurons

(**A**), Averaged traces (~100 events) of sEPSCs at +40 mV (upper traces) and -70 mV

(lower traces) in control, in DAAO (45 min, 0.2 units/ml) and in DAAO+D-serine (100

µM). At the right are superimposed traces for the three conditions. For each

condition, traces were recorded in the presence of picrotoxin (50 µM), CGP 55845 (1

µM) and strychnine (10 µM) to block GABA$_A$, GABA$_B$, and Glycine receptors,

respectively. Amplitude of NMDAR-mediated component was measured at +40mV

(vertical dashed lines). Note the decrease in amplitude of the averaged sEPSC in

DAAO at +40 mV.

(**B**), Averaged traces (~100 events) of sEPSCs at +40 mV and -70 mV in control, in

GO (45 min, 0.2 units/ml) and in GO+glycine (1 mM). Superimposed traces are at the

right.

(**C**), Summary bar graph of traces recorded as in (**A**). At +40 mV, control: 5.69 ± 1.43

pA; DAAO: 2.02 ± 0.36 pA; DAAO+D-serine: 3.28 ± 0.49 pA. NMDAR-mediated

current was decreased after treatment (Friedman test followed by Dunn's post hoc

test, **$P < 0.01$ versus control). NMDAR current was increased in DAAO+D-serine

(Friedman test followed by Dunn's post hoc test, *$P < 0.05$ versus DAAO) but was

not changed in comparison to control (Friedman test followed by Dunn's post hoc

test, ns for DAAO+D-serine versus control). At -70 mV, control: -8.84 ± 0.87 pA;

DAAO: -8.07 ± 0.71 pA; DAAO+D-serine: -8.48 ± 1.03 pA. AMPAR-mediated

currents were changed in neither condition (Anova 1 test followed by Tukey's post

hoc test). Bars represent mean ± SEM; points indicate data from individual

experiments. Data points are in control, n=7; DAAO, n=7; DAAO+D-serine, n=7.

(**D**), Summary bar graph of traces recorded as in (**B**). At +40 mV, control: 4.30 ± 0.43

pA; GO: 4.26 ± 0.48 pA; GO+glycine: 6.01 ± 0.74 pA. NMDAR current was not

changed in GO (Friedman test followed by Dunn's post hoc test, ns versus control).

NMDAR current was increased in GO+glycine (Friedman test followed by Dunn's

post hoc test, *$P < 0.05$ for GO+glycine versus GO and GO+glycine versus control).

At -70 mV, control: -10.09 ± 1.29 pA; GO: -8.80 ± 0.86 pA; GO+glycine: -7.69 ± 0.64

pA. AMPAR-mediated currents were changed in neither condition (Anova 1 test

followed by Tukey's post hoc test). Bars represent mean ± SEM; points indicate data

from individual experiments. Data points are in control, n=7; GO, n=7; GO+glycine,

n=7.

 Fig. 3. D-serine is the main co-agonist of NMDA receptors in STN-SNc

synapses

(**A**), Averaged traces (~30 events) of eEPSCs at +40 mV (upper traces) and -70 mV

(lower traces) in control, in DAAO and in DAAO+D-serine (100 µM). At the right are

superimposed traces from the three conditions. A focal double-barreled synaptic

stimulating theta electrode was used to activate afferent fibers originating from the

STN. For each condition, traces were recorded in the presence of picrotoxin (50 µM),

CGP 55845 (1 µM) and strychnine (10 µM). Amplitude of NMDAR-mediated

component was measured at +40mV (vertical dashed lines). In the square at the right

is an enlargement of the superimposed currents traces at -70 mV.

(**B**), Averaged traces (~30 events) of eEPSCs at +40 mV (upper traces) and -70 mV

(lower traces) in control, in GO and in GO+glycine (1 mM). At the right are

superimposed traces from the three conditions. In the square at the right is an

enlargement of the superimposed currents traces at -70 mV.

(**C**), Summary bar graph of experiments as in (**A**). At +40 mV, control: 42.07 ± 8.70

pA; DAAO, 25.10 ± 9.65 pA; DAAO+D-serine: 45.97 ± 15.06 pA. NMDAR-mediated

current was decreased after DAAO treatment (Anova 1 test followed by Tukey's post

hoc test, ***P* < 0.01 versus control). NMDAR current was increased in DAAO+D-

serine (Anova 1 test followed by Tukey's post hoc test, **P* < 0.05 versus DAAO) but

was not changed in comparison to control (Anova 1 test followed by Tukey's post hoc

test, ns for DAAO+D-serine versus control). At -70 mV, control: -85.19 ± 14.46 pA;

DAAO: -77.85 ± 12.51 pA; DAAO+D-serine: -90.78 ± 20.55 pA. AMPAR-mediated

currents were changed in neither conditions (Anova 1 test followed by Tukey's post

hoc test). Data points are in control, n=7; DAAO, n=7; DAAO+D-serine, n=5.

(**D**), Summary bar graph of traces recorded as in (**B**). At +40 mV, control: 27.90 ±

6.28 pA; GO: 25.91 ± 7.21 pA; GO+glycine: 43.17 ± 8.83 pA. NMDAR current was

not changed in GO (Anova 1 test followed by Tukey's post hoc test, ns versus

control). NMDAR current was increased in GO+glycine (Anova 1 test followed by

Tukey's post hoc test, **P* < 0.05 versus GO). NMDAR mediated current in

GO+glycine was similar to current in control (Friedman test followed by Dunn's post

hoc test, ns versus control). At -70 mV, control: -67.94 ± 14.34 pA; GO: -71.00 ±

18.98 pA; GO + glycine: -65.05 ± 18.00 pA. AMPAR-mediated currents were

changed in neither conditions (Anova 1 test followed by Tukey's post hoc test). Bars

represent mean ± SEM; points indicate data from individual experiments. Data points

are in control, n=7; GO, n=7; GO+glycine, n=5.

     Fig. 4. Subunit composition of synaptic NMDARs in nigral DA neurons

   (**A**), Averaged eEPSCs (~30 events) recorded from SNc DA neurons at +40 mV in

   control conditions (black traces), after incubation with tricine (gray trace) and

   subsequently after tricine+Zn^{2+} (light grey trace). A focal double-barreled stimulating

   theta electrode was placed rostrally to the SN to activate fibers from the STN. For

   each condition traces were recorded in the presence of picrotoxin (50 µM), CGP

   55485 (1 µM), CNQX (10 µM), and strychnine (10 µM). tricine was used at 10 mM

   and Zn^{2+} at 60 µM (to have 300 nM free Zn^{2+}).

(**B**), Bar chart showing the mean amplitude of NMDAR-eEPSCs (top) in control

conditions (dark grey bar), in tricine (grey bar) and in tricine+Zn^{2+} (light grey bar). At

+40 mV, control: 34.21 ± 2.80 pA; tricine: 33.48 ± 2.99 pA; tricine+Zn^{2+}: 32.40 ± 3.00

pA. NMDAR-mediated current was neither modified in tricine (Anova test followed by

Tukey's post hoc test, ns versus control) nor in tricine+Zn^{2+} (Anova test followed by

Tukey's post hoc test, ns for tricine+Zn^{2+} versus tricine and for tricine+Zn^{2+} versus

control). Bar chart showing the weighted decay time constant of the NMDAR-

eEPSCs (bottom) in control conditions (dark grey bar), in tricine (grey bar) and in

tricine+Zn^{2+} (light grey bar). Weighted decay, control: 84.02 ± 14.13 ms; tricine: 84.01

± 14.28 ms; tricine+Zn^{2+}: 84.99 ± 12.21 ms. Decay of NMDAR current was neither

modified in tricine (Anova test followed by Tukey's post hoc test, ns versus control)

nor in tricine+Zn^{2+} (Anova test followed by Tukey's post hoc test, ns for tricine+Zn^{2+}

versus tricine and for tricine+Zn^{2+} versus control). Bars represent mean ± SEM;

points indicate data from individual experiments. Data points are in control, n=8;

tricine, n=8; tricine+Zn^{2+}, n=8.

(**C**), Averaged eEPSCs (~30 events) recorded at +40 mV in control conditions (black

traces) and in Ro25-6981 (grey traces). Ro25-6981 was used at 1 µM.

(**D**), Bar chart showing the mean amplitude of NMDAR-eEPSCs (top) in control

conditions (dark grey bar) and in Ro25-6981 (grey bar). At +40 mV, control: 38.60 ±

5.71 pA; Ro25-6981: 23.42 ± 4.66 pA. NMDAR current amplitude was decreased in

Ro25-6981 (Paired T-test, ***$P < 0.001$ versus control). Bar chart showing the

weighted decay time constant of the NMDAR-eEPSCs (bottom) in control conditions

(dark grey bar) and in Ro25-6981 (grey bar). Weighted decay, control: 91.04 ± 17.93

ms; Ro25-6981: 92.51 ± 16.37 ms). NMDAR current decay was not changed in

Ro25-6981 (Paired T-test, ns versus control). Data points are in control, n=9; Ro25-

6981, n=9.

(**E**), Averaged sEPSCs (~30 events) recorded at +40 mV in control conditions (black

traces) and in DQP-1105 (grey traces,top). DQP-1105 was used at 20 µM.

(**F**), Bar chart showing the amplitude of the NMDAR-eEPSCs (top) in control

conditions (dark grey bar) and in DQP-1105 (grey bar). At +40 mV, control: 35.99 ±

3.19 pA; DQP-1105: 19.38 ± 4.95 pA). NMDAR current amplitude was decreased in

DQP-1105 (Paired T-test, **P < 0.01 versus control). Bar chart showing the decay of

the NMDAR-eEPSCs (bottom) in control conditions (dark grey bar) and in DQP-1105

(grey bar). Weighted decay, control: 68.84 ± 24.38 ms; DQP-1105: 76.05 ± 26.30

ms). NMDAR current decay was not changed in DQP-1105 (Paired T-test, ns versus

control). Data points are in control, n=7; DQP-1105, n=7.

Fig. 5. Glycine gates extrasynaptic NMDARs containing GluN2B and GluN2D in

nigral DA neurons

(**A**), Tonic NMDAR-mediated current in control conditions (top), during incubation of

DAAO (45 min, 0.2 units/ml; below middle) and in GO (45 min, 0.2 units/ml; bottom)

revealed by the current shift induced by the application of D-AP5 (50 µM) at a holding

potential of +40 mV. Dashed red line indicate amplitude of holding current before the

application of D-AP5. Tonic currents were recorded in the presence of CNQX (10

µM), picrotoxin (50 µM), CGP 55485 (1 µM) and strychnine (10 µM). Data in the

different conditions are from distinct experiments.

(**B**), Tonic NMDAR-mediated current in tricine (10 mM, top), in tricine+Zn^{2+} (60 µM,

second from top), in Ro25-6981 (1 µM, third from top) and in DQP-1105 (20 µM,

bottom) revealed by the current shift induced by the application of D-AP5 (50 µM) at

a holding potential of +40 mV. Dashed red line indicate amplitude of holding current

without D-AP5.

(**C**), Summary bar graph of the mean amplitude of the NMDAR-mediated tonic

current. Control: -26.47 ± 2.51 pA; DAAO: -21.26 ± 3.52 pA; GO: -12.06 ± 3.03 pA;

tricine: -22.60 ± 4.22 pA; tricine + Zn^{2+}: -23.10 ± 3.24 pA; Ro25-6981: -8.09 ± 2.35

pA; DQP-1105: -6.90 ± 3.45 pA. NMDAR-mediated tonic current was unchanged in

DAAO, in tricine and in tricine+ Zn^{2+} (Kruskal-Wallis test followed by a post hoc

Dunn's test, ns versus control). NMDAR-mediated tonic current was reduced in GO

(Kruskal-Wallis test followed by a post hoc Dunn's test, *P < 0.05 versus control), in

Ro25-6981 and in DQP-1105 (Kruskal-Wallis test followed by a post hoc Dunn's test,

**P < 0.01 versus control). Bars represent mean ± SEM; points indicate data from

individual experiments. Data points are in control, n=11; DAAO, n=10; GO, n=12;

tricine, n=11; tricine + Zn^{2+}, n=13; Ro25-6981, n=11; DQP-1105, n=11.

    Fig. 6. Tonic NMDAR-mediated current is mediated by extrasynaptic NMDARs

(**A**), Experimental protocol to block synaptic NMDARs using the NMDAR open

channel blocker MK-801 (left). Control electrically evoked EPSCs (eEPSCs) at V_{hold} =

+40 mV under low stimulation frequency (0.1 Hz) in voltage-clamp. Switch to current-

clamp (V_{hold} ~ -80 mV) for ~20 min during the application of MK-801 (20 µM).

Washout of MK-801 and switch back to V_{hold} = +40 mV. Stimulation was resumed

using a frequency of 5 Hz for 16 s (*45*, *75*). Note the very strong reduction of the

NMDAR-EPSC amplitude when stimulation was resumed. Neuron was kept at V_{hold} =

+40 mV and D-AP5 was applied. During the whole experiment, CNQX (10 µM),

picrotoxin (50 μM), CGP 55485 (1 μM) and strychnine (10 μM) were present in the

bath.

(**B**), Experimental design to block both synaptic NMDARs and extrasynaptic

NMDARs using MK-801 (left). During application of MK-801, holding potential was

kept at +40 mV and repetitive afferent fiber stimulation frequency was low (0.2 Hz).

Note the quasi absence of the tonic current after application of D-AP5.

(**C**), Summary bar graph of the peak amplitude of the tonic NMDAR-mediated

current. Control (without MK-801 treatment): -26.47 ± 2.51 pA; after MK-801

treatment to reduce synaptic NMDARs (syn): -21.84 ± 2.30 pA; after MK-801

treatment and continuous synaptic stimulation to block synaptic and extrasynaptic

NMDARs (syn+extrasyn): -10.64 ± 1.97 pA). Tonic NMDAR-mediated current was

not different after MK-801 (Anova test followed by a post hoc Tukey's test, ns versus

control) but was decreased after MK-801 and stimulation application (Anova test

followed by a post hoc Tukey's test, *$P < 0.05$ for MK-801+stimulation versus MK-

801 and *$P < 0.05$ for MK-801+stimulation versus control). Bars represent mean ±

SEM; points indicate data from individual experiments. Data points are in control,

n=11; +MK-801 (syn), n=9; +MK-801 (syn+extrasyn), n=7.

Fig. 7. Bursting activity is mediated by extrasynaptic NMDARs

(**A**), Blockade of synaptic NMDARs using MK-801. Control eEPSCs at V_{hold} = +40

mV under low stimulation frequency (0.1 Hz) in voltage-clamp. Switch to current-

clamp (V_{hold} = -80 mV) for ~20 min during the application of MK-801 (20 µM).

Washout of MK-801 and switch back to V_{hold} = +40 mV. Stimulation at 5 Hz for 16 s.

Bursting activity (right) recorded in current-clamp induced by application of NMDA

and apamin. In current-clamp to generate bursts, CNQX (10 µM), picrotoxin (50 µM),

CGP 55485 (1 µM) and strychnine (10 µM) were present in the bath. Note the

presence of spontaneous bursts after the blockade of synaptic NMDARs.

(**B**), Histogram for the distribution of interspike interval (ISI) for a DA neuron in control

(light grey) and after blockade of synaptic NMDARs ($_{syn}$NMDARs blocked, dark gray).

Note the similar distributions for the two conditions.

(**C**), Summary plot showing burst duration in control and after the blockade of

synaptic NMDARs. Control: 1.10 ± 0.18 s; $_{syn}$NMDARs blocked: 1.43 ± 0.29 ms. Burst

duration was not changed when synaptic NMDARs were blocked (Mann-Whitney

test, ns versus control).

(**D**), Summary plot showing action potential per burst in control and after the

blockade of synaptic NMDARs. Control: 2.77 ± 0.25; $_{syn}$NMDARs blocked: $2.20 \pm$

0.30. AP number was not changed after synaptic NMDAR blockade (unpaired T-test,

ns versus control).

(**E**), Summary plot showing burst frequency in control and after the blockade of

synaptic NMDARs. Control: 0.36 ± 0.05 Hz; $_{syn}$NMDARs blocked: 0.35 ± 0.08 Hz.

Burst frequency was not different after synaptic NMDAR blockade (unpaired T-test,

ns versus control). (**C-E**) Bars are mean $\pm$ SEM; points indicate data from individual

experiments. Data points are in control, n=8; $_{syn}$NMDARs blocked, n=6.

Fig. 8. Bursting activity requires glycine and glutamate spillover during high regime of synaptic transmission

(**A**), Voltage traces of action potential in a SNc DA during local synaptic stimulation at 10 Hz for 500 ms in control (left), in DAAO (45min, 0.2 units/ml, middle left), in GO (45min, 0.2 units/ml, middle right) and in GO (45min, 0.2 units/ml) + glycine (1 mM, right). Voltage traces were recorded in the presence of CNQX (10 µM), picrotoxin (50 µM), CGP 55485 (1 µM) and strychnine (10 µM).

(**B**), Voltage traces of action potential in a SNc DA during local synaptic stimulation at 50 Hz for 500 ms in control (left), in DAAO (middle left), in GO (middle right) and in GO + glycine (1 mM, right).

(**C**), Summary plot showing the mean number of action potential generated during local synaptic stimulation at 10 Hz (left). Control: 0.50 ± 0.17; DAAO: 0.67 ± 0.33; GO: 0.78 ± 0.15; GO + glycine: 1.00 ± 0.32. The number of APs was changed in

neither condition (Anova 2 followed by Tukey's post hoc Test, ns). AP number for

synaptic stimulation at 50 Hz (right). Control: 2.60 ± 0.31; DAAO: 3.17 ± 0.48; GO:

1.22 ± 0.28; GO + glycine: 2.40 ± 0.25. AP number was not different in DAAO and in

GO+glycine (Anova 2 followed by Tukey's post hoc Test, ns versus control) but was

decrease in GO (Anova 2 followed by Tukey's post hoc Test, **$P < 0.01$ for GO

versus control and ****$P < 0.0001$ for GO versus DAAO). AP number was increased

in GO+glycine in comparison to GO (Anova 2 followed by Tukey's post hoc Test, *P

< 0.05).

(**D**), Summary plot showing the intraburst frequency during local synaptic stimulation

at 10 Hz (left). Control: 0.36 ± 0,27 Hz; DAAO: 0.50 ± 0.50 Hz; GO: 0.22 ± 0.18 Hz;

GO + glycine: 0.75 ± 0.73 Hz. Intraburst frequency was changed in neither condition

(Anova 2 followed by Tukey's post hoc Test, ns). Intraburst frequency for stimulation

at 50 Hz (right). Control: 3.79 ± 0.66 Hz; DAAO, 5.49 ± 1.26 Hz; GO, 1.11 ± 0.44 Hz;

GO + glycine, 4.15 ± 0.33 Hz. Intraburst frequency was not different in DAAO and in

GO+glycine (Anova 2 followed by Tukey's post hoc Test, ns versus control) but was

decreased in GO (Anova 2 followed by Tukey's post hoc Test, **$P < 0.01$ for GO

versus control and ****$P < 0.0001$ for GO versus DAAO). Intraburst frequency was

increased in GO+glycine in comparison to GO (Anova 2 followed by Tukey's post hoc

Test, **$P < 0.01$).

(**E**), Summary plot showing the time to the first action potential during local synaptic

stimulation at 10 Hz (left). Control: 536.40 ± 104.60 ms; DAAO: 491.62 ± 112.93 ms;

GO: 483.43 ± 55.71 ms; GO+glycine: 451.97 ± 49.12 ms. Time to first AP was

changed in neither condition (Anova 2 followed by Tukey's post hoc Test, ns). Time

to first AP for stimulation at 50 Hz (right). Control: 104.60 ± 11.77 ms; DAAO: 112.93

± 19.85 ms; GO: 415.52 ± 33.76 ms; GO+glycine: 146.70 ± 20.61 ms. Time to first

AP was not different in DAAO and in GO+glycine (Anova 2 followed by Tukey's post

hoc Test, ns versus control) but was increased in GO (Anova 2 followed by Tukey's

post hoc Test, ***P < 0.001 for GO versus control and ****P < 0.0001 for GO versus

DAAO). Time to first AP was decreased in GO+glycine in comparison to GO (Anova

2 followed by Tukey's post hoc Test, *P < 0.05). (**C-E**) Bars are means ± SEM; points

indicate data from individual experiments. Data points are in control, n=10; DAAO,

n=6; GO, n=9; GO + glycine, n=5.

REFERENCES

1.   M. W. Howe, D. A. Dombeck, Rapid signalling in distinct dopaminergic axons during
locomotion and reward. *Nature* **535**, 505–510 (2016).
2.   J. A. da Silva, F. Tecuapetla, V. Paixão, R. M. Costa, Dopamine neuron activity before
action initiation gates and invigorates future movements. *Nature* **554**, 244–248 (2018).
3.   O. Hornykiewicz, "The discovery of dopamine deficiency in the parkinsonian brain" in
Parkinson's Disease and Related Disorders (Springer Vienna, Vienna, 2006), pp. 9–
15.
4.   F. G. Gonon, Nonlinear relationship between impulse flow and dopamine released by
rat midbrain dopaminergic neurons as studied by in vivo electrochemistry.
Neuroscience **24**, 19–28 (1988).
5.   R. Ammari, C. Lopez, H. Fiorentino, F. Gonon, C. Hammond, A mouse juvenile or
adult slice with preserved functional nigro-striatal dopaminergic neurons. *Neuroscience*
159, 3–6 (2009).
6.   J. E. Markowitz, W. F. Gillis, M. Jay, J. Wood, R. W. Harris, R. Cieszkowski, R. Scott,
D. Brann, D. Koveal, T. Kula, C. Weinreb, M. A. M. Osman, S. R. Pinto, N. Uchida, S.
W. Linderman, B. L. Sabatini, S. R. Datta, Spontaneous behaviour is structured by
reinforcement without explicit reward. *Nature* **614**, 108–117 (2023).
7.   K. Chergui, H. Akaoka, P. J. Charléty, C. F. Saunier, M. Buda, G. Chouvet,
Subthalamic nucleus modulates burst firing of nigral dopamine neurones via NMDA
receptors. [Preprint] (1994).
8.   M. Watabe-Uchida, L. Zhu, S. K. Ogawa, A. Vamanrao, N. Uchida, Whole-brain
mapping of direct inputs to midbrain dopamine neurons. *Neuron* **74**, 858–873 (2012).
9.   S. L. C. Brothwell, J. L. Barber, D. T. Monaghan, D. E. Jane, A. J. Gibb, S. Jones,
NR2B- and NR2D-containing synaptic NMDA receptors in developing rat substantia
nigra pars compacta dopaminergic neurones. *J Physiol* **586**, 739–50 (2008).
10.   S. A. Swanger, K. M. Vance, J. F. Pare, F. Sotty, K. Fog, Y. Smith, S. F. Traynelis,
NMDA receptors containing the GluN2D subunit control neuronal function in the
subthalamic nucleus. *Journal of Neuroscience* **35**, 15971–15983 (2015).
11.   F. Yi, S. Bhattacharya, C. M. Thompson, S. F. Traynelis, K. B. Hansen, Functional and
pharmacological properties of triheteromeric GluN1/2B/2D NMDA receptors. *Journal*
of Physiology **597**, 5495–5514 (2019).
12.   S. G. Brickley, C. Misra, M. H. S. Mok, M. Mishina, S. G. Cull-Candy, NR2B and
NR2D subunits coassemble in cerebellar Golgi cells to form a distinct NMDA receptor
subtype restricted to extrasynaptic sites. *J Neurosci* **23**, 4958–66 (2003).
13.   A. Momiyama, Distinct synaptic and extrasynaptic NMDA receptors identified in
dorsal horn neurones of the adult rat spinal cord. *Journal of Physiology* **523**, 621–628
(2000).
14.   S. C. Harney, D. E. Jane, R. Anwyl, Extrasynaptic NR2D-containing NMDARs are
recruited to the synapse during LTP of NMDAR-EPSCs. *Journal of Neuroscience* **28**,
11685–11694 (2008).
15.   N. A. Lozovaya, S. E. Grebenyuk, T. Sh. Tsintsadze, B. Feng, D. T. Monaghan, O. A.
Krishtal, Extrasynaptic NR2B and NR2D subunits of NMDA receptors shape
'superslow' afterburst EPSC in rat hippocampus. *J Physiol* **558**, 451–463 (2004).
16.   S. Jones, A. J. Gibb, Functional NR2B- and NR2D-containing NMDA receptor
channels in rat substantia nigra dopaminergic neurones. *J Physiol* **569**, 209–21 (2005).
17.   L. P. Wang, F. Li, D. Wang, K. Xie, D. Wang, X. Shen, J. Z. Tsien, NMDA receptors
in dopaminergic neurons are crucial for habit learning. *Neuron* **72**, 1055–1066 (2011).

18. L. S. Zweifel, J. G. Parker, C. J. Lobb, A. Rainwater, V. Z. Wall, J. P. Fadok, M.
Darvas, M. J. Kim, S. J. Mizumori, C. A. Paladini, P. E. Phillips, R. D. Palmiter,
Disruption of NMDAR-dependent burst firing by dopamine neurons provides selective
assessment of phasic dopamine-dependent behavior. *Proc Natl Acad Sci U S A* **106**,
7281–7288 (2009).
19. K. Otomo, J. Perkins, A. Kulkarni, S. Stojanovic, J. Roeper, C. A. Paladini, In vivo
patch-clamp recordings reveal distinct subthreshold signatures and threshold dynamics
of midbrain dopamine neurons. *Nat Commun* **11**, 1–15 (2020).
20. V. Seutin, S. W. Johnson, R. A. North, Apamin increases NMDA-induced burst-firing
of rat mesencephalic dopamine neurons. *Brain Res* **630**, 341–344 (1993).
21. G. Destreel, V. Seutin, D. Engel, Subsaturation of the N-methyl-D-aspartate receptor
glycine site allows the regulation of bursting activity in juvenile rat nigral dopamine
neurons. *Eur J Neurosci* **50**, 3454–3471 (2019).
22. M. E. Soden, G. L. Jones, C. A. Sanford, A. S. Chung, A. D. Güler, C. Chavkin, R.
Luján, L. S. Zweifel, Disruption of Dopamine Neuron Activity Pattern Regulation
through Selective Expression of a Human KCNN3 Mutation. *Neuron* **80**, 997–1009
(2013).
23. S. N. Blythe, J. F. Atherton, M. D. Bevan, Synaptic Activation of Dendritic AMPA and
NMDA Receptors Generates Transient High-Frequency Firing in Substantia Nigra
Dopamine Neurons In Vitro. *J Neurophysiol* **97**, 2837–2850 (2007).
24. J. W. Johnson, P. Ascher, Glycine potentiates the NMDA response in cultured mouse
brain neurons. [Preprint] (1987). https://doi.org/10.1038/325529a0.
25. N. W. Kleckner, R. Dingledine, Requirement for Glycine in Activation of NMDA-
Receptors Expressed in Xenopus Oocytes. *Science* **241**, 835–837 (1988).
26. M. C. Currás, B. S. Pallotta, Single-channel evidence for glycine and NMDA
requirement in NMDA receptor activation. *Brain Res* **740**, 27–40 (1996).
27. M. Le Bail, M. Martineau, S. Sacchi, N. Yatsenko, I. Radzishevsky, S. Conrod, K. Ait
Ouares, H. Wolosker, L. Pollegioni, J.-M. Billard, J.-P. Mothet, Identity of the NMDA
receptor coagonist is synapse specific and developmentally regulated in the
hippocampus. *Proc Natl Acad Sci U S A* **112**, E204-13 (2015).
28. T. Papouin, L. Ladépêche, J. Ruel, S. Sacchi, M. Labasque, M. Hanini, L. Groc, L.
Pollegioni, J. P. Mothet, S. H. R. Oliet, Synaptic and extrasynaptic NMDA receptors
are gated by different endogenous coagonists. *Cell* **150**, 633–646 (2012).
29. Y. Li, S. Sacchi, L. Pollegioni, A. C. Basu, J. T. Coyle, V. Y. Bolshakov, Identity of
endogenous NMDAR glycine site agonist in amygdala is determined by synaptic
activity level. *Nat Commun* **4**, 1760 (2013).
30. S. Neame, H. Safory, I. Radzishevsky, A. Touitou, F. Marchesani, M. Marchetti, S.
Kellner, S. Berlin, V. N. Foltyn, S. Engelender, J. M. Billard, H. Wolosker, The
NMDA receptor activation by D-serine and glycine is controlled by an astrocytic
Phgdh-dependent serine shuttle. *Proc Natl Acad Sci U S A* **116**, 20736–20742 (2019).
31. J. M. Wong, O. O. Folorunso, E. V. Barragan, C. Berciu, T. L. Harvey, J. T. Coyle, D.
T. Balu, J. A. Gray, Postsynaptic Serine Racemase Regulates NMDA Receptor
Function. *The Journal of Neuroscience* **40**, 9564–9575 (2020).
32. J.-P. Mothet, M. Le Bail, J.-M. Billard, Time and space profiling of NMDA receptor
co-agonist functions. *J Neurochem* **135**, 210–225 (2015).
33. K. Chergui, P. J. Charléty, H. Akaoka, C. F. Saunier, J. □L Brunet, M. Buda, T. H.
Svensson, G. Chouvet, Tonic Activation of NMDA Receptors Causes Spontaneous
Burst Discharge of Rat Midbrain Dopamine Neurons In Vivo. *Eur J Neurosci* **5**, 137–
144 (1993).

34. T. A. Hage, Y. Sun, Z. M. Khaliq, Electrical and Ca2+ signaling in dendritic spines of
substantia nigra dopaminergic neurons. *Elife* **5**, 1–26 (2016).
35. M. Jang, K. B. Um, J. Jang, H. J. Kim, H. Cho, S. Chung, M. K. Park, Coexistence of
glutamatergic spine synapses and shaft synapses in substantia nigra dopamine neurons.
Sci Rep **5**, 14773 (2015).
36. C. Bellone, R. A. Nicoll, Rapid Bidirectional Switching of Synaptic NMDA Receptors.
Neuron **55**, 779–785 (2007).
37. P. Paoletti, P. Ascher, J. Neyton, High-Affinity Zinc Inhibition of NMDA NR1–NR2A
Receptors. *Journal of Neuroscience* **17**, 5711–5725 (1997).
38. A. M. Vergnano, N. Rebola, L. P. Savtchenko, P. S. Pinheiro, M. Casado, B. L.
Kieffer, D. A. Rusakov, C. Mulle, P. Paoletti, Zinc dynamics and action at excitatory
synapses. *Neuron* **82**, 1101–1114 (2014).
39. R. S. Petralia, Y. X. Wang, F. Hua, Z. Yi, A. Zhou, L. Ge, F. A. Stephenson, R. J.
Wenthold, Organization of NMDA receptors at extrasynaptic locations. *Neuroscience*
167, 68–87 (2010).
40. K. Le Meur, M. Galante, M. C. Angulo, E. Audinat, Tonic activation of NMDA
receptors by ambient glutamate of non-synaptic origin in the rat hippocampus. *J
Physiol* **580**, 373–83 (2007).
41. A. R. Wild, M. Bollands, P. G. Morris, S. Jones, Mechanisms regulating spill-over of
synaptic glutamate to extrasynaptic NMDA receptors in mouse substantia nigra
dopaminergic neurons. *Eur J Neurosci* **42**, 2633–43 (2015).
42. N. O. Dalby, I. Mody, Activation of NMDA receptors in rat dentate gyrus granule cells
by spontaneous and evoked transmitter release. *J Neurophysiol* **90**, 786–97 (2003).
43. A. Z. Harris, D. L. Pettit, Extrasynaptic and synaptic NMDA receptors form stable and
uniform pools in rat hippocampal slices. *Journal of Physiology* **584**, 509–519 (2007).
44. J. E. Huettner, B. P. Bean, Block of N-methyl-D-aspartate-activated current by the
anticonvulsant MK-801: selective binding to open channels. *Proceedings of the
National Academy of Sciences* **85**, 1307–1311 (1988).
45. W. Koh, M. Park, Y. E. Chun, J. Lee, H. S. Shim, M. G. Park, S. Kim, M. Sa, J. Joo, H.
Kang, S.-J. Oh, J. Woo, H. Chun, S. E. Lee, J. Hong, J. Feng, Y. Li, H. Ryu, J. Cho, C.
J. Lee, Astrocytes Render Memory Flexible by Releasing D-Serine and Regulating
NMDA Receptor Tone in the Hippocampus. *Biol Psychiatry* **91**, 740–752 (2022).
46. D. dan Liu, Q. Yang, S. tian Li, Activation of extrasynaptic NMDA receptors induces
LTD in rat hippocampal CA1 neurons. [Preprint] (2013).
https://doi.org/10.1016/j.brainresbull.2012.12.003.
47. T. A. Hage, Z. M. Khaliq, Tonic Firing Rate Controls Dendritic Ca2+ Signaling and
Synaptic Gain in Substantia Nigra Dopamine Neurons. *Journal of Neuroscience* **35**,
5823–5836 (2015).
48. G. Beaudoin, J. Gomez, J. Perkins, J. Bland, A. Petko, C. Paladini, Cocaine selectively
reorganizes excitatory inputs to substantia Nigra pars compacta dopamine neurons.
Journal of Neuroscience **38**, 1151–1159 (2018).
49. A. Z. Harris, D. L. Pettit, Recruiting Extrasynaptic NMDA Receptors Augments
Synaptic Signaling. *J Neurophysiol* **99**, 524–533 (2008).
50. G. E. Hardingham, H. Bading, Synaptic versus extrasynaptic NMDA receptor
signalling: Implications for neurodegenerative disorders. *Nat Rev Neurosci* **11**, 682–
696 (2010).
51. M. T. Harnett, B. E. Bernier, K.-C. Ahn, H. Morikawa, Burst-timing-dependent
plasticity of NMDA receptor-mediated transmission in midbrain dopamine neurons.
Neuron **62**, 826–38 (2009).

52. R. E. Perszyk, J. O. DiRaddo, K. L. Strong, C. M. Low, K. K. Ogden, A. Khatri, G. A.
Vargish, K. A. Pelkey, L. Tricoire, D. C. Liotta, Y. Smith, C. J. McBain, S. F.
Traynelis, GluN2D-containing N-methyl-D-aspartate receptors mediate synaptic
transmission in hippocampal interneurons and regulate interneuron activity. *Mol*
Pharmacol **90**, 689–702 (2016).
53. I. Riebe, H. Seth, G. Culley, Z. Dósa, S. Radi, K. Strand, V. Fröjd, E. Hanse, Tonically
active NMDA receptors - a signalling mechanism critical for interneuronal excitability
in the CA1 stratum radiatum. *Eur J Neurosci* **43**, 169–178 (2016).
54. C. Rauner, G. Köhr, Triheteromeric NR1/NR2A/NR2B receptors constitute the major
N-methyl-D-aspartate receptor population in adult hippocampal synapses. *Journal of*
Biological Chemistry **286**, 7558–7566 (2011).
55. A. Panatier, D. T. Theodosis, J. P. Mothet, B. Touquet, L. Pollegioni, D. A. Poulain, S.
H. R. Oliet, Glia-Derived d-Serine Controls NMDA Receptor Activity and Synaptic
Memory. *Cell* **125**, 775–784 (2006).
56. L. Curcio, M. V. Podda, L. Leone, R. Piacentini, A. Mastrodonato, P. Cappelletti, S.
Sacchi, L. Pollegioni, C. Grassi, M. D'Ascenzo, Reduced d-serine levels in the nucleus
accumbens of cocaine-treated rats hinder the induction of NMDA receptor-dependent
synaptic plasticity. *Brain* **136**, 1216–1230 (2013).
57. P. Fossat, F. R. Turpin, S. Sacchi, J. Dulong, T. Shi, J. M. Rivet, J. V. Sweedler, L.
Pollegioni, M. J. Millan, S. H. R. Oliet, J. P. Mothet, Glial D-serine gates NMDA
receptors at excitatory synapses in prefrontal cortex. *Cerebral Cortex* **22**, 595–606
(2012).
58. J. S. Ferreira, T. Papouin, L. Ladépêche, A. Yao, V. C. Langlais, D. Bouchet, J.
Dulong, J. P. Mothet, S. Sacchi, L. Pollegioni, P. Paoletti, S. H. Oliet, L. Groc, Co-
agonists differentially tune GluN2B-NMDA receptor trafficking at hippocampal
synapses. *Elife* **6**, 1–22 (2017).
59. K. B. Hansen, L. P. Wollmuth, D. Bowie, H. Furukawa, F. S. Menniti, A. I.
Sobolevsky, G. T. Swanson, S. A. Swanger, I. H. Greger, T. Nakagawa, C. J. McBain,
V. Jayaraman, C. M. Low, M. L. Dell'Acqua, J. S. Diamond, C. R. Camp, R. E.
Perszyk, H. Yuan, S. F. Traynelis, Structure, Function, and Pharmacology of
Glutamate Receptor Ion Channels. [Preprint] (2021).
https://doi.org/10.1124/pharmrev.120.000131.
60. P. E. Chen, M. T. Geballe, E. Katz, K. Erreger, M. R. Livesey, K. K. O'toole, P. Le, C.
J. Lee, J. P. Snyder, S. F. Traynelis, D. J. A. Wyllie, Modulation of glycine potency in
rat recombinant NMDA receptors containing chimeric NR2A/2D subunits expressed in
Xenopus laevis oocytes. *Journal of Physiology* **586**, 227–245 (2008).
61. B. Cubelos, C. Giménez, F. Zafra, Localization of the GLYT1 glycine transporter at
glutamatergic synapses in the rat brain. *Cereb Cortex* **15**, 448–59 (2005).
62. J. G. Dopico, T. González-Hernández, I. M. Pérez, I. G. García, A. M. Abril, J. O.
Inchausti, M. Rodríguez Díaz, Glycine release in the substantia nigra: Interaction with
glutamate and GABA. *Neuropharmacology* **50**, 548–557 (2006).
63. C. Henneberger, L. Bard, C. King, A. Jennings, D. A. Rusakov, NMDA receptor
activation: Two targets for two co-agonists. [Preprint] (2013).
https://doi.org/10.1007/s11064-013-0987-2.
64. A. Drotos, Y. Herrera, R. Zarb, M. Roberts, GluN2D-containing NMDA receptors
enhance temporal integration in VIP neurons in the inferior colliculus. *bioRxiv*
preprint, 1–13 (2023).
65. N. Arnth-Jensen, D. Jabaudon, M. Scanziani, Cooperation between independent
hippocampal synapses is controlled by glutamate uptake. *Nat Neurosci* **5**, 325–331
(2002).

66. K. D. Oikonomou, S. M. Short, M. T. Rich, S. D. Antic, Extrasynaptic glutamate
receptor activation as cellular bases for dynamic range compression in pyramidal
neurons. *Front Physiol*, 1–22 (2012).

67. M. Herman, C. Jahr, Extracellular glutamate concentration in hippocampal slice. *J*
Neurosci **27**, 9736–41 (2007).

68. T. Suzuki, S. Kodama, C. Hoshino, T. Izumi, H. Miyakawa, A plateau potential
mediated by the activation of extrasynaptic NMDA receptors in rat hippocampal CA1
pyramidal neurons. *Eur J Neurosci* **28**, 521–534 (2008).

69. L. Yao, Y. Rong, X. Ma, H. Li, D. Deng, Y. Chen, S. Yang, T. Peng, T. Ye, F. Liang,
N. Xu, Q. Zhou, Extrasynaptic NMDA Receptors Bidirectionally Modulate Intrinsic
Excitability of Inhibitory Neurons. *Journal of Neuroscience* **42**, 3066–3079 (2022).

70. U. Heresco-Levy, S. Shoham, D. C. Javitt, Glycine site agonists of the N-methyl-d-
aspartate receptor and Parkinson's disease: A hypothesis. *Movement Disorders* **28**,
419–424 (2013).

71. C. H. Tsai, H. C. Huang, B. L. Liu, C. I. Li, M. K. Lu, X. Chen, M. C. Tsai, Y. W.
Yang, H. Y. Lane, Activation of N-methyl-D-aspartate receptor glycine site temporally
ameliorates neuropsychiatric symptoms of Parkinson's disease with dementia.
Psychiatry Clin Neurosci **68**, 692–700 (2014).

72. I. Frouni, E. Kim, J. Shaqfah, D. Bédard, C. Kwan, S. Belliveau, P. Huot, [3H]-NFPS
binding to the glycine transporter 1 in the hemi-parkinsonian rat brain. *Exp Brain Res*
242, 1203–1214 (2024).

73. H.-X. Wang, W.-J. Gao, Development of calcium-permeable AMPA receptors and
their correlation with NMDA receptors in fast-spiking interneurons of rat prefrontal
cortex. *J Physiol* **588**, 2823–38 (2010).

74. A. Kumar, O. Schiff, E. Barkai, B. W. Mel, A. Poleg-Polsky, J. Schiller, NMDA spikes
mediate amplification of inputs in the rat piriform cortex. *Elife* **7** (2018).

75. Q. Yang, G. Zhu, D. Liu, J. G. Ju, Z. H. Liao, Y. X. Xiao, Y. Zhang, N. Chao, J. Wang,
W. Li, J. H. Luo, S. T. Li, Extrasynaptic NMDA receptor dependent long-term
potentiation of hippocampal CA1 pyramidal neurons. *Sci Rep* **7**, 1–9 (2017).

76. S. Sacchi, S. Lorenzi, G. Molla, M. S. Pilone, C. Rossetti, L. Pollegioni, Engineering
the Substrate Specificity ofd-Amino-acid Oxidase. *Journal of Biological Chemistry*
277, 27510–27516 (2002).

77. V. Job, G. Molla, M. S. Pilone, L. Pollegioni, Overexpression of a recombinant
wild□type and His□tagged *Bacillus subtilis* glycine oxidase in *Escherichia coli*. *Eur J*
Biochem **269**, 1456–1463 (2002).

78. S. Fantinato, L. Pollegioni, M. Pilone S., Engineering, expression and purification of a
His-tagged chimeric D-amino acid oxidase from Rhodotorula gracilis. *Enzyme Microb*
Technol **29**, 407–412 (2001).

79. L. Frattini, E. Rosini, L. Pollegioni, M. S. Pilone, Analyzing the d-amino acid content
in biological samples by engineered enzymes. *Journal of Chromatography B* **879**,
3235–3239 (2011).

80. D. Punzo, F. Errico, L. Cristino, S. Sacchi, S. Keller, C. Belardo, L. Luongo, T. Nuzzo,
R. Imperatore, E. Florio, V. De Novellis, O. Affinito, S. Migliarini, G. Maddaloni, M.
J. Sisalli, M. Pasqualetti, L. Pollegioni, S. Maione, L. Chiariotti, A. Usiello, Age-
Related Changes in d-Aspartate Oxidase Promoter Methylation Control Extracellular
d-Aspartate Levels and Prevent Precocious Cell Death during Brain Aging. *The*
Journal of Neuroscience **36**, 3064–3078 (2016).

1220