Chromatin remodeling with combined FACT and BET inhibition disrupts oncogenic transcription in Diffuse Midline Glioma

Authors:

Holly Holliday[1,2]*, Aaminah Khan[1,2]*, Anahid Ehteda[1,2], Samuel E. Ross[3], Nisitha Jayatilleke[1], Anjana Gopalakrishnan[1], Xinyi Guo[1], Jie Liu[1], Hieu Nguyen[1], Alvin Lee[1], Yolanda Colino Sanguino[1,2], Marcel E. Dinger[3], Chelsea Mayoh[1,2], Benjamin S. Rayner[1,2], Maria Tsoli[1,2]**, David S. Ziegler[1, 2, 4]**

* ** equal contribution

Affiliations:

1. Children's Cancer Institute, Lowy Cancer Research Centre, UNSW Sydney, Sydney, NSW, Australia
2. School of Clinical Medicine, UNSW Sydney, Sydney, NSW, Australia
3. School of Life and Environmental Sciences, University of Sydney
4. Kid's Cancer Centre, Sydney Children's Hospital, Randwick, NSW, Australia

Keywords: FACT; BRD4; BET proteins; Diffuse Midline Glioma; DMG; DIPG; H3K27M; Epigenetics; Chromatin; Histones; CBL0137; JQ1; BET inhibitors; Epigenetic therapies; Pre-clinical

Running title: BETing on FACT in DMG

Corresponding authors: Professor David S. Ziegler, Kids Cancer Centre, Sydney Children's Hospital, High St, Randwick 2031 Australia, +61 (02) 9382 1730, d.ziegler@unsw.edu.au

Declaration of Interest statement: DSZ declares consulting/advisory board fees from Bayer, Astra Zeneca, Accendatech, Novartis, Day One, FivePhusion, Amgen, Alexion, and Norgine and research support from Accendatech. The remaining authors have declared that no competing interests exist.

Abstract

Aberrant epigenetic regulation is a hallmark of Diffuse Midline Glioma (DMG), an incurable pediatric brain tumor. The H3K27M driver histone mutation leads to transcriptional dysregulation, indicating that targeting the epigenome and transcription may be key therapeutic strategies against this highly aggressive cancer. One such target is the Facilitates Chromatin Transcription (FACT) histone chaperone. We found FACT to be enriched at developmental gene promoters, coinciding with regions of open chromatin and binding motifs of core DMG regulatory transcription factors. Furthermore, FACT interacted and co-localized with the Bromodomain and Extra-Terminal Domain (BET) protein BRD4 at promoters and enhancers, suggesting functional cooperation between FACT and BRD4 in DMG. *In vitro*, a combinatorial therapeutic approach using the FACT inhibitor CBL0137, coupled with BET inhibition revealed potent and synergistic cytotoxicity across a range of DMG cultures, with H3K27M-mutant cells demonstrating heightened sensitivity. These results were recapitulated *in vivo*, significantly extending survival in three independent orthotopic PDX models of DMG. Mechanistically, we show that CBL0137 treatment decreased chromatin accessibility, synergizing with BET inhibition to disrupt transcription, silencing several key oncogenes including *MYC, PDGFRA* and *MDM4*, as well as causing alterations to the splicing landscape. Combined, these data highlight the therapeutic promise of simultaneously targeting FACT and BRD4 in DMG, proposing a novel strategy for combating this devastating pediatric brain tumor.

Introduction

Diffuse midline glioma (DMG) is a highly aggressive brain tumor predominantly diagnosed in children under 10 years old. Recently classified as 'DMG K27-altered' by the 2021 World Health Organization, this designation recognizes the characteristic epigenetic dysregulation that arises from either the recurrent lysine-to-methionine driver mutations in Histone H3.1 and H3.3 genes (H3K27M), or overexpression of the oncohistone mimic EZHIP [1-7]. Both H3K27M and EZHIP inhibit Polycomb Repressive Complex 2 (PRC2) activity, resulting in a drastic reduction of Histone H3K27 trimethylation (H3K27me3), increased H3K27 acetylation (H3K27ac) and DNA hypomethylation. This culminates in an aberrant epigenetic landscape permissive to heightened transcriptional activity [8-14].

While direct therapeutic targeting of H3K27M has not been possible, targeting of epigenetic and transcription regulators has shown promise in several pre-clinical models of DMG (reviewed in [15, 16]). Among these regulators, the histone chaperone Facilitates Chromatin Transcription (FACT) comprising of two protein subunits; SSRP1 and SPT16, has garnered attention [17-19]. FACT plays a pivotal role in maintaining chromatin integrity by facilitating nucleosome turnover during transcription, DNA replication and repair. This chaperone is abundantly expressed in cancer cells, making it an attractive therapeutic target [20, 21].

The curaxin compound CBL0137 indirectly inhibits FACT by intercalating with DNA and trapping FACT on chromatin (c-trapping), effectively depleting the functional pool of this important chaperone [22-24]. We and others have demonstrated CBL0137 to be active against multiple tumor types, including neuroblastoma, glioblastoma, medulloblastoma, and diffuse intrinsic pontine glioma (DIPG)/DMG [19, 25-29]. This growing body of preclinical work has led to the advancement of CBL0137 to a Phase I/II clinical trial for pediatric patients with solid tumors, including DMG (NCT04870944). However, the aggressive nature of DMG necessitates combination therapies to address therapeutic resistance and reduce effective therapeutic doses. While initial studies combining CBL0137 with the pan-HDAC inhibitor panobinostat has shown promise against DMG pre-clinically [19], concerns over panobinostat's high toxicity, and limited efficacy, led to its withdrawal from clinical studies and highlights the need for alternative drug combinations [30].

Disruption of transcription has proven effective against many 'transcription addicted' malignancies, often achieved by targeting Bromodomain and Extra-Terminal Domain (BET) proteins such as BRD4 [31]. BRD4 recruits the transcriptional machinery to hyper-acetylated

chromatin at super-enhancers [32, 33], making it an attractive target against DMG, as

demonstrated in the preclinical setting ([34-39] and reviewed in [40]). JQ1, among other BET

inhibitors, has demonstrated pronounced efficacy in DMG, through transcriptional disruption

leading to cytostatis [35]. Subsequently, the clinically relevant BET inhibitor trotabresib (CC-

90010) has progressed to clinical trials for adults and children with brain cancer

(NCT04047303, NCT04324840, NCT03936465). However, modest anti-tumor activity

observed in adult monotherapy BET inhibitor trials necessitates the identification of

synergistic combinations [40].

Here we describe the promising efficacy of FACT and BET inhibition as an epigenetic

combination treatment in cell line and animal models of DMG. Mechanistically, this

epigenetic intervention induces changes to chromatin structure and transcriptional profiles.

Moreover, FACT inhibition with CBL0137 leads to chromatin closure at gene promoters,

which when combined with BET inhibition, results in widespread changes to the

transcriptome. Several key oncogenes vital for DMG growth, such as *MYC, PDGFRA,* and

MDM4 are directly regulated by FACT and BRD4, and effectively silenced by dual inhibition

of these targets. These findings suggest a new strategy for inhibiting multiple oncogenic and

otherwise 'untargetable' pathways in DMG.

Results

FACT and BRD4 are overexpressed, interact, and co-localize at active regulatory
elements in DMG

Our earlier work identified FACT inhibition with CBL0137 to be a promising epigenetic

therapy against DMG [19]. In order to further elucidate the genes bound and potentially

regulated by FACT, and its established interaction with the H3K27M oncohistone [19, 41],

we sought to map the genome-wide binding sites of the FACT subunit SPT16 in H3.3K27M-

mutant HSJD-DIPG007 cells using Cleavage Under Targets and Release Using Nuclease

(CUT&RUN). We identified 1,095 SPT16 peaks common to two independent replicates,

which were over-enriched at promoters of genes involved in migration, morphogenesis, and

development (FDR < 0.05) (**Fig 1A-B and Table S1A**). Furthermore, motifs for

developmental and epithelial-to-mesenchymal transition (EMT) transcription factors (TFs)

were enriched at SPT16 binding sites, including master regulators of DMG and

oligodendrocyte development TCF12 (p = 4.819e-14) and OLIG2 (p = 4.281e-9) [42, 43],

and SNAI1 (p = 3.58e-8), which is known to promote an invasive phenotype in malignant

gliomas [44, 45] (**Fig 1C**). This indicates that FACT facilitates transcription driven by these
critical transcription factors.

We next analyzed SPT16 occupancy at different regulatory elements and separately
determined the accessibility of these regions using Assay for Transposase-Accessible
Chromatin using sequencing (ATAC-seq). H3K27M and H3K27ac are known to incorporate
into heterotypic H3K27M-H3K27ac nucleosomes, which are bound by acetyl-binding BET
proteins at enhancer elements [36, 46], prompting us to analyze FACT colocalization with
these nucleosomes. Interestingly, SPT16, H3K27M and BRD4, were enriched at accessible
transcription start sites (TSSs), promoters marked by H3K4me3, and enhancers marked by
H3K27ac. Conversely, these proteins were absent from inaccessible H3K27me3-marked
heterochromatin (**Fig 1D and Fig S1A**). Furthermore, SPT16 was enriched at BRD4 binding
sites, and vice versa, suggesting that FACT and BRD4 occupy the same chromatin regions
(Fig 1E and Fig S1B).

In confirmation of the physical interaction between these proteins, an antibody targeting
BRD4 co-immunoprecipitated both FACT subunits (SSRP1 and SPT16) in DMG nuclear
extracts **(Fig 1F)**. However, BRD4 was absent in FACT IPs, which may be due to epitope
masking, where the IP antibody obstructed the surface of the complex required for its
interaction with BRD4, a common issue in co-IP assays [47]. FACT and BRD4 were in
complex with H3K27M and H3K27ac, in agreement with previous reports by us and others
demonstrating that FACT binds to H3K27M [19, 41] **(Fig 1F).** These biochemical and
genomic results imply that both FACT and BRD4 bind to, and play a role in the oncogenic
functioning of H3K27M in DMG. We next used CUT&RUN to further elucidate the interaction
of these two proteins, finding that SPT16 and BRD4 occupied unique and overlapping
genes, with 190 genes in common, which is a significant 24-fold over enrichment compared
to expectations by chance (hypergeometric p-value = 7.98e-212) **(Fig 1G and Table S1A).**
Genes occupied by both SPT16 and BRD4 were involved in transcription and chromatin
regulation **(Fig S1C),** including the tumor stem cell master regulator *SOX2* [48], the p53
negative regulator *MDM4* [49], and the catalytic subunit of the SWI/SNF chromatin
remodeling complex *SMARCA2* [50] **(Fig 1H)**.

Since DMG is heavily driven by epigenetic and transcription dysregulation, we hypothesized
that FACT and BET proteins, as in other cancers [19, 20, 51-54], would be upregulated in
DMG. Consistently, *SSRP1*, *SUPT16H*, and *BRD4* were overexpressed in DMG/High Grade
Glioma (HGG) patients from the ZERO childhood cancer program compared to normal fetal
brain (NFB) tissue [55, 56] with (fold-change 1.7 - 4.2; p<0.0001) **(Fig S1D).** These genes

were also upregulated (fold-change 1.5 – 2.7; p<0.0001) in H3K27M DMG cell lines
compared to non-malignant cell lines from the Childhood Cancer Model Atlas [57] **(Fig S1E).**
Furthermore, SSRP1 and BRD4 proteins were upregulated in a panel of DMG cell lines
compared to normal astrocytes, with an average of 19-fold and 8-fold increase in protein
expression for BRD4 and SSRP1 respectively (**Fig S1F**). In summary, FACT and BRD4 are
overexpressed, interact, and co-localize at transcriptionally active chromatin, along with
H3K27M, in the vicinity of key developmental genes required for DMG growth. This
prompted us to test the combination of FACT and BET inhibition as a therapeutic strategy.

Combined FACT and BET inhibition synergistically decrease DMG cell proliferation

Treatment with either of the BET inhibitors JQ1 or trotabresib resulted in decreased
proliferation of primary DMG cells, with 72h IC_{50} values ranging from 1-50 µM and 0.6-7 µM,
respectively, across a range of DMG cultures (**Fig 2A-B and Table S2-3**). Subsequent
investigation into the impact of FACT or BRD4 inhibition on DMG isogenic cell lines, with or
without the H3K27M mutation, revealed significantly increased sensitivity to single-agent
CBL0137, JQ1, and trotabresib in cells bearing the H3K27M mutation. Notably, significant
differences were observed in both IC_{50} and area under the curve (AUC) values between
wild-type and H3K27M mutant cultures, with a 5-fold increase in sensitivity observed in
H3K27M mutant compared to H3K27-KO (wild-type) DMG cells for CBL0137 and JQ1, and a
4-fold increase for trotabresib treatment (**Fig 2C, Fig S2A-C and Table S4**).
178
Next, we assessed the effect of combined inhibition of both FACT and BRD4. We found
potent efficacy of the combination of CBL0137 and JQ1/trotabresib in H3K27M mutant cell
lines, with no synergy observed in the KO cells (**Fig 2D-E**). This synergistic effect of the
combination therapy on cell proliferation was similarly observed in short term cytotoxicity
assays across three patient-derived DMG cell cultures, SU-DIPGVI, HSJD-DIPG007,
RA055, all H3K27M mutant (**Fig 2F and Fig S2D-G**). The combinations also led to
synergistic inhibition of colony formation, at concentrations of each compound ranging
between 25-50 nM for SU-DIPGVI, and 100-200 nM for HSJD-DIPG007 and RA055 (**Fig 2G**
and Fig S2H-K). Together, these data highlight the potential of combined FACT and BRD4
targeted inhibition as an effective strategy in H3K27M mutant DMG.
189
190 **Combined FACT and BET inhibition causes chromatin condensation at gene**
191 **promoters in DMG.**
192

Given that CBL0137 and BET inhibitors each target known epigenetic and transcriptional
regulators [17-19, 58], we next sought to determine the effect of these drugs on chromatin
structure and transcriptional regulation. To measure chromatin accessibility, ATAC-seq was
performed in two H3.3 K27M DMG cell lines (HSJD-DIPG007 and SU-DIPGVI) treated with
either CBL0137 or JQ1 as monotherapies or in combination after 4 hours to examine the
early and direct changes to chromatin at doses previously used in other mechanistic studies
[20, 29, 35, 59, 60]. Combinatorial treatment of DMG with CBL0137 and JQ1 led to a
decrease in ATAC-seq signal at transcription start sites (TSSs) (**Fig 3A)**, consistently
observed across independent replicates from both cell lines tested (**Fig S3A**). This effect
was largely driven by CBL0137, with Principal Component Analysis (PCA) revealing that the
CBL0137 treated samples clustered closely with combination treated samples (**Fig S3B**).
204
We next examined chromatin accessibility at DNA with SPT16 and BRD4 CUT&RUN peaks
in HSJD-DIPG007 cells, finding that both drugs alone, and in combination, decreased
chromatin accessibility at these genomic regions (**Fig 3B**). This suggests that FACT and
BRD4 are important for maintaining open chromatin, and that inhibition of these factors with
CBL0137 and JQ1 leads to chromatin closure/condensation at these sites. Further,
differential accessibility analysis revealed that combination treatment resulted in ~15-fold
more closed chromatin regions compared to open regions in both cell lines tested (SU-
DIPGVI: 41,457 regions closed vs. 2,620 regions opened; HSJD-DIPG007: 11,704 regions
closed vs. 828 regions opened, FDR<0.05, fold-change>2X) (**Fig 3C, Fig S3C-D, and Table
S1B**). There was a significant degree of overlap of the regions closed by the combination
treatment in both the models **(Fig 3D)** (hypergeometric p-value = 0.008). These regions
were enriched at proximal gene promoters (**Fig 3E**) with their annotated genes involved in
development, proliferation, and histone modification (**Fig 3F**). This indicates that combination
treatment leads to chromatin condensation at the promoters of genes involved in pathways
critical for DMG tumorigenesis. For example, the combination led to chromatin condensation
at the promotor of *NES* (**Fig 3G**)*,* encoding the neural progenitor marker Nestin, which is
essential for glioma development [61, 62]. Remarkably, combination treatment also
decreased chromatin accessibility near the promoters of *SUPT16H* (SPT16) and *BRD4* –
encoding the protein targets of CBL0137 and JQ1 (**Fig 3G**).
224
H3K27M disrupts the balance of repressive H3K27me3 and active H3K27ac histone
modifications, decreasing H3K27me3 and increasing H3K27ac [8]. We have previously
shown that CBL0137 can increase global H3K27me3 levels after 24h [19]. In order to assess
whether the immediate and direct chromatin changes in response to drug treatment related
to changes in trimethylation we next tested the effect at the shorter timepoint of 4h. Despite

evidence that FACT and BRD4 interact with H3K27M on chromatin (**Fig 1E-F**), inhibition of

these factors did not consistently affect global H3K27me3 or H3K27ac levels at this early

timepoint, observed through western blotting or ELISA assays (**Fig S3E-G**). Furthermore,

CBL0137 treatment did not significantly alter H3K27me3 levels or distribution at individual

gene loci using CUT&RUN (**Fig S3H-I**), together indicating that chromatin remodeling

induced by CBL0137 is not reliant on H3K27me3 restoration within this early timeframe.

These results are in line with the relatively slow turnover rates (3 days) for H3K27me3 [63].

CBL0137 and JQ1 cause widespread gene expression changes

To investigate the impact of chromatin condensation of CBL0137 and JQ1 treatment on

transcription, we performed RNA-seq on matching cells from the ATAC-seq experiments.

Differential gene expression analysis revealed that the combination treatment led to the

largest degree of gene expression changes in both cell lines (FDR<0.05, fold-change>1.5X),

with more genes becoming downregulated than upregulated (**Fig 4A and Table S1C**) when

comparing each treatment to the DMSO control. Consistently, multidimensional scaling of

RNA-seq samples revealed distinct clustering based on the drug treatments, with the

combination treatment responsible for the most variation in gene expression between the

cohorts (**Fig S4A**).

Analyses of the differentially expressed genes common to both DMG cell lines (**Fig 4B**)

indicated that genes downregulated in response to combination treatment were involved in

several processes critical for the sustained growth and proliferation of cancer cells, such as

RNA processing (RNA modification, tRNA modification, rRNA modification), ribosome

biogenesis, and DNA repair (**Fig S4B and Fig 4C**). Conversely, combination treatment led to

upregulation of stress (e.g. regulation of transcription in response to stress and the unfolded

protein response) and transmembrane transport pathways, reflecting an acute stress

response to the drug combination (**Fig S4B and Fig 4C**).

CBL0137 and JQ1 combination treatment represses transcription of key oncogenes

We next examined the relationship between chromatin occupancy and the drug-induced

alterations to chromatin accessibility and gene transcription through integration of the

CUT&RUN, ATAC-seq and RNA-seq datasets. Most of genes bound by SPT16 and BRD4

had decreased expression when these proteins were simultaneously inhibited, suggesting

that FACT and BRD4 directly activate a subset of the genes they occupy. This effect was

primarily driven by JQ1, as the transcriptional profile of the JQ1-treated samples resembled

that of the combination-treated samples (**Fig 5A**). As expected, the majority of genes with decreased promoter accessibility upon combination treatment also had decreased transcription with each single agent, exhibiting an intermediate level of expression between that of the control and combination treatment (**Fig 5B**). Together, this indicates that dual targeting of FACT and BRD4 leads to compaction of chromatin and to transcriptional silencing at specific loci.

Among the silenced genes were several key oncogenic drivers, for example *MYC, PDGFRA* and *MDM4*. *MYC* is an established transcriptional target of both FACT and BRD4 [26, 58]. Accordingly, we found SPT16 and BRD4 to be localized to the *MYC* promoter, marked by H3K4me3 and H3K27ac histone modifications (**Fig 5C**). Furthermore, this same region had decreased chromatin accessibility in combination-treated cells (**Fig 5C)** with a concomitant downregulation of both *MYC* mRNA (**Fig 5D and Fig S5A**) and protein expression following 24 hours (**Fig 5E and Fig S5B**). *PDGFRA* encodes another prominent super-enhancer associated glioma oncogene [35, 36]. SPT16, and BRD4 albeit to a lesser degree, were similarly localized to the *PDGFRA* promoter and putative upstream enhancer, with condensed chromatin at these loci, and repression on both the mRNA and protein level upon combination treatment (**Fig 5C-E and Fig S5A-B**). Another silenced oncogene was *MDM4*, which is a p53 negative regulator with additional p53-independent oncogenic roles [49, 64] (**Fig 5C-E and Fig S5A-B**). Interestingly, p53 protein was only upregulated by CBL0137, and only in the context of wildtype p53 HSJD-DIPG007 cells, but not in p53-mutant SU-DIPGVI cells **(Fig S5D-E)**. The same patterns of FACT/BRD4 co-localization, chromatin compaction, and decreased transcription were observed for *PTCH1,* an important member of the hedgehog signaling pathway shown to be elevated in DMG [45], and a consistent PTCH1 protein decrease was observed in SU-DIPGVI cells (**Fig S5E-G). Finally, *LARP1, an* RNA binding protein and mTOR effector involved in ribosome biosynthesis [65], and *KLHDC8A,* a super-enhancer driven glioma oncogene involved in hedgehog signaling [66], had similar epigenetic and transcriptional repression, however, these did not manifest in a significant decrease in protein expression (**Fig S5E-G).**

CBL0137 and JQ1 alter the splicing landscape in DMG

Given the downregulation of RNA processing genes in CBL0137 and JQ1 combination-treated cells (**Fig 4C**), and the fact that chromatin is known to regulate splicing through a co-transcriptional process [67], we next examined splicing in our RNA-seq experiments. Treatment with CBL0137 or JQ1 as single agents demonstrated limited differential splicing. However, compared to control cells, combination treatment led to hundreds of differential

splicing events with an average of 800 and 2000 events in HSJD-DIPG007 and SU-DIPGVI
cells, respectively (**Fig S6A**), with a proportion of these events occurring within the same
gene, resulting in approximately 400 unique genes in HSJD-DIPG007 cells and 1000 unique
genes in SU-DIPGVI cells (**Fig S6B and Table S1D**)**.** The most common form of alternative
splicing was intron retention, equating to 40-50% of events (**Fig S6C**). Next, we investigated
the potential regulation of splicing by drug-induced changes to chromatin architecture. We
observed a statistically significant overlap (hypergeometric p-values = 1.64e-10 for HSJD-
DIPG007 and 3.14e-56 for SU-DIPGVI) in genes which had both closed chromatin and
differential splicing, equating to 81 shared genes in HSJD-DIPG007 and 430 shared genes
in SU-DIPGVI cells. (**Fig S6D**). An example of intron retention is shown at *SPTBN2* (**Fig**
S6E). This gene also had chromatin condensation and reduced expression (**Fig S6E-F**) and
encodes for β-III spectrin which has been implicated as a driver in multiple cancer types [68-
70]. Interestingly, non-coding transcripts corresponding to retained introns for this gene have
been documented (**Fig S6G**) [71]. These findings indicate that epigenetic modulation with
CBL0137 and JQ1 disrupts the transcriptional splicing landscape in DMG.
319
320 **CBL0137 and JQ1 modulate the interferon response in DMG and in macrophages**
321
While our earlier experiments assessed the early and direct epigenetic effects of drug
treatment in DMG, we next assessed expression changes after an extended 48h treatment.
In keeping with the recognized ability of CBL0137 to invoke an interferon response [72], we
show that CBL0137 treatment for 48h elevated expression of interferon stimulated genes
(ISGs) in HSJD-DIPG007 cells, including statistically significant increases in *IRF1*, *OAS1*
and the expression of antigen presentation genes (*HLA-A, HLA-B, HLA-C)* (**Fig S7A**), whilst
concomitantly increasing expression of *IFNα, IFNβ, IFNγ*, and *IL-6* (**Fig S7B**). BRD4 has
been shown to play a role in the transcriptional regulation of ISGs [73]. Here, we found that
inhibition of BRD4 in DMG modulates both ISGs and cytokine mRNA expression, which is
largely recapitulated in DMG exposed to a combination of CBL0137 and JQ1 (**Fig S7A-B**).
332
Given that macrophages are the most prominent immune cell type in DMG [15], and that
macrophage polarization into M1 and M2 states is epigenetically regulated [74]**,** we next
assessed the ability of CBL0137 and JQ1 to modulate the inflammatory response within
cultured macrophages. Compared to untreated controls, macrophages exposed to CBL0137
demonstrated significantly increased expression in the pro-inflammatory M1 macrophage
markers *IL-1β, p53* and *iNOS* (**Fig S7C**) and increased expression of M2 markers *IRF3* and
HIF-1α (**Fig S7D**). Conversely, macrophages exposed to JQ1 responded with upregulation
of the M1 markers *IL-1β* and *IRF8*, with no significant increase in M2 marker expression. The

combination of CBL0137 and JQ1 resulted in the increased expression of the M1 markers
IL-1α, IL-1β, p53 and *IRF5*, whilst also increasing the M2 marker *HIF-1α*. Together, these
data indicate that this combinatorial therapeutic approach has the potential to positively
modulate the immune microenvironment.

Dual inhibition of FACT and BRD4 enhances survival in orthotopic models of DMG

To evaluate therapeutic efficacy, we utilized three orthotopic xenograft models using distinct
patient-derived DMG cells: SU-DIPGVI-LUC, RA055, and HSJD-DIPG007. These models
were chosen for their ability to faithfully replicate the diffuse infiltration characteristics
observed histologically in DMG tumors.

In the SU-DIPGVI-LUC model, single-agent treatments with CBL0137 and JQ1 exhibited
either negligible or limited anti-tumor effects, respectively. However, the combination
regimen, which was well-tolerated with no observed weight loss, significantly extended
survival compared to single-agent treatments and vehicle cohorts by delaying tumor growth,
as seen by xenogen imaging (median survival: vehicle: 73 days, CBL0137: 76 days, JQ1: 78
days, CBL0137/JQ1: 95.5 days; **Fig 6A-B**, **Fig S8A** and **Table S5**). Immunohistochemical
analysis of brain tissue collected following 4 weeks of combination treatment revealed a
significant reduction in cell proliferation, as evidenced by decreased Ki67 staining, and a
concurrent reduction in H3K27ac staining (**Fig 6C-D**).

These promising results were recapitulated in the highly aggressive RA055 model,
characterized by a median survival of 46 days in untreated mice post intracranial injection.
Treatment with the combination of CBL0137 and JQ1 significantly enhanced survival, with
approximately one-third of treated mice surviving until the experimental endpoint of 201 days
(**Fig 6E**, **Fig S8B** and **Table S6**). Consistent with the SU-DIPGVI-LUC model, cell
proliferation and H3K27ac immunohistochemistry staining were significantly reduced with
combination treatment compared to single-agent and untreated tissue samples (**Fig 6F and**
Fig S8C). We also observed a significant reduction in MYC and MDM4 staining in PDX
tumors treated with the combination, validating the *in vitro* mechanistic findings in an
independent *in vivo* model.

Extension in survival with combined FACT and BRD4 inhibition was also evident in the
HSJD-DIPG007 model, with the median survival of the combination-treated cohort extended
to 93.5 days compared to 70.5 days in the vehicle group (**Fig S8D-E** and **Table S7**). Taken
together, this combination strategy is effective against multiple aggressive DMG models.

Discussion

Despite hundreds of clinical trials, survival outcomes for DMG patients have not changed in
over three decades [15]. The current standard-of-care, radiotherapy, only delays tumor
progression and the majority of children still survive less than one year from diagnosis. Since
the discovery of the H3K27M mutation in 2012, efforts to target multiple layers of epigenetic
regulation, including DNA methylation [12], histone modification [75-78], chromatin readers
[35, 36], chromatin remodelers [79-81], and transcription machinery [35, 82], have been
ongoing. Due to DMG's aggressive and heterogenous nature, a monotherapy approach is
unlikely to suffice. Moreover, with multiple aberrant pathways, and undruggable oncogenic
drivers, even combination therapies that affect one or two pathways are unlikely to be
effective. Here we have uncovered a novel mechanism-anchored epigenetic combination
therapy - FACT and BET inhibition - which leads to simultaneous disruption of multiple
oncogenic pathways and demonstrates potent pre-clinical activity against DMG.

FACT disassembles and reassembles nucleosomes during cellular processes that require
DNA access, such as during transcription or replication. This overcomes the physical
hindrance of nucleosome-compacted DNA, allowing RNA Pol II or the replication machinery
to travel through chromatin unimpeded [22, 23]. Our findings indicate that FACT maintains
open chromatin at promoters and enhancers, supporting transcription driven by
developmental TFs (TCF12, OLIG2, SNAI1) in DMG. This parallels a recent study showing
FACT's role in maintaining open chromatin at regulatory elements to promote transcription
by core pluripotency TFs in embryonic stem cells [83].

BRD4 binds to acetylated chromatin via its tandem bromodomains and recruits positive
transcription elongation factor (P-TEFb) to active enhancers, which releases paused RNA
pol II, leading to transcriptional initiation and elongation [84]. Our findings suggest that FACT
and BRD4 cooperate at promoters and enhancers to drive transcription of multiple genes
vital for the growth and development of DMG, with blockade of these synergistically
disrupting the transcriptional landscape, both in terms of transcript abundance and splicing.
Combined targeting of FACT and BRD4 led to chromatin compaction and transcriptional
silencing of several notable oncogenes. For example, *MYC, PDGFRα*, and *KLHDC8A* are all
super enhancer-activated oncogenes involved in gliomagenesis [12, 66, 85]. Another
downstream target was *MDM4*, a negative regulator of p53, suggesting this treatment could
potentially restore p53 function in p53-WT DMGs [49]. *PTCH1* and *KLHDC8A* are both

involved in hedgehog signaling [66], which is notable given the hedgehog responsive cell-of-
origin of DMG [86], and that both CBL0137 and JQ1 synergistically antagonize oncogenic
hedgehog signaling in other pediatric brain cancers [87]. In sum, this drug intervention
simultaneously disarms multiple epigenetic-driven oncogenes in DMG.

The combination of CBL0137 with both BET inhibitors had higher efficacy against H3K27M
cells compared to isogenic cells lacking H3K27M. This could be explained by two
hypotheses: i) The drugs inhibit the function of H3K27M by targeting its co-factors (FACT
and BRD4), or ii) H3K27M's downstream epigenetic dysregulation sensitizes the cells to
FACT and BRD4 inhibition. The former is supported by the interaction and chromatin
localization between FACT and BRD4 with H3K27M, which is strengthened by an IP-mass
spectrometry study identifying SPT16 and SSRP1 as binding partners of H3K27M in DMG
mitotic extracts [41]. However, we did not observe marked changes in H3K27me3 levels at
early timepoints after CBL0137 treatment, indicating that H3K27M's ability to inhibit PRC2
was not impacted in a short time frame. It is possible that sustained treatment is required to
observe this effect, as we have previously shown that 24 hours of treatment with CBL0137
leads to increased global H3K27me3 [19]. Given the slow turnover rate of H3K27me3 (3
days) [63], extended timepoints with lower non-cytotoxic doses may be required to observe
changes in H3K27me3 patterning by CUT&RUN. In the second scenario, we propose that
the hypomethylated chromatin creates a dependency on FACT and BRD4 to maintain open
chromatin and high levels of transcription (see graphical abstract). CBL0137 and JQ1
treatment then leads to chromatin compaction at transcription start sites, disrupting the
expression of thousands of genes. This is analogous to recent studies targeting the
SWI/SNF chromatin remodeling complex, where the epigenetic reprogramming
characteristic of DMG creates a dependency on this complex to drive pathogenic gene
expression [79-81]. Notably, CBL0137 and JQ1 were still effective, albeit to a lesser degree,
against isogenic DMG cells lacking H3K27M, and have also shown synergy in other H3K27
WT pediatric brain malignancies, such as medulloblastoma and Atypical Teratoid/Rhabdoid
Tumor (AT/RT) cell lines [87], indicating potential broader therapeutic applications in
pediatric brain cancers beyond H3K27M DMG.

The three-dimensional (3D) organization of the genome is an additional layer of epigenetic
regulation and is emerging as a critical contributor to tumorigenesis in epigenetic driven
cancers such as DMG and Ependymoma [42, 88]. Both CBL0137 and BET inhibitors have
known roles in targeting genome topology, disrupting topologically associating domains
(TADs) and chromatin loops to alter enhancer-promoter communication [42, 60]. While this

epigenetic aspect was not addressed in this study, understanding the impact of combination
treatment on 3D genome architecture represents an exciting future direction.

We uncovered a role for combination treatment in regulating mRNA splicing. As splicing is
influenced by chromatin structure, we hypothesize that alternative splicing could arise due to
stalled transcription as RNA pol II attempts to travel through compacted chromatin, leading
to errors caused by the splicing machinery [67]. We propose a mechanism by which
changes to chromatin structure induced by CBL0137 renders the cells susceptible to
transcriptional disruption, which is achieved with the addition of JQ1. However, this
hypothesis warrants further exploration using nascent RNA-sequencing approaches or
mapping RNA pol II rate. Notably, epigenetic therapies are well known for their activation of
the immune response through various mechanisms [89]. Indeed, CBL0137 has been shown
to induce double stranded RNA, triggering a type I interferon response in various non-DMG
cancer models, and this can occur independently of FACT [27, 72, 90]. In our study,
sustained drug treatment revealed a heightened interferon response in DMG and highlights
the need to use immunocompetent models in future work.

The *in vivo* efficacy of the combined FACT and BRD4 inhibition in multiple orthotopic models
of DMG indicates its potential as a promising combinatorial treatment strategy. Employing an
alternative dosing regimen involving an increased dose of CBL0137 (40 mg/kg) paired with a
reduced dose of JQ1 (25 mg/kg) not only proved well-tolerated but also resulted in a
significantly enhanced survival response in highly aggressive DMG models. This
underscores the importance of exploring diverse dosing regimens for the development of
combination therapies.

In summary, our study provides compelling mechanism-based evidence for a promising
novel approach to treat DMG. FACT inhibitors and BET inhibitors are both presently being
tested in Phase I/II clinical trials for pediatric brain tumors as single agents (PEPN2111,
NCT03936465). Our results establish a strong rationale for a combination clinical trial to
improve treatment options for DMG patients.

Materials and Methods

Cell culture
DMG cells were maintained as neurospheres in Tumor Stem Cell (TSM) media consisting of
a 1:1 mixture of DMEM/F12 and Neurobasal medium (Invitrogen) supplemented with

Glutamax, Pyruvate, non-essential amino acids, HEPES buffer, antibiotic/antimycotic, B-27

(Invitrogen), Heparin (0.0002%), EGF (20 ng/mL), bFGF (20 ng/mL), PDGF-AA (10 ng/mL)

and PDGF-BB (10 ng/mL) (StemCell Technologies). SU-DIPGXIII K27M and K27M-KO

isogenic cells were maintained according to [12], adherently on flasks coated with poly-L-

ornithine (0.01%) (Sigma) and laminin (10 µg/mL) (Sigma), in Neurocult NS-A proliferation

media (StemCell Technologies) supplemented with heparin (0.0002%), EGF (20 ng/mL) and

bFGF (10 ng/mL) (StemCell Technologies). All lines were cultured at 37°C and 5% carbon

dioxide. Cells tested negative for mycoplasma contamination and were confirmed to match

their original tumors by STR profiling.

Human monocyte-derived macrophages (HMDM) were a kind gift from Prof. Kerry-Anne

Rye, Cardiometabolic Research Group, UNSW, following isolation and culture as previously

described [91]. In polarisation studies, HMDM were either primed for 48 h with 1 µg/mL

LPS and 20 ng/mL recombinant human IFNγ for the M1 phenotype, 20 ng/mL recombinant

human IL-4 and 20 ng/mL recombinant human IL-13 for the M2 phenotype, or treated for 48

h with 1µM CBL0137 or 1µM JQ1, alone or in combination, and total RNA extracted, and

cDNA generated for qPCR, as detailed below.

Co-immunoprecipitation (co-IP) assay

Co-IP was performed as described in [92]. Briefly, nuclei were isolated from 5 million cells

with hypotonic buffer (20 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, protease inhibitors

(Roche)) for 15 min on ice, followed by addition of IGEPAL (0.5%). Nuclei were collected by

centrifugation (5 min, 1000 *x g*, 4°C) then lysed in non-denaturing lysis buffer (50mM HEPES

pH 7.4, 140 mM NaCl, 3 mM $MgCl_2$, 1 mM EGTA, 1% Triton-X, 10% Glycerol, protease

inhibitors (Roche)). The nuclear lysate was passed 6 times through a 25-G needle to shear

the DNA, then incubated for 15 min on ice. Lysates were cleared by centrifuging at 13,000

rpm (top speed) for 15 min at 4°C to remove insoluble debris. The lysate supernatant was

then incubated with 15 µL protein A/G magnetic beads (Pierce) pre-conjugated with 2.5 µg

antibody (minimum 4 h, 4°C with rotation). Immunoprecipitation was carried out overnight at

4°C on a rotator. Magnetic beads were washed 3x with lysis buffer, and protein was eluted

with 2X Laemmli Sample Buffer diluted in lysis buffer, boiled at 85°C for 10 min, followed by

addition of DTT (100 mM). Western blotting to detect immunoprecipitated proteins was

performed as described below, except that VeriBlot (Abcam) was used for the secondary

antibody at 1:1000.

Western blotting

To extract total protein cells were incubated with RIPA buffer (CST) containing protease and
phosphatase inhibitors (Roche) for 15 min on ice with occasional vortexing, followed by
centrifugation at 13,000 rpm for 15 min at 4°C to remove debris. To extract histones, the
histone extraction kit (abcam) was used according to the manufacturer's protocol. Protein
concentration was determined using a BCA kit (Pierce). Protein samples (20-30 µg RIPA-
soluble proteins, or 3 µg for histones) were prepared with 1X Laemmli SDS loading dye then
boiled for 5 min at 95°C. BioRad 4-20% proTEAN pre-cast gels were used for gel
electrophoresis, and proteins were transferred for 1 h at 100 V onto 0.45 µm PVDF
membranes for RIPA-soluble proteins, or 0.45 µm nitrocellulose membranes for histones.
Membranes were blocked for 1 h in 5% milk diluted in Tris Buffered Saline with 0.01%
Tween-20 (TBS-T). Western blotting was carried out overnight with the indicated antibodies
(Table 1) diluted 5% Bovine Serum Albumin (BSA) in TBS-T. Following washing,
membranes were incubated in horseradish peroxidase- conjugated secondary antibody
(CST) diluted in 5% skim milk/TBS-T (1:2000) for 1 hr at room temperature. Membranes
were incubated with ECL substrate (Pierce) and bands imaged using a Bio-Rad ChemiDoc
imaging system. Densitometry was carried out using Image Lab software, with local
background subtraction, and normalization to the indicated housekeeping gene.
532
533 **ELISA**
534 Histone extracts were isolated as described above and 200 ng was used to analyze global
535 H3K27me3 and H3K27ac levels using ELISA kits according to the manufacturer's protocol
536 (EpiGenTek). Samples were run in technical duplicate wells, and the experiment repeated 3
537 independent times.
538
qPCR
Total RNA was extracted with an ISOLATE II RNA Mini Kit and first-strand cDNA generated
using a SensiFAST cDNA Synthesis Kit, with quantitative real-time PCR (qRT-PCR)
performed through the use of the SensiFAST SYBR No-ROX Kit (Meridian Bioscience),
using specific primers sequences (Table 2), synthesized by Integrated DNA Technologies.
Relative mRNA expression of genes of interest were normalized against the corresponding
house-keeping genes, using the delta-delta Ct method [93].
546
CUT&RUN
HSJD-DIPG007 neurospheres were collected by centrifugation at 800 rpm for 3 min and
dissociated into single cells with 1 mL Accutase (Sigma) for 5-10 minutes with occasional
pipetting, diluted with 5 mL TSM-B, and passed through a 40 µm cell strainer. Cells were
centrifuged at 1,200 rpm for 5 min and resuspended in complete media. Single cell

suspensions were counted using a haemocytometer and cell viability was confirmed to be
over 95%. CUT&RUN was performed using the EpiCypher CUTANA CUT&RUN kit
according to the manufacturer's instructions. Briefly, 500,000 viable cells per reaction were
washed and permeabilized with 0.01% Digitonin, which was determined to be the minimum
concentration for effective permeabilization. Cells were incubated with 0.5 µg antibody, or a
1:50 dilution for SPT16 (Table 1), on a nutator overnight at 4°C. Following washing, pAG-
MNase was added to the reactions and $CaCl_2$ was used to activate antibody-bound MNase
for targeted chromatin digestion and release of DNA fragments. The entire volume of purified
DNA (ranging from 1-10 ng) was used as input for NGS library preparation, according to the
EpiCypher CUTANA Library Prep Kit instructions. Library concentration was determined
using a Qubit Fluorometer (HS dsDNA) (Thermo) and library size determined using a 4200
Tapestation system on a high sensitivity D1000 ScreenTape (Agilent). Resulting libraries
were sequenced with 50-150bp paired-end reads, with a minimum of 7M reads per sample.
Two biological CUT&RUN replicates were performed.
566
CUT&RUN analysis
Quality control analysis of CUT&RUN reads was performed using fastQC (v0.11.9). Adapter
trimming was performed using cutadapt (v4.2) with parameters: --error-rate 0.1 --times 1 --
overlap 3 --action trim --minimum-length 17 --pair-filter any. Reads were aligned using
bowtie2 (v2.5.0) with -k 1 against the GRCh38 human reference genome. BAM files were
sorted, and optical duplicates marked using picard (v2.27.5) with parameters: --
DUPLICATE_SCORING_STRATEGY SUM_OF_BASE_QUALITIES --
OPTICAL_DUPLICATE_PIXEL_DISTANCE 100 --VALIDATION_STRINGENCY SILENT.
Marked duplicates were filtered and removed using samtools (v1.12). Peak calling was
performed using MACS2 (v2.2.7.1) using IgG as the control. Narrow peaks parameters: --
keep-dup all -p 1e-5 and broad peak using parameters: --keep-dup all -q 0.05 --broad --
broad-cutoff 0.05. BigWig coverage files were generated using deeptools bamCoverage
(v3.5.1) with parameters: --binSize 10 --normalizeUsing BPM --effectiveGenomeSize
2913022398. Broad peak settings were used for H3K27M and H3K27me3, and narrow
peaks for SPT16, BRD4, H3K27ac, H3K4me3. Peaks from biological replicates were
overlapped using BedTools [94].
583
BigWig coverage was visualized using DeepTools [95] at the following genomic regions:
hg38 Ensembl genes, H3K27me3 CUT&RUN peaks, H3K27me3 CUT&RUN peaks, SPT16
CUT&RUN peaks and BRD4 CUT&RUN peaks (all from this study) and H3K27ac ChIP-seq
peaks (GSE162976[42]), with blacklisted CUT&RUN regions excluded [96]. Integrated
genomics viewer (IGV) [97] was used to visualize BigWig coverage and MACS peak calls at

individual loci. Peaks were annotated using ChIPSeeker [98] and ClusterProfiler [99] was
used for Gene Ontology over-representation analysis of the annotated genes. To determine
if the overlap between the SPT16 and BRD4 associated genes was significantly over-
enriched a hypergeometric test was performed using the following parameters: number of
successes k = 190 (overlapping genes) sample size s = 478 (BRD4 peak genes common to
n=2 reps) number of successes in the population M = 1093 (SPT16 peak genes common to
n=2 reps) population size N = 67,319 (number of genes in hg38), where expected = (s*M)/N.
Transcription factor motif analysis was performed using the Cistrome SeqPos motif tool
[100]. Differential binding analysis for H3K27me3 comparing between DMSO and CBL0137
treated cells was carried out using CSAW as described [101], filtering on genomic windows
5-fold higher than background, with a significance threshold of FDR < 0.05 (n=2).
600

ATAC-seq
Neurospheres were treated for 4 h with CBL0137 and JQ1 (HSJD-DIPG007 1 µM CBL0137
1 µM JQ1; SU-DIPGVI 2 µM CBL0137 1 µM JQ1). DMSO was added to the control and
single drug treatments to equal concentration in all conditions (maximum 0.03% DMSO).
Transposition was performed as described in the OMNI-ATAC-seq protocol [102]. Briefly,
single cell suspensions from neurospheres were prepared as described for CUT&RUN and
washed in ice-cold PBS and cell viability was confirmed to be over 95%. Nuclei from 50,000
viable cells were isolated, permeabilized and transposed with 2.1 µL Tn5 (Illumina) for 30
min with mixing (1000 rpm). DNA was then purified, and PCR amplified (7-14 cycles) with
sequencing primers to generate libraries for sequencing. Library concentration was
determined using a Qubit Fluorometers (Thermo) (HS dsDNA) and library size determined
using a 4200 Tapestation system on a high sensitivity D1000 ScreenTape (Agilent). ATAC-
seq libraries were sequenced with a minimum of 50 million paired-end reads per sample and
read length 100bp. ATAC-seq experiments were performed in biological duplicate (HSJD-
DIPG007) or triplicate (SU-DIPGVI).
616

ATAC-seq analysis
ATAC-seq   reads   were   trimmed   with   Trimmomatic
(ILLUMINACLIP:Nextera.fa:2:30:10:8:TRUE   SLIDINGWINDOW:5:20   LEADING:3
TRAILING:3 MINLEN:20 CROP:130 HEADCROP:150). Trimmed reads were mapped with
bowtie2 using default settings [103] and the resulting BAM files deduplicated using
Sambamba markdup [104]. Deduplicated BAM files were then filtered to remove reads with
fragment sizes greater than 100bp and CPM BigWigs were generated using the DeepTools
bamCoverage [95].

BigWig coverage was visualized as heatmaps and profile plots using DeepTools [95] at the
following genomic regions: hg38 Ensembl genes, H3K27me3 CUT&RUN peaks, H3K27ac
ChIP-seq peaks, H3K27me3 CUT&RUN peaks, SPT16 CUT&RUN peaks and BRD4
CUT&RUN peaks (all from this study, except for H3K27ac from GSE162976[42]). PCA
clustering of ATAC-seq average signal over gene bodies was performed using DeepTools.
Differential accessibility analysis was carried out using CSAW as described [101], filtering on
genomic windows 5-fold higher than background. Regions with an FDR<0.05 were
considered significantly different, with 'opened' regions logFC > 1 and 'closed' regions logFC
< -1. Closed regions between the 2 models were overlapped using BedTools [94], annotated
using ChIPSeeker [98] and ClusterProfiler was used for Gene Ontology over-representation
analysis of the annotated genes [99]. BigWig coverage was visualized at individual gene loci
on IGV [97]. A hypergeometric test was used to test if the overlap in 'closed regions'
between the 2 models was significantly over-enriched (k = 6119 (overlap), s = 11704 (closed
in HSJD-DIPG007), M = 41,454 (closed in SU-DIPGXIII), N = 80,894 (number of windows
analyzed), where expected = (s*M)/N.

RNA-seq
Neurospheres were treated with CBL0137 and JQ1 for 4 h and collected for both ATAC-seq
and RNA-extraction. Total RNA was extracted from treated cells using the ISOLATE II RNA
Mini Kit from Meridian Bioscience. RNA concentration and purity was determined using a
NanoDrop Spectrophotometer and integrity measured using a 4200 Tapestation system.
cDNA libraries were prepared using the Illumina Stranded mRNA prep kit. RNA-seq libraries
were sequenced with 100bp paired-end reads, with a minimum of 50 million reads per
sample. Three biological replicates were performed for all RNA-seq experiments.
648

RNA-seq analysis
Illumina paired-end fastq files were aligned to the human genome GRCh38 and gencode
v41 using STAR (v2.7.8a) in two-pass mode with quantMode set to TranscriptomeSAM
[105]. Transcriptome alignment was quantified using RSEM (v1.3.3) rsem-calculate-
expression to obtain raw read counts, and normalized transcripts per million (TPM) and
fragments per kilobase million (FPKM) values [106]. Degust (10.5281/zenodo.3258932) was
used for differential expression analysis, using edgeR for normalization. Genes with low
expression were filtered out using a cutoff of 0.5 counts per million (CPM) in at least 3
samples. Genes with FDR<0.05 and absolute logFC>0.585 (1.5-fold) were considered
significantly differentially expressed. Multi-dimensional Scaling (MDS) clustering was
performed using Degust. ClusterProfiler was used for Gene Ontology over-representation
analysis of the differentially expressed genes [99]. A hypergeometric test was used to

determine the over enrichment of the overlapping up or downregulated genes between the 2

models (k = number of overlapping genes, s = number of genes in HSJD-DIPG007 model, M

= number of genes in HSJD-DIPG007 model, N = 20,000 coding genes in human genome),

where expected = (s*M)/N.

MINTIE [107] was used for differential splicing analysis using parameters assembler=soap,

min_logfc=2, min_cpm=0.1, fdr=0.05, and run_de_step=TRUE, with all 3 Vehicle replicates

used as control. BedTools [94] was used to determine the common events between the 3

biological replicates. BAM files were visualized on IGV [97] ensuring the same scale. A

hypergeometric test was used to test if the overlap in closed regions and differentially spliced

regions was over enriched (k = overlap, s = differentially spliced, M = closed regions,

population size N = 67319 (number of genes in hg38), where expected = (s*M)/N.

Gene expression analysis from patient samples and cell lines

Patient expression data for *SSRP1, SUPT16H* and *BRD4* was obtained from the ZERO

childhood cancer precision medicine program that had RNA-seq from 84 DMG/HGG patients

[55]. Patient expression data were also obtained from McGill University (11 normal fetal

brain samples) [56]. *SSRP1, SUPT16H* and *BRD4* expression in DMG and non-malignant

cell lines was downloaded from the Childhood Cancer Model Atlas [57].

Cell Viability Assays

Cells were cultured in 96-well plates at a density of 2,500-3,000 cells per well in a final

volume of 0.2 mL of cell culture media. The cells were subjected to specific treatments as

indicated in the study. Adherent cells had the drugs added 24 hours after plating, while

suspension cells received the drugs 72 hours after plating. After an incubation period of 72

hours, cell proliferation was evaluated using the resazurin assay (Sigma-Aldrich). The

resulting data was expressed as a percentage relative to the untreated cells.

Clonogenic Assays

Long-term clonogenic assays were conducted by plating 24-well plates with a base layer of

0.3 mL of 0.5% agar mixed with cell culture media. After 24 hours of incubation at 4°C, the

base layer was overlaid with a 0.4 mL layer consisting of cells (500-1500 cells/well), 0.33%

agar, cell culture media, and the drug at the indicated concentrations. The plates were then

incubated in a 5% CO_2 humidified atmosphere at 37°C. Following an incubation period of

either 2 or 4 weeks, depending on the specific cell line, the colonies were stained with 3-

(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) solution at 5 mg/mL After 1

hour, images of the colonies were captured using Image Lab Software. Colony quantification

was performed using Image J software, and the data were presented as a percentage
relative to untreated colonies.

Orthotopic model/drug treatment
The protocol for intracranial injections was implemented as described in the study conducted
by Khan et al. [108]. The timing of treatment initiation varied for each model with SU-
DIPGVI-Luc at 7 weeks, RA055 at 3 weeks. and HSJD-DIPG007 starting at 5 weeks post
intracranial surgery.

In the SU-DIPGVI-Luc and RA055 models, the drug treatments were as follows: CBL0137 -
dissolved in 5% dextrose, CBL0137 was administered intravenously at a dosage of 40
mg/kg/day. The treatment was given once every four days for a total of four weeks. JQ1 -
dissolved in DMSO and subsequently in a solution containing 0.9% saline with 5% PEG400
and 5% TWEEN-80, JQ1 was administered at a dosage of 25 mg/kg/day intraperitoneally.
The treatment was given five days a week over a period of four weeks.

In the HSJD-DIPG007 model, the drug treatments included: CBL0137 - dissolved in 5%
dextrose, CBL0137 was administered intravenously at a dosage of 20 mg/kg/day. The
treatment was given five days a week for a total of four weeks. JQ1 - Dissolved in DMSO
and subsequently in a solution containing 0.9% saline with 5% PEG400 and 5% TWEEN-80,
JQ1 was administered intraperitoneally at a dosage of 50 mg/kg/day. The treatment was
given five days a week for a duration of four weeks.

Throughout the study, the survival of the mice was closely monitored daily. Neurological and
clinical symptoms were assessed to detect signs of tumor development. Mice exhibiting
clinical signs of neurological decline, such as ataxia, circling, head tilting, with or without a
20% weight loss, were humanely euthanized. All animal experiments strictly adhered to the
guidelines set by the Animal Care and Ethics Committee of the University of New South
Wales and the Australian Code of Practice for the Care and Use of Animals for Scientific
Purposes. The experimental protocols used in this study were approved under protocol
number 22/46B.

Histological analysis of brain tissue
Brain samples were collected on the final day of treatment and fixed in a 10% formalin
neutral buffered solution obtained from Sigma-Aldrich. The fixed samples were then
embedded in paraffin wax, and 5-mm sections were cut and mounted on glass slides. To
prepare the sections for histological examination, dehydration was carried out.

Subsequently, the sections were stained with hematoxylin/eosin and Ki67, H3K27ac, MDM4,
and MYC (Table 1) using the BOND RX Fully Automated Research Stainer (Leica
biosystems) for histologic examination. Following staining, positive cells were identified and
quantified using Andy's Algorithms [109] on the image analysis pipeline on FIJI.
739

740 **Table 1. List of antibodies**
741

Target	Catalogue number	Application
SSRP1	CST 1341	WB
SSRP1	Biolegend 609710	IP
SPT16	CST 12191	WB, CUT&RUN
SPT16	Biolegend 607002	IP
BRD4	CST 13340	IP, WB
BRD4	EpiCypher 13-2003	CUT&RUN
ß-Actin	CST 8457	WB
IgG rabbit	EpiCypher 13-0042	CUT&RUN
IgG mouse	Santa Cruz 2025	IP
H3K4me3	EpiCypher 13-0041	CUT&RUN
H3K27me3	CST 9733	CUT&RUN, WB
H3K27ac	CST 8173	CUT&RUN, WB
H3K27ac	Abcam ab4729	IHC
H3K27M	CST 74829	CUT&RUN, WB
Histone H3	CST 9717	WB
C-MYC	CST 18583	WB
C-MYC	Abcam 32072	IHC
PDGFRα	CST 3164	WB
MDM4	BETHYL-A300-287	WB
MDM4	Abcam 244366	IHC
P53	CST 2527	WB
PTCH1	CST 2468	WB

LARP1	CST 14763	WB
KLHD8CA	nbp1-31181	WB
Ki67	Thermofisher RM-9106 SP6	IHC

744 **Table 2. Sequences of qPCR primers**

Gene	Forward	Reverse	GenBank
18s	GAGGATGAGGTGGAACGTGT	TCTTCAGTCGCTCCAGGTCT	NM_022551.2
Arg1	GTCTGTGGGAAAAGCAAGCG	TTGCCAAACTGTGGTCTCCG	NM_001244438.1
B2M	AGATGAGTATGCCTGCCGTG	GCGGCATCTTCAAACCTCCA	NM_004048.2
CD86	ACTAGCACAGACACACGGATG	CTTCAGAGGAGCAGCACCAGAG	NM_175862.4
CD206	CACGATCCGACCCTTCCTTG	GTCTCCGCTTCATGCCATTG	NM_002438.3
Egr2	TTTGTGCACCAGCTGTCTGA	TTGATCATGCCATCTCCGGC	NM_000399.5
FN-1	ACAAACACTAATGTTAATTGCCCA	CCTCCAGAGCAAAGGGCTTA	NM_212482.2
HIF1α	CTTTCCTTCTCTTCTCCGCGT	TTTCGACGTTCAGAACTTATCTTTT	NM_001530.4
HLA-A	CACAGACTGACCGAGTGGAC	GGATGGTGTGAGAACCGGC	NM_002116.8
HLA-B	GCAGTTGTGGTCATCGGAGC	TCCCTCCTTTTCCACCTGAACT	NM_005514.8
IFNα	ACTCATACACCAGGTCACGC	TGGTCATAGTTATAGCAGGGGTG	NM_024013.2
IFNβ	GCGACACTGTTCGTGTTGTC	GAGCTGTAGTGGAGAAGCACA	NM_002176.4
IFNγ	GAGTGTGGAGACCATCAAGGA	TGGACATTCAAGTCAGTTACCGAA	NM_000619
IL-1β	CAGGCTGCTCTGGGATTCTC	GTCCTGGAAGGAGCACTTCAT	NM_000576
IL-6	CCAGAGCTGTGCAGATGAGT	AGCTGCGCAGAATGAGATGA	M54894.1
IL-8	TCTGCAGCTCTGTGTGAAGG	TTCTCCACAACCCTCTGCAC	NM_000584.3
IL-29	CCCCCAAAAAGGAGTCCGC	CTCAGACACAGGTTCCCATCG	NM_172140.2
iNOS	CTGGCAAGCCCAAGGTCTAT	TCCCCGCAAACATAGAGGTG	NM_000625.4
IRF1	TCTTCCCTCTTCCACTCGGA	TGATGGGCATGTTGGCTCTG	NM_002198.3
IRF3	GACCTTCCATCGTAGGCCG	CTCTTGTTCACCCAGGCCAC	NM_001571.6
IRF5	GAAAGCGAGCTCGGACCC	GGTACTGGCAGCTGTTCACC	NM_032643.5
IRF7	TGTGCTGGCGAGAAGGC	TGGAGTCCAGCATGTGTGTG	NM_001572.5
IRF8	CAAACTAGGCGTGGCAACTG	AATCGTCCACAGAAGGCTCC	NM_001363907.1
MCP-1	AGCCACCTTCATTCCCCAAG	TTGGGTTTGCTTGTCCAGGT	X14768.1
OAS1	CTGGCACAAGAGGCTGAGG	CACTGTTGCTTTCAGCCAGC	NM_016816.4
OAS3	TGAACAAGGCCGTTGATACC	GCTGAAGAGCCACCCTTGA	NM_006187.4
p53	GTTCCGAGAGCTGAATGAGG	TCTGAGTCAGGCCCTTCTGT	AF098067

| SOCS1 | CACTTCCGCACATTCCGTTC | CAGTAGAATCCGCAGGCGTC | NM_003745.2 |
| TNFα | CCCATGTTGTAGCAAACCCTC | TATCTCTCAGCTCCACGCCA | NM_000594.3 |

Data Availability

Raw and processed NGS data is currently being uploaded onto the GEO repository.

Main figure legends

Figure 1. FACT and BRD4 expression and co-localization in DMG

A) Genomic distribution of SPT16 peaks (1,095 peaks common to 2 replicates) compared to
a background of ~1000 randomly shuffled genomic regions. **B)** Dot plot of over-represented
gene ontologies for genes associated with SPT16 peaks. **C)** Transcription factor motif
enrichment analysis and motif logos for SPT16 peaks. **D)** Heatmaps depicting SPT16,
BRD4, H3K27M and IgG CUT&RUN signal (red) and ATAC-seq signal (blue) at transcription
start sites, H3K4me3 promoter peaks, H3K27ac enhancer peaks (from [42]) and H3K27me3
heterochromatin peaks in HSJD-DIPG007 neurospheres. Heatmaps are ranked based on
ATAC-seq signal. Representative data from n=2 experiments. **E)** Profile plots depicting the
average SPT16 and BRD4 signal at SPT16 peaks (left) and BRD4 peaks (right) in HSJD-
DIPG007 cells. Representative data from n=2 experiments. **F)** Co-Immunoprecipitation and
western blotting in HSJD-DIPG007 nuclear extracts showing direct interactions between
BRD4 with the FACT subunits, H3K27M and H3K27ac. Representative data from n=3
experiments. **G)** Venn diagram depicting overlapping annotated genes from 2 independent
SPT16 and BRD4 CUT&RUN replicates. A hypergeometric test was used to determine over
enrichment. **H)** Gene tracks display input, SPT16 and BRD4 CUT&RUN signal at promoters
of developmental and cancer-related genes *SOX2, MDM4* and *SMARCA2.* Bars beneath
coverage tracks represent MACS2 peak calls.

Figure 2. Combinatorial FACT and BRD4 inhibition decreases cell proliferation and
colony formation in H3K27M mutant DMG cells.

Patient-derived DMG cell lines are sensitive to BRD4 inhibition via **A)** JQ1 and **B)**
trotabresib. **C)** Sensitivity to FACT and BRD4 inhibitors, CBL0137 and JQ1 and trotabresib
respectively, was significantly higher in isogenic SU-DIPGXIII H3K27M cells compared to
KO cells. Combination treatment of **D)** CBL0137 and JQ1 and **E)** CBL0137 and trotabresib
was synergistically effective in isogenic SU-DIPGXIII H3K27M cells, with no efficacy
observed in KO cells. SU-DIPGVI cells have, upon combined FACT and BRD4 inhibition,
synergistically decreased **F)** cell survival and **G)** form fewer colonies. Data is presented as

mean values ± SEM of three independent experiments. **p<0.01, ***p<0.001, ****p<0.0001.

P-values were calculated by **C)** two-tailed t-tests for H3K27M and H3K27M-KO cells and **G)**

2-way ANOVA for single and combination treatments. Combination index values were

calculated by Calusyn.

Figure 3. CBL0137 and JQ1 condense chromatin at gene promoters.

A) Heatmaps depicting ATAC-seq signal at transcription start sites in HSJD-DIPG007 (left)

and SU-DIPGVI (right) cells treated with DMSO, CBL0137 (1µM for HSJD-DIPG007 and

2µM for SU-DIPGVI), JQ1 (1µM) or the combination for 4 hours. Average signal is shown in

the profile plots above. Representative data from n=2 experiments for HSJD-DIPG007 and

n=3 experiments for SU-DIPGVI. **B)** Profile plots depicting the average ATAC-seq signal at

SPT16 peaks (top) and BRD4 peaks (bottom) in HSJD-DIPG007 cells with the indicated

treatments. **C)** Volcano plot showing differentially accessible regions in HSJD-DIPG007

cells. Blue data points represent significantly (FDR < 0.05, fold-change < -2) downregulated

"closed" regions, red data points represent significantly upregulated "opened" (FDR < 0.05,

fold-change > 2) regions, with the number of regions indicated. **D)** Venn diagram displaying

the overlap between the closed regions common to both cell lines. A hypergeometric test

was used to determine over enrichment. **E)** Genomic distribution of the overlapping closed

regions from panel D compared to background ATAC-seq regions. **F)** Dot plot of over-

represented gene ontologies in the genes associated with the overlap regions from panel D.

G) Gene tracks display ATAC-seq signal near promoters *NES*, *SUPT16H,* and *BRD4* in

HSJD-DIPG007 (upper) and SU-DIPGVI (lower) with the indicated treatments.

Representative replicate from n=2-3 experiments.

Figure 4. CBL0137 and JQ1 cause widespread gene expression changes

A) Volcano plots depicting the significantly differentially expressed genes in HSJD-DIPG007

cells (upper panels) and SU-DIPGVI cells (lower panels) treated with DMSO, CBL0137 (1µM

for HSJD-DIPG007 and 2µM for SU-DIPGVI), JQ1 (1µM) or the combination for 4 hours.

Blue data points represent significantly (FDR < 0.05, fold-change < -1.5) downregulated

genes, red data points represent significantly upregulated genes (FDR < 0.05, fold-change >

1.5), with the number of genes indicated. **B)** Venn diagrams displaying the intersecting

downregulated (blue) and upregulated (red) genes in combination-treated cells compared to

DMSO between the 2 DMG models. A hypergeometric test was used to determine over

enrichment. **C)** Enrichment map of over-represented gene ontologies in the up (left) and

downregulated (right) genes from the overlap regions from panel B.

Figure 5. Chromatin compaction by CBL0137 and JQ1 decreases expression of
oncogenes
A) Heatmaps depicting Z-scores for RNA-seq TPM values across the 4 indicated treatments
at genes associated with SPT16 and BRD4 peaks in HSJD-DIPG007 cells **B)** Heatmaps
depicting Z-scores for RNA-seq TPM values across the 4 indicated treatments at genes
associated with promoters (<1kb from TSS) with decreased accessibility in combination
treated HSJD-DIPG007 cells (logFC < -1.5, left) and SU-DIPGVI cells (logFC< -2.5 right)
(n=3). **C)** Gene tracks displaying H3K27ac ChIP-seq [42], H3K4me3, IgG, SPT16, BRD4
CUT&RUN (CR), and ATAC-seq signal across the 4 indicated treatments in HSJD-DIPG007
and SU-DIPGVI cells at the *MYC*, *PDGFRA* and *MDM4* loci. These promoters and putative
enhancers are occupied by SPT16 and BRD4 and decrease in accessibility in combination
treated cells. **D)** Bar charts displaying RNA-seq TPM values (mean +/- SEM; n=3) for *MYC*,
PDGFRA and *MDM4* in HSJD-DIPG007 cells treated for 4h with CBL0137 (1μM) and JQ1
(1μM). **E)** Representative MYC, PDGFRα and MDM4 western blots and bar charts
displaying densitometry (mean +/- SEM; n=3) normalised to the indicated housekeeper (β-
Actin or α-Tubulin) and expressed as a fold-change relative to the DMSO control. Cells were
treated for 24h. An ordinary one-way ANOVA with Sidak's multiple comparisons test was
used to test for significance between DMSO and each treatment, and the combination
treatment compared to each monotherapy.
833
Figure 6. Therapeutic efficacy of CBL0137 and JQ1 combination treatment in
orthotopic models of DMG.
Mice were intracranially injected with DMG cells and 3-7 weeks post injection, treatments
commenced. Mice were humanely euthanized when they displayed severe neurological
decline and/or weight loss or reached maximum holding time (MHT). **A)** Survival curve of
SU-DIPGVI-Luciferase with CBL0137/JQ1 treatment. Median survival of cohorts (days):
Vehicle (n=9) = 73, CBL0137 (n=9) = 76, JQ1 (n=6) = 78, CBL0137 + JQ1 (n=6) = 95.5.
Exact p-values listed in Supplementary Table 5. **B)** Xenogen imaging showed the effect of
CBL0137/JQ1 on SU-DIPGVI-Luciferase tumors. Immunohistochemistry staining for **C)** Ki67
and **D)** H3K27ac of DMG tumors post 4 weeks of treatment in SU-DIPGVI-Luciferase model.
Three images were taken from brain samples collected from three mice in each cohort.
Scale: 50μm . **E)** Survival curve of RA055 orthotopic model with CBL0137/JQ1 treatment.
Median survival of cohorts (days): Vehicle (n=12) = 46, CBL0137 (n=9) = 51, JQ1 (n=8) =
55, CBL0137 + JQ1 (n=9) = 66. Exact p-values listed in Supplementary Table 6.
Immunohistochemistry staining for **F)** Ki67, **G)** MYC and **H)** MDM4 of DMG tumors in the
RA055 model. Three images were taken from brain samples collected from three mice in
each cohort at the end of the treatment period. Scale: 50μm. **A, E)** Shaded areas indicate

treatment period of combination treatment in each model. Statistical analysis has been
performed using the Log-Rank (Mantel-Cox) with multiple test corrections applied.
C,D,F,G,H) Data is presented as mean values ± SEM. *p<0.05, **p<0.01, ***p<0.001,
****p<0.0001. p-values were calculated using one-way ANOVA for treated and untreated
cohorts.

Supplementary Table 1: Gene lists identified in CUT&RUN, ATAC-seq and RNA-seq
experiments.
A) Genes associated with SPT16 and BRD4 MACS2 peaks from two independent
CUT&RUN experiments in HSJD-DIPG007 cells. Genes from individual replicates, combined
replicates, and overlapping genes between SPT16 and BRD4 combined replicates are listed.
B) Unique genes associated with ATAC-seq CSAW differentially accessible regions (DAR)
with FDR<0.05 and fold-change >2x = opened, fold-change < -2x = closed in HSJD-
DIPG007 and SU-DIPGVI cells treated with CBL0137 (1μM HSJD-DIPG007 and 2μM SU-
DIPGVI) and JQ1 (1μM) for 4h. **C)** RNA-seq significantly differentially expression genes with
FDR<0.05 and fold-change >1.5x = UP, fold-change < -1.5x = DOWN in HSJD-DIPG007
and SU-DIPGVI cells treated with CBL0137 (1μM HSJD-DIPG007 and 2μM SU-DIPGVI) and
JQ1 (1μM) for 4h. **D)** Genes with differential splicing common to 3 independent replicates in
HSJD-DIPG007 and SU-DIPGVI treated with CBL0137 (1μM HSJD-DIPG007 and 2μM SU-
DIPGVI) and JQ1 (1μM) in combination for 4h.

Acknowledgments

We would like to express our gratitude to Michelle Monje, Angel Montero Carcaboso, and
Esther Hulleman for their generous provision of the SU-DIPG, HSJD-DIPG, and VUMC-
DIPG10 cells, respectively, which were instrumental in the success of this study. We thank
Nada Jabado and Brian Krug for providing the isogenic H3K27M KO SU-DIPGXIII models
and Yunjia Zhang, Oluwaseyi Samson Ogunmodede, and Kerry-Anne Rye for providing the
HMDM cells. We also acknowledge the Core Services Group at Children's Cancer Institute
for their valuable technical support throughout the research. We extend our appreciation to
the Flow Cytometry Core Facility, BRIL Facility, and BMSF Facility (Russel Pickford) at
UNSW assistance in flow cytometry, imaging, and pharmacokinetic testing, respectively,
which greatly contributed to our data analysis and interpretation. We also acknowledge the
Garvan Institute Histopathology Facility for their services. Furthermore, we would like to
acknowledge the Zero Childhood Cancer Program and the Tumour Bank teams at CCI for
their significant contributions to the establishment of primary DIPG and normal astrocyte

cultures, which were indispensable for our investigations. The authors thank the Children's
Cancer Institute Animal Facility for providing support to this study. We would like to thank
Nathan Dahl and Lays Sobral for technical advice regarding CUT&RUN and ATAC-seq. This
work has generously been funded by the Kids Cancer Alliance, We Love You Connie
Foundation Grant, Levi's Project, CINSW Fellowship (HH) (2022/ECF1425), The Kids
Cancer Project Col Reynolds Fellowship (AK), NHMRC Synergy Grant (2019056), Cancer
Institute NSW Translational Program Grant (2019/TPG2037), NHMRC Investigator grant
(GNT2017898), and DIPG Collaborative: The Cure Starts Now (2018). Graphical abstract
created with BioRender.com.

Author Contributions

897 **Conceptualization**: MT, DSZ. **Methodology:** HH, AK, JL. **Formal Analysis:** HH, AK, AE,
898 NJ, SER, XG, HN, CM, BR. **Investigation:** HH, AK, AE, AG, HN, AL, BR, MT. **Data
899 Curation:** HH, AK, CM, XG, NJ, SER. **Visualization:** HH, AK, SER, YCS, BR. **Writing –
900 Original Draft:** HH, AK. **Writing – Review & Editing:** All. **Supervision:** CM, BR, MT, MD,
901 DSZ. **Project Administration:** HH, AK. **Funding Acquisition:** HH, BR, MT, DSZ.

Declaration of Interests

904 DSZ declares consulting/advisory board fees from Bayer, Astra Zeneca,
905 Accendatech, Novartis, Day One, FivePhusion, Amgen, Alexion, and Norgine and
906 research support from Accendatech. The remaining authors have declared that no
907 competing interests exist.

References

1.   Louis, D.N., et al., *The 2021 WHO Classification of Tumors of the Central Nervous
System: a summary.* Neuro-Oncology, 2021. **23**(8): p. 1231-1251.
2.   Schwartzentruber, J., et al., *Driver mutations in histone H3.3 and chromatin
remodelling genes in paediatric glioblastoma.* Nature, 2012. **482**(7384): p. 226-231.
3.   Khuong-Quang, D.A., et al., *K27M mutation in histone H3.3 defines clinically and
biologically distinct subgroups of pediatric diffuse intrinsic pontine gliomas.* Acta
Neuropathol, 2012. **124**(3): p. 439-47.
4.   Wu, G., et al., *Somatic histone H3 alterations in pediatric diffuse intrinsic pontine
gliomas and non-brainstem glioblastomas.* Nat Genet, 2012. **44**(3): p. 251-3.
5.   Castel, D., et al., *Histone H3 wild-type DIPG/DMG overexpressing EZHIP extend the
spectrum diffuse midline gliomas with PRC2 inhibition beyond H3-K27M mutation.*
Acta Neuropathologica, 2020. **139**(6): p. 1109-1113.

6.   Pratt, D., et al., *Diffuse intrinsic pontine glioma-like tumor with EZHIP expression and*
923   *molecular features of PFA ependymoma.* Acta Neuropathologica Communications,
924   2020. **8**(1): p. 37.

925   7.   Ajuyah, P., et al., *Histone H3-wild type diffuse midline gliomas with H3K27me3 loss*
926   *are a distinct entity with exclusive EGFR or ACVR1 mutation and differential*
927   *methylation of homeobox genes.* Sci Rep, 2023. **13**(1): p. 3775.

928   8.   Lewis, P.W., et al., *Inhibition of PRC2 activity by a gain-of-function H3 mutation found*
929   *in pediatric glioblastoma.* Science, 2013. **340**(6134): p. 857-61.

930   9.   Piunti, A., et al., *CATACOMB: An endogenous inducible gene that antagonizes*
931   *H3K27 methylation activity of Polycomb repressive complex 2 via an H3K27M-like*
932   *mechanism.* Science Advances, 2019. **5**(7): p. eaax2887.

933   10.   Bender, S., et al., *Reduced H3K27me3 and DNA hypomethylation are major drivers*
934   *of gene expression in K27M mutant pediatric high-grade gliomas.* Cancer Cell, 2013.
935   **24**(5): p. 660-72.

936   11.   Harutyunyan, A.S., et al., *H3K27M induces defective chromatin spread of PRC2-*
937   *mediated repressive H3K27me2/me3 and is essential for glioma tumorigenesis.*
938   Nature Communications, 2019. **10**(1): p. 1262.

939   12.   Krug, B., et al., *Pervasive H3K27 Acetylation Leads to ERV Expression and a*
940   *Therapeutic Vulnerability in H3K27M Gliomas.* Cancer cell, 2019. **35**(5): p. 782-
941   797.e8.

942   13.   Nagaraja, S., et al., *Histone Variant and Cell Context Determine H3K27M*
943   *Reprogramming of the Enhancer Landscape and Oncogenic State.* Mol Cell, 2019.
944   **76**(6): p. 965-980.e12.

945   14.   Brien, G.L., et al., *Simultaneous disruption of PRC2 and enhancer function underlies*
946   *histone H3.3-K27M oncogenic activity in human hindbrain neural stem cells.* Nature
947   Genetics, 2021.

948   15.   Hayden, E., et al., *Therapeutic Targets in Diffuse Midline Gliomas-An Emerging*
949   *Landscape.* Cancers (Basel), 2021. **13**(24).

950   16.   Groves, A. and T.M. Cooney, *Epigenetic programming of pediatric high-grade*
951   *glioma: Pushing beyond proof of concept to clinical benefit.* Front Cell Dev Biol, 2022.
952   **10**: p. 1089898.

953   17.   Orphanides, G., et al., *FACT, a factor that facilitates transcript elongation through*
954   *nucleosomes.* Cell, 1998. **92**(1): p. 105-16.

955   18.   Orphanides, G., et al., *The chromatin-specific transcription elongation factor FACT*
956   *comprises human SPT16 and SSRP1 proteins.* Nature, 1999. **400**(6741): p. 284-8.

957   19.   Ehteda, A., et al., *Dual targeting of the epigenome via FACT complex and histone*
958   *deacetylase is a potent treatment strategy for DIPG.* Cell Rep, 2021. **35**(2): p.
959   108994.

960   20.   Garcia, H., et al., *Facilitates Chromatin Transcription Complex Is an "Accelerator" of*
961   *Tumor Transformation and Potential Marker and Target of Aggressive Cancers.* Cell
962   Reports, 2013. **4**(1): p. 159-173.

963   21.   Gurova, K., et al., *Structure and function of the histone chaperone FACT - Resolving*
964   *FACTual issues.* Biochimica et biophysica acta. Gene regulatory mechanisms, 2018:
965   p. S1874-9399(18)30159-7.

966   22.   Safina, A., et al., *FACT is a sensor of DNA torsional stress in eukaryotic cells.*
967   Nucleic Acids Res, 2017. **45**(4): p. 1925-1945.

968   23.   Chang, H.-W., et al., *Mechanism of FACT removal from transcribed genes by*
969   *anticancer drugs curaxins.* Science advances, 2018. **4**(11): p. eaav2131-eaav2131.

970   24.   Neefjes, J., et al., *Chromatin as an old and new anticancer target.* Trends Cancer,
971   2024.

972   25.   Gasparian, A.V., et al., *Curaxins: anticancer compounds that simultaneously*
973   *suppress NF-κB and activate p53 by targeting FACT.* Science translational medicine,
974   2011. **3**(95): p. 95ra74-95ra74.

975   26.   Carter, D.R., et al., *Therapeutic targeting of the MYC signal by inhibition of histone*
976   *chaperone FACT in neuroblastoma.* Sci Transl Med, 2015. **7**(312): p. 312ra176.

27.   Xiao, L., et al., *Dual targeting of chromatin stability by the curaxin CBL0137 and histone deacetylase inhibitor panobinostat shows significant preclinical efficacy in neuroblastoma.* Clin Cancer Res, 2021.

28.   Tallman, M.M., et al., *The small molecule drug CBL0137 increases the level of DNA damage and the efficacy of radiotherapy for glioblastoma.* Cancer Letters, 2021. **499**: p. 232-242.

29.   Wang, J., et al., *Effective inhibition of MYC-amplified group 3 medulloblastoma by FACT-targeted curaxin drug CBL0137.* Cell Death Dis, 2020. **11**(12): p. 1029.

30.   Monje, M., et al., *Phase I trial of panobinostat in children with diffuse intrinsic pontine glioma: A report from the Pediatric Brain Tumor Consortium (PBTC-047).* Neuro Oncol, 2023. **25**(12): p. 2262-2272.

31.   Bradner, J.E., D. Hnisz, and R.A. Young, *Transcriptional Addiction in Cancer.* Cell, 2017. **168**(4): p. 629-643.

32.   Shi, J. and C.R. Vakoc, *The mechanisms behind the therapeutic activity of BET bromodomain inhibition.* Molecular cell, 2014. **54**(5): p. 728-736.

33.   Lovén, J., et al., *Selective inhibition of tumor oncogenes by disruption of super-enhancers.* Cell, 2013. **153**(2): p. 320-34.

34.   Taylor, I.C., et al., *Disrupting NOTCH Slows Diffuse Intrinsic Pontine Glioma Growth, Enhances Radiation Sensitivity, and Shows Combinatorial Efficacy With Bromodomain Inhibition.* J Neuropathol Exp Neurol, 2015. **74**(8): p. 778-90.

35.   Nagaraja, S., et al., *Transcriptional Dependencies in Diffuse Intrinsic Pontine Glioma.* Cancer cell, 2017. **31**(5): p. 635-652.e6.

36.   Piunti, A., et al., *Therapeutic targeting of polycomb and BET bromodomain proteins in diffuse intrinsic pontine gliomas.* Nature Medicine, 2017. **23**(4): p. 493-500.

37.   Zhang, Y., et al., *Combination of EZH2 inhibitor and BET inhibitor for treatment of diffuse intrinsic pontine glioma.* Cell Biosci, 2017. **7**: p. 56.

38.   Wiese, M., et al., *Combined treatment with CBP and BET inhibitors reverses inadvertent activation of detrimental super enhancer programs in DIPG cells.* Cell Death & Disease, 2020. **11**(8): p. 673.

39.   Watanabe, J., et al., *BET bromodomain inhibition potentiates radiosensitivity in models of H3K27-altered diffuse midline glioma.* J Clin Invest, 2024.

40.   Groves, A., J. Clymer, and M.G. Filbin, *Bromodomain and Extra-Terminal Protein Inhibitors: Biologic Insights and Therapeutic Potential in Pediatric Brain Tumors.* Pharmaceuticals, 2022. **15**(6): p. 665.

41.   Bočkaj, I., et al., *The H3.3K27M oncohistone affects replication stress outcome and provokes genomic instability in pediatric glioma.* PLoS genetics, 2021. **17**(11): p. e1009868-e1009868.

42.   Wang, J., et al., *Epigenomic landscape and 3D genome structure in pediatric high-grade glioma.* Science Advances, 2021. **7**(23): p. eabg4126.

43.   Yu, Y., et al., *Olig2 targets chromatin remodelers to enhancers to initiate oligodendrocyte differentiation.* Cell, 2013. **152**(1-2): p. 248-61.

44.   Han, S.P., et al., *SNAI1 is involved in the proliferation and migration of glioblastoma cells.* Cell Mol Neurobiol, 2011. **31**(3): p. 489-96.

45.   Meel, M.H., et al., *Signaling pathways and mesenchymal transition in pediatric high-grade glioma.* Cell Mol Life Sci, 2018. **75**(5): p. 871-887.

46.   Furth, N., et al., *H3-K27M-mutant nucleosomes interact with MLL1 to shape the glioma epigenetic landscape.* Cell Rep, 2022. **39**(7): p. 110836.

47.   Falsone, S.F., B. Gesslbauer, and A.J. Kungl, *Coimmunoprecipitation and proteomic analyses.* Methods Mol Biol, 2008. **439**: p. 291-308.

48.   Weina, K. and J. Utikal, *SOX2 and cancer: current research and its implications in the clinic.* Clin Transl Med, 2014. **3**: p. 19.

49.   Marine, J.C. and A.G. Jochemsen, *MDMX (MDM4), a Promising Target for p53 Reactivation Therapy and Beyond.* Cold Spring Harb Perspect Med, 2016. **6**(7).

50. Centore, R.C., et al., *Mammalian SWI/SNF Chromatin Remodeling Complexes:*
Emerging Mechanisms and Therapeutic Strategies. Trends Genet, 2020. **36**(12): p.
936-950.
51. Hsieh, F.-K., et al., *FACT in Cell Differentiation and Carcinogenesis.* Oncotarget,
2011. **2**(11): p. 830-832.
52. Sui, S., et al., *Ferritinophagy is required for the induction of ferroptosis by the*
bromodomain protein BRD4 inhibitor (+)-JQ1 in cancer cells. Cell Death & Disease,
2019. **10**(5): p. 331.
53. Lu, L., et al., *Inhibition of BRD4 suppresses the malignancy of breast cancer cells via*
regulation of Snail. Cell Death & Differentiation, 2020. **27**(1): p. 255-268.
54. Boyson, S.P., et al., *Functional Roles of Bromodomain Proteins in Cancer.* Cancers
(Basel), 2021. **13**(14).
55. Wong, M., et al., *Whole genome, transcriptome and methylome profiling enhances*
actionable target discovery in high-risk pediatric cancer. Nature Medicine, 2020.
26(11): p. 1742-1753.
56. Jessa, S., et al., *Stalled developmental programs at the root of pediatric brain*
tumors. Nature genetics, 2019. **51**(12): p. 1702-1713.
57. Sun, C.X., et al., *Generation and multi-dimensional profiling of a childhood cancer*
cell line atlas defines new therapeutic opportunities. Cancer Cell, 2023. **41**(4): p. 660-
677.e7.
58. Delmore, Jake E., et al., *BET Bromodomain Inhibition as a Therapeutic Strategy to*
Target c-Myc. Cell, 2011. **146**(6): p. 904-917.
59. Nesher, E., et al., *Role of Chromatin Damage and Chromatin Trapping of FACT in*
Mediating the Anticancer Cytotoxicity of DNA-Binding Small-Molecule Drugs. Cancer
Res, 2018. **78**(6): p. 1431-1443.
60. Kantidze, O.L., et al., *The anti-cancer drugs curaxins target spatial genome*
organization. Nat Commun, 2019. **10**(1): p. 1441.
61. Krupkova, O., Jr., et al., *Nestin expression in human tumors and tumor cell lines.*
Neoplasma, 2010. **57**(4): p. 291-8.
62. Wang, Q., et al., *Nestin Is Required for Spindle Assembly and Cell-Cycle*
Progression in Glioblastoma Cells. Mol Cancer Res, 2021. **19**(10): p. 1651-1665.
63. Zee, B.M., et al., *In vivo residue-specific histone methylation dynamics.* J Biol Chem,
2010. **285**(5): p. 3341-50.
64. Wohlberedt, K., et al., *Mdm4 supports DNA replication in a p53-independent fashion.*
Oncogene, 2020. **39**(25): p. 4828-4843.
65. Tcherkezian, J., et al., *Proteomic analysis of cap-dependent translation identifies*
LARP1 as a key regulator of 5'TOP mRNA translation. Genes Dev, 2014. **28**(4): p.
357-71.
66. Lee, D., et al., *Superenhancer activation of KLHDC8A drives glioma ciliation and*
hedgehog signaling. J Clin Invest, 2023. **133**(2).
67. Alló, M., et al., *Chromatin and alternative splicing.* Cold Spring Harb Symp Quant
Biol, 2010. **75**: p. 103-11.
68. Zhou, X., et al., *SPTBN2 Promotes the Progression of Thyroid Cancer by*
Accelerating G1/S Transition and Inhibiting Apoptosis. Dis Markers, 2022. **2022**: p.
2562595.
69. Wu, C., et al., *SPTBN2, a New Biomarker of Lung Adenocarcinoma.* Front Oncol,
2021. **11**: p. 754290.
70. Yang, L. and Y. Gu, *SPTBN2 regulates endometroid ovarian cancer cell proliferation,*
invasion and migration via ITGB4⬚mediated focal adhesion and ECM receptor
signalling pathway. Exp Ther Med, 2023. **25**(6): p. 277.
71. Harrison, P.W., et al., *Ensembl 2024.* Nucleic Acids Research, 2023. **52**(D1): p.
D891-D899.
72. Leonova, K., et al., *TRAIN (Transcription of Repeats Activates INterferon) in*
response to chromatin destabilization induced by small molecules in mammalian
cells. Elife, 2018. **7**.

73.   Bachu, M., A. Dey, and K. Ozato, *Chromatin Landscape of the IRF Genes and Role*
1086   *of the Epigenetic Reader BRD4.* J Interferon Cytokine Res, 2016. **36**(7): p. 470-5.
1087   74.   Yang, X., et al., *Epigenetic regulation of macrophage polarization by DNA*
1088   *methyltransferase 3b.* Mol Endocrinol, 2014. **28**(4): p. 565-74.
1089   75.   Hashizume, R., et al., *Pharmacologic inhibition of histone demethylation as a therapy*
1090   *for pediatric brainstem glioma.* Nat Med, 2014. **20**(12): p. 1394-6.
1091   76.   Mohammad, F., et al., *EZH2 is a potential therapeutic target for H3K27M-mutant*
1092   *pediatric gliomas.* Nature Medicine, 2017. **23**(4): p. 483-492.
1093   77.   Grasso, C.S., et al., *Functionally defined therapeutic targets in diffuse intrinsic*
1094   *pontine glioma.* Nature Medicine, 2015. **21**(6): p. 555-559.
1095   78.   Balakrishnan, I., et al., *Senescence Induced by BMI1 Inhibition Is a Therapeutic*
1096   *Vulnerability in H3K27M-Mutant DIPG.* Cell Rep, 2020. **33**(3): p. 108286.
1097   79.   Mo, Y., et al., *Epigenome programing by H3.3K27M mutation creates a dependence*
1098   *of pediatric glioma on SMARCA4.* Cancer Discov, 2022.
1099   80.   Panditharatna, E., et al., *BAF complex maintains glioma stem cells in pediatric*
1100   *H3K27M-glioma.* Cancer Discov, 2022.
1101   81.   Mota, M., et al., *Targeting SWI/SNF ATPases in H3.3K27M diffuse intrinsic pontine*
1102   *gliomas.* Proc Natl Acad Sci U S A, 2023. **120**(18): p. e2221175120.
1103   82.   Dahl, N.A., et al., *Super Elongation Complex as a Targetable Dependency in Diffuse*
1104   *Midline Glioma.* Cell Rep, 2020. **31**(1): p. 107485.
1105   83.   Klein, D.C., et al., *FACT regulates pluripotency through proximal and distal regulation*
1106   *of gene expression in murine embryonic stem cells.* BMC Biol, 2023. **21**(1): p. 167.
1107   84.   Jang, M.K., et al., *The bromodomain protein Brd4 is a positive regulatory component*
1108   *of P-TEFb and stimulates RNA polymerase II-dependent transcription.* Mol Cell,
1109   2005. **19**(4): p. 523-34.
1110   85.   Rahme, G.J., et al., *Modeling epigenetic lesions that cause gliomas.* Cell, 2023.
1111   **186**(17): p. 3674-3685.e14.
1112   86.   Monje, M., et al., *Hedgehog-responsive candidate cell of origin for diffuse intrinsic*
1113   *pontine glioma.* Proc Natl Acad Sci U S A, 2011. **108**(11): p. 4453-8.
1114   87.   Mo, J., et al., *Inhibition of the FACT Complex Targets Aberrant Hedgehog Signaling*
1115   *and Overcomes Resistance to Smoothened Antagonists.* Cancer Research, 2021.
1116   **81**(11): p. 3105.
1117   88.   Okonechnikov, K., et al., *3D genome mapping identifies subgroup-specific*
1118   *chromosome conformations and tumor-dependency genes in ependymoma.* Nat
1119   Commun, 2023. **14**(1): p. 2300.
1120   89.   Jones, P.A., et al., *Epigenetic therapy in immune-oncology.* Nat Rev Cancer, 2019.
1121   **19**(3): p. 151-161.
1122   90.   Somers, K., et al., *Potent antileukemic activity of curaxin CBL0137 against MLL-*
1123   *rearranged leukemia.* Int J Cancer, 2020. **146**(7): p. 1902-1916.
1124   91.   Rayner, B.S., et al., *Role of hypochlorous acid (HOCl) and other inflammatory*
1125   *mediators in the induction of macrophage extracellular trap formation.* Free Radic
1126   Biol Med, 2018. **129**: p. 25-34.
1127   92.   Holliday, H., et al., *Inhibitor of Differentiation 4 (ID4) represses mammary*
1128   *myoepithelial differentiation via inhibition of HEB.* iScience, 2021. **24**(2): p. 102072.
1129   93.   Livak, K.J. and T.D. Schmittgen, *Analysis of Relative Gene Expression Data Using*
1130   *Real-Time Quantitative PCR and the 2−ΔΔCT Method.* Methods, 2001. **25**(4): p. 402-
1131   408.
1132   94.   Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing*
1133   *genomic features.* Bioinformatics, 2010. **26**(6): p. 841-2.
1134   95.   Ramírez, F., et al., *deepTools: a flexible platform for exploring deep-sequencing*
1135   *data.* Nucleic Acids Res, 2014. **42**(Web Server issue): p. W187-91.
1136   96.   Nordin, A., et al., *The CUT&RUN suspect list of problematic regions of the genome.*
1137   Genome Biol, 2023. **24**(1): p. 185.
1138   97.   Robinson, J.T., et al., *Integrative genomics viewer.* Nat Biotechnol, 2011. **29**(1): p.
1139   24-6.

98. Yu, G., L.G. Wang, and Q.Y. He, *ChIPseeker: an R/Bioconductor package for ChIP*
peak annotation, comparison and visualization. Bioinformatics, 2015. **31**(14): p.
2382-3.
99. Yu, G., et al., *clusterProfiler: an R package for comparing biological themes among*
gene clusters. Omics, 2012. **16**(5): p. 284-7.
100. Liu, T., et al., *Cistrome: an integrative platform for transcriptional regulation studies.*
Genome Biol, 2011. **12**(8): p. R83.
101. Lun, A.T. and G.K. Smyth, *csaw: a Bioconductor package for differential binding*
analysis of ChIP-seq data using sliding windows. Nucleic Acids Res, 2016. **44**(5): p.
e45.
102. Corces, M.R., et al., *An improved ATAC-seq protocol reduces background and*
enables interrogation of frozen tissues. Nat Methods, 2017. **14**(10): p. 959-962.
103. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat
Methods, 2012. **9**(4): p. 357-9.
104. Tarasov, A., et al., *Sambamba: fast processing of NGS alignment formats.*
Bioinformatics, 2015. **31**(12): p. 2032-4.
105. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013.
29(1): p. 15-21.
106. Li, B. and C.N. Dewey, *RSEM: accurate transcript quantification from RNA-Seq data*
with or without a reference genome. BMC Bioinformatics, 2011. **12**: p. 323.
107. Cmero, M., et al., *MINTIE: identifying novel structural and splice variants in*
transcriptomes using RNA-seq data. Genome Biol, 2021. **22**(1): p. 296.
108. Khan, A., et al., *Dual targeting of polyamine synthesis and uptake in diffuse intrinsic*
pontine gliomas. Nat Commun, 2021. **12**(1): p. 971.
109. Law, A.M.K., et al., *Andy's Algorithms: new automated digital image analysis*
pipelines for FIJI. Sci Rep, 2017. **7**(1): p. 15717.
1166

Diffuse Midline Glioma H3K27M mutant

FACT and BRD4 support transcriptionally active chromatin in H3K27M DMG

BRD4

FACT

SSRP1 SPT16

ac

Pol II

H3K27M

oncogenic mRNA

e.g. *PDGFRA, MYC, MDM4, PTCH1, LARP1, KLDHC8A*

Combined inhibition leads to chromatin condensation and perturbed transcription

CBL0137 + JQ1

BRD4

ac

SSRP1 SPT16

- repression of oncogenic transcripts
- alternatively spliced transcripts

Figure 1

Figure 2

Figure 3

A HSJD-DIPG007   SU-DIPGVI

B HSJD-DIPG007

C HSJD−DIPG007 − Combination vs. DMSO

11,704 "closed"   828 "opened"

D HSJD-DIPG007 Closed regions   SU-DIPGVI Closed regions

5,752   6,119   35,338

E Feature Distribution

Commonly closed

HSJD-DIPG007 background

SU-DIPGVI background

Feature:
Promoter (<=1kb)
Promoter (1-2kb)
Promoter (2-3kb)
5' UTR
3' UTR
1st Exon
Other Exon
1st Intron
Other Intron
Downstream (<=300)
Distal Intergenic

F GSEA on commonly closed genes

commonly closed genes

axon development
histone modification
axonogenesis
Wnt signaling pathway
cell-cell signaling by wnt
actin filament organization
cell junction assembly
regulation of neuron projection development
regulation of cell morphogenesis
regulation of small GTPase mediated signal transduction
spindle organization
neural precursor cell proliferation

G

chr1:156,666,763-156,679,407   chr14:21,380,874-21,386,980   chr19:15,273,531-15,296,773

NES   SUPT16H   XR_001750627.1   BRD4   BRD4

Figure 4

Figure 5

Figure 6