Chromatin remodeling with combined FACT and BET inhibition

disrupts oncogenic transcription in Diffuse Midline Glioma

Authors: Holly Holliday[1,2]*, Aaminah Khan[1,2]*, Anahid Ehteda[1,2], Hieu Nguyen[1], Samuel E.

Ross[3], Nisitha Jayatilleke[1], Anjana Gopalakrishnan[1], Eyden Wang[1], Yolanda Colino

Sanguino[1,2], Daisy Kavanagh[4,5], Xinyi Guo[1,6], Jie Liu[1], David Lawrence[1,2], Claire X. Sun[6,7],

Rebecca Lehmann[1,2], Chi Kin Ip[1,2], Alvin Lee[1], Laura Rangel-Sanchez[4,8], Wenyan Li[1], Robert

Salomon[1,2], Ron Firestein[6,7], Robert J Weatheritt[4,5], Fatima Valdes-Mora[1,2], Marcel E.

Dinger[3], Timothy N. Phoenix[9], Chelsea Mayoh[1,2], Benjamin S. Rayner[1,2], Maria Tsoli[1,2]**,

David S. Ziegler[1, 2, 10]**

*, ** equal contribution

Affiliations:

1. Children's Cancer Institute, Lowy Cancer Research Centre, UNSW Sydney, Sydney,
NSW, Australia
2. School of Clinical Medicine, UNSW Sydney, Sydney, NSW, Australia
3. School of Life and Environmental Sciences, University of Sydney, NSW, Australia
4. Garvan Institute of Medical Research, Sydney, NSW, Australia
5. School of Biotechnology and Biomolecular Sciences, UNSW Sydney, Sydney, NSW,
Australia
6. Centre for Cancer Research, Hudson Institute of Medical Research, Clayton,
Australia
7. Department of Medicine, School of Clinical Sciences, Monash University, Clayton,
Australia
8. School of Biomedical Engineering, Faculty of Engineering and Information
Technology, University of Technology Sydney, Sydney, NSW Australia
9. Division of Pharmaceutical Sciences, James L. Winkle College of Pharmacy,
University of Cincinnati, Cincinnati, OH, USA
10. Kid's Cancer Centre, Sydney Children's Hospital, Randwick, NSW, Australia

Keywords: FACT; BRD4; BET proteins; Diffuse Midline Glioma; DMG; DIPG; H3K27M;

Epigenetics; Chromatin; Histones; CBL0137; JQ1; BET inhibitors; Epigenetic therapies; Pre-

clinical

Corresponding authors: Professor David S. Ziegler, Kids Cancer Centre, Sydney

Children's Hospital, High St, Randwick 2031 Australia, +61 (02) 9382 1730,

d.ziegler@unsw.edu.au

Declaration of Interest statement: DSZ declares consulting/advisory board fees from

Bayer, Astra Zeneca, Accendatech, Novartis, Day One, FivePhusion, Amgen, Alexion, and

Norgine and research support from Accendatech. The remaining authors have declared that

no competing interests exist.

Abstract

Aberrant epigenetic regulation is a hallmark of Diffuse Midline Glioma (DMG), an incurable pediatric brain tumor. The H3K27M driver histone mutation leads to transcriptional dysregulation, indicating that targeting the epigenome and transcription may be key therapeutic strategies against this highly aggressive cancer. One such target is the Facilitates Chromatin Transcription (FACT) histone chaperone. We found FACT to be enriched at developmental gene promoters, coinciding with regions of open chromatin and binding motifs of core DMG regulatory transcription factors. Furthermore, FACT interacted and co-localized with the Bromodomain and Extra-Terminal Domain (BET) protein BRD4 at promoters and enhancers, suggesting functional cooperation between FACT and BRD4 in DMG. *In vitro*, a combinatorial therapeutic approach using the FACT inhibitor CBL0137, coupled with BET inhibition revealed potent and synergistic cytotoxicity across a range of DMG cultures. These results were recapitulated *in vivo*, significantly extending survival in three independent orthotopic PDX models of DMG. Mechanistically, we show that CBL0137 treatment decreased chromatin accessibility, synergizing with BET inhibition to cause broad transcriptional collapse, silencing several key oncogenes including *MYC, PDGFRA*, *MDM4* and *SOX2*, as well as causing alterations to the splicing landscape. Notably, this combination also elicited immune-related effects, including activation of the interferon response and antigen presentation mechanisms in DMG cells and induction of an activated state in macrophages and T cells, as demonstrated in an immunocompetent setting with spatial transcriptomics. Altogether, our data highlights the therapeutic promise of simultaneously targeting FACT and BET proteins in DMG, offering a dual tumor-intrinsic and immune-mediated strategy for combating this devastating pediatric brain tumor.

Introduction

Diffuse midline glioma (DMG) is a highly aggressive brain tumor predominantly diagnosed in children under 10 years old. Recently classified as 'DMG K27-altered' by the 2021 World Health Organization, this designation recognizes the characteristic epigenetic dysregulation that arises from either the recurrent lysine-to-methionine driver mutations in Histone H3.1 and H3.3 genes (H3K27M), or overexpression of the oncohistone mimic EZHIP [1-7]. Both H3K27M and EZHIP inhibit Polycomb Repressive Complex 2 (PRC2) activity, leading to a global loss of Histone H3K27 trimethylation (H3K27me3), increased H3K27 acetylation (H3K27ac) and DNA hypomethylation. These changes establish an aberrant epigenetic landscape permissive to heightened transcriptional activity [8-14].

While direct therapeutic targeting of H3K27M has not been possible, targeting of epigenetic and transcription regulators has shown promise in several pre-clinical models of DMG (reviewed in [15, 16]). Among these regulators, the histone chaperone Facilitates Chromatin Transcription (FACT) comprising of two protein subunits; SSRP1 and SPT16, has emerged as a compelling target [17-19]. FACT plays a pivotal role in maintaining chromatin integrity by facilitating nucleosome turnover during transcription, DNA replication and repair. This chaperone is abundantly expressed in cancer cells, making it an attractive therapeutic target [20, 21].

The curaxin compound CBL0137 indirectly inhibits FACT by intercalating with DNA and trapping FACT on chromatin (a process known as 'c-trapping'), effectively depleting the functional pool of this important chaperone [22-24]. We and others have demonstrated CBL0137 to be active against multiple tumor types, including neuroblastoma, glioblastoma, medulloblastoma, and diffuse intrinsic pontine glioma (DIPG)/DMG [19, 25-29]. This growing body of preclinical work has led to the advancement of CBL0137 to a Phase I/II clinical trial for pediatric patients with solid tumors, including DMG (NCT04870944). However, the aggressive nature of DMG necessitates combination therapies to address therapeutic resistance and reduce effective therapeutic doses. While initial studies combining CBL0137 with the pan-HDAC inhibitor panobinostat has shown promise against DMG pre-clinically [19], concerns over panobinostat's high toxicity, and limited efficacy, led to its withdrawal from clinical studies and highlights the need for alternative combination strategies [30].

Disruption of transcription has proven effective against many 'transcription addicted' malignancies, often achieved by targeting Bromodomain and Extra-Terminal Domain (BET)

proteins such as BRD4 [31]. BRD4 recruits the transcriptional machinery to hyper-acetylated

chromatin at super-enhancers [32, 33], making it an attractive target against DMG, as

demonstrated in the preclinical setting ([34-39] and reviewed in [40]). The tool compound

JQ1, among other BET inhibitors, has demonstrated pronounced efficacy against DMG,

through transcriptional disruption leading to cytostatis [35]. While next-generation BET

inhibitors, including trotabresib (CC-90010), have entered clinical trials for brain tumors

(NCT04047303, NCT04324840, NCT03936465)[40], their modest efficacy as single agents

underscores the need for improved BET-targeting compounds and rational combination

strategies.

Here we describe the promising efficacy of FACT and BET inhibition as a dual epigenetic

therapy in cell line and animal models of DMG. Mechanistically, this epigenetic intervention

induces chromatin compaction at gene promoters, and leads to widespread transcriptional

collapse. Notably, oncogenes such as *MYC, PDGFRA, MDM4* and *SOX2* are directly

regulated by FACT and BRD4, being silenced through dual inhibition. Additionally, we

observe aberrant splicing, upregulation of interferon genes and the antigen presentation

machinery, and changes in immune cell transcriptional signatures and phenotypes.

Together, our findings provide proof-of-principle evidence for combining FACT and BET

inhibition to co-target key oncogenic drivers and otherwise 'untargetable' pathways in DMG,

and modulating the tumor-immune landscape in DMG for effective tumor clearance. While

further optimization of BET inhibitors will be critical for clinical translation, these data

establish a promising foundation for future therapeutic development.

Results

FACT and BRD4 are overexpressed, interact, and co-localize at active regulatory

elements in DMG

To elucidate the genes bound and potentially regulated by FACT, we mapped the genome-

wide binding sites of the FACT subunit SPT16 in H3.3K27M-mutant HSJD-DIPG007 and

SU-DIPGVI, as well as H3-WT VUMC-DIPG10 cells using Cleavage Under Targets and

Release Using Nuclease (CUT&RUN). We identified 4,259 peaks in HSJD-DIPG007 cells,

1,792 peaks in SU-DIPGVI cells, and 707 peaks in VUMC-DIPG10 cells (**Table S1A-B).**

SPT16 peaks were over-enriched at proximal gene promoters in all cell lines (**Fig 1A**).

SPT16-bound genes were enriched for mesenchymal differentiation across all lines, with

transcription factor binding and GTPase signaling uniquely enriched in the histone mutant

cells (**Fig 1B**). We focused subsequent analyses on the H3K27M lines cell lines, due to

fewer peaks with weaker enrichment in VUMC-DIPG10 cells (**Fig S1A**). E-box motifs and
motifs for EMT transcription factors were enriched at SPT16 binding sites, including key
DMG regulators TCF12 and OLIG2 [41-43], and the invasive glioma drivers SNAI1/2 [44, 45]
(**Fig 1C**). These findings suggest that FACT promotes transcriptional programs driven by
lineage-defining and EMT transcription factors in DMG.

We examined SPT16 occupancy at promoters genome wide, and assessed chromatin
accessibility at these regions using Assay for Transposase-Accessible Chromatin using
sequencing (ATAC-seq). The FACT complex is known to interact with H3K27M [19, 46].
Given that H3K27M and H3K27ac are known to incorporate into heterotypic H3K27M-
H3K27ac nucleosomes, bound by acetyl-binding BET proteins at enhancer elements [36,
47], we investigated FACT colocalization with these nucleosomes. Transcription start sites
(TSSs) were sorted by SPT16 enrichment and divided into two clusters which corresponded
to highly accessible TSSs and less accessible TSSs. SPT16, BRD4 and H3K27M were
enriched at accessible TSSs, which were also marked by the active histone modifications
H3K4me3 and H3K27ac (**Fig 1D and Fig S1B**). Examples of SPT16 binding to the TSS of
three DMG oncogenes (*MYC, PDGFRA,* and *MDM4*) are shown (**Fig 1E**).

To explore SPT16 and BRD4 co-occupancy, we visualized their enrichment at peaks
classified as either unique to each protein, or common sites bound by both proteins. As
expected, both SPT16 and BRD4 showed robust enrichment at the common peaks. Notably,
each protein was also enriched at peaks unique to the other when compared to the IgG
negative control, indicating co-localization of FACT and BRD4 at shared chromatin regions
(**Fig 1F and S1C**). To confirm the physical interaction between these proteins, we performed
endogenous co-immunoprecipitation of the FACT subunits, BRD4, and H3K27M from
nuclear extracts of three H3K27M cell lines (HSJD-DIPG007, SU-DIPGVI and SU-DIPGXIII).
BRD4 co-immunoprecipitated the FACT subunits consistent with previous reports in other
cell types [48-50] **(Fig 1G and Fig S1D)**. Furthermore, FACT and BRD4 were found to
interact with H3K27M, consistent with previous studies [19, 36, 46] **(Fig 1G and Fig S1D).**
Notably, FACT was only able to co-precipitate BRD4 and H3K27M in two of three cell lines
tested. This discrepancy may be due to transient and possibly indirect interactions between
FACT, BRD4, and H3K27M, making it challenging to reproducibly capture, especially when
examining endogenous protein levels under native conditions. Genes bound by SPT16 and
BRD4 were involved in transcription, chromatin remodeling and p53 signaling (**Fig S1E**).
Several example genes co-bound by FACT and BRD4 are shown at genes encoding the
BET protein *BRD2,* OPC lineage transcription factors *OLIG2 and SOX2*, and survival gene
BCL2L1 (**Fig 1H**).

Since DMG is heavily driven by epigenetic and transcription dysregulation, we hypothesized
that FACT and BET proteins, shown to be upregulated in other cancers [19, 20, 51-54],
would also be elevated DMG. Consistently, *BRD4, SSRP1* and *SUPT16H*, were significantly
overexpressed in DMG patients from the ZERO Childhood Cancer Program compared to
normal brain tissue [55, 56] (**Fig S1F**). However, their expression was not higher in DMG
than in other pediatric brain tumors, suggesting a broad role for these proteins in diverse
malignancies. These genes were also upregulated in H3K27M-DMG cell lines compared to
non-malignant cells, but not higher than in other pediatric cancer lines [57] (**Fig S1G**).
Furthermore, SSRP1 and BRD4 proteins were upregulated in a panel of DMG cell lines
compared to normal astrocytes (**Fig S1H**). Finally, whole genome CRISPR screening
revealed that *BRD4*, *SSRP1*, and *SUPT16H* are essential for cell viability (gene effect < -
0.5) of HGG cell lines from the Childhood Cancer Model Atlas, regardless of their histone
mutant status (**Fig S1I**). In summary, FACT and BRD4 are overexpressed, interact, and co-
localize at transcriptionally active chromatin, in the vicinity of key genes required for growth
and development. This prompted us to test the combination of FACT and BET inhibition as a
therapeutic strategy.

Combined FACT and BET inhibition synergistically decrease DMG cell proliferation

Treatment with either of the BET inhibitors JQ1 or trotabresib resulted in decreased
proliferation of primary DMG cells, with 72h IC_{50} values ranging from 1-50µM and 0.6-7µM,
respectively, across a range of DMG cultures (**Fig 2A-B and Table S2-3**). Further testing on
isogenic cell lines, with or without the H3K27M mutation, revealed significantly increased
sensitivity to single-agent CBL0137, JQ1, and trotabresib in H3K27M cells compared to
H3K27M KO cells. Notably, significant differences were observed in both IC_{50} and area
under the curve (AUC) values with different H3K27M mutational status, with a 5-fold
increase in sensitivity observed in H3K27M mutant compared to H3K27-KO DMG cells for
CBL0137 and JQ1, and a 4-fold increase for trotabresib treatment (**Fig 2C, Fig S2A and
Table S4**). These findings suggest that H3K27M-driven chromatin dysregulation increases
dependence on FACT and BRD4.

Next, we assessed the effect of combined inhibition of both FACT and BRD4 in the isogenic
histone-mutant and WT DMG cells. We found potent efficacy of the combination of CBL0137
and either BET inhibitor (JQ1 or trotabresib) in SU-DIPGXIII H3K27M mutant cell lines, and
this synergy was lost upon deletion of the H3K27M mutation (**Fig 2D-E**). A similar synergistic
effect between CBL0137 and the BET inhibitors was observed in four additional histone-

mutant, patient-derived DMG cultures (H3.3 K27M: SU-DIPGVI, HSJD-DIPG007, RA055;
H3.1 K27M: P005401) (**Fig 2F and Fig S2B-D**). However, we also found the combination to
be synergistic against histone wild-type (WT) HGG/DMG cultures (P016802, VUMC-
DIPG10, and AUS-DIPG17) (**Fig S2E-G).** These findings suggest that while H3K27M directly
contributes to dependency on FACT and BRD4, alternative oncogenic drivers in histone WT
tumors, such as MYCN amplification or CDKN2A/B loss, can create similar vulnerabilities to
chromatin-targeting therapies (**Table S5).**
214
The combinations also led to synergistic inhibition of colony formation, at concentrations of
each compound ranging between 100-200nM for HSJD-DIPG007 and RA055, and as low as
25-50nM for SU-DIPGVI (**Fig 2G-H and Fig S2H**). Lower drug concentrations were used in
clonogenic assays to model sustained, sub-lethal exposure over time, in contrast to higher
doses in short-term viability assays. Analysis of cell cycle distribution using flow cytometry
revealed that combined FACT and BET inhibition induced G0-G1 arrest, reducing the
proportion of cells in S, G2, and M phases, an effect driven by BET inhibition (**Fig. 2I-J, Fig.**
S2J-K). Additionally, Western blotting showed increased levels of cleaved PARP and
cleaved Caspase-3 following combination treatment, indicative of apoptotic cell death (**Fig.**
2K). Together, these data highlight the potential of combined FACT and BRD4 targeted
inhibition as an effective strategy in DMG.
226
227 **Combined FACT and BET inhibition causes chromatin condensation at gene**
228 **promoters in DMG.**
229
Given that CBL0137 and BET inhibitors target epigenetic and transcriptional regulators [17-
19, 58], we next sought to determine the effect of these drugs on chromatin accessibility and
transcription. To measure chromatin accessibility, ATAC-seq was performed in two H3.3
K27M DMG cell lines (HSJD-DIPG007 and SU-DIPGVI) treated with either CBL0137 or JQ1
as monotherapies or in combination after 4 hours to examine the early and direct changes to
chromatin at doses previously used in other mechanistic studies [20, 29, 35, 59, 60].
Combinatorial treatment of DMG with CBL0137 and JQ1 led to a decrease in ATAC-seq
signal at TSSs, consistently observed across independent replicates from both cell lines
tested (**Fig S3A-B**). This effect was more driven by CBL0137, with Principal Component
Analysis (PCA) revealing that the CBL0137 treated samples clustered closely with
combination treated samples (**Fig S3C-D**).
241
We next examined chromatin accessibility at SPT16 and BRD4 CUT&RUN peaks, finding
that both drugs alone, and in combination, decreased chromatin accessibility at these

genomic regions (**Fig 3A and Fig S3E-F**), with the combination significantly decreasing
chromatin accessibility at FACT and BRD4 binding sites more than each single agent in SU-
DIPGVI cells, an effect more strongly driven by CBL0137 (**Fig 3A, Fig S3F**). This suggests
that FACT and BRD4 are important for maintaining open chromatin, and that inhibition of
these factors with CBL0137 and JQ1 leads to chromatin closure/condensation at these sites.
Furthermore, differential accessibility analysis revealed that combination treatment resulted
in ~15-fold more closed chromatin regions compared to open regions in both cell lines tested
(HSJD-DIPG007: 11,704 regions closed vs. 828 regions opened; SU-DIPGVI: 41,457
regions closed vs. 2,620 regions opened; FDR < 0.05, |LFC| > 1) (**Fig 3B, Fig S3G-H**).
There was a significant degree of overlap in the regions closed by the combination treatment
in both the models **(Fig 3C)** (hypergeometric p-value = 0.008). These regions were enriched
at proximal gene promoters (**Fig 3D**) with their associated genes involved in pathways
critical for DMG tumorigenesis such as development, proliferation, and histone modification
(**Table S1C** and **Fig 3E**). For example, combination treatment led to chromatin condensation
at the promotor of the FACT-bound oncogenes *MYC*, *PDGFRA*, and *MDM4* (**Fig 3F).**
259
The H3K27M oncohistone hinders the spreading of H3K27me3 from PRC2-landing sites,
resulting in a decrease in H3K27me3 genome wide [2-4, 11]. We have previously shown that
CBL0137 increases global H3K27me3 levels and that this is potentiated by the HDACi
panobinostat [19]. Consistently, in this study, we observed an increase in global H3K27me3
levels induced by CBL0137 through western blotting (**Fig S3I-J**), and at specific loci using
CUT&RUN (p-val < 0.05) (**Fig S3K**). However, this increase was not further enhanced by
JQ1 (**Fig S3I-J**), suggesting that the mechanism by which the CBL0137 interacts with JQ1 is
distinct from its mechanism of synergy with panobinostat [19].
268
269 **Combined CBL0137 and JQ1 treatment leads to broad transcriptional silencing**
270
To assess the impact of chromatin condensation caused by CBL0137 and JQ1 treatment on
transcription, we performed nascent RNA-seq using TimeLapse-seq [61]. Unlike
conventional RNA-seq, where the effects of transcriptional inhibition can be masked by
stable pre-existing transcripts, TimeLapse-seq enables the specific identification of newly
synthesized mRNAs directly regulated by treatment [61]. Notably, the impact of each drug on
nascent transcripts was far more extensive than on total RNA (**Fig 4A, Fig S4A and Table**
S1D). Combination treatment had the most widespread effect compared to each
monotherapy, with over 60% of all detected genes being significantly (FDR<0.05, LFC< -1)
downregulated (9,159/13,153 for HSJD-DIPG007 and 8,930/13,919 for SU-DIPGVI),

indicating that CBL0137 and JQ1 induce a broad suppression of transcription (**Fig 4A and**
Fig S4A).

To investigate the nature of the combination response we overlapped the genes
downregulated by each treatment compared to DMSO and found that a large subset (38-
50%) of repressed genes were unique to the combination treatment, indicating that the
combined action of CBL0137 and JQ1 are required to silence these targets (**Fig 4B**). To
determine if the combination effect was additive or synergistic, we compared the fold-change
in gene expression between the combination and single agents. If the combination were
synergistic in silencing gene expression, the magnitude of fold-change in the combination
treatment would exceed the sum of the fold-changes observed with each single agent.
However, although the combination strongly correlated with the additive effect (R^2 = 0.68–
0.77; **Fig 4C, Fig S4B**), it did not surpass it. This suggests that, while the combination
suppresses more genes overall, the effect on individual genes is additive rather than
synergistic.

Examining individual genes revealed notable top upregulated genes induced by the
combination, related to stress (*GADD45B, EGR1, UBB*), neuronal differentiation (*TUBB3*),
and antigen presentation and interferon response (*HLA-E, IRF1/7, SOCS1*) (**Fig 4D, Fig**
S4C, Table S1D) with GADD45B and TUBB3 protein induction confirmed by western blot
(**Fig S4D**). Conversely, downregulated genes included several histone genes and
oligodendrocyte progenitor cell transcription factors (*OLIG1, SOX1/2*) (**Fig 4D, Fig S4C,**
Table S1D). Gene set enrichment analysis of the genes downregulated by combination
treatment revealed several critical processes involved in cancer cell growth and proliferation
such as DNA replication, DNA repair, histone modification, RNA modification, ribosome
biogenesis, and splicing (**Fig 4E** and **Fig S4E**). Mass spectrometry of CBL0137-treated
DMG cells showed concordant downregulation of proteins involved in translation and RNA
processing (**Fig S4F** and **Table S1G**). Integration of CUT&RUN with nascent RNA-seq
identified genes whose transcription is directly sustained by FACT and BRD4. Genes
associated with FACT and/or BRD4 binding exhibited strong transcriptional repression
following combination treatment (**Fig 4F, Fig S4G**), including critical DMG stem/progenitor
regulators OLIG2 and SOX1/2.

CBL0137 and JQ1 combination treatment represses transcription of key oncogenes

Among the directly downregulated targets were key oncogenes, including *MYC, PDGFRA,*
MDM4, and *SOX2* (**Fig 4F, Fig 5A**), which also displayed reduced chromatin accessibility at

FACT and BRD4-bound promoters/enhancers (**Fig S5A**). Transcriptional and protein

repression of these targets was validated in independent RNA-seq and western blot

experiments (**Fig 5B-C, Fig S5B-C, Table S1E**). MYC is an established transcriptional target

of both FACT and BRD4 [26, 58], PDGFRA is a known super-enhancer associated glioma

oncogene [35, 36], SOX2 is a potent stem cell transcription factor [62], and MDM4 is a p53

negative regulator with additional oncogenic roles [63, 64]. Given MDM4's role as a p53

inhibitor, we assessed p53 levels and observed CBL0137-mediated p53 protein upregulation

selectively in p53-wildtype HSJD-DIPG007 cells, but not in p53-mutant SU-DIPGVI cells (**Fig

S5E**).

We next investigated whether depleting MYC, MDM4, and PDGFRA could further enhance

the anti-proliferative action of the drug combination. To this end, we performed siRNA

knockdowns (KD) for each gene individually and in combination (triple KD) in SF8628

H3K27M DMG cells, achieving 40–70% knockdown (**Fig. S5F**), which was comparable to

the transcriptional repression observed with combination drug treatment (**Fig. S5G**). The

synergy between CBL0137 and JQ1 was confirmed in these cells (**Fig. S5H**). Treatment with

CBL0137 and JQ1 resulted in a significant reduction in cell proliferation, and this effect was

potentiated with depletion of *MYC*, *MDM4*, and *PDGFRA* (**Fig. 5D**). We next evaluated a

triple combination of CBL0137 and JQ1 with the clinically relevant MDM2/4 inhibitor

sulanemadlin. As expected, sulanemadlin exhibited high potency in p53 wild-type cells, while

isogenic HSJD-DIPG007 cells lacking p53 were more than 200-fold resistant [65] (**Fig. S5I**).

Similarly, p53-mutant DMG cell lines (SU-DIPGVI and SU-DIPGXVII) also displayed

resistance (**Fig. S5I**), confirming p53 restoration as the primary mechanism of action for

sulanemadlin in DMG. Cytotoxicity assays demonstrated that sulanemadlin synergized with

CBL0137 and JQ1 in both p53 wild-type HSJD-DIPG007 cells and in p53-mutant SU-DIPGVI

cultures (**Fig. 5E-F, Fig. S5J**). These findings uncover a novel therapeutic combination

involving MDM inhibition, identified through our investigation into the mechanisms of

CBL0137 and JQ1.

CBL0137 and JQ1 alter the splicing landscape in DMG

Since CBL0137 and JQ1 led to downregulation of splicing genes in our nascent RNA-seq

experiments (**Fig 4E**, **Fig S4E, Fig S6A**), and since chromatin structure is known to

influence splicing through a co-transcriptional process [66], we next examined alternative

splicing in an independent RNA-seq experiment. Nascent RNA-seq was performed after 2

hours of treatment, and splicing was analyzed at 4 hours to allow time for splicing to occur.

Single-agent treatment resulted in limited differential splicing. However, combination

treatment led to hundreds of differential splicing events compared to control cells, with an
average of 800 events in HSJD-DIPG007 cells and 2000 events in SU-DIPGVI cells. Some
of these splicing events occurred within the same gene, affecting approximately 400 unique
genes in HSJD-DIPG007 and 1000 unique genes in SU-DIPGVI (**Fig. S6B** and **Table S1F**).
The most common form of alternative splicing was intron retention, accounting for 40-50% of
events (**Fig S6C**). Examples of intron retention are shown at three loci, which also exhibited
condensed chromatin in the vicinity of the retained introns (**Fig S6D**). These findings suggest
that, in addition to disrupting chromatin dynamics and transcription, CBL0137 and JQ1 may
also alter the splicing landscape in DMG.

Dual inhibition of FACT and BRD4 enhances survival in orthotopic models of DMG

To evaluate the therapeutic efficacy of combined FACT and BET inhibition, we used three
orthotopic xenograft models using distinct patient-derived DMG cells: SU-DIPGVI-LUC,
HSJD-DIPG007, and RA055. The dosing schedule (CBL0137 every 4 days and JQ1 five
days a week) was based on the drug's pharmacokinetics, tolerability, and prior studies.
CBL0137's relatively longer half-life (~6h) supports infrequent dosing, while JQ1's shorter
half-life (~1h) necessitates more frequent administration [19, 36, 67-69].

In the SU-DIPGVI-LUC model, single-agent treatments with CBL0137 and JQ1 exhibited
either negligible or limited anti-tumor effects, respectively. However, the combination
regimen significantly extended survival compared to single-agent treatments and vehicle
cohorts by delaying tumor growth, as seen by xenogen imaging (median survival: vehicle: 73
days, CBL0137: 76 days, JQ1: 78 days, CBL0137/JQ1: 95.5 days; **Fig 6A-B** and **Table S6**).
The combination regimen was well tolerated, with no observed weight loss or overt
hematological or biochemical toxicities over the four-week treatment period (**Fig S7A-C,**
Table S9). We have previously demonstrated that CBL0137 can penetrate the blood–brain
barrier (BBB) [19], and here show that JQ1 is likewise detectable in the brain, as measured
by liquid chromatography–tandem mass spectrometry (LC–MS/MS) (**Fig S7D**).
Immunohistochemical analysis of brain tissue collected following 4 weeks of combination
treatment revealed a significant reduction in cell proliferation, as evidenced by decreased
Ki67 staining, and a concurrent reduction in H3K27ac staining (**Fig 6C-D**). Extension in
survival with combined FACT and BET inhibition was also evident in the HSJD-DIPG007
model, with the median survival of the combination-treated cohort extended to 93.5 days
compared to 70.5 days in the vehicle group (**Fig 6E, Fig S7B** and **Table S7**).

These promising results were recapitulated in the highly aggressive RA055 model,
characterized by a median survival of 46 days in untreated mice post intracranial injection.
Treatment with the combination of CBL0137 and JQ1 significantly enhanced survival, with
approximately one-third of treated mice surviving until the experimental endpoint of 201 days
(**Fig 6F**, **Fig S7D** and **Table S8**). Consistent with the SU-DIPGVI-LUC model, Ki67 cell
proliferation and H3K27ac immunohistochemistry staining were significantly reduced with
combination treatment compared to single-agent and untreated tissue samples (**Fig 6G-H**).
We also observed a significant reduction in MDM4 and MYC positive cells in PDX tumors
treated with the combination, validating our *in vitro* mechanistic findings in an independent *in*
vivo model (**Fig 6I-J**). Taken together, this combination strategy is effective against multiple
aggressive patient-derived DMG models. Furthermore, the combination of CBL0137 with the
clinically-relevant BET inhibitor ZEN-3694 [70, 71] was also well tolerated *in vivo*, with no
evidence of weight loss or toxicity (**Fig S7F-G, Table S10**), and pharmacokinetic analysis
showed that ZEN-3694 reaches the brain at levels comparable to plasma (**Fig. S7H**). These
results support the feasibility of translating combined FACT and BET inhibition into a
clinically actionable strategy for the treatment of DMG.

CBL0137 and JQ1 influence immune-associated pathways in DMG and tumor-
infiltrating immune cells
Interferon signaling plays a key role in anti-tumor immunity and involves upregulation of the
antigen presentation machinery (reviewed in [72, 73]), and epigenetic therapies can enhance
expression of the antigen presentation machinery in other cancers [64][74-77]. Similarly, in
DMG cells we observed that combination treatment upregulated transcription of interferon
response genes (*IRF1, IRF3, IRF7, SOCS1*) and MHC-I antigen presentation genes (*HLA-A,*
HLA-B, HLA-C, HLA-E) (**Fig 7A and Fig S8A**). as well as positive enrichment of the 'antigen
processing and presentation of exogenous peptide antigen' pathway (**Fig 7B**). To
complement our acute RNA-seq study, we examined expression changes following an
extended 48-hour treatment using qPCR. These results confirmed that CBL0137 elevates
expression of interferon and antigen presentation genes (*IRF1, HLA-A, IFNA, IFNB,* and
OAS1) (**Fig S8B**). Finally, flow cytometry confirmed increased in HLA-ABC and HLA-E
surface expression in response to combination treatment (**Fig 7C and Fig S8C**).
416
Given the prominence of immunosuppressive macrophages in DMG [15], and the epigenetic
regulation of their polarization [78], we next assessed whether CBL0137 and JQ1 modulate
the inflammatory response in cultured human monocyte-derived macrophages
(HMDMs).Treatment increased mRNA expression of pro-inflammatory M1 marker genes
(*IRF3, IRF8, IFNB, HIF1A, p53 and IL1B)* [79] (**Fig S8D**). Using flow cytometry, we show

that combination treatment increased surface expression of both the M1 marker CD86, a T

cell co-stimulatory molecule, and the M2 marker CD206, implicated in phagocytosis and

antigen cross-presentation [80] (**Fig 7D**). Moreover, combination-treated HMDMs displayed

increased phagocytic activity in a fluorescent bead uptake assay **(Fig S8E)**.

We investigated the effects of CBL0137 and JQ1 in an *in vivo* immunocompetent model.

C57Bl6 mice were orthotopically engrafted with syngeneic H3.3 K27M glioma cells as

previously described [73, 74], and tumors were allowed to establish. Following three weeks

of treatment, we performed *in situ* spatial transcriptomic profiling using the Xenium Prime

5000-gene panel. Single cells were clustered and annotated based on marker gene

expression and spatial localization (**Fig. S8F–H, Table S1H**), with a focus on tumor cells and

infiltrating immune cells. We identified three distinct tumor populations characterized by

elevated expression of astrocyte lineage markers (cluster 3: *Gfap, Sox9*), oligodendrocyte

lineage markers (cluster 9: *Sox10, Pdgfra, Mag*), and proliferation-associated genes (cluster

11: *Mki67, Top2a, Plk1*) (**Fig. S8H-I, Table S1I).** Among CD45+ (*Ptprc*+) immune cells, we

identified tumor-associated macrophages (TAMs; cluster 6: *Cd68, Itgam, Cx3cr1*) and

infiltrating T cells (cluster 24: *Il2rb, Cd3d/g, Cd4, Cd8a/b1, Gzma/b*) (**Fig. S8H, Table S1H).**

Combination treatment depleted both tumor and tumor-associated immune cell populations

(**Fig. 7E–F**) and altered the cellular composition within the tumor microenvironment (**Fig.**

7G). Notably, there was a decrease in the tumor-to-immune cell ratio, as well as a relative

increase in T cell abundance within the tumor following combination therapy **(Fig. 7G)**. To

investigate transcriptional changes, we subclustered each cell type and performed

differential gene expression analysis comparing vehicle- and combination-treated cells. In

the tumor compartment (comprising all three tumor clusters), we observed downregulation of

chromatin associated genes, transcriptional factors, and stem cell maintenance genes (**Fig.**

S8J). Leading edge genes driving enrichment included *Myc* and *Sox2* (**Fig. S8K, Table**

S1J), consistent with our *in vitro* findings which showed that FACT and BRD4 inhibition

epigenetically silence these critical oncogenic drivers. Furthermore, the top upregulated

gene was the interferon stimulated gene *Ifi27* (**Fig. S8K, Table S1J),** further supporting the

induction of an interferon response in tumor cells by combination therapy.

In TAMs, there was positive enrichment of genes involved in phagocytosis (including *Ighm*

and *C3*), antigen presentation, and T cell activation – hinting at tumor cell engulfment and

antigen cross-presentation to T cells [81-83] (**Fig 7H, J, Table S1J**). Furthermore, T cells

displayed transcriptional signatures consistent with infiltration and activation, including

pathways related to cellular extravasation, T cell selection and immune activation. This

included *Ptprc* and *Dock2* (**Fig 7I, K, Table S1J**). CD45, encoded by *Ptprc*, plays a critical
role in T cell activation by initiating T cell receptor signaling, and DOCK2 is involved in
formation of T cell immune synapses [84, 85]. Finally, immune cells were localized within
close proximity to endothelial cells, together with their transcriptional profiles, this supports
the notion of extravasation from blood vessels into the tumor. Visualization of individual
transcripts for phagocytic and T cell activation genes (*Ighm, C3, Ptprc* and *Dock2*) confirmed
increased expression within TAMs and T cells upon combination treatment (**Fig. 7L**).
466
467 Taken together, combination treatment exerts coordinated effects on both glioma and
468 immune compartments – suppressing transcriptional stem cell programs and enhancing
469 antigen presentation and interferon signaling in tumor cells, while simultaneous
470 reprogramming TAMs towards a more phagocytic and immunostimulatory state and
471 promoting T cell activation.
472

Discussion

Despite hundreds of clinical trials, survival outcomes for DMG patients have not changed in
over three decades [15]. The current standard-of-care, radiotherapy, only delays tumor
progression and the majority of children still survive less than one year from diagnosis. Since
the discovery of the H3K27M mutation in 2012, efforts to target multiple layers of epigenetic
regulation, including DNA methylation [12], histone modification [86-89], chromatin readers
[35, 36], chromatin remodelers [90-92], and transcription machinery [35, 93], have been
ongoing. Given DMG's aggressive, heterogeneous nature and multiple undruggable
oncogenic pathways, monotherapies are unlikely to be effective. Here we have uncovered a
novel mechanism-anchored epigenetic combination therapy - FACT and BET inhibition -
which simultaneously disrupts multiple oncogenic pathways and demonstrates potent pre-
clinical activity against DMG.
485
FACT disassembles and reassembles nucleosomes during cellular processes that require
DNA access, such as during transcription or replication. This overcomes the physical
hindrance of nucleosome-compacted DNA, allowing RNA Pol II or the replication machinery
to travel through chromatin unimpeded [22, 23]. Our findings indicate that FACT maintains
open chromatin at promoters and enhancers, supporting transcription driven by
developmental TFs (TCF12, OLIG2, SNAI1) in DMG. This is consistent with a recent study
showing that FACT preserves open chromatin at regulatory elements to promote
transcription by core pluripotency factors in embryonic stem cells [94]. BRD4 binds to

acetylated chromatin via its tandem bromodomains and recruits positive transcription
elongation factor (P-TEFb) to active enhancers, releasing paused RNA pol II, leading to
transcriptional initiation and elongation [95]. Combined targeting of FACT and BRD4 led to
chromatin compaction and widespread transcriptional silencing, including several notable
oncogenes. For example, *MYC and PDGFRα*, are both super enhancer-activated oncogenes
involved in gliomagenesis [12, 96]. Another downstream target was *MDM4*, a negative
regulator of p53, suggesting this treatment could potentially restore p53 function in p53-WT
DMGs [63]. Thus, this drug intervention simultaneously disarms multiple epigenetic-driven
oncogenes in DMG. However, it is unlikely that the cytotoxic effects arise solely from
silencing several genes. Rather, we propose that the global disruption of transcriptional
output – a fundamental stressor in rapidly dividing cells – is the key driver of cell death.

While off-target effects cannot be fully excluded, the consistent phenotypic and
transcriptomic responses across multiple models suggest that the therapeutic effects are
primarily driven by on-target inhibition of transcriptional regulation. Model-specific differences
in transcriptional response and drug sensitivity may reflect intrinsic features such as p53
status, with HSJD-DIPG007 (p53 WT) showing more JQ1-driven changes and SU-DIPGVI
(p53 mutant) more CBL0137-driven effects.

The combination of CBL0137 with either of the BET inhibitors tested had higher efficacy
against H3K27M cells compared to isogenic cells lacking H3K27M. This could be explained
by two non-mutually exclusive hypotheses: i) Direct inhibition of H3K27M function by
targeting co-factors like FACT and BRD4, or ii) increased dependency on these factors due
to downstream epigenetic reprogramming. The first hypothesis is supported by prior studies
and our data showing chromatin localization and co-immunoprecipitation of FACT and BRD4
with H3K27M. An IP-mass spectrometry study previously identified SPT16 and SSRP1 as
H3K27M interactors in mitotic DMG extracts [46], and additional studies have reported
interactions between H3K27M and both FACT and BRD4 [19, 36, 46]. Alternatively, the
hypomethylated chromatin creates a dependency on FACT and BRD4 to maintain open
chromatin and high levels of transcription. Treatment with CBL0137 and JQ1 then leads to
chromatin compaction at transcription start sites, leading to transcriptional collapse. This
mechanism mirrors recent studies targeting the SWI/SNF chromatin remodeling complex,
where the epigenetic reprogramming characteristic of DMG creates a dependency on this
complex to drive pathogenic gene expression [90-92]. Notably, CBL0137 and JQ1 were still
effective against histone WT DMG, and both FACT and BRD4 emerged as critical
dependencies in pHGG cell lines irrespective of H3 mutation status. The synergy of this
combination has also been observed in other H3 WT pediatric brain tumors, namely

medulloblastoma and Atypical Teratoid/Rhabdoid Tumor (AT/RT) cell lines [97], indicating
potential broader therapeutic applications in pediatric brain cancers beyond H3K27M DMG.
533

We also uncovered a role for this combination treatment in alternative splicing, possibly
through downregulation of the splicing machinery. As splicing is influenced by chromatin
structure, we hypothesize that alternative splicing may also result from transcriptional
stalling. Specifically, RNA pol II may encounter increased resistance when transcribing
through compacted chromatin, leading to splicing errors [66], particularly in the context of
reduced expression of splicing genes. This hypothesis warrants further exploration, for
example by mapping RNA pol II elongation rates. 3D genome architecture is an emerging
contributor to tumorigenesis in epigenetically driven cancers, including DMG and
ependymoma [41, 98]. Both CBL0137 and BET inhibitors can disrupt topologically
associated domains (TADs) and enhancer–promoter loops [41, 60]. Although not explored in
this study, examining the impact of combination therapy on genome topology represents an
exciting avenue for future investigation.
546

The *in vivo* efficacy of the combined FACT and BRD4 inhibition in multiple orthotopic models
of DMG indicates its potential as a promising combinatorial treatment strategy. Employing an
alternative dosing regimen involving an increased dose of CBL0137 (40 mg/kg) paired with a
reduced dose of JQ1 (25 mg/kg) not only proved well-tolerated but also resulted in a
significantly enhanced survival response in highly aggressive DMG models. This
underscores the importance of exploring diverse dosing regimens for the development of
combination therapies.
554

Epigenetic therapies have been shown to activate an immune response through various
mechanisms [99]. For example, FACT, HDAC and DNMT inhibitors, have been shown to
trigger an interferon response and enhance expression of the antigen presentation
machinery in other cancers [64][74-77]. While BET inhibitors, such as JQ1, have been
associated with suppression of interferon signaling due to downregulation of key interferon-
stimulated genes in glioblastoma [100], we observed transcriptional activation of the
interferon pathway in DMG cells treated with the combination of CBL0137 and JQ1. This
suggests CBL0137 may counteract the interferon-suppressive effects of BET inhibition,
possibly through induction of double-stranded RNA (dsRNA), which is a known trigger of the
type I interferon response. Indeed, CBL0137 has been shown to induce dsRNA and activate
antiviral pathways in various non-DMG cancer model [27, 101, 102]. This is particularly
relevant given our observation of increased intron retention, which may generate
neoantigens [103, 104]. When presented via MHC-I, these can enhance tumor immune

visibility. Consistent with this, we detected increased MHC-I surface expression on DMG
cells following treatment. Together, these findings support a model whereby chromatin
disruption by CBL0137 induces splicing errors and/or dsRNA, leading to interferon activation
and increased antigen presentation.
572
In macrophages, we observed upregulation of both M1- and M2-like markers, suggesting
that the drug combination influences the plasticity of TAMs, which are known to adopt mixed
or transitional states [105, 106]. In an immunocompetent mouse model, we observed
decreased tumor burden and immune rewiring consistent with our *in vitro* findings. This
included TAM reprogramming toward a phagocytic phenotype, as well as T cells exhibiting
heightened infiltration and activation signatures. Altogether, our data indicates that this
combination elicits both tumor-intrinsic and immune-mediated effects. By simultaneously
disrupting oncogenic transcriptional programs and enhancing immune detection, CBL0137
and BET inhibition may offer a dual therapeutic strategy for DMG and other immunologically
'cold' pediatric brain tumors, particularly in combination with immunotherapies such as CAR
T cells.
584
Our data nonetheless highlights the therapeutic potential of this combinatorial approach of
BETi and FACTi. CBL0137 is currently under clinical investigation in pediatric DMG
(PEPN2111), and our findings provide a strong mechanistic rationale for future combinations
with transcriptional or chromatin-disrupting agents to enhance therapeutic efficacy. Together,
this work lays a foundation for novel strategies to exploit the epigenetic dependencies of
DMG and improve outcomes for children with this devastating disease.

Materials and Methods

Cell culture
DMG cells were maintained as neurospheres in Tumor Stem Cell (TSM) media consisting of
a 1:1 mixture of DMEM/F12 and Neurobasal medium (Invitrogen) supplemented with
Glutamax, Pyruvate, non-essential amino acids, HEPES buffer, antibiotic/antimycotic, B-27
(Invitrogen), Heparin (0.0002%), EGF (20 ng/mL), bFGF (20 ng/mL), PDGF-AA (10 ng/mL)
and PDGF-BB (10 ng/mL) (StemCell Technologies). SU-DIPGXIII K27M and K27M-KO
isogenic cells were maintained according to [12], adherently on flasks coated with poly-L-
ornithine (0.01%) (Sigma) and laminin (10 µg/mL) (Sigma), in Neurocult NS-A proliferation
media (StemCell Technologies) supplemented with heparin (0.0002%), EGF (20 ng/mL) and
bFGF (10 ng/mL) (StemCell Technologies). SF8628 cells were maintained adherently in
DMEM supplemented with 10% FBS and 2mM L-Glutamine. All lines were cultured at 37°C

and 5% carbon dioxide. Cells tested negative for mycoplasma contamination and were
confirmed to match their original tumors by STR profiling.

Human monocyte-derived macrophages (HMDMs) were a kind gift from Prof. Kerry-Anne
Rye, Cardiometabolic Research Group, UNSW, following isolation and culture as previously
described [107], maintained in RPMI + 10% human serum with the media being replaced
every 3 days. In polarisation studies, HMDM were either primed for 48 h with 1 µg/mL LPS
(Sigma) and 20 ng/mL recombinant human IFNγ (Sigma) for the M1 phenotype, 20 ng/mL
recombinant human IL-4 (Sigma) and 20 ng/mL recombinant human IL-13 (Sigma) for the
M2 phenotype.

CUT&RUN
Neurospheres were collected by centrifugation at 800 rpm for 3 min and dissociated into
single cells with 1 mL Accutase (Sigma) for 5-10 minutes with occasional pipetting, diluted
with 5 mL TSM-B, and passed through a 40 µm cell strainer. Cells were centrifuged at 1,200
rpm for 5 min and resuspended in complete media. Single cell suspensions were counted
using a haemocytometer and cell viability was confirmed to be over 95%. CUT&RUN was
performed using the EpiCypher CUTANA CUT&RUN kit according to the manufacturer's
instructions. Briefly, 500,000 viable cells per reaction were washed and permeabilized with
0.01% Digitonin. Cells were incubated with 0.5 µg antibody, or a 1:50 dilution for SPT16
(**Table 1**), on a nutator overnight at 4°C. Following washing, pAG-MNase was added to the
reactions and $CaCl_2$ was used to activate antibody-bound MNase for targeted chromatin
digestion and release of DNA fragments. The entire volume of purified DNA (ranging from 1-
10 ng) was used as input for NGS library preparation, according to the EpiCypher CUTANA
Library Prep Kit instructions. Library concentration was determined using a Qubit
Fluorometer (HS dsDNA) (Thermo) and library size determined using a 4200 Tapestation
system on a high sensitivity D1000 ScreenTape (Agilent). Resulting libraries were
sequenced with 50-150bp paired-end reads, with a minimum of 7M reads per sample at
Ramaciotti Centre for Genomics (UNSW Sydney, Australia). Two biological CUT&RUN
replicates were performed.

CUT&RUN analysis
Quality control analysis of CUT&RUN reads was performed using fastQC (v0.11.9). Adapter
trimming was performed using cutadapt (v4.2) with parameters: --error-rate 0.1 --times 1 --
overlap 3 --action trim --minimum-length 17 --pair-filter any. Reads were aligned using
bowtie2 (v2.5.0) with -k 1 against the GRCh38 human reference genome. BAM files were
sorted, and optical duplicates marked using picard (v2.27.5) with parameters: --

DUPLICATE_SCORING_STRATEGY   SUM_OF_BASE_QUALITIES   --

OPTICAL_DUPLICATE_PIXEL_DISTANCE 100 --VALIDATION_STRINGENCY SILENT.

Marked duplicates were filtered and removed using samtools (v1.12). Peak calling was

performed using MACS2 (v2.2.7.1) using IgG as the control. Narrow peaks parameters: --

keep-dup all -p 1e-5 and broad peak using parameters: --keep-dup all -q 0.05 --broad --

broad-cutoff 0.05. BigWig coverage files were generated using deeptools bamCoverage

(v3.5.1) with parameters: --binSize 10 --normalizeUsing BPM --effectiveGenomeSize

2913022398. Broad peak settings were used for H3K27M and H3K27me3, and narrow

peaks for SPT16, BRD4, H3K27ac, H3K4me3. Peaks from biological replicates were

merged using BedTools [108].

BigWig coverage was visualized using DeepTools [109] at hg38 Ensembl genes with

blacklisted CUT&RUN regions excluded [110], or at SPT16 and BRD4 MACS2 peaks.

Integrated genomics viewer (IGV) [111] was used to visualize BigWig coverage at individual

loci. Peaks were annotated using ChIPSeeker [112] and ClusterProfiler [113] was used for

Gene Ontology over-representation analysis of the annotated genes. Transcription factor

motif analysis was performed using the Cistrome SeqPos motif tool [114]. Differential binding

analysis for H3K27me3 comparing between DMSO and CBL0137 treated cells was carried

out using CSAW as described [115], filtering on genomic windows 5-fold higher than

background, with a significance threshold of FDR < 0.05 (n=2).

Co-immunoprecipitation (co-IP) assay

Co-IP was performed as described in [116]. Briefly, nuclei were isolated from 5 million cells

with hypotonic buffer (20 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, protease inhibitors

(Roche)) for 15 min on ice, followed by addition of IGEPAL (0.5%; Sigma-Aldrich). Nuclei

were collected by centrifugation (5 min, 1000 *x g*, 4°C) then lysed in non-denaturing lysis

buffer (50 mM HEPES pH 7.4, 140 mM NaCl, 3 mM $MgCl_2$, 1 mM EGTA, 1% Triton-X, 10%

Glycerol, protease inhibitors (Roche)). The nuclear lysate was passed 6 times through a 25-

G needle to shear the DNA, then incubated for 15 min on ice. Lysates were cleared by

centrifuging at 13,000 rpm (top speed) for 15 min at 4°C to remove insoluble debris. The

lysate supernatant was then incubated with 15 µL protein A/G magnetic beads (Pierce) pre-

conjugated with 2.5 µg antibody (minimum 4 h, 4°C with rotation). Immunoprecipitation was

carried out overnight at 4°C on a rotator. Magnetic beads were washed 3x with lysis buffer,

and protein was eluted with 2X Laemmli Sample Buffer diluted in lysis buffer, boiled at 85°C

for 10 min, followed by addition of DTT (100 mM; Sigma-Aldrich). Western blotting to detect

immunoprecipitated proteins was performed as described below, except that VeriBlot

(Abcam) was used for the secondary antibody at 1:1000.

Western blotting

To extract total protein cells were incubated with RIPA buffer (CST) containing protease and
phosphatase inhibitors (Roche) for 15 min on ice with occasional vortexing, followed by
centrifugation at 13,000 rpm for 15 min at 4°C to remove debris. BioRad 4-20% proTEAN
pre-cast gels were used for gel electrophoresis, and proteins were transferred for 1 h at 100
V onto 0.45 μm PVDF membranes for RIPA-soluble proteins, or 0.45 μm nitrocellulose
membranes for histones. Membranes were blocked for 1 h in 5% milk diluted in Tris Buffered
Saline with 0.01% Tween-20 (TBS-T). Western blotting was carried out overnight with the
indicated antibodies (Table 1) diluted 5% Bovine Serum Albumin (BSA; Sigma-Aldrich) in
TBS-T. Following washing, membranes were incubated in horseradish peroxidase-
conjugated secondary antibody (CST) diluted in 5% skim milk/TBS-T (1:2000) for 1 hr at
room temperature. Membranes were incubated with ECL substrate (Pierce) and bands
imaged using a Bio-Rad ChemiDoc imaging system. Densitometry was carried out using
Image Lab software, with local background subtraction, and normalization to the indicated
housekeeping gene.

Gene expression analysis from patient samples and cell lines

Patient expression data for *SSRP1, SUPT16H* and *BRD4* was obtained from the ZERO
childhood cancer precision medicine program [55]. Normal brain expression data was
obtained from McGill University [56]. Integrated RNA-seq expression data were batch
corrected using ComBat-Seq from the Bioconductor package sva (v3.48.0). *SSRP1,*
SUPT16H and *BRD4* expression in DMG and non-malignant cell lines was downloaded from
the Childhood Cancer Model Atlas [57].

CRISPR dependency in HGG cell lines

The CRISPR dependency scores for BRD4, SSRP1 and SUPT16H are obtained from
Childhood Cancer Model Atlas (https://vicpcc.org.au/dashboard; CCMA [57]). The results are
part of the whole-genome scale CRISPR screens for n=24 high grade glioma cell lines
following the protocol described in [57, 117]. Data was obtained under the data transfer
agreement with the CCMA team. Briefly, the primary cells were engineered with the Cas12
endonuclease and Blasticidin antibiotic resistance-expressing genes (Lenti-AsCpf1-Blast:
1xNLS-Cas12a lentiviral expression construct; Addgene #84750). The whole genome
libraries used were Human CRISPR Knockout Humagne Set C and Set D Pooled Library
(Addgene #172650 and #172651). Pooled knockout genetic screens were conducted over
28 and 35 days at a multiplicity of infection (MOI) of 0.3. The gene dependency score was
determined using the chronos method [118], and scaling was performed across the whole

CCMA cohort using the negative control genes and core essential genes (Achilles Common

Essential Controls) from the depmap portal (https://depmap.org/portal; [119]).

Cell Viability Assays

Cells were cultured in 96-well plates at a density of 2,500–3,000 cells per well. For adherent

cells, drugs were added 24 hours after plating; for suspension cells, treatment was initiated

72 hours post-plating. After a 72-hour incubation period, cell proliferation was assessed

using the resazurin assay (Sigma-Aldrich), with results expressed as a percentage relative to

untreated controls. For combination treatments with CBL0137 (from Prof Andrei Gudkov and

Prof Katerina Gurova) and BET inhibitors (JQ1 [Selleck S7110] and trotabresib [Synonym:

CC-90010; MedChemExpress HY-137573]), cells were exposed to serial dilutions of each

drug, and synergy was calculated using CalcuSyn, where CI values < 1 indicate synergy.

For triple combination treatments with CBL0137, JQ1, and Sulanemadlin (Synonym: ALRN-

6924; MedChemExpress HY-P4210), cells were treated for 96 hours in a matrix format, and

BLISS synergy scores was assessed using SynergyFinder [120].

Clonogenic Assays

Long-term clonogenic assays were conducted by plating 24-well plates with a base layer of

0.3 mL of 0.5% agar mixed with cell culture media. After 24 hours of incubation at 4°C, the

base layer was overlaid with a 0.4 mL layer consisting of cells (500-1500 cells/well), 0.33%

agar, cell culture media, and the drug at the indicated concentrations. The plates were then

incubated in a 5% CO_2 humidified atmosphere at 37°C. Following an incubation period of

either 2 or 4 weeks, depending on the specific cell line, the colonies were stained with 3-

(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) solution (Sigma-Aldrich) at

5 mg/mL. After 1 hour, images of the colonies were captured using Image Lab Software.

Colony quantification was performed using Image J software, and the data were presented

as a percentage relative to untreated colonies.

Cell cycle analysis

DMG cells were treated for 48 hours with CBL0137, JQ1, or trotabresib at cell line-specific

concentrations. For SU-DIPGVI cells, 1 µM for each drug. For RA055, the doses were

CBL0137 (0.4 µM), JQ1 (0.1 µM), and trotabresib (0.2 µM). HSJD-DIPG007 cells were

treated with CBL0137 (0.2 µM), JQ1 (1 µM), and trotabresib (1 µM). Cell cycle analysis of

treated DMG cells was performed as previously described [121].

ATAC-seq

Neurospheres were treated for 4 h with CBL0137 and JQ1 (HSJD-DIPG007 1 µM CBL0137
1 µM JQ1; SU-DIPGVI 2 µM CBL0137 1 µM JQ1). DMSO was added to the control and
single drug treatments to equal concentration in all conditions (maximum 0.03% DMSO).
Cells were collected for both ATAC-seq and RNA-extraction. Tn5 Transposition was
performed as described in the OMNI-ATAC-seq protocol [122]. Briefly, single cell
suspensions from neurospheres were prepared as described for CUT&RUN and washed in
ice-cold PBS and cell viability was confirmed to be over 95%. Nuclei from 50,000 viable cells
were isolated, permeabilized and transposed with 2.1 µL Tn5 (Illumina) for 30 min with
mixing (1000 rpm). DNA was then purified, and PCR amplified (7-14 cycles) with sequencing
primers to generate libraries for sequencing. Library concentration was determined using a
Qubit Fluorometers (Thermo) (HS dsDNA) and library size determined using a 4200
Tapestation system on a high sensitivity D1000 ScreenTape (Agilent). ATAC-seq libraries
were sequenced with a minimum of 50 million paired-end reads per sample and read length
100bp. ATAC-seq experiments were performed in biological duplicate (HSJD-DIPG007) or
triplicate (SU-DIPGVI).
765
ATAC-seq analysis
ATAC-seq reads were trimmed with Trimmomatic
(ILLUMINACLIP:Nextera.fa:2:30:10:8:TRUE   SLIDINGWINDOW:5:20   LEADING:3
TRAILING:3 MINLEN:20 CROP:130 HEADCROP:150). Trimmed reads were mapped with
bowtie2 using default settings [123] and the resulting BAM files deduplicated using
Sambamba markdup [124]. Deduplicated BAM files were then filtered to remove reads with
fragment sizes greater than 100bp and CPM BigWigs were generated using the DeepTools
bamCoverage [109].

BigWig coverage was visualized as heatmaps and profile plots using DeepTools [109] at
hg38 Ensembl genes and at SPT16 peaks and BRD4 peaks. PCA clustering of ATAC-seq
average signal over gene bodies was performed using DeepTools. Differential accessibility
analysis was carried out using CSAW as described [115], filtering on genomic windows 5-
fold higher than background. Regions with an FDR<0.05 were considered significantly
different, with 'opened' regions logFC > 1 and 'closed' regions logFC < -1. Closed regions
between the 2 models were overlapped using BedTools [108], annotated using ChIPSeeker
[112] and ClusterProfiler was used for Gene Ontology over-representation analysis of the
annotated genes [113]. BigWig coverage was visualized at individual gene loci on IGV [111].
A hypergeometric test was used to test if the overlap in 'closed regions' between the 2
models was significantly over-enriched (k = 6119 (overlap), s = 11704 (closed in HSJD-

DIPG007), M = 41,454 (closed in SU-DIPGXIII), N = 80,894 (number of windows analyzed),

where expected = (s*M)/N.

RNA-seq

DMG cells were treated with CBL0137 and JQ1 for 4 h as for ATAC-seq. Cells were

collected for both ATAC-seq and RNA-extraction at the same time. Total RNA was extracted

from treated cells using the ISOLATE II RNA Mini Kit from Meridian Bioscience. RNA

concentration and purity was determined using a NanoDrop Spectrophotometer and integrity

measured using a 4200 Tapestation system. cDNA libraries were prepared using the

Illumina Stranded mRNA prep kit. RNA-seq libraries were sequenced with 100bp paired-end

reads, with a minimum of 50 million reads per sample. Three biological replicates were

performed for all RNA-seq experiments.

RNA-seq analysis

Illumina paired-end fastq files were aligned to the human genome GRCh38 and gencode

v41 using STAR (v2.7.8a) in two-pass mode with quantMode set to TranscriptomeSAM

[125]. Transcriptome alignment was quantified using RSEM (v1.3.3) rsem-calculate-

expression to obtain raw read counts, and normalized transcripts per million (TPM) and

fragments per kilobase million (FPKM) values [126]. Degust (10.5281/zenodo.3258932) was

used for differential expression analysis, using edgeR for normalization. Genes with low

expression were filtered out using a cutoff of 0.5 counts per million (CPM) in at least 3

samples. Genes with FDR<0.05 and absolute logFC>0.585 (1.5-fold) were considered

significantly differentially expressed. Multi-dimensional Scaling (MDS) clustering was

performed using Degust. ClusterProfiler was used for Gene Ontology over-representation

analysis of the differentially expressed genes [113].

MINTIE [127] was used for differential splicing analysis using parameters assembler=soap,

min_logfc=2, min_cpm=0.1, fdr=0.05, and run_de_step=TRUE, with all 3 Vehicle replicates

used as control. BedTools [108] was used to determine the common events between the 3

biological replicates. BAM files were visualized on IGV [111] ensuring the same scale.

TimeLapse RNA-seq

DMG cells were treated with CBL0137 and JQ1 (HSJD-DIPG007 1 µM CBL0137 1 µM JQ1;

SU-DIPGVI 2 µM CBL0137 1 µM JQ1) for 2 h. At the same time as drug treatment, 4-

Thiouridine (4sU) (ab143718) was added to a final concentration of 100 µM. Cells without

4sU or drug were used as a control. RNA was isolated using a modified version of the

RNeasy protocol (Qiagen) as described in [61]. RNA (2 µg) was then subjected to Timelapse

chemistry to convert 4sU to C [61]. Libraries were prepared using the Illumina Stranded
Total RNA plus RIboZero Plus kit and sequenced on 1 lane of a NovaSeq X Plus 10B
2x150bp, with a minimum 50 million reads per sample. Three biological replicates were
performed for TimeLapse RNA-seq experiments.

TimeLapse RNA-seq analysis
Reads were quality and adapter trimmed using trim-galore 0.6.10 using options –quality 20 -j
4 –paired. Trimmed reads were aligned to the human reference genome hg38 with star [125]
2.7.10 using options --runThreadN 23 --runMode alignReads --readFilesCommand zcat --
outReadsUnmapped Fastx --outSAMattributes nM MD –outFileNamePrefix --
peOverlapNbasesMin 10 --outFilterMismatchNmax 999 --outFilterMultimapNmax 1 --
alignEndsType EndToEnd --outFilterMismatchNoverReadLmax 0.3 --outSAMtype BAM
SortedByCoordinate and indexed using samtools 1.6. BAM files were converted to CIT
format using the gedi Bam2CIT function. 4sU induced conversions were called using
GrandSlam [128] using options -plot -full -D -allGenes -trim5p 5 -trim3p 5. The gene-wise
estimates for the new-to-total ratio were input into GrandR [129]. Nascent differential
expression was performed using the PairwiseDESeq2 function, with normalization with
respect to total. P values were adjusted using Benjamini–Hochberg correction and a cut-off
of p.adj < 0.05 and |LFC| > 1 were used to call differentially expressed genes.

Label-free Total Proteomics Analysis
Following 24 h drug treatment (IC50 doses), cell pellets were washed 2x in PBS and
incubated in Lysis Buffer (Cell Signalling, #98030) for 30min on ice with occasional vortexing
followed by centrifugation at 13,000 rpm for 15 min at 4°C. Protein concentration was
measured with Pierce BCA Protein Assay kit following the manufacturer's instructions
(ThermoScientific, #23225). From each sample, 50µg protein was used for total proteomics
analysis. Mass spectrometry analysis was performed at the Bioanalytical Mass Spectrometry
Facility at UNSW, Sydney, Australia. Protein masses were detected by an Orbitrap Fusion
Lumos Tribrid mass spectrometer (Thermo Scientific, Bremen, Germany) as described
previously [130].

Protein peaks were identified using MaxQuant Software and the Andromeda Algorithm
version 2.4.13.0 [131] with the following standard settings: oxidation and acetyl (Protein N-
term) as variable modifications and carbamidomethyl as fixed modification. Digestion mode
was set to trypsin/P with a maximum of 2 missed cleavages. The main search peptide
tolerance was +/- 4.5ppm and the isotope match tolerance was set to 2ppm. Protein peaks
were searched against the reference human proteome (UniProt, UP000005640). Label-free

Quantification (LFQ) was used with standard parameters (minimum ratio 2 and maximum
ratio 5) to identify proteins.

Subsequent downstream analysis on the proteomics dataset was performed using Perseus
Software (v.2.0.11) to identify differential protein expression [132]. Unnecessary and
incorrect protein identifications were filtered out from the data set that were classified as
potential contaminants, reverse or only identified by site. The data was then log transformed
with base parameter 2, non-valid values were filtered out, and missing values were replaced
from normal distribution with 0.3 width and 1.8 downshift standard parameters. Two-sample
T-test was used to compare expression levels between vehicle and drug-treated samples.
Proteins with a -log p value ≥ 2 were included in downstream pathway analysis. Gene Set
Enrichment Pathway analysis was performed in R (v.4.4.2) using the ClusterProfiler package
[113].

siRNA transfection

SF8628 cells were seeded at a density of 10,000 cells/cm^2 in 24 well plates in antibiotic free
media. The following day, siRNA constructs (Dharmacon) were transfected into the cells
using DharmaFECT-1 transfection reagent according to the manufacturer's instructions at a
concentration of 20nM. siRNA used were ON-TARGETplus Non-targeting Control Pool (D-
001810), ON-TARGETplus MYC (4609) SMARTpool, ON-TARGETplus MDM4 (4194)
SMARTpool, and ON-TARGETplus PDGFRA (5156) SMARTpool. For simultaneous
knockdown of all 3 genes, total siRNA concentration of 60nM was used. Culture media was
changed the following day.

qPCR
Total RNA was extracted using the ISOLATE II RNA Mini Kit (Meridian Bioscience), and first-
strand cDNA was synthesised using the SensiFAST cDNA Synthesis Kit (Meridian
Bioscience). Quantitative PCR (qPCR) was performed with the SensiFAST SYBR No-ROX
Kit (Meridian Bioscience) using primer sequences listed in **Table 2,** synthesised by
Integrated DNA Technologies using a QuantStudio 3 qPCR machine. Relative mRNA
expression levels of genes of interest were normalised to corresponding housekeeping
genes using the ΔΔCt method [133]. qPCR experiments were performed in biological
triplicate.

Timelapse microscopy proliferation assay
SF8628 cells were transfected with siRNA in duplicate wells, and the following day, the
media was changed, and cells were treated with CBL0137 (0.2µM) and JQ1 (0.5µM). Cells

were grown for 6 days, and phase-contrast confluence was monitored by time-lapse
microscopy every 6 h using an Incuyte S3 Live Cell Analysis System. Incucyte experiments
were repeated three times.

Orthotopic model/drug treatment

The protocol for intracranial injections was implemented as described in the study conducted
by Khan et al. [121]. The timing of treatment initiation varied for each model with SU-
DIPGVI-Luc at 7 weeks, RA055 at 3 weeks. and HSJD-DIPG007 starting at 5 weeks post
intracranial surgery.

In the SU-DIPGVI-Luc and RA055 models, the drug treatments were as follows: CBL0137 -
dissolved in 5% dextrose, CBL0137 was administered intravenously at a dosage of 40
mg/kg/day. The treatment was given once every four days for a total of four weeks. JQ1 -
dissolved in DMSO and subsequently in a solution containing 0.9% saline with 5% PEG400
and 5% TWEEN-80, JQ1 was administered at a dosage of 25 mg/kg/day intraperitoneally.
The treatment was given five days a week over a period of four weeks.

In the HSJD-DIPG007 model, the drug treatments included: CBL0137 - dissolved in 5%
dextrose, CBL0137 was administered intravenously at a dosage of 20 mg/kg/day. The
treatment was given five days a week for a total of four weeks. JQ1 - Dissolved in DMSO
and subsequently in a solution containing 0.9% saline with 5% PEG400 and 5% TWEEN-80,
JQ1 was administered intraperitoneally at a dosage of 50 mg/kg/day. The treatment was
given five days a week for a duration of four weeks.

For the immunocompetent model the protocol for intracranial injections was implemented as
described above with 5-7 weeks old female C57BL6 mice, using IUE 24B7 cells that harbor
H3.3K27M mutation [134, 135]. The timing of treatment initiation was 18 days post
intracranial surgery. The drug treatments were as follows: CBL0137 - dissolved in 5%
dextrose, CBL0137 was administered intravenously at a dosage of 40 mg/kg/day. The
treatment was given once every four days for a total of three weeks. JQ1 - dissolved in
DMSO and subsequently in a solution containing 0.9% saline with 5% PEG400 and 5%
TWEEN-80, JQ1 was administered at a dosage of 50 mg/kg/day intraperitoneally. The
treatment was given five days a week over three weeks. In combination group, CBL0137
was administered before JQ1. Throughout the study, the survival of the mice was closely
monitored daily for neurological and clinical symptoms as described. Brain samples were
collected at the end of the treatment period.

Throughout the study, the survival of the mice was closely monitored daily. Neurological and
clinical symptoms were assessed to detect signs of tumor development. Mice exhibiting
clinical signs of neurological decline, such as ataxia, circling, head tilting, with or without a
20% weight loss, were humanely euthanized. All animal experiments strictly adhered to the
guidelines set by the Animal Care and Ethics Committee of the University of New South
Wales and the Australian Code of Practice for the Care and Use of Animals for Scientific
Purposes. The experimental protocols used in this study were approved under protocol
number 22/46B.

Hematologic and Biochemical Analysis

Hematologic and biochemical analyses were performed as previously described by Khan et
al [121]. Briefly, healthy female BALB/c mice (12-14 weeks old) were treated four weeks with
either vehicle or the indicated compounds, with three mice per treatment cohort. The study
included six experimental groups: vehicle control (buffer A administered intraperitoneally and
5% dextrose intravenously), CBL0137 (40 mg/kg intravenously), JQ1 (25 mg/kg
intraperitoneally), ZEN3694 (Selleckchem E1517) (50 mg/kg orally), CBL0137 + JQ1
combination, and CBL0137 + ZEN3694 combination. ZEN3694 was dissolved in a vehicle
consisting of 5% DMSO, 40% PEG300, 5% Tween 80, and 50% ddH$_2$O. JQ1 was dissolved
in DMSO to prepare a 10X stock solution, which was subsequently diluted with 9 volumes of
buffer A. CBL0137 was dissolved in 5% dextrose. CBL0137 was administered once every
four days via intravenous injection. JQ1 and ZEN3694 were administered on a 5 days on, 2
days off schedule via intraperitoneal for JQ1 and by oral gavage for ZEN3694. Vehicle
control animals received intravenous 5% dextrose once every four days and intraperitoneal
buffer A on weekdays to match the treatment schedules of the active compounds. Weekly
tail vein blood samples were collected for hematological and biochemical analysis
throughout the treatment period. For hematology, blood was collected in EDTA-containing
tubes (Sarstedt, cat # 20.1339.100) (15 μL) and diluted by a factor of 2 with PBS before
loading to the machine to avoid clot formation, then analyzed using a Mindray BC-5000
hematology analyzer. For biochemistry, blood samples were pooled from each cage and
collected in lithium heparin microtainer tubes (Sarstedt, cat # 20.1345.100) (100 μL), then
analyzed using an Abaxis VetScan Biochemistry Analyser with disposable rotors (Zoetis,
Australia; Comprehensive diagnostic profile cat # 10023220). On the final day of treatment,
animals were euthanised and samples collected. Blood was obtained via cardiac puncture
for hematology, biochemistry, and plasma isolation.

LC-MS/MS determination of drug concentration in blood and tissue samples

Mice were treated with the indicated compounds as described in the previous section. Brain
tissue samples (100 mg) were suspended in 900 µL methanol + 0.1% formic acid and
homogenized for compound extraction. Following homogenization, samples were agitated at
4°C for 30 minutes, then centrifuged at 17,000g. The resulting supernatant was collected for
mass spectrometry analysis. Plasma samples (50-100 µL) were mixed with 9 volumes of ice-
cold acetonitrile for protein precipitation. After incubation on ice for 5 minutes, samples were
centrifuged at 17,000g and the supernatant collected for mass spectrometry analysis. All
mass spectrometry analyses, including calibration curve preparation and sample analysis,
were performed by the Mass Spectrometry Facility at the University of New South Wales
(UNSW BMSF).
979

Immunohistochemistry analysis of brain tissue
Brain samples were collected on the final day of treatment and fixed in a 10% formalin
neutral buffered solution (Sigma-Aldrich). The fixed samples were then embedded in paraffin
wax, and 5 µm sections were cut and mounted on glass slides. To prepare the sections for
histological examination, dehydration was carried out using standard procedures.
Subsequently, the sections were stained with hematoxylin/eosin and Ki67, H3K27ac, MDM4,
and MYC (**Table 1**) using the BOND RX Fully Automated Research Stainer (Leica
biosystems) for histological examination. Following staining, positive cells were identified
and quantified using Andy's Algorithms [136] on the image analysis pipeline on FIJI.
989

Surface marker analysis by flow cytometry
DMG cells were treated with CBL0137 and JQ1 (HSJD-DIPG007 1 µM CBL0137 1 µM JQ1;
SU-DIPGVI 2 µM CBL0137 1 µM JQ1) for 24 hours. Following treatment, cells were
harvested, blocked with FcR blocking reagent for 10 min at 4°C, and stained with HLA-
ABC-VioBlue and HLA-E-PE antibodies (1:50 dilution; Table 1) for an additional 10 min at
4°C. Washed cells were analyzed on a MACSQuant VYB Flow Cytometer and data
processed using FlowJo software. Experiments were performed in at least three biological
replicates.
998

Human monocyte derived macrophages were treated in technical duplicate for the following
groups: M0 (media only), M1 (proinflammatory, LPS 1 ng/mL and IFNγ 20 ng/mL), M2 (anti-
inflammatory, IL-4 20 ng/mL and IL-13 20 ng/mL), CBL1037 1 µM, JQ1 1 µM, and
combination of the two drugs. After 48 hours, media containing treatment in plates were
aspirated and replaced with warm Gey's Balanced Salt Solution. Cells were then stained
with CD86-FITC (1:50) and CD206-APC (1:50) (BD Biosciences) (**Table 1**) as per
manufacturer's protocol. After 1 hour of staining, cells were then washed with Gey's

Balanced Salt Solution and harvested using cell scrapers into FACS tubes for flow
cytometry. Analysis was performed using FlowJo. Experiments were performed in biological
triplicate.

1009

Phagocytosis Assay

pHrodo™ BioParticles™ conjugate (Cat. A10010, Thermo Fisher) was diluted in Gey's
solution to a final volume of 2 mL per vial. The suspension was transferred to a 20 mL glass
Schott bottle and sonicated in an ultrasonic water bath at room temperature for 10 minutes
to ensure uniform particle dispersion. For each well of HMDM cells seeded in a 12-well plate,
100 µL of the bead suspension was added, maintaining the manufacturer-recommended
1:10 ratio. No-cell background controls were similarly prepared. Following bead addition,
cells were incubated at 37°C for 2–3 hours. Fluorescence imaging was performed using a
Floid Cell Imaging Station (Thermo Fisher Scientific) with the RFP filter. Fluorescent signal
per cell was quantified using ImageJ. A cell mask was generated, and total fluorescence
intensity was divided by the number of cells per field of view. The experiment was performed
in duplicate.

1022

Xenium In situ spatial profiling

Formalin-fixed brains were sectioned at 5 µm and mounted onto Xenium slides, with two
brains per slide. Xenium Prime In Situ (10x Genomics) was performed according to the
manufacturer's instructions using the 5000 pre-designed gene panel and cell segmentation
staining. Briefly, priming oligos were hybridized to the target RNA, followed by site-specific
cleavage using RNase treatment to release the RNA strand from the hybridized oligo. The
5000-gene panel was then added to the sections to hybridize with the RNA, followed by
ligation to generate circular DNA probes, which were subsequently amplified via rolling circle
amplification. After a series of ethanol washes, the tissue was blocked and stained with the
cell segmentation stain and DAPI. Slides were then loaded into the Xenium Analyzer for
imaging and decoding. Spatial Transcriptomics Services for this study were provided by the
Garvan Genomics Platform at the Garvan Institute of Medical Research.

1035

Xenium In situ spatial profiling analysis

Xenium output from the four treatment conditions were merged into a single Seurat object
(Seurat v5.2.1). Cells with transcript counts <10 and above the 98[th] percentile were filtered,
and data were normalized with the median transcript count. The Seurat object was down
sampled (25,000 cells per sample) using SketchData and LeverageScore to retain rare cell
populations for subsequent scaling and clustering. From the subsetted data, variable
features (n = 2000 genes) were identified using FindVariableFeatures, data were scaled

using ScaleData and Dimensionality reduction performed using RunPCA. Batch correction was performed using Harmony Integration, followed by UMAP projection and clustering in Harmony space. Clustering was then projected to all cells. Integrated clusters were exported for visualization using Xenium Explorer (v3.0). Marker genes for each cluster were identified using FindAllMarkers. Tumour and immune clusters of interest were subset and re-analyzed with feature selection, scaling, PCA, Harmony integration, and UMAP clustering. Differential expression analysis between vehicle and combination treatment groups in select clusters was performed using FindMarkers. Genes expressed in fewer than 15% of cells in either group were excluded prior to GSEA using ClusterProfiler.

Table 1. List of antibodies

Target	Catalogue number	Application
SSRP1	CST 13421	WB
SSRP1	Biolegend 609710	IP
SPT16	CST 12191	WB, CUT&RUN
SPT16	Biolegend 607002	IP
BRD4	CST 13440	IP, WB
BRD4	EpiCypher 13-2003	CUT&RUN
ß-Actin	CST 8457	WB
IgG rabbit	EpiCypher 13-0042	CUT&RUN
IgG mouse	Santa Cruz 2025	IP
H3K4me3	EpiCypher 13-0041	CUT&RUN
H3K27me3	CST 9733	CUT&RUN, WB
H3K27ac	Abcam ab4729	IHC
H3K27M	CST 74829	CUT&RUN, WB
PARP	CST 9532	WB
Cleaved PARP	CST 5625	WB
Caspase 3	CST 9662	WB
Cleaved Caspase 3	CST 9664	WB
Histone H3	CST 9717	WB
C-MYC	CST 18583	WB

C-MYC	Abcam 32072	IHC
PDGFRα	CST 3164	WB
MDM4	BETHYL-A300-287	WB
MDM4	Abcam 244366	IHC
P53	CST 2527	WB
GADD45B	Thermo PA5-43160	WB
SOX2	CST 3579	WB
TUBB3	CST 5568	WB
Ki67	Thermofisher RM-9106 SP6	IHC
CD206-APC	BD 550889	Flow
CD86-FITC	BD 555657	Flow
HLA-ABC-VioBlue	Miltenyi 130-120-435	Flow
HLA-E- PE-Vio 615	Miltenyi 130-117-403	Flow

1054

1055

1056 **Table 2. Sequences of qPCR primers**

Gene	Forward	Reverse	GenBank
RPS18	GAGGATGAGGTGGAACGTGT	TCTTCAGTCGCTCCAGGTCT	NM_022551.2
B2M	AGATGAGTATGCCTGCCGTG	GCGGCATCTTCAAACCTCCA	NM_004048.2
HIF1α	CTTTCCTTCTCTTCTCCGCGT	TTTCGACGTTCAGAACTTATCTTTT	NM_001530.4
HLA-A	CACAGACTGACCGAGTGGAC	GGATGGTGTGAGAACCGGC	NM_002116.8
IFNA	ACTCATACACCAGGTCACGC	TGGTCATAGTTATAGCAGGGGTG	NM_024013.2
IFNB	GCGACACTGTTCGTGTTGTC	GAGCTGTAGTGGAGAAGCACA	NM_002176.4
IL-1B	CAGGCTGCTCTGGGATTCTC	GTCCTGGAAGGAGCACTTCAT	NM_000576
IRF1	TCTTCCCTCTTCCACTCGGA	TGATGGGCATGTTGGCTCTG	NM_002198.3
IRF3	GACCTTCCATCGTAGGCCG	CTCTTGTTCACCCAGGCCAC	NM_001571.6
IRF8	CAAACTAGGCGTGGCAACTG	AATCGTCCACAGAAGGCTCC	NM_001363907.1
OAS1	CTGGCACAAGAGGCTGAGG	CACTGTTGCTTTCAGCCAGC	NM_016816.4
p53	GTTCCGAGAGCTGAATGAGG	TCTGAGTCAGGCCCTTCTGT	AF098067
MYC	TACAACACCCGAGCAAGGAC	CTAACGTTGAGGGGCATCGT	NM_002467.6
MDM4	CCTGGACAAATCAATCAGGTACG	CCATATGCTGCTCCTGCTGA	NM_002393.5
PDGFRA	TGTCATTTGGAAACAGAAACCGA	AAACGCTCCAGACCCCAAG	NM_006206.6

1057

 Data Availability

 Raw and processed NGS data has been uploaded onto the GEO repository. CUT&RUN
 (GSE272130), ATAC-Seq (GSE272129), nascent RNA-seq (GSE299509), RNA-seq
 (GSE272131), and Xenium (GSE299434).

 Figure legends

 Figure 1. FACT and BRD4 expression and co-localization in DMG
 A) Genomic distribution of SPT16 peaks in HSJD-DIPG007 (DIPG7), SU-DIPGVI (DIPGVI),
 and VUMC-DIPG10 (DIPG10) compared to a background of ~1000 randomly shuffled
 genomic regions. **B)** Over-represented gene ontologies for genes associated with SPT16
 peaks. **C)** Transcription factor motif enrichment analysis and motif logos for SPT16 peaks in
 HSJD-DIPG007 and SU-DIPGVI cells. **D)** Heatmaps depicting SPT16, BRD4, H3K27M,
 H3K4me3, IgG CUT&RUN signal and ATAC-seq signal at transcription start sites (TSS) in
 HSJD-DIPG007 neurospheres. Public H3K27ac ChIP-seq data was downloaded from [41].
 Heatmaps are ranked based on SPT16 signal and separated into two k-means clusters.
 Average signal is shown in the profile plots. Representative data from n=2 experiments. **E)**
 Gene tracks display SPT16, BRD4, H3K27M, H3K4me3, IgG CUT&RUN and H3K27ac[41]
 ChIP-seq signal at TSSs of cancer-related genes *MYC, PDGFRA* and *MDM4* in HSJD-
 DIPG007 and SU-DIPGVI cells. **F)** Profile plots and heatmaps depicting the average SPT16,
 BRD4 and IgG CUT&RUN at SPT16 + BRD4 common peaks (orange), SPT16 unique peaks
 (pink), BRD4 unique peaks (blue) in HSJD-DIPG007 cells. **G)** Co-Immunoprecipitation and
 western blotting in HSJD-DIPG007 and SU-DIPGXIII nuclear extracts showing interactions
 between BRD4 with the FACT subunits and H3K27M. **H)** Gene tracks displaying SPT16 and
 BRD4 CUT&RUN signal at the TSSs of *BRD2, OLIG2, SOX2,* and *BCL2L1.* See related
 Figure S1.

 Figure S1. FACT and BRD4 expression and co-localization in DMG. Related to Figure
 1.
 A-B) Profile plots and heatmaps depicting SPT16, BRD4, H3K27M, H3K4me3, IgG
 CUT&RUN signal in VUMC-DIPG10 (**A**) and SU-DIPGVI (**B**) neurospheres. Heatmaps are
 ranked based on SPT16 signal and separated into two k-means clusters. Representative
 data from n=2 experiments. For SU-DIPGVI, H3K27ac CUT&RUN from [47] and ATAC-seq
 signal is shown. **C)** Profile plots and heatmaps depicting the average SPT16, BRD4 and IgG
 CUT&RUN signal at SPT16 + BRD4 common peaks (orange), SPT16 unique peaks (pink),
 BRD4 unique peaks (blue) in SU-DIPGVI cells. **D**) Co-Immunoprecipitation and western

blotting in SU-DIPGVI nuclear extracts showing direct interactions between BRD4 with the
FACT subunits and H3K27M. **E**) Dot plot of over-represented gene ontologies for genes
associated with SPT16 peaks, BRD4 peaks, and SPT16+BRD4 common peaks in HSJD-
DIPG007 cells. **F**) Expression of FACT subunit genes *SSRP1* and *SUPT16H*, and *BRD4* in
DMG and other CNS tumors compared to normal fetal and adult brain. RNA-seq data for
tumor samples were obtained from the ZCC clinical trial [55] and normal brain samples from
McGill University for NFB [56]. **G**) Expression of FACT subunit genes *SSRP1* and
SUPT16H, and *BRD4* in H3K27M DMG cell lines compared to other pediatric cancer cell
lines and to non-malignant cell lines. DMG samples were compared to all other samples
using independent two-sample t-tests. **H)** Western blot for SSRP1 and BRD4 in a panel of
DMG cell lines compared to Normal Human Astrocytes (NHA). **I**) CRISPR screen density
plot showing gene dependency scores for *BRD4*, *SSRP1*, and *SUPT16H* across glioma
subtypes. Gene effect score distributions are shown for H3G34-mutant Diffuse Hemispheric
Glioma (green), H3K27M DMG (pink), and H3 wild-type high-grade glioma (blue). A
threshold of < –0.5 (dashed line) indicates gene dependency.
1107
Figure 2. In vitro therapeutic efficacy of combinatorial FACT and BRD4 inhibition
A-B) Dose-response curves representing cell viability in a panel of patient-derived DMG cell
lines treated with JQ1 **(A)** and trotabresib **(B)** following 72 hours. **C**) Dose response curves
in isogenic SU-DIPGXIII cells (+/- H3K27M) and bar plots depicting IC50 values for
CBL0137, JQ1 and trotabresib. **D-E**) Cell viability curves in SU-DIPGXIII isogenic cells
treated with a combination of CBL0137 and JQ1 (**D**) or trotabresib (**E**). (**F**) Dose-response
curves for SU-DIPGVI cells treated with CBL0137 combined with JQ1 and trotabresib. All
dose response curves are represented as percentage viability compared to untreated
controls, mean ± SEM (n = 3). **G-H**) Colony formation assays in HSJD-DIPG007 cells treated
with the indicated drugs for 2 weeks. Data is presented as mean ± SEM (n=4).
Representative images shown from the 0.2μM concentration. (**I-J**) Flow cytometry cell cycle
distributions in SU-DIPGVI cells treated with CBL0137 combined with JQ1 (**I**) or trotabresib
(**J**) for 48 hours. (n=3). (**K**) Western blot analysis of apoptosis markers in SU-DIPGVI cells
treated with CBL0137 and JQ1 for 24 hours. Representative blot shown (n=3). P-values
were calculated using a two-tail t-test (**C**) or an ANOVA test with Tukey's multiple
comparisons test for single and combination treatments (**G-J**). * p<0.05, **p<0.01,
***p<0.001, ****p<0.0001. Combination index (CI) values were calculated using Calcusyn,
with synergistic CI values indicated. See related Figure S2.
1126
Figure S2. In vitro therapeutic efficacy of combinatorial FACT and BRD4 inhibition.
Related to Figure 2.

A) Area under the curve (AUC) values for dose-response curves comparing SU-DIPGXIII
isogenic cells -/+ H3K27M. P-values were calculated using a two-tail t-test. (**B-D**)
Combination dose-response curves for Histone mutant HSJD-DIPG007 (**B**), RA055 (**C**), and
P005401 (**D**) cells treated for 72 hours (**E-G**) Combination dose-response curves for Histone
wildtype P016802 (**E**), VUMC-DIPG10 (**F**), and AUS-DIPG17 (**G**) cells treated for 72 hours.
All dose response curves are represented as percentage viability compared to untreated
controls, mean ± SEM (n = 3). Combination index (CI) values were calculated using
Calcusyn, with synergistic CI values indicated. (**H-I**) Colony formation assays in RA055 (**H**)
and SU-DIPGVI (**I**) cells treated with the indicated drugs for 2 or 4 weeks. Data is presented
as mean ± SEM (n=4). (**J-K**) Flow cytometry cell cycle distributions in RA055 (**J**) and HSJD-
DIPG007 (**K**) cells treated with CBL0137 combined with JQ1 or trotabresib for 48 hours
(n=3). Significance was calculated using a one-way ANOVA with Tukey's multiple
comparisons test for single and combination treatments. * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$,
****$p < 0.0001$.

1143

Figure 3. CBL0137 and JQ1 condense chromatin at gene promoters.
A) Profile and box plots displaying ATAC-seq signal at SPT16 peaks (left) and BRD4 peaks
(right) in DMG cells treated with DMSO, CBL0137 (1µM for HSJD-DIPG007 and 2µM for SU-
DIPGVI), JQ1 (1µM) or the combination for 4 hours. A Wilcox test was used to test
significance between treatment groups. **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. Representative
data from n=2 experiments for HSJD-DIPG007 and n=3 experiments for SU-DIPGVI with
remaining replicates shown in Fig S3E-F. **B)** Volcano plot showing differentially accessible
regions. Blue data points represent significantly (FDR < 0.05, LFC < -1) downregulated
"closed" regions, red data points represent significantly upregulated "opened" (FDR < 0.05,
LFC > 1) regions, with the number of regions indicated. **C)** Venn diagram displaying the
overlap between the closed regions common to both cell lines. A hypergeometric test was
used to determine over enrichment. **D)** Genomic distribution of the overlapping closed
regions from panel C compared to background ATAC-seq genomic windows. **E**) Dot plot of
over-represented gene ontologies in the genes associated with the overlap regions from
panel C. **F)** Gene tracks display ATAC-seq signal at promoters and putative enhancers of
MYC, PDGFRA, and MDM4. H3K27ac (from [41] [47]) and H3K4me3 (this study) enrichment
is also shown. Representative replicate from n=2-3 experiments. See also Fig S3.

1161

Figure S3. CBL0137 and JQ1 condense chromatin at gene promoters. Related to
Figure 3.
A-B) Heatmaps depicting ATAC-seq signal at transcription start sites in HSJD-DIPG007
(n=2) (**A**) and SU-DIPGVI (n=3) (**B**) cells treated with DMSO, CBL0137 (1µM for HSJD-

DIPG007 and 2µM for SU-DIPGVI), JQ1 (1µM) or the combination for 4 hours. Average
signal is shown in the profile plots above. **C-D)** Principal component analysis (PCA) plots
showing the clustering of samples based on signal at TSS in HSJD-DIPG007 (**C**) and SU-
DIPGVI (**D**). **E-F)** Box plots displaying ATAC-seq signal at SPT16 peaks (left) and BRD4
peaks (right) in DMG cells treated with DMSO, CBL0137 (1µM for HSJD-DIPG007 and 2µM
for SU-DIPGVI), JQ1 (1µM) or the combination for 4 hours. A Wilcox test was used to test
significance between treatment groups. * $p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$. **G-H**)
Bar charts depicting the number of significantly differentially accessible regions (DARs). Blue
represents significantly (FDR < 0.05, LFC < -1) downregulated "closed" regions, red data
represents significantly upregulated "opened" (LFC > 1). **I**) Western blot for H3K27me3, total
H3, and β-actin in HSJD-DIPG007 cells treated for 24 hours with the indicated treatment
groups. **J)** Densitometry bar graphs (mean +/- SEM; n=3) normalized to β-actin and total H3,
and expressed as a fold-change relative to the DMSO control. An unpaired t-test was used
to calculate significance. * $p<0.05$. **K**) H3K27me3 CUT&RUN volcano plot displaying CSAW
differential binding analysis in HSJD-DIPG007 cells treated with 1µM CBL0137 for 4h.
Significant threshold p-val<0.05 (n=2).

1182

Figure 4. CBL0137 and JQ1 induce a broad suppression of transcription
A) Total RNA (upper) and nascent RNA (lower) MA plots showing significantly differentially
expressed genes in HSJD-DIPG007 cells treated with CBL0137 (1µM), JQ1 (1µM), or the
combination. Blue points represent significantly (FDR < 0.05, LFC < -1) downregulated
genes, red points represent significantly upregulated genes (FDR < 0.05, LFC >1) with the
number of genes indicated. Bars indicate the proportion of DE genes. **B)** Proportional Venn
diagrams illustrating the overlap of downregulated nascently transcribed genes in cells
treated with single agents and combination therapy. **C**) Correlations of log fold-change in
nascently transcribed genes between combination therapy, each monotherapy, and the sum
of both monotherapies in HSJD-DIPG007 cells. **D**) Nascent RNA-seq heatmap displaying
top 30 up and downregulated genes DMSO vs Combination treated HSJD-DIPG007 cells. **E)**
Enrichment map of over-represented gene ontologies among nascently transcribed genes
downregulated by combination treatment in HSJD-DIPG007 cells, with high-level processes
highlighted in labelled boxes. **F**) Volcano plot showing differential expression of nascently
transcribed genes associated with SPT16 and BRD4 peaks in HSJD-DIPG007 cells. Key
genes are labelled. See also Fig S4.

1199

1200 **Figure S4. CBL0137 and JQ1 induce a broad suppression of transcription. Related to**
1201 **Fig 4.**

A) Total RNA (upper) and nascent RNA (lower) MA plots showing significantly differentially expressed genes in SU-DIPGVI cells treated with CBL0137 (2μM), JQ1 (1μM), or the combination for 4 h. Blue points represent significantly (FDR < 0.05, LFC < -1) downregulated genes, red points represent significantly upregulated genes (FDR < 0.05, LFC >1) with the number of genes indicated. Bars indicate the proportion of DE genes. **B)** Correlations of log fold-change in nascently transcribed genes between combination therapy, each monotherapy, and the sum of both monotherapies in SU-DIPGVI cells. **C)** Nascent RNA-seq heatmap displaying top 30 up and downregulated genes DMSO vs Combination treated SU-DIPGVI cells. **D**) GADD45B and TUBB3 western blot from treated SU-DIPGVI cells after 24 h. representative blot from 3 independent experiments. **E**) Enrichment map of over-represented gene ontologies among nascently transcribed genes downregulated by combination treatment in SU-DIPGVI cells, with high-level processes highlighted in labelled boxes. **F**) Over-represented ontologies of significantly downregulated proteins identified by mass spectrometry in HSJD-DIPG007 (left) and SU-DIPGVI (right) cells treated with CBL0137 at IC50 doses. **G**) Volcano plot showing differential expression of nascently transcribed genes associated with SPT16 and BRD4 peaks in SU-DIPGVI cells.

Figure 5. Chromatin compaction by CBL0137 and JQ1 decreases expression of oncogenes

A) Nascent RNA-seq heatmap displaying *MYC, MDM4, PDGFRA,* and *SOX2* genes in the indicated treatment groups for HSJD-DIPG007 and SU-DIPGVI cells. **B**) RNA-seq TPM values (mean +/- SEM; n=3) for *MYC, PDGFRA*, *MDM4* and *SOX2* in HSJD-DIPG007 cells treated for 4 h with CBL0137 and JQ1. **C**) Representative MYC, PDGFRα, MDM4 and SOX2 western blots and bar charts displaying densitometry (mean +/- SEM; n=3) normalized to β-Actin or α-Tubulin and expressed as a fold-change relative to the DMSO control. 24 h treatment. PDGFRα and MDM4 were run on the same gel, and use the same loading control. **D**) Proliferation of SF8628 cells measured by time-lapse microscopy every 6 h for 6 days. Cells were transfected with non-targeting siRNA or siRNA targeting MYC, MDM4 and PDGFRA (tripleKD) and treated with DMSO or the combination of CBL0137 (0.2μM) and JQ1 (0.5μM). **E**) Cytotoxicity assay comparing CBL0137+JQ1, sulanemadlin, and the combination of all 3 drugs in HSJD-DIPG007 cells after 96 h and Bliss synergy score matrix comparing CBL0137+JQ1 and sulanemadlin. Significance was calculated using a one-way ANOVA with Sidak's multiple comparisons test for single and combination treatments. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure S5. Chromatin compaction by CBL0137 and JQ1 decreases expression of oncogenes. Related to Figure 5.

A) Gene tracks displaying H3K4me3, H3K27ac [41, 47], IgG, SPT16, BRD4 CUT&RUN, and

ATAC-seq signal across the 4 indicated treatments in HSJD-DIPG007 and SU-DIPGVI cells

at the *MYC, PDGFRA, MDM4, SOX2* and *SUPT16H* loci. **B**) RNA-seq TPM values (mean +/-

SEM; n=3) for *MYC, PDGFRA, MDM4, SOX2,* and *SUPT16H* in SU-DIPGVI cells treated for

4 h with CBL0137 and JQ1 **C**) Representative MYC, PDGFRα, MDM4, SOX2 and SPT16

western blots and bar charts displaying densitometry (mean +/- SEM; n=3) normalized to β-

Actin and expressed as a fold-change relative to the DMSO control. 24 h treatment.

PDGFRα and MDM4 were run on the same gel, and use the same loading control. SOX2

was run on same gel as Fig S4D and use the same loading control. **D**) RNA-seq TPM values

(mean +/- SEM; n=3) for TP53 in the indicated cell lines. 4h treatment. **E**) Representative

p53 western blots and bar charts displaying densitometry (mean +/- SEM; n=3) normalized

to β-Actin and expressed as a fold-change relative to the DMSO control. 24 h treatment. **F**)

qPCR bar plots illustrating gene knockdown of MYC, PDGFRA, and MDM4 in SF8628 cells.

Cells were transfected with siRNA targeting each gene individually or in combination (triple

knockdown) for 24 hours. **G**) qPCR bar plots illustrating relative expression of MYC,

PDGFRA and MDM4 in SF8628 cells treated with DMSO or the combination of CBL0137

(0.6 µM) and JQ1 (1 µM) for 24 h. **H**) cytotoxicity assay in SF8628 DMG cells treated with

CBL0137 and JQ1. Combination index values were calculated with Calcusyn. **I**)

Sulanemadlin toxicity assay in HSJD-DIPG007 cells with and without TP53, SU-DIPGVI (p53

mutant), and SU-DIPGXVII (p53 mutant) cells. **J**) Cytotoxicity assay comparing

CBL0137+JQ1, sulanemadlin, and the combination of all 3 drugs in SU-DIPGVI cells after 96

h and Bliss synergy score matrix comparing CBL0137+JQ1 and sulanemadlin. **(B-F)**

Significance was calculated using a one-way ANOVA with Sidak's multiple comparisons test

for single and combination treatments. For **G)** Statistical significance was determined using

an unpaired student's t-test. * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

1264

Figure S6. Combination treatment leads to alternative splicing in DMG cells

A) GSEA enrichment plot showing negative enrichment (downregulation) of the spliceosomal

snRNP assembly pathway in SU-DIPGVI cells (DMSO vs Combination). **B-C**) Bar graph

displaying the total number of splicing events (upper) or uniquely spliced genes (lower) in

HSJD-DIPG007 and SU-DIPGVI cells with the indicated treatments. Significance was

calculated using a one-way ANOVA with Dunnet's multiple comparisons test comparing

single and combination treatments. **p<0.01, ***p<0.001, ****p<0.0001. **C**) Pie charts

depicting the breakdown of differential splicing events common to 3 replicates for the

combination vs DMSO condition, in each model. **D**) Gene tracks display ATAC-seq signal

(upper tracks) and RNA-seq coverage (lower tracks) in DMSO control and combination

treated HSJD-DIPG007 and SU-DIPGVI cells. Representative tracks for n=2-3 experiments

for each model.

1277

Figure 6. Therapeutic efficacy of CBL0137 and JQ1 combination treatment in

orthotopic models of DMG.

A) Treatment scheme and survival curve of SU-DIPGVI-Luciferase with CBL0137/JQ1

treatment. Median survival of cohorts (days): Vehicle (n=9) = 73, CBL0137 (n=9) = 76, JQ1

(n=6) = 78, CBL0137 + JQ1 (n=6) = 95.5. Exact p-values listed in Supplementary Table 6.

B) Xenogen luminescence imaging of SU-DIPGVI-Luciferase tumor burden and averaged

luminescence signal measured over time in each cohort. **C-D**) IHC staining and

quantification for Ki67 (**C**) and H3K27ac (**D**) of SU-DIPGVI-LUC tumors after 4 weeks of

treatment (n=3). **E**) treatment schema and survival curve for the HSJD-DIPG007 model with

CBL0137/JQ1 treatment. Median survival of cohorts (days): Vehicle (n=12) = 70.5, CBL0137

(n=12) = 82, JQ1 (n=12) = 86, CBL0137 + JQ1 (n=12) = 93.5. Exact p-values listed in

Supplementary Table 7. **F**) Treatment scheme and survival curve of RA055 model with

CBL0137/JQ1 treatment. Median survival of cohorts (days): Vehicle (n=12) = 46, CBL0137

(n=9) = 51, JQ1 (n=8) = 55, CBL0137 + JQ1 (n=9) = 66. Exact p-values listed in

Supplementary Table 8. **F-J**) Immunohistochemistry staining of RA055 tumors for Ki67 (**G**),

H3K27ac (**H**), MDM4 (**I**), and MYC (**J**). Three images were taken from brain samples

collected from three mice in each cohort at the end of the treatment period. Scale: 50µm. For

survival curves, statistical analysis was performed using the Log-Rank (Mantel-Cox) with

multiple test corrections applied. For IHC analysis data is presented as mean values ± SEM.

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. p-values were calculated using one-way

ANOVA with Tukey's multiple comparisons test for treated and untreated cohorts. See also

Fig S7.

1300

Figure S7. Therapeutic efficacy of CBL0137 and JQ1 combination treatment in

orthotopic models of DMG related to Figure 6.

A-B) Weights of mice before, during and after treatment period for the SU-DIPGVI-Luc (**A**)

and HSJD-DIPG007 model (**B**). Data for individual mice is shown in (**A**) and averaged mice

are shown in (**B**). (**C**) Hematologic measurements from non-tumor-bearing mice. A two-way

ANOVA with Dunnett's multiple comparisons test was used to assess significance between

vehicle and treatment groups (n = 3 per cohort). *p < 0.05, **p < 0.01, ***p < 0.001, ****p <

0.0001.(**D**) Pharmacokinetic analysis of JQ1 in brain and plasma following 4 weeks. (**E**) IHC

staining for Ki67 on brain samples from endpoint surviving mice in RA055 model

demonstrating positive engraftment. (**F**) Treatment scheme for CBL0137 and ZEN-3694

administration, and weights of mice before, during and after treatment period. (**G**)

Hematologic measurements from non-tumor-bearing mice. A two-way ANOVA with
Dunnett's multiple comparisons test was used to assess significance between vehicle and
treatment groups (n = 3 per cohort). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. **(H)**
Pharmacokinetic analysis of ZEN-3694 in brain and plasma following 4 weeks **(I-K)** Whole
brain scans of H&E stained brains from the SU-DIPGVI-LUC model (**I**), RA055 model **(J)** and
HSJD-DIPG007 model (**K**). Representative images shown. Scale = 2000µM.

1318

Figure 7. CBL0137 and JQ1 influence immune-associated pathways in DMG and
tumor-infiltrating immune cells
A) Nascent RNA-seq heatmap displaying interferon and antigen presentation genes in the
indicated treatment groups for SU-DIPGVI cells. **B**) GSEA enrichment plot showing positive
enrichment (upregulation) of the antigen presentation pathway in SU-DIPGVI cells (DMSO
vs Combination). **C)** Flow cytometry histograms for HLA-ABC and HLA-E in treated SU-
DIPGVI cells (24h). Representative data from at least 3 independent experiments. Median
Fluorescence Intensity (MFI) values are shown. **D**) Flow cytometry histograms for CD86 and
CD206 in human monocyte-derived macrophages (HMDM) cells in M1 or M2 polarization
conditions or treated with CBL0137 and JQ1 for 48h. Representative data from 3
independent experiments (3 donors) shown. MFI values were quantified as above. P-values
for MFI comparisons **(C,D)** were calculated using a one-way ANOVA with Sidak's multiple
comparison test. data is presented as mean values ± SEM. *p<0.05, **p<0.01. **E**)
Normalized Xenium single cell UMAP plot of immunocompetent murine glioma model treated
with CBL0137 and JQ1 for 3 weeks. UMAP is split into the 4 treatments cohorts. Tumor-
associated clusters are indicated in the dashed circles. **F**) Xenium Explorer images of
immunocompetent model. Upper panels: whole brain sagittal slices with tumor cells
(Astrocyte, oligodendrocyte lineage and proliferative subtypes) and immune cells (tumor
associated macrophage; TAM, and T cells) overlaid onto the immunofluorescence image.
Scale = 5000µM. Middle panels: Higher magnification insets of tumor region showing the
distribution of tumor and immune clusters that are highlighted in panel E. Scale = 1000 µM.
Lower panel: higher magnification insets displaying immune clusters only. Scale = 200µM.
G) Composition of each tumor-associated cluster relative to the total of all tumor-associated
clusters. **H-I**) GSEA enrichment plots comparing combination to vehicle (red = upregulated,
blue = downregulated) in TAMs (**H**) and T cells (**I**). **J-K**) Violin plots for select genes
significantly upregulated in the vehicle vs combination treated TAMs (*Ighm* and *C3*) (**J**) and
T cells (*Ptprc* and *Dock2*) (**K**). **L**) Xenium Explorer images of tumor, immune and endothelial
cells with individual transcripts represented by dots. Scale = 100µM. See also Fig S8.

1347

Figure S8. CBL0137 and JQ1 influence immune-associated pathways in DMG and tumor-infiltrating immune cells, related to Figure 7.

A) Nascent RNA-seq heatmap displaying interferon and antigen presentation genes in the indicated treatment groups for HSJD-DIPG007 cells. **B)** qPCR analysis of interferon and inflammatory genes in HSJD-DIPG007 cells treated for 48h with CBL0137 (0.5µM) or JQ1 (1µM). (n=3). **C)** Flow cytometry histograms for HLA-ABC and HLA-E in treated SU-DIPG007 cells (24h). Representative data from at least 3 independent experiments. Median Fluorescence Intensity (MFI) values are shown. **D)** qPCR analysis of human monocyte-derived macrophage (HMDM) expression M1 proinflammatory genes following treatment with 1µM CBL0137 or 1µM JQ1, alone or in combination for 48 h, compared to untreated macrophages (M0) and macrophages exposed to M1 and M2 priming conditions. N=3. **E)** Fluorescent bead-uptake phagocytosis assay of HMDMs in M1 or M2 priming conditions or treated with CBL0137 and JQ1 for 48 hours quantification of fluorescent signal/cell calculated from 2 wells. P-values were calculated using a one-way ANOVA test with Dunnett's multiple comparison test. Data is presented as mean values ± SEM. $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$. **F)** Xenium single cell UMAP projection of 4 integrated samples. Major clusters are annotated based on lineage markers. **G)** Vehicle treated brain slice stained with DAPI (white) with tumor (clusters 3, 9, 11) and immune (clusters 6, 24) associated clusters overlaid. **H)** Dot plot displaying marker gene expression across the 35 UMAP clusters. **I)** Volcano plots displaying significantly differentially expressed genes in each of the 3 tumor clusters compared to the other 2 tumor clusters. **J)** GSEA enrichment plots comparing combination to vehicle (red = upregulated, blue = downregulated) in tumor cells (clusters 3+9+11). **K)** Violin plots for select genes (*Myc, Sox2,* and *Ifi27)* significantly differentially expressed genes in the vehicle vs combination treated tumor cells.

Supplementary Table 1: Gene lists from CUT&RUN, ATAC-seq, RNA-seq, proteomics, and spatial in situ experiments. (All lists are ranked by FDR significance.)

A) Chromosome coordinates for SPT16 and BRD4 MACS2 peaks from merged CUT&RUN replicates in HSJD-DIPG007, SU-DIPGVI, and VUMC-DIPG10 DMG cells. **B)** Genes associated with SPT16 and BRD4 peaks in HSJD-DIPG007, SU-DIPGVI, and VUMC-DIPG10 DMG cells. **C)** Genes linked to differentially accessible regions (DARs) identified by ATAC-seq CSAW analysis (FDR < 0.05, LFC > 1 = opened; LFC < −1 = closed) in HSJD-DIPG007 and SU-DIPGVI cells treated with CBL0137 (1 µM for HSJD-DIPG007, 2 µM for SU-DIPGVI) or JQ1 (1 µM) for 4 h. **D)** Total and nascent RNA-seq differentially expressed genes (FDR < 0.05, |LFC| > 1) in HSJD-DIPG007 and SU-DIPGVI cells treated with CBL0137 or JQ1 for 2 h. **E)** RNA-seq differentially expressed genes (FDR < 0.05, |LFC| > 1) in HSJD-DIPG007 and SU-DIPGVI cells treated with CBL0137 or JQ1 for 4 h. **F)** Genes with

differential splicing events common across three independent replicates in HSJD-DIPG007 and SU-DIPGVI cells treated with CBL0137 and JQ1 (4 h). **G)** Differentially expressed proteins identified by proteomics in HSJD-DIPG007 and SU-DIPGVI cells treated with CBL0137 (0.4 µM and 0.9 µM, respectively) for 24 h. **H)** Xenium spatial transcriptomics marker genes differentially expressed (FDR < 0.05, LFC > 1) in each cluster compared to all clusters. **I)** Xenium marker genes differentially expressed between tumor clusters (FDR < 0.05, LFC > 1). **J)** Xenium differentially expressed genes comparing combination therapy versus vehicle in tumor, TAM, and T cell clusters (FDR < 0.05, |LFC| > 0.5).

Acknowledgments

We would like to express our gratitude to Michelle Monje, Angel Montero Carcaboso, and Esther Hulleman for their generous provision of the SU-DIPG, HSJD-DIPG, and VUMC-DIPG10 cells, respectively, which were instrumental in the success of this study. We thank Nada Jabado and Brian Krug for providing the isogenic H3K27M KO SU-DIPGXIII models and Yunjia Zhang, Oluwaseyi Samson Ogunmodede, and Kerry-Anne Rye for providing the HMDM cells. We thank Katerina Gurova and Andrei Gudkov for providing the CBL0137 drug. We also acknowledge the Core Services Group at Children's Cancer Institute for their valuable technical support throughout the research. We extend our appreciation to the Flow Cytometry Core Facility, BRIL Facility, and BMSF Facility (Russel Pickford) at UNSW assistance in flow cytometry, imaging, and pharmacokinetic testing, respectively, which greatly contributed to our data analysis and interpretation. We also acknowledge the Garvan Institute Histopathology Facility and Garvan Genomics Platform for their services. Furthermore, we would like to acknowledge the Zero Childhood Cancer Program and the Tumour Bank teams at CCI for their significant contributions to the establishment of primary DMG cultures, which were indispensable for our investigations. The authors thank the Children's Cancer Institute Animal Facility for providing support to this study. We would like to thank Nathan Dahl and Lays Sobral for technical advice regarding CUT&RUN and ATAC-seq, and John Reeves and Quy Xiao Xuan Lin for technical support for spatial analyses. We would like to also acknowledge Vyoma Patel for proofreading this manuscript. This work has generously been funded by the Kids Cancer Alliance, We Love You Connie Foundation Grant, Levi's Project, CINSW Fellowship (HH) (2022/ECF1425), The Kids Cancer Project Col Reynolds Fellowship (AK), NHMRC Synergy Grant (2019056), Cancer Institute NSW Translational Program Grant (2019/TPG2037), NHMRC Investigator grant (GNT2017898), and DIPG Collaborative: The Cure Starts Now (2018). Graphical abstract and treatment schemes created with BioRender.com.

1420

Author Contributions

Conceptualization: MT, DSZ. **Methodology:** HH, AK, JL. **Formal Analysis:** HH, AK, AE, NJ, SER, XG, HN, CM, BR, DK, CS. **Investigation:** HH, AK, AE, AG, HN, AL, BR, MT, EW, DL, RL, KI, WL, RS, LRS. **Data Curation:** HH, AK, CM, NJ, SER, CXS. **Visualization:** HH, AK, SER, YCS, BR. **Writing – Original Draft:** HH, AK. **Writing – Review & Editing:** All. **Supervision:** RF, RJW, FVM, TNP, CM, BR, MT, MD, DSZ. **Project Administration:** HH, AK. **Funding Acquisition:** HH, BR, MT, DSZ.

Declaration of Interests

DSZ declares consulting/advisory board fees from Bayer, Astra Zeneca, Accendatech, Novartis, Day One, FivePhusion, Amgen, Alexion, and Norgine and research support from Accendatech. The remaining authors have declared that no competing interests exist.

References

1. Louis, D.N., et al., *The 2021 WHO Classification of Tumors of the Central Nervous System: a summary.* Neuro-Oncology, 2021. **23**(8): p. 1231-1251.
2. Schwartzentruber, J., et al., *Driver mutations in histone H3.3 and chromatin remodelling genes in paediatric glioblastoma.* Nature, 2012. **482**(7384): p. 226-231.
3. Khuong-Quang, D.A., et al., *K27M mutation in histone H3.3 defines clinically and biologically distinct subgroups of pediatric diffuse intrinsic pontine gliomas.* Acta Neuropathol, 2012. **124**(3): p. 439-47.
4. Wu, G., et al., *Somatic histone H3 alterations in pediatric diffuse intrinsic pontine gliomas and non-brainstem glioblastomas.* Nat Genet, 2012. **44**(3): p. 251-3.
5. Castel, D., et al., *Histone H3 wild-type DIPG/DMG overexpressing EZHIP extend the spectrum diffuse midline gliomas with PRC2 inhibition beyond H3-K27M mutation.* Acta Neuropathologica, 2020. **139**(6): p. 1109-1113.
6. Pratt, D., et al., *Diffuse intrinsic pontine glioma-like tumor with EZHIP expression and molecular features of PFA ependymoma.* Acta Neuropathologica Communications, 2020. **8**(1): p. 37.
7. Ajuyah, P., et al., *Histone H3-wild type diffuse midline gliomas with H3K27me3 loss are a distinct entity with exclusive EGFR or ACVR1 mutation and differential methylation of homeobox genes.* Sci Rep, 2023. **13**(1): p. 3775.
8. Lewis, P.W., et al., *Inhibition of PRC2 activity by a gain-of-function H3 mutation found in pediatric glioblastoma.* Science, 2013. **340**(6134): p. 857-61.
9. Piunti, A., et al., *CATACOMB: An endogenous inducible gene that antagonizes H3K27 methylation activity of Polycomb repressive complex 2 via an H3K27M-like mechanism.* Science Advances, 2019. **5**(7): p. eaax2887.
10. Bender, S., et al., *Reduced H3K27me3 and DNA hypomethylation are major drivers of gene expression in K27M mutant pediatric high-grade gliomas.* Cancer Cell, 2013. **24**(5): p. 660-72.

11. Harutyunyan, A.S., et al., *H3K27M induces defective chromatin spread of PRC2-mediated repressive H3K27me2/me3 and is essential for glioma tumorigenesis.* Nature Communications, 2019. **10**(1): p. 1262.
12. Krug, B., et al., *Pervasive H3K27 Acetylation Leads to ERV Expression and a Therapeutic Vulnerability in H3K27M Gliomas.* Cancer cell, 2019. **35**(5): p. 782-797.e8.
13. Nagaraja, S., et al., *Histone Variant and Cell Context Determine H3K27M Reprogramming of the Enhancer Landscape and Oncogenic State.* Mol Cell, 2019. **76**(6): p. 965-980.e12.
14. Brien, G.L., et al., *Simultaneous disruption of PRC2 and enhancer function underlies histone H3.3-K27M oncogenic activity in human hindbrain neural stem cells.* Nature Genetics, 2021.
15. Hayden, E., et al., *Therapeutic Targets in Diffuse Midline Gliomas-An Emerging Landscape.* Cancers (Basel), 2021. **13**(24).
16. Groves, A. and T.M. Cooney, *Epigenetic programming of pediatric high-grade glioma: Pushing beyond proof of concept to clinical benefit.* Front Cell Dev Biol, 2022. **10**: p. 1089898.
17. Orphanides, G., et al., *FACT, a factor that facilitates transcript elongation through nucleosomes.* Cell, 1998. **92**(1): p. 105-16.
18. Orphanides, G., et al., *The chromatin-specific transcription elongation factor FACT comprises human SPT16 and SSRP1 proteins.* Nature, 1999. **400**(6741): p. 284-8.
19. Ehteda, A., et al., *Dual targeting of the epigenome via FACT complex and histone deacetylase is a potent treatment strategy for DIPG.* Cell Rep, 2021. **35**(2): p. 108994.
20. Garcia, H., et al., *Facilitates Chromatin Transcription Complex Is an "Accelerator" of Tumor Transformation and Potential Marker and Target of Aggressive Cancers.* Cell Reports, 2013. **4**(1): p. 159-173.
21. Gurova, K., et al., *Structure and function of the histone chaperone FACT - Resolving FACTual issues.* Biochimica et biophysica acta. Gene regulatory mechanisms, 2018: p. S1874-9399(18)30159-7.
22. Safina, A., et al., *FACT is a sensor of DNA torsional stress in eukaryotic cells.* Nucleic Acids Res, 2017. **45**(4): p. 1925-1945.
23. Chang, H.-W., et al., *Mechanism of FACT removal from transcribed genes by anticancer drugs curaxins.* Science advances, 2018. **4**(11): p. eaav2131-eaav2131.
24. Neefjes, J., et al., *Chromatin as an old and new anticancer target.* Trends Cancer, 2024.
25. Gasparian, A.V., et al., *Curaxins: anticancer compounds that simultaneously suppress NF-κB and activate p53 by targeting FACT.* Science translational medicine, 2011. **3**(95): p. 95ra74-95ra74.
26. Carter, D.R., et al., *Therapeutic targeting of the MYC signal by inhibition of histone chaperone FACT in neuroblastoma.* Sci Transl Med, 2015. **7**(312): p. 312ra176.
27. Xiao, L., et al., *Dual targeting of chromatin stability by the curaxin CBL0137 and histone deacetylase inhibitor panobinostat shows significant preclinical efficacy in neuroblastoma.* Clin Cancer Res, 2021.
28. Tallman, M.M., et al., *The small molecule drug CBL0137 increases the level of DNA damage and the efficacy of radiotherapy for glioblastoma.* Cancer Letters, 2021. **499**: p. 232-242.
29. Wang, J., et al., *Effective inhibition of MYC-amplified group 3 medulloblastoma by FACT-targeted curaxin drug CBL0137.* Cell Death Dis, 2020. **11**(12): p. 1029.
30. Monje, M., et al., *Phase I trial of panobinostat in children with diffuse intrinsic pontine glioma: A report from the Pediatric Brain Tumor Consortium (PBTC-047).* Neuro Oncol, 2023. **25**(12): p. 2262-2272.
31. Bradner, J.E., D. Hnisz, and R.A. Young, *Transcriptional Addiction in Cancer.* Cell, 2017. **168**(4): p. 629-643.

32.   Shi, J. and C.R. Vakoc, *The mechanisms behind the therapeutic activity of BET*
bromodomain inhibition. Molecular cell, 2014. **54**(5): p. 728-736.
33.   Lovén, J., et al., *Selective inhibition of tumor oncogenes by disruption of super-*
enhancers. Cell, 2013. **153**(2): p. 320-34.
34.   Taylor, I.C., et al., *Disrupting NOTCH Slows Diffuse Intrinsic Pontine Glioma Growth,*
Enhances Radiation Sensitivity, and Shows Combinatorial Efficacy With
Bromodomain Inhibition. J Neuropathol Exp Neurol, 2015. **74**(8): p. 778-90.
35.   Nagaraja, S., et al., *Transcriptional Dependencies in Diffuse Intrinsic Pontine Glioma.*
Cancer cell, 2017. **31**(5): p. 635-652.e6.
36.   Piunti, A., et al., *Therapeutic targeting of polycomb and BET bromodomain proteins*
in diffuse intrinsic pontine gliomas. Nature Medicine, 2017. **23**(4): p. 493-500.
37.   Zhang, Y., et al., *Combination of EZH2 inhibitor and BET inhibitor for treatment of*
diffuse intrinsic pontine glioma. Cell Biosci, 2017. **7**: p. 56.
38.   Wiese, M., et al., *Combined treatment with CBP and BET inhibitors reverses*
inadvertent activation of detrimental super enhancer programs in DIPG cells. Cell
Death & Disease, 2020. **11**(8): p. 673.
39.   Watanabe, J., et al., *BET bromodomain inhibition potentiates radiosensitivity in*
models of H3K27-altered diffuse midline glioma. J Clin Invest, 2024.
40.   Groves, A., J. Clymer, and M.G. Filbin, *Bromodomain and Extra-Terminal Protein*
Inhibitors: Biologic Insights and Therapeutic Potential in Pediatric Brain Tumors.
Pharmaceuticals, 2022. **15**(6): p. 665.
41.   Wang, J., et al., *Epigenomic landscape and 3D genome structure in pediatric high-*
grade glioma. Science Advances, 2021. **7**(23): p. eabg4126.
42.   Yu, Y., et al., *Olig2 targets chromatin remodelers to enhancers to initiate*
oligodendrocyte differentiation. Cell, 2013. **152**(1-2): p. 248-61.
43.   Talley, M.J., et al., *Distinct requirements for Tcf3 and Tcf12 during oligodendrocyte*
development in the mouse telencephalon. Neural Dev, 2023. **18**(1): p. 5.
44.   Han, S.P., et al., *SNAI1 is involved in the proliferation and migration of glioblastoma*
cells. Cell Mol Neurobiol, 2011. **31**(3): p. 489-96.
45.   Meel, M.H., et al., *Signaling pathways and mesenchymal transition in pediatric high-*
grade glioma. Cell Mol Life Sci, 2018. **75**(5): p. 871-887.
46.   Bočkaj, I., et al., *The H3.3K27M oncohistone affects replication stress outcome and*
provokes genomic instability in pediatric glioma. PLoS genetics, 2021. **17**(11): p.
e1009868-e1009868.
47.   Furth, N., et al., *H3-K27M-mutant nucleosomes interact with MLL1 to shape the*
glioma epigenetic landscape. Cell Rep, 2022. **39**(7): p. 110836.
48.   Lambert, J.P., et al., *Interactome Rewiring Following Pharmacological Targeting of*
BET Bromodomains. Mol Cell, 2019. **73**(3): p. 621-638.e17.
49.   Shu, S., et al., *Synthetic Lethal and Resistance Interactions with BET Bromodomain*
Inhibitors in Triple-Negative Breast Cancer. Mol Cell, 2020. **78**(6): p. 1096-1113.e8.
50.   Zhou, D., et al., *FACT subunit SUPT16H associates with BRD4 and contributes to*
silencing of interferon signaling. Nucleic Acids Res, 2022. **50**(15): p. 8700-8718.
51.   Hsieh, F.-K., et al., *FACT in Cell Differentiation and Carcinogenesis.* Oncotarget,
2011. **2**(11): p. 830-832.
52.   Sui, S., et al., *Ferritinophagy is required for the induction of ferroptosis by the*
bromodomain protein BRD4 inhibitor (+)-JQ1 in cancer cells. Cell Death & Disease,
2019. **10**(5): p. 331.
53.   Lu, L., et al., *Inhibition of BRD4 suppresses the malignancy of breast cancer cells via*
regulation of Snail. Cell Death & Differentiation, 2020. **27**(1): p. 255-268.
54.   Boyson, S.P., et al., *Functional Roles of Bromodomain Proteins in Cancer.* Cancers
(Basel), 2021. **13**(14).
55.   Wong, M., et al., *Whole genome, transcriptome and methylome profiling enhances*
actionable target discovery in high-risk pediatric cancer. Nature Medicine, 2020.
26(11): p. 1742-1753.

56. Jessa, S., et al., *Stalled developmental programs at the root of pediatric brain
tumors.* Nature genetics, 2019. **51**(12): p. 1702-1713.
57. Sun, C.X., et al., *Generation and multi-dimensional profiling of a childhood cancer
cell line atlas defines new therapeutic opportunities.* Cancer Cell, 2023. **41**(4): p. 660-
677.e7.
58. Delmore, Jake E., et al., *BET Bromodomain Inhibition as a Therapeutic Strategy to
Target c-Myc.* Cell, 2011. **146**(6): p. 904-917.
59. Nesher, E., et al., *Role of Chromatin Damage and Chromatin Trapping of FACT in
Mediating the Anticancer Cytotoxicity of DNA-Binding Small-Molecule Drugs.* Cancer
Res, 2018. **78**(6): p. 1431-1443.
60. Kantidze, O.L., et al., *The anti-cancer drugs curaxins target spatial genome
organization.* Nat Commun, 2019. **10**(1): p. 1441.
61. Schofield, J.A., et al., *TimeLapse-seq: adding a temporal dimension to RNA
sequencing through nucleoside recoding.* Nat Methods, 2018. **15**(3): p. 221-225.
62. Takahashi, K. and S. Yamanaka, *Induction of pluripotent stem cells from mouse
embryonic and adult fibroblast cultures by defined factors.* Cell, 2006. **126**(4): p. 663-
76.
63. Marine, J.C. and A.G. Jochemsen, *MDMX (MDM4), a Promising Target for p53
Reactivation Therapy and Beyond.* Cold Spring Harb Perspect Med, 2016. **6**(7).
64. Wohlberedt, K., et al., *Mdm4 supports DNA replication in a p53-independent fashion.*
Oncogene, 2020. **39**(25): p. 4828-4843.
65. Xu, C., et al., *TP53 wild-type/PPM1D mutant diffuse intrinsic pontine gliomas are
sensitive to a MDM2 antagonist.* Acta Neuropathol Commun, 2021. **9**(1): p. 178.
66. Alló, M., et al., *Chromatin and alternative splicing.* Cold Spring Harb Symp Quant
Biol, 2010. **75**: p. 103-11.
67. Filippakopoulos, P., et al., *Selective inhibition of BET bromodomains.* Nature, 2010.
468(7327): p. 1067-1073.
68. Lock, R., et al., *Initial testing (stage 1) of the curaxin CBL0137 by the pediatric
preclinical testing program.* Pediatr Blood Cancer, 2017. **64**(4).
69. Sartor, G.C., et al., *Epigenetic Readers of Lysine Acetylation Regulate Cocaine-
Induced Plasticity.* J Neurosci, 2015. **35**(45): p. 15062-72.
70. Aggarwal, R.R., et al., *A Phase Ib/IIa Study of the Pan-BET Inhibitor ZEN-3694 in
Combination with Enzalutamide in Patients with Metastatic Castration-resistant
Prostate Cancer.* Clin Cancer Res, 2020. **26**(20): p. 5338-5347.
71. Kharenko, O.A., et al., *Combination of ZEN-3694 with CDK4/6 inhibitors reverses
acquired resistance to CDK4/6 inhibitors in ER-positive breast cancer.* Cancer Gene
Ther, 2022. **29**(6): p. 859-869.
72. Zhou, F., *Molecular mechanisms of IFN-gamma to up-regulate MHC class I antigen
processing and presentation.* Int Rev Immunol, 2009. **28**(3-4): p. 239-60.
73. Chhipa, A.S., et al., *The curious case of type I interferon signaling in cancer.* Biochim
Biophys Acta Rev Cancer, 2024. **1879**(6): p. 189204.
74. Chen, M., et al., *Stimulation of an anti-tumor immune response with "chromatin-
damaging" therapy.* Cancer Immunol Immunother, 2021. **70**(7): p. 2073-2086.
75. Zhang, M., et al., *BET inhibition triggers antitumor immunity by enhancing MHC class
I expression in head and neck squamous cell carcinoma.* Mol Ther, 2022. **30**(11): p.
3394-3413.
76. Magner, W.J., et al., *Activation of MHC class I, II, and CD40 gene expression by
histone deacetylase inhibitors.* J Immunol, 2000. **165**(12): p. 7017-24.
77. Luo, N., et al., *DNA methyltransferase inhibition upregulates MHC-I to potentiate
cytotoxic T lymphocyte responses in breast cancer.* Nat Commun, 2018. **9**(1): p. 248.
78. Yang, X., et al., *Epigenetic regulation of macrophage polarization by DNA
methyltransferase 3b.* Mol Endocrinol, 2014. **28**(4): p. 565-74.
79. Wang, T., et al., *HIF1α-Induced Glycolysis Metabolism Is Essential to the Activation
of Inflammatory Macrophages.* Mediators Inflamm, 2017. **2017**: p. 9029327.

80. Modak, M., et al., *CD206+ tumor-associated macrophages cross-present tumor*
antigen and drive antitumor immunity. JCI Insight, 2022. **7**(11).
81. Litvack, M.L., M. Post, and N. Palaniyar, *IgM promotes the clearance of small*
particles and apoptotic microparticles by macrophages. PLoS One, 2011. **6**(3): p.
e17223.
82. Weinstein, J.R., et al., *IgM-Dependent Phagocytosis in Microglia Is Mediated by*
Complement Receptor 3, Not Fcα/μ Receptor. J Immunol, 2015. **195**(11): p. 5309-17.
83. Muntjewerff, E.M., L.D. Meesters, and G. van den Bogaart, *Antigen Cross-*
Presentation by Macrophages. Front Immunol, 2020. **11**: p. 1276.
84. McNeill, L., et al., *The differential regulation of Lck kinase phosphorylation sites by*
CD45 is critical for T cell receptor signaling responses. Immunity, 2007. **27**(3): p.
425-37.
85. Ji, L., et al., *Insights from DOCK2 in cell function and pathophysiology.* Front Mol
Biosci, 2022. **9**: p. 997659.
86. Hashizume, R., et al., *Pharmacologic inhibition of histone demethylation as a therapy*
for pediatric brainstem glioma. Nat Med, 2014. **20**(12): p. 1394-6.
87. Mohammad, F., et al., *EZH2 is a potential therapeutic target for H3K27M-mutant*
pediatric gliomas. Nature Medicine, 2017. **23**(4): p. 483-492.
88. Grasso, C.S., et al., *Functionally defined therapeutic targets in diffuse intrinsic*
pontine glioma. Nature Medicine, 2015. **21**(6): p. 555-559.
89. Balakrishnan, I., et al., *Senescence Induced by BMI1 Inhibition Is a Therapeutic*
Vulnerability in H3K27M-Mutant DIPG. Cell Rep, 2020. **33**(3): p. 108286.
90. Mo, Y., et al., *Epigenome programing by H3.3K27M mutation creates a dependence*
of pediatric glioma on SMARCA4. Cancer Discov, 2022.
91. Panditharatna, E., et al., *BAF complex maintains glioma stem cells in pediatric*
H3K27M-glioma. Cancer Discov, 2022.
92. Mota, M., et al., *Targeting SWI/SNF ATPases in H3.3K27M diffuse intrinsic pontine*
gliomas. Proc Natl Acad Sci U S A, 2023. **120**(18): p. e2221175120.
93. Dahl, N.A., et al., *Super Elongation Complex as a Targetable Dependency in Diffuse*
Midline Glioma. Cell Rep, 2020. **31**(1): p. 107485.
94. Klein, D.C., et al., *FACT regulates pluripotency through proximal and distal regulation*
of gene expression in murine embryonic stem cells. BMC Biol, 2023. **21**(1): p. 167.
95. Jang, M.K., et al., *The bromodomain protein Brd4 is a positive regulatory component*
of P-TEFb and stimulates RNA polymerase II-dependent transcription. Mol Cell,
2005. **19**(4): p. 523-34.
96. Rahme, G.J., et al., *Modeling epigenetic lesions that cause gliomas.* Cell, 2023.
186(17): p. 3674-3685.e14.
97. Mo, J., et al., *Inhibition of the FACT Complex Targets Aberrant Hedgehog Signaling*
and Overcomes Resistance to Smoothened Antagonists. Cancer Research, 2021.
81(11): p. 3105.
98. Okonechnikov, K., et al., *3D genome mapping identifies subgroup-specific*
chromosome conformations and tumor-dependency genes in ependymoma. Nat
Commun, 2023. **14**(1): p. 2300.
99. Jones, P.A., et al., *Epigenetic therapy in immune-oncology.* Nat Rev Cancer, 2019.
19(3): p. 151-161.
100. Gusyatiner, O., et al., *BET inhibitors repress expression of interferon-stimulated*
genes and synergize with HDAC inhibitors in glioblastoma. Neuro Oncol, 2021.
23(10): p. 1680-1692.
101. Leonova, K., et al., *TRAIN (Transcription of Repeats Activates INterferon) in*
response to chromatin destabilization induced by small molecules in mammalian
cells. Elife, 2018. **7**.
102. Somers, K., et al., *Potent antileukemic activity of curaxin CBL0137 against MLL-*
rearranged leukemia. Int J Cancer, 2020. **146**(7): p. 1902-1916.
103. Smart, A.C., et al., *Intron retention is a source of neoepitopes in cancer.* Nat
Biotechnol, 2018. **36**(11): p. 1056-1058.

104. Kwok, D.W., et al., *Tumour-wide RNA splicing aberrations generate actionable public*
neoantigens. Nature, 2025. **639**(8054): p. 463-473.
105. Müller, S., et al., *Single-cell profiling of human gliomas reveals macrophage*
ontogeny as a basis for regional differences in macrophage activation in the tumor
microenvironment. Genome Biol, 2017. **18**(1): p. 234.
106. Wei, J., et al., *Immune biology of glioma-associated macrophages and microglia:*
functional and therapeutic implications. Neuro Oncol, 2020. **22**(2): p. 180-194.
107. Rayner, B.S., et al., *Role of hypochlorous acid (HOCl) and other inflammatory*
mediators in the induction of macrophage extracellular trap formation. Free Radic
Biol Med, 2018. **129**: p. 25-34.
108. Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing*
genomic features. Bioinformatics, 2010. **26**(6): p. 841-2.
109. Ramírez, F., et al., *deepTools: a flexible platform for exploring deep-sequencing*
data. Nucleic Acids Res, 2014. **42**(Web Server issue): p. W187-91.
110. Nordin, A., et al., *The CUT&RUN suspect list of problematic regions of the genome.*
Genome Biol, 2023. **24**(1): p. 185.
111. Robinson, J.T., et al., *Integrative genomics viewer.* Nat Biotechnol, 2011. **29**(1): p.
24-6.
112. Yu, G., L.G. Wang, and Q.Y. He, *ChIPseeker: an R/Bioconductor package for ChIP*
peak annotation, comparison and visualization. Bioinformatics, 2015. **31**(14): p.
2382-3.
113. Yu, G., et al., *clusterProfiler: an R package for comparing biological themes among*
gene clusters. Omics, 2012. **16**(5): p. 284-7.
114. Liu, T., et al., *Cistrome: an integrative platform for transcriptional regulation studies.*
Genome Biol, 2011. **12**(8): p. R83.
115. Lun, A.T. and G.K. Smyth, *csaw: a Bioconductor package for differential binding*
analysis of ChIP-seq data using sliding windows. Nucleic Acids Res, 2016. **44**(5): p.
e45.
116. Holliday, H., et al., *Inhibitor of Differentiation 4 (ID4) represses mammary*
myoepithelial differentiation via inhibition of HEB. iScience, 2021. **24**(2): p. 102072.
117. DeWeirdt, P.C., et al., *Optimization of AsCas12a for combinatorial genetic screens in*
human cells. Nat Biotechnol, 2021. **39**(1): p. 94-104.
118. Dempster, J.M., et al., *Chronos: a cell population dynamics model of CRISPR*
experiments that improves inference of gene fitness effects. Genome Biol, 2021.
22(1): p. 343.
119. Tsherniak, A., et al., *Defining a Cancer Dependency Map.* Cell, 2017. **170**(3): p. 564-
576.e16.
120. Ianevski, A., A.K. Giri, and T. Aittokallio, *SynergyFinder 3.0: an interactive analysis*
and consensus interpretation of multi-drug synergies across multiple samples.
Nucleic Acids Res, 2022. **50**(W1): p. W739-w743.
121. Khan, A., et al., *Dual targeting of polyamine synthesis and uptake in diffuse intrinsic*
pontine gliomas. Nat Commun, 2021. **12**(1): p. 971.
122. Corces, M.R., et al., *An improved ATAC-seq protocol reduces background and*
enables interrogation of frozen tissues. Nat Methods, 2017. **14**(10): p. 959-962.
123. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat
Methods, 2012. **9**(4): p. 357-9.
124. Tarasov, A., et al., *Sambamba: fast processing of NGS alignment formats.*
Bioinformatics, 2015. **31**(12): p. 2032-4.
125. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013.
29(1): p. 15-21.
126. Li, B. and C.N. Dewey, *RSEM: accurate transcript quantification from RNA-Seq data*
with or without a reference genome. BMC Bioinformatics, 2011. **12**: p. 323.
127. Cmero, M., et al., *MINTIE: identifying novel structural and splice variants in*
transcriptomes using RNA-seq data. Genome Biol, 2021. **22**(1): p. 296.

128. Jürges, C., L. Dölken, and F. Erhard, *Dissecting newly transcribed and old RNA using GRAND-SLAM.* Bioinformatics, 2018. **34**(13): p. i218-i226.

129. Rummel, T., L. Sakellaridi, and F. Erhard, *grandR: a comprehensive package for nucleotide conversion RNA-seq data analysis.* Nat Commun, 2023. **14**(1): p. 3559.

130. Harney, D.J., et al., *Proteomic Analysis of Human Plasma during Intermittent Fasting.* J Proteome Res, 2019. **18**(5): p. 2228-2240.

131. Cox, J. and M. Mann, *MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.* Nat Biotechnol, 2008. **26**(12): p. 1367-72.

132. Tyanova, S., et al., *The Perseus computational platform for comprehensive analysis of (prote)omics data.* Nat Methods, 2016. **13**(9): p. 731-40.

133. Livak, K.J. and T.D. Schmittgen, *Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method.* Methods, 2001. **25**(4): p. 402-408.

134. du Chatinier, A., et al., *Generation of immunocompetent syngeneic allograft mouse models for pediatric diffuse midline glioma.* Neurooncol Adv, 2022. **4**(1): p. vdac079.

135. Patel, S.K., et al., *Generation of diffuse intrinsic pontine glioma mouse models by brainstem-targeted in utero electroporation.* Neuro Oncol, 2020. **22**(3): p. 381-392.

136. Law, A.M.K., et al., *Andy's Algorithms: new automated digital image analysis pipelines for FIJI.* Sci Rep, 2017. **7**(1): p. 15717.

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7