Object-translocation induces event coding in the hippocampus

Ya-Li Duan[1,3,4], Shu-Yi Hu[1,3,4], Xiao-Fan Ge[1,2], Jia-Li Long[1,3], Chu Deng[1,3], Yu-Ming Sun[1,3], Cheng-Ji Li[1,2], Rong Zhang[1,2], Xing Cai[1,2] & Li Lu[1,2,5]

1. Key Laboratory of Genetic Evolution & Animal Models, and Key Laboratory of Animal Models & Human Disease Mechanisms of Yunnan Province, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650201, China
2. National Research Facility for Phenotypic & Genetic Analysis of Model Animals (Primate Facility), Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650107, China
3. Kunming College of Life Science, University of Chinese Academy of Sciences, Kunming, Yunnan 650204, China
4. Equal contribution
5. Corresponding author

Word counts: A148-I638- R2867-D1179-M3183-F2547-R71

Abstract

Episodic memory refers to the recollection of personally experienced events anchored in spatial contexts. Salient objects within a navigation environment serve as both landmarks and event cues; however, the nuanced integration of these elements during episodic memory formation is not well understood. The mnemonic functions of the hippocampal CA1 region are partially organized along its transverse axis, with proximal and distal subregions preferentially processing spatial and non-spatial information, respectively. Based on comprehensive analysis of neuronal responses along the CA1 proximodistal axis to manipulations of environmental cues and object features, we identified an event-like population coding within CA1 following object displacement. In addition, the neuronal responses at both initial and relocated object locations were shaped by distinct convergent inputs. These findings highlight the specific roles that objects may play in spatial navigation and memory encoding, providing novel insights into the selective processing of object-related information within the entorhinal-hippocampal network.

Keywords:

episodic memory, object, landmark, event, CA1 transverse axis, population coding

Significance Statement

Salient objects within a navigation environment serve as both landmarks and event cues critical for episodic memory. However, the enigmatic contribution of objects to the "where" and "what" components of episodic memory remains poorly understood. Here we comprehensively analyzed the neuronal responses in the hippocampal CA1 region along its transverse axis, to manipulations of environmental cues and object features. Our results revealed an event-like population coding within CA1 following object displacement, suggesting a minimal landmark role these objects play during memory formation. This study provides novel insights into the selective and flexible processing of object-related information within the entorhinal-hippocampal network.

Introduction

The hippocampus is crucial for the formation of episodic memory, representing personally experienced events within specific spatial frameworks [1]. In rodents, the collective localized firing of hippocampal neurons, known as place cells, generates a cognitive map of the spatial environment, providing a structural basis for linking episodic memories [2]. Place cells are capable of reorganizing their firing patterns, a process termed remapping, to accommodate environmental changes or novel experiences. This dynamic reorganization facilitates the decorrelation of overlapping neural signals, thereby preventing interference between distinct memories [3, 4]. When objects are introduced into the navigation space, hippocampal place cells undergo substantial changes in activity, including modifications in the size, number, and spatial positioning of their receptive fields [5-10]. Among these diverse responses, some hippocampal cells fire specifically at the location of an object [5], while others fire at a fixed distance and direction from the object, independent of its identity [8]. These unique behaviors, observed in "landmark-vector cells" and "object cells", likely reflect adaptations to landmark shifts critical for spatial representation and to local object-related events encountered during navigation. This diversity suggests a multifaceted role of salient objects in spatial mapping and memory encoding. However, the complex interplay between landmark and event-related information in the hippocampus complicates the dissection of their individual contributions to

the "where" and "what" aspects of episodic memory, with the precise role of objects in these

processes still unclear.

The anatomical and functional segregation within the entorhinal-hippocampal network offers

critical insights into the complexity of episodic memory formation. The entorhinal cortex (EC)

integrates multimodal sensory information and conveys highly processed signals to the

hippocampus [11]. Within the EC, the medial region (MEC) contains several specialized cell

types, including grid cells [12, 13], aperiodic spatial cells [14], head direction cells [15], border

cells [16], object-vector cells [17], and speed cells [18], which collaboratively encode

landmark-based distance and directional information, dynamically mapping the spatial location

of the animal within the environment [19, 20]. In contrast, the lateral EC (LEC) exhibits

minimal spatial selectivity in open fields [21, 22], with neurons predominantly representing

non-spatial aspects of the animal's experience, such as objects [23, 24], odors [25, 26], time

[27], memory traces [24], and egocentric positioning relative to landmarks [28], providing the

hippocampus with a temporal progression of events [2]. Inputs from the EC are partially

segregated along the CA1 transverse axis, where MEC neurons primarily project to the

proximal CA1 (pCA1), and LEC neurons preferentially target the distal CA1 (dCA1) [29, 30].

Immediate-early gene expression analyses have indicated distinct mnemonic functions along

the CA1 proximodistal axis, with dCA1 preferentially processing non-spatial information, such

as time, odor, and objects, across various behavioral tasks, while pCA1 is more engaged in tasks

requiring spatial memory [31-36]. Electrophysiological studies have further demonstrated a

stronger spatial modulation in pCA1 [37, 38] and a more coherent response to conflicting

landmarks in dCA1 [39], as well as coordinated activity between the LEC and dCA1 in odor-

associated memory encoding [25, 40]. These findings suggest that the entorhinal-hippocampal

network processes landmark and event information with distinct population coding in pCA1

and dCA1, offering a potential resolution to the enigmatic role of objects in self-localization

and event integration within episodic memory.

In this study, we aimed to distinguish the "where" and "what" components of local object

memory by analyzing place cell activity across the proximodistal CA1 axis as rats navigated an

open field with or without objects. Our findings revealed a complex population response to both

the objects themselves and the environmental changes resulting from their placement.

Manipulating the identity and location of one object elicited stronger population responses in

dCA1 than in pCA1, suggesting that the newly introduced objects were primarily processed as

events rather than landmarks within the entorhinal-hippocampal network. These results

highlight the complexity of information integration that occurs during spatial navigation and

memory formation.

Results

Object-induced local and remote population responses in CA1

To investigate whether CA1 responses to discrete objects varied along the proximodistal axis,

we recorded place cell activity across CA1 subregions (Figure S1) when two identical familiar

objects were introduced to a recording box during trials 2 and 3 of a four-trial experimental

paradigm (Figure S2A–B). To enable cross-animal comparisons, recording positions were

normalized for each tetrode, dividing CA1 into three equal-sized bands: pCA1, middle CA1

(mCA1), and dCA1. While the introduction of objects did not significantly alter overall rat

behavior (Figure S2C), CA1 pyramidal cells showed substantial changes in firing patterns,

primarily through altered firing rates around the objects (Figure 1A). Notably, dCA1 exhibited

the lowest proportion of place cells (Figures S2D & S3) but the highest proportion of rate-

modified neurons, while mCA1 contained the highest proportion of stable cells (Figure 1B).

Given that hippocampal place cells collectively signal the presence of objects at a population

level [10], we analyzed object-related population coding across CA1 subregions using

population vector correlation (PVC) to capture composite distribution changes in firing location

and rate [41]. Consistent with prior studies [10], trial 4 exhibited a pronounced hysteresis effect

(Figure S2E), prompting exclusion of these data from analysis to avoid confounding carryover

effects. PVC analysis revealed high correlation between trials 2 and 3 (matching conditions) in

all CA1 subregions. As expected, correlations decreased between trials 1 and 2 (before and after

object placement), with the most substantial changes in PVC occurring near the objects (Figure

1C). Additionally, dispersed activity shifts distant from the objects were detected in both pCA1

and dCA1, indicating that these responses may not directly reflect the landmark or event

features of the objects, but rather environmental changes triggered by object placement. The

reduction in PVC gradually diminished with object distance from mCA1, consistent with

previous findings [10]; however, pCA1 and dCA1 showed either sustained or increased PVC

for regions distant from the objects (Figure 1D). Notably, a remote population response to

objects was not observed in the CA3 region, which projects heavily to CA1 (Figure S2F–H),

suggesting that direct inputs from the entorhinal cortex may shape this remote response to

objects in CA1.

Overall population responses to object introduction differed significantly across CA1

subregions, with the most pronounced PVC reduction observed in pCA1 (Figure S2E&I). To

clarify object-induced activity changes, we defined an "object region" as the area within a 20

cm radius around the center of each object, following the distance used in a prior study [10].

While parallel changes in PVC within the object region were observed across CA1 subregions,

mCA1 cells displayed minimal activity changes outside the region of interest (ROI) following

object placement (Figures 1E and S2J). Thus, the introduction of familiar objects elicited a

profound population response in the CA1 region, integrating multiple object-related

information components. Neurons in mCA1 demonstrated a comparable local response to the

objects but exhibited the weakest remote population reorganization compared to neurons in

both pCA1 and dCA1.

V-shaped population coding of non-spatial information along CA1 transverse axis

To assess whether the remote responses to object placement in pCA1 and dCA1 were indeed

driven by changes in non-spatial cues within the arena, we recorded CA1 neuronal activity

during a color-reversal task, a conventional non-spatial manipulation paradigm involving the

navigation of a square box with alternating black or white walls (Figure S4A–B). As anticipated

from prior work [41-43], rate remapping, characterized by alterations in firing rates within place

fields while maintaining consistent firing locations [4], was induced in CA1 (Figure 2A). A

linear decrease in spatial modulation from pCA1 to dCA1 in the open field was observed

(Figures S4–S5), consistent with earlier findings [37, 38]. We first examined population coding

of color across CA1 subregions by comparing PVCs between conditions. While population

vectors (PVs) correlated strongly between matching conditions in all CA1 subregions,

correlations were weaker between black and white wall configurations. Differences in PVC

between conditions with reversed wall colors and matching wall colors were significant across

subregions (Figures 2B & S6A). Similar to the remote population response to objects, mCA1

exhibited the smallest change in PVC between condition pairs (Figure 2B–C).

We next examined rate coding in individual place cells, finding that the proportions of rate-

remapped cells between black and white conditions differed across CA1 bands, with mCA1

containing the lowest percentage of rate-remapped neurons (Figure 2D). Rate remapping was

quantified using two metrics: spatial correlation (SC), reflecting the degree of rate redistribution

for each cell when firing location remains constant, and rate difference (RD), representing the

normalized overall change in firing rate irrespective of the firing location of the cell. SC

distributions closely resembled PVC distributions along the CA1 transverse axis, with

significant SC reductions due to color change in pCA1 and dCA1 compared to mCA1 (Figure

2E). Consistent with the observed SC pattern, RD values also differed significantly between

condition pairs and across CA1 subregions, with pCA1 and dCA1 showing more pronounced

rate redistribution than mCA1 upon wall color reversal (Figure 2F, left). Notably, pCA1

displayed a significantly higher RD increase between condition pairs (Figure 2F, right). The

minimal rate remapping in mCA1 was not due to individual animal variations, as similar

patterns emerged when analyses were restricted to simultaneously recorded pCA1, mCA1, and

dCA1 neurons (Figure S6B–C). A similar outcome was obtained when the weighted RD for

individual place fields (RD_f) was computed (Figure S6D–F).

To visualize population response patterns across the CA1 transverse axis, mean SC and RD

values were plotted by tetrode location, revealing a V-shaped distribution from the proximal to

distal end of CA1 for both measures. Analyses of proximal and distal tetrodes independently

showed significant correlations within each half for SC and RD (Figure 2G–H), with quadratic

polynomial curve fitting providing the best explanation for changes along the entire axis. A

similar pattern was observed when RD_f was plotted as a function of tetrode location (Figure

S6G). To further verify activity change patterns along the CA1 transverse axis, a naïve Bayes

decoder was trained on place cell activity from trials with white walls (trials 2 and 3) to predict

rat location in a black-walled environment (trial 1). For experiments with over 30

simultaneously recorded place cells across subregions, decoding accuracy was highest in the

mCA1 subregion (Figure S6H), indicating minimal activity change between the black and white

configurations.

To summary, hippocampal CA1 activity during the color-reversal task demonstrated a V-shaped

population response along the transverse axis under non-spatial cue manipulation. Firing rate

redistribution was minimum near the middle of the axis and increased progressively towards

both the proximal and distal ends. This pattern of population coding closely resembled the

remote response observed in the CA1 region to object placement, suggesting that the latter may

also be driven by non-spatial environmental changes.

Extensive reorganization of pCA1 activity in response to landmark changes

Given that the MEC and LEC selectively convey landmark and event-related information to the

proximal and distal regions of CA1, respectively [2], we hypothesized that landmark changes

would elicit stronger population responses in pCA1, while event changes would lead to elevated

activity alterations in dCA1. Thus, we analyzed population activity along the CA1 transverse

axis in two rooms with distinct spatial configurations and distal landmarks (Figure S7A–B).

Global remapping, characterized by the engagement of a second population of place cells to

represent a distinct spatial map in the hippocampus [4], was evident at the individual cell level

based on significant changes in the firing location and rate within the recording box (Figure

3A), in line with previous reports [41-43]. PVCs were high for trials within the same room

across all subregions but decreased substantially between trials in different rooms (Figures 3B

& S7C). The reduction in PVC was region-dependent, with pCA1 exhibiting the greatest

decrease, mCA1 showing a moderate decrease, and dCA1 demonstrating the smallest decrease

(Figure 3C), suggesting that pCA1 was more sensitive to changes in salient landmarks

compared to the distal regions of CA1.

To further examine spatial sensitivity at the level of individual place cells, we analyzed

population overlap across CA1 subregions, which remained consistent (Figure S7D–E). The

SC of individual place cells, which is sensitive to rate map reorganization, was high for repeated

trials within the same room in all CA1 subregions but decreased markedly between rooms

(Figure 3D, left), indicating considerable shifts in firing location across rooms. Between-room

SC tended to be higher in mCA1 and dCA1 than in pCA1 (Figure S7F). The decrease in SC

with room changes was most pronounced in pCA1 and declined progressively along the

proximodistal axis, as reflected by a strong linear correlation between SC decline and tetrode

location along CA1 (Figure 3D, right). Quadratic polynomial curve fitting provided minimal

additional explanatory power, indicating a predominantly linear gradient from pCA1 to dCA1

(Figure 3E). Similar patterns emerged when using RD to assess neuronal activity levels (Figure

S7G–H). Overall, place cell representations were most orthogonalized in pCA1 when salient

landmarks were altered, with orthogonality decreasing monotonically along the transverse axis,

consistent with previous research [39]. Furthermore, SC variations were significantly correlated

with spatial tuning degree (Figure S8), implying a significant contribution of direct MEC inputs

to landmark coding within pCA1.

Enhanced rate coding in dCA1 in response to object replacement

To assess whether the dCA1 region exhibited increased sensitivity to novel events within the

environment, we replaced one of the familiar objects with a novel one (Figure S9A),

maintaining the landmark properties of the object [17]. Overall, the rats showed consistent

behavior before and after object replacement (Figure S9B). A substantial subset of place cells

adjusted their firing rates in response to the novel object (Figure 4A), with dCA1 exhibiting the

highest proportion of cells responsive to object identity (Figures 4B & S9C). To control for

potential hysteresis effects observed in trial 4 (Figure S9D), our analysis focused on firing rate

shifts between trials 1 and 2 (object identity switched) and between trials 2 and 3 (object identity

unchanged).

Due to minimal changes in activity patterns, no significant regional differences in SC changes

were detected across trials in the novel-object task (Figure S9E). The overall changes in firing

rate for individual place cells were marginally but significantly enhanced in trial pairs before

and after object replacement, with the degree of enhancement strongly dependent on the

location of place cells. Neurons in pCA1 exhibited the lowest increase in RD, followed by

mCA1 and dCA1 (Figure 4C). RD enhancements were significantly correlated with recording

positions (Figure 4D). The linear changes in RD were not related to sample size (Figure S9F)

or exploratory behavior around objects (Figure S9G). Consistent patterns were obtained for RD_f

scores calculated for individual place fields (Figure S9H–I).

The ensemble activity of all pyramidal neurons, as measured by PVC, demonstrated a marginal

decrease following object replacement, with a significant interaction between novelty and CA1

subregions (Figures 4E & S9D). Cell assemblies within pCA1 exhibited the smallest response

to the novel object (Figure 4F), while Bayesian decoding showed that context prediction

accuracy was higher for cells in mCA1 than in dCA1 (Figure S9J). Unlike the object-placement

task, population activity reorganization in response to the novel object was less pronounced and

less localized across CA1 subregions (Figure 4E). The regional differences in response were

primarily attributed to subtle differences in activity changes around the object, with ensemble

activity remaining stable outside the object region (Figures 4G & S9K). In addition, CA3

showed minimal response to object replacement (Figure S9L–M). In conclusion, place cells in

dCA1 displayed greater firing rate redistribution following object replacement, likely due to

preferential input from object-identity-responsive cells in the LEC and potentially from the

perirhinal cortex (PER) [22, 23, 44, 45].

Object-translocation elicited event-like population coding in CA1

Based on the differential encoding of landmarks and events in pCA1 and dCA1 regions, we

next examined whether a relocated familiar object within a known environment would

contribute to spatial mapping or be encoded as an event. To test this, we translocated a familiar

object to a different quadrant of the recording box (Figure S10A). If the object served as a

landmark, we anticipated increased activity in pCA1; alternatively, if it functioned as an event,

a stronger population response would be expected in dCA1. Despite variations in individual

responses to object translocation, the overall discrimination index remained largely unchanged

(Figure S10B). A substantial subset of CA1 place cells adjusted their firing patterns, with many

neurons displaying new firing fields around the relocated object (current-location coding, CLC),

or near its original position (original-location coding, OLC), reminiscent of object-trace cells

reported in the LEC [24] (Figure 5A). Notably, although the dCA1 region had the lowest

proportion of place cells (Figure S10C), it contained the highest proportion of OLC and CLC

neurons, as well as neurons with emergent or disappearing fields outside the region around the

relocated object (ROI) (Figure 5B). A minority of cells maintained consistent firing at a fixed

distance and direction relative to the relocated object, with or without trace activity remaining

in its initial location after relocation, similar to previously reported object-vector cells (OVCs)

in the MEC [17]. However, these OVC-like neurons were rare (<2.2% for all subregions) and

did not differ across CA1 subregions (Figure 5B).

As anticipated from the distribution of relocation-responsive cell types, the dCA1 region

displayed a stronger population response to object relocation compared to its proximal

counterpart. The SC of individual place cells was high when objects were in the same location

(trials 2 vs 3) but slightly reduced after object relocation (trials 1 vs 2) across all CA1 subregions.

The reduction in SC varied significantly by subregion, with the smallest decrease in pCA1 and

the largest decrease in dCA1 (Figure 5C). A significant correlation was observed between SC

change and tetrode location (Figure 5D). The increase in RD between trial pairs mirrored the

SC decrease across CA1 subregions and increased progressively from pCA1 to dCA1 (Figure

5E). A significant correlation was also found between the shift in RD for individual tetrodes

and their recording positions (Figure 5F). These spatially graded changes along the CA1

transverse axis were not influenced by object exploration (Figure S10D), sample size (Figure

S10E–F), or inter-animal variations (Figure S10G). Consistent findings were observed when

RD_f scores were calculated for individual place fields (Figure S10H–I). Correlations between

corresponding PVs across spatial bins displayed similar changes to those observed in SC

(Figures 5G & S10J). Notably, the decline in PVC between trial pairs progressively increased

along the CA1 transverse axis, from pCA1 to dCA1 (Figure 5H). Bayesian decoding also

revealed a marked improvement in position predictions within proximally located CA1 regions

(Figure S10K). Furthermore, significant reductions in PVC were detected for bins within the

ROI in the distal two-thirds of CA1, while pCA1 remained minimally unaffected (Figures 5I &

S10L). Collectively, the minimal reorganization in pCA1 alongside the pronounced response in

dCA1 suggests that relocated objects were not assimilated into the spatial map as fixed

landmarks but were instead encoded as distinct events encountered during navigation.

Distinct inputs shape responses at original and relocated object locations

Given that object relocation primarily elicited event coding within the CA1 region, we next

examined the contributions of specific neuronal subpopulations to this process by

systematically removing each subgroup from the overall population analysis. Excluding OVC-

like neurons, which were more responsive to landmarks, led to a graded increase in population

response to object relocation (Figure 6A). The decline in PVC around the relocated object

before and after relocation increased progressively from pCA1 to dCA1 (Figure 6B). To assess

differential processing at the original and current object locations, we analyzed population

responses at these positions separately and found patterns consistent with those observed within

the combined ROI data (Figure 6C).

Further removal of OLC neurons led to significant changes at original location but not current

location (Figure 6D–E). Notably, the population response to object removal at original object

location was predominantly driven by a small fraction of OLC neurons in pCA1 (3.08% of cells

contributed to 94.61% of activity changes), while a larger fraction of OLC neurons in dCA1

(8.08%) contributed considerably less to activity change (17.44%) (Figure 6F), suggesting the

rapid involvement of other neuronal subgroups in representing past experience from pCA1 to

dCA1. Conversely, excluding CLC neurons led to only mild reductions in PVC change at the

current location (Figure 6G). Although activity changes showed a graded increase (Figure 6H),

the relative reduction in population response due to CLC neuron exclusion remained low

(28.56%–35.37%) and was consistent along the transverse axis (Figure 6I), suggesting that

population coding for the displaced object was stable across CA1. The profound activity change

observed in CA3 at current location, coupled with a weaker response at the original location

(Figure S11A–B), suggests that distinct population coding at original and current object

locations in CA1 may be partially influenced by CA3 inputs. Excluding neurons with field

changes outside the ROI minimally affected responses at either original or current object

location (Figure S11C–D). Overall, event coding in CA1 appears to be shaped by signals from

multiple sources, with CA3 inputs primarily affecting responses at current location and other

inputs influencing responses at original location.

Discussion

The integration of salient objects during episodic memory formation remains a subject of

ongoing debate. Here, we performed a comprehensive analysis of population-level responses

across the CA1 proximodistal axis to evaluate the encoding of newly introduced objects in the

recording environment, alongside traditional spatial and non-spatial manipulations. Results

revealed that object placement triggered localized firing rate changes near the objects and

elicited population-level responses similar to those elicited by non-spatial environmental

changes (Figures 1–2). Furthermore, by altering spatial configuration and object identity, we

validated the segregated encoding of landmarks and events in pCA1 and dCA1 (Figures 3–4).

Notably, our final experiment indicated that pCA1 neurons exhibited minimal activity changes

in response to object translocation, while more distal neurons along the transverse axis

displayed diverse responses to the displaced object, suggesting that the object contributed

minimally to self-localization within an established spatial map (Figures 5–6). Collectively,

these findings suggest that neutral objects, when introduced into a familiar environment, are

primarily processed as transient events rather than integrated landmarks for navigation.

Experience-dependent integration of landmark and event information

Our results challenge the traditional perspective that object-location memory (OLM) relies

predominantly on the MEC over the LEC [46]. While impairments in OLM have been linked

to extensive lesions in both the MEC and LEC [47], or complete MEC lesions and inactivation

of MEC stellate cells [48-50], LEC-specific lesions appear to have no effect on OLM

performance [51, 52]. These discrepancies can be explained by the Hebbian plasticity theory,

which posits that the formation of a cognitive map with accurate positional information is

highly dependent on repeated experience with landmarks [53]. In standard object-location tasks,

brief habituation may not allow for the complete development of a cognitive map, leading to

the incorporation of later introduced objects as prominent landmarks. In our study, however,

objects were introduced after weeks of intensive training, resulting in event coding tied to a

well-established spatial map referencing other established landmarks. The capacity for

landmark changes to prompt immediate grid cell reorganization in the MEC, one synapse

upstream from the hippocampus, may stem from modified firing patterns in entorhinal object-

vector cells that convey allocentric vector information about landmarks [17, 54]. However,

these MEC-driven changes showed minimal reflection in the CA1 region during our object-

relocation experiments (Figure 5), indicating that selective and flexible processing of MEC

inputs in the hippocampus may depend on context-specific mechanisms.

The enhanced population response to object relocation observed in dCA1 likely arises from

convergent inputs from multiple brain regions. Within the LEC, subpopulations of neurons are

tuned to fire at specific object locations or at certain egocentric distances and directions relative

to objects [23, 28]. Object-trace cells within the LEC, which fire exclusively at locations

previously occupied by an object, may convey past experiences to dCA1 [24]. These object-

location-related and experience-dependent activities are evident in the pronounced activity

changes observed at both the original and current object locations in dCA1 (Figure 6).

Furthermore, the PER, which projects to the LEC and dCA1, contains object-responsive cells

similar to those found in the LEC [29, 44, 45]. Additional object-place signals may also

originate from subicular dCA1-projecting neurons, which modulate place cell responses to

object displacement [55]. Recently identified subicular corner cells [56] may further contribute

to this heterogeneity, although their back-projection to dCA1 remains to be determined.

Direct entorhinal inputs shape rate and global remapping patterns in CA1

Our study revealed a distinct V-shaped rate remapping pattern within CA1 in both the color-

reversal and object-placement tasks (Figures 1–2). The pronounced rate redistribution observed

in both pCA1 and dCA1 suggests contributions from both the MEC and LEC to hippocampal

rate coding. Certain LEC neurons are known to undergo remapping in response to contextual

changes in the environment, and lesions in the LEC have been shown to partially impair

hippocampal rate coding [22, 42], supporting the hypothesis that the LEC provides modulated

non-spatial inputs critical for hippocampal rate remapping [57]. In contrast, MEC grid cells

typically maintain stable firing locations within a given spatial context but exhibit reliable rate

redistribution in response to non-spatial manipulations [58-60]. Precise modulation of MEC

activity has been linked to hippocampal remapping [61-63], indicating that rate redistribution

in MEC cells may directly influence place cell firing. The degree of rate remapping along CA1

may therefore reflect the relative dominance of MEC and LEC inputs across the transverse axis.

Neurons in pCA1 or dCA1, predominantly receiving inputs from either the MEC or LEC,

exhibit pronounced rate remapping, whereas mCA1 neurons, receiving balanced inputs from

both the MEC and LEC, display weaker rate remapping due to dynamic interaction between

their competing signals.

Our findings further support the established view that spatial tuning and global remapping in

the CA1 region exhibit a gradient decrease from the proximal to distal end (Figures 3 & S4, S5,

S8) [37-39]. Global remapping in the rat hippocampus is accompanied by a realignment of

activity patterns in grid, head direction, and border cells within the MEC [16, 64], which

appears to minimize spatial information overlap and potentially enhance distinct spatial

representations in pCA1. Conversely, neurons in the LEC, which encode shared non-spatial

features, likely maintain their activity levels, contributing to a more generalized ensemble

coding in dCA1 [36, 65]. This gradient in spatial selectivity and global remapping in the CA1

region reflects differential MEC and LEC inputs, with pCA1 predominantly receiving spatial

signals from the MEC and dCA1 primarily receiving non-spatial signals from the LEC [66].

Differential representation of contextual cues along CA1 transverse axis

We also identified three distinct cue representation patterns along the proximodistal axis of CA1,

which may contribute to hippocampal mnemonic functions. These patterns comprise a gradient

of decreasing spatial information processing, a V-shaped response to non-spatial environmental

changes, and an increase in the representation of local events. These variations likely stem from

selective inputs from specialized neuronal populations in the entorhinal cortex (Figure S12) [17,

19, 22-24, 28]. Our findings suggest the need for further investigation into the causal dynamics

linking selective upstream inputs to specific downstream representations. Future theoretical and

experimental endeavors should aim to elucidate the precise mechanisms that govern these

interactions, providing crucial insights into the neural basis of spatial navigation, episodic

memory, and the broader cognitive functions of the hippocampus.

Limitations of the study

This study has two primary limitations. First, CA1 neuronal activity was recorded using a

hyperdrive with movable tetrodes, constraining our capacity to capture activity from a large

population of neurons across CA1 subregions simultaneously in individual rats. This restriction

may have introduced inter-animal variability, potentially confounding the interpretation of our

results. Second, the complexity in CA1 extends to its radial organization into superficial and

deep sublayers, each with distinct characteristics and functions [67]. Our tetrode-based

methodology was unable to differentiate between these sublayers. As such, future studies

should incorporate two-photon calcium imaging and multi-shank probe recording to monitor

neural activity within specific sublayers. These advanced techniques could provide further

insights into laminar contributions to functional heterogeneity along the CA1 transverse axis,

leading to a more nuanced understanding of landmark and/or event coding within the

hippocampus.

Methods

Subjects

The experiments were conducted at the Kunming Institute of Zoology, Chinese Academy of

Sciences (Kunming, China) in compliance with the guidelines for the care and use of laboratory

animals. The study received approval from the Institutional Animal Care and Use Committee

of the Kunming Institute of Zoology (KIZ-IACUC-RE-2021-06-019). Data were collected from

22 male Long-Evans rats (3–4 months old, 500–600 g at the time of implantation). Five rats

from a previous study [43] contributed proximal-to-middle CA1 data. The remaining 17 rats

were implanted with a multi-tetrode "hyperdrive" targeting the right hippocampus. The animals

were maintained on a reversed 12-hour light/dark cycle (lights off from 9.00 am to 9.00 pm)

under standard laboratory conditions (temperature: 20–23 °C, relative humidity: 40%–60%).

Following surgery, the rats were housed individually, and experimental procedures were

conducted during the dark phase. During the experimental period, the animals were mildly food

restricted to maintain their weight at 85%–90% of free-feeding body weight.

Electrode preparation

The multi-tetrode "hyperdrive", consisting of 18 independently movable tetrodes made of 17

μm polyimide-coated platinum-iridium (90%–10%) wire (California Fine Wires, USA), was

prepared as described previously [68]. The electrode tips were plated to reduce impedance to

150–250 kΩ at 1 kHz (Biomega, Bio-Signal Technologies, China) and sterilized with ethylene

oxide (Sanqiang Medical, China) within 48 hours before implantation.

Surgery

Rats were anesthetized with isoflurane (air flow: 0.8–1.0 L/min, 0.5–3% isoflurane vol/vol

mixed with oxygen), with the concentration adjusted based on physiological monitoring (RWD

Life Science, China). Upon induction of anesthesia, Metacam (2 mg/mL, 1 mg/kg body weight),

Baytril (50 mg/mL, 5 mg/kg), and atropine (0.5 mg/mL, 0.1 mg/kg) were administered

subcutaneously. The animal's head was then fixed in a stereotaxic frame (RWD Life Science,

China). Local anesthetic (2% lidocaine, 200 μL) was applied under the skin before making the

incision. The skull and dura were removed, followed by immediate hyperdrive implantation.

Tetrodes were inserted at the cortical surface above the right hippocampus (CA1:

anteroposterior (AP) from 3.2 to 5.6 mm, mediolateral (ML) from 1.0 to 4.5 mm relative to

bregma; CA3: AP: 4.2 to 4.9 mm, ML: 2.4 to 4.0 mm). The hyperdrive was secured to the skull

with jewelers' screws and dental cement. Grounding was achieved by connecting two screws

above the cerebellum to the hyperdrive ground, while two reference tetrodes were positioned

in an inactive area of the cortex for differential recording.

Electrophysiological recordings

For the first 3–4 weeks post-surgery, the tetrodes were advanced daily until they reached the

dorsal CA1 pyramidal cell layer, characterized by large-amplitude theta-modulated complex-spike activity. On the recording day, the electrodes were kept stationary to ensure stable recordings. Broadband neural signals were captured from the hyperdrive using a multichannel data acquisition system (Zeus, Bio-Signal Technologies, China). Unit activity was amplified ($\times$ 192) and band-pass filtered (300 to 7 500 Hz). Spike waveforms that exceeded the -50 μV threshold were time-stamped and digitized at 30 kHz for 1 ms. Local field potential (LFP) signals, one per tetrode, were recorded in the 0.3–300 Hz frequency band at a sampling rate of 1 000 Hz. The animal's position was monitored in real-time by tracking light-emitting diodes (LEDs) on the headstage at 50 Hz using a Cyclops video tracking system (Bio-Signal Technologies, China).

Behavioral paradigms

Each task consisted of four 10-minute exploration trials within the recording box, interspersed with five 5-minute rest periods in a flowerpot placed on a pedestal outside the box. The exploration trials were designed in an A-B-B-A configuration, where A and B represented distinct test conditions. To encourage exploration, cookie crumbs were randomly scattered throughout the recording box. One week after surgery, the mildly food-restricted rats were trained daily in the color-reversal task in two separate rooms until they could traverse the entire arena within 10 minutes in all trials. Once the tetrodes reached the CA1 cell layer, the animal underwent sequential testing and recording in the color-reversal, two-room, and three object-related tasks.

Test environments

Recording boxes (100 cm $\times$ 100 cm) consisted of a metal frame and four individually exchangeable plastic walls (black on one side, white on the other; 100 cm $\times$ 50 cm). Cue cards (white cue card for black walls and vice versa; 50 cm $\times$ 30 cm) were placed at the center of one wall. Distal background cues were obscured by curtains encircling the recording box (120 cm $\times$ 120 cm) for the color-reversal and object-related tasks but not for the two-room task. The pedestal with the flowerpot was positioned between the test box and experimenter, outside the curtains. The recording rooms for the two-room task were similar in size but featured different

511 spatial arrangements.

Color-reversal task

514 As detailed previously, this task was designed to elicit rate remapping in the hippocampus [41-

515 43]. Rats were tested in the recording box at a fixed location over four trials daily, with trials 1

516 and 4 conducted with black walls and trials 2 and 3 conducted with white walls. The floor of

517 the recording box was cleaned with 75% ethanol after each trial.

Two-room task

520 This task was designed to evoke global remapping in the hippocampus [41, 43]. Rats were

521 tested in similar recording boxes with black walls located at fixed locations in two separate

522 recording rooms, with trials 1 and 4 conducted in room A and trials 2 and 3 conducted in room

523 B. The floor of the recording box was cleaned with 75% ethanol after each trial.

Tasks with objects

Object-placement task: Animals were tested in the black recording box at a fixed location over

four trials, with two identical familiar objects (glass bottles filled with white stones) placed in

quadrants 3 and 4 of the box for trials 2 and 3. Novel-object task: Animals were tested in the

black recording box containing two objects over four trials, with the familiar object in quadrant

4 replaced by a novel object in trials 2 and 3. Object-relocation task: Animals were tested in the

black recording box containing two familiar objects over four trials, with the object in quadrant

4 relocated to quadrant 1 for trials 2 and 3. For each task, the floor of the recording box and the

objects were thoroughly cleaned with 75% ethanol before each trial.

Occupation time and discrimination index

536 Occupation time (OT) and discrimination index (DI) were employed to quantify preference for

537 objects in the object-placement task and for relocated/novel objects in the object-relocation and

538 novel-object tasks. OT was defined as the proportion of time spent around the objects relative

539 to the total time spent in the recording box, and DI was calculated as:

$$540 \qquad DI = \frac{T_r - T_u}{T_r + T_u};$$

where T_r is the time spent around the relocated/replaced object and T_u is the time spent around

the unchanged object. A higher DI value indicates a greater preference for the relocated/novel

object.

Histology and recording sites

Following the completion of electrophysiological recordings, the electrodes were maintained

in their positions. The rats were subsequently euthanized and perfused with formalin for brain

extraction. Frozen brain sections (40 μm thick) were cut on a cryostat (KEDEE, KD-2950,

China), and stained using cresyl-violet (Sigma Aldrich, USA, CAS: 10510-54-0, Cat# C5042-

10G). the Nissl-stained brain sections were imaged using a bright-field microscope (2×/0.06, 4

×/0.13 NA objectives, COIC, UB103i, China), with UOPView software (v2.0). The location of

each tetrode tip was then identified and recorded. Hippocampal subregions were delineated

based on established criteria [29, 37]. ImageJ (v1.52a) was used to estimate the normalized

position along the proximodistal axis of the CA1 region for each recording site. Hand-drawn

lines tracing the CA1 pyramidal cell layer were extended from the CA2/CA1 border to the

CA1/subiculum border, passing through the recording site on each section. The distance from

the CA2/CA1 border to the recording site was measured and normalized by the total length of

the CA1 region on the same section. Based on these normalized positions, recording sites were

categorized into proximal (pCA1, 0–0.33), middle (mCA1, 0.33–0.67), and distal (dCA1, 0.67–

1) subregions.

Spike sorting and cell classification

Spike sorting was manually performed using the MClust graphical cluster-cutting tool in

Matlab (v4.4, A.D. Redish). The process involved the analysis of two-dimensional projections

of the multidimensional parameter space, consisting of waveform amplitudes, waveform

energies, and waveform peak-valley differences. Waveform and autocorrelation functions were

also employed as tools for separation and classification. Clustering was applied on the entire

dataset of each experiment, with the spikes subsequently divided into four exploration trials.

Putative excitatory cells were distinguished from putative interneurons and bypassing axons

based on spike width (peak-to-trough waveform durations greater than 250 μs), average rate

(less than 5 Hz), and occasional presence of bursts. Only cells classified as putative excitatory

neurons were included in subsequent analyses. Cells recorded repeatedly in the same task were

excluded from analyses.

Rate maps and place fields

Position estimates were derived from tracking the LEDs on the headstage connected to the

hyperdrive. Data were filtered to include only epochs with instantaneous running speeds

between 2.5 and 100 cm/s. The mean firing rate for each cell was calculated by summing the

total number of spikes and dividing it by the time spent in that trial. Place field analyses were

conducted exclusively for cells with a mean firing rate of 0.10 Hz or higher, defined as active

cells. Spatial firing rate distributions (place fields) for each well-isolated neuron were generated

by summing the spikes within specific location bins (5 cm × 5 cm), dividing by the duration of

time spent by the rat in that location, and applying Gaussian smoothing centered on each bin,

as described previously [43]. A place field was defined as a contiguous region of at least 225

cm^2 (nine or more 5 × 5 cm bins) where the firing rate exceeded 20% of the peak rate and the

peak firing rate of the area was 1 Hz or greater. The number of non-overlapping place fields

was estimated for each cell.

Neuron subpopulations in the object-relocation task

Neurons classified as object-vector cell-like were characterized by their exclusive activation at

a specific distance and direction relative to the displaced object, both before and after its

relocation. Furthermore, the center of the receptive field must be situated within a region 20 cm

around the object. Neurons designated as original and current location coding cells were

identified by the presence or absence of at least one receptive field at the object's original or

current location (within a 20 cm radius), respectively.

Theta modulation

The theta modulation index (TMI) was computed to assess the temporal modulation of firing

within the theta frequency range (7–10 Hz) of each cell, as described previously [69]. A

temporal autocorrelogram was constructed by tallying the occurrences of spikes within each 5-

ms bin over a range of 0 to 500 ms from a reference spike at time 0, then dividing by the total

trial length to determine the rate of occurrence for each bin. The TMI was calculated as:

$$TMI = \frac{P-T}{P+T};$$

where P and T represent the theta modulation peak (mean of bins 100–140 ms) and trough

(mean of time bins 50–70 ms), respectively.

Spatial information, coherence, and stability

The spatial information rate in bits per second (SI_r) and spatial information content in bits per

spike (SI_c) were calculated as described in previous research [70]:

$$SI_r = \sum_i p_i \lambda_i \log_2 \frac{\lambda_i}{\lambda}; \ \ SI_c = \sum_i p_i \frac{\lambda_i}{\lambda} \log_2 \frac{\lambda_i}{\lambda};$$

where λ_i is the mean firing rate of a unit in the i-th bin, λ is the overall mean firing rate, and p_i

is the probability of the animal being in the i-th bin (occupancy in the i-th bin / total recording

time). Pyramidal neurons with SI_c scores above the 95[th] percentile of the shuffled distribution

were classified as place cells.

Spatial coherence was estimated as the first-order spatial autocorrelation of the unsmoothed

place field map, i.e., mean correlation between the firing rate of each bin and average firing

rate in the eight adjacent bins.

Spatial stability was determined by comparing the firing pattern in the first half of each trial to

that in the second half. The rate map for each half was smoothed and binned into 5 cm × 5 cm

matrices, and the firing rates in the bins of the two half-trial maps were correlated for each trial.

Spatial correlation (SC)

Firing patterns were compared across trials using a SC procedure. Each map was smoothed and

binned into 5 × 5 cm matrices, and the firing rates in the pixels of the two maps were correlated

for each cell. SC between two inactive trials (firing rates below 0.1 Hz) was excluded to avoid

artifacts. For the two-room task, rate maps were rotated in 90° steps to identify the maximum

SC score when compared across rooms.

Rate difference (RD) and rate difference by field (RD$_f$)

The change in overall firing rate across trials was calculated as RD:

634
$$RD = \left| \frac{FR_1 - FR_2}{FR_1 + FR_2} \right|;$$

635   where FR_1 is the mean firing rate of a cell in one trial and FR_2 is the mean firing rate of the

636   same cell in a second trial, with RD = 1 indicating that the cell was active in one trial only and

637   RD = 0 indicating that the firing rates in the two trials were the same.

639   The RD$_f$ was computed to mitigate the neutralization effect across individual place fields from

640   the same cell:

641
$$RD_f = \sum_i w_i RD_i \, ;$$

where RD_i is the rate difference of the i-th place field and w_i is the weight of the i-th place field

estimated from its firing rate. RD$_f$ was not computed when changes in firing location occurred.

Population vector correlation (PVC)

Rate vectors were constructed for the entire population of cells recorded under each condition,

including those with firing rates below 0.1 Hz and spatial information content below 0.9586

bits/spike. This process involved arranging all rate maps for all cells in an x-y-z stack, with x

and y denoting the two spatial dimensions (20 × 20 bins) and z representing the cell-identity

index [41]. For the two-room task, rate maps were rotated analogously to the SC analyses

performed for each cell. The distribution of bin rates along the z axis for a specific x-y location

represented the composite population vector for that location. Population vectors from

corresponding positions were then correlated across pairs of trials. Normalized PVC change

was calculated as:

655
$$Normalized \; \Delta PVC = \frac{\Delta PVC \; for \; all \; cells - \Delta PVC \; without \; certain \; neuron \; population}{\Delta PVC \; for \; all \; cells}.$$

Shuffling

Shuffling of spatial information was conducted on a cell-by-cell basis, as described previously [37]. Each CA1 cell recorded from the color-reversal task underwent 100 permutations. In each permutation, the entire sequence of spikes fired by the cell was time-shifted along the animal's path by a random interval ranging from 20 seconds to the total trial length minus 20 seconds. The end of the trial was wrapped back to the beginning to allow for circular displacements. A rate map was generated for each permutation, and the distribution of spatial information values across all permutations for all CA1 cells was determined. A putative excitatory neuron was classified as a place cell if its spatial information content exceeded the 95[th] percentile of the shuffled distribution (0.9586 bits/spike).

For each CA1 band, a control distribution for SC analyses was established using a random permutation procedure. This procedure involved all active maps (mean firing rates higher than 0.10 Hz) from all cells recorded within that band [43]. For each permutation, two distinct active maps from different cells were randomly selected and correlated in 90° rotation steps to achieve the highest correlation score. In total, 10 000 permutations were performed for each CA1 band. This method preserved the rate maps within each band while eliminating connections between maps. The distribution of random correlations from all 10 000 permutations was then calculated, and the 95[th] percentile was determined.

Bayesian decoding

A Bayesian decoding algorithm was employed to convert ensemble spiking activity into positional information for each experiment. The decoding process used a 2 000 ms sliding time window, advancing by 200 ms with each step. The probability of a rat being at position x, given the number of spikes n from each unit recorded within the time window, was estimated using Bayes' rule [71]:

$$P(x \mid n) = \frac{P(n|x)P(x)}{P(n)};$$

The probabilistic likelihood $P(n|x)$ was calculated using the probability mass function of the Poisson distribution to determine the likelihood of observing a specific number of spikes given

686   the activity pattern of a neuron:

687
$$P(n \mid x) = \frac{e^{-\lambda}\lambda^n}{n!};$$

where λ is the expected number of spikes based on the tuning curve and n is the actual observed

number of spikes within the given time window. The prior probability of position $P(x)$ was set

to the frequency of the animal's presence in each state within the training data. The normalizing

constant $P(n)$ ensured that the posterior probability $P(x|n)$ summed to 1. For visualization and

maximum decoding performance estimation, only experiments with over 30 simultaneously

recorded place cells in both CA1 bands were considered. Downsampling was applied to

equalize cell numbers and repeated three times to determine the mean decoding accuracy value.

For position decoding, 90% of the data from the four trials served as the training set, while the

remaining 10% was used as the test set. This cross-validation procedure was repeated 10 times

(10-fold validation). For context decoding, data from trials 2 and 3 formed the training set, with

90% of trial 1 data serving as the test set. This procedure was also repeated 10 times. Decoding

accuracy was expressed as:

700
$$R^2 = 1 - \frac{\Sigma_i(\hat{y}_i - y_i)^2}{\Sigma_i(y_i - \bar{y})^2};$$

701   Where $\hat{y}_i$ and y_i are the predicted and true positions of the i-th time bin, respectively, and $\bar{y}$ is

702   the mean position value.

704   **Fitting**

705   The EzyFit Toolbox in Matlab (v2.44) was used to compute curve fitting for the average data

706   obtained from the tetrodes. The explained variance was calculated as:

707
$$R^2 = \frac{\Sigma_i(\hat{y}_i - \bar{y})^2}{\Sigma_i(y_i - \bar{y})^2};$$

where $\hat{y}_i$ is the predicted value based on the curve fitting and normalized position of each tetrode,

y_i is the average tetrode data, and $\bar{y}$ is the mean value across all tetrodes.

Statistical procedures

Results are presented as medians and interquartile ranges (IQR) due to the non-normal

distribution of most data. The Kruskal-Wallis test was used to compare the firing properties of

cells across CA1 bands. The Mann-Whitney U test was employed to compare two groups of

unrelated measures, while the Wilcoxon signed-rank test and paired *t*-test were employed to compare two related measures. Regional differences in remapping between two conditions were assessed using two-way repeated-measures ANOVA with a 3×2 factor design, considering condition pairs (matching or different) as within-subject factors and CA1 bands as between-subjects factors. Two-way repeated-measures ANOVA was also used when comparing data across recording experiments. The Pearson chi-square test and Fisher's exact test were applied to compare frequency distributions across groups, and Pearson correlation was calculated to measure the linear correlation between two variables. The binomial test was applied to determine whether the observed frequency differed from the expected frequency. Holm-Bonferroni *post-hoc* tests were used to identify significant differences between pairs. The significance level was set at $P < 0.05$ for two-tailed tests. All statistical analyses were conducted using SPSS v25 (IBM, USA).

Editing

Kimi AI was used exclusively for language refinement. The authors reviewed and edited the output as needed and take full responsibility for the content of the publication.

Acknowledgments

The authors express gratitude to Yi-Qing Hui, Yi-Fan Ye, Chang-Lin Shen, and Yi-Ling Shi for their technical assistance in animal care, single-unit isolation and Matlab coding. They also thank Edvard Moser and May-Britt Moser for providing the Behavioral Neurology Toolbox and data from five animals, and Cheng Wang, Xin-Jian Li, Dun Mao, Hai-Bing Xu, and Yong-Gang Yao for their valuable suggestions on the manuscript.

Author Contributions

LL designed the experiments and analyses; YLD, SYH, XFG, JLL, CD, CJL, RZ, and XC performed the experiments; YLD, SYH, and LL conducted the analyses; YMS carried out the Bayesian decoding. YLD, XC, and LL wrote the manuscript. All authors contributed to discussion and interpretation of the results.

Funding

This research was supported by the Science and Technology Innovation (STI) 2030-Major Projects (2022ZD0205000 to LL), National Natural Science Foundation of China (31970963 to LL), Yunnan Revitalization Talent Support Program (Yunling Scholar Project to LL), Yunnan Province (202305AH340006 to LL), and Yunnan Fundamental Research Projects (202101AT070289 to XC).

Conflict of interest statement

The authors declare no competing financial interests.

Code accessibility

All custom-written codes used in the analyses are available upon request.

Data availability

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Supplementary material

Supplementary material is accessible in the online version of this paper.

References

1. Tulving, E., *Episodic memory: from mind to brain.* Annu Rev Psychol, 2002. **53**: p. 1-25.

2. Sugar, J. and M.B. Moser, *Episodic memory: Neuronal codes for what, where, and when.* Hippocampus, 2019. **29**(12): p. 1190-1205.

3. Colgin, L.L., E.I. Moser, and M.B. Moser, *Understanding memory through hippocampal remapping.* Trends Neurosci, 2008. **31**(9): p. 469-77.

4. Latuske, P., et al., *Hippocampal Remapping and Its Entorhinal Origin.* Front Behav

Neurosci, 2017. **11**: p. 253.

5.   Rivard, B., et al., *Representation of objects in space by two classes of hippocampal*
pyramidal cells. Journal of General Physiology, 2004. **124**(1): p. 9-25.

6.   Manns, J.R. and H. Eichenbaum, *A cognitive map for object memory in the hippocampus.*
Learn Mem, 2009. **16**(10): p. 616-24.

7.   Burke, S.N., et al., *The influence of objects on place field expression and size in distal*
hippocampal CA1. Hippocampus, 2011. **21**(7): p. 783-801.

8.   Deshmukh, S.S. and J.J. Knierim, *Influence of local objects on hippocampal*
representations: Landmark vectors and memory. Hippocampus, 2013. **23**(4): p. 253-67.

9.   Vandrey, B., S. Duncan, and J.A. Ainge, *Object and object-memory representations across*
the proximodistal axis of CA1. Hippocampus, 2021. **31**(8): p. 881-896.

10.   Nagelhus, A., et al., *Object-centered population coding in CA1 of the hippocampus.*
Neuron, 2023. **111**(13): p. 2091-2104 e14.

11.   Canto, C.B., F.G. Wouterlood, and M.P. Witter, *What does the anatomical organization of*
the entorhinal cortex tell us? Neural Plast, 2008. **2008**: p. 381243.

12.   Hafting, T., et al., *Microstructure of a spatial map in the entorhinal cortex.* Nature, 2005.
436(7052): p. 801-806.

13.   Ouchi, A. and S. Fujisawa, *Predictive grid coding in the medial entorhinal cortex.* Science,
2024. **385**(6710): p. 776-784.

14.   Fyhn, M., et al., *Spatial representation in the entorhinal cortex.* Science, 2004. **305**(5688):
p. 1258-64.

15.   Sargolini, F., et al., *Conjunctive representation of position, direction, and velocity in*
entorhinal cortex. Science, 2006. **312**(5774): p. 758-62.

16.   Solstad, T., et al., *Representation of geometric borders in the entorhinal cortex.* Science,
2008. **322**(5909): p. 1865-8.

17.   Hoydal, O.A., et al., *Object-vector coding in the medial entorhinal cortex.* Nature, 2019.
568(7752): p. 400-404.

18.   Kropff, E., et al., *Speed cells in the medial entorhinal cortex.* Nature, 2015. **523**(7561): p.
419-24.

19.   Tukker, J.J., et al., *Microcircuits for spatial coding in the medial entorhinal cortex.* Physiol

Rev, 2022. **102**(2): p. 653-688.

20. Long, J.L. and L. Lu, *Dynamic coding in the hippocampus during navigation.* Zool Res, 2022. **43**(6): p. 1023-1025.

21. Hargreaves, E.L., et al., *Major dissociation between medial and lateral entorhinal input to dorsal hippocampus.* Science, 2005. **308**(5729): p. 1792-4.

22. Huang, X., et al., *Distinct spatial maps and multiple object codes in the lateral entorhinal cortex.* Neuron, 2023. **111**(19): p. 3068-3083 e7.

23. Deshmukh, S.S. and J.J. Knierim, *Representation of non-spatial and spatial information in the lateral entorhinal cortex.* Front Behav Neurosci, 2011. **5**: p. 69.

24. Tsao, A., M.B. Moser, and E.I. Moser, *Traces of experience in the lateral entorhinal cortex.* Curr Biol, 2013. **23**(5): p. 399-405.

25. Igarashi, K.M., et al., *Coordination of entorhinal-hippocampal ensemble activity during associative learning.* Nature, 2014. **510**(7503): p. 143-7.

26. Leitner, F.C., et al., *Spatially segregated feedforward and feedback neurons support differential odor processing in the lateral entorhinal cortex.* Nat Neurosci, 2016. **19**(7): p. 935-44.

27. Tsao, A., et al., *Integrating time from experience in the lateral entorhinal cortex.* Nature, 2018. **561**(7721): p. 57-62.

28. Wang, C., et al., *Egocentric coding of external items in the lateral entorhinal cortex.* Science, 2018. **362**(6417): p. 945-949.

29. Cappaert, N.L., N.M. van Strien, and M.P. Witter, *Hippocampal Formation*, in *The Rat Nervous System*, G. Paxinos, Editor. 2015, Academic Press. p. 511-573.

30. Nilssen, E.S., et al., *Neurons and networks in the entorhinal cortex: A reappraisal of the lateral and medial entorhinal subdivisions mediating parallel cortical pathways.* Hippocampus, 2019. **29**(12): p. 1238-1254.

31. Nakamura, N.H., et al., *Proximodistal segregation of nonspatial information in CA3: preferential recruitment of a proximal CA3-distal CA1 network in nonspatial recognition memory.* J Neurosci, 2013. **33**(28): p. 11506-14.

32. Beer, Z., et al., *The memory for time and space differentially engages the proximal and distal parts of the hippocampal subfields CA1 and CA3.* PLoS Biol, 2018. **16**(8): p.

832 e2006100.

33. Flasbeck, V., et al., *Spatial information is preferentially processed by the distal part of CA3: Implication for memory retrieval.* Behav Brain Res, 2018. **354**: p. 31-38.

34. Ito, H.T. and E.M. Schuman, *Functional division of hippocampal area CA1 via modulatory gating of entorhinal cortical inputs.* Hippocampus, 2012. **22**(2): p. 372-87.

35. Nakazawa, Y., et al., *Memory retrieval along the proximodistal axis of CA1.* Hippocampus, 2016. **26**(9): p. 1140-8.

36. Hartzell, A.L., et al., *Transcription of the immediate-early gene Arc in CA1 of the hippocampus reveals activity differences along the proximodistal axis that are attenuated by advanced age.* J Neurosci, 2013. **33**(8): p. 3424-33.

37. Henriksen, E.J., et al., *Spatial representation along the proximodistal axis of CA1.* Neuron, 2010. **68**(1): p. 127-37.

38. Oliva, A., et al., *Spatial coding and physiological properties of hippocampal neurons in the Cornu Ammonis subregions.* Hippocampus, 2016. **26**(12): p. 1593-1607.

39. Deshmukh, S.S., *Distal CA1 Maintains a More Coherent Spatial Representation than Proximal CA1 When Local and Global Cues Conflict.* J Neurosci, 2021. **41**(47): p. 9767-9781.

40. Li, Y., et al., *A distinct entorhinal cortex to hippocampal CA1 direct circuit for olfactory associative learning.* Nat Neurosci, 2017. **20**(4): p. 559-570.

41. Leutgeb, S., et al., *Independent codes for spatial and episodic memory in hippocampal neuronal ensembles.* Science, 2005. **309**(5734): p. 619-23.

42. Lu, L., et al., *Impaired hippocampal rate coding after lesions of the lateral entorhinal cortex.* Nat Neurosci, 2013. **16**(8): p. 1085-93.

43. Lu, L., et al., *Topography of place maps along the CA3-to-CA2 axis of the hippocampus.* Neuron, 2015. **87**(5): p. 1078-92.

44. Burke, S.N., et al., *Representation of three-dimensional objects by the rat perirhinal cortex.* Hippocampus, 2012. **22**(10): p. 2032-44.

45. Deshmukh, S.S., J.L. Johnson, and J.J. Knierim, *Perirhinal cortex represents nonspatial, but not spatial, information in rats foraging in the presence of objects: comparison with lateral entorhinal cortex.* Hippocampus, 2012. **22**(10): p. 2045-58.

46. Aggleton, J.P. and A.J.D. Nelson, *Distributed interactive brain circuits for object-in-place*
memory: A place for time? Brain Neurosci Adv, 2020. **4**: p. 2398212820933471.

47. Parron, C. and E. Save, *Comparison of the effects of entorhinal and retrosplenial cortical*
lesions on habituation, reaction to spatial and non-spatial changes during object
exploration in the rat. Neurobiology of Learning and Memory, 2004. **82**(1): p. 1-11.

48. Rodo, C., F. Sargolini, and E. Save, *Processing of spatial and non-spatial information in*
rats with lesions of the medial and lateral entorhinal cortex: Environmental complexity
matters. Behavioural Brain Research, 2017. **320**: p. 200-209.

49. Van Cauter, T., et al., *Distinct roles of medial and lateral entorhinal cortex in spatial*
cognition. Cereb Cortex, 2013. **23**(2): p. 451-9.

50. Tennant, S.A., et al., *Stellate Cells in the Medial Entorhinal Cortex Are Required for*
Spatial Learning. Cell Reports, 2018. **22**(5): p. 1313-1324.

51. Wilson, D.I., et al., *Lateral entorhinal cortex is critical for novel object-context*
recognition. Hippocampus, 2013. **23**(5): p. 352-66.

52. Wilson, D.I., et al., *Lateral entorhinal cortex is necessary for associative but not*
nonassociative recognition memory. Hippocampus, 2013. **23**(12): p. 1280-90.

53. Ocko, S.A., et al., *Emergent elasticity in the neural code for space.* Proc Natl Acad Sci U
S A, 2018. **115**(50): p. E11798-e11806.

54. Wen, J.H., et al., *One-shot entorhinal maps enable flexible navigation in novel*
environments. Nature, 2024.

55. Sun, Y., et al., *CA1-projecting subiculum neurons facilitate object-place learning.* Nat
Neurosci, 2019. **22**(11): p. 1857-1870.

56. Sun, Y., et al., *Subicular neurons encode concave and convex geometries.* Nature, 2024.
627(8005): p. 821-829.

57. Renno-Costa, C., J.E. Lisman, and P.F. Verschure, *The mechanism of rate remapping in*
the dentate gyrus. Neuron, 2010. **68**(6): p. 1051-8.

58. Diehl, G.W., et al., *Grid and Nongrid Cells in Medial Entorhinal Cortex Represent Spatial*
Location and Environmental Features with Complementary Coding Schemes. Neuron,
2017. **94**(1): p. 83-92 e6.

59. Ismakov, R., et al., *Grid Cells Encode Local Positional Information.* Curr Biol, 2017.

27(15): p. 2337-2343 e3.

60. Marozzi, E., et al., *Purely Translational Realignment in Grid Cell Firing Patterns*
Following Nonmetric Context Change. Cereb Cortex, 2015. **25**(11): p. 4619-27.

61. Miao, C., et al., *Hippocampal Remapping after Partial Inactivation of the Medial*
Entorhinal Cortex. Neuron, 2015. **88**(3): p. 590-603.

62. Rueckemann, J.W., et al., *Transient optogenetic inactivation of the medial entorhinal*
cortex biases the active population of hippocampal neurons. Hippocampus, 2016. **26**(2):
p. 246-60.

63. Kanter, B.R., et al., *A Novel Mechanism for the Grid-to-Place Cell Transformation*
Revealed by Transgenic Depolarization of Medial Entorhinal Cortex Layer II. Neuron,
2017. **93**(6): p. 1480-1492 e6.

64. Fyhn, M., et al., *Hippocampal remapping and grid realignment in entorhinal cortex.*
Nature, 2007. **446**(7132): p. 190-4.

65. Leutgeb, S., et al., *Distinct ensemble codes in hippocampal areas CA3 and CA1.* Science,
2004. **305**(5688): p. 1295-1298.

66. Knierim, J.J., J.P. Neunuebel, and S.S. Deshmukh, *Functional correlates of the lateral and*
medial entorhinal cortex: objects, path integration and local-global reference frames.
Philos Trans R Soc Lond B Biol Sci, 2014. **369**(1635): p. 20130369.

67. Soltesz, I. and A. Losonczy, *CA1 pyramidal cell diversity enabling parallel information*
processing in the hippocampus. Nat Neurosci, 2018. **21**(4): p. 484-493.

68. Lu, L., et al., *Construction of an Improved Multi-Tetrode Hyperdrive for Large-Scale*
Neural Recording in Behaving Rats. J Vis Exp, 2018(135).

69. Cacucci, F., et al., *Theta-modulated place-by-direction cells in the hippocampal formation*
in the rat. J Neurosci, 2004. **24**(38): p. 8265-77.

70. Skaggs, W.E., et al., *Theta phase precession in hippocampal neuronal populations and the*
compression of temporal sequences. Hippocampus, 1996. **6**(2): p. 149-72.

71. Glaser, J.I., et al., *Machine Learning for Neural Decoding.* eNeuro, 2020. **7**(4).

Figures

Figure 1

Figure 1. Object-placement induced complex population response in CA1

A. Representative firing rate distribution maps (rate maps) of neurons recorded from proximal, middle, and distal bands of CA1, showing responses to familiar objects placed in the recording box. Maps for the same cell across four trials are arranged in rows, with animal and cell IDs indicated on the left. Color is scaled to maximum peak rate of four trials. Text above each map indicates peak rate of that trial. Circles represent objects. Obj−, trial without objects; Obj+, trial with two familiar objects.

B. Distribution of cells in response to object placement across three CA1 bands. Stable cells had rate changes less than 50%. Moderately remapped cells had 0.5–3-fold increase in rate. Strongly remapped cells had rate changes higher than 3 folds. Numbers indicate active place cells in each category. Proportion of cells with firing rate changes: pCA1, 57.94%, mCA1, 50.41%, and dCA1, 60.58%; chi-square test, $\chi^2 = 15.640$, $P < 0.001$; Holm-Bonferroni *post hoc* tests, ***: $P < 0.001$.

C. Heatmaps showing PVC of all putative pyramidal neurons recorded from each CA1 subregion between trials with or without objects (top two rows), as well as changes in PVC

between condition pairs (bottom row). Heatmap bins correspond to spatial bins in rate maps.

Color scales are indicated on the right of each row. Small circles represent objects, large circles

indicate object region.

D. Degree of changes in PVC as a function of distance to the objects (mean $\pm$ SEM) in each

CA1 subregion. Two-way ANOVA, distance: $F_{(18, 1\,143)} = 58.526$, $P < 0.001$; band: $F_{(2, 1}$

$_{143)} = 37.619$, $P < 0.001$; distance $\times$ band: $F_{(36, 1\,143)} = 5.278$, $P < 0.001$.

E. Boxplots showing degree of changes in PVC between trials pairs, for spatial bins inside and

outside the object region, with outliers omitted. Bins inside object region: median (25%–75%

percentile), pCA1, 0.199 (0.171–0.228), mCA1, 0.215 (0.174–0.285), and dCA1, 0.185 (0.154–

0.239). Bins outside box region: pCA1, 0.096 (0.065–0.150), mCA1, 0.063 (0.026–0.107),

dCA1, 0.075 (0.042–0.130); Kruskal-Wallis test, H = 66.740, $P < 0.001$, 960 bins; Holm-

Bonferroni *post hoc* tests, **: $P < 0.01$; ***: $P < 0.001$.

Figure 2

A.

R0388
Trial 1 | Trial 2 | Trial 3 | Trial 4
B | W | W | B
Path

R0175
Trial 1 | Trial 2 | Trial 3 | Trial 4
B | W | W | B
Path

pCA1
T09C1 — 1.13 Hz, 10.98 Hz, 10.52 Hz, 5.70 Hz
T09C2 — 0.95 Hz, 11.05 Hz, 12.17 Hz, 8.03 Hz
T09C3 — 3.03 Hz, 4.96 Hz, 2.80 Hz, 1.44 Hz
T10C1 — 4.62 Hz, 5.89 Hz, 5.27 Hz, 6.70 Hz
T10C2 — 10.34 Hz, 12.94 Hz, 14.00 Hz, 16.09 Hz

mCA1
T08C1 — 10.81 Hz, 12.11 Hz, 11.61 Hz, 10.83 Hz
T08C2 — 11.44 Hz, 9.69 Hz, 7.21 Hz, 14.15 Hz
T16C1 — 11.17 Hz, 6.67 Hz, 7.29 Hz, 7.10 Hz
T16C2 — 6.91 Hz, 7.12 Hz, 6.29 Hz, 8.58 Hz
T16C3 — 9.56 Hz, 6.53 Hz, 5.34 Hz, 5.08 Hz

mCA1
T01C1 — 15.77 Hz, 13.18 Hz, 10.18 Hz, 10.62 Hz
T01C2 — 6.05 Hz, 5.12 Hz, 7.39 Hz, 7.15 Hz
T12C1 — 5.07 Hz, 6.85 Hz, 6.82 Hz, 6.40 Hz
T12C2 — 6.57 Hz, 13.37 Hz, 10.40 Hz, 9.45 Hz
T13C1 — 2.31 Hz, 3.18 Hz, 4.33 Hz, 2.65 Hz

dCA1
T03C1 — 7.28 Hz, 7.21 Hz, 8.35 Hz, 5.70 Hz
T03C2 — 0.10 Hz, 8.09 Hz, 9.91 Hz, 12.89 Hz
T04C1 — 3.43 Hz, 7.22 Hz, 7.71 Hz, 8.24 Hz
T15C1 — 4.05 Hz, 7.36 Hz, 4.88 Hz, 8.48 Hz
T15C2 — 5.30 Hz, 6.32 Hz, 5.79 Hz, 5.17 Hz

0 Hz — Max

B. pCA1 mCA1 dCA1 — Same, B vs W, ΔPVC

C. Color-reversal induced ΔPVC — Δ PV correlation — pCA1, mCA1, dCA1 — *** *** ***

E. Spatial correlation — Same, B vs W — pCA1, mCA1, dCA1 — Δ spatial correlation — *** * ***

F. Rate difference — Same, B vs W — pCA1, mCA1, dCA1 — Δ rate difference — ***

D. Strong, Moderate, Stable — Frequency
pCA1: 217, 389, 297
mCA1: 122, 424, 424
dCA1: 140, 328, 281
* *** **

G. Spatial correlation — Δ spatial correlation — Normalized recording position — pro. dis. — R² = 0.162, R² = 0.141 (p), R² = 0.140 (d)

H. Rate difference — Δ rate difference — Normalized recording position — pro. dis. — R² = 0.308, R² = 0.204 (p), R² = 0.096 (d)

Figure 2. V-shaped population response to reversed wall color across CA1 subdivisions

A. Representative rate maps from simultaneously recorded CA1 neurons in two separate color-reversal experiments. Left: pCA1 and mCA1 place cells recorded from rat R0388. Right: mCA1 and dCA1 place cells recorded from rat R0175. Gray lines indicate trajectories of animal in each trial. B, black; W, white version of the same box. Note, mCA1 showed minimum response to the color change.

B. Heatmaps showing PVC of all putative pyramidal neurons recorded from each CA1 subregion between color conditions (top two rows), and changes in PVC between condition pairs (bottom row) in the color-reversal task. Note, mCA1 exhibited the lowest activity changes. Reduction in PVC is stronger for spatial bins away from arena walls, where the color was

manipulated.

C. Degree of changes in PVC between condition pairs, with outliers omitted. Place cells in

mCA1 exhibited least activity change. pCA1: 0.108 (0.085–0.147), mCA1: 0.031 (0.005–

0.054), and dCA1: 0.052 (0.022–0.094); Kruskal-Wallis test, H = 541.211, $P < 0.001$, 1 200

bins, Holm-Bonferroni *post hoc* tests, ***: $P < 0.001$.

D. Distribution of rate remapped neurons between black and white color conditions across three

CA1 bands. Stable cells had rate changes less than 50%. Moderately remapped cells had 0.5–

3-fold increase in rate. Strongly remapped cells had rate changes higher than 3 folds. Numbers

indicate active place cells in each category. Proportion of strongly remapped cells: pCA1,

24.03%, mCA1, 12.58%, and dCA1, 18.69%, chi-square test, $\chi^2 = 41.214$, $P < 0.001$; Holm-

Bonferroni *post hoc* tests, *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$.

E. Left: distributions of SC in CA1 subregions between matching (Same) and color reversed

(B vs W) conditions. Horizontal line indicates median, and thick vertical bar indicates

interquartile range (IQR, 25%–75% percentile) of each subregion. Two-way repeated-measures

ANOVA, color: F (1, 2 619) = 262.113, $P < 0.001$; band: F (2, 2 619) = 26.679, $P < 0.001$; color

× band: F (2, 2 619) = 28.724, $P < 0.001$. Right: changes in SC between condition pairs, with

outliers omitted. pCA1: 0.032 (−0.003–0.106), mCA1: 0.010 (−0.020–0.049), and dCA1: 0.025

(−0.011–0.091); Kruskal-Wallis test, 2 622 cells, H = 68.492, $P < 0.001$; Holm-Bonferroni *post*

hoc tests, *: $P < 0.05$; ***: $P < 0.001$.

F. Similar to **E** but showing changes in RD across CA1 subdivisions. Left: color: F (1, 2 619)

= 288.514, $P < 0.001$; band: F (2, 2 619) = 20.373, $P < 0.001$; color × band: F (2, 2 619) =

41.298, $P < 0.001$. Right: pCA1: 0.012 (−0.005–0.140), mCA1: 0.001 (−0.006–0.047), and

dCA1: 0.001 (−0.010–0.048); H = 38.540, $P < 0.001$; *post hoc* tests, ***: $P < 0.001$.

G. Changes in SC between condition pairs as a function of tetrode position. Each cross

corresponds to average data of a single tetrode. Blue curve, linear regression; red curve,

quadratic polynomial regression. Explained variances (R^2) are indicated with the same color

code. pro., proximal end; dis., distal end. p, proximal half; d, distal half. Proximal half: r =

−0.375, $P = 0.002$, Pearson correlation, 63 tetrodes; distal half: r = 0.374, $P = 0.007$, 50 tetrodes.

H. Similar to **G** but showing correlations between tetrode position and changes in RD between

condition pairs. Proximal half: r = −0.452, $P < 0.001$; distal half: r = 0.314, $P = 0.029$.

Figure 3

Figure 3. Proximal CA1 was more sensitive to changes in landmarks

A. Simultaneously recorded CA1 neurons in two rats exhibiting global remapping in the two-room task. Left: pCA1 and mCA1 place cells recorded from rat R0172. A, Room A; B, Room B. Right: mCA1 and dCA1 place cells recorded from rat R0175. Note, redistribution of firing locations between different rooms.

B. Heatmaps showing PVCs of all putative pyramidal neurons recorded from each subregion in the same room (top row) and between rooms (middle row), and changes in PVC between room pairs (bottom row).

C. Degree of changes in PVC between condition pairs, with outliers omitted. pCA1: 0.678 (0.609–0.731), mCA1: 0.452 (0.367–0.535), and dCA1: 0.388 (0.334–0.441); Kruskal-Wallis test, H = 679.871, $P < 0.001$, 1 200 bins, Holm-Bonferroni *post hoc* tests, ***: $P < 0.001$.

D. Left: distribution of SC in CA1 subdivisions in the same room, and in different rooms (Room A vs B). Two-way repeated-measures ANOVA, room: F (1, 758) = 1 175.415, $P < 0.001$; band: F (2, 758) = 13.538, $P < 0.001$; room × band: F (2, 758) = 27.266, $P < 0.001$. Right: change in SC between condition pairs, with outliers omitted. pCA1, 0.455 (0.256–0.747), mCA1, 0.354 (0.151–0.527), and dCA1, 0.287 (0.095–0.504); Kruskal-Wallis test, 761 cells, H = 41.735, $P < 0.001$; Holm-Bonferroni *post hoc* tests, ***: $P < 0.001$.

E. Pearson correlation between tetrode position and changes in SC across condition pairs (54 tetrodes). Each cross corresponds to average data of a single tetrode. Blue curve, linear regression; red curve, quadratic polynomial regression.

Figure 4

Figure 4. Distal CA1 preferentially represented changes in object-identity

A. Representative neurons in proximal, middle, and distal bands of CA1, showing remapping

to the novel object. Circle and star represent familiar and novel objects, respectively. Fam, trial

with two familiar objects; Nov, trial with right object replaced with a novel object.

B. Distribution of neurons in response to the novel object across three CA1 bands. Proportion of cells responsive to object-replacement: pCA1, 37.84%, mCA1, 41.15%, and dCA1, 53.02%; chi-square test, $\chi^2 = 24.602$, $P < 0.001$; Holm-Bonferroni *post hoc* tests, ***: $P < 0.001$.

C. Left: distributions of RD in CA1 subregions between trials with a novel object (Same) and trials before and after object replacement (Obj F vs N). Two-way repeated-measures ANOVA, novelty F (1, 1 448) = 58.170, $P < 0.001$; band: F (2, 1 448) = 10.527, $P < 0.001$; novelty × band: F (2, 1 448) = 6.169, $P = 0.002$. Right: changes in RD between condition pairs, with outliers omitted. pCA1, 0.008 (−0.101–0.120), mCA1, 0.036 (−0.064–0.162), and dCA1, 0.054 (−0.085–0.225); Kruskal-Wallis test, 1 451 cells, H = 13.604, $P = 0.001$; Holm-Bonferroni *post hoc* tests, *: $P < 0.05$; ***: $P < 0.001$.

D. Pearson correlation between tetrode position and changes in RD between condition pairs (67 tetrodes). Each cross corresponds to average data of a single tetrode. Blue curve, linear regression; red curve, quadratic polynomial regression.

E. Heatmaps showing PVCs of all putative pyramidal neurons recorded from each subregion between trials (top two rows), as well as changes in PVC between trial pairs (bottom row) in the object-replacement task. Small circle and star represent familiar and novel objects, respectively; large circle indicates region of the novel object.

F. Degree of changes in overall PVC between trial pairs across CA1 bands, with outliers omitted. pCA1, 0.031 (0.006–0.065), mCA1, 0.040 (0.023–0.061), and dCA1, 0.047 (0.014–0.079); Kruskal-Wallis test, H = 19.995, $P < 0.001$, 1,200 bins; Holm-Bonferroni *post hoc* tests, **: $P \leq 0.01$; ***: $P \leq 0.001$.

G. Degree of changes in PVC inside and outside ROI between trial pairs across CA1 bands, with outliers omitted. Bins inside ROI: pCA1, 0.005 (−0.014–0.028), mCA1, 0.034 (0.021–0.046), and dCA1, 0.070 (0.053–0.082); Kruskal-Wallis test, H = 61.612, $P < 0.001$, 120 bins. Bins outside ROI: pCA1, 0.035 (0.009–0.069), mCA1, 0.041 (0.023–0.064), and dCA1, 0.044 (0.012–0.078); H = 5.689, $P = 0.058$, 1 080 bins. Holm-Bonferroni *post hoc* tests, n.s. not significant; **: $P \leq 0.01$; ***: $P \leq 0.001$.

Figure 5

A.

pCA1 · mCA1 · dCA1 panels

B. Proportion of cells

C. Spatial correlation

D.

E. Rate difference

F.

G. pCA1 mCA1 dCA1

H. Overall ΔPVC

I.

1044

Figure 5. Object-translocation induced stronger population response in dCA1

A. Representative neurons in proximal, middle, and distal bands of CA1, showing responses to

object displacement. Circles represent objects. Fam, objects in familiar locations; Reloc, trial

with right object relocated.

B. Distribution of neurons in response to object translocation across three CA1 bands.

Proportion of cells that showed field changes to the translocation: pCA1, 29.52%, mCA1,

36.65%, and dCA1, 54.42%; chi-square test, $\chi^2 = 61.433$, $P < 0.001$; Holm-Bonferroni *post hoc*

tests, ***: $P < 0.001$. Subgroups: OVC like neurons, Fisher's exact test, $P = 0.338$; other

relocation responsive cells, chi-square test, all $P < 0.05$.

C. Left: distribution of SC in CA1 subregions between trials with relocated objects (Same obj

location) and trials before and after object relocation (Obj location A vs B). Two-way repeated-

measures ANOVA, relocation: F (1, 1 483) = 153.986, $P < 0.001$; band: F (2, 1 483) = 21.400,

$P < 0.001$; relocation × band: F (2, 1 483) = 4.917, $P = 0.007$. Right: degree of changes in SC

between condition pairs, with outliers omitted. pCA1, 0.024 (−0.026–0.121), mCA1, 0.049

(−0.021–0.175), and dCA1, 0.069 (−0.027–0.233); Kruskal-Wallis test, 1 486 cells, H = 10.123,

$P = 0.006$; Holm-Bonferroni *post hoc* tests, **: $P < 0.01$.

D. Pearson correlation between recording position and changes in SC between condition pairs

(69 tetrodes). Each cross corresponds to average data of a single tetrode. Blue curve, linear

regression; red curve, quadratic polynomial regression.

E. Similar to **C** but showing changes in RD across CA1 subdivisions. Left: relocation: F (1,

1 483) = 147.858, $P < 0.001$; band: F (2, 1 483) = 20.702, $P < 0.001$; relocation × band: F (2,

1 483) = 2.609, $P = 0.074$. Right: pCA1, 0.044 (−0.073–0.205), mCA1, 0.057 (−0.057–0.221),

and dCA1, 0.112 (−0.056–0.308); H = 9.346, $P = 0.009$; *post hoc* tests, *: $P < 0.05$.

F. Similar to **D** but showing correlations between tetrode position and changes in RD between

condition pairs.

G. Heatmaps showing PVCs of all putative pyramidal neurons recorded from each subregion

between trials (top two rows), as well as changes in PVC between trial pairs (bottom row) in

object-relocation task. Small circles represent objects, large circles indicate target region.

H. Degree of changes in overall PVC between trial pairs across CA1 bands, with outliers

omitted. pCA1, 0.055 (0.014–0.111), mCA1, 0.068 (0.030–0.127), dCA1, 0.077 (0.043–0.138);

Kruskal-Wallis test, H = 44.493, $P < 0.001$, 1 200 bins; Holm-Bonferroni *post hoc* tests, ***: P

≤ 0.001.

I. Degree of changes in PVC inside and outside ROI between trial pairs across CA1 bands, with

outliers omitted. Bins inside ROI: pCA1, 0.081 (0.054–0.114), mCA1, 0.212 (0.147–0.269),

and dCA1, 0.225 (0.155–0.303); Kruskal-Wallis test, H = 125.736, $P < 0.001$, 240 bins. Bins

outside ROI: pCA1, 0.043 (0.009–0.106), mCA1, 0.049 (0.026–0.091), and dCA1, 0.063

(0.038–0.101); H = 17.077, $P < 0.001$, 960 bins. Holm-Bonferroni *post hoc* tests, *: $P \leq 0.05$;

***: $P \leq 0.001$.

Figure 6

1083

Figure 6. Distinct contributions of neuron subpopulations during event coding

A. Heatmaps showing PVCs of place cells excluding OVC-like neurons recorded from each subregion between trials (top two rows), as well as changes in PVC between trial pairs (bottom row) in object-relocation task. Small circles represent objects, large circles indicate target region.

B. Degree of changes in PVC within ROI between trial pairs across CA1 bands, with outliers omitted. pCA1, 0.075 (0.043–0.110), mCA1, 0.127 (0.081–0.187), dCA1, 0.244 (0.183–0.308); Kruskal-Wallis test, H = 128.394, $P < 0.001$, 240 bins; Holm-Bonferroni *post hoc* tests, ***: $P < 0.001$.

C. Degree of changes in PVC at original and current locations between trial pairs across CA1 bands, with outliers omitted. Original location: pCA1, 0.089 (0.058–0.125), mCA1, 0.130 (0.091–0.201), and dCA1, 0.285 (0.229–0.319); Kruskal-Wallis test, H = 71.317, $P < 0.001$, 120 bins. Current location: pCA1, 0.069 (0.022–0.090), mCA1, 0.108 (0.069–0.170), and dCA1, 0.198 (0.147–0.291); H = 64.233, $P < 0.001$. Holm-Bonferroni *post hoc* tests, **: $P < 0.01$; ***:

$P \leq 0.001$.

D. Heatmaps showing changes in PVC for population of place cells without OLC neurons

between trial pairs, as well as reductions in PVC change (compared to Fig. 6A bottom panel)

after exclusion for CA1 subregions. Circles indicate ROI.

E. Reductions in PVC at original and current locations after exclusion of OLC neurons across

CA1 bands, with outliers omitted. Original location: pCA1, 0.077 (0.038–0.141), mCA1, 0.080

(0.052–0.105), and dCA1, 0.045 (0.020–0.077); Kruskal-Wallis test, H = 15.605, $P < 0.001$,

120 bins. Current location: pCA1, 0.008 (0.003–0.021), mCA1, −0.002 (−0.010–0.008), and

dCA1, −0.001 (−0.008–0.002). Holm-Bonferroni *post hoc* tests, **: $P < 0.01$; ***: $P < 0.001$.

F. Normalized PVC change after OLC neuron exclusion at original location across CA1 bands,

with outliers omitted. pCA1, 0.901 (0.417–1.000), mCA1, 0.547 (0.416–0.622), and dCA1,

0.163 (0.077–0.245); Kruskal-Wallis test, H = 71.904, $P < 0.001$, 120 bins. Holm-Bonferroni

post hoc tests, ***: $P < 0.001$.

G–I. Similar to **D**–**F**, but show changes in PVC for population of place cells without CLC

neurons (**G**), reductions in PVC (**H**. OL: pCA1, −0.0003 (−0.0007–0.0002), mCA1, 0.001

(−0.003–0.005), and dCA1, 0.011 (−0.001–0.025). CL: pCA1, 0.021 (0.010–0.031), mCA1,

0.034 (0.023–0.050), and dCA1, 0.056 (0.033–0.092); Kruskal-Wallis test, H = 34.125, $P <$

0.001, 120 bins), and normalized PVC change at current location (**I**. pCA1, 0.290 (0.183–0.404),

mCA1, 0.389 (0.178–0.523), and dCA1, 0.270 (0.107–0.630); H = 0.658, $P = 0.720$). Holm-

Bonferroni *post hoc* tests, n.s.: not significant; *: $P < 0.05$; **: $P < 0.01$, ***: $P < 0.001$.