1 ZNHIT1-dependent H2A.Z deposition at meiotic prophase I

2 underlies pachytene gene expression and meiotic progression during

3 male meiosis

5 Shenfei Sun[1,2,5,*], Yamei Jiang[2,5], Ning Jiang[2,5], Qiaoli Zhang[2], Hongjie Pan[3], Fujing

6 Huang[2], Xinna Zhang[1], Yuxuan Guo[1], Xiaoyu You[2], Kai Gong[2], Wei Wei[2], Hanmin Liu[4],

7 Zhenju Song[2], Yuanlin Song[2], Xiaofang Tang[2], Miao Yu[2], Runsheng Li[3], Xinhua Lin[1,2,4*]

9 Affiliations

[1]State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock,

Institutes of Biomedical Sciences, School of Life Sciences, Inner Mongolia University,

Hohhot 010070, China;

[2]State Key Laboratory of Genetic Engineering, Greater Bay Area Institute of Precision

Medicine (Guangzhou), School of Life Sciences, Zhongshan Hospital, Fudan University,

Shanghai 200438, China;

[3]National Health Commission (NHC) Key Laboratory of Reproduction Regulation,

Shanghai Institute for Biomedical and Pharmaceutical Technologies, Shanghai 200032,

China;

[4]The Joint Laboratory for Lung Development and Related Diseases of West China Second

University Hospital, Sichuan University and School of Life Sciences of Fudan University,

Chengdu 610041, China;

[5]These authors contributed equally to this work.

[*]Correspondence to: xlin@fudan.edu.cn (X.L.); shenfei_sun@fudan.edu.cn (S.S.).

Keywords: male meiosis; pachytene gene expression; meiotic progression; ZNHIT1;
H2A.Z

Abstract

Accurate meiotic progression is important for gamete formation and the generation of
genetic diversity. However, little is known about the identity of chromatin regulators that
underlie mammalian meiosis in vivo. Here, we identify the multifaceted functions of the
chromatin remodeler ZNHIT1 in governing meiosis. The expression of *Znhit1* gradually
increases during the meiotic prophase, and *Znhit1* knockout in spermatocytes results in
arrested pachytene development, impaired DNA double-strand break repair, and defective
homologous recombination. Single-cell RNA sequencing and transcriptome analysis
reveal that *Znhit1* loss dysregulates meiotic transcriptional programs at the pachytene
stage. Chromatin immunoprecipitation data show that ZNHIT1 is needed for the
incorporation of the histone variant H2A.Z into pachytene chromatin. Moreover, we
found that H2A.Z cooperates with the transcription factor A-MYB to co-bind DNA
elements and control gene activity. Our findings provide insights into the regulatory
mechanisms governing meiotic progression and highlight ZNHIT1 as a critical regulator
of meiotic progression.
44

Introduction

Meiosis is a fundamental and conserved process that plays a crucial role in gamete formation and the generation of genetic diversity. By undergoing one round of DNA replication and two rounds of cell division, the meiotic process ensures the production of haploid cells, each with a unique combination of genetic materials (Handel and Schimenti, 2010; Lascarez-Lagunas et al., 2020). Any disruptions to the normal progression of meiosis would have significant consequences, such as aneuploidy, infertility, spontaneous abortion, and congenital diseases (Hassold and Hunt, 2001). Therefore, understanding the molecular mechanisms underlying meiosis will provide valuable insights for the diagnosis and treatment of reproductive and developmental diseases.

Multiple chromosomal events occur during meiotic prophase I, including homologous chromosome pairing and synapsis, DNA double-strand break (DSB) formation and repair, and homologous recombination (HR) (Baudat et al., 2013; Hunter, 2015; Zickler and Kleckner, 2023). In this process, homolog pairing and synapsis coincide with SPO11-mediated genome-wide formation of DSBs (Bergerat et al., 1997; Keeney et al., 1997). Following the formation of DSBs, the DNA undergoes resection, leading to the generation of single-stranded DNA overhangs. These overhangs are then coated by RPA, DMC1, and RAD51, facilitating the production of recombination intermediates. Subsequently, these intermediates are processed and resolved, ultimately forming either meiotic crossovers or non-crossovers (Gray and Cohen, 2016; San Filippo et al., 2008; Symington, 2014). The tightly coordinated timing and spatial arrangement of these meiotic events are of utmost importance for proper germ cell development.

There is transcriptional awakening during male meiotic prophase I, in which the meiotic genome becomes transcriptionally active during the zygotene-to-pachytene transition (Alexander et al., 2023; Ernst et al., 2019; Green et al., 2018; Rabbani et al., 2022; Turner, 2015). This process, referred to as pachytene genome activation (PGA) in this context, is responsible for the expression of numerous genes and plays an essential role in controlling meiotic and post-meiotic events. Previous studies have emphasized the significance of the transcription factors (TFs) A-MYB, BRDT, and TCFL5 in activating transcription during pachytene (Alexander et al., 2023; Bolcun-Filas et al., 2011; Cecchini et al., 2023; Gaucher et al., 2012; Li et al., 2013; Maezawa et al., 2020; Manterola et al., 2018; Ozata et al., 2020; Yu et al., 2021). Additionally, extensive chromatin remodeling takes place in spermatocytes during meiotic prophase I, involving histone variant exchange, histone modifications, and high-order genome rearrangement (Kota and Feil,

2010; Wang et al., 2017; Zheng and Xie, 2019). It has been known that

SETDB1-mediated H3K9 trimethylation (H3K9me3) is required for sex chromosome

transcription silencing (a process called meiotic sex chromosome inactivation, MSCI) and

male meiotic procession (Hirota et al., 2018). However, the specific involvement of

chromatin regulators in gene activation on autosomes remains poorly understood.

Zinc finger HIT-type containing 1 (ZNHIT1), an evolutionarily conserved subunit of the

SRCAP chromatin remodeling complex, acts as a key regulator for the histone variant

H2A.Z deposition to control gene expression and cell fate determination (Cai et al., 2005;

Cuadrado et al., 2007; Feng et al., 2022). Recent studies have shown that ZNHIT1 is

involved in a wide range of developmental processes, including muscle and lens

differentiation, heart development, lung branching, and adult tissue stem cell maintenance

(Cuadrado et al., 2010; Lu et al., 2022; Sun et al., 2020; Wei et al., 2022; Xu et al., 2021;

Zhao et al., 2019). In our previous study, *Znhit1* deletion in early male germ cells

impaired the mitosis-to-meiosis transition by regulating *Meiosin* expression (Sun et al.,

2022). However, the in vivo function of ZNHIT1 in regulating meiotic progression

remains unknown.

To delve into the function of ZNHIT1 in meiosis, we analyzed *Znhit1* expression and

observed upregulation specifically during meiotic prophase. Here, we examine the role of

ZNHIT1 in multiple meiotic events and show that *Znhit1* deletion in spermatocytes

causes defects of pachytene gene expression, thereby resulting in meiotic developmental

arrest. Hence, our study highlights the essential role of ZNHIT1 in ensuring meiotic

progression.

Results

Znhit1 expression is upregulated during meiotic prophase and *Znhit1* knockout in spermatocytes disrupts spermatogenesis

To identify potential chromatin regulators in meiosis, we first queried for chromatin factors highly expressed during the zygotene stage based on published scRNA-seq data (Fig S1A and Table S1) (Chen et al., 2018). Quantitative analysis showed that the chromatin remodeler *Znhit1* was among the most highly expressed factors (Fig 1A, B, S1B, C). RNA in situ hybridization validated that *Znhit1* expression in male germ cells decreased after meiotic initiation, followed by a gradual increase in spermatocytes from the zygotene stage to the metaphase stage (Fig 1C). These results suggest that ZNHIT1 is a potential chromatin regulator of meiotic progression.

To investigate whether ZNHIT1 regulates meiotic progression, we generated spermatocyte-specific *Znhit1* knockout mice (*Znhit1*^{fl/fl}; *Stra8-cre*, referred to as *Znhit1*-sKO). In the first wave of spermatogenesis, Ngn3-negative prospermatogonia directly differentiate into A2 spermatogonia(Law et al., 2019; Rabbani et al., 2022; Yoshida et al., 2006; Yoshida et al., 2004). With Stra8-cre inducing recombination in A1 spermatogonia and preleptotene spermatocytes separately(Lin et al., 2017), it is feasible to generate mice with spermatocyte-specific gene knockout (Fig 1D). RNA in situ experiments on testis sections confirmed the deletion of *Znhit1* mRNA in spermatocytes (Fig 1E, S1D). Compared with littermate controls, *Znhit1*-sKO male mice had smaller testes (Fig S1E, F). PAS-histological and PNA-fluorescent staining illuminated the absence of round spermatids and elongated spermatids in *Znhit1*-sKO testes (Fig 1F, G), indicating that *Znhit1* knockout in spermatocytes disrupted spermatogenesis.

Znhit1 deletion leads to meiotic arrest at the pachytene stage

To study spermatocyte development in *Znhit1*-sKO testes, we co-stained SYCP3, a marker of primary spermatocytes, and HSPA2, a testis-specific HSP70 family member that accumulates from the pachytene stage onward. Both control and *Znhit1*-sKO testis sections exhibited SYCP3$^+$HSPA2$^+$ primary spermatocytes (Fig S2A). Phospho-histone H3 (Thr3) (pH3) is a marker of mitotic and meiotic metaphase cells, and *Znhit1* deletion resulted in the complete absence of pH3$^+$ metaphase spermatocytes within the seminiferous tubule lumen (Fig S2B). These results suggest that *Znhit1* deletion arrests spermatocytes in meiotic prophase I.

To pinpoint the onset of meiotic failure, we performed immunostaining against SYCP3 in

spermatocyte chromosome spreads. The wild-type testes displayed typical spermatocytes

at consecutive substages, including leptotene, zygotene, pachytene, diplotene, and

diakinesis (Fig 2A). However, spermatocytes at the diplotene and diakinesis stages were

rarely observed in *Znhit1*-sKO testes (Fig 2B, C). We also examined the expression of

H1T, a middle pachytene marker, and no difference in H1T expression was observed in

Znhit1$^{-/-}$ spermatocytes (Fig S2C). *Znhit1* knockout resulted in increased

TUNEL-positive spermatocytes, suggesting that these defective spermatocytes were

eliminated by the apoptotic pathway (Fig 2D, E). We further performed scRNA-seq

analysis of postnatal day 35 (P35) testes from control and *Znhit1*-sKO. Uniform manifold

approximation and projection (UMAP) showed that germ cells normally progressed

through all meiotic stages and successfully gave rise to spermatids in control groups (Fig

2F). By contrast, in the *Znhit1* knockout group, late pachytene spermatocytes decreased

significantly, and only very few subsequent germ cell types were observable. *Pou5f2* is

expressed explicitly at the diplotene stage, and we found a lack of *Pou5f2*$^{+}$ diplotene

spermatocytes in *Znhit1*-sKO testes (Fig 2G). These results indicate that *Znhit1* deletion

leads to meiotic arrest at the pachytene stage.

ZNHIT1 is required for DSB repair and meiotic recombination

To investigate the impact of ZNHIT1 on meiotic events, we first performed

immunostaining of SYCP1 and SYCP3, the essential components of the synaptonemal

complex, to examine whether ZNHIT1 regulates homologous synapses. Pachytene nuclei

with fully synapsed autosomes were observed in spermatocytes from the control (Fig. 3A).

In *Znhit1*-sKO testes, while autosomal synapsis appeared grossly normal, abnormalities

were present at the terminal ends of autosomes. During the pachytene stage, the X and Y

chromosomes undergo pairing and synapses between the short pseudoautosomal regions

(PARs). We found an increase in the percentage of unsynapsed X-Y chromosomes in

Znhit1-sKO spermatocytes (Fig 3B). Thus, deletion of *Znhit1* results in impaired

chromosomal synapsis.

Next, we examined the programmed formation and repair of DSBs by co-staining SYCP3

and γH2AX, a DSB marker, on spermatocyte chromosome spreads. In both control and

Znhit1-sKO spermatocytes, γH2AX signals were evident in the leptotene and zygotene

stages (Fig 3C). However, in *Znhit1*-deficient pachynema, we still observed diffuse

γH2AX signals on the autosomes, while in control pachynema, γH2AX signals only

accumulated on the X-Y chromosomes. These results demonstrate that ZNHIT1 is

required for completing DSB repair.

To understand how ZNHIT1 regulates DSB repair, we performed an immunostaining

analysis using the markers RPA2 and RAD51, the essential factors involved in DSB

repair. We found an increase in RPA2 and RAD51 counts in both zygonema and

pachynema of *Znhit1*-deficient testes (Fig 3D, E, and S3A, B), indicating that *Znhit1*

deletion delayed recombinational repair, resulting in defective resolution of autosomal

DSBs in *Znhit1*$^{-/-}$ pachynema. The formation and repair of DSBs play crucial roles in

initiating and facilitating meiotic recombination, respectively. To examine the

consequences of unrepaired DSBs on meiotic recombination in *Znhit1* mutants, we

performed an immunostaining analysis against MLH1, a marker of meiotic crossover

formation. Indeed, we found an almost absence of MLH1 foci in *Znhit1*$^{-/-}$ pachytene

spermatocytes compared with controls (Fig 3F and S3C), indicating that *Znhit1* deletion

severely disrupted meiotic crossover formation.

scRNA-seq analysis reveals that *Znhit1* deletion impairs meiotic transcriptional

programs

We next investigated whether *Znhit1* deletion impacts meiotic gene expression programs.

To address this, we isolated testicular cells from control and *Znhit1*-sKO mice at P16 and

performed scRNA-seq analysis. scRNA-seq analysis identified 6 distinct germ cell types

based on the expression of canonical stage-specific markers (Chen et al., 2018), including

undifferentiated spermatogonia (*Zbtb16*), differentiating spermatogonia (*Kit*),

preleptotene spermatocytes (*Nacad*), leptotene spermatocytes (*Meiob*), zygotene

spermatocytes (*Psma8*), and pachytene spermatocytes (*Piwil1*) (Fig 4A and S4A, B). We

then analyzed the average expression of gene sets dynamically changed during meiotic

transitions, including genes upregulated or downregulated at the leptotene-to-preleptotene

(Lep/PreL), zygotene-to-leptotene (Zyg/Lep), and pachytene-to-zygotene (Pac/Zyg)

stages (Table S2). While *Znhit1* deletion had minimal effects on genes activated or

repressed during early meiotic transitions (PreL→Lep and Lep→Zyg), it specifically and

severely impaired the transcriptional programs at the zygotene-to-pachytene transition.

Specifically, the average expression of 1,560 genes was activated at the pachytene stage

(referred to as pachytene-activated genes, or PGA genes), while *Znhit1* deletion

downregulated these PGA genes (Fig 4B). Functional enrichment analysis revealed that

these PGA genes were primarily involved in DNA repair, cilium organization, and

spermatid development, consistent with the biological process of germ cell development

(Table S3).

Moreover, we identified differentially expressed genes (DEGs) in each spermatocyte type

between the control and *Znhit1*-sKO groups. The number of DEGs gradually increased

from the preleptotene to the pachytene stage, with downregulated genes predominating

(Fig 4C and Table S4). In particular, *Znhit1$^{-/-}$* pachytene spermatocytes exhibited 2,785

downregulated genes, among which were most of the pachytene-activated genes (1,094

out of 1,560, 70.1%). Functional enrichment analysis revealed that upregulated genes

following *Znhit1* deletion were associated with apoptotic pathways, supporting an

observation that *Znhit1* deletion resulted in germ cell clearance through apoptosis (Fig

4D). Moreover, downregulated GO terms included germ cell development and DNA

recombination (Fig 4E and Table S5). Interestingly, recent studies showed that cilium

organization was essential for spermatocyte development (Mytlis et al., 2022), and *Znhit1*

deletion reduced the expression of cilium-related genes in zygotene and pachytene

spermatocytes (Fig 4E). We further conducted bulk RNA-seq experiments with P14

control and *Znhit1*-sKO testicular cells and identified 882 DEGs (812 downregulated

genes and 70 upregulated genes) in *Znhit1*-sKO testicular cells (Fig S5A, and Table S6).

Through integrative analysis of bulk and single-cell RNA-seq data, we found that the

expression of these 812 genes was activated during the zygotene-to-pachytene transition,

and *Znhit1* deletion significantly reduced the expression of these genes (Fig S5B). We

also analyzed the expression of XY-linked genes. As shown in Figure 4F, while only

modest numbers of XY-DEGs were detected at preleptotene and leptotene, a striking

accumulation of upregulated XY-DEGs was observed at zygotene (120 genes) and

pachytene (119 genes). This aberrant activation of XY-linked genes directly reveals a

failure of Meiotic Sex Chromosome Inactivation (MSCI) in *Znhit1*-knockout

spermatocytes. Together, these data indicate that ZNHIT1 is essential for the regulation of

meiotic transcriptional programs.

Consistent with meiotic recombination defects in *Znhit1* mutants, genes associated with

homologous recombination were repressed following *Znhit1* deletion, including *Ccnb1ip1*,

Rnf212, *Spo16*, *Ankrd31*, and *Terb1* (Boekhout et al., 2019; Papanikos et al., 2019; Qiao

et al., 2014; Rao et al., 2017; Reynolds et al., 2013; Shibuya et al., 2014; Wang et al.,

2019; Zhang et al., 2019) (Fig S5C, D).

Znhit1 deletion impairs H2A.Z incorporation into pachytene chromatin

We then asked how ZNHIT1 regulates transcription. ZNHIT1 is a subunit of the SRCAP

complex that facilitates the incorporation of the histone variant H2A.Z by replacing H2A

(Fig S6A)(Yu et al., 2024). Immunostaining against H2A.Z showed a gradual increase in

H2A.Z accumulation on autosomes from the leptotene to the diakinesis stage, while the

H2A.Z signal was markedly low on the X-Y body (Fig 5A). *Znhit1* loss decreased H2A.Z

staining on the pachytene cells, suggesting that ZNHIT1 is required for maintaining

H2A.Z integrity during meiosis.

To profile H2A.Z target genes, we performed chromatin immunoprecipitation followed

by sequencing (ChIP-seq) against H2A.Z in P14 testicular cells. Using the ChromHMM

model to annotate H2A.Z signals with meiocyte chromatin states defined by published

epigenetic markers (Spruce et al., 2020), we found that H2A.Z sites were primarily

enriched in chromatin state 1 and state 2 (promoters and enhancers) but were less

enriched in recombination hotspots (Fig 5B, C), consistent with previously published

results (Spruce et al., 2020). Quantitative analysis showed significant H2A.Z reduction at

37.9% (23,926 of 63,109) of H2A.Z-bound genome sites following *Znhit1* deletion (Fig

5D). Comparison analysis with bulk RNA-seq data identified 480 downregulated DEGs

with decreased H2A.Z signals (Fig 5E). ~41.3 and 37.8% of the downregulated H2A.Z

peaks occupied promoters and enhancers, respectively. At promoters, these

downregulated H2A.Z preferentially occupied active promoters (H3K4me3 only), while

H2A.Z occupied three types of enhancers: active, poised, and primed enhancers (Fig 5F,

and S6B, C). These findings indicate that *Znhit1* deletion impairs H2A.Z incorporation

into pachytene chromatin.

H2A.Z deposition cooperates with A-MYB to regulate transcription

We further asked how ZNHIT1/H2A.Z deposition facilitates lineage-specific gene

expression. We performed TF motif analysis using active promoters/enhancers with

decreased H2A.Z signals. This analysis identified several MYB family TFs, including

MYB, A-MYB, and B-MYB (Fig 6A). Gene regulatory network analysis conducted by

SCENIC further identified A-MYB (encoded by the *Mybl1* gene) as the core regulatory

TF in spermatocytes (Fig 6B). We reanalyzed RNA-seq data in P14 testes from

A-MYB-deficient mice (Li et al., 2013) and found that A-MYB-deficient DEGs

correlated with those observed in *Znhit1*-deficient testicular cells (Fig 6C). Although

Znhit1 deletion didn't affect *Mybl1* mRNA expression, transcriptome analyses showed

that the expression of A-MYB target genes was significantly reduced in *Znhit1*$^{-/-}$

pachytene spermatocytes (Fig 6D, E, and S7A). These results suggest that *Znhit1*$^{-/-}$ and

Mybl1$^{-/-}$ pachytene spermatocytes share common gene expression alterations.

Next, we compared the genome binding sites of H2A.Z and A-MYB. Utilizing published

A-MYB ChIP-seq data from P14 testicular cells (Li et al., 2013), we identified 6,088

A-MYB binding signals, with 78.1% (4755) of these coinciding with H2A.Z peaks (Fig

6F). Annotation of genomic features showed that approximately 66.52% and 12.02% of

these overlapping sites occupied active promoters and enhancers, respectively,

representing a stronger enrichment compared to other regulatory elements (Fig 6G).

Genes associated with H2A.Z and A-MYB binding included essential HR-related genes,

such as *Ccnb1ip1* (Fig 6H). Moreover, we showed that *Znhit1* deletion resulted in a

marked reduction in H2A.Z signals and nascent transcription identified by KAS-seq

signals (Fig 6I). We further linked KAS□seq signals with gene expression profiles, and

found that *Znhit1* depletion caused a global reduction in KAS□seq signals, especially at

promoters of downregulated genes (Fig S8A), indicating defective transcription bubble

formation. In comparison, genes with increased expression showed low KAS□seq signals

in both control and mutant groups, likely reflecting indirect regulation. These results

further support the essential role of ZNHIT1 in transcriptional regulation.

Previous studies have shown the fundamental role of A-MYB in activating meiotic

enhancers (Maezawa et al., 2020). We found that *Znhit1* deletion resulted in reduced

KAS-seq signals at active enhancers (Fig 6I). Interestingly, the chromatin accessibility

detected by ATAC-seq at active enhancers increased after *Znhit1* deletion, suggesting that

Znhit1 deletion results in dysregulated chromatin state. These findings suggest that

ZNHIT1/H2A.Z deposition is essential for enhancer activation.

Together, these results demonstrate the critical role of the ZNHIT1/H2A.Z/A-MYB axis

in governing meiotic transcriptional programs.

Discussion

Meiosis plays a crucial role in the production of haploid gametes. Any disturbances during meiotic progress are known to lead to infertility and congenital diseases. Here we delineate the multifaceted functions of the chromatin remodeler ZNHIT1 in regulating meiotic progression. Our study provides convincing data to show that the ZNHIT1/H2A.Z/A-MYB axis controls spermatocyte development, meiotic recombination, and PGA. Thus, these findings underscore the significance of specific chromatin structures in governing appropriate transcriptional reprogramming and precise meiotic progression.

The role of H2A.Z in DNA injury and repair has been extensively studied (Colino-Sanguino et al., 2022; Dong et al., 2014; Xu et al., 2012; Yamada et al., 2018), but its function in programmed DSB formation and repair during mammalian meiosis remains unknown. To address this question, we generated *Znhit1* conditional knockout mice specifically during male meiosis to examine the role of ZNHIT1-mediated H2A.Z deposition in the formation and repair of programmed DSB. One new observation in this study is that *Znhit1* loss is dispensable for DSB formation but necessary for DNA break repair during meiosis. Intriguingly, unlike the observations in plant meiosis but consistent with Christopher L. Baker's group observation in mouse, we confirmed that H2A.Z does not directly bind DNA recombination sites in mammalian spermatocytes (Choi et al., 2013; Spruce et al., 2020; Wang and Copenhaver, 2018). Moreover, we showed that *Znhit1* deletion delays recombinational repair but has limited impacts on DNA resection, ultimately causing defective meiotic crossover formation. Importantly, our study demonstrates that ZNHIT1 is essential for the expression of meiotic recombination-related genes, such as *Ccnb1ip1* and *Rnf212*, implying that ZNHIT1/H2A.Z modulates meiotic recombination through transcriptional regulation, rather than a direct regulatory effect of ZNHIT1 on recombination machinery. Therefore, our data reveal a different mechanism underlying meiotic recombination between plants and mammals.

Pachytene spermatocytes utilize a highly specific mechanism of meiotic surveillance, the pachytene checkpoint, to prevent aneuploidy formation by removing abnormal germ cells with incomplete chromosome synapsis and defective homologous recombination (Huang and Roig, 2023; Roeder and Bailis, 2000; Subramanian and Hochwagen, 2014). Two checkpoint pathways exist in male meiosis: one responding to unrepaired DSBs and the

other triggered by MSCI failure (Royo et al., 2013). However, whether epigenetic

regulation is involved in meiotic surveillance is not well defined. Previous studies have

reported that upon stimulation by unrepaired DSBs, the MRE11-ATM-CHK2 pathway is

activated to eliminate aberrant germ cells (Marcet-Ortega et al., 2017). Our study found

that *Znhit1* deletion impairs pachytene development and XY-linked gene repression,

ultimately causing meiotic pachytene arrest and apoptosis. Therefore, these findings

demonstrate that ZNHIT1 acts as an important chromatin factor involved in the regulation

of the pachytene checkpoint.

It has long been noticed that in meiotic prophase I, chromatin that has not yet completed

chromosomal synapses is transcriptionally inactive, known as meiotic silencing of

unsynapsed chromatin (MSUC) (Turner et al., 2005). When germ cells enter the

pachytene stage, a large number of protein-coding genes and non-coding RNAs begin to

be actively expressed, known as PGA. Previous studies have shown that transcription

factors A-MYB and BRDT are involved in transcriptional activation during meiotic

prophase, but how PGA takes place in the chromatin context is unclear. In this study, we

found that the expression of the chromatin remodeler Znhit1 is specifically upregulated

during the zygotene-to-pachytene transition. We also observed that H2A.Z deposition is

enriched in the autosomes but less in the sex chromosomes at the pachytene stage. *Znhit1*

deletion in spermatocytes repressed pachytene gene activation globally and reduced

chromosome-wide H2A.Z deposition, supporting a central role of ZNHIT1 in PGA

regulation and chromatin remodeling. It has been known that A-MYB is a transcription

factor that controls pachytene transcription activation. We found that *Znhit1*-deficient

spermatocytes phenocopied abnormal meiotic phenotypes observed in A-MYB mutants,

such as X-Y synapsis failure, impaired DSB repair, and defective HR (Alexander et al.,

2023; Bolcun-Filas et al., 2011; Li et al., 2013; Maezawa et al., 2020). Our results also

revealed that ZNHIT1/H2A.Z cooperates with A-MYB to regulate PGA gene activation.

Therefore, our study illuminates the molecular mechanisms underlying the fundamental

question of how PGA is regulated.

The histone variant H2A.Z is enriched at gene promoters and regulatory regions, yet there

is ongoing debate about its role in transcriptional regulation. A recent paper reported that

H2A.Z knockout in post-mitotic muscle cells has limited effects on gene expression

(Belotti et al., 2020). In this study, we utilized the meiotic prophase as a model system

(where DNA replication does not occur) to study H2A.Z's function in transcriptional

regulation. We found that ZNHIT1-mediated H2A.Z deposition, independent of DNA

replication, is indispensable for the transcriptional activation of a large number of meiotic

genes. One explanation of this conflicting phenomenon is that H2A.Z dynamics, but not

stable H2A.Z accumulation is essential for priming transcriptional changes.

Through the integration of functional and molecular evidence, our findings establish the

critical involvement of ZNHIT1-dependent chromatin remodeling in the orchestration of

meiotic progression and coordination of various meiotic processes, such as HR and PGA.

Furthermore, we pinpoint the pivotal role played by the ZNHIT1/H2A.Z/A-MYB axis in

driving transcriptional reprogramming during meiotic prophase. Taken together, this

study deepens our understanding of the interplay between epigenetic regulation and

mammalian meiosis.

Materials and Methods

Animals. *Znhit1*^{fl/fl} mice have been previously described (Zhao et al., 2019) and are

available from the Model Animal Research Center of Nanjing University (MARC,

Nanjing, China). The *Stra8-cre* knock-in mouse line was kindly provided by Dr.

Ming-Han Tong(Lin et al., 2017). All mice were maintained on the C57BL/6J background.

Germ cell-specific *Znhit1* knockout mice (*Znhit1*^{fl/fl}; *Stra8-cre*) were obtained by crossing

Znhit1^{fl/+}; *Stra8-cre* mice with *Znhit1*^{fl/fl} mice. All mice were housed in the SPF

(Specific-Pathogen-Free) animal facility with standard 12 h light/dark cycles and standard

temperature (22 to 24□). All mice were provided with ad libitum access to standard

laboratory food and water. All experiments in this study were performed following

relevant guidelines and approved by the Animal Care and Use Committee of Fudan

University.

Histological and immunohistochemical analysis. Testes were fixed in modified

Davidson's fixative as previously described (Latendresse et al., 2002), embedded in

paraffin, and sectioned. For periodic acid-Schiff (PAS)-hematoxylin staining, 5 µm testis

sections were deparaffinized, rehydrated, and stained with Schiff's reagent and

hematoxylin solution. For immunofluorescent staining, 5 µm testis sections were

retrieved by sodium citrate antigen retrieval buffer (pH 6.0) and blocked with 5% BSA in

PBS for 30 min at room temperature. The sections were later incubated overnight at 4°C

with primary antibodies as follows: mouse anti-SYCP3 (Abcam, ab97672), rabbit

anti-HSPA2 (Abcam, ab108416), rabbit anti-pH3 (Millipore, H0412), rabbit anti-H1T

(Invitrogen, PA5-51200), or lectin PNA (Sigma□Aldrich, L7381). On the following day,

secondary fluorescein-conjugated antibodies and DAPI (Sigma□Aldrich, D9542) were

added for 1 h, followed by Fluoromount-G mounting (Southern Biotech, 0100-01).

Images were analyzed using the confocal microscope.

TUNEL staining was carried out using the DeadEnd^{TM} Fluorometric TUNEL System

(Promega, G3250) according to the manufacturer's instructions.

Immunostaining of spermatocyte chromosome spreads. Spermatocyte chromosome

spreads were prepared as previously described (Alavattam et al., 2018). Briefly, the

clumps of seminiferous tubules were transferred to the hypotonic extraction buffer (30

mM Tris base, 17 mM trisodium citrate, 5 mM EDTA, 50 mM sucrose, 5 mM

dithiothreitol (DTT), and 1× protease inhibitor, pH 8.2) and incubated on ice for 1.5 h.

The clumps of tubules were then transferred to 30 µL of ice-cold 100 mM sucrose and

mashed gently using tweezers to obtain a cell suspension. An additional 30 µL of ice-cold

100 mM sucrose was added to the cell suspension and mixed several times. Positively

charged slides were incubated in the ice-cold fixation solution (2% paraformaldehyde, 0.1%

Triton X-100, and 0.02% SDS, pH 9.2) for 3 min. 30 µL of the diluted cell suspension

was applied to the slides and incubated in humid chambers at room temperature for 2 h.

The slides were washed with 0.4% Photo-Flo 200 and stored at -80□.

For immunostaining of spermatocyte chromosome spreads, slides were washed with PBS

and blocked with 5% BSA in PBS for 30 min at room temperature. The slides were

stained with primary antibodies as follows: rabbit anti-SYCP3 (Abcam, ab15093), mouse

anti-SYCP3 (Abcam, ab97672), rabbit anti-SYCP1 (Abcam, ab15090), mouse

anti-γH2AX (BioLegend, 613401), rabbit anti-RAD51 (Proteintech, 14961-1-AP), rabbit

anti-RPA2 (Proteintech, 10412-1-AP), rabbit anti-MLH1 (Cell Signaling Technology,

3515T), rabbit anti-H2A.Z (Abcam, ab4174).

Znhit1 **in situ hybridization.** Testis sections (5 µM) at the indicated times were prepared

for *Znhit1* in situ hybridization with the RNAscope kit (Advanced Cell Diagnostics,

323100) according to the manufacturer's instructions.

Quantitative reverse transcription PCR (RT-qPCR). Total RNA was isolated using the

RNeasy Mini-plus Kit (Qiagen, 74134) and then reverse-transcribed into complementary

DNA (cDNA) with the GoScript Reverse Transcription System (Promega, A5003). cDNA

was used as the template for the quantitative PCR assay using 2×SYBR Green qPCR

Master Mix (Bimake, B21202). Quantitative PCR primers are listed in Table S7.

RNA-seq library generation and sequencing. Total RNA from fresh testes was isolated

and mRNA was purified with magnetic beads (Vazyme, N401-01). Then, mRNA was

fragmented and processed to generate RNA-seq libraries (Vazyme, NR605-01). Over 40

million reads were obtained per sample using the Illumina NovaSeq platform for 2

independent biological replicates.

Single-cell RNA sequencing (scRNA-seq). P16 and P35 control and *Znhit1*-sKO testes

were digested by collagenase IV and trypsin at 37□ for 10 min to obtain testicular cell

suspensions. scRNA-seq libraries were constructed using a 10x Genomics kit and

sequenced on the Illumina platform.

ChIP-seq library generation and sequencing. ChIP-seq library generation was

performed as previously described. Briefly, P14 control and *Znhit1*-sKO testes were

digested by collagenase IV and trypsin at 37□ for 10 min, crosslinked with 1%

formaldehyde for 10 min, and quenched with glycine. The cells were lysed and sheared

with a Bioruptor Plus machine for 20 min. Then, 2 μg of anti-H2A.Z antibody (Abcam,

ab4174) was added to the sonicated chromatin and incubated overnight at 4□. The

following day, 20 μL of protein G beads were added and incubated for 2 h at 4□. The

beads were washed and reverse-crosslinked for 4 h at 65□. DNA was extracted using

phenol-chloroform and subjected to library construction using the VAHTS Universal

DNA Library Prep Kit for Illumina (Vazyme, ND607-01) according to the manufacturer's

instructions. Over 30 million reads were obtained per sample using the Illumina NovaSeq

platform for 2 independent biological replicates.

KAS-seq library and ATAC-seq library generation and sequencing. Testes were

digested by collagenase IV and trypsin at 37□ for 10 min. Pachytene cells were sorted

using fluorescence-activated cell sorting (FACS) as previously described (Long et al.,

2017). For KAS-seq, pachytene cells were labeled with N3-kethoxal, and DNA was

isolated using the DNA Clean and Concentrator kit (Zymo, D4013) and subjected to

library construction using Q5 high-fidelity DNA polymerase (New England Biolabs,

M0544S). Genomic DNA was fragmented using Tn5 transposase and subjected to library

construction. For ATAC-seq, pachytene nuclei were isolated and fragmented with Tn5

transposase. DNA was isolated using the DNA Clean and Concentrator kit (Zymo, D4013)

and subjected to library construction using Q5 high-fidelity DNA polymerase (New

England Biolabs, M0544S). For high-throughput sequencing, over 30 million reads were

obtained per sample using the Illumina NovaSeq platform for 2 independent biological

replicates.

RNA-seq data analysis. We performed RNA-seq analysis as described previously

(Pertea et al., 2016). Briefly, after removing adapters using Cutadapt (v2.5) (Kechin et al.,

2017), paired-end reads were aligned to the annotated mouse transcripts (mm10 Gencode

vM23 release) using Hisat2 (v2.2.1) (Kim et al., 2015; Kim et al., 2019). Gene expression

levels were calculated using StringTie (v2.2.1) (Pertea et al., 2015). Read counts were

calculated using a Python script (prepDE.py3) provided by the StringTie development

team. Differentially expressed genes were identified using the R package DESeq2

(v1.38.3) (Love et al., 2014). Genes with read counts > 50 in at least one sample were

kept for further analysis. A given gene was considered to significantly changed if the

adjusted P value (padj) was < 0.05, the P value was < 0.01, and the fold-change was≥ 2.

Gene Set Enrichment Analysis (GSEA) was carried out using GSEA software

(Subramanian et al., 2005). GO analysis was performed using clusterProfiler (v4.6.2) (Wu

et al., 2021).

scRNA-seq data analysis. FASTQ files were run through CellRanger (v7.1.0) software

with default parameters for de-multiplexing, aligning reads with STAR software to mm10,

and counting unique molecular identifiers (UMIs). As input files of the Seurat R package

(v4.4.0), the filtered gene expression matrices were then used for downstream analyses

(Butler et al., 2018). Low-quality cells were filtered (expressing < 500 or >6,000 unique

gene counts and >15% mitochondrial reads). Principal component analysis was performed

on SCT-transformed data using 3,000 variable genes. The top 50 principal components

were used for clustering and visualized using the UMAP algorithm in the Seurat R

package. The "FindAllMarkers" function of the Seurat R package was used to calculate

cluster-specific genes. Marker genes for each cluster are shown in Table S8.

The "FindMarkers" function of the Seurat R package was used to identify differentially

expressed genes (DEGs) for spermatocytes (preleptotene, leptotene, zygotene, and

pachytene). Only those with $|\text{'avg\_logFC'}| > 0.25$ and 'p_val_adj' < 0.05 were considered

as DEGs. For the transcriptional regulatory network analysis, the raw count matrix was

used by the pyscenic (v0.12.1) workflow using default parameters (Aibar et al., 2017).

Then, the output loom file was opened by the SCENIC R package (v1.3.1), and AUC

values of regulons were extracted for visualization of downstream transcription factor

activity.

ChIP-seq and KAS-seq data analysis. For analyzing ChIP-seq and KAS-seq data, we

used the ENCODE ChIP-seq pipeline (v2.2.1,

https://github.com/ENCODE-DCC/chip-seq-pipeline2). Raw reads were cleaned by

Cutadapt (v2.5) (Kechin et al., 2017). After removing adapters, clean reads were aligned

against the mouse mm10 genome using bwa (v0.7.17) (Li and Durbin, 2009). Then, sam

files were converted to bam files using samtools (v1.9) (Li et al., 2009). PCR duplicates

were removed using Picard (v2.20.7) (https://broadinstitute.github.io/picard/). Histone

ChIP-seq peaks (such as H2A.Z and histone marks) and KAS-seq peaks were called using

MACS2 (v2.2.4) (Zhang et al., 2008), while transcription factor ChIP-seq peaks (such as

A-MYB) were called using the R packages spp (v1.15.5)

(https://github.com/hms-dbmi/spp). The conservative narrow peaks were used for

downstream analysis.

For visualization, bam files were converted to bigWig files using deepTools (v3.3.1)

(Ramirez et al., 2014), and bigWig files were used to calculate tag density under 50 bp

resolution. In addition, CPM was used to normalize the number of reads per bin. We used

the R package DiffBind (v3.8.4)

(http://bioconductor.org/packages/release/bioc/html/DiffBind.html) for differential peak

analysis and the R package rGREAT (v2.1.11) (Gu and Hubschmann, 2023) for peak

annotation. We used ComputeMatrix and plotHeatmap of deepTools to generate count

matrices and heatmaps, respectively. H2A.Z enrichment within chromatin states was

annotated using the ChromHMM model (Ernst and Kellis, 2012; Spruce et al., 2020). TF

motif enrichment was calculated using HOMER (v4.11) (Heinz et al., 2010). We used

deepTools to draw Pearson's correlation scatter plots for the correlation between

replicates of each experiment.

For promoter and enhancer type analysis, H3K27ac ChIP-seq data in wild-type pachytene

spermatocytes were downloaded from the published data (GSE107398) (Adams et al.,

2018); ChIP-seq data for H3K4me1, H3K4me3, and H3K27me3 in wild-type pachytene

spermatocytes were downloaded from the published data (GSE132446) (Chen et al.,

2020).

ATAC-seq data analysis. ATAC-seq data were analyzed using a standardized ENCODE

ATAC-seq pipeline (v2.2.2, https://github.com/ENCODE-DCC/atac-seq-pipeline). The

mouse reference genome (mm10) was used in the pipeline. Trimming, mapping, and

duplicate-removing were performed as the ENCODE pipeline suggested using Cutadapt

(v2.5) (Kechin et al., 2017), Bowtie2 (v2.3.4.3) (Langmead et al., 2009), and Picard

(v2.20.7) (https://broadinstitute.github.io/picard/) respectively. MACS2 (v2.2.4) (Zhang

et al., 2008) was used for peak calling.

Statistics and reproducibility. Statistical analyses were performed using R and Prism 9.

Data are presented as means ± s.d. unless otherwise indicated. The two-tailed, unpaired,

Student's t-test, Mann–Whitney test, or one-way or two-way ANOVA were performed to

analyze statistical significance.

Supplementary Materials

This manuscript file includes Figs. S1 to S9 and Tables S1 and S8.

Reference

Adams, S.R., Maezawa, S., Alavattam, K.G., Abe, H., Sakashita, A., Shroder, M., Broering,

T.J., Sroga Rios, J., Thomas, M.A., Lin, X., *et al.* (2018). RNF8 and SCML2 cooperate to

regulate ubiquitination and H3K27 acetylation for escape gene activation on the sex

chromosomes. PLoS genetics *14*, e1007233.

Aibar, S., Gonzalez-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H., Hulselmans, G.,

Rambow, F., Marine, J.C., Geurts, P., Aerts, J., *et al.* (2017). SCENIC: single-cell regulatory

network inference and clustering. Nature methods *14*, 1083-1086.

Alavattam, K.G., Abe, H., Sakashita, A., and Namekawa, S.H. (2018). Chromosome Spread

Analyses of Meiotic Sex Chromosome Inactivation. Methods in molecular biology (Clifton, NJ)

1861, 113-129.

Alexander, A.K., Rice, E.J., Lujic, J., Simon, L.E., Tanis, S., Barshad, G., Zhu, L., Lama, J.,

Cohen, P.E., and Danko, C.G. (2023). A-MYB and BRDT-dependent RNA Polymerase II

pause release orchestrates transcriptional regulation in mammalian meiosis. Nature

communications *14*.

Baudat, F., Imai, Y., and de Massy, B. (2013). Meiotic recombination in mammals: localization

and regulation. Nature reviews Genetics *14*, 794-806.

Belotti, E., Lacoste, N., Simonet, T., Papin, C., Padmanabhan, K., Scionti, I., Gangloff, Y.G.,

Ramos, L., Dalkara, D., Hamiche, A., *et al.* (2020). H2A.Z is dispensable for both basal and

activated transcription in post-mitotic mouse muscles. Nucleic acids research *48*, 4601-4613.

Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P.C., Nicolas, A., and Forterre, P. (1997).

An atypical topoisomerase II from Archaea with implications for meiotic recombination. Nature

386, 414-417.

Boekhout, M., Karasu, M.E., Wang, J., Acquaviva, L., Pratto, F., Brick, K., Eng, D.Y., Xu, J.,

Camerini-Otero, R.D., Patel, D.J., *et al.* (2019). REC114 Partner ANKRD31 Controls Number,

Timing, and Location of Meiotic DNA Breaks. Molecular cell *74*, 1053-1068 e1058.

Bolcun-Filas, E., Bannister, L.A., Barash, A., Schimenti, K.J., Hartford, S.A., Eppig, J.J.,

Handel, M.A., Shen, L., and Schimenti, J.C. (2011). A-MYB (MYBL1) transcription factor is a

master regulator of male meiosis. Development *138*, 3319-3330.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nature

biotechnology *36*, 411-420.

Cai, Y., Jin, J., Florens, L., Swanson, S.K., Kusch, T., Li, B., Workman, J.L., Washburn, M.P.,

Conaway, R.C., and Conaway, J.W. (2005). The mammalian YL1 protein is a shared subunit

of the TRRAP/TIP60 histone acetyltransferase and SRCAP complexes. The Journal of

biological chemistry *280*, 13665-13670.

Cecchini, K., Biasini, A., Yu, T., Saflund, M., Mou, H., Arif, A., Eghbali, A., Colpan, C.,

Gainetdinov, I., de Rooij, D.G., *et al.* (2023). The transcription factor TCFL5 responds to

A-MYB to elaborate the male meiotic program in mice. Reproduction *165*, 183-196.

Chen, Y., Lyu, R.T., Rong, B.W., Zheng, Y.X., Lin, Z., Dai, R.F., Zhang, X., Xie, N.N., Wang,

S.Q., Tang, F.C., *et al.* (2020). Refined spatial temporal epigenomic profiling reveals intrinsic

connection between PRDM9-mediated H3K4me3 and the fate of double-stranded breaks. Cell

research *30*, 256-268.

Chen, Y., Zheng, Y., Gao, Y., Lin, Z., Yang, S., Wang, T., Wang, Q., Xie, N., Hua, R., Liu, M.,

et al. (2018). Single-cell RNA-seq uncovers dynamic processes and critical regulators in

mouse spermatogenesis. Cell research *28*, 879-896.

Choi, K., Zhao, X., Kelly, K.A., Venn, O., Higgins, J.D., Yelina, N.E., Hardcastle, T.J.,

Ziolkowski, P.A., Copenhaver, G.P., Franklin, F.C.*, et al.* (2013). Arabidopsis meiotic

crossover hot spots overlap with H2A.Z nucleosomes at gene promoters. Nature genetics *45*,

1327-1336.

Colino-Sanguino, Y., Clark, S.J., and Valdes-Mora, F. (2022). The H2A.Z-nuclesome code in

mammals: emerging functions. Trends in genetics : TIG *38*, 273-289.

Cuadrado, A., Corrado, N., Perdiguero, E., Lafarga, V., Munoz-Canoves, P., and Nebreda,

A.R. (2010). Essential role of p18Hamlet/SRCAP-mediated histone H2A.Z chromatin

incorporation in muscle differentiation. The EMBO journal *29*, 2014-2025.

Cuadrado, A., Lafarga, V., Cheung, P.C., Dolado, I., Llanos, S., Cohen, P., and Nebreda, A.R.

(2007). A new p38 MAP kinase-regulated transcriptional coactivator that stimulates

p53-dependent apoptosis. The EMBO journal *26*, 2115-2126.

Dong, S., Han, J., Chen, H., Liu, T., Huen, M.S., Yang, Y., Guo, C., and Huang, J. (2014). The

Human SRCAP Chromatin Remodeling Complex Promotes DNA-End Resection. Current

biology : CB *24*, 2097-2110.

Ernst, C., Eling, N., Martinez-Jimenez, C.P., Marioni, J.C., and Odom, D.T. (2019). Staged

developmental mapping and X chromosome transcriptional dynamics during mouse

spermatogenesis. Nature communications *10*, 1251.

Ernst, J., and Kellis, M. (2012). ChromHMM: automating chromatin-state discovery and

characterization. Nature methods *9*, 215-216.

Feng, Y., Zhang, Y., Lin, Z., Ye, X., Lin, X., Lv, L., Lin, Y., Sun, S., Qi, Y., and Lin, X. (2022).

Chromatin remodeler Dmp18 regulates apoptosis by controlling H2Av incorporation in

Drosophila imaginal disc development. PLoS genetics *18*, e1010395.

Gaucher, J., Boussouar, F., Montellier, E., Curtet, S., Buchou, T., Bertrand, S., Hery, P.,

Jounier, S., Depaux, A., Vitte, A.L., *et al.* (2012). Bromodomain-dependent stage-specific male

genome programming by Brdt. The EMBO journal *31*, 3809-3820.

Gray, S., and Cohen, P.E. (2016). Control of Meiotic Crossovers: From Double-Strand Break

Formation to Designation. Annu Rev Genet *50*, 175-210.

Green, C.D., Ma, Q., Manske, G.L., Shami, A.N., Zheng, X., Marini, S., Moritz, L., Sultan, C.,

Gurczynski, S.J., Moore, B.B., *et al.* (2018). A Comprehensive Roadmap of Murine

Spermatogenesis Defined by Single-Cell RNA-Seq. Developmental cell *46*, 651-667 e610.

Gu, Z., and Hubschmann, D. (2023). rGREAT: an R/bioconductor package for functional

enrichment on genomic regions. Bioinformatics *39*.

Handel, M.A., and Schimenti, J.C. (2010). Genetics of mammalian meiosis: regulation,

dynamics and impact on fertility. Nature reviews Genetics *11*, 124-136.

Hassold, T., and Hunt, P. (2001). To err (meiotically) is human: the genesis of human

aneuploidy. Nature reviews Genetics *2*, 280-291.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,

Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell identities. Molecular

cell *38*, 576-589.

Hirota, T., Blakeley, P., Sangrithi, M.N., Mahadevaiah, S.K., Encheva, V., Snijders, A.P.,

Ellnati, E., Ojarikre, O.A., de Rooij, D.G., Niakan, K.K., *et al.* (2018). SETDB1 Links the Meiotic

DNA Damage Response to Sex Chromosome Silencing in Mice. Developmental cell *47*,

645-+.

Huang, Y., and Roig, I. (2023). Genetic control of meiosis surveillance mechanisms in

mammals. Front Cell Dev Biol *11*, 1127440.

Hunter, N. (2015). Meiotic Recombination: The Essence of Heredity. Cold Spring Harbor

perspectives in biology *7*.

Kechin, A., Boyarskikh, U., Kel, A., and Filipenko, M. (2017). cutPrimers: A New Tool for

Accurate Cutting of Primers from Reads of Targeted Next Generation Sequencing. J Comput

Biol *24*, 1138-1143.

Keeney, S., Giroux, C.N., and Kleckner, N. (1997). Meiosis-specific DNA double-strand breaks

are catalyzed by Spo11, a member of a widely conserved protein family. Cell *88*, 375-384.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low

memory requirements. Nature methods *12*, 357-360.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology *37*,

907-915.

Kota, S.K., and Feil, R. (2010). Epigenetic transitions in germ cell development and meiosis.

Developmental cell *19*, 675-686.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome biology *10*, R25.

Lascarez-Lagunas, L., Martinez-Garcia, M., and Colaiacovo, M. (2020). SnapShot: Meiosis -

Prophase I. Cell *181*, 1442-1442 e1441.

Latendresse, J.R., Warbrittion, A.R., Jonassen, H., and Creasy, D.M. (2002). Fixation of testes

and eyes using a modified Davidson's fluid: comparison with Bouin's fluid and conventional

Davidson's fluid. Toxicol Pathol *30*, 524-533.

Law, N.C., Oatley, M.J., and Oatley, J.M. (2019). Developmental kinetics and transcriptome

dynamics of stem cell specification in the spermatogenic lineage. Nature communications *10*,

2787.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics *25*, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map

format and SAMtools. Bioinformatics *25*, 2078-2079.

Li, X.Z., Roy, C.K., Dong, X., Bolcun-Filas, E., Wang, J., Han, B.W., Xu, J., Moore, M.J.,

Schimenti, J.C., Weng, Z.*, et al.* (2013). An ancient transcription factor initiates the burst of

piRNA production during early meiosis in mouse testes. Molecular cell *50*, 67-81.

Lin, Z., Hsu, P.J., Xing, X., Fang, J., Lu, Z., Zou, Q., Zhang, K.J., Zhang, X., Zhou, Y., Zhang,

T.*, et al.* (2017). Mettl3-/Mettl14-mediated mRNA N6-methyladenosine modulates murine

spermatogenesis. Cell research.

Long, J., Huang, C., Chen, Y., Zhang, Y., Shi, S., Wu, L., Liu, Y., Liu, C., Wu, J., and Lei, M.

(2017). Telomeric TERB1-TRF1 interaction is crucial for male meiosis. Nat Struct Mol Biol *24*,

1073-1080.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome biology *15*, 550.

Lu, J., An, J., Wang, J., Cao, X., Cao, Y., Huang, C., Jiao, S., Yan, D., Lin, X., and Zhou, X.

(2022). Znhit1 Regulates p21Cip1 to Control Mouse Lens Differentiation. Investigative

ophthalmology & visual science *63*, 18.

Maezawa, S., Sakashita, A., Yukawa, M., Chen, X., Takahashi, K., Alavattam, K.G., Nakata, I.,

Weirauch, M.T., Barski, A., and Namekawa, S.H. (2020). Super-enhancer switching drives a

burst in gene expression at the mitosis-to-meiosis transition. Nat Struct Mol Biol *27*, 978-988.

Manterola, M., Brown, T.M., Oh, M.Y., Garyn, C., Gonzalez, B.J., and Wolgemuth, D.J. (2018).

BRDT is an essential epigenetic regulator for proper chromatin organization, silencing of sex

chromosomes and crossover formation in male meiosis. PLoS genetics *14*, e1007209.

Marcet-Ortega, M., Pacheco, S., Martinez-Marchal, A., Castillo, H., Flores, E., Jasin, M.,

Keeney, S., and Roig, I. (2017). p53 and TAp63 participate in the recombination-dependent

pachytene arrest in mouse spermatocytes. PLoS genetics *13*, e1006845.

Mytlis, A., Kumar, V., Qiu, T., Deis, R., Hart, N., Levy, K., Masek, M., Shawahny, A., Ahmad, A.,

Eitan, H.*, et al.* (2022). Control of meiotic chromosomal bouquet and germ cell morphogenesis

by the zygotene cilium. Science, eabh3104.

Ozata, D.M., Yu, T., Mou, H., Gainetdinov, I., Colpan, C., Cecchini, K., Kaymaz, Y., Wu, P.H.,

Fan, K., Kucukural, A.*, et al.* (2020). Evolutionarily conserved pachytene piRNA loci are highly

divergent among modern humans. Nat Ecol Evol *4*, 156-168.

Papanikos, F., Clement, J.A.J., Testa, E., Ravindranathan, R., Grey, C., Dereli, I., Bondarieva,

A., Valerio-Cabrera, S., Stanzione, M., Schleiffer, A.*, et al.* (2019). Mouse ANKRD31

Regulates Spatiotemporal Patterning of Meiotic Recombination Initiation and Ensures

Recombination between X and Y Sex Chromosomes. Molecular cell *74*, 1069-1085 e1011.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level

expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nature

protocols *11*, 1650-1667.

Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T., and Salzberg, S.L.

(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.

Nature biotechnology *33*, 290-295.

Qiao, H., Prasada Rao, H.B., Yang, Y., Fong, J.H., Cloutier, J.M., Deacon, D.C., Nagel, K.E.,

Swartz, R.K., Strong, E., Holloway, J.K., *et al.* (2014). Antagonistic roles of ubiquitin ligase

HEI10 and SUMO ligase RNF212 regulate meiotic recombination. Nature genetics *46*,

194-199.

Rabbani, M., Zheng, X., Manske, G.L., Vargo, A., Shami, A.N., Li, J.Z., and Hammoud, S.S.

(2022). Decoding the Spermatogenesis Program: New Insights from Transcriptomic Analyses.

Annu Rev Genet *56*, 339-368.

Ramirez, F., Dundar, F., Diehl, S., Gruning, B.A., and Manke, T. (2014). deepTools: a flexible

platform for exploring deep-sequencing data. Nucleic acids research *42*, W187-191.

Rao, H.B., Qiao, H., Bhatt, S.K., Bailey, L.R., Tran, H.D., Bourne, S.L., Qiu, W., Deshpande, A.,

Sharma, A.N., Beebout, C.J., *et al.* (2017). A SUMO-ubiquitin relay recruits proteasomes to

chromosome axes to regulate meiotic recombination. Science *355*, 403-407.

Reynolds, A., Qiao, H.Y., Yang, Y., Chen, J.K., Jackson, N., Biswas, K., Holloway, J.K.,

Baudat, F., de Massy, B., Wang, J., *et al.* (2013). RNF212 is a dosage-sensitive regulator of

crossing-over during mammalian meiosis. Nature genetics *45*, 269-278.

Roeder, G.S., and Bailis, J.M. (2000). The pachytene checkpoint. Trends in genetics : TIG *16*,

395-403.

Royo, H., Prosser, H., Ruzankina, Y., Mahadevaiah, S.K., Cloutier, J.M., Baumann, M.,

Fukuda, T., Hoog, C., Toth, A., de Rooij, D.G., *et al.* (2013). ATR acts stage specifically to

regulate multiple aspects of mammalian meiotic silencing. Genes & development *27*,

1484-1494.

San Filippo, J., Sung, P., and Klein, H. (2008). Mechanism of eukaryotic homologous

recombination. Annual review of biochemistry *77*, 229-257.

Shibuya, H., Ishiguro, K., and Watanabe, Y. (2014). The TRF1-binding protein TERB1

promotes chromosome movement and telomere rigidity in meiosis. Nature cell biology *16*,

145-156.

Spruce, C., Dlamini, S., Ananda, G., Bronkema, N., Tian, H., Paigen, K., Carter, G.W., and

Baker, C.L. (2020). HELLS and PRDM9 form a pioneer complex to open chromatin at meiotic

recombination hot spots. Genes & development *34*, 398-412.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,

Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., *et al.* (2005). Gene set enrichment

analysis: a knowledge-based approach for interpreting genome-wide expression profiles.

Proceedings of the National Academy of Sciences of the United States of America *102*,

15545-15550.

Subramanian, V.V., and Hochwagen, A. (2014). The meiotic checkpoint network: step-by-step

through meiotic prophase. Cold Spring Harbor perspectives in biology *6*, a016675.

Sun, S., Jiang, N., Jiang, Y., He, Q., He, H., Wang, X., Yang, L., Li, R., Liu, F., Lin, X., *et al.*

(2020). Chromatin remodeler Znhit1 preserves hematopoietic stem cell quiescence by

determining the accessibility of distal enhancers. Leukemia *34*, 3348-3358.

Sun, S., Jiang, Y., Zhang, Q., Pan, H., Li, X., Yang, L., Huang, M., Wei, W., Wang, X., Qiu, M.,

et al. (2022). Znhit1 controls meiotic initiation in male germ cells by coordinating with Stra8 to

activate meiotic gene expression. Developmental cell *57*, 901-913 e904.

Symington, L.S. (2014). End resection at double-strand breaks: mechanism and regulation.

Cold Spring Harbor perspectives in biology *6*.

Turner, J.M. (2015). Meiotic Silencing in Mammals. Annu Rev Genet *49*, 395-412.

Turner, J.M., Mahadevaiah, S.K., Fernandez-Capetillo, O., Nussenzweig, A., Xu, X., Deng,

C.X., and Burgoyne, P.S. (2005). Silencing of unsynapsed meiotic chromosomes in the mouse.

Nature genetics *37*, 41-47.

Wang, L., Xu, Z., Khawar, M.B., Liu, C., and Li, W. (2017). The histone codes for meiosis.

Reproduction *154*, R65-R79.

Wang, Y., Chen, Y., Chen, J., Wang, L., Nie, L., Long, J., Chang, H., Wu, J., Huang, C., and

Lei, M. (2019). The meiotic TERB1-TERB2-MAJIN complex tethers telomeres to the nuclear

envelope. Nature communications *10*, 564.

Wang, Y., and Copenhaver, G.P. (2018). Meiotic Recombination: Mixing It Up in Plants. Annu

Rev Plant Biol *69*, 577-609.

Wei, W., Tang, X., Jiang, N., Ni, C., He, H., Sun, S., Yu, M., Yu, C., Qiu, M., Yan, D., *et al.*

(2022). Chromatin remodeler Znhit1 controls bone morphogenetic protein signaling in

embryonic lung tissue branching. The Journal of biological chemistry *298*, 102490.

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., *et al.*

(2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. Innovation

(Camb) *2*, 100141.

Xu, M., Yao, J., Shi, Y., Yi, H., Zhao, W., Lin, X., and Yang, Z. (2021). The SRCAP chromatin

remodeling complex promotes oxidative metabolism during prenatal heart development.

Development *148*.

Xu, Y., Ayrapetov, M.K., Xu, C., Gursoy-Yuzugullu, O., Hu, Y., and Price, B.D. (2012). Histone

H2A.Z controls a critical chromatin remodeling step required for DNA double-strand break

repair. Molecular cell *48*, 723-733.

Yamada, S., Kugou, K., Ding, D.Q., Fujita, Y., Hiraoka, Y., Murakami, H., Ohta, K., and

Yamada, T. (2018). The histone variant H2A.Z promotes initiation of meiotic recombination in

fission yeast. Nucleic acids research *46*, 609-620.

Yoshida, S., Sukeno, M., Nakagawa, T., Ohbo, K., Nagamatsu, G., Suda, T., and Nabeshima,

Y. (2006). The first round of mouse spermatogenesis is a distinctive program that lacks the

self-renewing spermatogonia stage. Development *133*, 1495-1505.

Yoshida, S., Takakura, A., Ohbo, K., Abe, K., Wakabayashi, J., Yamamoto, M., Suda, T., and

Nabeshima, Y. (2004). Neurogenin3 delineates the earliest stages of spermatogenesis in the

mouse testis. Developmental biology *269*, 447-458.

Yu, J., Sui, F., Gu, F., Li, W., Yu, Z., Wang, Q., He, S., Wang, L., and Xu, Y. (2024). Structural

insights into histone exchange by human SRCAP complex. Cell Discov *10*, 15.

Yu, T., Fan, K., Ozata, D.M., Zhang, G., Fu, Y., Theurkauf, W.E., Zamore, P.D., and Weng, Z.

(2021). Long first exons and epigenetic marks distinguish conserved pachytene piRNA

clusters from other mammalian genes. Nature communications *12*, 73.

Zhang, Q., Ji, S.Y., Busayavalasa, K., and Yu, C. (2019). SPO16 binds SHOC1 to promote

homologous recombination and crossing-over in meiotic prophase I. Sci Adv *5*, eaau9780.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,

Myers, R.M., Brown, M., Li, W.*, et al.* (2008). Model-based analysis of ChIP-Seq (MACS).

Genome biology *9*, R137.

Zhao, B., Chen, Y., Jiang, N., Yang, L., Sun, S., Zhang, Y., Wen, Z., Ray, L., Liu, H., Hou, G.*,*

et al. (2019). Znhit1 controls intestinal stem cell maintenance by regulating H2A.Z

incorporation. Nature communications *10*, 1071.

Zheng, H., and Xie, W. (2019). The role of 3D genome organization in development and cell

differentiation. Nature reviews Molecular cell biology.

Zickler, D., and Kleckner, N. (2023). Meiosis: Dances Between Homologs. Annu Rev Genet.

Acknowledgments: We thank Chuan He for N3-kethoxal and Christopher L. Baker for

meiotic chromatin state data. **Funding:** This work was supported by grants from the

National Key Research and Development Program of China (2022YFA0806200,

2018YFC1003500, 2018YFA0800100, 2021YFC2501800), the National Natural Science

Foundation of China (32192400, 81971443, 32300702, 32350710191), the Science and

Technology Major Project of Inner Mongolia Autonomous Region of China to the State

Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock

(2020ZD0008). Shenfei Sun was supported by the fellowship of China Postdoctoral

Science Foundation (2022M720797) and the Postdoctoral Fellowship Program (Grade B)

of China Postdoctoral Science Foundation (GZB20230161).

Author Contributions: S.S. and X.L. conceived and designed the study; S.S. performed

most of the experiments with the help of Y.J., Q.Z., H.P., F.H., X.Z., Y.G., X.Y., K.G.,

W.W., H.L., Z.S., Y.S., X.T., M.Y., and R.L.; Y.J. and N.J. performed the bioinformatics

analysis; X.L. supervised the work and acquired the funding support; and S.S. and X.L.

wrote the manuscript, with contributions from all authors. **Competing interests:** The

authors declare that they have no competing interests. **Data and materials availability:**

The NGS data generated in this study were deposited in the NCBI SRA database under

accession numbers SRP467214 (RNA-seq and ChIP-seq data) and SRP467448

(ATAC-seq and KAS-seq data).

Figure 1 | *Znhit1* deletion in spermatocytes disrupts spermatogenesis. (**A**) Schematic

representation of the male germline development in mice. (**B**) Heatmap showing the

clustering of 47 chromatin factors based on the transcription levels in leptotene and

zygotene spermatocytes. (**C**) *Znhit1* in situ hybridization in postnatal day 60 (P60) testis

sections. Scale bar, 20 μm. Sg, spermatogonia; pL, preleptotene; L, leptotene; Z, zygotene;

P, pachytene; D, diplotene; MI, metaphase; R, round spermatids; E, elongating spermatids.

(**D**) Schematic diagram showing the onset of Stra8-cre expression in the male germline

development. (**E**) *Znhit1* in situ hybridization in P42 testis sections. Scale bar, 50 μm. (**F**)

Histological testicular sections in control or *Znhit1*-sKO testis sections at the indicated

times. Scale bar, 50 μm. (**G**) Immunostaining of PNA in testis sections from control or

Znhit1-sKO mice at the indicated times. rST, round spermatids; eST, elongating

spermatids. Scale bar, 20 μm. All images are representative of n = 3 mice per genotype.

Figure 2 | *Znhit1* deletion leads to meiotic pachytene arrest. (**A-C**) Immunostaining of

SYCP3 in spermatocyte chromosome spreads of control (**A**) or *Znhit1*-sKO mice (**B**).

Quantitative data are shown in (**C**). Scale bar, 10 μm. (**D** and **E**) TUNEL staining in testis

sections from P14 control or *Znhit1*-sKO mice. Quantitative data are shown in (**E**). Scale

bar, 40 μm. All images are representative of n = 3 mice per genotype. (**F**) UMAP plot

showing the annotated cells captured from P35 control and *Znhit1*-sKO testicular cells.

(**G**) UMAP plot showing the expression of *Znhit1* and *Pou5f2* in control P35 control and

Znhit1-sKO testicular cells. Data are presented as the mean ± s.d. *** $p < 0.001$.

Figure 3 | *Znhit1* deletion impairs meiotic recombination. (**A** and **B**) Immunostaining

of SYCP3 and SYCP1 in spermatocyte chromosome spreads of control or *Znhit1*-sKO

mice. Arrows indicate autosomal regions with abnormal synapsis. X, X chromosome; Y,

Y chromosome; PAR, pseudoautosomal region. Quantitative data are shown in (**B**). Scale

bar, 10 μm. (**C**) Immunostaining of SYCP3 and γH2AX in spermatocyte chromosome

spreads of control or *Znhit1*-sKO mice. Scale bar, 10 μm. (**D** and **E**) Immunostaining of

SYCP3 and RAD51 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice.

Quantitative data are shown in (**E**). Scale bar, 10 μm. (**F**) Immunostaining of SYCP3 and

MLH1 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Scale bar, 10

μm. All images are representative of n = 3 mice per genotype. Data are presented as the

mean ± s.d. *** $p < 0.001$.

Figure 4 | *Znhit1* deletion impairs meiotic transcriptional programs. (**A**) UMAP plot

showing the annotated cells captured from P16 control and *Znhit1*-sKO testicular cells. (**B**)

Violin plots showing the average expression level of upregulated or downregulated genes

between two consecutive spermatocyte stages. PreL, preleptotene; Lep, leptotene; Zyg,

zygotene; Pac, pachytene. (**C**) Volcano plot showing the distribution of upregulated and

downregulated DEGs for each cell type in spermatocytes between control and

Znhit1-sKO testicular cells. (**D** and **E**) Representative shared Gene Ontology (GO) terms

of upregulated and downregulated DEGs at consecutive spermatocyte stages. (**F**) Volcano

plot showing differentially expressed XY-linked genes in scRNA-seq data following

Znhit1 deletion.

Figure 5 | ZNHIT1 regulates H2A.Z binding on promoter and enhancer regions. (**A**)

Immunostaining of SYCP3 and H2A.Z in spermatocyte chromosome spreads of control

or *Znhit1*-sKO mice. The dashed circle indicates X-Y chromosomes. Scale bar, 10 μm. (**B**)

Representative tracks of END-seq, DMC1-SSDS, and H2A.Z ChIP-seq in wild-type,

control, or *Znhit1*-sKO testicular cells. (**C**) Heatmaps of chromatin states produced by

ChromHMM and showing different enrichment for H2A.Z in control or *Znhit1*-sKO

testicular cells. (**D**) Heatmaps and mean plots of H2A.Z ChIP-seq signal in control or

Znhit1-sKO testicular cells (n = 2). (**E**) Venn diagram showing the overlap between

decreased H2A.Z-bound genes and downregulated genes in *Znhit1*-sKO testicular cells.

(**F**) Heatmaps and mean plots of downregulated H2A.Z ChIP-seq signals surrounding

different types of promoters and enhancers.

Figure 6 | ZNHIT1/H2A.Z/A-MYB axis regulates chromatin state and gene

expression. (**A**) Motif analysis of active promoters or active enhancers with decreased

H2A.Z binding for putative transcription factor (TF)-binding sites using the HOMER

database. (**B**) Ridge map showing the A-MYB activity (AUC) at each cell type. (**C**)

Scatter plot with marginal histograms comparing the fold change of DEGs from

Mybl1-related RNA-seq data with *Znhit1*-related RNA-seq data in P14 testicular cells. (**D**)

GSEA of RNA-seq data for the control and *Znhit1*-sKO testicular cells. Selected gene sets

encoded products related to A-MYB target genes. (**E**) Violin plots showing the expression

of A-MYB target genes at consecutive spermatocyte stages. (**F**) Histogram showing

H2A.Z and A-MYB binding sites. (**G**) Pie chart showing the distribution of H2A.Z and

A-MYB co-binding sites. (**H**) Representative H2A.Z ChIP-seq tracks, KAS-seq signals,

and RNA-seq signals of the *Ccnb1ip1* locus in the control and *Znhit1*-sKO testicular cells,

compared with A-MYB ChIP-seq peaks. (**I**) Heatmaps and mean plots of H2A.Z

ChIP-seq, KAS-seq, and ATAC-seq showing signal changes within A-MYB and H2A.Z

co-binding active promoters or active enhancers. ******** $p < 0.0001$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Materials for

ZNHIT1-dependent H2A.Z deposition at meiotic prophase I underlies

pachytene gene expression and meiotic progression during male meiosis

Shenfei Sun *et al.*

907
908
*Corresponding author. Email: Xinhua Lin, xlin@fudan.edu.cn; Shenfei Sun
909
shenfei_sun@fudan.edu.cn
910

This PDF file includes:

    Figs. S1 to S9

Other Supplementary Material for this manuscript includes the following:

    Tables. S1 to S8

Figure S1. *Znhit1* expression during meiotic progression and

spermatocyte-conditional deletion of *Znhit1* with Stra8-cre. (**A** and **B**) Venn diagram

and heatmaps showing chromatin regulators highly expressed during leptotene and

zygotene spermatocytes. (**C**) Violin plot showing *Znhit1* transcription level during

spermatogenesis. Gene expression profiles derived from published scRNA-seq data (Chen

et al, 2018). (**D**) *Znhit1* in situ hybridization in P14 testis sections. Scale bar, 20 μm. (**E**)

Testis size comparison between control and *Znhit1*-sKO mice at P35. (**F**) Hematoxylin

and eosin (H&E) - stained testis sections from control and *Znhit1*-sKO mice at P60. Scale

bar, 50 μm.

937
938

Figure S2. Spermatocyte development in *Znhit1*-sKO testes. (**A**) Immunostaining of
SYCP3 and HSPA2 in testis sections of control or *Znhit1*-sKO mice at P35. Scale bar, 20
μm. (**B**) Immunostaining of SYCP3 and pH3 in testis sections of control or *Znhit1*-sKO
mice at P35. Scale bar, 20 μm. (**C**) Immunostaining of H1T in testis sections from control
and *Znhit1*-sKO mice at P35. Scale bar, 20 μm.

947 **Figure S3. ZNHIT1 is required for meiotic recombination.** (**A** and **B**) Immunostaining

948 of SYCP3 and RPA2 in spermatocyte chromosome spreads of control or *Znhit1*-sKO

949 mice. Quantitative data are shown in (**B**). Scale bar, 10 μm. (**C**) Plots showing

950 quantitative data of MLH1 foci of control or *Znhit1*-sKO mice. Data are presented as the

951 mean ± s.d. *** $p < 0.001$

952
953

955 **Figure S4. Single-cell RNA-seq of testicular cells.** (**A**) Dot plot showing the relative

956 expression and the percentage of cells expressing selected markers across scRNA-seq

957 clusters. (**B**) UMAP plots showing the normalized expression of the indicated genes in

958 scRNA-seq data.

959
960

Figure S5. Transcriptomic analysis of testicular cells. (**A**) Heatmap showing

differentially expressed genes (DEGs) in P14 testicular cells between control and

Znhit1-sKO mice. (**B**) Violin plots showing the average expression level of upregulated or

downregulated genes between P14 control and *Znhit*1-sKO testicular cells. PreL,

preleptotene; Lep, leptotene; Zyg, zygotene; Pac, pachytene. (**C** and **D**) Bar charts

showing HR-related gene expression in P14 control and *Znhit*1-sKO testes. Data from

bulk RNA-seq are shown in (**C**). Validated data from RT-qPCR are shown in (**D**). Ctrl,

control; sKO, *Znhit1*-sKO.

970
971
972

Figure S6. H2A.Z is enriched at meiotic gene promoters and enhancers. (**A**)

Schematic representation of the components of the SRCAP chromatin remodeling

complex. (**B**) Heatmaps showing H2A.Z ChIP-seq signals surrounding different types of

promoters. (**C**) Heatmaps showing H2A.Z ChIP-seq signals surrounding different types of

enhancers.

Figure S7. *Znhit1* deletion downregulates the expression of A-MYB target genes. (A)

Violin plot showing the normalized expression of *Mybl1* in control and *Znhit1*-sKO

spermatocytes at consecutive stages. (**B**) Cluster annotation on the basis of the expression

of A-MYB target genes in control or *Znhit1*-sKO testicular cells.

985
986

988   **Figure S8. KAS-seq analysis.** (**A**) Mean plots of KAS-seq showing signal changes

989   within indicated genes.

990
991

Figure S9. Reproducibility of replicates in this study. (**A**) Scatter plots showing H2A.Z
enrichments in P14 control or *Znhit1*-sKO testes for two independent H2A.Z ChIP-seq
replicates. Pearson correlation coefficients are indicated. (**B**) Scatter plots showing
KAS-seq signals in P14 control or *Znhit1*-sKO pachytene cells for two independent
H2A.Z ChIP-seq replicates. Pearson correlation coefficients are indicated. (**C**) Scatter
plots showing ATAC-seq signals in P14 control or *Znhit1*-sKO pachytene cells for two
independent H2A.Z ChIP-seq replicates. Pearson correlation coefficients are indicated.

1000

1001