H2A.Z deposition at meiotic prophase I underlies homologous recombination and pachytene genome activation during male meiosis

Shenfei Sun[1,5,*], Yamei Jiang[1,5], Ning Jiang[1,5], Qiaoli Zhang[1,4], Hongjie Pan[2], Fujing Huang[1], Xinna Zhang[4], Yuxuan Guo[4], Xiaoyu You[1], Kai Gong[1], Wei Wei[1], Hanmin Liu[3], Zhenju Song[1], Yuanlin Song[1], Xiaofang Tang[1], Miao Yu[1], Runsheng Li[2], Xinhua Lin[1,3,6,*]

Affiliations

[1]State Key Laboratory of Genetic Engineering, Greater Bay Area Institute of Precision Medicine (Guangzhou), School of Life Sciences, Zhongshan Hospital, Fudan University, Shanghai, 200438, China;

[2]National Health Commission (NHC) Key Laboratory of Reproduction Regulation, Shanghai Institute for Biomedical and Pharmaceutical Technologies, Shanghai, 200032, China;

[3]The Joint Laboratory for Lung Development and Related Diseases of West China Second University Hospital, Sichuan University and School of Life Sciences of Fudan University, Chengdu, 610041, China;

[4]State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock, School of Life Sciences, Inner Mongolia University, Hohhot, 010070, China;

[5]These authors contributed equally to this work;

[6]Lead contact.

[*]Correspondence to: xlin@fudan.edu.cn (X.L.); shenfei_sun@fudan.edu.cn (S.S.).

Keywords: male meiosis; pachytene genome activation; homologous recombination; Znhit1; chromatin remodeling;

Abstract

Accurate meiotic progression is important for gamete formation and the generation of
genetic diversity. However, little is known about the identity of chromatin regulators that
underlie mammalian meiosis in vivo. Here, we identify the multifaceted functions of the
chromatin remodeler Znhit1 in governing meiosis. We observe a gradual increase in
Znhit1 expression during the meiotic prophase. Znhit1 deficiency in spermatocytes results
in arrested pachytene development, impaired DNA double-strand break repair, and
defective homologous recombination. Single-cell RNA sequencing and transcriptome
analysis reveal that Znhit1 loss downregulates the transcription of pachytene genome
activation (PGA) genes globally. Chromatin immunoprecipitation data show that Znhit1 is
needed for the incorporation of the histone variant H2A.Z into pachytene chromatin.
Moreover, we find that H2A.Z cooperates with the transcription factor A-MYB to co-bind
DNA elements and control enhancer activity. Our findings provide insights into the
regulatory mechanisms governing meiotic progression and highlight Znhit1 as a critical
regulator of meiotic recombination and PGA.
43

Introduction

Meiosis is a fundamental and conserved process that plays a crucial role in gamete formation and the generation of genetic diversity. By undergoing one round of DNA replication and two rounds of cell division, the meiotic process ensures the production of haploid cells, each with a unique combination of genetic materials (Handel and Schimenti, 2010; Lascarez-Lagunas et al., 2020). Any disruptions to the normal progression of meiosis would have significant consequences, such as aneuploidy, infertility, spontaneous abortion, and congenital diseases (Hassold and Hunt, 2001). Therefore, understanding the molecular mechanisms underlying meiosis will provide valuable insights for the diagnosis and treatment of reproductive and developmental diseases.

Multiple chromosomal events occur during meiotic prophase I, including homologous chromosome pairing and synapsis, DNA double-strand break (DSB) formation and repair, and homologous recombination (HR) (Baudat et al., 2013; Hunter, 2015; Zickler and Kleckner, 2023). In this process, homolog pairing and synapsis coincide with Spo11-mediated genome-wide formation of DSBs (Bergerat et al., 1997; Keeney et al., 1997). Following the formation of DSBs, the DNA undergoes resection, leading to the generation of single-stranded DNA overhangs. These overhangs are then coated by RPA, DMC1, and RAD51, facilitating the production of recombination intermediates. Subsequently, these intermediates are processed and resolved, ultimately forming either meiotic crossovers or non-crossovers (Gray and Cohen, 2016; San Filippo et al., 2008; Symington, 2014). The tightly coordinated timing and spatial arrangement of these meiotic events are of utmost importance for proper germ cell development.

There is transcriptional awakening during male meiotic prophase I, in which the quiescent meiotic genome becomes transcriptionally active during the zygotene-to-pachytene transition (Ernst et al., 2019; Green et al., 2018; Rabbani et al., 2022; Turner, 2015). This process, referred to as pachytene genome activation (PGA), is responsible for the expression of numerous genes and plays an essential role in controlling meiotic and post-meiotic events. Previous studies have emphasized the significance of the transcription factors (TFs) A-MYB, BRDT, and TCFL5 in activating transcription during pachytene (Alexander et al., 2023; Bolcun-Filas et al., 2011; Cecchini et al., 2023; Gaucher et al., 2012; Li et al., 2013; Maezawa et al., 2020; Manterola et al., 2018; Ozata et al., 2020; Yu et al., 2021). Additionally, extensive chromatin remodeling takes place in spermatocytes during meiotic prophase I, involving histone variant exchange, histone modifications, and high-order genome rearrangement (Kota and Feil, 2010; Wang et al.,

2017; Zheng and Xie, 2019). It has been known that SETDB1-mediated H3K9

trimethylation (H3K9me3) is required for sex chromosome transcription silencing (a

process called meiotic sex chromosome inactivation, MSCI) and male meiotic procession

(Hirota et al., 2018). However, the specific involvement of chromatin regulators in gene

activation on autosomes remains poorly understood.

Zinc finger HIT-type containing 1 (Znhit1), an evolutionarily conserved subunit of the

SRCAP chromatin remodeling complex, acts as a key regulator for the histone variant

H2A.Z deposition to control gene expression and cell fate determination (Cai et al., 2005;

Cuadrado et al., 2007; Feng et al., 2022). Recent studies have shown that Znhit1 is

involved in a wide range of developmental processes, including muscle and lens

differentiation, heart development, lung branching, and adult tissue stem cell maintenance

(Cuadrado et al., 2010; Lu et al., 2022; Sun et al., 2020; Wei et al., 2022; Xu et al., 2021;

Zhao et al., 2019). In our most recent study, we deleted Znhit1 in early male germ cells

and found that Znhit1 is required for the mitosis-to-meiosis transition by controlling

Meiosin expression (Sun et al., 2022). However, the in vivo function of Znhit1 in

regulating meiotic progression remains unknown.

To delve into the function of Znhit1 in meiosis, we analyzed Znhit1 expression and

observed upregulated Znhit1 expression specifically during meiotic prophase. Here, we

examine the role of Znhit1 in multiple meiotic events and show that the deletion of Znhit1

in spermatocytes causes impaired HR and defective PGA, thereby resulting in meiotic

developmental arrest. Hence, our study highlights the essential role of Znhit1 in ensuring

meiotic progression, implicating its potential as a therapeutic target for meiosis-related

diseases.

Results

***Znhit1* expression is upregulated during meiotic prophase and *Znhit1* knockout in**
spermatocytes disrupts spermatogenesis

To identify potential chromatin regulators in meiosis, we first queried for chromatin
factors highly expressed during the zygotene stage based on published transcription data
(Fig S1A and Table S1) (Chen et al., 2018). Quantitative analysis showed that the
chromatin remodeler Znhit1 was among the most highly expressed factors (Figs 1A, 1B,
S1B, and S1C). RNA in situ hybridization validated that *Znhit1* expression in male germ
cells decreased after meiotic initiation, followed by a gradual increase in spermatocytes
from the zygotene stage to the metaphase stage (Fig 1C). These results suggest that
Znhit1 is a potential chromatin regulator of meiotic progression.

To investigate whether Znhit1 regulates meiotic progression, we generated
spermatocyte-specific *Znhit1* knockout mice (*Znhit1*$^{fl/fl}$; *Stra8-cre*, referred to as
Znhit1-sKO). In the first wave of spermatogenesis, Ngn3-negative prospermatogonia
directly differentiate into A2 spermatogonia(Law et al., 2019; Rabbani et al., 2022;
Yoshida et al., 2006; Yoshida et al., 2004). With Stra8-cre inducing recombination in A1
spermatogonia and preleptotene spermatocytes separately(Lin et al., 2017), it is feasible
to generate mice with spermatocyte-specific gene knockout (Fig 1D). RNA in situ
experiments on testis sections confirmed *Znhit1* mRNA deletion in spermatocytes (Figs
1E and S1D). Compared with littermate controls, *Znhit1*-sKO male mice had smaller
testes (Fig S1E and S1F). PAS-histological and PNA-fluorescence staining illuminated
the absence of round spermatids and elongated spermatids in *Znhit1*-sKO testes (Fig 1F
and 1G), indicating that *Znhit1* knockout in spermatocytes disrupted spermatogenesis.

Znhit1 deletion leads to meiotic arrest at the pachytene stage

To dissect spermatocyte development in *Znhit1*-sKO testes, we co-stained SYCP3, a
marker of primary spermatocytes, and HSPA2, a testis-specific HSP70 family member
that accumulates from the pachytene stage onward. As shown in Figure 2A, both control
and *Znhit1*-sKO testis sections exhibited SYCP3^{+}HSPA2^{+} primary spermatocytes.
Phospho-histone H3 (Thr3) (pH3) is a marker of mitotic and meiotic metaphase cells, and
Znhit1 deletion caused the complete absence of pH3^{+} metaphase spermatocytes within the
cavity of seminiferous tubules (Fig 2B). These results suggest that Znhit1 deletion causes
spermatocyte arrest at meiotic prophase I.

To pinpoint the onset of meiotic failure, we performed immunostaining against SYCP3 in

spermatocyte chromosome spreads. The wild-type testes displayed typical spermatocytes

at consecutive substages, including leptotene, zygotene, pachytene, diplotene, and

diakinesis (Fig 2C). However, spermatocytes at the diplotene and diakinesis stages were

rarely observed in *Znhit1*-sKO testes (Fig 2D and 2E). We also examined the expression

of H1T, a middle pachytene marker, and no difference in H1T expression was observed in

Znhit1$^{-/-}$ spermatocytes (Fig S2A). Moreover, *Znhit1* knockout resulted in increased

TUNEL-positive spermatocytes, suggesting that these defective spermatocytes were

eliminated by the apoptotic pathway (Fig 2F and 2G). These results indicate that Znhit1

deletion leads to meiotic arrest at the pachytene stage.

Znhit1 is required for DSB repair and meiotic recombination

To investigate the impact of Znhit1 on meiotic events, we first performed immunostaining

of SYCP1 and SYCP3, the essential components of the synaptonemal complex, to

examine whether Znhit1 regulated homologous synapses. As shown in Figure 3A,

pachytene nuclei with fully synapsed autosomes were observed in spermatocytes from

control and *Znhit1*-sKO testes. During the pachytene stage, the X and Y chromosomes

undergo pairing and synapses between the short pseudoautosomal regions (PARs). We

found an increase in the percentage of unsynapsed X-Y chromosomes in *Znhit1*-sKO

spermatocytes (Fig 3B). Thus, deletion of Znhit1 results in impaired sex chromosomal

synapsis.

Next, we examined the programmed formation and repair of DSBs by co-staining SYCP3

and γH2AX, a DSB marker, on spermatocyte chromosome spreads. In both control and

Znhit1-sKO spermatocytes, γH2AX signals were evident in the leptotene and zygotene

stages (Fig 3C). However, in Znhit1-deficient pachynema, we still observed diffuse

γH2AX signals on the autosomes, while in control pachynema, γH2AX signals only

accumulated on the X-Y chromosomes. Moreover, we detected the presence of another

DSB marker, phospho-ATM (pATM), on the autosomes in *Znhit1*-sKO pachytene

spermatocytes (Fig 3D). Together, these results demonstrate that Znhit1 is required for

completing DSB repair.

To understand how Znhit1 regulates DSB repair, we performed an immunostaining

analysis using the markers RPA2 and DMC1, the essential factors involved in DSB repair

by coating resected DSBs. We found an increase in RPA2 and DMC1 counts in both

zygonema and pachynema of *Znhit1*-deficient testes (Fig 3E, 3F, and S3A), indicating

that Znhit1 deletion delayed recombinational repair, resulting in defective resolution of

autosomal DSBs in $Znhit1^{-/-}$ pachynema. The formation and repair of DSBs play crucial

roles in initiating and facilitating meiotic recombination, respectively. To examine the

consequences of unrepaired DSBs on meiotic recombination in Znhit1 mutants, we

performed an immunostaining analysis against MLH1, a marker of meiotic crossover

formation. Indeed, we found an almost absence of MLH1 foci in $Znhit1^{-/-}$ pachytene

spermatocytes compared with controls (Fig 3G and 3H), indicating that the loss of Znhit1

severely disrupted meiotic crossover formation.

Collectively, these results argue that Znhit1 is essential for multiple meiotic events,

including sexual chromosome synapsis, DSB repair, and homologous recombination.

scRNA-seq analysis reveals abnormal pachytene-to-diplotene transition caused by

***Znhit1* knockout**

We then asked how Znhit1 regulates spermatocyte development. We isolated testicular

cells from control and *Znhit1*-sKO mice at postnatal 16 (P16) and performed scRNA-seq

analysis. After quality control, a total of 2,4164 cells (12,212 cells from control, 11,952

cells from *Znhit1*-sKO) were captured, yielding 2,473 genes per cell. As visualized via

uniform manifold approximation and projection (UMAP), the single-cell atlas comprised

germ cells at various stages and major somatic cell types, including Sertoli cells ($Sox9^+$),

Leydig cells ($Hsd3b1^+$), peritubular myoid cells ($Acta2^+$), and macrophages ($Adgre1^+$)

(Figs 4A, 4B and S4A, and Table S2). The scRNA-seq data identified 6 germ cell types

based on the expression of specific markers, including undifferentiated spermatogonia

($Zbtb16^+$), differentiating spermatogonia (Kit^+), preleptotene spermatocytes ($Nacad^+$),

leptotene spermatocytes ($Meiob^+$), zygotene spermatocytes (Gml^+), and pachytene

spermatocytes ($Ldhc^+$) (Fig 4B and 4C). Examination of germ cell scRNA-seq profiles

via unsupervised clustering revealed decreased pachytene spermatocytes after *Znhit1*

deletion (Figs 4D, 4E, and S4B). *Pou5f2* is expressed explicitly at the diplotene stage, and

we found a lack of $Pou5f2^+$ diplotene spermatocytes in *Znhit1*-sKO testes (Fig 4F). These

results indicate that Znhit1 is required for male germ cell development beyond the

pachytene stage.

Znhit1 deletion impairs meiotic transcriptional activation

Next, we asked whether meiotic transcriptional activation was affected by Znhit1 deletion. We analyzed changes in gene expression between consecutive spermatocyte stages and identified 4,603 genes that showed significant changes (Fig 5A and Table S3). Of the changed genes between zygotene spermatocytes and pachytene spermatocytes, 1,560 genes were upregulated (referred to as pachytene-activated genes or PGA genes). Moreover, we identified differentially expressed genes (DEGs) in each spermatocyte type between the control and *Znhit1*-sKO groups. The number of DEGs gradually increased from the preleptotene stage to the pachytene stage, with downregulated genes being predominant (Fig 5B and Table S4). In particular, $Znhit1^{-/-}$ pachytene spermatocytes exhibited 2,785 downregulated genes, among which were most of the pachytene-activated genes (1,094 out of 1,560, 70.1%; Figs 5B-5D, and S5A). Gene Ontology (GO) enrichment analysis revealed that upregulated genes following Znhit1 deletion were associated with apoptotic pathways, supporting an observation that Znhit1 deletion resulted in germ cell clearance through apoptosis. Moreover, downregulated GO terms included germ cell development and DNA recombination (Fig 5E and Table S5). Interestingly, recent studies showed that cilium organization was essential for spermatocyte development (Mytlis et al., 2022), and Znhit1 deletion reduced the expression of cilium-related genes in zygotene and pachytene spermatocytes (Fig 5E).

We further conducted bulk RNA-seq experiments with P14 control and *Znhit1*-sKO testicular cells and identified 882 DEGs (812 downregulated genes and 70 upregulated genes) in *Znhit1*-sKO testicular cells (Figs 5F and S6A, and Table S6). Through integrative analysis of bulk and single-cell RNA-seq data, we found that the expression of these 812 genes was activated during the zygotene-to-pachytene transition and Znhit1 deletion significantly reduced the expression of these genes (Figs 5G, 5H, and S5B). The downregulated genes following Znhit1 loss were enriched for germ cell-related GO terms, such as cilium formation and spermatid development (Fig S6B). Together, these data argue that Znhit1 is required for meiotic transcriptional activation.

Consistent with meiotic recombination defects in Znhit1 mutants, gene set enrichment analysis (GSEA) revealed that the expression of HR-related genes was globally downregulated following Znhit1 loss (Fig 5I and Table S7). The top genes associated with this function included *Ccnb1ip1*, *Rnf212*, *Spo16*, *Ankrd31*, and *Terb1* (Boekhout et al., 2019; Papanikos et al., 2019; Qiao et al., 2014; Rao et al., 2017; Reynolds et al., 2013; Shibuya et al., 2014; Wang et al., 2019; Zhang et al., 2019) (Figs 5J, S6C, and S6D). As

231 expected, the Rnf212-dependent sumoylation level decreased in *Znhit1*$^{-/-}$ spermatocytes

232 (Fig 5K). These results suggest that Znhit1 regulates meiotic recombination by activating

233 HR-related genes.

Znhit1 deletion impairs H2A.Z incorporation into pachytene chromatin

We then asked how Znhit1 regulates transcription. Znhit1 is a subunit of the SRCAP

complex that facilitates the incorporation of the histone variant H2A.Z by replacing H2A

(Fig S7A). Immunostaining against H2A.Z showed a gradual increase in H2A.Z

accumulation on autosomes from the leptotene to the diakinesis stage, while the H2A.Z

signal was markedly low on the X-Y body (Fig 6A). Znhit1 loss decreased H2A.Z

staining on the pachytene cells, suggesting that Znhit1 is required for maintaining H2A.Z

integrity during meiosis.

To probe the H2A.Z targets, we performed chromatin immunoprecipitation followed by

sequencing (ChIP-seq) against H2A.Z in P14 testicular cells. Using the ChromHMM

model to annotate H2A.Z signals with meiocyte chromatin states defined by published

epigenetic markers (Spruce et al., 2020), we found that H2A.Z sites were primarily

enriched in chromatin state 1 and state 2 (promoters and enhancers) but were less

enriched in recombination hotspots (Fig 6B and 6C), consistent with previously published

results (Spruce et al., 2020). Quantitative analysis showed significant H2A.Z binding

reduction at 37.9% (23,926 of 63,109) of H2A.Z-bound genome sites following Znhit1

deletion (Fig 6D). Comparison analysis with bulk RNA-seq data identified 480

downregulated DEGs with decreased H2A.Z signals (Fig 6E). ~41.3 and 37.8% of the

downregulated H2A.Z peaks occupied promoters and enhancers, respectively. At

promoters, these downregulated H2A.Z preferentially occupied active promoters

(H3K4me3 only), while H2A.Z occupied three types of enhancers: active, poised, and

primed enhancers (Figs 6F, S7B, and S7C). These findings indicate that Znhit1 deletion

impairs H2A.Z incorporation into pachytene chromatin.

H2A.Z deposition cooperates with A-MYB to regulate transcription

We further asked how Znhit1/H2A.Z deposition facilitates lineage-specific gene

expression. We performed TF motif analysis using active promoters/enhancers with

decreased H2A.Z signals. This analysis identified several MYB family TFs, including

MYB, A-MYB, and B-MYB (Fig 7A). Transcriptional network analysis conducted by

SCENIC further identified A-MYB (encoded by *Mybl1* gene) as the core regulatory TF in

spermatocytes (Fig 7B). We reanalyzed RNA-seq data in P14 testes from

A-MYB-deficient mice (Li et al., 2013) and found that A-MYB-deficient DEGs

correlated with those observed in Znhit1-deficient testicular cells (Fig 7C). Although

Znhit1 deletion didn't affect *Mybl1* mRNA expression, transcriptome analyses showed

that the expression of A-MYB target genes was significantly reduced in *Znhit1*$^{-/-}$

pachytene spermatocytes (Figs 7D, 7E, S8A, and S8B). These results suggest that *Znhit1*$^{-}$

$^{/-}$ and *Mybl1*$^{-/-}$ pachytene spermatocytes share common gene expression alterations.

Next, we compared the genome binding sites of H2A.Z and A-MYB. Utilizing published

A-MYB ChIP-seq data from P14 testicular cells (Li et al., 2013), we identified 6,088

A-MYB binding signals, with 78.1% (4755) of these coinciding with H2A.Z peaks (Fig

7F). Annotation of genomic features showed that approximately 66.52% and 12.02% of

these overlapping sites occupied active promoters and enhancers, respectively,

representing a stronger enrichment compared to other regulatory elements (Fig 7G).

Genes associated with H2A.Z and A-MYB binding included essential HR-related genes,

such as *Ccnb1ip1* (encode the Hei10 protein) (Fig 7H). Moreover, we showed that Znhit1

deletion resulted in a marked reduction in H2A.Z signals and nascent transcription

identified by KAS-seq signals (Fig 7I and 7J).

Previous studies have shown the fundamental role of A-MYB in activating meiotic

enhancers (Maezawa et al., 2020). Enhancer activity is linked to an increased formation

of enhancer single-stranded DNA (ssDNA), which can be evaluated using KAS-seq (Wu

et al., 2020). We found that the removal of Znhit1 resulted in abolished ssDNA at active

enhancers (Fig 7I and 7J). Interestingly, the chromatin accessibility detected by

ATAC-seq at active enhancers increased after Znhit1 deletion. These findings suggest

that Znhit1/H2A.Z deposition is essential for enhancer activation.

Together, these results demonstrate the critical role of the Znhit1/H2A.Z/A-MYB axis in

governing transcriptional activation and enhancer activity.

Discussion

Meiosis plays a crucial role in the production of haploid gametes. Any disturbances

during meiotic progress are known to lead to infertility and congenital diseases. Here we

delineate the multifaceted functions of the chromatin remodeler Znhit1 in regulating

meiotic progression. Our study provides convincing data to show that the

Znhit1/H2A.Z/A-MYB axis controls spermatocyte development, meiotic recombination,

and PGA. Thus, these findings underscore the significance of specific chromatin

structures in governing appropriate transcriptional reprogramming and precise meiotic

recombination.

The role of H2A.Z in DNA injury and repair has been extensively studied

(Colino-Sanguino et al., 2022; Dong et al., 2014; Xu et al., 2012; Yamada et al., 2018),

but its function in programmed DSB formation and repair during mammalian meiosis

remains unknown. To address this question, we generated Znhit1 conditional knockout

mice specifically during male meiosis to examine the role of Znhit1-mediated H2A.Z

deposition in the formation and repair of programmed DSB. One new observation in this

study is that Znhit1 loss is dispensable for DSB formation but necessary for DNA break

repair during meiosis. Intriguingly, unlike the observations in plant meiosis, we found that

H2A.Z does not directly bind DNA recombination sites in mammalian spermatocytes

(Choi et al., 2013; Spruce et al., 2020; Wang and Copenhaver, 2018). Moreover, we

showed that Znhit1 deletion delays recombinational repair, but has limited impacts on

DNA resection, ultimately causing defective meiotic crossover formation. Importantly,

our study demonstrates that Znhit1 is essential for the expression of meiotic

recombination-related genes, such as *Ccnb1ip1* and *Rnf212*, implying that Znhit1/H2A.Z

modulates meiotic recombination through transcriptional regulation. Therefore, our data

reveals a different mechanism underlying meiotic recombination between plants and

mammals.

Pachytene spermatocytes utilize a highly specific mechanism of meiotic surveillance, the

pachytene checkpoint, to prevent aneuploidy formation by removing abnormal germ cells

with incomplete chromosome synapsis and defective homologous recombination (Huang

and Roig, 2023; Roeder and Bailis, 2000; Subramanian and Hochwagen, 2014). Two

checkpoint pathways exist in male meiosis: one responding to unrepaired DSBs and the

other triggered by MSCI failure (Royo et al., 2013). However, whether epigenetic

regulation is involved in meiotic surveillance is not well defined. Previous studies have

reported that upon stimulation by unrepaired DSBs, the MRE11-ATM-CHK2 pathway is

activated to eliminate aberrant germ cells (Marcet-Ortega et al., 2017). Our study found

that Znhit1 deletion impairs the removal of ATM phosphorylation signals on autosomes

in pachynema, ultimately causing meiotic pachytene arrest and apoptosis. Therefore,

these findings demonstrate that Znhit1 acts as an important chromatin factor involved in

the regulation of the pachytene checkpoint.

It has long been noticed that in meiotic prophase I, chromatin that has not yet completed

chromosomal synapses is transcriptionally inactive, known as meiotic silencing of

unsynapsed chromatin (MSUC) (Turner et al., 2005). When germ cells enter the

pachytene stage, a large number of protein-coding genes and non-coding RNAs begin to

be actively expressed, known as PGA. Previous studies have shown that transcription

factors A-MYB and BRDT are involved in transcriptional activation during meiotic

prophase, but how PGA takes place in the chromatin context is unclear. In this study, we

found that the expression of the chromatin remodeler Znhit1 is specifically upregulated

during the zygotene-to-pachytene transition. We also observed that H2A.Z deposition is

enriched in the autosomes but less in the sex chromosome at the pachytene stage. Znhit1

deletion in spermatocytes repressed pachytene gene activation globally and reduced

chromosome-wide H2A.Z deposition, supporting a central role of Znhti1 in PGA

regulation and chromatin remodeling. It has been known that A-MYB is a transcription

factor that controls pachytene transcription activation. We found that Znhit1-deficient

spermatocytes phenocopied abnormal meiotic phenotypes in A-MYB mutants, such as

X-Y synapsis failure, impaired DSB repair, and defective HR (Alexander et al., 2023;

Bolcun-Filas et al., 2011; Li et al., 2013; Maezawa et al., 2020). Our results also revealed

that Znhit1/H2A.Z cooperates with A-MYB to regulate PGA gene activation. Therefore,

our study illuminates the molecular mechanisms underlying the fundamental question of

how PGA is regulated.

The histone variant H2A.Z is enriched at gene promoters and regulatory regions, but there

is ongoing debate regarding its involvement in transcriptional regulation. A recent paper

reported that H2A.Z knockout in post-mitotic muscle cells has limited effects on gene

expression (Belotti et al., 2020). In this study, we utilized the meiotic prophase as a model

system (where DNA replication does not occur) to study H2A.Z's function in

transcriptional regulation. We found that Znhit1-mediated H2A.Z deposition, independent

of DNA replication, is indispensable for the transcriptional activation of a large number

of meiotic genes. One explanation of this conflicting phenomenon is that H2A.Z

dynamics but not stable H2A.Z accumulation is essential for priming transcriptional

changes.

Through the integration of functional and molecular evidence, our findings establish the

critical involvement of Znhit1-dependent chromatin remodeling in the orchestration of

meiotic progression and coordination of various meiotic processes, such as HR and PGA.

Furthermore, we pinpoint the pivotal role played by the Znhit1/H2A.Z/A-MYB axis in

driving transcriptional reprogramming during meiotic prophase. Taken together, this

study deepens our understanding of the interplay between epigenetic regulation and

mammalian meiosis.

Materials and Methods

Animals. *Znhit1*$^{fl/fl}$ mice have been previously described (Zhao et al., 2019) and are available from the Model Animal Research Center of Nanjing University (MARC, Nanjing, China). The *Stra8-cre* knock-in mouse line was kindly provided by Dr. Ming-Han Tong(Lin et al., 2017). All mice were maintained on the C57BL/6J background. Germ cell-specific *Znhit1* knockout mice (*Znhit1*$^{fl/fl}$; *Stra8-cre*) were obtained by crossing *Znhit1*$^{fl/+}$; *Stra8-cre* mice with *Znhit1*$^{fl/fl}$ mice. All mice were housed in the SPF (Specific-Pathogen-Free) animal facility with standard 12 h light/dark cycles and standard temperature (22 to 24°C). All mice were provided with ad libitum access to standard laboratory food and water. All experiments in this study were performed following relevant guidelines and approved by the Animal Care and Use Committee of Fudan University.

Histological and immunohistochemical analysis. Testes were fixed in modified Davidson's fixative as previously described (Latendresse et al., 2002), embedded in paraffin, and sectioned. For periodic acid-Schiff (PAS)-hematoxylin staining, 5 μm testis sections were deparaffinized, rehydrated, and stained with Schiff's reagent and hematoxylin solution. For immunofluorescent staining, 5 μm testis sections were retrieved by sodium citrate antigen retrieval buffer (pH 6.0) and blocked with 5% BSA in PBS for 30 min at room temperature. The sections were later incubated overnight at 4°C with primary antibodies as follows: mouse anti-SYCP3 (Abcam, ab97672), rabbit anti-HSPA2 (Abcam, ab108416), rabbit anti-pH3 (Millipore, H0412), rabbit anti-H1T (Invitrogen, PA5-51200), or lectin PNA (Sigma–Aldrich, L7381). On the following day, secondary fluorescein-conjugated antibodies and DAPI (Sigma–Aldrich, D9542) were added for 1 h, followed by Fluoromount-G mounting (Southern Biotech, 0100-01). Images were analyzed using the confocal microscope.

TUNEL staining was carried out using the DeadEnd^{TM} Fluorometric TUNEL System (Promega, G3250) according to the manufacturer's instructions.

Immunostaining of spermatocyte chromosome spreads. Spermatocyte chromosome spreads were prepared as previously described (Alavattam et al., 2018). Briefly, the clumps of seminiferous tubules were transferred to the hypotonic extraction buffer (30 mM Tris base, 17 mM trisodium citrate, 5 mM EDTA, 50 mM sucrose, 5 mM dithiothreitol (DTT), and 1× protease inhibitor, pH 8.2) and incubated on ice for 1.5 h. The clumps of tubules were then transferred to 30 μL of ice-cold 100 mM sucrose and mashed gently using tweezers to obtain a cell suspension. An additional 30 μL of ice-cold 100 mM sucrose was added to the cell suspension and mixed several times. Positively charged slides were incubated in the ice-cold fixation solution (2% paraformaldehyde, 0.1% Triton X-100, and 0.02% SDS, pH 9.2) for 3 min. 30 μL of the diluted cell suspension was applied to the slides and incubated in humid chambers at room temperature for 2 h. The slides were washed with 0.4% Photo-Flo 200 and stored at -80°C.

For immunostaining of spermatocyte chromosome spreads, slides were washed with PBS and blocked with 5% BSA in PBS for 30 min at room temperature. The slides were stained with primary antibodies as follows: rabbit anti-SYCP3 (Abcam, ab15093), mouse anti-SYCP3 (Abcam, ab97672), rabbit anti-SYCP1 (Abcam, ab15090), mouse anti-γH2AX (BioLegend, 613401), rabbit anti-pATM (Millipore, 05-740), rabbit anti-RAD51 (Proteintech), rabbit anti-RPA2 (Proteintech, 10412-1-AP), rabbit anti-MLH1 (Cell Signaling Technology, 3515T), rabbit anti-H2A.Z (Abcam, ab4174).

***Znhit1* in situ hybridization.** Testis sections (5 μM) at the indicated times were prepared for *Znhit1* in situ hybridization with the RNAscope kit (Advanced Cell Diagnostics, 323100) according to the manufacturer's instructions.

Quantitative reverse transcription PCR (RT-qPCR). Total RNA was isolated using the RNeasy Mini-plus Kit (Qiagen, 74134) and then reverse-transcribed into complementary DNA (cDNA) with the GoScript Reverse Transcription System (Promega, A5003). cDNA was used as the template for the quantitative PCR assay using 2×SYBR Green qPCR Master Mix (Bimake, B21202). Quantitative PCR primers are listed in Table S8.

RNA-seq library generation and sequencing. Total RNA from fresh testes was isolated
and mRNA was purified with magnetic beads (Vazyme, N401-01). Then, mRNA was
fragmented and processed to generate RNA-seq libraries (Vazyme, NR605-01). Over 40
million reads were obtained per sample using the Illumina NovaSeq platform for 2
independent biological replicates.

Single-cell RNA sequencing (scRNA-seq). P16 control and *Znhit1*-sKO testes were
digested by collagenase IV and trypsin at 37°C for 10 min to obtain testicular cell
suspensions. scRNA-seq libraries were constructed using a $10 \times$ Genomics kit and
sequenced on the Illumina platform.

ChIP-seq library generation and sequencing. ChIP-seq library generation was
performed as previously described. Briefly, P14 control and *Znhit1*-sKO testes were
digested by collagenase IV and trypsin at 37°C for 10 min, crosslinked with 1%
formaldehyde for 10 min, and quenched with glycine. The cells were lysed and sheared
with a Bioruptor Plus machine for 20 min. Then, 2 μg of anti-H2A.Z antibody (Abcam,
ab4174) was added to the sonicated chromatin and incubated overnight at 4°C. The
following day, 20 μL of protein G beads were added and incubated for 2 h at 4°C. The
beads were washed and reverse-crosslinked for 4 h at 65°C. DNA was extracted using
phenol-chloroform and subjected to library construction using the VAHTS Universal
DNA Library Prep Kit for Illumina (Vazyme, ND607-01) according to the manufacturer's
instructions. Over 30 million reads were obtained per sample using the Illumina NovaSeq
platform for 2 independent biological replicates.

KAS-seq library and ATAC-seq library generation and sequencing. Testes were
digested by collagenase IV and trypsin at 37°C for 10 min. Pachytene cells were sorted
using fluorescence-activated cell sorting (FACS) as previously described (Long et al.,
2017). For KAS-seq, pachytene cells were labeled with N3-kethoxal, and DNA was
isolated using the DNA Clean and Concentrator kit (Zymo, D4013) and subjected to
library construction using Q5 high-fidelity DNA polymerase (New England Biolabs,
M0544S). Genomic DNA was fragmented using Tn5 transposase and subjected to library

construction. For ATAC-seq, pachytene nuclei were isolated and fragmented with Tn5

transposase. DNA was isolated using the DNA Clean and Concentrator kit (Zymo, D4013)

and subjected to library construction using Q5 high-fidelity DNA polymerase (New

England Biolabs, M0544S). For high-throughput sequencing, over 30 million reads were

obtained per sample using the Illumina NovaSeq platform for 2 independent biological

replicates.

RNA-seq data analysis. We performed RNA-seq analysis as described previously

(Pertea et al., 2016). Briefly, after removing adapters using Cutadapt (v2.5) (Kechin et al.,

2017), paired-end reads were aligned to the annotated mouse transcripts (mm10 Gencode

vM23 release) using Hisat2 (v2.2.1) (Kim et al., 2015; Kim et al., 2019). Gene expression

levels were calculated using StringTie (v2.2.1) (Pertea et al., 2015). Read counts were

calculated using a Python script (prepDE.py3) provided by the StringTie development

team. Differentially expressed genes were identified using the R package DESeq2

(v1.38.3) (Love et al., 2014). Genes with read counts > 50 in at least one sample were

kept for further analysis. A given gene was considered to significantly change if the

adjusted P value (padj) was < 0.05, the P value was < 0.01, and the fold-change was ≥ 2.

Gene Set Enrichment Analysis (GSEA) was carried out using GSEA software

(Subramanian et al., 2005). GO analysis was performed using clusterProfiler (v4.6.2) (Wu

et al., 2021).

scRNA-seq data analysis. FASTQ files were run through CellRanger (v7.1.0) software

with default parameters for de-multiplexing, aligning reads with STAR software to mm10,

and counting unique molecular identifiers (UMIs). As input files of the Seurat R package

(v4.4.0), the filtered gene expression matrices were then used for downstream analyses

(Butler et al., 2018). Low-quality cells were filtered (expressing < 500 or >6,000 unique

gene counts and >15% mitochondrial reads). Principal component analysis was performed

on SCT-transformed data using 3,000 variable genes. The top 50 principal components

were used for clustering and visualized using the UMAP algorithm in the Seurat R

package. The "FindAllMarkers" function of the Seurat R package was used to calculate

cluster-specific genes. Marker genes for each cluster are shown in Table S2.

The "FindMarkers" function of the Seurat R package was used to identify differentially

expressed genes (DEGs) for spermatocytes (preleptotene, leptotene, zygotene, and

pachytene). Only those with |'avg_logFC'| > 0.25 and 'p_val_adj' < 0.05 were considered

as DEGs. For the transcriptional regulatory network analysis, the raw count matrix was

used by the pyscenic (v0.12.1) workflow using default parameters (Aibar et al., 2017).

Then, the output loom file was opened by the SCENIC R package (v1.3.1), and AUC

values of regulons were extracted for visualization of downstream transcription factor

activity.

ChIP-seq and KAS-seq data analysis. For analyzing ChIP-seq and KAS-seq data, we

used the ENCODE ChIP-seq pipeline (v2.2.1,

https://github.com/ENCODE-DCC/chip-seq-pipeline2). Raw reads were cleaned by

Cutadapt (v2.5) (Kechin et al., 2017). After removing adapters, clean reads were aligned

against the mouse mm10 genome using bwa (v0.7.17) (Li and Durbin, 2009). Then, sam

files were converted to bam files using samtools (v1.9) (Li et al., 2009). PCR duplicates

were removed using Picard (v2.20.7) (https://broadinstitute.github.io/picard/). Histone

ChIP-seq peaks (such as H2A.Z and histone marks) and KAS-seq peaks were called using

MACS2 (v2.2.4) (Zhang et al., 2008), while transcription factor ChIP-seq peaks (such as

A-Myb) were called using the R packages spp (v1.15.5)

(https://github.com/hms-dbmi/spp). The conservative narrow peaks were used for

downstream analysis.

For visualization, bam files were converted to bigWig files using deepTools (v3.3.1)

(Ramirez et al., 2014), and bigWig files were used to calculate tag density under 50 bp

resolution. In addition, CPM was used to normalize the number of reads per bin. We used

the R package DiffBind (v3.8.4)

(http://bioconductor.org/packages/release/bioc/html/DiffBind.html) for differential peak

analysis and the R package rGREAT (v2.1.11) (Gu and Hubschmann, 2023) for peak

annotation. We used ComputeMatrix and plotHeatmap of deepTools to generate count

matrices and heatmaps, respectively. H2A.Z enrichment within chromatin states was

annotated using the ChromHMM model (Ernst and Kellis, 2012; Spruce et al., 2020). TF

motif enrichment was calculated using HOMER (v4.11) (Heinz et al., 2010). We used

deepTools to draw Pearson's correlation scatter plots for the correlation between

replicates of each experiment.

For promoter and enhancer type analysis, H3K27ac ChIP-seq data in wild-type pachytene

spermatocytes were downloaded from the published data (GSE107398) (Adams et al.,

2018); ChIP-seq data for H3K4me1, H3K4me3, and H3K27me3 in wild-type pachytene

spermatocytes were downloaded from the published data (GSE132446) (Chen et al.,

2020).

ATAC-seq data analysis. ATAC-seq data were analyzed using a standardized ENCODE

ATAC-seq pipeline (v2.2.2, https://github.com/ENCODE-DCC/atac-seq-pipeline). The

mouse reference genome (mm10) was used in the pipeline. Trimming, mapping, and

duplicate-removing were performed as the ENCODE pipeline suggested using Cutadapt

(v2.5) (Kechin et al., 2017), Bowtie2 (v2.3.4.3) (Langmead et al., 2009), and Picard

(v2.20.7) (https://broadinstitute.github.io/picard/) respectively. MACS2 (v2.2.4) (Zhang

et al., 2008) was used for peak calling.

Statistics and reproducibility. Statistical analyses were performed using R and Prism 9.

Data are presented as means ± s.d. unless otherwise indicated. The two-tailed, unpaired,

Student's t-test, Mann–Whitney test, or one-way or two-way ANOVA were performed to

analyze statistical significance.

Supplementary Materials

This manuscript file includes Figs. S1 to S9 and Tables S1 and S8.

Reference

Adams, S.R., Maezawa, S., Alavattam, K.G., Abe, H., Sakashita, A., Shroder, M., Broering,

T.J., Sroga Rios, J., Thomas, M.A., Lin, X*., et al.* (2018). RNF8 and SCML2 cooperate to

regulate ubiquitination and H3K27 acetylation for escape gene activation on the sex

chromosomes. PLoS genetics *14*, e1007233.

Aibar, S., Gonzalez-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H., Hulselmans, G.,

Rambow, F., Marine, J.C., Geurts, P., Aerts, J*., et al.* (2017). SCENIC: single-cell regulatory

network inference and clustering. Nature methods *14*, 1083-1086.

Alavattam, K.G., Abe, H., Sakashita, A., and Namekawa, S.H. (2018). Chromosome Spread

Analyses of Meiotic Sex Chromosome Inactivation. Methods in molecular biology (Clifton, NJ)

1861, 113-129.

Alexander, A.K., Rice, E.J., Lujic, J., Simon, L.E., Tanis, S., Barshad, G., Zhu, L., Lama, J.,

Cohen, P.E., and Danko, C.G. (2023). A-MYB and BRDT-dependent RNA Polymerase II

pause release orchestrates transcriptional regulation in mammalian meiosis. Nature

communications *14*.

Baudat, F., Imai, Y., and de Massy, B. (2013). Meiotic recombination in mammals: localization

and regulation. Nature reviews Genetics *14*, 794-806.

Belotti, E., Lacoste, N., Simonet, T., Papin, C., Padmanabhan, K., Scionti, I., Gangloff, Y.G.,

Ramos, L., Dalkara, D., Hamiche, A*., et al.* (2020). H2A.Z is dispensable for both basal and

activated transcription in post-mitotic mouse muscles. Nucleic acids research *48*, 4601-4613.

Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P.C., Nicolas, A., and Forterre, P. (1997).

An atypical topoisomerase II from Archaea with implications for meiotic recombination. Nature

386, 414-417.

Boekhout, M., Karasu, M.E., Wang, J., Acquaviva, L., Pratto, F., Brick, K., Eng, D.Y., Xu, J.,

Camerini-Otero, R.D., Patel, D.J., *et al.* (2019). REC114 Partner ANKRD31 Controls Number,

Timing, and Location of Meiotic DNA Breaks. Molecular cell *74*, 1053-1068 e1058.

Bolcun-Filas, E., Bannister, L.A., Barash, A., Schimenti, K.J., Hartford, S.A., Eppig, J.J.,

Handel, M.A., Shen, L., and Schimenti, J.C. (2011). A-MYB (MYBL1) transcription factor is a

master regulator of male meiosis. Development *138*, 3319-3330.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nature

biotechnology *36*, 411-420.

Cai, Y., Jin, J., Florens, L., Swanson, S.K., Kusch, T., Li, B., Workman, J.L., Washburn, M.P.,

Conaway, R.C., and Conaway, J.W. (2005). The mammalian YL1 protein is a shared subunit

of the TRRAP/TIP60 histone acetyltransferase and SRCAP complexes. The Journal of

biological chemistry *280*, 13665-13670.

Cecchini, K., Biasini, A., Yu, T., Saflund, M., Mou, H., Arif, A., Eghbali, A., Colpan, C.,

Gainetdinov, I., de Rooij, D.G., *et al.* (2023). The transcription factor TCFL5 responds to

A-MYB to elaborate the male meiotic program in mice. Reproduction *165*, 183-196.

Chen, Y., Lyu, R.T., Rong, B.W., Zheng, Y.X., Lin, Z., Dai, R.F., Zhang, X., Xie, N.N., Wang,

S.Q., Tang, F.C., *et al.* (2020). Refined spatial temporal epigenomic profiling reveals intrinsic

connection between PRDM9-mediated H3K4me3 and the fate of double-stranded breaks. Cell

research *30*, 256-268.

Chen, Y., Zheng, Y., Gao, Y., Lin, Z., Yang, S., Wang, T., Wang, Q., Xie, N., Hua, R., Liu, M.,

et al. (2018). Single-cell RNA-seq uncovers dynamic processes and critical regulators in

mouse spermatogenesis. Cell research *28*, 879-896.

Choi, K., Zhao, X., Kelly, K.A., Venn, O., Higgins, J.D., Yelina, N.E., Hardcastle, T.J.,

Ziolkowski, P.A., Copenhaver, G.P., Franklin, F.C.*, et al.* (2013). Arabidopsis meiotic

crossover hot spots overlap with H2A.Z nucleosomes at gene promoters. Nature genetics *45*,

1327-1336.

Colino-Sanguino, Y., Clark, S.J., and Valdes-Mora, F. (2022). The H2A.Z-nuclesome code in

mammals: emerging functions. Trends in genetics : TIG *38*, 273-289.

Cuadrado, A., Corrado, N., Perdiguero, E., Lafarga, V., Munoz-Canoves, P., and Nebreda,

A.R. (2010). Essential role of p18Hamlet/SRCAP-mediated histone H2A.Z chromatin

incorporation in muscle differentiation. The EMBO journal *29*, 2014-2025.

Cuadrado, A., Lafarga, V., Cheung, P.C., Dolado, I., Llanos, S., Cohen, P., and Nebreda, A.R.

(2007). A new p38 MAP kinase-regulated transcriptional coactivator that stimulates

p53-dependent apoptosis. The EMBO journal *26*, 2115-2126.

Dong, S., Han, J., Chen, H., Liu, T., Huen, M.S., Yang, Y., Guo, C., and Huang, J. (2014). The

Human SRCAP Chromatin Remodeling Complex Promotes DNA-End Resection. Current

biology : CB *24*, 2097-2110.

Ernst, C., Eling, N., Martinez-Jimenez, C.P., Marioni, J.C., and Odom, D.T. (2019). Staged

developmental mapping and X chromosome transcriptional dynamics during mouse

spermatogenesis. Nature communications *10*, 1251.

Ernst, J., and Kellis, M. (2012). ChromHMM: automating chromatin-state discovery and

characterization. Nature methods *9*, 215-216.

Feng, Y., Zhang, Y., Lin, Z., Ye, X., Lin, X., Lv, L., Lin, Y., Sun, S., Qi, Y., and Lin, X. (2022).

Chromatin remodeler Dmp18 regulates apoptosis by controlling H2Av incorporation in

Drosophila imaginal disc development. PLoS genetics *18*, e1010395.

Gaucher, J., Boussouar, F., Montellier, E., Curtet, S., Buchou, T., Bertrand, S., Hery, P.,

Jounier, S., Depaux, A., Vitte, A.L.*, et al.* (2012). Bromodomain-dependent stage-specific male

genome programming by Brdt. The EMBO journal *31*, 3809-3820.

Gray, S., and Cohen, P.E. (2016). Control of Meiotic Crossovers: From Double-Strand Break

Formation to Designation. Annu Rev Genet *50*, 175-210.

Green, C.D., Ma, Q., Manske, G.L., Shami, A.N., Zheng, X., Marini, S., Moritz, L., Sultan, C.,

Gurczynski, S.J., Moore, B.B.*, et al.* (2018). A Comprehensive Roadmap of Murine

Spermatogenesis Defined by Single-Cell RNA-Seq. Developmental cell *46*, 651-667 e610.

Gu, Z., and Hubschmann, D. (2023). rGREAT: an R/bioconductor package for functional

enrichment on genomic regions. Bioinformatics *39*.

Handel, M.A., and Schimenti, J.C. (2010). Genetics of mammalian meiosis: regulation,

dynamics and impact on fertility. Nature reviews Genetics *11*, 124-136.

Hassold, T., and Hunt, P. (2001). To err (meiotically) is human: the genesis of human

aneuploidy. Nature reviews Genetics *2*, 280-291.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,

Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell identities. Molecular

cell *38*, 576-589.

Hirota, T., Blakeley, P., Sangrithi, M.N., Mahadevaiah, S.K., Encheva, V., Snijders, A.P.,

Ellnati, E., Ojarikre, O.A., de Rooij, D.G., Niakan, K.K.*, et al.* (2018). SETDB1 Links the Meiotic

DNA Damage Response to Sex Chromosome Silencing in Mice. Developmental cell *47*,

645-+.

Huang, Y., and Roig, I. (2023). Genetic control of meiosis surveillance mechanisms in

mammals. Front Cell Dev Biol *11*, 1127440.

Hunter, N. (2015). Meiotic Recombination: The Essence of Heredity. Cold Spring Harbor

perspectives in biology *7*.

Kechin, A., Boyarskikh, U., Kel, A., and Filipenko, M. (2017). cutPrimers: A New Tool for

Accurate Cutting of Primers from Reads of Targeted Next Generation Sequencing. J Comput

Biol *24*, 1138-1143.

Keeney, S., Giroux, C.N., and Kleckner, N. (1997). Meiosis-specific DNA double-strand breaks

are catalyzed by Spo11, a member of a widely conserved protein family. Cell *88*, 375-384.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low

memory requirements. Nature methods *12*, 357-360.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology *37*,

907-915.

Kota, S.K., and Feil, R. (2010). Epigenetic transitions in germ cell development and meiosis.

Developmental cell *19*, 675-686.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome biology *10*, R25.

Lascarez-Lagunas, L., Martinez-Garcia, M., and Colaiacovo, M. (2020). SnapShot: Meiosis -

Prophase I. Cell *181*, 1442-1442 e1441.

Latendresse, J.R., Warbrittion, A.R., Jonassen, H., and Creasy, D.M. (2002). Fixation of testes

and eyes using a modified Davidson's fluid: comparison with Bouin's fluid and conventional

Davidson's fluid. Toxicol Pathol *30*, 524-533.

Law, N.C., Oatley, M.J., and Oatley, J.M. (2019). Developmental kinetics and transcriptome

dynamics of stem cell specification in the spermatogenic lineage. Nature communications *10*,

2787.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinformatics *25*, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map

format and SAMtools. Bioinformatics *25*, 2078-2079.

Li, X.Z., Roy, C.K., Dong, X., Bolcun-Filas, E., Wang, J., Han, B.W., Xu, J., Moore, M.J.,

Schimenti, J.C., Weng, Z.*, et al.* (2013). An ancient transcription factor initiates the burst of

piRNA production during early meiosis in mouse testes. Molecular cell *50*, 67-81.

Lin, Z., Hsu, P.J., Xing, X., Fang, J., Lu, Z., Zou, Q., Zhang, K.J., Zhang, X., Zhou, Y., Zhang,

T.*, et al.* (2017). Mettl3-/Mettl14-mediated mRNA N6-methyladenosine modulates murine

spermatogenesis. Cell research.

Long, J., Huang, C., Chen, Y., Zhang, Y., Shi, S., Wu, L., Liu, Y., Liu, C., Wu, J., and Lei, M.

(2017). Telomeric TERB1-TRF1 interaction is crucial for male meiosis. Nat Struct Mol Biol *24*,

1073-1080.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome biology *15*, 550.

Lu, J., An, J., Wang, J., Cao, X., Cao, Y., Huang, C., Jiao, S., Yan, D., Lin, X., and Zhou, X.

(2022). Znhit1 Regulates p21Cip1 to Control Mouse Lens Differentiation. Investigative

ophthalmology & visual science *63*, 18.

Maezawa, S., Sakashita, A., Yukawa, M., Chen, X., Takahashi, K., Alavattam, K.G., Nakata, I.,

Weirauch, M.T., Barski, A., and Namekawa, S.H. (2020). Super-enhancer switching drives a

burst in gene expression at the mitosis-to-meiosis transition. Nat Struct Mol Biol *27*, 978-988.

Manterola, M., Brown, T.M., Oh, M.Y., Garyn, C., Gonzalez, B.J., and Wolgemuth, D.J. (2018).

BRDT is an essential epigenetic regulator for proper chromatin organization, silencing of sex

chromosomes and crossover formation in male meiosis. PLoS genetics *14*, e1007209.

Marcet-Ortega, M., Pacheco, S., Martinez-Marchal, A., Castillo, H., Flores, E., Jasin, M.,

Keeney, S., and Roig, I. (2017). p53 and TAp63 participate in the recombination-dependent

pachytene arrest in mouse spermatocytes. PLoS genetics *13*, e1006845.

Mytlis, A., Kumar, V., Qiu, T., Deis, R., Hart, N., Levy, K., Masek, M., Shawahny, A., Ahmad, A.,

Eitan, H.*, et al.* (2022). Control of meiotic chromosomal bouquet and germ cell morphogenesis

by the zygotene cilium. Science, eabh3104.

Ozata, D.M., Yu, T., Mou, H., Gainetdinov, I., Colpan, C., Cecchini, K., Kaymaz, Y., Wu, P.H.,

Fan, K., Kucukural, A.*, et al.* (2020). Evolutionarily conserved pachytene piRNA loci are highly

divergent among modern humans. Nat Ecol Evol *4*, 156-168.

Papanikos, F., Clement, J.A.J., Testa, E., Ravindranathan, R., Grey, C., Dereli, I., Bondarieva,

A., Valerio-Cabrera, S., Stanzione, M., Schleiffer, A.*, et al.* (2019). Mouse ANKRD31

Regulates Spatiotemporal Patterning of Meiotic Recombination Initiation and Ensures

Recombination between X and Y Sex Chromosomes. Molecular cell *74*, 1069-1085 e1011.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level

expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nature

protocols *11*, 1650-1667.

Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T., and Salzberg, S.L.

(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.

Nature biotechnology *33*, 290-295.

Qiao, H., Prasada Rao, H.B., Yang, Y., Fong, J.H., Cloutier, J.M., Deacon, D.C., Nagel, K.E.,

Swartz, R.K., Strong, E., Holloway, J.K.*, et al.* (2014). Antagonistic roles of ubiquitin ligase

HEI10 and SUMO ligase RNF212 regulate meiotic recombination. Nature genetics *46*,

194-199.

Rabbani, M., Zheng, X., Manske, G.L., Vargo, A., Shami, A.N., Li, J.Z., and Hammoud, S.S.

(2022). Decoding the Spermatogenesis Program: New Insights from Transcriptomic Analyses.

Annu Rev Genet *56*, 339-368.

Ramirez, F., Dundar, F., Diehl, S., Gruning, B.A., and Manke, T. (2014). deepTools: a flexible

platform for exploring deep-sequencing data. Nucleic acids research *42*, W187-191.

Rao, H.B., Qiao, H., Bhatt, S.K., Bailey, L.R., Tran, H.D., Bourne, S.L., Qiu, W., Deshpande, A.,

Sharma, A.N., Beebout, C.J.*, et al.* (2017). A SUMO-ubiquitin relay recruits proteasomes to

chromosome axes to regulate meiotic recombination. Science *355*, 403-407.

Reynolds, A., Qiao, H.Y., Yang, Y., Chen, J.K., Jackson, N., Biswas, K., Holloway, J.K.,

Baudat, F., de Massy, B., Wang, J.*, et al.* (2013). RNF212 is a dosage-sensitive regulator of

crossing-over during mammalian meiosis. Nature genetics *45*, 269-278.

Roeder, G.S., and Bailis, J.M. (2000). The pachytene checkpoint. Trends in genetics : TIG *16*,

395-403.

Royo, H., Prosser, H., Ruzankina, Y., Mahadevaiah, S.K., Cloutier, J.M., Baumann, M.,

Fukuda, T., Hoog, C., Toth, A., de Rooij, D.G.*, et al.* (2013). ATR acts stage specifically to

regulate multiple aspects of mammalian meiotic silencing. Genes & development *27*,

1484-1494.

San Filippo, J., Sung, P., and Klein, H. (2008). Mechanism of eukaryotic homologous

recombination. Annual review of biochemistry *77*, 229-257.

Shibuya, H., Ishiguro, K., and Watanabe, Y. (2014). The TRF1-binding protein TERB1

promotes chromosome movement and telomere rigidity in meiosis. Nature cell biology *16*,

145-156.

Spruce, C., Dlamini, S., Ananda, G., Bronkema, N., Tian, H., Paigen, K., Carter, G.W., and

Baker, C.L. (2020). HELLS and PRDM9 form a pioneer complex to open chromatin at meiotic

recombination hot spots. Genes & development *34*, 398-412.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,

Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S.*, et al.* (2005). Gene set enrichment

analysis: a knowledge-based approach for interpreting genome-wide expression profiles.

Proceedings of the National Academy of Sciences of the United States of America *102*,

15545-15550.

Subramanian, V.V., and Hochwagen, A. (2014). The meiotic checkpoint network: step-by-step

through meiotic prophase. Cold Spring Harbor perspectives in biology *6*, a016675.

Sun, S., Jiang, N., Jiang, Y., He, Q., He, H., Wang, X., Yang, L., Li, R., Liu, F., Lin, X.*, et al.*

(2020). Chromatin remodeler Znhit1 preserves hematopoietic stem cell quiescence by

determining the accessibility of distal enhancers. Leukemia *34*, 3348-3358.

Sun, S., Jiang, Y., Zhang, Q., Pan, H., Li, X., Yang, L., Huang, M., Wei, W., Wang, X., Qiu, M.*,*

et al. (2022). Znhit1 controls meiotic initiation in male germ cells by coordinating with Stra8 to

activate meiotic gene expression. Developmental cell *57*, 901-913 e904.

Symington, L.S. (2014). End resection at double-strand breaks: mechanism and regulation.

Cold Spring Harbor perspectives in biology *6*.

Turner, J.M. (2015). Meiotic Silencing in Mammals. Annu Rev Genet *49*, 395-412.

Turner, J.M., Mahadevaiah, S.K., Fernandez-Capetillo, O., Nussenzweig, A., Xu, X., Deng,

C.X., and Burgoyne, P.S. (2005). Silencing of unsynapsed meiotic chromosomes in the mouse.

Nature genetics *37*, 41-47.

Wang, L., Xu, Z., Khawar, M.B., Liu, C., and Li, W. (2017). The histone codes for meiosis.

Reproduction *154*, R65-R79.

Wang, Y., Chen, Y., Chen, J., Wang, L., Nie, L., Long, J., Chang, H., Wu, J., Huang, C., and

Lei, M. (2019). The meiotic TERB1-TERB2-MAJIN complex tethers telomeres to the nuclear

envelope. Nature communications *10*, 564.

Wang, Y., and Copenhaver, G.P. (2018). Meiotic Recombination: Mixing It Up in Plants. Annu

Rev Plant Biol *69*, 577-609.

Wei, W., Tang, X., Jiang, N., Ni, C., He, H., Sun, S., Yu, M., Yu, C., Qiu, M., Yan, D.*, et al.*

(2022). Chromatin remodeler Znhit1 controls bone morphogenetic protein signaling in

embryonic lung tissue branching. The Journal of biological chemistry *298*, 102490.

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.*, et al.*

(2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. Innovation

(Camb) *2*, 100141.

Wu, T., Lyu, R., You, Q., and He, C. (2020). Kethoxal-assisted single-stranded DNA

sequencing captures global transcription dynamics and enhancer activity in situ. Nature

methods *17*, 515-523.

Xu, M., Yao, J., Shi, Y., Yi, H., Zhao, W., Lin, X., and Yang, Z. (2021). The SRCAP chromatin

remodeling complex promotes oxidative metabolism during prenatal heart development.

Development *148*.

Xu, Y., Ayrapetov, M.K., Xu, C., Gursoy-Yuzugullu, O., Hu, Y., and Price, B.D. (2012). Histone

H2A.Z controls a critical chromatin remodeling step required for DNA double-strand break

repair. Molecular cell *48*, 723-733.

Yamada, S., Kugou, K., Ding, D.Q., Fujita, Y., Hiraoka, Y., Murakami, H., Ohta, K., and

Yamada, T. (2018). The histone variant H2A.Z promotes initiation of meiotic recombination in

fission yeast. Nucleic acids research *46*, 609-620.

Yoshida, S., Sukeno, M., Nakagawa, T., Ohbo, K., Nagamatsu, G., Suda, T., and Nabeshima,

Y. (2006). The first round of mouse spermatogenesis is a distinctive program that lacks the

self-renewing spermatogonia stage. Development *133*, 1495-1505.

Yoshida, S., Takakura, A., Ohbo, K., Abe, K., Wakabayashi, J., Yamamoto, M., Suda, T., and

Nabeshima, Y. (2004). Neurogenin3 delineates the earliest stages of spermatogenesis in the

mouse testis. Developmental biology *269*, 447-458.

Yu, T., Fan, K., Ozata, D.M., Zhang, G., Fu, Y., Theurkauf, W.E., Zamore, P.D., and Weng, Z.

(2021). Long first exons and epigenetic marks distinguish conserved pachytene piRNA

clusters from other mammalian genes. Nature communications *12*, 73.

Zhang, Q., Ji, S.Y., Busayavalasa, K., and Yu, C. (2019). SPO16 binds SHOC1 to promote

homologous recombination and crossing-over in meiotic prophase I. Sci Adv *5*, eaau9780.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,

Myers, R.M., Brown, M., Li, W*., et al.* (2008). Model-based analysis of ChIP-Seq (MACS).

Genome biology *9*, R137.

Zhao, B., Chen, Y., Jiang, N., Yang, L., Sun, S., Zhang, Y., Wen, Z., Ray, L., Liu, H., Hou, G*.,*

et al. (2019). Znhit1 controls intestinal stem cell maintenance by regulating H2A.Z

incorporation. Nature communications *10*, 1071.

Zheng, H., and Xie, W. (2019). The role of 3D genome organization in development and cell

differentiation. Nature reviews Molecular cell biology.

Zickler, D., and Kleckner, N. (2023). Meiosis: Dances Between Homologs. Annu Rev Genet.

Acknowledgments: We thank Chuan He for N3-kethoxal and Christopher L. Baker for

meiotic chromatin state data. **Funding:** This work was supported by grants from the

National Key Research and Development Program of China (2022YFA0806200,

2018YFC1003500, 2018YFA0800100, 2021YFC2501800), the National Natural Science

Foundation of China (32192403, 81971443, 32300702), the Science and Technology

Major Project of Inner Mongolia Autonomous Region of China to the State Key

Laboratory of Reproductive Regulation and Breeding of Grassland Livestock

(2020ZD0008), and the China Postdoctoral Science Foundation (2022M720797).

Author Contributions: S.S. and X.L. conceived and designed the study; S.S. performed

most of the experiments with the help of Y.J., Q.Z., H.P., F.H., X.Z., Y.G., X.Y., K.G.,

W.W., H.L., Z.S., Y.S., X.T., M.Y., and R.L.; Y.J. and N.J. performed the bioinformatics

analysis; X.L. supervised the work and acquired the funding support; and S.S. and X.L.

wrote the manuscript, with contributions from all authors. **Competing interests:** The

authors declare that they have no competing interests. **Data and materials availability:**

The NGS data generated in this study were deposited to the NCBI SRA database under

accession numbers SRP467214 (RNA-seq and ChIP-seq data) and SRP467448

(ATAC-seq and KAS-seq data).

Figure 1 | Znhit1 deletion in spermatocytes disrupts spermatogenesis. (**A**) Schematic

representation of the male germline development in mice. (**B**) Heatmap showing the

clustering of 47 epigenetic factors based on the transcription levels in leptotene and

zygotene spermatocytes. (**C**) *Znhit1* in situ hybridization in postnatal day 60 (P60) testis

sections. Scale bar, 20 μm. Sg, spermatogonia; pL, preleptotene; L, leptotene; Z, zygotene;

P, pachytene; D, diplotene, MI, metaphase; R, round spermatids; E, elongating spermatids.

(**D**) Schematic diagram showing the onset of Stra8-cre expression in the male germline

development. (**E**) *Znhit1* in situ hybridization in P42 testis sections. Scale bar, 50 μm. (**F**)

Histological testicular sections in control or *Znhit1*-sKO testis sections at the indicated

times. Scale bar, 50 μm. (**G**) Immunostaining of PNA in testis sections from control or

Znhit1-sKO mice at the indicated times. rST, round spermatids; eST, elongating

spermatids. Scale bar, 20 μm. All images are representative of n = 3 mice per genotype.

Figure 2 | Znhit1 deletion leads to meiotic pachytene arrest. (**A**) Immunostaining of

SYCP3 and HSPA2 in testis sections of P35 control or *Znhit1*-sKO mice. Scale bar, 20

μm. (**B**) Immunostaining of SYCP3 and pH3 in testis sections of P35 control or

Znhit1-sKO mice. Scale bar, 20 μm. (**C**-**E**) Immunostaining of SYCP3 in spermatocyte

chromosome spreads of control (**C**) or *Znhit1*-sKO mice (**D**). Quantitative data are shown

in (**E**). Scale bar, 10 μm. (**F** and **G**) TUNEL staining in testis sections from P14 control or

Znhit1-CKO mice. Quantitative data are shown in (**G**). Scale bar, 40 μm. All images are

representative of n = 3 mice per genotype. Data are presented as the mean ± s.d. ******* *p* <

0.001.

Figure 3 | Znhit1 controls meiotic recombination. (**A** and **B**) Immunostaining of SYCP3 and SYCP1 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. X, X chromosome; Y, Y chromosome; PAR, pseudoautosomal region. Quantitative data are shown in (**B**). Scale bar, 10 μm. (**C**) Immunostaining of SYCP3 and γH2AX in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Scale bar, 10 μm. (**D**) Immunostaining of SYCP3 and pATM in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Scale bar, 10 μm. (**E** and **F**) Immunostaining of SYCP3 and RAD51 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Quantitative data are shown in (**F**). Scale bar, 10 μm. (**G** and **H**) Immunostaining of SYCP3 and MLH1 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Quantitative data are shown in (**H**). Scale bar, 10 μm. All images are representative of n = 3 mice per genotype. Data are presented as the mean ± s.d. ******* *p* < 0.001.

Figure 4 | Spermatocyte transcriptome analysis using scRNA-seq. (**A**) UMAP plot

showing the annotated germ cells captured from both control and *Znhit1*-sKO testicular

cells. (**B**) Expression of selected markers identifying major somatic cell types and germ

cell types on the UMAP plot. UMAP plot showing the expression patterns of *Znhit1* in

control or *Znhit1*-sKO testicular cells. (**C**) Heatmap showing the top marker genes of each

cell cluster from scRNA-seq data. U-Sg, undifferentiated spermatogonia; D-Sg,

differentiating spermatogonia; PreL, preleptotene; Lep, leptotene; Zyg, zygotene; Pac,

pachytene. (**D**) Top: UMAP plot showing the annotated germ cells captured from control

or *Znhit1*-sKO testicular cells. Down: UMAP plot showing the expression patterns of

Znhit1 in control or *Znhit1*-sKO testicular cells. (**E**) Bar plot showing the percentage of

major spermatocyte types for control or *Znhit1*-sKO testicular cells. (**F**) UMAP plot

showing the expression patterns of *Pou5f2* in control or *Znhit1*-sKO germ cells.

Figure 5 | Znhit1 deletion impairs meiotic transcriptional activation. (**A**) Line graph

showing DEGs (upregulation or downregulation) in two consecutive spermatocyte stages.

PreL, preleptotene; Lep, leptotene; Zyg, zygotene; Pac, pachytene. (**B**) Volcano plot

showing the distribution of upregulated and downregulated DEGs for each cell type in

spermatocytes between control or *Znhit1*-sKO testicular cells. (**C**) Cluster annotation on

the basis of the expression of pachytene-activated genes in control or *Znhit1*-sKO

testicular cells. (**D**) Venn plots showing the number of shared genes between

pachytene-activated genes and DEGs in *Znhit1*-sKO pachytene spermatocytes. (**E**)

Representative shared Gene Ontology (GO) terms of upregulated (upper) and

downregulated (lower) DEGs at consecutive spermatocyte stages. (**F**) Volcano plot

showing differentially expressed genes (DEGs) in bulk RNA-seq following Znhit1

deletion (n = 2). (**G**) Heatmap depicting the expression of the indicated genes at the

zygotene and pachytene stages in the scRNA-seq data. (**H**) Cluster annotation on the basis

of the expression of downregulated genes in bulk RNA-seq between control and

Znhit1-sKO testicular cells. (**I**) GSEA of RNA-seq data for the control and *Znhit1*-sKO

testicular cells. Selected gene sets encoded products related to homologous recombination.

(**J**) Violin plots showing downregulated DEGs of Znhit1-deficient zygotene cells and

pachytene cells related to homologous recombination. (**K**) Immunostaining of SYCP3 and

SUMO2/3 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Scale bar,

10 μm. cluster genes. ****** $p < 0.01$, ******* $p < 0.001$, ******** $p < 0.0001$.

Figure 6 | Znhit1 regulates H2A.Z binding on promoter and enhancer regions. (**A**) Immunostaining of SYCP3 and H2A.Z in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. The dashed circle indicates X-Y chromosomes. Scale bar, 10 μm. (**B**) Representative tracks of END-seq, DMC1-SSDS, and H2A.Z ChIP-seq in wild-type, control, or *Znhit1*-sKO testicular cells. (**C**) Heatmaps of chromatin states produced by ChromHMM and showing different enrichment for H2A.Z in control or *Znhit1*-sKO testicular cells. (**D**) Heatmaps and mean plots of H2A.Z ChIP-seq signal in control or *Znhit1*-sKO testicular cells (n = 2). (**E**) Venn diagram showing the overlap between decreased H2A.Z-bound genes and downregulated genes in *Znhit1*-sKO testicular cells.

886 (**F**) Heatmaps and mean plots of downregulated H2A.Z ChIP-seq signals surrounding

887 different types of promoters and enhancers.

Figure 7 | Znhit1/H2A.Z/A-MYB axis regulates enhancer activation and gene

expression. (**A**) Motif analysis of active promoters or active enhancers with decreased

H2A.Z binding for putative transcription factor (TF)-binding sites using the HOMER

database. (**B**) Ridge map showing the A-MYB activity (AUC) at each cell type. (**C**)

Scatter plot with marginal histograms comparing the fold change of DEGs from

A-MYB-related RNA-seq data with Znhit1-related RNA-seq data in P14 testicular cells.

(**D**) GSEA of RNA-seq data for the control and *Znhit1*-sKO testicular cells. Selected gene

sets encoded products related to A-MYB target genes. (**E**) Violin plots showing the

expression of A-MYB target genes at consecutive spermatocyte stages. (**F**) Histogram

showing H2A.Z and A-MYB binding sites. (**G**) Pie chart showing the distribution of

H2A.Z and A-MYB co-binding sites. (**H**) Representative H2A.Z ChIP-seq tracks,

KAS-seq signals, and RNA-seq signals of the *Ccnb1ip1* locus in the control and

Znhit1-sKO testicular cells, compared with A-MYB ChIP-seq peaks. (**I** and **J**) Heatmaps

and mean plots of H2A.Z ChIP-seq, KAS-seq, and ATAC-seq showing signal changes

within A-MYB and H2A.Z co-binding active promoters or active enhancers. **** $p <$

0.0001.

Supplementary Materials for

H2A.Z deposition at meiotic prophase I underlies homologous

recombination and pachytene genome activation during male meiosis

Shenfei Sun *et al.*

916
917
918   *Corresponding author. Email: Xinhua Lin, xlin@fudan.edu.cn; Shenfei Sun
919   shenfei_sun@fudan.edu.cn

This PDF file includes:

925
926
927   Figs. S1 to S9

Other Supplementary Material for this manuscript includes the following:

930
931
932   Tables. S1 to S8

Figure S1. Znhit1 expression during meiotic progression and spermatocyte-specific
deletion of Znhit1 with Stra8-cre. (**A** and **B**) Venn diagram and heatmaps showing
chromatin regulators highly expressed during leptotene and zygotene spermatocytes. (**C**)
Violin plot showing Znhit1 transcription level during spermatogenesis. Gene expression
profiles derived from published data (Chen et al, 2018). (**D**) Znhit1 in situ hybridization
in P14 testis sections. (**E**) Testis size comparison between control and Znhit1-sKO mice
at P35. (**F**) Hematoxylin and eosin (H&E) - stained testis sections from control and
Znhit1-sKO mice at P60.

Figure S2. H1T⁺ pachytene spermatocytes are detected in *Znhit1*-sKO testes. (A)

Immunostaining of H1T in testis sections from control and *Znhit1*-sKO mice at P35.

Scale bar, 20 μm.

953 **Figure S3. Znhit1 controls meiotic recombination.** (**A**) Immunostaining of SYCP3 and

954 RPA2 in spermatocyte chromosome spreads of control or *Znhit1*-sKO mice. Scale bar, 10

955 μm.

956
957

A

B

959 **Figure S4. Single-cell RNA-seq of testicular cells.** (**A**) Dot plot showing the relative

960 expression and the percentage of cells expressing selected markers across scRNA-seq

961 clusters. (**B**) Violin plot showing the normalized expression of *Znhit1* in control and

962 *Znhit1*-sKO spermatocytes at consecutive stages.

963
964

Figure S5. Single-cell transcriptomic analysis of meiotic genome activation. (**A**) Violin plots showing the average expression level of upregulated or downregulated genes between two consecutive spermatocyte stages. (**B**) Violin plots showing the average expression level of upregulated or downregulated genes between P14 control and *Znhit*1-sKO testicular cells. PreL, preleptotene; Lep, leptotene; Zyg, zygotene; Pac, pachytene.

972
973

975 **Figure S6.** differentially expressed genes (DEGs) in P14 testicular cells between control

976 and Znhit1-sKO mice. (**B**) Gene ontology (GO) analysis of DEGs showing changed

977 biological processes. (**C** and **D**) Bar charts showing HR-related gene expression in P14

978 control and *Znhit*1-sKO testes. Data from bulk RNA-seq are shown in (**C**). Validated data

979 from RT-qPCR are shown in (**D**). Ctrl, control; sKO, *Znhit1*-sKO.

980
981

Figure S7. H2A.Z is enriched at meiotic gene promoters and enhancers. (**A**)
Schematic representation of the components of the SRCAP chromatin remodeling
complex. (**B**) Heatmaps showing H2A.Z ChIP-seq signals surrounding different types of
promoters. (**C**) Heatmaps showing H2A.Z ChIP-seq signals surrounding different types of
enhancers.

988
989

991 **Figure S8. Znhit1 deletion downregulates the expression of A-MYB target genes.** (A)

992 Violin plot showing the normalized expression of *Mybl1* in control and *Znhit1*-sKO

993 spermatocytes at consecutive stages. (B) Cluster annotation on the basis of the expression

994 of A-MYB target genes in control or *Znhit1*-sKO testicular cells.

995
996

Figure S9. Reproducibility of replicates in this study. (**A**) Scatter plots showing H2A.Z

enrichments in P14 control or *Znhit1*-sKO testes for two independent H2A.Z ChIP-seq

replicates. Pearson correlation coefficients are indicated. (**B**) Scatter plots showing

KAS-seq signals in P14 control or *Znhit1*-sKO pachytene cells for two independent

H2A.Z ChIP-seq replicates. Pearson correlation coefficients are indicated. (**C**) Scatter

plots showing ATAC-seq signals in P14 control or *Znhit1*-sKO pachytene cells for two

independent H2A.Z ChIP-seq replicates. Pearson correlation coefficients are indicated.

1005
1006

1007