MULTIPLE DISTINCT TIMESCALES OF RAPID SENSORY ADAPATION IN THE THALAMOCORTICAL CIRCUIT

Abbreviated title: The timescales of sensory adaptation

Yi Juin Liew[1, 2, *], Elaida D Dimwamwa[1, *], Nathaniel C Wright[1],
Yong Zhang[3, 4], Garrett B Stanley[1]

[1] Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA 30332, USA

[2] Joint PhD Program in Biomedical Engineering, Georgia Institute of Technology – Emory University – Peking University, Atlanta, GA 30332, USA

[3] Neuroscience Research Institute and Department of Neurobiology, School of Basic Medical Sciences, Peking University, Key Laboratory for Neuroscience, Ministry of Education of China and National Health Commission of the People's Republic of China, Beijing 100083, China

[4] PKU-IDG/McGovern Institute for Brain Research, Beijing, 100871, China.

[*] Indicates equal contribution

Correspondence: Professor Garrett B Stanley
Coulter Department of Biomedical Engineering
Georgia Institute of Technology & Emory University
313 Ferst Drive, Atlanta GA 30332-0535 USA
Phone: 404-385-5037, Fax: 404-385-5044
Email: garrett.stanley@bme.gatech.edu

Number of pages: 32
Number of figures: 8
Number of words in Abstract: 225
Numbef of words in Introduction: 700
Number of words in Discussion: 1648

Keywords: thalamocortical; rapid adaptation; thalamic synchrony; vibrissa

Disclosures: The authors declare no competing financial interests.

Acknowledgements: This work was supported by NIH National Institute of Neurological Disorders and Stroke (NINDS) BRAIN Grant R01NS104928 and NINDS BRAIN Grant RF1NS128896. YJL was supported by a Georgia Tech-Emory-PKU Global Biomedical Engineering Fellowship. EDD was supported by a National Science Foundation Graduate Research Fellowship and the Howard Hughes Medical Institute through the James H. Gilliam Fellowships for Advanced Study program. YZ was supported by STI2030-Major Projects 2022ZD0204900. The authors thank Eunji Cheong and Cheong lab members at Yonsei University for helpful conversations at various stages of this work.

ABSTRACT

Numerous studies have shown that neuronal representations in sensory pathways are far from static but are instead strongly shaped by the complex properties of the sensory inputs they receive. Adaptation dynamically shapes the neural signaling that underlies our perception of the world, yet remains poorly understood. We investigated rapid adaptation across timescales from hundreds of milliseconds to seconds through simultaneous multi-electrode recordings from the ventro-posteromedial nucleus of the thalamus (VPm) and layer 4 of the primary somatosensory cortex (S1) in anesthetized mice in response to controlled, persistent whisker stimulation. Observations in VPm and S1 reveal a degree of adaptation that progresses through the pathway. Signatures of two distinct timescales of rapid adaptation in the firing rates of both thalamic and cortical neuronal populations were revealed, also reflected in the synchrony of the thalamic population and in the thalamocortical synaptic efficacy that was measured in putatively monosynaptically connected thalamocortical pairs. Controlled optogenetic activation of VPm further demonstrated that the longer timescale adaptation observed in S1 is likely inherited from slow decreases in thalamic firing rate and synchrony. Despite the degraded sensory responses, adaptation resulted in a shift in coding strategy that favors theoretical discrimination over detection across the observed timescales of adaptation. Overall, although multiple mechanisms contribute to rapid adaptation at distinct timescales, they support a unifying framework on the role of adaptation in sensory processing.

Significance Statement: Although the perceptual effects of persistent sensory stimulation have been known for centuries, the rapid sensory adaptation of the underlying neural signaling to these persistent inputs are not well understood. Here, we present evidence for two distinct timescales of adaptation over several seconds across the thalamocortical circuit in mice. We identify both the overall level of neural activity and the corresponding population synchrony of the thalamic inputs to primary somatosensory cortex as key role players shaping the cortical adaptation.

INTRODUCTION

The natural sensory environment that we experience consists of inputs that span a wide range of temporal and spatial scales. As sensory signals move from the periphery to the sensory cortex, sensory representations are progressively transformed via the specific characteristics of the circuitry at each stage and the nature of the anatomical projections from region to region (Maravall et al., 2007, 2013; Adibi and Lampl, 2021). Adaptation has been well-documented across all sensory modalities in a wide variety of organisms (Whitmire and Stanley, 2016), thought to overcome changes in stimulus statistics via multiple mechanisms that modulate neuronal response properties (Adibi et al., 2013; Latimer et al., 2019). It has long been asserted that the brain's ability to adapt to dynamic inputs at each stage in the processing hierarchy is a fundamental function that is critical for survival, providing a functional advantage by shifting the coding strategy in an ethological advantageous manner (Barlow, 1961). While adaptation has been proposed as a unifying theoretical framework that may integrate numerous disparate experimental observations (Drew and Abbott, 2006), the large majority of experimental studies of neuronal activity in the face of sensory adaptation focus on a specific single timescale of adaptation (Chung et al., 2002; Khatri et al., 2004; Kheradpezhouh et al., 2017). What is not well understood is the evolution of adaptation along different timescales across brain regions in the processing hierarchy, and the subsequent functional implications.

In the rodent whisker-mediated somatosensory pathway, previous studies have shown that the degree of sensory adaptation increases as neural signals traverse from periphery to cortex (Ahissar et al., 2000; Khatri et al., 2004; Ganmor et al., 2010; Mohar et al., 2013, 2015). A particularly important stage of the sensory signaling pathway is the thalamocortical circuit, where highly divergent thalamic afferents synapse on to both excitatory and inhibitory neurons in the cortex, initiating thalamic-driven feedforward inhibition that has profound effects on sensory representations (Simons and Carvell, 1989; Gabernet et al., 2005; Cruikshank et al., 2007; Wang et al., 2010a). Thalamocortical synapses have been shown to exhibit short-term depression, modulating the relative role of the projecting thalamic input (Chung et al., 2002; Castro-Alamancos, 2004a; Katz et al., 2006). However, thalamic neurons themselves also exhibit a moderate degree of adaptation, resulting in a withdrawal of the primary drive to cortex. Recent work at the population level also suggests that adaptive changes in thalamic population spiking synchrony play an important role in mediating cortical adaptation through the highly timing-sensitive integration by thalamo-recipient cortical neurons (Wang et al., 2010b; Wright et al., 2021). Given the complexity of the thalamocortical circuit, it is likely that multiple mechanisms are engaged in the transformation of sensory representations throughout the pathway. How this myriad of elements collectively evolves across timescales remains unknown.

Here, we investigated in detail the effects of persistent tactile stimuli on neuronal population activity across the thalamocortical circuit, and specifically the relative roles of aspects of the thalamic input in shaping the cortical response. We conducted multi-site, multi-electrode extracellular recordings of spiking activity from the ventro-posteromedial nucleus of the thalamus (VPm) and layer 4 of the whisker response region of primary somatosensory cortex (S1) in

anesthetized mice. We presented controlled, repetitive, punctate whisker deflection trains to
individual whiskers and quantified the adaptive effects on single-unit spiking activity in VPm and
putative excitatory and inhibitory S1 neurons, as well as in a subset of putative synaptically-
connected thalamocortical pairs. In both the cortex and the thalamus, we uncovered evidence of
two distinct timescales in the rapid adaptation dynamics over a timecourse of seconds that
exhibited differential functional properties across the thalamocortical circuit. We observed an
evolution in the relative roles of individual mechanisms across these timescales, with cortical
adaptation at the longer timescale appearing to primarily reflect the adaptation of thalamic drive
and synchrony. As a result, from the perspective of an ideal observer of spiking activity, adaptation
in both of the distinct timescales also continually shifts towards better discrimination of whisker
deflection velocity at the expense of detection. These results collectively suggest a complex
interplay of adaptive elements at the population level that evolves on distinct timescales, and
position the thalamic population activity as a key role player in shaping the cortical representation.

METHODS

Animals
Experiments were conducted using 10 adult C57BL/6J and 14 adult NR133 x Ai32 (NR133 Cre-recombinase driver line crossed with Ai32 mice for expression of channelrhodopsin in the VPm; Bolus et al., 2020; Gerfen et al., 2013) mice of both sexes (14 male, 10 female) aged between 11 and 44 weeks (mean: 20 weeks). All procedures were approved by the Georgia Institute of Technology Institutional Animal Care and Use Committee and followed guidelines established by the National Institutes of Health.

Head-plate implantation
Head-plate implantation was performed three to seven days before electrophysiological recordings were conducted. A lightweight custom metal (titanium or stainless steel) head-plate was implanted on the skull of the mouse for head fixation and improved stability during recording, in accordance with a previously described protocol (Borden et al., 2017; Liew et al., 2021; Wright et al., 2021; Dimwamwa et al., 2023). During this survival surgical preparation, the animal was sedated with 5% vaporized isoflurane and anesthesia was maintained with 2-3% isoflurane for the remainder of the procedure. An opioid and non-steroidal anti-inflammatory analgesic were administered post-operatively (SR-Buprenorphine 0.8 - 1 mg/kg, SC and Ketoprofen 5-10 mg/kg, IP). The animal's eyes were covered with ophthalmic eye ointment (Puralube Vet Ointment) to prevent dehydration. Body temperature was monitored and maintained at 37 °C. After sterilizing the skin above the skull and applying topical anesthetic (lidocaine 2%, 0.05 ml, max 7mg/kg, SC), we removed the skin and scraped off periosteum and any conjunctive tissue from the skull. We gently separated tissue and muscles close to the lateral edge of the skull using a scalpel blade (Glass Van & Technocut, size no. 15), leaving sufficient room for head-plate attachment, away from the targeted recording areas. Especially critical for targeting of the ventro-posteromedial nucleus of the thalamus (VPm), the skull surface was leveled during head fixation by adjusting the animal's head position to minimize the relative height difference (< 100 um) between bregma and lambda (Franklin and Paxinos, 2008). The metal headplate was secured over the intact skull with C&B-Metabond (Parkell Prod Inc) and the rest of the attachment site and edges of the skin incision were secured with skin adhesive (Vetbond, 3M). The final head-plate and dental acrylic structure created a well for the containment of Ringer's solution (in mM: 135 NaCl, 5 KCl, 5 HEPES, 1 MgCl2-6H2O, 1.8 CaCl2-2H2Ol pH 7.3) for future imaging and electrophysiological recording sessions. We applied a thin layer of transparent Metabond dental acrylic over both hemispheres to prevent inflammation and covered the exposed skull with a silicone elastomer (Kwik-Cast, World Precision Instruments).

Intrinsic signal optical imaging of the mouse S1
Intrinsic signal optical imaging of whisker-evoked responses in mouse primary somatosensory cortex was performed through the skull under 1-1.2% isoflurane anesthesia to functionally generate a map of the precise location of S1 barrel columns (Lefort et al., 2009; Pala and Petersen, 2015). The imaging was conducted in a separate session to the anesthetized electrophysiology, either within the same session as head-plate implantation or in a separate session. The skull was

thinned using a dental drill (0.1 mm diameter bit, Komet, USA) until blood vessels became visible under warm Ringer's solution. The Ringer's was then covered with a glass coverslip (thickness 0.13-0.17mm, Fisherbrand). A reference image of the blood vasculature was captured under green illumination (530nm; M530F1 LED, Thorlabs). Repetitive whisker stimuli (4 seconds of stimuli at 10 Hz; sawtooth waveform; 1000 deg/s) were delivered to individual whiskers (see "Sensory stimulus section" below) while imaging S1 under red illumination (625 nm; M625F1 LED, Thorlabs). Images were acquired using a CCD Camera (MiCam02HR, SciMedia) at 100 Hz. Using previously published methods, we recorded intrinsic signals from at least three columns and estimated the location of the unmapped columns by overlaying a template S1 barrel map reconstructed from histology (Liew et al., 2021). At the end of the imaging session, we applied transparent Metabond dental acrylic directly on the S1 site to protect the thinned skull from contamination and infection and sealed the exposed skull with a silicone elastomer (Kwik-Cast, World Precision Instruments).

Anesthetized electrophysiology
Animals were sedated with 5% vaporized isoflurane and maintained with 2-3% isoflurane. Their eyes were covered ophthalmic eye ointment and the silicone elastomer removed from the skull surface. The skull above the VPm and S1 was carefully thinned with a dental drill (0.1 mm diameter bit, Komet, USA), while frequently irrigating with Ringer's solution frequently to remove debris and prevent overheating. When the skull was thin enough to easily puncture, we made small craniotomies above VPm and S1. The craniotomies were constantly irrigated with Ringer's solution to maintain the integrity of the brain tissue.

The animals were then maintained with 1-2% isoflurane for the electrophysiological recordings. Either in separate sessions or in the same session for paired recordings, 32-channel silicon probes with the 'Poly-3' configuration (A1x32-Poly3-5mm-25s-177, A1x32-Poly3-6mm-50s-177, or A1x32-Poly3-10mm-50s-177, Neuronexus, Ann Arbor, MI) were independently lowered into the VPm and S1. All signals were acquired using the Tucker Davis Technologies RZ2 Bioprocessor (Alachua, FL). Neuronal signals were amplified, bandpass filtered (500 Hz – 5 kHz), and digitized at 24,414.0625 Hz per channel. Stimulus waveforms and other continuous data were digitized at 10 kHz per channel.

For VPm recordings, the electrode was typically positioned 1.8 mm caudal and 1.8 mm lateral to bregma and lowered to a depth of 3 – 3.5 mm. If VPm could not be functionally located using this method, we also targeted VPm using coordinates 1.0 - 1.2 mm from the B1 or Beta columns of S1 as identified via the intrinsic signal optical imaging map.

For S1 recordings, a small craniotomy was created at the desired location based on the intrinsic signal optical imaging map, being careful to remain distant from blood vessels. The electrode was positioned at a 35 ° angle from the vertical axis (parallel to S1) and cortical neurons were recorded at a stereotaxic depth of 300-600 um from the cortical surface, corresponding to layer 4.

208 After each recording session, the probes were removed, exposed tissue was covered with
209 agarose, and the well was sealed with Kwik-Cast. Subsequent recording sessions were done
210 every other day. One recording session was conducted per day and two to three recording
211 sessions were conducted for each mouse using either the original craniotomy or additional
212 craniotomies targeting a different cortical column and barreloid.
213
Sensory stimulus
Single whiskers of interest were identified by manually deflecting individual whiskers while
monitoring the extracellular signal to determine the whisker that evoked the largest response in
each brain region being recorded. Controlled, single whisker stimulation was delivered to the
identified whiskers of interest in the rostral-caudal plane using a computer-controlled galvo-motor
(galvanometer optical scanner model 6210H, Cambridge Technology). Whiskers were trimmed
~12 mm from the face. The whisker was fed into the insertion hole at the end of an extension tube
(inner diameter: ~200-300 μm um, length: 15mm) that was connected to the rotor of the
galvanometer stimulator. The stimulus probe was positioned 10 mm from the Vibrissal pad. The
range of motion of the galvanometer was ± 20°, with a bandwidth of 200 Hz. The galvanometer
system was controlled using a custom developed hardware/software system (MATLAB Real-Time
Simulink System, MathWorks or the Real- time experiment Interface application (http://rtxi.org/),
with a 1 kHz sampling rate. The inter-trial interval (time between end of one trial and beginning of
the next) for all stimuli was at least 15 seconds, to minimize any residual effects from the previous
trial.
229
230 *Adapting stimulus.* To induce rapid sensory adaptation, a train of punctate whisker deflections
231 was presented at 10 Hz for 10 seconds in the rostral-caudal plane using the galvanometer as
232 described above. Each punctate deflection consisted of a 300 deg/s sawtooth waveform with an
233 exponential rise and decay lasting 17 ms. The reported velocity is defined by the average velocity
234 over the 8.5 ms rising phase (Wang et al., 2010).
235
For the assessment of velocity sensitivity and the detectability and discriminability from the
perspective of an ideal observer, we presented sawtooth deflections delivered at one of five
velocities: 50, 300, 450, 600, 1200 deg/s. The deflections were presented under three different
conditions: "Control" condition with no preceding adapting stimulus or "Adapted" conditions with
either a preceding 1.5 second adapting stimulus or 10 second adapting stimulus. Isolated
deflections were delivered 100 msec following the end of the adapting train. All stimulus conditions
were randomized across trials, and at least 20 trials per stimulus condition were typically obtained
for different velocities.
244
Weak sinusoidal stimulus. To assess monosynaptic connectivity between VPm and S1 neurons,
the thalamocortical circuit was probed using a weak, desynchronizing sinusoidal whisker stimulus
to elevate baseline firing rates in recorded neurons (Liew et al., 2021). Specifically, 2 - 4 Hz weak
sinusoidal deflections of 4 deg amplitude and 4 sec duration (Bruno and Simons, 2002; Bruno
and Sakmann, 2006; Wang et al., 2010b) were delivered for approximately 200-500 trials to obtain

at least 2000 spikes for the cross-correlation analysis. Occasionally, the first 0.5 - 1 seconds of
the trials were eliminated due to high firing rates at the onset of stimulus presentation.
252
Optogenetic stimulation of neural activity
A subset of recordings was performed on NR133 x Ai32 transgenic mice to directly activate VPm
neurons, bypassing sensory activation from the periphery. In these experiments, a 32-channel
silicon opto-electrode with an attached 105 um fiber (A1x32-Poly3-5 mm-25s-177-OA32LP
Neuronexus, Ann Arbor, MI) coupled to a 200 μm optic fiber (M95L01, Thorlabs) and a 470 nm
LED (M470F3, Thorlabs) was lowered into VPm. The optogenetic stimulation was designed to
emulate the adaptation induced by the sensory stimulation described above. The optogenetic
adapting stimulus consisted of 470 nm LED pulses that were 17 ms in duration and were delivered
to VPm cell bodies for 10 seconds at 10 Hz.
262
Post-mortem histology
To verify the recording sites in VPm and S1, a subset of animals was perfused after the paired
recording experiments. To label each probe recording track, the probe was slowly retracted along
its axis of entry at the conclusion of recording and a few drops of DiI (2.5 mg/mL, in ethanol) were
applied on the probe. The electrode was then inserted into the same penetration site along the
same axis and the probe was left in the brain for at least 15 minutes. The animal was then
euthanized with an overdose of sodium pentobarbital (euthasol, 0.1 mL at 390 mg/mL), a
transcardial perfusion was performed, the brain was extracted, and the brain was fixed in 4%
paraformaldehyde (PFA, Electron Microscopy Sciences) overnight. The brain was sliced in 100
um coronal sections. The slices were incubated with DAPI (2 mM in PBS, AppliChem) for 15 min
before being mounted on slides with a DABCO solution (1,4-Diazabicyclo[2.2.2]octane, Sigma).
The slices were imaged with a confocal microscope (Laser Scanning Microscope 900, Zeiss,
Germany) to verify the overlap between the DiI stained recording track and the CO-stain.
276
Analytical Methods
Spike Sorting. Spike sorting was performed offline using KiloSort2.5 (Pachitariu et al., 2024)
followed by manual curation of neuronal clusters in Phy2 (Rossant and Harris, 2013). Well-
isolated single units were classified using the following criteria: mean waveform signal-to-noise
ratio greater than 3 and less than 1% and 3% of all spikes violating the 2 ms refractory period for
cortical and VPm neurons, respectively (Liew et al., 2021; Dimwamwa et al., 2023).
283
Analysis of sensory response. Whisker evoked activity was quantified within the neural response
window for each neuron, defined as a 30 ms window following stimulus onset. Evoked response
amplitudes were characterized from the trial-to-trial spike count in the neural response window
following each deflection. When applicable, evoked responses were normalized to the maximum
response to the first three stimuli. The normalized evoked response amplitude was defined as the
spike count in the response window divided by the maximum response amplitude evoked by the
first three deflections in the adapting trains. The adaptation ratio was computed for each neuron
as the average spike count elicited by the last three deflections, normalized to the averaged
response to the first three deflections. First-spike latencies were defined as the average time of

8

the first spike within the response window across trials in which there was a spike within the 30
ms response widow (Storchi et al., 2012).

VPm neuron validation. It is important to take great care to distinguish different thalamic nuclei
from one another, particularly in the mouse where the regions are very small and close together.
Thus, all analyses only considered neurons that were responsive to the deflection of at least one
whisker. Whisker responsiveness was determined by performing the Wilcoxon signed rank test
on the post-stimulus spiking in the 30 ms neuronal response window compared to an equivalent
duration pre-stimulus window. Then, a combination of measures was used to classify whisker
responsive neurons as originating from the VPm. First, for isolated (non-adapting) stimuli, we
quantified the first-spike latency. Second, we quantified the shift in first-spike latency response to
adapting stimuli, defined as the time difference between the first-spike latency to the first and last
adapting stimuli (Masri et al., 2008). Third, we calculated the response reliability, defined as the
percentage of trials where a response was detected within 20 ms in response to each deflection
of the repetitive stimulation (8 – 10 Hz; Mainen and Sejnowski 1995). Only neurons with an
average first-spike latency of less than 12 ms, latency shift of less than 20 ms, and response
reliability of more than 20% were included (Liew et al., 2021; Wright et al., 2021; Borden et al.,
2022; Dimwamwa et al., 2023). Then, each neuron was assigned a primary whisker based on the
whisker that elicited the largest peak and smallest latency in the neural response window as
quantified from the LFP on the channel to which the neuron belonged. The LFP was obtained by
bandpass filtering the raw neuronal signal (0.5 – 200 Hz) on each channel. The sensory response
of each neuron was only analyzed based on trials in which its primary whisker was being deflected.

Cortical neuron classification. S1 layer 4 is the major thalamocortical recipient of VPm projections,
where a diversity of cell types receive the feedforward synaptic inputs (Bruno and Simons, 2002).
Here, cortical neurons were classified as regular-spiking (RS) or fast-spiking (FS) based on the
time interval between the trough and peak of the mean spike waveform, with a cutoff of 0.4 ms
separating the broad RS neurons from the narrow FS neurons (Barthó et al., 2004).

Determining topographic alignment. A VPm-S1 neuron pair was verified to be topographically
aligned when they shared the whisker that elicits the maximum response to isolated stimuli (both
spike evoked response and LFP amplitude) and the mean spike latency difference in the sensory
response between VPm and S1 neuron pair was between 1-5 ms.

Characterization of the adaptation dynamics with exponential models. The adaptive reduction in
response amplitudes was characterized by either single or bi-exponential models applied to the
normalized evoked response or synchrony over the 10 second adapting stimulus period.
Normalization was to the maximum response/synchrony for the first three stimuli.

For a single exponential fit, the following exponential function was used:

334
$$y(t) = a * e^{-\lambda * t}$$

336 where y(t) is the normalized evoked response at time t, λ is the time constant of adaptation, and
337 a is a scaling factor.
338
339 For the bi-exponential fit, the following function was used:
340
341 $$y(t) = a * e^{-\lambda_1 * t} + b * e^{-\lambda_2 * t}$$
342
where y(t) is the normalized evoked response at time t, λ_1 and λ_2 are the time constants of two
timescales of adaptation, and a and b are scaling factors (Patterson et al., 2013). All parameters
were optimized using the 'fmincon' function in MATLAB. The optimization aimed to minimize the
sum of squared differences between the observed data points and the model predictions. We
imposed a constraint requiring that the second time constant be at least $\frac{1}{\sqrt{10}}$ times larger than the
first time constant, empirically determined to separate timescales. The goodness of fit of each
model was quantified by assessing the correlation coefficient (R^2) between the experimental
measures and the fit.
351
352 To visualize the exponential decay of the adaptation response, we examined the data within a
353 semi-log plot to identify any exponential components present in the data. Each straight-line
354 segment on the plot corresponds to an exponential component. We then performed linear
355 regression on each segment to determine its slope, which represents the time constant of the
356 exponential component.
357
358 *Analysis of velocity sensitivity.* To measure the sensitivity of thalamocortical neurons to sensory
359 stimulus velocity, the whisker response was quantified as a function of increasing deflection
360 velocities: 50, 125, 300, 600, 900, 1200 deg/s. These stimuli were presented either in isolation
361 (Control) or immediately following adapting stimuli (1.5 sec or 10 sec), all randomly interleaved.
362
Spike cross-correlation analysis. Spike cross-correlation analysis was performed to infer
monosynaptic connectivity and compute synaptic efficacy for paired recordings in topographically
aligned thalamocortical regions, as well as to compute VPm population synchrony. For each spike
in a 'reference' neuron, a cross-correlogram (CCG) histogram of the relative times of the spikes
from a 'target' neuron was generated within a 25 msec window before and after each reference
spike. All CCGs were constructed using 0.5 ms bins.
369
Inferring monosynaptic connectivity. Using a previously developed method for monosynaptic
connectivity inference, in paired and topographically aligned recordings, the thalamocortical
circuit was externally driven with a weak whisker stimulus (see Methods: Weak sinusoidal
stimulus; Liew et al., 2021). For each topographically aligned VPm-S1 pair, a CCG of the S1
neuron's spiking relative to the VPm neurons spiking during the weak whisker stimulus was
constructed, referred to as the raw CCG. To correct for correlated, stimulus-locked spiking, a
shuffle-corrected (SC) spike cross-correlogram was generated. Specifically, a shuffled CCG was
generated by randomly shuffling the trials of the VPm and S1 neuron spiking activity relative to

each other. The SC CCG was then generated by subtracting the shuffled CCG from the raw CCG.
This effectively corrects for any elements of the CCG that are due to the stimulus. To conclude
that a neuronal pair was 'connected', two criteria were adopted that were expanded from previous
studies (Reid and Alonso, 1995; Swadlow and Gusev, 2001; Bruno and Sakmann, 2006) based
on both the raw and shuffle-corrected cross-correlogram: (Criterion 1) a notable sharp,
millisecond-fine peak observed within a narrow lag of 1 - 4 ms after a VPm spike that is the
maximum in the raw CCG; and (Criterion 2) this fast 'monosynaptic peak' is significant or still
present after accounting for (subtracting) stimulus-induced correlations in the SC CCG (Liew et
al., 2021).
387

388   *Measuring synaptic efficacy*. To quantify the effects of adaptation on synaptic transmission across
389   the thalamocortical junction, we measured synaptic efficacy, quantified as the number of spikes
390   within a 1 - 3 ms window of the SC CCG divided by the total number of VPm spikes used in the
391   analysis (Swadlow and Gusev, 2001). Synaptic efficacy was computed per stimulus in the
392   adapting train using an 80 ms post-stimulus response window.
393

Measuring VPm synchrony. To measure the level of synchronization in the VPm, synchrony was
computed for all pairs of simultaneously recorded VPm neurons that responded to the same
principal whisker. Synchrony was quantified using the total number of spikes within the central ±
7.5 ms window of the CCG generated from stimulus-evoked spiking in the 30 ms response window
divided by the total number of spikes in the raw CCG (Borden et al., 2017; Wright et al., 2021).
When applicable, normalized synchrony was normalized to the maximum response to the first
three stimuli.
401

Receiver operator characteristic (ROC) analysis. To evaluate the functional consequences of the
effects of adaptation over timescales, the theoretical detectability of sensory features was
assessed for significantly responsive RS and FS neurons by applying ideal observer analysis
(Wang et al., 2010b; Ollerenshaw et al., 2014; Whitmire and Stanley, 2016). The noise distribution
was generated from baseline activity, or the spike count 30 ms before the stimulus onset in each
trial. For each cortical neuron response to a stimulus at a given velocity, the signal came from a
probability distribution estimated using the across-trial mean (μ) and standard deviation (σ) of the
spike count per stimulus in the 30 ms neural response window. A parameterized 'population'
distribution was constructed for each neuron using a Gamma distribution, Γ (N*α, Θ), where α =
μ^2/σ^2, $\Theta = \sigma^2/\mu$, and N is the assumed number of identical neurons that share response properties
similar to the experimentally recorded neuron. Results were reported using N = 10 but results
were qualitatively similar for N = 1, 5, and 20. 1000 samples (trials) were drawn from the
'population' distribution.
415

ROC curves were generated by sliding the threshold across the noise and signal distributions,
generating a range of false alarm rates versus hit rates that reflect the degree of overlap of the
distributions. The area under the ROC curve (AUROC) was used as a performance metric for
theoretical detectability. Note that higher AUROC values indicate better performance as an
AUROC value of 1 corresponds to zero overlap between signal and noise distributions (an ideal

detector). An AUROC value of 0.5 corresponds to complete overlap between distributions, hence
operating at chance.
423

For theoretical discriminability, the ability of an ideal observer to discriminate between five
different velocities (50, 300, 450, 600, 1200 deg/s) was quantified based on the observed S1
activity. S1 recordings of L4 RS and L4 FS neurons with at least 20 stimulus trials per velocity
condition were used to generate a pool of 205 L4 RS neurons and 146 L4 FS neurons. Stimulus
trials, comprising a minimum of 20 trials per condition, were amalgamated from randomly sampled
cortical neurons. These datasets were employed to construct spike count distributions, which
were subsequently parameterized using a Gamma distribution. The resulting distributions were
input into Linear Discriminant Analysis (LDA) classifiers using MATLAB's 'fitcdiscr' command to
evaluate theoretical discrimination performance. This process was iterated 1000 times for each
stimulus condition. We conceptualize the ideal observer as an LDA decoder employing maximum
likelihood (ML) estimation to allocate an observed response to a specific stimulus velocity. The
accuracy of this classification process forms a performance matrix. Given the five stimulus
velocities employed here, the performance matrix is a 5x5 matrix where the (i,j) element
represents the probability of assigning the observed response to stimulus s_j when the actual
stimulus is s_i. The theoretical discrimination performance was quantified as the sum of all diagonal
elements of the performance matrix. Given that the five velocities were presented with equal
probability, achieving chance discrimination corresponds to a probability of 0.2.
441

442   *Statistical analyses.* All comparisons of measures of the neuronal response across stimulus
443   conditions were assessed with the two-sided Wilcoxon signed rank test with a Bonferroni
444   correction to correct for multiple comparisons.
445

446   **Experimental design**
447   This study consists of a within-subject design such that all comparisons are between the same
448   set of neurons, representative of multiple animals of both sexes (see Animals above).
449

450   **Data accessibility**
451   Data will be made available upon publication of this article.
452

RESULTS

Differential rapid sensory adaptation in VPm and S1

To assess the effects of adaptive sensory inputs on thalamocortical activity, we recorded extracellular neuronal spiking in the vibrissa lemniscal pathway of lightly isoflurane-anesthetized, head-fixed mice using multi-channel silicon probes. We measured the activity of populations of well-isolated, single neurons in the VPm of the thalamus as well as layer 4 of the primary somatosensory "barrel" cortex (S1 L4). S1 neurons were further parsed into putative excitatory regular spiking (RS) and putative inhibitory fast spiking (FS) neurons based on the trough-to-peak time of the spike waveform (Figure 1A; see Methods). We assessed the neuronal responses to a 10 Hz pattern of deflections of a single whisker on the contralateral side of the face using a computer-controlled actuator for up to 10 seconds (each deflection is a transient, 300 deg/s sawtooth whisker deflection; see Methods).

The grand-averaged peri-stimulus time histograms (PSTHs) in Figure 1B show the effects of sensory adaptation induced by 10 Hz repetitive stimulation on the spiking activity of example VPm, S1 L4 RS, and L4 FS neurons. As qualitatively observed for these example neurons, the sensory response of most recorded neurons in each neuronal population decreased over the timecourse of the repetitive stimulation. However, some neurons exhibit no overall change in stimulus-evoked spiking, and others an increase, as demonstrated in VPm by the overlaid PSTHs of the 1st and 100th stimuli in the adapting stimulus train in Figure 1C. All sensory responsive VPm and S1 L4 neurons were therefore parsed into three distinct categories - increased firing, decreased firing, and no change in firing - based on the cumulative spiking evoked within 30 ms of the 1st and 100th deflection of the adapting stimulus. We found that 66%, 95%, and 83% of VPm, RS, and FS neurons, respectively, decreased their sensory response between the 1st and 100th deflection (Figure 1D).

Despite some variability across neurons, we found that most responsive neurons across these regions responded most vigorously to the first few deflections in the stimulus train and adapted strongly to repetitive sensory stimuli. This is evident in the normalized population-averaged evoked response, calculated for each deflection in the train as the trial-averaged number of spikes in the 30 ms response window following the deflection (Figure 1E). The sensory responses were normalized by the un-adapted response, or the maximum response to the first three deflections, hereafter referred to as the control response. For each population, the response amplitude was reduced between 0 and 1.5 seconds, with a more gradual reduction in responsiveness from 1.5 and 10 seconds. Hence, throughout the rest of the study, we exclusively analyzed neurons whose sensory responses decreased between the control and the last stimulus in the 10 second train.

As further evidenced in Figure 1E, L4 RS and FS neurons adapted to a greater degree than VPm neurons in terms of their normalized response amplitude. Consistent with previous studies (Simons and Carvell, 1989; Gabernet et al., 2005; Heiss et al., 2008; Wright et al., 2021), the adaptation was even more pronounced in the RS population compared to both the VPm and the FS population when considering the adaptive decrease relative to the control sensory response.

13

To quantify this, we calculated the adaptation ratio, or the ratio of the mean spike count in
response to the last three stimuli in the repetitive stimulus train divided by the mean spike count
in response to the first three stimuli. For the first 1.5 seconds of the adapting train, the adaptation
ratios were 0.70, 0.56, and 0.65 for the VPm, RS, and FS populations, respectively. At the 10
second time point in the adapting train, the adaptation ratios decreased to 0.48, 0.34, and 0.41
(Figure 1F). Given these observations, we sought to further investigate the adaptation dynamics
in both S1 neuron subtypes and VPm thalamus at three time points of interest: the control time
point as well as both the 1.5 and 10 second time points.
503
Multiple timescales underlie the adaptation dynamics of cortical neurons in response to
repetitive sensory stimulation
Figures 2A and 2B show the grand-averaged PSTHs for the population of L4 RS and FS neurons,
respectively (note the different vertical scales for RS versus FS). Generally, both cortical sub-
populations generated short-latency, transient spiking in response to the control stimulus. There
was a qualitatively observable increase in latency and decrease in spiking response as a function
of the duration of the adapting stimulus. Indeed, the mean first spike latency for the L4 RS
population increased from 12.2 msec to 19.3 msec between the Control and 1.5 second time point,
with a further increase to 20.2 msec at the 10 second time point, corresponding to a 58% and 5%
increase in latency between the time points. For the L4 FS population, the first spike latency
increased from 10.35 msec to 16.51 msec to 18.09 msec, corresponding to a 59% and 10%
increase in latency between the time points (Figure 2C). Similarly, the mean evoked response at
the three time points of interest decreased from 0.50 to 0.22 to 0.14 spks/stim for the L4 RS
population, corresponding to a 56% and 36% decrease in spiking response between the time
points and from 0.72 to 0.41 to 0.30 spks/stim for the L4 FS population, corresponding to a 43%
and 27% decrease in spiking response between the time points (Figure 2D).
520
Given the non-saturating nature of the observed adaptation dynamics, we quantified the
timescales of the decay in responsiveness of the cortical populations by fitting exponential models
to the normalized population-averaged mean evoked response as a function of stimulus duration
(see Methods). Although adaptation is widely characterized as a simple, single exponential decay
(Chung et al., 2002; Khatri et al., 2004; Patterson et al., 2013; Kheradpezhouh et al., 2017), we
found that the declining profile of the adaptation responses over the 10 second repetitive
stimulation was better characterized with bi-exponentials (RS: R^2 = 0.99; FS: R^2 = 0.99) compared
to single exponential functions (RS: R^2 = 0.70; FS: R^2 = 0.72; Figures 2E & 2F). This trend was
even more evident when plotted on semi-log axes, where the plot reveals two strikingly clear
linear regimes with negative slopes, suggesting two distinct timescales of exponential decay
(figure insets).
532
533   **Multiple timescales underlie the adaptation dynamics of thalamic firing rate and population**
534   **synchrony in response to repetitive sensory stimulation**
535   The clear multi-timescale adaptation dynamics observed in S1 opens the possibility of multiple
536   factors contributing to these observed cortical dynamics. A likely important contributor is the
537   spiking patterns of the VPm neurons that provide the direct input to the cortical layer 4 neurons,

14

in both the mean firing rate and population synchrony of this highly convergent input. Towards
characterizing the adaptation dynamics in VPm, Figure 3A shows the grand-averaged PSTHs for
the VPm population in response to the adapting sensory stimulus at the three time points of
interest. Like S1 neurons, VPm neurons generated short-latency, transient spiking in response to
the Control stimulus with shifts in latency and spiking response from the Control to the 1.5 second
and 10 second time points. When quantifying these shifts for the population, the mean first spike
latency increased from 9.16 to 12.88 to 14.01 msec for the three time points of interest (Figure
3B). These shifts in latency correspond to a 40% increase from Control to the 1.5 second time
point, and a further increase of 8.75% between the 1.5 second the 10 second time points.
Additionally, the mean evoked response decreased from 0.81 to 0.52 spks/stim between the
Control and 1.5 second time point, with an additional decrease to 0.38 spks/stim from the 1.5
second to the 10 second time point (Figure 3C), corresponding to 36% and 27% decreases in
sensory response, respectively. Taken together, adaptation dynamics in the VPm also did not
saturate between 1.5 to 10 seconds, but rather continued to progress throughout the duration of
the adapting stimulus.
553
To characterize the timescale of the decay in responsiveness of VPm neurons, we again fit with
exponential models the normalized population-averaged mean evoked response as a function of
stimulus duration. We found that the declining profile of the adaptation response over the 10
second repetitive stimulation was better characterized with a bi-exponential (R^2 = 0.99) compared
to a single-exponential function (R^2 = 0.85; Figure 3D). Again, the bi-exponential nature of the
decay became strikingly clear when plotting the normalized evoked responses on semi-log axes,
highlighting two linear regions (figure insets).
561
A number of studies have shown that S1 can be highly sensitive to changes in the synchrony of
the convergent thalamic inputs (Pinto et al., 2000; Swadlow and Gusev, 2001; Bruno and
Sakmann, 2006; Wang et al., 2010b, 2010a; Ollerenshaw et al., 2014). We thus sought to also
characterize the adaptation profile of VPm synchrony. We quantified the synchrony between
every pair of simultaneously recorded neurons having maximal responsiveness to the same
whisker (i.e. likely in the same thalamic barreloid). Synchrony was computed by generating a
cross-correlogram (CCG) and measuring the occurrence of coincident spikes within a +/- 7.5 ms
window, normalized by the number of spikes used in the analysis (see Methods, Figure 3E). This
normalization accounts for the overall firing rate which is important given that synchrony and firing
rate naturally covary. First examining changes in synchrony in response to the adapting stimuli at
the three time points of interest, the VPm population synchrony decreased from 0.89 to 0.81 to
0.76 (unitless quantity, see Methods, Figure 3F). Further, as with the mean firing rate, the
synchrony evolved in a bi-exponential manner (R^2 = 0.92) and was not as well fit with a single
exponential (R^2 = 0.75; Figure 3G).
576
Multiple timescales underlie the adaptation dynamics of thalamocortical synaptic efficacy
in response to repetitive stimulation
The above analyses suggest a relationship between cortical L4 neurons and the VPm neurons
from which the L4 neurons receive direct sensory input, but do not measure this relationship

explicitly. To evaluate the dynamics of the transformation across the thalamocortical synapse, we
next examined the effects of the adapting stimulus on synaptic efficacy of putatively connected
S1 pairs recorded simultaneously using previously established methods for identifying
monosynaptic connectivity in-vivo (Liew et al., 2021). The left panel of Figure 4A shows raster
plots of example topographically aligned VPm, L4 RS, and L4 FS neurons that were
simultaneously recorded in response to a weak, sinusoidal input stimulus. Note that the overall
firing rate of all the neurons was elevated but not generally stimulus-locked; this is critical for
identifying putatively connected pairs. The spiking evoked by this weak sinusoidal stimulus was
used to generate raw CCGs between every possible pair of topographically aligned VPm-L4 RS
neurons. Shuffled CCGs were generated by randomizing the trials of the VPM and S1 spiking
activity relative to each other, effectively correcting for any stimulus-induced elements of the CCG.
Finally, shuffle-corrected (SC) CCGs were generated by subtracting the shuffled CCGs from the
raw CCGs (see Methods). As is the case for the SC CCGs of the example pairs of neurons shown
in the right panel of Figure 4A , a narrow peak within the 1-3 msec window with prominence
greater than 3.5 standard deviations of the shuffled CCG suggests monosynaptic connectivity
(see Methods; Liew et al., 2021). Subsequent analyses were restricted to putatively connected
thalamocortical neurons.
598
Responses to the 10 Hz adapting stimulus for the same example VPm, L4 RS, and L4 FS neurons
in Figure 4A are shown in Figure 4B. For each of the three time points of interest, we assessed
the synaptic efficacy of each connected thalamocortical pair, defined as the probability of an S1
neuron spike following a spike of the pre-synaptic VPm neuron. More specifically, using the
spiking activity in a 80 msec window following each of the time points of interest in the repetitive
stimulus, synaptic efficacy was calculated as the number of synchronous spikes in the SC CCG
within the 1-3 ms window following each VPm spike divided by the total number of spikes in the
SC CCG (Swadlow and Gusev, 2001).
607
Figures 4C and 4D show a clear drop in the peak of the grand-averaged SC CCG generated from
the spiking responses at the three time points of interest in the 10 Hz repetitive stimulation for
VPm-L4 RS and VPm-L4 FS connected pairs, respectively. Indeed, synaptic efficacy decreased
across the three time points for both VPm-L4 RS and VPm-L4 FS, summarized in Figures 4E and
4F. Note that the apparent decrease in synaptic efficacy from 1.5 to 10 seconds for the VPm-L4
FS pairs was not statistically significant. When plotting the synaptic efficacy over the timecourse
of stimulation, we observe a decaying trend, with an early rapid drop of 81% and 37% at the 1.5
second time point compared to control for the VPm-L4 RS and VPm-L4 FS pairs, respectively.
Following this early drop, synaptic efficacy gradually declined during the latter part of stimulation,
corresponding to a 50% and 5% drop at the 10 second time point compared to the 1.5 second
time point for the VPm-L4 RS and VPm-L4 FS pairs, respectively. Here again, we observe two
distinct timescales, as evident in the semi-log plot insets (Figures 4G and 4H). It is important to
note that since the above measures of synaptic efficacy were obtained during sensory inputs that
synchronize the thalamic population, the efficacy measures could be influenced by the overall
drive and synchronization of the thalamic input, as well as depression at the thalamocortical
synapse (see Discussion).

624
Direct optogenetic drive of VPm thalamus eliminates longer timescale adaptation in VPm
In our next set of experiments, we sought to explore the relationship across the thalamocortical
junction through a more direct manipulation of VPm itself. Specifically, we optogenetically
stimulated VPm neurons in NR133 x Ai32 mice which enables selective expression of
Channelrhodopsin (ChR2) in VPm neurons (Figure 5A). Using an optrode lowered into VPm, we
presented pulsatile optogenetic stimulation of identical frequency and duration to the sensory
stimulation: 10 Hz for 10 seconds. The LED pulse waveform was designed to elicit a response
that mimics the control VPm sensory response (Figure 5B; see Methods).
633
As shown by the grand-averaged PSTHs in Figure 5C, repetitive optogenetic stimulation
produced a decrease in response in VPm neurons between the control and 1.5 second time point
in the repetitive optogenetic stimulation that was similar to that observed for adaptation with the
sensory stimulus (Figure 3A). However, in contrast to the sensory adaptation, qualitatively there
was a less clear difference in response between the 1.5 and 10 second time points. In response
to the optogenetic stimulation at the three time points of interest, the evoked VPm population firing
rate decreased from the Control to the 1.5 second condition but did not further decrease between
the 1.5 second and 10 second conditions (Figure 5D). We characterized the timescale of the
decay in responsiveness of VPm neurons by again fitting with exponential models the normalized
population-averaged mean evoked optogenetic response as a function of stimulus duration. This
time, we found that the declining profile was better characterized with a single exponential (R^2 =
0.70). Naturally, due to the added degree of freedom, the R^2 value for a bi-exponential is larger
(R^2 = 0.71). However, this increase in R^2 is marginal and negligible. The single exponential nature
of the decay becomes strikingly clear when plotting the normalized evoked responses on semi-
log axes, highlighting only one linear region with a slope less than 0, suggesting a single
exponential with a constant offset (Figure 5E).
650
Similarly, in response to repetitive optogenetic stimulation, VPm synchrony decreased from
Control to the 1.5 second adapted conditions (also similar to that for sensory adaptation – Figure
3F), but no differences were detected between the 1.5 second and 10 second adapted conditions
(Figure 5F). In addition, the timecourse of VPm synchrony to repetitive optogenetic stimulation
was well fit with a single exponential and not improved with a second exponential, further
evidenced by the semi-log insets (Figure 5G; 1-exp: R^2 = 0.48; 2-exp: R^2 = 0.48).
657
Optogenetic drive of VPm results in a loss of longer timescale adaptation in S1
Having designed a condition that eliminates the longer timescale adaptation in thalamus such that
VPm synchrony and firing rate rapidly decayed before stabilizing for the remainder of the repetitive
stimulus, we then considered the dynamics in S1. Figures 6A and 6B show the grand-averaged
cortical responses to optogenetic stimulation of VPm. The L4 RS and FS neurons adapted
strongly to the repetitive optogenetic stimuli. At the three time points of interest, both the L4 RS
and FS neuron populations' firing rate decreased from the Control to the 1.5 second time point
but did not further decrease from the 1.5 second to 10 second time point (Figures 6C and 6D,
respectively). As evident in the normalized population-averaged mean evoked response in both

17

cortical populations, the reduction in cortical response over the 10 second timecourse was well characterized by a single timescale, with marginal gain for a second exponential (Figure 6E and 6F; 1-exp: R^2 = 0.78 for L4 RS and R^2 = 0.81 for L4 FS; 2-exp: R^2 = 0.78 for L4 RS and R^2 = 0.83 for L4 FS). When visualizing the timecourse on semi-log axes for both L4 RS and FS populations, a distinct pattern emerges: a single linear region is followed by a relatively flat line. This observation indicates that the decay in evoked responses occurred within a single timescale.

Through these control experiments, we find that optogenetic drive of VPm that eliminates the longer timescale adaptation in both firing rate and synchrony in the VPm neuron population also eliminates the longer timescale adaptation in S1 layer 4 neurons. This suggests that the previous observations of the longer timescale adaptation in S1 neurons during sensory drive are inherited from VPm thalamus, consistent with a growing notion of VPm as the source of much of the adaptation observed in S1 (Wright et al., 2021).

Adaptation over multiple timescales degrades stimulus detection for an ideal observer of cortical activity
Having explored thalamocortical activity across multiple timescales of sensory adaptation, we next sought to investigate the functional consequences of these timescales. In previous studies of the vibrissa pathway, we have shown that repetitive stimulation lasting as brief as 1-2 seconds is sufficient to not only reduce the overall responsiveness of the cortical population but shift the neuronal responses towards an adaptive trade-off that favors sensory discrimination over detection (Wang et al., 2010b; Ollerenshaw et al., 2014). To determine if this holds true for longer timescales of adaptation, we probed cortical responses either in the presence or absence of a preceding adapting stimulus and leveraged signal detection theory to ascertain the ability of an ideal observer to detect a signal (stimulus present) versus noise (stimulus absent; Figure 7A). Although the results in Figure 2 regarding the decrease in the magnitude of the sensory evoked cortical response would suggest a corresponding decrease in detectability, this is ultimately determined by both the magnitude of the response and the corresponding variability, relative to ongoing cortical activity. To directly test this, the cortical response to a 300 deg/s punctate stimulus and the corresponding detectability were evaluated in isolation (Control) and in the presence of preceding 10 Hz adapting stimuli of 1.5 second or 10 second durations.

To quantify cortical sensory responses under the signal detection framework, we constructed parameterized probability distributions of the signal and noise spike counts resulting from the probe stimulus preceded by the Control and adapting stimulus conditions. The signal distributions were estimated by drawing samples from gamma distribution fits of the observed cortical spike counts within a 30 ms post-stimulus response window for a 300 deg/s stimulus. The noise distributions were estimated from baseline activity 30 ms prior to stimulus onset (see Methods). We find that adaptation pushed the signal distribution of the evoked response closer to the noise distribution at each time point of interest for both L4 RS and FS populations (Figures 7B & 7C, respectively).

To predict the potential impact of adaptation on stimulus detectability, we quantified the extent to
which an ideal observer of cortical spiking activity accurately detects the presence of a signal from
background noise when the signal is preceded by the Control and adapted conditions. Thus, we
applied a sliding threshold on the parametric distributions to generate a Receiver Operator
Characteristics (ROC) curve which shows the hit rate vs the false alarm rate for an ideal observer
of the neuronal activity. For ROC curves, a perfect diagonal indicates that the ideal observer is
operating at chance, and deviations above the diagonal indicate improved performance. Figures
7D & 7E show the ROC curves for an example L4 RS and FS neuron, respectively, for each of
the three stimulus conditions. For both populations, adapting stimuli preceding the probe stimulus
brought the ROC curve closer to the diagonal, reflective of a loss in theoretical detection
performance. This loss was greatest when the probe stimulus was preceded by 10 seconds of an
adapting stimulus.
721
To quantify this loss in theoretical detection performance, we measured the area under the ROC
curve (AUROC) for each neuron. An AUROC value of 1 corresponds to zero overlap between
signal and noise distribution (an ideal detector). An AUROC value of 0.5 corresponds to complete
overlap between distributions, hence the ideal observer operating at chance. We found that the
detectability of both RS and FS neurons significantly decreased when preceded by Control versus
1.5 second adapting stimuli, and further decreased when preceded by 10 second adapting stimuli
(Figures 7F & 7G). Consistent with the observation in Figure 1F that RS neurons had lower
adaptation ratios than FS neurons, we found that the theoretical detectability was generally lower
for RS neurons as compared to FS neurons.
731
Adaptation over multiple timescales enhances stimulus discrimination for an ideal
observer of cortical activity
We next evaluated the functional consequences of adaptation on stimulus discrimination across
multiple timescales by presenting probe stimuli of varying velocities (50, 300, 450, 600, or 1200
deg/s), preceded by either Control, 1.5 second, or 10 second adapting stimuli. First, in terms of
the impact of adaptation on the velocity sensitivity properties in the cortex, adaptation attenuated
the response magnitude across all velocities for cortical populations, although differently for
different velocities. The velocity sensitivity was further reduced for both the RS and FS
populations when the probe stimuli were preceded by the 10-sec10 second adapted stimulus
compared to the 1.5 second adapted stimulus (Figures 8A and 8B).
742
Taking the perspective of an ideal observer, we examined the extent to which adaptation over
multiple timescales affects discrimination between whisker deflections at different velocities.
Shown in Figures 8C and 8D are parametric fits of the experimentally observed spike count
distributions for the RS and FS populations, respectively. Qualitatively, the populations' spike
counts were relatively high in the Control conditions, but with highly overlapped and unordered
distributions. However, while increasing pre-stimulus adaptation attenuated the population spikes
counts, the distributions became more separated and ordered (Figure 8C & 8D).
750

To quantify this effect, we compared the performance of the ideal observer in discriminating
between different deflection velocities when preceded by Control, 1.5 second and 10 second
adapting stimuli. For each stimulus condition, we randomly selected 25 neurons from each L4 RS
and L4 FS population and combined stimulus trials to form sensory response distributions of the
cortical responses to various velocities (parameterized gamma distributions, see Methods). To
assess discrimination performance, we implemented a holdout validation approach, reserving
data from 20% of trials for testing while using the remaining 80% of trials for training. Linear
discriminant analysis (LDA) classifiers were then applied to infer the most likely velocity presented
based on the observed cortical response, and discrimination performance was evaluated by the
accuracy of the ideal observer in making correct stimulus classifications (see Methods). This
process was iterated 1000 times to comprehensively assess discrimination performance for each
stimulus condition (Control, 1.5 second, and 10 second adapted) across various combinations of
neurons and trials. Given that all five velocities were presented with equal probability, a value of
0.2 reflects the ideal observer operating at chance. We found that increasing adaptation
significantly improved overall discrimination performance in both RS and FS populations (Figure
8E & 8F), across the early and later stages of the adaptation. Taken together, our results suggest
that adaptation over multiple timescales continues to facilitate better discrimination performance
by reducing the overlap between stimulus distributions by trading off the overall sensitivity to
repetitive stimulation.
770
771

DISCUSSION

In this study, we discovered two clear and distinct timescales of rapid adaptation across the thalamocortical circuit over the course of seconds. Several other studies have examined the properties of adaptation over a range of timepoints using various durations of repetitive stimulation (Dragoi et al., 2000; Patterson et al., 2013; Latimer et al., 2019), white noise stimulation, (Fairhall et al., 2000; Lundstrom et al., 2010) and modeling approaches (Latimer and Fairhall, 2020). Here, the ability to record simultaneously from the thalamic VPm population as well as both the cortical L4 RS and FS populations allowed us to precisely examine transformations of adaptation across the thalamocortical junction through the spiking response amplitudes, thalamic population synchrony, and synaptic efficacy.

Numerous studies have identified potential circuit mechanisms that could underlie the differential adaptation from thalamus to cortex. Potential drivers of the observed rapid cortical sensory adaptation include depression of thalamocortical (TC) synapses (Chung et al., 2002) and/or synaptic depression/facilitation of excitatory/inhibitory intracortical (IC) projections (Beierlein and Connors, 2002; Reig et al., 2006), and neuronal population mechanisms such as thalamic firing rate and population synchrony, all of which may collectively shape the IC balance of excitation and inhibition. One prominent view is that short-term depression of thalamocortical synapses plays a dominant role in this type of cortical adaptation (Chung et al., 2002; Castro-Alamancos, 2004b; Katz et al., 2006). This was experimentally demonstrated using selective depression of TC and IC synapses via electrical stimulation in simultaneous, paired extracellular recordings under anesthesia (Chung et al., 2002). However, the rapid synaptic depression of the TC synapse reported in Chung et al., 2002 does not rule out the contribution of thalamic population synchrony observed here in shaping S1 adaptation, which was not measured in the relevant studies. Further, subsequent studies suggest that thalamocortical desynchronization plays perhaps an even more prominent role than short-term changes in the TC or IC synapses (Wright et al., 2021). In fact, many lines of evidence have demonstrated the profound effect that thalamic spike timing and synchrony has on S1 processing either under anesthesia (Pinto et al., 2000; Wang et al., 2010b; Whitmire and Stanley, 2016) or wakefulness (Wright et al., 2021; Borden et al., 2022). In the context of this study, the optogenetic control experiments do not rule out the possibility of TC synaptic depression playing a role in the fast timescale adaptation, but previous findings (Wright et al., 2021) suggest that this role may be overshadowed by the adaptive reductions in thalamic firing rate and synchrony that are also clearly present in the adaptation at fast timescales (0-1.5 seconds). What the optogenetic control experiments here do show is that the monotonic slow desynchronization of the thalamic population beyond the first few seconds (2-10 seconds) seems to play a dominant role in modulating cortical adaptation at longer timescales, with little or no influence from TC synaptic depression on this timescale. These results together suggest the possibility of a complex interaction between distinct mechanisms, and that multiple mechanisms are likely co-activated over distinct periods of time to contribute to the cellular dynamics that we observed.

It is important to note that the adaptive shift in excitation-inhibition (E-I) balance in the cortex has
also been explored extensively. The differential adaptation of the putative excitatory and inhibitory
cortical neurons observed here likely shapes overall cortical excitability and the integration
properties of ascending sensory signals from thalamus. Importantly, this effect has been shown
during wakefulness, but the extent of the differential adaptation is strongly shaped by the nature
of the adapting stimulus itself (Wright et al., 2021). The more profound depression of intracortical
inhibitory synapses as compared to excitatory synapses during persistent stimulation has been
shown to skew the E-I balance towards excitation (Gabernet et al., 2005; Heiss et al., 2008). This
reduction in feedforward inhibition was shown to expand the cortical window of integration and
affect how inputs are summed across multiple timescales of adaptation. This mechanism could
potentially regulate the effects of the thalamic desynchronization observed here, offsetting the
loss of thalamic synchrony through a loss in cortical sensitivity to thalamic synchrony. However,
in previous work, we have shown that the desynchronization observed in thalamus during
adaptation outweighs the changes in cortical integration (Wang et al., 2010b).

This study highlights the important contributions of the rapid adaptation of thalamus in shaping
cortical adaptation. What is the origin of the adaptative properties observed in sensory thalamus?
For the fastest timescale component of the VPm adaptation, given the high firing rates of neurons
in the brainstem (Ahissar et al., 2000), it is likely that depression of the synaptic inputs from
brainstem to thalamus play an important role in the rapid decrease in thalamic firing rate, as well
as the desynchronization in the thalamic population at this timescale. One important finding here
was that the longer timescale adaptation observed in S1 appears to be largely inherited from VPm.
When VPm thalamus was optogenetically driven, this eliminated the slow decrease in thalamic
firing and synchronization (Figure 5), which subsequently eliminated the slower, later phase of
adaptation in S1 (Figure 6). One distinct possibility is that the direct optogenetic drive of VPm
overcomes intrinsic properties of the thalamic neurons that are responsible for the slow drift. In
brain slice experiments, thalamic neurons have been shown to exhibit a slow after-
hyperpolarization that persists for several seconds after controlled depolarization (Goaillard and
Vincent, 2002). The light activation of Channelrhodopsin in the thalamic neurons likely overcomes
these intrinsic cell properties, eliminating the slow after-hyperpolarization and the corresponding
decrease in firing rate and synchronization of the thalamic population. However, further
experiments are needed to more fully understand this process in-vivo and to better understand
the interaction between intrinsic properties of thalamic neurons and the ascending input from the
brainstem.

Given the nature of how single empirical phenomena often arise from multiple contributing
biophysical mechanisms, studying adaptation across multiple timescales is not without its
challenges. In this study, we were able to capture multiple timescales of adaptation using a 10 Hz
repetitive stimulation over 10 seconds. It is important to note, however, that adaptation is a change
in function that takes time to develop and also dissipate. It has been shown that the temporal
scales over which adaptive dynamics build up and recover are distinct, and also highly dependent
upon the duration of exposure to repetitive stimulation (Dragoi et al., 2000; Chung et al., 2002;
Patterson et al., 2013). In general, brief exposure produces adaptation that develops and recovers

rapidly (Bonds, 1991; Nelson, 1991; Müller et al., 1999), while more prolonged stimulation can
result in slower and more lasting forms of changes (Greenlee et al., 1991). Additionally, thalamic
and cortical neurons also exhibit diverse timescales of responsiveness to repeated stimulation
(Chung et al., 2002). Therefore, as a rule of thumb, an inter-trial interval of at least 4 times the
duration of adapting stimulus train (10 seconds) was utilized in this study, while acquiring at least
100 trials of repeated measurements of the adaptation dynamics. As a control, we also assessed
the difference between the magnitude of the response to the first deflection of the adapting train
at the beginning and end of a stimulus block. We did not observe a significant difference in this
measure, suggesting that the majority of the neurons returned to the baseline condition prior to
the initiation of the next trial, although longitudinal effects cannot be completely ruled out. For a
subset of the data, we also probed the recovery for each thalamic and cortical neuron and found
that most of the cells showed a full recovery to the baseline condition.
869
Given that this study was performed in the somatosensory pathway of isoflurane-anesthetized
mice, it is important to acknowledge some of the limitations in the findings. The large majority of
in-vivo studies of the relative role of short-term depression of thalamocortical and intracortical
synapses have been conducted under anesthesia (Heiss et al., 2008; Chung et al., 2019), and
the current study cannot rule out the potential influence of TC synaptic depression in the early,
fast timescale component of adaptation. In contrast, the adaptive changes in synchronous
thalamic firing have recently been reported to play a more prominent role than synaptic
depression in the TC and IC synapse under wakefulness (Wright et al., 2021). One possibility is
that the overlap between distinct mechanisms during wakefulness is smaller compared to the
condition under anesthesia. It was reported that the thalamocortical circuit may be in an elevated
baseline firing state during wakefulness (Castro-Alamancos, 2004a), hence the suggestion that
TC synapses might already be depressed in this state. When TC synaptic depression is fully
engaged during the high thalamic firing rates associated with wakefulness, thalamic
desynchronization may dominate and exert an even greater effect on regulating the nature of
information conveyed downstream to cortical targets. However, to fully disentangle these effects
and the relative contribution from each mechanism, further techniques would need to be
employed, such as selective optogenetic manipulation of TC and CC synaptic depression
combined with simultaneous, multi-site recordings, beyond the scope of the current study.
888
Finally, the results here suggest that although multiple mechanisms may interact and transition
dynamically at various timescales, adaptation over the timecourse that we examined in this study
improve theoretical discrimination at the expense of detection. This similarity in function supports
the possibility of a unifying framework across disparate mechanisms and timescales, which is to
allow the brain to maintain a flexible representation in response to stimulus statistics that span a
wide range in our natural environment. Moreover, these results further support Barlow's efficient
coding hypothesis such that the reduction in responsiveness of neuronal response in the TC
circuit is not a negative attribute that signifies a loss of sensitivity to stimuli, but instead a
mechanism that could mediate an adaptive trade-off in coding strategy to put the brain, in this
case the primary sensory cortex, in a state that optimizes neural coding efficiency (Barlow, 1961).
899

REFERENCES:

Adibi M, Clifford CWG, Arabzadeh E (2013) Informational basis of sensory adaptation: entropy and single-spike efficiency in rat barrel cortex. J Neurosci Off J Soc Neurosci 33:14921–14926.

Adibi M, Lampl I (2021) Sensory Adaptation in the Whisker-Mediated Tactile System: Physiology, Theory, and Function. Front Neurosci 15:770011.

Ahissar E, Sosnik R, Haidarliu S (2000) Transformation from temporal to rate coding in a somatosensory thalamocortical pathway. Nature 406:302–306.

Barlow H (1961) Possible Principles Underlying the Transformations of Sensory Messages. Sens Commun 1.

Barthó P, Hirase H, Monconduit L, Zugaro M, Harris KD, Buzsáki G (2004) Characterization of neocortical principal cells and interneurons by network interactions and extracellular features. J Neurophysiol 92:600–608.

Beierlein M, Connors BW (2002) Short-term dynamics of thalamocortical and intracortical synapses onto layer 6 neurons in neocortex. J Neurophysiol 88:1924–1932.

Bonds AB (1991) Temporal dynamics of contrast gain in single cells of the cat striate cortex. Vis Neurosci 6:239–255.

Borden PY, Ortiz AD, Waiblinger C, Sederberg AJ, Morrissette AE, Forest CR, Jaeger D, Stanley GB (2017) Genetically expressed voltage sensor ArcLight for imaging large scale cortical activity in the anesthetized and awake mouse. Neurophotonics 4:031212.

Borden PY, Wright NC, Morrissette AE, Jaeger D, Haider B, Stanley GB (2022) Thalamic bursting and the role of timing and synchrony in thalamocortical signaling in the awake mouse. Neuron 110:2836-2853.e8.

Bruno RM, Sakmann B (2006) Cortex is driven by weak but synchronously active thalamocortical synapses. Science 312:1622–1627.

Bruno RM, Simons DJ (2002) Feedforward Mechanisms of Excitatory and Inhibitory Cortical Receptive Fields. J Neurosci 22:10966–10975.

Castro-Alamancos MA (2004a) Absence of Rapid Sensory Adaptation in Neocortex during Information Processing States. Neuron 41:455–464.

Castro-Alamancos MA (2004b) Dynamics of sensory thalamocortical synaptic networks during information processing states. Prog Neurobiol 74:213–247.

Chung JE, Joo HR, Fan JL, Liu DF, Barnett AH, Chen S, Geaghan-Breiner C, Karlsson MP, Karlsson M, Lee KY, Liang H, Magland JF, Pebbles JA, Tooker AC, Greengard LF, Tolosa VM, Frank LM (2019) High-Density, Long-Lasting, and Multi-region Electrophysiological Recordings Using Polymer Electrode Arrays. Neuron 101:21-31.e5.

Chung S, Li X, Nelson SB (2002) Short-term depression at thalamocortical synapses
contributes to rapid adaptation of cortical sensory responses in vivo. Neuron 34:437–
446.

Cruikshank SJ, Lewis TJ, Connors BW (2007) Synaptic basis for intense thalamocortical
activation of feedforward inhibitory cells in neocortex. Nat Neurosci 10:462–468.

Dimwamwa ED, Pala A, Chundru V, Wright NC, Stanley GB (2023) Dynamic corticothalamic
modulation of the somatosensory thalamocortical circuit during wakefulness. Available
at: http://biorxiv.org/lookup/doi/10.1101/2023.07.18.549491 [Accessed April 1, 2024].

Dragoi V, Sharma J, Sur M (2000) Adaptation-induced plasticity of orientation tuning in adult
visual cortex. Neuron 28:287–298.

Drew PJ, Abbott LF (2006) Models and Properties of Power-Law Adaptation in Neural Systems.
J Neurophysiol 96:826–833.

Fairhall A, Lewen G, Bialek W, van Steveninck R (2000) Multiple Timescales of Adaptation in a
Neural Code. In: Advances in Neural Information Processing Systems. MIT Press.

Franklin K, Paxinos G (2008) The mouse brain in stereotaxic coordinates, compact. The coronal
plates and diagrams. Elsevier Academic Press.

Gabernet L, Jadhav SP, Feldman DE, Carandini M, Scanziani M (2005) Somatosensory
integration controlled by dynamic thalamocortical feed-forward inhibition. Neuron
48:315–327.

Ganmor E, Katz Y, Lampl I (2010) Intensity-Dependent Adaptation of Cortical and Thalamic
Neurons Is Controlled by Brainstem Circuits of the Sensory Pathway. Neuron 66:273–
286.

Goaillard J-M, Vincent P (2002) Serotonin suppresses the slow afterhyperpolarization in rat
intralaminar and midline thalamic neurones by activating 5-HT7 receptors. J Physiol
541:453–465.

Greenlee MW, Georgeson MA, Magnussen S, Harris JP (1991) The time course of adaptation
to spatial contrast. Vision Res 31:223–236.

Heiss JE, Katz Y, Ganmor E, Lampl I (2008) Shift in the Balance between Excitation and
Inhibition during Sensory Adaptation of S1 Neurons. J Neurosci 28:13320–13330.

Katz Y, Heiss JE, Lampl I (2006) Cross-Whisker Adaptation of Neurons in the Rat Barrel Cortex.
J Neurosci 26:13363–13372.

Khatri V, Hartings JA, Simons DJ (2004) Adaptation in Thalamic Barreloid and Cortical Barrel
Neurons to Periodic Whisker Deflections Varying in Frequency and Velocity. J
Neurophysiol 92:3244–3254.

Kheradpezhouh E, Adibi M, Arabzadeh E (2017) Response dynamics of rat barrel cortex
neurons to repeated sensory stimulation. Sci Rep 7:11445.

Latimer KW, Barbera D, Sokoletsky M, Awwad B, Katz Y, Nelken I, Lampl I, Fairhall AL, Priebe
NJ (2019) Multiple Timescales Account for Adaptive Responses across Sensory
Cortices. J Neurosci 39:10019–10033.

Latimer KW, Fairhall AL (2020) Capturing Multiple Timescales of Adaptation to Second-Order
Statistics With Generalized Linear Models: Gain Scaling and Fractional Differentiation.
Front Syst Neurosci 14.

Lefort S, Tomm C, Floyd Sarria J-C, Petersen CCH (2009) The excitatory neuronal network of
the C2 barrel column in mouse primary somatosensory cortex. Neuron 61:301–316.

Liew YJ, Pala A, Whitmire CJ, Stoy WA, Forest CR, Stanley GB (2021) Inferring thalamocortical
monosynaptic connectivity in vivo. J Neurophysiol 125:2408–2431.

Lundstrom BN, Fairhall AL, Maravall M (2010) Multiple timescale encoding of slowly varying
whisker stimulus envelope in cortical and thalamic neurons in vivo. J Neurosci Off J Soc
Neurosci 30:5071–5077.

Maravall M, Alenda A, Bale MR, Petersen RS (2013) Transformation of adaptation and gain
rescaling along the whisker sensory pathway. PloS One 8:e82418.

Maravall M, Petersen RS, Fairhall AL, Arabzadeh E, Diamond ME (2007) Shifts in coding
properties and maintenance of information transmission during adaptation in barrel
cortex. PLoS Biol 5:e19.

Masri R, Bezdudnaya T, Trageser JC, Keller A (2008) Encoding of Stimulus Frequency and
Sensor Motion in the Posterior Medial Thalamic Nucleus. J Neurophysiol 100:681–689.

Mohar B, Ganmor E, Lampl I (2015) Faithful Representation of Tactile Intensity under Different
Contexts Emerges from the Distinct Adaptive Properties of the First Somatosensory
Relay Stations. J Neurosci 35:6997–7002.

Mohar B, Katz Y, Lampl I (2013) Opposite Adaptive Processing of Stimulus Intensity in Two
Major Nuclei of the Somatosensory Brainstem. J Neurosci 33:15394–15400.

Müller JR, Metha AB, Krauskopf J, Lennie P (1999) Rapid adaptation in visual cortex to the
structure of images. Science 285:1405–1408.

Nelson SB (1991) Temporal interactions in the cat visual system. III. Pharmacological studies of
cortical suppression suggest a presynaptic mechanism. J Neurosci Off J Soc Neurosci
11:369–380.

Ollerenshaw DR, Zheng HJV, Millard DC, Wang Q, Stanley GB (2014) The Adaptive Trade-Off
between Detection and Discrimination in Cortical Representations and Behavior. Neuron
81:1152–1164.

Pachitariu M, Sridhar S, Pennington J, Stringer C (2024) Spike sorting with Kilosort4 | Nature
Methods. Nat Methods.

Pala A, Petersen CCH (2015) In Vivo Measurement of Cell-Type-Specific Synaptic Connectivity
and Synaptic Transmission in Layer 2/3 Mouse Barrel Cortex. Neuron 85:68–75.

Patterson CA, Wissig SC, Kohn A (2013) Distinct Effects of Brief and Prolonged Adaptation on
Orientation Tuning in Primary Visual Cortex. J Neurosci 33:532–543.

Pinto DJ, Brumberg JC, Simons DJ (2000) Circuit dynamics and coding strategies in rodent
somatosensory cortex. J Neurophysiol 83:1158–1166.

Reid RC, Alonso JM (1995) Specificity of monosynaptic connections from thalamus to visual
cortex. Nature 378:281–284.

Reig R, Gallego R, Nowak LG, Sanchez-Vives MV (2006) Impact of Cortical Network Activity on
Short-term Synaptic Depression. Cereb Cortex 16:688–695.

Simons DJ, Carvell GE (1989) Thalamocortical response transformation in the rat vibrissa/barrel
system. J Neurophysiol 61:311–330.

Storchi R, Bale MR, Biella GEM, Petersen RS (2012) Comparison of latency and rate coding for
the direction of whisker deflection in the subcortical somatosensory pathway. J
Neurophysiol 108:1810–1821.

Swadlow HA, Gusev AG (2001) The impact of "bursting" thalamic impulses at a neocortical
synapse. Nat Neurosci 4:402–408.

Wang H-P, Spencer D, Fellous J-M, Sejnowski TJ (2010a) Synchrony of Thalamocortical Inputs
Maximizes Cortical Reliability. Science 328:106–109.

Wang Q, Webber RM, Stanley GB (2010b) Thalamic synchrony and the adaptive gating of
information flow to cortex. Nat Neurosci 13:1534–1541.

Whitmire CJ, Stanley GB (2016) Rapid Sensory Adaptation Redux: A Circuit Perspective.
Neuron 92:298–315.

Wright NC, Borden PY, Liew YJ, Bolus MF, Stoy WM, Forest CR, Stanley GB (2021) Rapid
Cortical Adaptation and the Role of Thalamic Synchrony during Wakefulness. J Neurosci
41:5421–5439.

1032

1033

 Figure captions:

1035

Figure 1. Differential rapid sensory adaptation in VPm and S1.
A. 32-channel silicon probe recordings were performed in the ventro-posteromedial nucleus of
the thalamus (VPm) and in layer 4 (L4) of the primary somatosensory cortex (S1) in lightly
isoflurane-anesthetized mice. The mean +/- sem waveforms of example well-isolated VPm,
putative excitatory L4 regular spiking (RS), and putative inhibitory L4 fast spiking (FS) typical of
each population are shown in green, orange, and pink, respectively. RS and FS cortical neurons
were identified based on the trough to peak time of their spike waveforms. Precise deflections
were delivered to single whiskers for sensory input using a computer-controlled actuator.
Adaptation was induced using a train of sawtooth punctate sensory stimuli (300 deg/s) delivered
at 10 Hz for 10 sec (see Methods). **B.** Peri-stimulus time histograms (PSTHs) of the example
neurons from A in response to 10 Hz repetitive sensory stimulation. **C.** Overlaid PSTHs of three
example VPm neurons' sensory responses to the 1st and 100th sensory stimulus in the 10 Hz
repetitive stimulus train (solid and dashed lines, respectively), each with differing response
profiles. The stimulus is delivered at time 0 sec in each plot. **D.** The distribution of responsive
neurons from each population that exhibited a decrease, an increase, or no change in sensory
response between the 1st and 100th sensory stimulus. Change was determined using the Wilcoxon
signed rank test on the per-trials spike count in the 30 ms neural response window for the 1st vs
100th stimulus in the repetitive train. **E.** Grand-averaged mean +/- sem evoked response for the
population of VPm, L4 RS, and L4 FS neurons to each stimulus in the 10 second repetitive
stimulus train, normalized to the maximum response of the first three stimuli (see Methods). **F.**
The mean +/- sem adaptation ratio for each neuron population after 1.5 and 10 sec of repetitive
stimulation, defined as the ratio of the mean spike count for the last three stimuli divided by the
mean spike count for the first three stimuli. (1.5 sec adapted: VPm: 0.70 ± 0.02, RS: 0.56 ± 0.02;
FS: 0.65 ± 0.02; VPm vs RS: p = 2.60e-07, VPm vs FS: p = 6.82e-02, RS vs FS: p = 3.96e-04;
10 sec adapted: VPm: 0.48 ± 0.02, RS: 0.34 ± 0.01; FS: 0.41 ± 0.02; VPm vs RS: p = 1.52e-12,
VPm vs FS: p = 8.25e-04, RS vs FS: p = 1.35e-03. All statistical comparisons were performed
using the Wilcoxon rank-sum test with a Bonferroni correction; *** indicates p < 3.3e-04, **
indicates p < 3.3e-03, * indicates p < 1.67e-02).

1064

Figure 2. Multiple timescales underlie the adaptation dynamics of cortical neurons in
response to repetitive sensory stimulation.
A. Grand-averaged PSTHs of S1 layer 4 RS neurons to the 10 Hz repetitive stimulus at the three
time points of interest: Control (1st stimulus in the repetitive train), 1.5 sec adapted (15th stimulus),
and 10 sec adapted (100th stimulus; note that only sensory-responsive neurons whose sensory
response significantly decreased between the Control and 10 sec adapted stimuli were included).
B. Same as A, but for the L4 FS population. **C.** Mean ± sem first-spike latency of the L4 RS and
FS populations to repetitive sensory stimuli at the three time points of interest (RS: Control: 12.21
± 0.15 msec, 1.5 sec adapted: 19.31 ± 0.19 msec, 10 sec adapted: 20.22. ± 0.20 msec; Control
versus 1.5 sec adapted: p = 6.40e-40, 1.5 sec adapted versus 10 sec adapted: p = 6.62e-07,
Control versus 10 sec adapted: p = 5.85e-40; FS: Control: 10.35 ± 0.13 msec, 1.5 sec adapted:
16.51 ± 0.21 msec, 10 sec adapted: 18.09 ± 0.25 msec; Control versus 1.5 sec adapted: p =
1.04e-31, 1.5 sec adapted versus 10 sec adapted: p = 5.31e-19, Control versus 10 sec adapted:
p = 6.26e-33; all statistical comparisons were performed using the Wilcoxon signed-rank test with
a Bonferroni correction; *** indicates p < 3.3e-04). **D.** Mean ± sem evoked firing rate to repetitive
sensory stimuli at the three time points of interest for the L4 RS and FS populations (RS: Control:
0.50 ± 0.02 spks/stim, 1.5 sec adapted: 0.22 ± 0.01 spks/stim, 10 sec adapted: 0.14 ± 0.01
spks/stim ; Control versus 1.5 sec adapted: p = 1.65e-35, 1.5 sec adapted versus 10 sec adapted:

p = 3.36e-31, Control versus 10 sec adapted: p = 7.77e-40; FS: Control: 0.72 ± 0.03 spks/stim,
1.5 sec adapted: 0.41 ± 0.0.03 spks/stim, 10 sec adapted: 0.30 ± 0.02 spks/stim ; Control versus
1.5 sec adapted: p = 7.56e-26, 1.5 sec adapted versus 10 sec adapted: p = 5.48e-28, Control
versus 10 sec adapted: p = 7.58e-31; all statistical comparisons were performed using the
Wilcoxon signed-rank test with a Bonferroni correction; *** indicates p < 3.3e-04). **E**. The
normalized mean evoked response for the L4 RS population over the timecourse of repetitive
stimulation overlaid with a single and a bi-exponential model fit (purple and blue dashed lines,
respectively; see Methods). The evoked response is normalized per neuron by the maximum
response to the first three stimuli before taking the average across all neurons. Inset: semi-log
plot of the normalized evoked response. The black lines denote a linear fit over two timescales,
suggesting bi-exponential behavior (τ_1 = 0.93 sec, τ_2 = 37.98 sec). **F**. Same as E, but for the L4
FS population (τ_1 = 0.92 sec, τ_2 = 40.53 sec).
1095
Figure 3. Multiple timescales underlie the adaptation dynamics of thalamic firing rate and
synchrony in response to repetitive sensory stimulation.
A. Grand-averaged PSTHs of VPm neurons to the 10 Hz repetitive stimulus at the three time
points of interest: Control (1[st] stimulus in the repetitive train), 1.5 sec adapted (15[th] stimulus), and
10 sec adapted (100[th] stimulus; note that only responsive neurons whose sensory response
significantly decreased between the Control and 10 sec adapted stimuli were included). **B**. Mean
± sem first-spike latency of the VPm population to repetitive sensory stimuli at the three time
points of interest (Control: 9.16 ± 0.15 ms, 1.5 sec adapted: 12.88 ± 0.20 ms, 10 sec adapted:
14.01 ± 0.21 ms; Control versus 1.5 sec adapted: p = 1.63e-31, 1.5 sec adapted versus 10 sec
adapted: p = 1.39e-17, Control versus 10 sec adapted: p = 1.41e-31; all statistical comparisons
were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates p
< 3.3e-04). **C**. Mean ± sem evoked firing rate of the VPm population to repetitive sensory stimuli
at the three time points of interest (Control: 0.81 ± 0.05 spks/stim, 1.5 sec adapted: 0.52 ± 0.03
spks/stim, 10 sec adapted: 0.38 ± 0.02 spks/stim. Control versus 1.5 sec adapted: p = 1.15e-13;
1.5 sec adapted versus 10 sec adapted: p = 1.44e-23, Control versus 10 sec adapted: p = 1.43e-
27, all statistical comparisons were performed using the Wilcoxon signed-rank test with a
Bonferroni correction; *** indicates p < 3.3e-04). **D**. The normalized mean evoked response for
the VPm population over the timecourse of repetitive stimulation overlaid with a single and a bi-
exponential model fit (purple and blue dashed lines, respectively). The evoked response is
normalized per neuron by the maximum response to the first three stimuli before taking the
average across all neurons. Inset: semi-log plot of the normalized evoked response. The black
lines denote a linear fit over two timescales, suggesting bi-exponential behavior (τ_1 = 0.79 sec, τ_2
= 29.65 sec). **E.** VPm population synchrony is measured by generating pairwise cross-
correlograms (CCGs) and quantifying the area in the +/-7.5 msec window normalized by the area
under the entire CCG (see Methods). **F.** Mean ± sem synchrony of the VPm population to
repetitive stimuli at the three time points of interest (Control: 0.89 ± 4.01e-03, 1.5 sec adapted:
0.81 ± 4.30e-03, 10 sec adapted: 0.76 ± 5.52e-03. Control versus 1.5 sec adapted: p = 6.90e-92;
1.5 sec adapted versus 10 sec adapted: p = 1.52e-20, Control versus 10 sec adapted: p = 3.90e-
107, all statistical comparisons were performed using the Wilcoxon signed-rank test with a
Bonferroni correction; *** indicates p < 3.3e-04). **G.** The normalized mean evoked synchrony for
the VPm population over the timecourse of repetitive stimulation overlaid with a single and a bi-
exponential model fit (purple and blue dashed lines, respectively). The synchrony is normalized
per pair of neurons by the maximum synchrony in response to the first three stimuli before taking
the average across all pairs of neurons. Inset: semi-log plot of the normalized population
synchrony. The black lines denote the linear fit over two timescales, suggesting bi-exponential
behavior (τ_1 = 0.66 sec, τ_2 = 185.55 sec).

1132

1133

Figure 4. Multiple timescales underlie the adaptation dynamics of thalamocortical synaptic efficacy in response to repetitive sensory stimulation

A. Left: Spike raster plots of a simultaneously recorded VPm, L4 RS, and L4 FS neuron in response to a weak sinusoidal whisker stimulus (4 Hz, mean velocity: 25-50 deg/s) in order to elevate spiking activity without driving correlated, stimulus-locked spiking across neurons. Right: The shuffle-corrected (SC) cross-correlograms (CCGs) for the L4 RS (top) and FS (bottom) neuron, each relative to the VPm neuron's spiking. The SC CCGs are generated by subtracting the shuffled CCGs from the raw CCGs. Raw CCGs are generated using the spikes elicited by the sinusoidal whisker stimulus and the shuffled CCGs are generated by randomizing the trials of the VPM and S1 spiking activity relative to each other. The gray lines denote 3.5 standard deviations of the shuffled distribution. A peak in the SC CCGs in the 1-3 ms window (window highlighted in light brown) indicates that these pairs of neurons are putative monosynaptically connected (Liew et al. 2021). **B**. Spike rasters of the putative monosynaptically connected neurons from A in response to 10 Hz repetitive sensory stimulus (300 deg/s). **C**. The grand-averaged Control (left), 1.5 sec adapted (middle), and 10 sec adapted (right) SC (black) and raw (purple) CCGs for all putative monosynaptically connected VPm-L4 RS neuron pairs. The 1-4 ms window used in subsequent analysis is highlighted in light brown. **D**. Same as C, but for all putative monosynaptically connected VPm-L4 FS pairs. **E**. Mean ± sem synaptic efficacy for all putative monosynaptically connected VPm-L4 RS pairs to repetitive sensory stimuli at the three time points of interest. Given the noisy nature of synaptic efficacy measurements, the control datapoint represents the average efficacy across first the stimuli, the 1.5-sec adapted datapoint represents the average efficacy across the 14^{th}-16^{th} stimuli (1.4-1.6 sec), and the 10-sec adapted timepoint represents the average efficacy across the 98^{th}-100^{th} stimuli (9.8-10 sec; Control: 2.1e-02 ± 1.7e-03, 1.5 sec adapted: 8.6e-03 ± 9.4e-04, 10 sec adapted: 6.4e-03 ± 8.1e-04. Control versus 1.5 sec adapted: p = 1.2e-11; 1.5 sec adapted versus 10 sec adapted: p = 1.2e-03, Control versus 10 sec adapted: p = 2.2e-15, all comparisons were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates p < 3.3e-04, ** indicates p < 3.3e-03). Synaptic efficacy is defined for each adaptation time point as the number of synchronous spikes within the 1-3 ms window divided by the number of VPm spikes contributing to the analysis (see Methods). **F**. Same as E, but for all putative monosynaptically connected VPm-L4 FS pairs (Control: 4.1e-02 ± 2.2e-03, 1.5 sec adapted: 3.2e-02 ± 1.9e-03, 10 sec adapted: 2.7e-02 ± 1.9e-03. Control versus 1.5 sec adapted: p = 4.3e-05; 1.5 sec adapted versus 10 sec adapted: p = 3.8e-02, Control versus 10 sec adapted: p = 1.6e-08, all comparisons were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates p < 3.3e-04). **G.** The normalized mean synaptic efficacy for all putative monosynaptically connected VPm-L4 RS pairs over the timecourse of repetitive sensory stimulation overlaid with a single and a bi-exponential model fit (blue and purple dashed lines, respectively). The efficacy is normalized per neuron by the maximum response to the first three stimuli before taking the average across all neurons. Inset: semi-log plot of the normalized population synaptic efficacy. The purple and blue lines denote the linear fit over two timescales, suggesting bi-exponential behavior (τ_1 = 0.22 sec, τ_2 = 34.1 sec). **H.** Same as G, but for all putative monosynaptically connected VPm-L4 FS pairs (τ_1 = 0.20 sec, τ_2 = 56.2 sec).

Figure 5. Direct optogenetic drive of VPm thalamus eliminates longer timescale adaptation in VPm

A. ChR2-expressing VPm neurons in lightly isoflurane-anesthetized NR133xAi32 mice were activated with 10 Hz repetitive LED pulses to selectively control properties of the VPm inputs while recording VPm and cortical L4 activity using 32-channel silicon probe. The histological section

shows expression of ChR2 tagged with eYFP (green) in the VPm and the co-localized electrode track fluorescently labeled with DiI (red) amongst DAPI labelled neurons (blue). **B.** The mean ± sem firing rate of the VPm population to control sensory and opto stimulation (Sensory: 0.6 ± 5.6e-2 spks/stim, Opto: 0.6 ± 0.8.2e-2 spks/stim, p = 0.37, the comparison was performed using the Wilcoxon signed-rank test). **C**. Grand-averaged PSTHs for VPm neurons to 10 Hz opto stimulation of VPm at the three time points of interest. **D.** Mean ± sem evoked response of the VPm population to repetitive optogenetic stimuli at the three time points of interest (Control: 0.62 ± 0.09 spks/stim, 1.5 sec adapted: 0.44 ± 0.06 spks/stim, 10 sec adapted: 0.45 ± 0.06 spks/stim. Control versus 1.5 sec adapted: p = 1.72e-04; 1.5 sec adapted versus 10 sec adapted: p = 7.17e-01, Control versus 10 sec adapted: p = 2.50e-04, all statistical comparisons were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates $p < 3.3e-04$). **E.** Population-averaged evoked response for the VPm population over the timecourse of repetitive optogenetic stimulation overlaid with a single exponential model fit. Inset: semi-log plot of the normalized evoked response; the black lines denote the linear fit over one timescale plus a constant, suggesting single exponential behavior (τ_1 = 0.36 sec). **F.** Mean ± sem synchrony of the VPm population to repetitive optogenetic stimuli at the three time points of interest (Control: 0.87 ± 1.80e-03, 1.5 sec adapted: 0.79 ± 1.88e-03, 10 sec adapted: 0.79 ± 1.76e-03. Control versus 1.5 sec adapted: p = 1.45e-07; 1.5 sec adapted versus 10 sec adapted: p = 3.48e-01, Control versus 10 sec adapted: p = 8.08e-08, all statistical comparisons were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates $p < 3.3e-04$). **G.** Population-averaged synchrony of the VPm population over the timecourse of repetitive optogenetic stimulation overlaid with a single exponential model fit. Inset: semi-log plot of the normalized evoked response; the black lines denote the linear fit over one timescale plus a constant, suggesting single exponential behavior (τ_1 = 0.82 sec).

Figure 6. Optogenetic drive of VPm results in a single timecourse of adaptation in S1
A. Grand-averaged PSTHs of L4 RS neurons in response to 10 Hz repetitive optogenetic stimulation of VPm at the three timepoints of interest. **B.** Same as A but for L4 FS neurons. **C.** Mean ± sem evoked response of the L4 RS population to repetitive optogenetic stimuli of VPm at the three time points of interest (Control: 1.07e-01 ± 9.30e-03, 1.5 sec adapted: 6.16e-02 ± 4.52e-03, 10 sec adapted: 5.59e-02 ± 3.49e-03. Control versus 1.5 sec adapted: p = 3.89e-13; 1.5 sec adapted versus 10 sec adapted: p = 2.63e-01, Control versus 10 sec adapted: p = 1.76e-15, all statistical comparisons were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates $p < 3.3e-04$). **D.** Same as C, but for the L4 FS population (Control: 2.09e-01 ± 2.33e-02, 1.5 sec adapted: 9.46e-02 ± 9.58e-03, 10 sec adapted: 8.87e-02 ± 8.84e-03. Control versus 1.5 sec adapted: p = 3.64e-09; 1.5 sec adapted versus 10 sec adapted: p = 3.99e-01, Control versus 10 sec adapted: p = 2.87e-11, all statistical comparisons were performed using the Wilcoxon signed-rank test with a Bonferroni correction; *** indicates $p < 3.3e-04$). **E.** Population-averaged evoked response for the L4 RS population over the timecourse of repetitive optogenetic stimulation of VPm overlaid with a single exponential model fit. Inset: semi-log plot of the normalized evoked response; the black lines denote the linear fit over one timescale plus a constant, suggesting single exponential behavior (τ_1 = 0.39 sec). **F.** Same as E, but for the L4 FS population (τ_1 = 0.47 sec).

Figure 7. Adaptation degrades stimulus detection for an ideal observer of cortical activity
A. Cortical responses to sensory inputs of 300 deg/s were probed either in the presence or absence of a preceding adapting stimulus. **B**. Parametric signal (stimulus present) and noise (stimulus absent) population spike probability distributions for L4 RS neurons when the probe stimulus is preceded by the Control (top), 1.5 sec adapted (middle), or 10 sec adapted (bottom)

stimulus. Parametric fits were generated using a Gamma distribution on the L4 RS population
response (see Methods). **C**. Same as in B, but for the L4 FS neuron population. **D**. Receiver
operator characteristic (ROC) curves of the sensory response of an example L4 RS neuron when
the stimulus is preceded by the Control, 1.5 sec adapting, or 10 sec adapting stimuli. **E**. Same as
D, but for the L4 FS population. **F**. Mean ± sem performance of an ideal observer of L4 RS activity
in detecting the signal versus noise in conditions preceded by Control, 1.5 sec adapting, and 10
sec adapting stimuli. The performance is quantified by the area under the receiver operator
characteristic curve (AUROC) performed across neurons; AUROC for control: $0.96 \pm 6.36\text{e-}03$,
1.5 sec adapted: $0.81 \pm 1.31\text{e-}02$, 10 sec adapted: $0.74 \pm 1.18\text{e-}02$; Control versus 1.5 sec
adapted: $p = 5.05\text{e-}23$, 1.5 sec versus 10 sec: $p = 5.31\text{e-}07$, Control versus 10 sec adapted: $p =$
$5.68\text{e-}29$; all statistical comparisons were performed using the Wilcoxon signed-rank test with a
Bonferroni correction; *** indicates $p < 3.3\text{e-}04$). **G**. Same as F but for L4 FS neurons (AUROC
for control: $0.96 \pm 8.04\text{e-}03$, 1.5 sec adapted: $0.88 \pm 1.45\text{e-}02$, 10 sec adapted: $0.81 \pm 1.63\text{e-}02$;
Control versus 1.5 sec adapted: $p = 4.76\text{e-}10$, 1.5 sec versus 10 sec: $p = 4.76\text{e-}09$, Control versus
10 sec adapted: $p = 1.37\text{e-}17$), all statistical comparisons were performed using the Wilcoxon
signed-rank test with a Bonferroni correction; *** indicates $p < 3.3\text{e-}04$).
1246
Figure 8. Adaptation enhances stimulus discrimination for an ideal observer of cortical
activity.
A. Stimulus response curves of the L4 RS population response to probe stimuli of varying
velocities preceded by Control, 1.5 sec adapting, or 10 sec adapting stimuli. **B.** Same as A, but
for the L4 FS population. **C.** Parametric distribution of L4 RS sensory responses to probe stimuli
of varying velocities when the probe stimulus is preceded by the Control (top), 1.5 sec adapting
(middle), or 10 sec adapting (bottom) stimuli. Parametric fits are generated using a Gamma
distribution of the RS population response (see Methods). **D**. Same as C but for the L4 FS
population. **E**. The performance of an ideal observer of L4 RS activity in discriminating between
sensory stimuli of varying velocities in conditions preceded by Control, 1.5 sec adapting, and 10
sec adapting stimuli (mean +/- 95% confidence intervals from re-sampling neurons with
replacement; see Methods). The performance is quantified by the accuracy of an LDA classifier
in discriminating between all velocities (Control: $0.33 \pm 3.1\text{e-}03$, 1.5 sec adapted: $0.35 \pm 4.7\text{e-}03$,
10 sec adapted: $0.36 \pm 4.3\text{e-}03$. Control versus 1.5 sec adapted: $p = 1.7\text{e-}02$, 1.5 sec versus 10
sec, $p = 1.7\text{e-}02$, Control versus 10 sec adapted: $p = 1.7\text{e-}02$; all statistical comparisons of
confidence intervals were performed with a Bonferroni correction). **F**. Same as E but for the FS
population (mean +/- 99.9% confidence intervals from re-sampling neurons with replacement;
Control: $0.34 \pm 5.7\text{e-}03$, 1.5 sec adapted: $0.41 \pm 7.3\text{e-}03$, 10 sec adapted: $0.43 \pm 6.9\text{e-}03$; Control
versus 1.5 sec adapted: $p = 3.3\text{e-}04$, 1.5 sec versus 10 sec, $p = 3.3\text{e-}04$, Control versus 10 sec
adapted: $p = 3.3\text{e-}04$).
1267

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8