Low Kit expression identifies Hematopoietic Stem Cell subsets with enhanced lymphoid potential in mice and humans

Harold K. Elias[1,2], Sneha Mitra[3], Marina B. da Silva[1,14], Adhithi Rajagopalan[1,14], Brianna Gipson[1], Nicole Lee[1], Anastasia I. Kousa[1], Mohamed A.E. Ali[4], Simon Grassman[1], Xiaoqun Zhang[1], Susan DeWolf [6], Melody Smith[5], Hana Andrlova[1], Kimon V. Argyropoulos[1], Roshan Sharma[7], Teng Fei[8], Joseph C Sun[1], Cynthia E. Dunbar[9], Christopher Y Park[4], Christina S. Leslie[3], Avinash Bhandoola[10], and Marcel R.M. van den Brink[1,11,12,13]

Affiliations
[1]Department of Immunology, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[2]National Institutes of Health (NIH), Bethesda, MD, 20892, USA, USA
[3]Computational and Systems Biology, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[4]Department of Pathology, New York University Grossman School of Medicine, New York, NY, 10016, USA
[5]Division of Blood and Marrow Transplantation and Cellular Therapy, Department of Medicine, Stanford University School of Medicine, Stanford, CA, 94305
[6]Leukemia Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[7]Single-cell Analytics and Innovation Lab, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[8]Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[9]Translational Stem Cell Biology Branch, National Heart, Lung, and Blood Institute, NIH, Bethesda, MD, 20892, USA
[10]Laboratory of Genome Integrity, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, MD, 20892, USA
[11]Department of Immunology and Microbial Pathogenesis, Weill Cornell Medical College; New York, NY
[12]Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA.

[13]Lead Contact
[14]These authors contributed equally
Correspondence: eliash@mskcc.org (HKE); vandenbm@mskcc.org (MRMvdB)

ABSTRACT

Hematopoietic stem cells (HSCs) with multilineage potential are critical for effective T cell reconstitution and restoration of the adaptive immune system after allogeneic Hematopoietic Cell Transplantation (allo-HCT). Our current understanding of lymphoid-poised HSC subsets is limited due to a lack of reliable markers. Through comprehensive molecular profiling of HSCs, we identify Kit^{lo} HSCs- a subset with enhanced lymphoid potential, which declines with age. Using a preclinical allo-HCT model, we demonstrate that Kit^{lo} HSCs exhibit superior lymphoid progenitor output, enhanced thymic recovery, and T-cell reconstitution resulting in improved T cell responses to infection post-HCT. Furthermore, Kit^{lo} HSCs with augmented BM lymphopoiesis mitigate age-associated thymic alterations, thus enhancing T-cell recovery in middle-aged hosts. Chromatin profiling revealed that Kit^{lo} HSCs exhibit higher activity of lymphoid-specifying transcription factors (TFs), including *Zbtb1*. Deletion of *Zbtb1* in Kit^{lo} HSCs, diminished their T-cell potential, while reinstating *Zbtb1* in megakaryocytic-biased Kit^{hi} HSCs rescued T-cell potential, *in vitro* and *in vivo*. Finally, we discover an analogous Kit^{lo} HSC subset with enhanced lymphoid potential in human bone marrow. Our results demonstrate that Kit^{lo} HSCs identify HSC subsets with enhanced lymphoid potential mediated by an underlying epigenetic program.

KEYWORDS: Hematopoietic stem cells; lymphoid differentiation; epigenetic priming; thymus; immune reconstitution; immunosenescence.

INTRODUCTION

T lymphocytes play a crucial role in the adaptive immune response. A highly diverse T cell pool is required to recognize and eliminate foreign pathogens while also maintaining self-tolerance. The thymus continuously produces T cells, but with advancing age, its regenerative capacity declines,[1] leading to decreased thymic output and compromised T cell diversity. Furthermore, the thymus is also vulnerable to acute damage from infections, cancer therapies, and conditioning regimens for allogeneic hematopoietic cell transplantation (allo-HCT), which results in prolonged T cell lymphopenia, thereby increasing the risk of infections and cancer relapse, contributing to transplant-related complications and mortality.[2] Likely for these reasons, early T-cell reconstitution is a positive prognostic indicator of allo-HCT outcomes.[3,4]

Post-transplant T cell reconstitution requires a steady supply of hematopoietic stem cells (HSCs)-derived lymphoid progenitors[5] as well as thymic recovery. Prior research has demonstrated that thymic integrity is dependent on thymocyte-stromal crosstalk, particularly in the thymic epithelial cell (TEC) compartment.[6-8] Age-associated reduced lymphoid potential originating at the level of HSCs,[9] coupled with thymic decline[1] contribute to delayed immune recovery. Therefore, identifying reliable markers to prospectively isolate HSC subsets that augment T lymphopoiesis and promote thymic regeneration is an unmet clinical need. Despite efforts to identify markers for HSCs with multilineage potential, the functional heterogeneity within phenotypic HSCs poses a challenge. More importantly, these markers are unknown within the context of the lymphoid-producing human hematopoietic stem cell hierarchy.

Single-cell transplantation with tracking and sequencing of progeny[10-15] have uncovered heterogeneity in self renewal capacity as well as lineage output among reconstituting HSCs-

highlighting HSCs with multilineage versus lineage-biased potential. These studies showed that,

while only a fraction of HSCs generate "balanced" multilineage output, the majority exhibit a

diverse range of lineage potential following transplantation- varying in their contributions and

reconstitution kinetics for each lineage, thereby showcasing their inherent biases. These lineage

biases are intrinsically stable and maintained following transplantation[16-18]. At the molecular

level, these cell autonomous features are mediated by epigenetic configuration and are divergent

from their transcriptional state.[13,19] Although this suggests the existence of a highly organized

and predictable framework for lineage-restricted fates of long-term self-renewing HSCs, the

molecular identities, gene regulatory networks and functional implications of these differences

governing lymphoid fate decisions remain largely unexplored.

Here, we detail the T-cell potential within the long-term reconstituting Kit^{lo} HSC subset, which

decreases with aging. We delineate the pivotal role of this population in orchestrating thymic

regeneration and immune reconstitution in both young and aged hosts. Leveraging integrated

single cell transcriptional and epigenetic profiling, we uncover the distinct epigenetic signature

of Kit^{lo} HSCs, highlighting the presence of lymphoid-specifying transcription factors like Zbtb1

and their pivotal role in enhancing T-cell lymphoid potential. Additionally, we identify an

analogous HSC subset in human bone marrow and demonstrate their enhanced lymphoid

potential.

METHODS

Detailed description can be found in Supplemental Information

Mice and Cell lines

Young (6-10 weeks), aged (18-24 months), and middle-aged (14-16 months) C57BL/6J, B6.SJL-

PtprcaPepcb/BoyJ,   BALB/cJ,   FVB-Tg   (RAG2-EGFP)1Mnz/J,   and   B6(C)-

Gt(ROSA)26Sor$^{em1.1(CAG-cas9*,-EGFP)Rsky/J}$ mice were acquired from Jackson Labs or National

Institute on Aging. S17, mouse and human MS5-DLL4, and MS5 stromal cells were used for *in*

vitro assays.

Peripheral Blood and Organ Analyses

120   Transplant chimerism was tracked in peripheral blood. Bone Marrrow, Thymi and spleens were

121   analyzed by flow cytometry and ELISA at the indicated time points.

Single Cell Multiome ATAC and Gene Expression Cell Preparation

124   Single Cell Multiome ATAC + Gene Expression was performed with the 10X genomics system

125   using Chromium Next GEM Single Cell Multiome Reagent Kit A (catalog no. 1000282) and

126   ATAC Kit A (catalog no. 1000280) following Chromium Next GEM Single Cell Multiome

127   ATAC + Gene Expression Reagent Kits User Guide and demonstrated protocol - Nuclei

128   Isolation for Single Cell Multiome ATAC + Gene Expression Sequencing.

Data and Code Availability

131   Multiome single cell RNA-and- ATAC sequencing data generated for this paper is accessible

132   through Gene Expression Omnibus (GEO) Series accession number GSE246464. No new

133   algorithms were developed for this paper. All analysis code and input for reproducing figures is

134   available at GitHub repository (https://github.com/HElias01/AgingHSC-Multiome.git).

RESULTS

Reduced Lymphoid Output in Aged HSCs Impairs Thymic Recovery

We utilized a pre-clinical allo-HCT model to analyze immune reconstitution of young and old donors. To evaluate if HSCs with differences in lymphoid progenitor production could impact thymic recovery, we transplanted Lin−Sca-1+Kit+ (LSK) CD34−CD150+CD48− cells LT-HSCs from young adult (2-mo) or old (24-mo) C57BL/6 mice with rescue BM cells (CD45.1) into lethally irradiated young BALB/cJ recipients (**Figure S1A**). Because HSC-derived thymic progeny emerge within 5-6 weeks,[20] we established an 8-week harvest timepoint. This interval allowed sufficient time for HSC-derived progenitors to repopulate the thymus, thereby providing a better understanding of thymic regeneration dynamics and the efficacy of HSC transplantation in promoting thymic reconstitution.

Consistent with prior reports, eight weeks post-HCT, we found that young HSCs supported balanced lineage reconstitution, while old HSCs exhibited significantly impaired T-cell reconstitution in the peripheral blood (PB), despite comparable reconstitution patterns at four weeks (**Figures S1B-S1E**). Furthermore, we observed greater multilineage potential of young HSCs, including lymphoid and myeloid reconstitution (LMPP/MPP4 and CLP, **Figures S1F and S1G**), when compared to old HSCs. Furthermore, young HSC recipients had significantly greater thymic cellularity and reconstitution of precursor and mature thymocytes than recipients of old HSCs (**Figures S1H-S1J**). In conclusion, old HSCs have significantly less multi-lineage potential, but especially T lymphoid potential than young HSCs in young allo-HCT recipients.

Aging alters Multilineage Potential of HSCs

Recent reports have shown remarkable functional heterogeneity of HSC[21-26] subsets, highlighting

discordance between current HSC phenotypic definitions to their molecular identities[11-13]. To

identify HSCs with an enriched multilineage program and gain insights into their transcriptional

and epigenetic framework, we performed multiome single-cell RNA and ATAC sequencing on

hematopoietic stem cells (LT-HSCs: Lineage$^-$CD34$^-$CD48$^-$CD150$^+$Sca-1$^+$c-Kit$^+$) from 2-mo

(young) or 24-mo (old) female mice.

To define HSC subsets, we performed unsupervised Leiden cluster analysis on the multiome

scRNA-seq data from our young cohort (**Figure S2A-S2B**). Using nomenclature and signatures

derived from previous studies [10,11,13,27-30], we mapped all unsupervised clusters into five main

subsets (**Figure S2C**): Quiescent HSC (q-HSC), Platelet-biased HSC (Mgk-HSC), Multilineage

HSC (MLin-HSC), Proliferative HSC (p-HSC), and a novel Intermediate HSC (Int-HSC) that

showed overlapping features with Mgk-HSC, MLin-HSC and p-HSC subsets (**Data Table 1,**

Figure S2D). To identify HSCs with multilineage potential, we compared multilineage enriched

MLin-HSC versus low output q-HSC subsets in young HSCs and queried for previously

described bonafide HSC markers (*Hlf, Kit, Neo1, Procr, CD244a, Pdzk1ip1, Ly9, Hoxb5*). *Hlf*

and *Kit* had significantly higher expression in q-HSCs compared to MLin-HSCs (**Figure 1A**).

Hlf has been previously implicated in HSC quiescence,[31-33] whereas *Kit* enriches for platelet-

biased HSCs.[34] To further analyze the composition of HSCs with varying Kit expression, we

defined Kit high and low HSC populations (Kit^{hi} and Kit^{lo} HSCs) based on the top and bottom

20% of *Kit* gene expression levels. We observed lower Kit expression in MLin-HSCs (**Figure**

1B) and increased representation of q-HSC and MLin-HSCs within the Kit^{lo} HSCs (**Figures 1C**

and S2E), consistent with prior reports of increased self-renewal capacity and multipotency in

Kit^{lo} HSCs[34,35].

To assess for age-associated changes, we first compared the transcriptional profile of young and old HSCs. In our differential gene analysis, we found that HSCs from young mice had higher expression of genes related to cell cycle (*Ccnd3*, *Cdc25a*, *Cdk1)*, and cell polarization (*Arhgap15*); conversely, old HSCs were enriched for expression of genes related to quiescence (*Egr1*, *Junb, Jun*, *Jund*, *Fos*), platelet differentiation (*vWf*, *Clu*, *Itgb3, Fhl1 and Tgm2*), and myeloid differentiation (*Selp, Neo1*) (**Data Table 2**). Next to investigate age-related shifts in HSC cluster composition, we employed *ingest()* from scanpy to query the old HSC transcriptomic data using the "trained" young reference data for HSC subsets (**Figure S2F**). Subsequently, we generated a UMAP of our young and old datasets (**Figures S2G and 1D**) to visualize the post-ingestion results (**Figures 1E**). While all HSC subsets were present in young and old cohorts (**Figure 1F**), we notably observed an increased representation of MLin-HSCs, p-HSCs, and Int-HSCs in young HSCs. Conversely q-HSCs and Mgk-HSCs were predominantly in old HSCs (**Figures S2H- S2J**). These findings suggest shifts in the composition of HSC subsets with aging, indicating a reduction in multilineage potential accompanied by an increase in platelet bias.

Given the increased frequency of Mgk-HSCs in old HSCs and prior work[34] demonstrating that Kit^{hi} HSCs identify platelet-biased HSCs (**Figures S2F- S2I**), we hypothesized this could be attributed to an expansion in Kit^{hi} HSCs. We observed significantly higher *Kit* expression (**Figures 1G-1H and S2K**) in old HSCs, with an age-dependent reduction in the proportion of Kit^{lo} HSCs accompanied by an increase in the frequency of Kit^{hi} HSCs (**Figure 1I**). Further substantiating our transcriptional findings with flow cytometry analysis of young and old bone

204 marrow, we noted a significant expansion of Kit[hi] HSCs, accompanied by a concomitant

205 decrease in Kit[lo] HSCs in old HSCs (**Figures S2L and 1J**), consistent with our prior report[36].

206 Together, these findings indicate that the frequency of multilineage Kit[lo] HSCs declines with age.

208 Young Kit[lo] HSCs Exhibit Enhanced *In Vitro* and *In Vivo* Lymphoid Potential

Next, we sought to determine the lymphoid potential of the HSC Kit subsets. To this end, we

used the S17 lymphoid progenitor assay[37] to assess for lymphoid progenitor output of purified

Kit[hi] and Kit[lo] HSCs from young mice (**Figure 2A**). Prior studies have elucidated the existence of

functional heterogeneity among phenotypic Common Lymphoid Progenitors (CLPs)

characterized by Ly6D expression,[38] resulting in the identification of two distinct subsets:

Absolute Lymphocyte Progenitors (ALPs), also referred to as functional CLPs, and B-cell biased

Lymphocyte Progenitors (BLPs). Following 12-day co-culture, we observed that young Kit[lo]

HSCs generated more CLPs and ALPs than Kit[hi] HSCs (**Figure 2B**). To further examine their T-

cell differentiation potential, we used murine artificial thymic organoids (mATO, BM-ATO)[39-41]

(**Figure 2A**). Following an 8-week culture, we noted a substantial increase in overall ATO

output from young Kitl[o] HSCs in comparison to Kit[hi] HSCs (**Figure 2C**).

To further evaluate the reconstituting potential of Kit[lo] HSCs, we competitively transplanted Kit[lo]

or Kit[hi] HSCs from young C57BL/6 (CD45.2) mice with rescue BM cells (CD45.1) into lethally

irradiated young BALB/cJ recipients (**Figure 2D**). Eight weeks after allo-HCT, we found that

Kit[lo] HSCs generated significantly better B- and T-cell reconstitution, while myeloid

reconstitution was comparable (**Figures 2E, S3A-S3B**). Concordant with a prior report of better

self-renewal ability[34], we found that Kit[lo] HSCs demonstrated increased LT HSC reconstitution

(**Figure 2F**). In support of multilineage potential, we observed that Kit[lo] HSC recipients had

higher lymphoid progenitor reconstitution (including LMPP and CLP, **Figures 2F and 2G**),

along with comparable myeloid precursor output (including MPP3, CMP, GMP, MEP, **Figures**

2F and S3C).

To evaluate whether enhanced lymphoid precursor output could contribute to thymic

reconstitution, we concurrently evaluated the recipient thymi. We found that Kit[lo] recipients

demonstrated enhanced thymic recovery, characterized by significantly higher thymic cellularity

and reconstitution of precursor and mature thymocytes (**Figures 2H-2J**). Notably, in support of a

critical role for thymocyte-stromal crosstalk in preserving thymic architecture[6-8], we also found

higher thymic epithelial cell (TEC), specifically medullary TEC, and endothelial cell recovery in

Kit[lo] reconstituted thymi (**Figures 2K and S3D**). We next examined donor-derived lymphoid

progenitors for differential expression of molecules, involved in homing to the thymus (CCR7,

CCR9, and PSGL1)[42].We observed significantly higher expression of CCR7 on Kit[lo]-derived

lymphoid progenitors (CLP) (**Figure S3E**). In addition, we analyzed thymic supernatants and

found no differences in levels of thymopoietic ligands, IL22[43] and RANKL[44] (**Figure S3F**).

Finally, we found significantly higher secondary lymphoid organ (spleen) reconstitution in Kit[lo]

recipients (**Figures S3G and S3H**). To evaluate long-term reconstitution, we performed

additional harvests at 20-weeks following HCT. We observed similar trends in higher BM

lymphopoiesis, thymic reconstitution, and thymic stromal recovery in Kit[lo] recipients (**Figure**

S4).

To assess thymic output, we used the RAG2-GFP[45] model, which allows the analysis of recent

thymic emigrants (RTEs) (**Figure 2L**). We observed higher numbers of PB GFP+ T cells in Kit[lo]

recipients **(Figure 2M)**. Additionally, to assess whether this increased number PB T cells in Kit[lo]

recipients results in increased functionality, (**Figure 2N**), we analyzed response to ovalbumin

engineered *Listeria Monocytogenes* (LM-OVA) and observed a significant increase in OT1+

CD8+ T cells (**Figure 2O**).

A key element in the post-HCT recovery of thymic function is the influx of bone marrow-

derived lymphoid progenitors, also known as thymic seeding progenitors (TSPs).[5] Furthermore,

the significance of restoring the bone marrow-thymus axis in mitigating thymic involution has

been underscored through various strategies aimed at augmenting BM-lymphopoiesis, including

sex-steroid ablation[46,47] and administration of Ghrelin,[48] among other approaches.[49] We therefore

hypothesized that robust BM lymphopoiesis driven by Kit^{lo} HSCs could potentially counteract

age-related alterations within the thymic microenvironment, consequently ameliorating the

decline in T-cell output. To this end, we competitively transplanted Kit^{lo} or Kit^{hi} HSCs from

young C57BL/6 mice (CD45.2) with rescue BM cells (CD45.1) into lethally irradiated 14-

month-old (Middle-aged) BALB/cJ recipients (**Figure 2P**). Sixteen weeks post-transplantation,

we observed in Kit^{lo} vs Kit^{hi} recipients significantly higher numbers of peripheral T– and B-cells

(**Figures 2Q, S3I, and S3J**). This was attributed to increased numbers of Kit^{lo} HSC-derived

lymphoid progenitors (including LMPP/MPP4 and CLPs, **Figures S3K and S3L**), while

numbers of myeloid precursors were comparable (MPP3, CMP, GMP, and MEP, **Figures S3K**

and S3M). In our parallel analysis of the thymi of Kit^{lo} vs Kit^{hi} recipients, we observed increased

numbers of thymocytes and TECs (**Figures 2R-2T and S3N**). Collectively, these findings

establish the potential of Kit^{lo} HSCs for lymphopoiesis and to improve thymic recovery and

peripheral T cell reconstitution in recipients of various ages.

Kit^{lo} HSCs are Epigenetically Programmed for Lymphoid Potential

HSCs can have a divergence between transcriptional and epigenetic lineage priming,[13,22,50] and

epigenetic priming predicting differential lymphoid output.[13,19] To identify the gene regulatory

network orchestrating the lymphoid potential differences in HSCs, we analyzed our single-cell

ATAC-seq (scATAC-seq) component from our multiome dataset (**Figure S5D**). Through

unsupervised clustering, we identified 6 clusters, including young and old HSCs (**Figures 3A**

and 3B). Distinct HSC subsets identified via scRNA-seq analyses (**Figure 1B**) are situated in

varied regions on the scATAC-seq UMAP (**Figures 3C and 3D**). Notably, MLin-HSC and Kit[lo]

HSC intersect with Cluster 3 from the scATAC-seq clusters (**Figure 3E**).

We characterized the differentially accessible regions (DARs) of chromatin that defined the HSC

subsets and found 2816 DARs in cluster C3 (MLin-HSC and Kit[lo] enriched) cluster compared to

cluster C6 (q-HSC and Kit[hi] enriched) (**Figure S5F**). Next to determine the differential activity

and accessibility of lymphoid-specifying transcription factors (TFs) in the identified Kit[hi] and

Kit[lo] HSC subsets, we analyzed chromatin accessibility variation within TF motifs using

chromVAR and gene accessibility. Compared to Kit[hi] HSCs, we found increased gene

accessibility and enrichment for Zinc finger motifs, specifically *Zbtb1 and Zbtb7a,* in Kit[lo]

DARs, which are implicated in lymphoid differentiation program[13,51-53] (**Figures 3E-3F**).

In summary, we found that Kit[lo] HSCs exhibit an epigenetic program, emphasizing an

enrichment of lymphoid-specifying transcription factors, in support of their heightened lymphoid

potential.

***Zbtb1* regulates T-cell potential in phenotypic HSCs**

Among members of the ZBTB gene family, *Zbtb7a* is a critical regulator of B versus T lineage

fate decision[52], while *Zbtb1* plays a pivotal role in T-cell development[53-55]. Furthermore, given

the role of *Zbtb1* in lymphoid reconstitution[53], we hypothesized that Zbtb1 loss impairs the

lymphoid potential of Kit[lo] HSCs. To this end, we used the CRISPR-Cas9 system to generate

Zbtb1-deficient Kit[lo] HSCs (**Figure 4A**) and confirmed decrease in Zbtb1 by flow-cytometric

analysis (**Figure S6A**). *Zbtb1*-deficient Kit[lo] HSCs showed significantly reduced *in vitro* T-cell

potential (**Figure 4B**) and impaired lymphoid reconstitution *in vivo* (**Figure 4C**). Next, we

questioned whether reinstating *Zbtb1* expression in megakaryocytic-biased, Kit[hi] HSCs is

sufficient to rescue their lymphoid defect. To test our hypothesis, we overexpressed *Zbtb1* cDNA

in Kit[hi] HSCs (**Figure 4D and S6B**) and we observed a rescue of their T-cell potential when

compared to Kit[hi] controls, thus partially phenocopying Kit[lo] HSCs (**Figures 4E**) as well as

improved lymphoid reconstitution *in vivo* (**Figure 4F**).

Previous studies have identified Zbtb1 targets which facilitate Notch-mediated T-cell

differentiation in lymphoid progenitors[56] and support chromatin remodeling following DNA

damage[57]. Given *Zbtb1* motif enrichment and enhanced T-cell differentiation in Kit[lo] HSCs, we

hypothesized that Zbtb1 targets mediating lymphoid differentiation gene programs are also

enriched in Kit[lo] HSCs. To this end, we queried Kit[lo] enriched genes for Notch1-interacting

proteins[58]. We observed ~50% overlap of Notch1-partners enriched in Kit[lo] HSCs, of which 60%

were Zbtb1-targets identified in a previous CHIP-seq study[56] (**Figure 4G**). Furthermore, we

identified an additional subset of Zbtb1 targets (33%) that were not associated with the Notch1

interactome (**Data Table 3**). To further query for non-Notch related biological pathways, we

performed pathway analysis and identified a strong enrichment for gene sets related to chromatin

remodeling, RNA stability, and (**Figure 4H, Data Table 4**).

These data demonstrate that Zbtb1 contributes to an epigenetic regulatory program that facilitates

T-cell differentiation in primitive Kit^{lo} HSCs likely through instructive and permissive

mechanisms.

Old Kit^{lo} HSCs Exhibit Enhanced *In Vitro* and *In Vivo* Lymphoid Potential

To determine whether Kit^{lo} HSCs retain their lymphoid potential with age, we evaluated the

lymphoid potential of old Kit subsets. Remarkably, even in the case of old mice, we noted a

substantial increase in lymphoid progenitor and ATO output from old Kit^{lo} HSCs compared to

Kit^{hi} HSCs (**Figures 5A-5B**), although modest when compared to the young HSCs. This finding

is consistent with prior reports of proliferative defects in aged T-cell precursors contributing to

decreased thymopoiesis.[59,60]

Next, to test their *in vivo* potential, we competitively transplanted either Kit^{lo} or Kit^{hi} HSCs from

old C57BL/6 (CD45.2) mice with rescue BM (CD45.1) cells into lethally irradiated young

BALB/cJ recipients (**Figure 5C**). At eight weeks post-transplantation, we noted significantly

higher numbers of T cells in the peripheral blood of Kit^{lo} vs Kit^{hi} recipients. (**Figures 5D and**

S7A). Consistent with our experiments using HSCs from young donors, we observed in the BM

of Kit^{lo} vs Kit^{hi} recipients significantly higher numbers of LT HSCs, LMPPs and CLPs (**Figures**

5E-5F) with comparable myeloid potential (**Figures 5E and S7B**). Furthermore, mirroring our

findings in HSCs from young mice, we observed in recipients of old Kit^{lo} HSCs enhanced

overall thymic cellularity, thymocyte reconstitution, and recovery of TECs and endothelial cells

(**Figures 5G-5I and S7C-S7D**), increased CD4+ and CD8+ T cells in the spleen (**Figures S7E**

and S7F), and higher RTE output (**Figures 5J-5K**). At 20-weeks post-HCT, we observed similar

trends of enhanced reconstitution in old Kit^{lo} HSC recipients, confirming their long-term

reconstituting potential (**Figure S7H-S7O**). Next to examine whether old HSC subsets with

preserved lymphoid could augment aged thymic recovery, we competitively transplanted middle-

aged recipients with old Kit^{lo} or Kit^{hi} donors (**Figure 5L**). We observed increased CLPs in the

BM and peripheral T cells, but a modest, non-significant increase in thymic cellularity in Kit^{lo}

recipients (**Figures 5M-5O, and S7G**). Similar to our *Zbtb1* functional studies with young Kit^{hi}

HSCs, we observed a rescue in T-cell potential in *Zbtb1*-overexpressed old Kit^{hi} HSCs (**Figure**

5P).

Finally, to directly compare the per cell functionality of Kit^{lo} HSCs in the context of aging, we

generated mixed chimeras combining Kit^{lo} HSCs cells from both young and old mice in an equal

competition transplant (**Figure S8A**). Consistent with our *in vitro* findings (**Figures 5A-5B**), we

found an overall decrease in lymphoid reconstitution within the PB, BM, and thymus with old

Kit^{lo} HSCs when compared to their young counterparts (**Figures S8B-S8G**). Pathway analysis

revealed decreased enrichment of chromatin remodeling related gene sets (**Figure S8H, Data**

Table 5). Additionally, we noted decreased accessibility and expression of *Zbtb1 (***Figures S8I-**

S8K*)* in Old Kit^{lo} HSCs, indicating a impaired epigenetic program for lymphoid differentiation.

These findings demonstrate that old Kit^{lo} HSCs exhibit preserved lymphoid potential and

contribute to thymic recovery, albeit with reduced output.

Kit^{lo} Human BM HSCs Subsets show Higher Lymphoid Potential

Informed by a recent study that mapped the molecular profiles of human thymic seeding

progenitors (TSPs) to their transcriptional counterparts within the HSC compartment,[61] we next

sought to identify a comparable human HSC subset with lymphoid potential. Interrogating a

CITE-Seq dataset[62] generated from young and old human bone marrow (BM) samples (**Figures**

6A-6C, S9A), we mapped our mouse Kit^{lo} gene signature onto human HSCs and noted a

significant enrichment in young BM (**Figures 6D and 6E**). We observed lower protein

expression of KIT (CD117) on young HSCs by using antibody-derived tag (ADT) reads (**Figure**

S9B). To validate our in-silico findings, we next assessed KIT expression in human BM samples

(young: 20-40 years; middle-age/ old: >40 years). In our FACS analysis, we found that

phenotypic HSCs (pHSCs: CD34+CD38-CD10-CD45RA-CD90+) showed the lowest KIT

expression (**Figure S9C**), with higher KIT levels on old pHSCs (**Figure 6F**). Moreover, we

noted a statistically significant association between age and the composition of KIT subsets in

pHSCs, indicating an increase in KIT^{hi} and a decrease in KIT^{lo} HSCs, aligning with our

observations in mice. (**Figure 6G**).

To evaluate the T-cell potential of these human HSC subsets, we generated artificial human

thymic organoids (huATOs) using purified KIT^{hi} or KIT^{lo} HSCs from human BM samples

(**Figures S9D and 6H**; **Data Table 6**). In concordance with our mouse *in vitro* studies,

following 8-10 weeks of culture, we observed significantly higher output of precursor and

mature T-cells from KIT^{lo} HSCs- derived ATOs compared to KIT^{hi} HSCs (**Figure 6I**). To further

evaluate the multilineage potential of KIT HSC subsets, we performed a lympho-myeloid

differentiation assay, as previously described[63]. While we observed comparable myeloid

differentiation potential, Kit^{lo} HSCs exhibited significantly more B cell and a non-significant

increase in NK cell differentiation potential (**Figure 6J**).

Collectively, these results identify a previously uncharacterized subset of human BM HSCs,

KIT^{lo} HSCs – which exhibit enhanced lymphoid potential.

<u>DISCUSSION</u>

Recent studies have uncovered significant heterogeneity within the HSC compartment and their

role in hematopoiesis. This heterogeneity has implications for disease development and

treatment. Of note, certain aspects of HSC heterogeneity challenge the established hematopoietic

models, with lineage biases and lineage-restricted cells impacting self-renewal dynamics.[64,65]

Decisions regarding lympho-myeloid fate are established at multiple levels of the hematopoietic

hierarchy: HSCs,[16,17,66] MPP, LMPP, and more mature GMPs and CLPs. At the level of HSCs,

the process of epigenetic imprinting confers a distinct advantage for lymphoid priming within

multilineage HSCs, as well as maintaining these cell autonomous features imposed at steady

state, even under conditions of stress.[13,19]

Here, we report on the enhanced lymphoid potential of the BM Kit HSC subset, Kit[lo] HSCs,

observed in both mice and humans. Through combining multiomic single-cell sequencing and

functional analyses utilizing a preclinical allo-HCT model, we demonstrate augmented potential

of Kit[lo] HSCs for lymphopoiesis and to support thymic recovery, independent of age. During the

early post-transplant period, we noted a relative increase in DN3 thymocytes relative to other

thymocyte precursor subsets. This observation is consistent with previous studies reporting that

DN3 thymocytes undergo compensatory expansion to support early post-transplant

thymopoiesis, with minimal contribution from T-cell precursor thymocytes (ETP and DN2)[5,67,68].

Further investigation into mechanisms driving these compensatory processes and its implications

for immune function is critical to our understanding of post-transplant immune reconstitution.

Mechanistically, Kit[lo] HSCs exhibit differential expression and activity of lymphoid-specifying

transcription factors (TFs), such as *Zbtb1*, indicative of their epigenetic program towards a

lymphoid fate. Studies in mouse models have shown that disruptions in *Zbtb1* result in T-cell

developmental defects and severe combined immunodeficiency phenotypes[53,54], along with

competitive disadvantages in B- and NK-cell compartments [53]. In our study, we identify *Zbtb1* as

an essential transcription factor governing lymphoid specification in phenotypic HSCs and

facilitating T-cell differentiation by reprogramming lineage biased Kit^{hi} HSCs. Additionally, we

observed functional differences within Kit^{lo} HSCs in the context of aging, specifically a

competitive disadvantage in Old Kit^{lo} HSCs, potentially attributable to their differences in *Zbtb1*

expression. Consistent with prior research, our study suggests *Zbtb1* modulates Notch-

dependent[56] and cellular survival programs[24,69]. However, further mechanistic studies are needed

to elucidate the precise transcriptional mechanisms underlying this regulation.

KIT has previously been reported as a reliable cross-species marker to fractionate phenotypic

HSCs in Cord Blood (CB)[70]. This study[70] showed reduced multipotency of KIT^{int} HSCs,

contradicting a prior report that showed the long-term reconstituting potential of CD34+KIT^{lo}

human BM cells[71]. In our study, combining transcriptional, immunophenotypic, and functional

analyses of human BM samples we identified a KIT^{lo} subset within HSCs with enhanced

lymphoid potential- demonstrating their translational relevance in human biology. In contrast to

the CB study[70], our observation of multipotency in KIT^{lo} HSCs within the BM suggests an

ontogeny specific relevance of Kit signaling. Although further functional characterization is

warranted to definitively establish their clinical significance.

Methods

Animal use

Young mice (6-10 weeks of age): C57BL/6J (CD45.2/H-2Kb), B6.SJL-PtprcaPepcb/BoyJ

(CD45.1), BALB/cJ (H-2Kd), and B6(C)-Gt(ROSA)26Sor[em1.1(CAG-cas9*,-EGFP)Rsky/J] (Rosa26[Cas9] KI

) mice were purchased from The Jackson Laboratory. Aged C57BL/6 (CD45.2/H-2Kb) mice (23-

24 months of age) were obtained from National Institute of Aging (Baltimore, MD). Aged mice

(ranging between 18-24 months of age) correlate with humans ranging between 56-69 years of

age. We obtained middle-aged mice (ranging between 14-16 months of age) by initially

purchasing young BALB/cJ (H-2Kd) mice from Jackson Laboratories (JAX) and subsequently

allowing them to age under controlled conditions in our facility. Middle-aged mice (ranging

between 14-16 months of age) correlate with humans ranging between 40-60 years of age.

RAG2-EGFP-CD45.1 chimeric mice were generated by crossing FVB-Tg (RAG2-EGFP)1Mnz/J

(JAX# 005688) and B6.SJL-PtprcaPepcb/BoyJ (CD45.1, JAX# 002014). Young OT-1-CD45.2

(Va2+Vb2+/ CD45.2/H2-Kb) were obtained from Dr. Andrea Schietinger. For consistency all

experiments were carried out using only female mice. These mice were allowed to acclimatize in

our vivarium for at least 10-14 days before experiments. Mice were euthanized with CO2 gas

inhalation and were maintained under pathogen-free conditions according to an MSKCC

IACUC-approved protocol.

Cell lines

To provide a microenvironment that supports lymphoid progenitor in vitro, we cocultured

purified HSCs from young and old B6 bone marrow with the S17 stromal cells[37], which were

originally obtained from K. Dorshkind (UCLA). To support in vitro T-cell differentiation, we co-

cultured

purified HSCs from young and old B6 bone marrow and human BM with MS5-mDLL4 or MS5-

hDLL4 stromal cells [39], which were obtained from Dr. Gay Crooks (UCLA). Subsequently, both

cell lines were maintained in our laboratory as described in the original references. Details of the

differentiation cultures are given below under HSC Lymphoid Differentiation Assays.

Cell Preparation and Staining

Femurs, tibias, pelvises, and the spine were dissected from euthanized mice and cleaned of

muscles and connective tissue on ice in 1X PBS. Bone marrow cells were collected by crushing

all the bones with a sterile mortar and pestle in FACS buffer: 1X PBS containing 2% Fetal

Bovine Serum (Hyclone) and 2 mM EDTA. BM cells were enriched for c-Kit+ cells using

mouse CD117 magnetic microbeads (Miltenyi Biotec) per manufacturer's protocol.

Bone Marrow analysis and Flow-cytometric Isolation of Hematopoietic Stem Cells

BM cells were stained with fluorochrome-conjugated antibodies as well as propidium iodide

(Molecular Probes) or Zombie Aqua™ Fixable Viability Kit (Biolegend). All monoclonal

antibodies were purchased from Biolegend and eBiosciences. The monoclonal antibodies

included Phycoerythrin-Cyanine 7 (PE-Cy7) antibodies to lineage (Lin) markers (CD3ε, CD4,

CD8a, B220, Gr-1, Mac-1 and Ter119), Brilliant Violet 711 (BV711) or Brilliant Violet 605 to

Sca-1, APC/Cyanine7 to cKit, FITC antibody to CD34, PE antibody to CD150 (SLAM), Pacific

Blue to CD48, APC to CD45.2 and PerCP/Cyanine5.5 to CD45.1. Cells were analyzed or sorted

using a FACS LSR II UV or Aria II cell sorter (BD Biosciences). For all experiments

(transplantation and in vitro assays) in which Kit^{hi}, Kit^{lo} or Kit HSCs were purified, they were

double sorted to ensure >95% purity.

To calculate bone marrow cellularity and frequency of donor-derived hematopoietic precursors,

two femurs and two tibias of primary recipient (BALB/cJ) were flushed into FACS buffer at 8-

or 20-weeks post-transplant. Collected cells were then incubated with red blood cell lysis buffer

(ACK lysis buffer, Thermo Fisher Scientific) for 8 minutes and then washed twice with

PBS/2.5% FBS. Cells were resuspended and then stained in PBS/2.5% fetal calf serum with

fluorochrome-conjugated antibodies (Biolegend, BD, eBiosciences) against lineage (Lin)

markers (CD3ε, CD4, CD8a, B220, Gr-1, Mac-1 and Ter119) (PE-Cy5), Sca-1 (BV711), c-Kit

(APC-Cyanine7), CD150/ SLAM (BV605), CD135/Flt3 (BV421 or PE-Cy5), CD127/IL7Ra

(APC), CD48 (BV510), CD16/32 (AF-700), CD34 (FITC), CD45.2 (BUV395), CD45.1 (PE-

Cy7 or AF-700), H-2Kd (PE) and H-2Kb (BV650 or BV786). For additional thymic adhesion

molecule analyses, antibodies against CCR7 (BV421), CCR9 (FITC) and PSGL1 (PE) were used

for chimerism studies. Following a 30-45 incubation on ice, cells were washed in in PBS/2.5%

fetal calf serum. Finally, propidium iodide (Molecular Probes) was added as a viability dye

before acquiring the data. Cells were analyzed or sorted using a FACS LSR II UV (BD

Biosciences).

Transplantation experiments

A. For competitive transplantation assays, double-FACS sorted Kit^{hi}, Kit^{lo} HSCs cells from

young or old CD45.2 mice were mixed with unfractionated BMMCs from young CD45.1 mice.

Cells were transplanted via retro-orbital sinus injections into lethally irradiated young (6-8-week-

old) or middle aged (14-16-month-old) BALB/cJ recipient mice (9 Gy, single dose, using a 79

X-Ray source) under isoflurane anesthesia, within 1h post-irradiation. **B. For Kit^{lo} equal**

competition experiment, 375 double-FACS sorted Kit^{lo} HSCs cells from both young

(CD45.1/CD45.2) and old CD45.2 mice were combined with $5x10^5$ unfractionated BMMCs from

young CD45.1 mice and transplanted into lethally irradiated 6-8-week-old BALB/cJ recipient

mice. **C. For RTE experiments**, 750 double-FACS sorted Kit^{hi}, Kit^{lo} HSCs cells from young

and old RAG2-GFP (CD45.1) mice were mixed with $5x10^5$ unfractionated BMMCs from young

CD45.2 mice and transplanted into lethally irradiated 6-8-week-old BALB/cJ recipient mice. **C.**

OT-1 Transplant and Adoptive T-cell transfer for functional assessment, 750 Kit^{hi} and Kit^{lo}

HSCs were double FACS-sorted from 8–10-week-old OT-1 mice and competitively transplanted

with $5x10^5$ unfractionated competitor BMMCs from (CD45.1) into lethally irradiated young (6-

8-week-old) BALB/cJ (H-2Kd) recipients. Eight-week post-transplant (BMT), spleens of

primary recipients were brought into single-cell suspension and 1/10 of splenocytes were

adoptively transferred into young (8-10wk) C57BL/6 secondary recipients. 4 hours later, mice

were infected with 5.000 CFU L. monocytogenes bacteria expressing chicken ovalbumin (LM-

OVA) to evaluate functional CD8+T cell response to primary infection. On day 8 post-infection,

spleens of infected mice were analyzed via flow cytometry. Counting beads were used to

calculate absolute cell numbers.

Peripheral blood analysis

Peripheral blood samples were collected in 50 mM EDTA solution (Thermo Fisher Scientific)

via retro-orbital sinus bleeds. Thereafter PB was incubated with red blood cell lysis buffer (ACK

lysis buffer, Thermo Fisher Scientific) for 8 minutes and then washed twice with PBS/2.5% fetal

bovine serum. Cells were resuspended and then stained in PBS/2.5% fetal calf serum with

antibodies against CD3 (AF700), B220 (PE-Cy7), Gr-1 (APC), Mac-1 (APC), NK1-1 (BV605),

CD45.2 (BUV395), CD45.1 (BV421), H-2Kd (PE) and H-2Kb (FITC) for chimerism studies for

30 mins followed by wash in in PBS/2.5% FBS. Finally, propidium iodide (Molecular Probes)

was added as a viability dye before acquiring the data. Cells were analyzed using a FACS LSR II

UV and LSR Fortessa X50 (BD Biosciences). To generate absolute T-cell counts, complete

blood counts, including differentials, were obtained using a ProCyte Dx Hematology Analyzer

(IDEXX).

Thymi Harvests: FACS and ELISA analysis

All steps were performed on ice unless indicated. Primary recipient (BALB/cJ) thymi at 8 or 20

wks post-transplant were excised and enzymatically digested following an adapted protocol.[43]

Briefly, thymi were mechanically dissociated into ca. 2 mm pieces. Tissue pieces were incubated

with a digestion buffer (RPMI, 10% FCS, 62.5 um/mL liberase TM, 0.4 mg/ml DNase I) twice

for 30 min at 37 C. Between incubation steps, supernatant containing dissociated cells was

transferred to 50 mL conical tubes equipped with 100 um filter.

For thymic immunophenotypic analysis, cells were first incubated with Fc block solution (anti-

CD16/CD32 antibody) for 10 min on ice. Solution was discarded and cells were stained in

PBS/2.5% fetal calf serum with fluorochrome-conjugated antibodies against lineage (Lin)

markers (CD19, CD11b, NK1-1, TCRγδ, Gr-1, Ter119) (Biotin), Streptavidin (PE-TexasRed),

CD4 (PE-Cy7), CD8 (BV711), CD25 (BV510), CD44 (AF700), c-Kit (APC), CD135/Flt3

(BV421), CD45.2 (BUV395), CD45.1 (BV605), H-2Kd (PE) and H-2Kb (FITC). for chimerism

studies. To characterize the CD45- compartment, thymic cells were stained with antibodies

against UEA-1 (FITC), 6C3 (PE), EPCAM (BV605), PDGFRa (BV421), MHC-II (APC), CD31

(PE-Cy7), CD45 (BUV395) and Ter-119 (PE-Cy5.5) for 30 mins followed by wash in in

PBS/2.5% FBS. Antibodies were purchased from either (Biolegend and eBiosciences), except

Ulex europaeus agglutinin 1 (UEA-1), conjugated to FITC, was purchased from Vector

Laboratories (Burlingame, CA). Finally, 7-AAD (Molecular Probes) was added as a viability dye

before acquiring the data. Flow Cytometric analysis was performed on FACS LSR II UV (BD

Biosciences).

For all ELISA experiments, thymi were suspended by mechanical dissociation. The resultant

supernatants were quantified using mouse cytokine (IL-22 and RANKL) specific ELISA kits

from R&D Systems and read on a Spark Multimode Microplate Reader (TECAN).

Spleen Harvest and Analysis

All steps were performed on ice unless indicated. Primary recipient (BALB/cJ) spleens at 8- or

20-weeks post-transplant were excised and crushed, filtered through a 70uM strainer

(pluriSelect), intermittently washed with PBS/2.5% fetal bovine serum and collected in a 50ml

conical tube. Subsequently, spleen pellet was incubated in red blood cell lysis buffer (ACK lysis

buffer, Thermo Fisher Scientific) for 8 minutes and then washed twice with PBS/2.5% fetal

bovine serum. Prior to staining, an aliquot was set aside for counts with the Nexcelom

Cellometer K2. For spleen immunophenotypic analysis, cells were first incubated with Fc block

solution (anti-CD16/CD32 antibody) for 10 min on ice. Solution was discarded and stained in

PBS/2.5% fetal calf serum with fluorochrome-conjugated antibodies against lineage (Lin)

markers (CD19, CD11b, NK1-1, TCRγδ, Gr-1, Ter119) (Biotin), Streptavidin (PE-TexasRed),

CD4 (PE-Cy7), CD8 (BV711), CD62L (BV510), CD44 (AF700), CD45.2 (BUV395), CD45.1

(BV605), H-2Kd (PE) and H-2Kb (FITC). for chimerism studies.

<u>**HSC Lymphoid Differentiation Assays**</u>

A. S17 Progenitor Assay

Lymphoid progenitor production from young and old B6 bone marrow was measured by mixing

FACS purified 200 Kit[hi] and Kit[lo] HSCs with 50,000 S17 stromal cells[37] in 1.5 mL of

methylcellulose (MC) medium. MC medium was prepared by supplementing a-MEM with 30%

heat-inactivated FCS, 1% methylcellulose (STEMCELL Technologies), 5x10-5 M 2ME, 2 mM

L-glutamine, 50 µg/mL gentamicin, 100 U/mL streptomycin, 100 µg/mL penicillin, 0.1 mM

MEM vitamins, 0.1 mM nonessential amino acids, 1 mM sodium pyruvate, 20 ng/mL stem cell

factor, 20 ng/mL Flt3L ligand, and 50 ng/mL IL-7 (all from Biosource). The mixture was plated

in non-tissue-culture treated 3.5-cm2 dishes (Becton Dickinson). Following 12 days of culture,

the contents of the plates were harvested, cells were enumerated, and examined for production of

lymphoid progenitors by flow cytometry. All cultures were placed at 37C, 5% CO_2 humidified

incubators until processing.

B. Mouse and Human ATO Assay

Prior to setting up the ATO experiments, the MS5 stromal cell lines were thawed and expanded

in culture for at least 7 days. The cell lines and ATOs were set up and maintained as initially

described.[39]

For the mouse ATO system, FACS purified 100 Kit[hi] and Kit[lo] HSC cells from young and old

mice were combined with 150k MS5-mDll4 stromal cells to form each murine ATO. The media

(DMEM/F-12, 50x B27, 100x Glutamax, 100x Pen/Strep, and 1000x Ascorbic acid) was

changed twice a week, with beta-mercaptoethanol (1000x) and cytokines (5ng/mL murine IL-7,

5ng/mL Flt3L, and 5ng/mL SCF) added fresh each time.

For the human ATO system, healthy human bone marrow mononuclear cells were purchased

from BIOIVT (Johnson City, TN) for FACS sorting of Kit[hi] and Kit[lo] HSCs within phenotypic

HSCs (CD34$^+$CD38$^-$CD10$^-$CD45RA$^-$CD90$^+$). The sorted cells were then aggregated with MS5-

hDll4 (50-150 HSCs and 150k MS5-hDll4 cells per ATO) and seeded on culture inserts

(Millipore Sigma). The human ATO media (RPMI 1640, 25x B27, 100x Glutamax, 100x

Pen/Strep, 1000x Ascorbic Acid) was refreshed twice a week, supplemented with 2.5ng/mL

recombinant human IL-7, 5ng/mL recombinant human Flt3L (Peprotech), and 5ng of human

SCF was used only for the first week of differentiation.

For analysis via FACS, the ATO was mechanically disrupted, filtered through a 70uM strainer

(pluriSelect), and counted with the Nexcelom Cellometer K2 prior to staining. Harvest

timepoints varied, dependent on the initial primary cell population (mATO: 6-8-weeks; huATO:

8-10-weeks).

C. Lympho-myeloid Differentiation Assay

As previously described[63], MS-5 cells were seeded onto 0.1% gelatin-coated 24-well plate at an

initiating density of $2\,\times\,10^4$/well in α-MEM medium (Thermo Fisher) supplemented with 10%

FBS (Gibco), 1% penicillin-streptomycin (Hyclone) and 1% GlutaMAXTM (Gibco). Twenty-

four hours after plating of MS-5 stroma, 200 cells were added into each well in the presence of

0.1 μM DuP-697 (Cayman Chemicals), 20 ng/ml SCF (Peprotech), 10 ng/ml G-CSF

(Peprotech), 10 ng/ml FLT3L (Peprotech), 10 ng/ml IL-2 (Peprotech), 10 ng/ml IL-15 (Sino

Biological). Cultures were maintained for 4 weeks with weekly half media changes (with 2x

cytokines). Cocultures were transferred onto fresh MS-5 stroma every two weeks through

40 μm filter to remove the stromal cells. All the cells in each well were harvested and analyzed

by flow cytometry at week 4. The antibodies used to read the lineage outputs in this assay were

anti-human CD45-BV480, anti-human CD14-BUV805, anti-human CD15-BV421, anti-human

CD19-FITC, anti-human CD56-BUV496, and anti-human CD235a-PE-Cy7.

Lentiviral Production

For deletion experiments, scramble and Zbtb1 guide RNAs (sgRNA#1:

GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAA

AGTGGCACCGAGTCGGTGC;   sgRNA#2:

GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAA

AGTGGCACCGAGTCGGTGC) were cloned into U6-EFS-mCherry (VectorBuilder), while

Control and mouse Zbtb1 [NM_178744.3] were individually cloned into CMV-EFS-EGFP

(VectorBuilder) for over-expression experiments and utilized for lentivirus generation.

Lentivirus was generated by cotransfecting the above plasmids containing either sgRNAs or

cDNA, pMD2.G (Addgene #12259), and psPAX2 (Addgene#12260) into HEK 293□T cells.

SFEM Media was replaced 6h post-transfection, and the virus suspension was harvested 36-48 h

post-media change. The supernatant was collected and used for lentiviral transductions.

HSC Lentiviral Transduction

As previously outlined sorted Kit^{lo} and Kit^{hi} HSCs from young, aged B6, and young Rosa26^{Cas9}

KI mice were cultured overnight in SFEM (STEMCELL Technologies) media supplemented

with murine cytokines (50□ng/ml SCF, 10□ng/ml TPO) in single-wells of a 96-well plate (TC

treated). The cells were transduced with lentiviral suspensions by spinfection in retronectin

(Takara Bio)-coated plates for 90 mins. Post spinfection, cytokine-supplemented fresh media

was added to the transduced wells. Twenty-four hours later, cells were either sorted for RFP and

GFP positivity (for deletion experiments), or GFP expression (for over-expression experiments)

using the BD Aria instrument. KO and OE was confirmed by flow cytometric analysis with

antibodies against Zbtb1 (AlexaFluor647, Signalway Antibody Cat#C47476-AF647).

Single Cell Multiome ATAC and Gene Expression Cell Preparation

Single Cell Multiome ATAC + Gene Expression was performed with the 10X genomics system

using Chromium Next GEM Single Cell Multiome Reagent Kit A (catalog no. 1000282) and

ATAC Kit A (catalog no. 1000280) following Chromium Next GEM Single Cell Multiome

ATAC + Gene Expression Reagent Kits User Guide and demonstrated protocol - Nuclei

Isolation for Single Cell Multiome ATAC + Gene Expression Sequencing. Briefly, cells

(viability 95%) were lysed for 4min and resuspended in Diluted Nuclei Buffer (10x Genomics,

PN- 2000207). Lysis efficiency and nuclei concentration was evaluated on Countess II automatic

cell counter by trypan blue staining. Nuclei were loaded per transposition reaction, with a

targeting recovery between 1,000 and 10,000 nuclei after encapsulation. After transposition

reaction nuclei were encapsulated and barcoded. Next-generation sequencing libraries were

constructed following User Guide, which were sequenced on an Illumina NovaSeq 6000 system.

SEQUENCING DATA PROCESSING

Single Cell RNA-seq

i) Preprocessing and downstream data analysis

FASTQ files were processed using the 10x Cell Ranger package (v7.01). The Cell Ranger

generated filtered_feature_bc_matrix.h5 files were processed following the guidelines on the

shunPykeR GitHub repository(https://github.com/kousaa/shunPykeR), an assembled pipeline of

publicly available single cell analysis packages put in coherent order, that allows for data

analysis in a reproducible manner and seamless usage of Python and R code. Genes that were not

expressed in any cell and ribosomal and hemoglobin genes were removed from downstream

analysis. Each cell was then normalized to a total library size of 10,000 reads and gene counts

were log-transformed using the log(X+1) formula, in which log denotes the natural logarithm.

Principal component analysis (components =20) was applied to reduce noise prior to data

clustering. To select the optimal number of principal components to retain for each dataset, the

knee point (eigenvalues smaller radius of curvature) was used. Leiden clustering[72] (resolution =

0.9) was used to identify clusters within the PCA-reduced data.

Quality of the single cells (**Fig. S2A-S2B**) was computationally assessed based on total counts,

number of genes, mitochondrial and ribosomal fraction per cell, with low total counts, low

number of genes ($\leq$1000) and high mitochondrial content ($\geq$0.2) as negative indicators of cell

quality. Cells characterized by more than one negative indicator were considered as "bad"

quality cells. Although cells were negatively sorted prior to sequencing for the CD45 marker, a

small amount of non-hematopoietic cells (expressing no Ptprc), were detected within our dataset.

To remove bad quality cells and contaminants in an unbiased way, we assessed them in a cluster

basis rather than individually. Leiden clusters with a "bad" quality profile and/or a high number

of contaminating cells were removed. Finally, cells marked as doublets by scrublet[73] were also

filtered out. Overall, a total of 2325 cells, representing ~8.4% of all our data, was excluded from

further analysis (see Figure S2 for per sample metrics). After removal of these cells, we

calculated highly variable genes (HVG=3000) and re-performed PCA with unsupervised

clustering analysis (components =20), followed by batch effect correction across all samples

using harmony[74] to assist annotation of cell type subsets within the dataset.

ii) Defining Kit^{lo} and Kit^{hi} HSC subsets

Owing to the significant drop-out effect in scRNA-seq data,[75] it is virtually impossible to

ascertain if a cell has low/mid (and sometimes high) Kit expression versus no expression. In

order to methodically deal with this, we performed _data denoising and_ imputation using the

MAGIC (Markov Affinity-based Graph Imputation of Cells) method,[76] to denoise distinguish

between dropouts and genuinely low gene expression values. Post-MAGIC imputation, we

proceeded to set the thresholds for *Kit*, specifically, we calculated the 20^{th} and 80^{th} percentile

expression value across all cells. Subsequently, cells were classified into Kit^{lo} (*Kit* expression

below the 20^{th} percentile), Kit^{hi} (*Kit* expression above the 80^{th} percentile), and Kit^{mid} (*Kit*

expression between the 20^{th} and 80^{th} percentile). This approach helped to enhance the quality of

the data by denoising and imputing the missing values, thereby providing a more accurate

representation of the gene expression landscape.

iii) Differential expression analysis

Differential expression analysis for comparisons of interest was performed with MAST (Model-

based Analysis of Single-cell Transcriptomics) using the likelihood ratio test.[77] In all cases,

differentially expressed genes were considered statistically significant if the FDR-adjusted p-

value was less than 0.05. To select maximally specific genes per cell type/subset, we ran MAST

for all pairwise cluster comparisons (each versus the rest) and retained the top 10 DEGs for a

given cluster (FDR<=0.05, sorted by decreasing coefficient). To generate the unique cluster

signatures per se, we used the *sc.tl.score_genes*() function from *scanpy* [78] that calculates

averaged scores based on the cluster specific genes (scores are subtracted with a randomly

sampled reference gene set).

iv) Differential abundance (DA) analysis

DA HSC subset across age groups was identified by sampling neighborhoods of cells from a k-

nearest neighbors (k-NN) graph and looking for enrichment of either age in each neighborhood

as implemented in MiloR.[79] MiloR is a graph-based statistical method to compute differential

cellular abundances in neighborhoods of cells. The 15 batch-corrected latent dimensions from

scanpy were used for MiloR (v.0.99.19) k-NN graph construction (k□=□35) and neighborhood

indexing (proportion□=□0.1). DA testing was performed with generalized linear models,

including age as covariates (neighborhoods significant if spatial corrected FDR□<□0.25).

Single- Cell ATAC sequencing

Preprocessing, dimensionality reduction, clustering

Single-cell ATAC-seq from young and old mice were aligned to the mm10 genome and we

processed the cellranger output file, fragments.tsv, with ArchR (v.1.0.2)[80] was used for

downstream analysis. We performed QC filtering on scATAC-seq using ArchR with the default

parameters of *createArrowFiles*() (**Fig. S5**). We retained cells containing at least 1,000 and at

most 100,000 fragments. Next, we filtered out the cells that did not pass QC in the corresponding

scRNA-seq data, thus retaining 11493 cells common to both scRNA-seq and scATAC-seq. We

then performed dimensionality reduction using iterative latent semantic indexing (LSI) on the top

25,000 variable features from the tile matrix to get a reduced dimensionality of 30 components

with *addIterativeLSI*() function in ArchR. To generate visualizations, we employed the

addUMAP() function in ArchR with the following settings: nNeighbors□=□20;

minDist□=□0.1. For clustering, we utilized the *addClusters*() function in ArchR, specifying the

parameters as follows: method $=$ 'Seurat'; knnAssign $=$ 10, and maxClusters $=$ 10. Next,

we used Harmony to perform batch effect correction.

Peak-calling and TF motif accessibility scoring

Each individual sample was pseudobulked for peak calling with MACS2[81] peak caller and

iterative peak overlapping removal within ArchR (using default settings). We subsequently

added motif annotations using *addMotifAnnotations*() with the CisBP motif database and

computed chromVAR[82] deviations for each single cell with *addDeviationsMatrix*(). To identify

differentially   accessible   motifs   within   each   group   of   interest,   we   applied   the

rank_genes_groups() function in *scanpy*. With the following settings: method $=$ 'wilcoxon'

and corr_method = 'benjamini-hochberg' and performed the analysis on the chromVAR zscore

matrix.

Processing Human Single-Cell CITE-seq

We analyzed the young human bone marrow CITE-seq from Sommarin et al.[62] using Seurat.[83]

Authors provided the peak matrix aligned to the hg38 genome. We used the sample BM_34 for

which the authors provided the cell type annotations. We processed the scRNA-seq and CITE-

seq data for young and old bone marrow using Seurat. The hash-tagged cells were demultiplexed

using *HTODemux*() from Seurat. The cell type annotations were provided by the authors for

young bone marrow samples labeled yBM1_hpc and yBM2_hpc. We predicted cell type

annotations for the old bone marrow cells by label transfer using *FindTransferAnchors*() and

TransferData() functions. We obtained the Kit^{lo} gene signature from our mouse scRNAseq data

through differential gene expression analysis. To extend this signature to our human bone

marrow HSPC dataset, we used Ensembl[84] Biomart to identify human orthologues of the mouse

genes. Subsequently, we scored the Kit^{lo} gene signature in human scBM HSCs using Seurat's

(AddModuleScore) function.

Data and Code Availability

Multiome single cell RNA-and- ATAC sequencing data generated for this paper is available

upon request from the corresponding author.

Statistical Analysis

Data were processed in GraphPad Prism 10.0 software. Statistical comparisons between 2 groups

were performed with either the nonparametric unpaired Mann-Whitney U test, two-way

ANOVA, ComBat batch effect correction[85] or Wilcoxon rank sum test with continuity correction

as indicated.

ACKNOWLEDGEMENTS

We thank Dr. Andrea Schietinger for the gift of mice; Ronan Chaligné, Kalina Belcheva, and

Anthony Michaels for help; Dr. Gay Crooks, and Julia Gerscheimer (UCLA) for sharing the

mATO and huATO systems and assistance with troubleshooting; Kenneth Dorshkind (UCLA)

for providing S17 cells; Parashar Dhopala and Göran Karlsson (Lund University) for providing

the human BM CITE-seq raw matrices.[62]

FUNDING SOURCES

This research was supported by National Cancer Institute award numbers, R35-CA284024, P01-

CA023766, R01-CA228308, and P30 CA008748 MSK Cancer Center Support Grant/Core

Grant; National Heart, Lung, and Blood Institute (NHLBI) award number R01-HL164902;

National Institute on Aging award number P01-AG052359, and Tri Institutional Stem Cell

Initiative. Additional funding was received from The Lymphoma Foundation, The Susan and

Peter Solomon Family Fund, The Solomon Microbiome Nutrition and Cancer Program, Cycle

for Survival, Parker Institute for Cancer Immunotherapy, Paula and Rodger Riney Multiple

Myeloma Research Initiative, Starr Cancer Consortium, and Seres Therapeutics. HKE receives

research support from NHLBI. SM receives research support from Marie-Josée Kravis

fellowship in quantitative biology. SG receives research support from CRI / Donald J. Gogel

Postdoctoral Fellowship (CRI #3934). SDW reports research funding from the MSK Leukemia

SPORE Career Enhancement Program (NIH/NCI P50 CA254838-01), the MSK Gerstner

Physician Scholar Program, and the Parker Institute for Cancer Immunotherapy. MS receives

research support from the following funding sources: Burroughs Wellcome Fund (PDEP),

Damon Runyon Cancer Research Foundation (Clinical Investigator Award), V Foundation (V

Scholar), Emerson Collective, and the National Institutes of Health (NHLBI K08 HL156082-

01A1). KA receives research support from the Gerstner Physician Scholar program.

AUTHOR CONTRIBUTIONS

HKE, MRMvdB conceived and designed the study. HKE, MRMvdB acquired funding. HKE,

MBdS, AR, BG, NL, MAEA, SG planned research activities. HKE curated data and performed

initial analysis. AIK and SM performed initial computational analysis of single-cell RNA- seq

data, and single-cell ATAC- seq and human CITE-seq data, respectively. RS provided code and

assisted HKE with downstream data/sequence computation and statistical analyses. HKE

analyzed flow cytometric data. HKE, MBdS, AR, BG, NL, MAEA, XZ, HA, SG, MS, SDW,

KV, CYP, CED, and JCS planned and performed experiments. CSL, AB, MRMvdB supervised

study. HKE, SM, and MBdS prepared figures and visualized data. HKE, MRMvdB wrote the

initial draft. All authors reviewed and approved the manuscript.

DECLARATION OF INTERESTS

<u>**REFERENCES**</u>

1.   Hale JS, Boursalian TE, Turk GL, Fink PJ. Thymic output in aged mice. *Proc Natl Acad*
Sci U S A. 2006;103(22):8447-8452.
2.   van den Brink M, Uhrberg M, Jahn L, DiPersio JF, Pulsipher MA. Selected biological
issues affecting relapse after stem cell transplantation: role of T-cell impairment, NK cells and
intrinsic tumor resistance. *Bone Marrow Transplant*. 2018;53(8):949-959.
3.   de Koning C, Prockop S, van Roessel I, et al. CD4+ T-cell reconstitution predicts
survival outcomes after acute graft-versus-host-disease: a dual-center validation. *Blood*.
2021;137(6):848-855.

4. van Roessel I, Prockop S, Klein E, et al. Early CD4+ T cell reconstitution as predictor of
outcomes after allogeneic hematopoietic cell transplantation. *Cytotherapy*. 2020;22(9):503-510.
5. Zlotoff DA, Zhang SL, De Obaldia ME, et al. Delivery of progenitors to the thymus
limits T-lineage reconstitution after bone marrow transplantation. *Blood*. 2011;118(7):1962-
1970.
6. van Ewijk W, Shores EW, Singer A. Crosstalk in the mouse thymus. *Immunol Today*.
1994;15(5):214-217.
7. Poliani PL, Facchetti F, Ravanini M, et al. Early defects in human T-cell development
severely affect distribution and maturation of thymic stromal cells: possible implications for the
pathophysiology of Omenn syndrome. *Blood*. 2009;114(1):105-108.
8. Pala F, Oguz C, Corsino C, et al. Profound abnormalities in thymic epithelial cells in
Rag1 hypomorphic mice: implications for immune reconstitution after stem cell transplantation.
The Journal of Immunology. 2022;208(1_Supplement):167.102-167.102.
9. Elias HK, Bryder D, Park CY. Molecular mechanisms underlying lineage bias in aging
hematopoiesis. *Semin Hematol*. 2017;54(1):4-11.
10. Rodriguez-Fraticelli AE, Weinreb C, Wang SW, et al. Single-cell lineage tracing unveils
a role for TCF15 in haematopoiesis. *Nature*. 2020;583(7817):585-589.
11. Dong F, Hao S, Zhang S, et al. Differentiation of transplanted haematopoietic stem cells
tracked by single-cell transcriptomic analysis. *Nat Cell Biol*. 2020;22(6):630-639.
12. Carrelha J, Meng Y, Kettyle LM, et al. Hierarchically related lineage-restricted fates of
multipotent haematopoietic stem cells. *Nature*. 2018;554(7690):106-111.
13. Meng Y, Carrelha J, Drissen R, et al. Epigenetic programming defines haematopoietic
stem cell fate restriction. *Nat Cell Biol*. 2023;25(6):812-822.
14. Morita Y, Ema H, Nakauchi H. Heterogeneity and hierarchy within the most primitive
hematopoietic stem cell compartment. *J Exp Med*. 2010;207(6):1173-1182.
15. Yamamoto R, Wilkinson AC, Ooehara J, et al. Large-Scale Clonal Analysis Resolves
Aging of the Mouse Hematopoietic Stem Cell Compartment. *Cell Stem Cell*. 2018;22(4):600-607
e604.
16. Benz C, Copley MR, Kent DG, et al. Hematopoietic stem cell subtypes expand
differentially during development and display distinct lymphopoietic programs. *Cell Stem Cell*.
2012;10(3):273-283.
17. Challen GA, Boles NC, Chambers SM, Goodell MA. Distinct hematopoietic stem cell
subtypes are differentially regulated by TGF-beta1. *Cell Stem Cell*. 2010;6(3):265-278.
18. Naik SH, Perie L, Swart E, et al. Diverse and heritable lineage imprinting of early
haematopoietic progenitors. *Nature*. 2013;496(7444):229-232.
19. Yu VWC, Yusuf RZ, Oki T, et al. Epigenetic Memory Underlies Cell-Autonomous
Heterogeneous Behavior of Hematopoietic Stem Cells. *Cell*. 2016;167(5):1310-1322 e1317.
20. Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid
progenitors in mouse bone marrow. *Cell*. 1997;91(5):661-672.
21. Flohr Svendsen A, Yang D, Kim K, et al. A comprehensive transcriptome signature of
murine hematopoietic stem cell aging. *Blood*. 2021;138(6):439-451.
22. Grover A, Sanjuan-Pla A, Thongjuea S, et al. Single-cell RNA sequencing reveals
molecular and functional platelet bias of aged haematopoietic stem cells. *Nat Commun*.
2016;7:11075.

23. Herault L, Poplineau M, Mazuel A, Platet N, Remy E, Duprez E. Single-cell RNA-seq
reveals a concomitant delay in differentiation and cell cycle of aged hematopoietic stem cells.
BMC Biol. 2021;19(1):19.
24. Kim YW, Zara G, Kang H, et al. Integration of single-cell transcriptomes and biological
function reveals distinct behavioral patterns in bone marrow endothelium. *Nat Commun*.
2022;13(1):7235.
25. Konturek-Ciesla A, Dhapola P, Zhang Q, et al. Temporal multimodal single-cell profiling
of native hematopoiesis illuminates altered differentiation trajectories with age. *Cell Rep*.
2023;42(4):112304.
26. Safi F, Dhapola P, Warsi S, et al. Concurrent stem- and lineage-affiliated chromatin
programs precede hematopoietic lineage restriction. *Cell Rep*. 2022;39(6):110798.
27. Giladi A, Paul F, Herzog Y, et al. Single-cell characterization of haematopoietic
progenitors and their trajectories in homeostasis and perturbed haematopoiesis. *Nat Cell Biol*.
2018;20(7):836-846.
28. Cabezas-Wallscheid N, Buettner F, Sommerkamp P, et al. Vitamin A-Retinoic Acid
Signaling Regulates Hematopoietic Stem Cell Dormancy. *Cell*. 2017;169(5):807-823 e819.
29. Lauridsen FKB, Jensen TL, Rapin N, et al. Differences in Cell Cycle Status Underlie
Transcriptional Heterogeneity in the HSC Compartment. *Cell Rep*. 2018;24(3):766-780.
30. Cabezas-Wallscheid N, Klimmeck D, Hansson J, et al. Identification of regulatory
networks in HSCs and their immediate progeny via integrated proteome, transcriptome, and
DNA methylome analysis. *Cell Stem Cell*. 2014;15(4):507-522.
31. Wahlestedt M, Ladopoulos V, Hidalgo I, et al. Critical Modulation of Hematopoietic
Lineage Fate by Hepatic Leukemia Factor. *Cell Rep*. 2017;21(8):2251-2263.
32. Komorowska K, Doyle A, Wahlestedt M, et al. Hepatic Leukemia Factor Maintains
Quiescence of Hematopoietic Stem Cells and Protects the Stem Cell Pool during Regeneration.
Cell Rep. 2017;21(12):3514-3523.
33. Lehnertz B, Chagraoui J, MacRae T, et al. HLF expression defines the human
hematopoietic stem cell state. *Blood*. 2021;138(25):2642-2654.
34. Shin JY, Hu W, Naramura M, Park CY. High c-Kit expression identifies hematopoietic
stem cells with impaired self-renewal and megakaryocytic bias. *J Exp Med*. 2014;211(2):217-
231.
35. Grinenko T, Arndt K, Portz M, et al. Clonal expansion capacity defines two consecutive
developmental stages of long-term hematopoietic stem cells. *J Exp Med*. 2014;211(2):209-215.
36. Elias HK, Shin JY, Ali MA, et al. Identification of Old HSCs with Preserved Self-
Renewal and Long-Term Reconstitution Potential. *Blood*. 2019;134:2479.
37. Montecino-Rodriguez E, Kong Y, Casero D, Rouault A, Dorshkind K, Pioli PD.
Lymphoid-Biased Hematopoietic Stem Cells Are Maintained with Age and Efficiently Generate
Lymphoid Progeny. *Stem Cell Reports*. 2019;12(3):584-596.
38. Inlay MA, Bhattacharya D, Sahoo D, et al. Ly6d marks the earliest stage of B-cell
specification and identifies the branchpoint between B-cell and T-cell development. *Genes Dev*.
2009;23(20):2376-2381.
39. Seet CS, He C, Bethune MT, et al. Generation of mature T cells from human
hematopoietic stem and progenitor cells in artificial thymic organoids. *Nat Methods*.
2017;14(5):521-530.
40. Montel-Hagen A, Sun V, Casero D, et al. In Vitro Recapitulation of Murine
Thymopoiesis from Single Hematopoietic Stem Cells. *Cell Rep*. 2020;33(4):108320.

41. Zhou W, Gao F, Romero-Wolf M, Jo S, Rothenberg EV. Single-cell deletion analyses show control of pro-T cell developmental speed and pathways by Tcf7, Spi1, Gata3, Bcl11a, Erg, and Bcl11b. *Sci Immunol*. 2022;7(71):eabm1920.

42. Zlotoff DA, Sambandam A, Logan TD, Bell JJ, Schwarz BA, Bhandoola A. CCR7 and CCR9 together recruit hematopoietic progenitors to the adult thymus. *Blood*. 2010;115(10):1897-1905.

43. Dudakov JA, Hanash AM, Jenq RR, et al. Interleukin-22 drives endogenous thymic regeneration in mice. *Science*. 2012;336(6077):91-95.

44. Lopes N, Vachon H, Marie J, Irla M. Administration of RANKL boosts thymic regeneration upon bone marrow transplantation. *EMBO Mol Med*. 2017;9(6):835-851.

45. Yu W, Nagaoka H, Jankovic M, et al. Continued RAG expression in late stages of B cell development and no apparent re-induction after immunization. *Nature*. 1999;400(6745):682-687.

46. Goldberg GL, Sutherland JS, Hammet MV, et al. Sex steroid ablation enhances lymphoid recovery following autologous hematopoietic stem cell transplantation. *Transplantation*. 2005;80(11):1604-1613.

47. Goldberg GL, Alpdogan O, Muriglan SJ, et al. Enhanced immune reconstitution by sex steroid ablation following allogeneic hemopoietic stem cell transplantation. *J Immunol*. 2007;178(11):7473-7484.

48. Dixit VD, Yang H, Sun Y, et al. Ghrelin promotes thymopoiesis during aging. *J Clin Invest*. 2007;117(10):2778-2790.

49. Spadaro O, Youm Y, Shchukina I, et al. Caloric restriction in humans reveals immunometabolic regulators of health span. *Science*. 2022;375(6581):671-677.

50. Mansson R, Hultquist A, Luc S, et al. Molecular evidence for hierarchical transcriptional lineage priming in fetal and adult stem cells and multipotent progenitors. *Immunity*. 2007;26(4):407-419.

51. Shin B, Hosokawa H, Romero-Wolf M, et al. Runx1 and Runx3 drive progenitor to T-lineage transcriptome conversion in mouse T cell commitment via dynamic genomic site switching. *Proc Natl Acad Sci U S A*. 2021;118(4).

52. Maeda T, Merghoub T, Hobbs RM, et al. Regulation of B versus T lymphoid lineage fate decision by the proto-oncogene LRF. *Science*. 2007;316(5826):860-866.

53. Siggs OM, Li X, Xia Y, Beutler B. ZBTB1 is a determinant of lymphoid development. *J Exp Med*. 2012;209(1):19-27.

54. Punwani D, Simon K, Choi Y, et al. Transcription factor zinc finger and BTB domain 1 is essential for lymphocyte development. *J Immunol*. 2012;189(3):1253-1264.

55. Zhang X, Lu Y, Cao X, Zhen T, Kovalovsky D. Zbtb1 prevents default myeloid differentiation of lymphoid-primed multipotent progenitors. *Oncotarget*. 2016;7(37):58768-58778.

56. Koizumi M, Kama Y, Hirano KI, et al. Transcription factor Zbtb1 interacts with bridging factor Lmo2 and maintains the T-lineage differentiation capacity of lymphoid progenitor cells. *J Biol Chem*. 2022;298(11):102506.

57. Kim H, Dejsuphong D, Adelmant G, et al. Transcriptional repressor ZBTB1 promotes chromatin remodeling and translesion DNA synthesis. *Mol Cell*. 2014;54(1):107-118.

58. Yatim A, Benne C, Sobhian B, et al. NOTCH1 nuclear interactome reveals key regulators of its transcriptional activity and oncogenic function. *Mol Cell*. 2012;48(3):445-458.

59. Min H, Montecino-Rodriguez E, Dorshkind K. Reduction in the developmental potential of intrathymic T cell progenitors with age. *J Immunol*. 2004;173(1):245-250.

60. Srinivasan J, Vasudev A, Shasha C, et al. The initial age-associated decline in early T-
cell progenitors reflects fewer pre-thymic progenitors and altered signals in the bone marrow and
thymus microenvironments. *Aging Cell*. 2023.
61. Cordes M, Cante-Barrett K, van den Akker EB, et al. Single-cell immune profiling
reveals thymus-seeding populations, T cell commitment, and multilineage development in the
human thymus. *Sci Immunol*. 2022;7(77):eade0182.
62. Sommarin MNE, Dhapola P, Safi F, et al. Single-Cell Multiomics Reveals Distinct Cell
States at the Top of Human Hematopoietic Hierarchy. *bioRxiv*. 2021:2021.2004.2001.437998.
63. Karamitros D, Stoilova B, Aboukhalil Z, et al. Single-cell analysis reveals the continuum
of human lympho-myeloid progenitor cells. *Nat Immunol*. 2018;19(1):85-97.
64. Notta F, Zandi S, Takayama N, et al. Distinct routes of lineage development reshape the
human blood hierarchy across ontogeny. *Science*. 2016;351(6269):aab2116.
65. Velten L, Haas SF, Raffel S, et al. Human haematopoietic stem cell lineage commitment
is a continuous process. *Nat Cell Biol*. 2017;19(4):271-281.
66. Dykstra B, Kent D, Bowie M, et al. Long-term propagation of distinct hematopoietic
differentiation programs in vivo. *Cell Stem Cell*. 2007;1(2):218-229.
67. Paiva RA, Sousa AGG, Ramos CV, et al. Self-renewal of double-negative 3 early
thymocytes enables thymus autonomy but compromises the beta-selection checkpoint. *Cell Rep*.
2021;35(2):108967.
68. Maillard I, Schwarz BA, Sambandam A, et al. Notch-dependent T-lineage commitment
occurs at extrathymic sites following bone marrow transplantation. *Blood*. 2006;107(9):3511-
3519.
69. Cao X, Lu Y, Zhang X, Kovalovsky D. Zbtb1 Safeguards Genome Integrity and Prevents
p53-Mediated Apoptosis in Proliferating Lymphoid Progenitors. *J Immunol*. 2016;197(4):1199-
1211.
70. Cosgun KN, Rahmig S, Mende N, et al. Kit regulates HSC engraftment across the
human-mouse species barrier. *Cell Stem Cell*. 2014;15(2):227-238.
71. Kawashima I, Zanjani ED, Almaida-Porada G, Flake AW, Zeng H, Ogawa M. CD34+
human marrow cells that express low levels of Kit protein are enriched for long-term marrow-
engrafting cells. *Blood*. 1996;87(10):4136-4142.
72. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-
connected communities. *Sci Rep*. 2019;9(1):5233.
73. Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets
in Single-Cell Transcriptomic Data. *Cell Syst*. 2019;8(4):281-291 e289.
74. Korsunsky I, Millard N, Fan J, et al. Fast, sensitive and accurate integration of single-cell
data with Harmony. *Nat Methods*. 2019;16(12):1289-1296.
75. Kharchenko PV, Silberstein L, Scadden DT. Bayesian approach to single-cell differential
expression analysis. *Nature methods*. 2014;11(7):740-742.
76. van Dijk D, Sharma R, Nainys J, et al. Recovering Gene Interactions from Single-Cell
Data Using Data Diffusion. *Cell*. 2018;174(3):716-729 e727.
77. Finak G, McDavid A, Yajima M, et al. MAST: a flexible statistical framework for
assessing transcriptional changes and characterizing heterogeneity in single-cell RNA
sequencing data. *Genome Biol*. 2015;16:278.
78. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol*. 2018;19(1):15.

79.   Dann E, Henderson NC, Teichmann SA, Morgan MD, Marioni JC. Differential
abundance testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol*.
2022;40(2):245-253.
80.   Granja JM, Corces MR, Pierce SE, et al. ArchR is a scalable software package for
integrative single-cell chromatin accessibility analysis. *Nat Genet*. 2021;53(3):403-411.
81.   Zhang Y, Liu T, Meyer CA, et al. Model-based analysis of ChIP-Seq (MACS). *Genome
Biol*. 2008;9(9):R137.
82.   Schep AN, Wu B, Buenrostro JD, Greenleaf WJ. chromVAR: inferring transcription-
factor-associated accessibility from single-cell epigenomic data. *Nat Methods*. 2017;14(10):975-
978.
83.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell
data. *Cell*. 2021;184(13):3573-3587 e3529.
84.   Cunningham F, Allen JE, Allen J, et al. Ensembl 2022. *Nucleic Acids Res*.
2022;50(D1):D988-D995.
85.   Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
using empirical Bayes methods. *Biostatistics*. 2007;8(1):118-127.
86.   Fang Z, Liu X, Peltz G. GSEApy: a comprehensive package for performing gene set
enrichment analysis in Python. *Bioinformatics*. 2023;39(1).
1021

FIGURE LEGENDS

Figure 1: Aged HSCs Exhibit Loss of Multilineage Potential

(**A-H**) Multiome single-cell RNA and ATAC sequencing was performed on HSC (Lineage-Sca-

1+cKit+ CD34-CD48-Flt3-CD150+) cells isolated from 2-mo (young) or 24-mo (old) female

C57BL/6 mice. nTOTAL =12,350 cells. (**A**) Volcano plot of DGE analysis between MLin-HSC

vs. q-HSC subsets in young HSCs, highlighting bonafide HSC markers. (**B**) Violin plot for

imputed *Kit* gene expression by annotated HSC subsets (as described in **Fig. S2C**. (**C**) Frequency

of computationally defined HSC subtypes in Kit^{hi} and Kit^{lo} subsets in young HSCs. (**D-E**)

Combined Uniform manifold approximation and projection (UMAP) of young and old HSCs

after Harmony batch correction annotated by age (**D**) and annotated HSC subsets (q-HSC:

Quiescent HSCs; Mgk-HSCs: Platelet-biased HSCs; Mlin-HSC: Multilineage HSCs; p-HSC:

Proliferative HSCs; Int-HSC: Intermediate HSCs) (**E**). (**F**) Frequency of HSC subtypes in young

and old HSCs. (**G**) Heatmap showing gene expression of bonafide HSC markers ordered by

increasing Kit expression. **(H)** Violin plot for imputed Kit gene expression by age. Statistical

analysis was performed using the Wilcoxon test. **(I)** Scaled change in frequency for each Kit

HSC subset with age. **(J)** FACS analysis showing frequency of Kit^{hi} and Kit^{lo} HSCs by age. All

data are from n=6 mice/group. Error bars represent mean ± SEM. **P<0.01, ***P<0.001. P

values calculated by two-way ANOVA.

1040

Figure 2: Young Kit^{lo} HSCs Exhibit Enhanced *In Vitro* and *In Vivo* Lymphoid Potential

(A) Experimental schema to evaluate *in vitro* lymphoid progenitor and T-cell differentiation

potential of Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 2-mo (young) C57BL/6 mice using the S17

lymphoid assay and murine artificial thymic organoid (mATO) respectively. Enumeration of

absolute number of lymphoid progenitor cells following 14 days in S17 lymphoid assay (B) and

absolute number of T-cell subsets following 8-weeks of culture in mATOs **(C).** Aggregated data

across 4-5 independent experiments. **(D)** Experimental schema for competitive allogeneic HCT

(allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 2-mo (young) C57BL/6 mice with

competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ mice transplanted into

lethally irradiated 7-week-old (young) BALB/cJ recipients. **(E-K)** Eight weeks after competitive

HCT, **(E)** frequency of donor-derived chimerism (CD45.2/ H-2Kb) of mature lineages in the

peripheral blood (PB, CD45+, Myeloid cell: Gr1+CD11b+; B-cell: B220+; T-cell: CD3+), **(F-G)**

Enumeration of absolute number of donor-derived cells in the BM for LSK cells (LT HSC:

Lineage$^-$Sca-1$^+$cKit$^+$ CD34$^-$CD48$^-$Flt3$^-$CD150$^+$ **;** ST HSC: Lineage$^-$Sca-1$^+$cKit$^+$ CD48$^-$Flt3$^-$

CD150$^-$**;** MPP2: Lineage$^-$Sca-1$^+$cKit$^+$ CD48$^+$Flt3$^-$CD150$^-$; MPP3: Lineage$^-$Sca-1$^+$cKit$^+$

CD48$^+$Flt3$^-$CD150$^+$; MPP4/ LMPP: Lineage$^-$Sca-1$^+$cKit$^+$ Flt3$^+$CD150$^-$) **(F),** Common Lymphoid

Progenitors cells (CLP: Lineage$^-$IL7Ra$^+$Flt3$^+$Sca$^{mid/lo}$ Kit^{lo}) **(G). (H-K)** Post-HCT thymi analysis

for total thymic cellularity **(H)**, **(I-J)** enumeration of absolute number of donor-derived cells for

T-cell precursors (ETP: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^+$ CD25$^-$Kit$^+$;DN2: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^+$

CD25$^+$; DN3: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^-$ CD25$^+$) **(I)**, Mature T-cells (DP: Lineage$^-$ CD4$^+$ CD8$^+$;

SP4: Lineage$^-$ CD4$^+$ CD8$^-$; SP8: Lineage$^-$ CD4$^-$ CD8$^+$) **(J),** and analysis of thymic CD45-

compartment for Thymic Epithelial Cells (TEC: CD45$^-$ EpCAM$^+$), Cortical TEC (cTEC: CD45$^-$

EpCAM$^+$UEA-1^{lo} 6C3^{hi} MHCII$^{hi/lo}$), and Medullary TEC (mTEC: CD45$^-$ EpCAM$^+$UEA-1^{hi}

6C3^{lo} MHCII$^{hi/lo-}$) **(K)**. All data are from n=10-11 mice/group. **(L)** Experimental schema to

evaluate thymic function following competitive allo-HCT using young HSCs. **(M)** Frequency of

donor-derived Recent Thymic Emigrants (RTE: CD3+GFP+) in the PB at 8 weeks post-HCT.

All data are from n=5-6 mice/group. **(N)** Experimental schema to investigate functional response

of differential Kit-expressing HSC-derived T-cells in secondary recipients to *L. monocytogenes*

expressing chicken ovalbumin (LM-OVA) infection. **(O)** Spleen analysis of secondary recipients

for frequency of OT1-derived chimerism of CD8+ T cells (OT1+CD8+: Va2$^+$Vb5$^+$H2-K^{b}CD8$^+$)

7 days post-infection. All data are from n=9-10 mice/group. **(P)**   Experimental schema for

competitive allogeneic HCT (allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 2-mo

(young) C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-

PtprcaPepcb/BoyJ mice transplanted into lethally irradiated 14-mo (Middle-aged) BALB/cJ

recipients. **(Q)** Frequency of donor-derived T-cell chimerism at the indicated time-points. **(R-T)**

Sixteen weeks after competitive HCT enumeration of total thymic cellularity **(R)**, donor-derived

T-cell precursor thymocyte subsets as defined in Figure 3F **(S)**, donor-derived mature T-cell

thymocyte subsets as defined in Figure 3G **(T).** All data are from n=6-7 mice/group. Error bars

represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. P values calculated by

nonparametric unpaired Mann-Whitney U test. Panels **A**, **D**, **L**, **N**, **and P** were created using

BioRender.

1082

Figure 3: Kit^{lo} HSCs are Epigenetically Programmed for Lymphoid Potential

(A) UMAP of 6 unsupervised HSC clusters identified by tile-based scATAC-seq analysis after

Harmony batch correction. **(B-D)** Annotated by age **(B)**, HSC subsets **(C)**, and Kit subsets **(D)** on

the scATAC-seq UMAP. **(E)** scATAC-seq cluster composition by HSC subsets (**left**) and Kit

subsets (**right**). (**F-G**) Matrix plots showing motif enrichment identified by ChromVAR (**F**) and

gene accessibility (**G**) for lymphoid-specifying transcription factors for Kit subsets in young and

old HSCs.

1090

Figure 4: *Zbtb1* regulates T-cell potential in phenotypic HSCs

(A) Experimental schema for T-cell differentiation assay and competitive allogeneic HCT with

Zbtb1-deficient Kit^{lo} HSCs generated with 2-mo young Rosa26Cas9-eGFP KI mice. **(B)**

Frequency of T-cell subsets following 6-weeks of culture in mATOs. Aggregated data across 3

independent experiments. (C) Frequency of donor-derived lineage chimerism of control or *Zbtb1*

KO Kit^{lo} HSCs in the PB at 6 weeks following competitive HCT. All data are from n=5-6

mice/group. (D) Experimental schema for T-cell differentiation assay and competitive allogeneic

HCT with *Zbtb1*-OE Kit^{hi} HSCs from young mice. (E) Frequency of T-cell subsets following 6-

weeks of culture in mATO of young Kit^{hi} HSCs. Aggregated data across 3 independent

experiments. (F) Frequency of donor-derived lineage chimerism of control or *Zbtb1* OE young

Kit^{hi} HSCs in the PB at 6-weeks following competitive HCT. All data are from n=5-6

mice/group. (G) Previously identified Zbtb1-targets[56] and Notch1-interacting proteins[58] enriched

in Kit^{lo} HSCs. (H) Pathway enrichment analysis for non-Notch1 Zbtb1-targets enriched in Kit^{lo}

HSCs by *gseapy()*[86] using GO_Biological_Process_2023 libraries. Bubbleplot showing

representative pathways. Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001. P

values calculated by nonparametric unpaired Mann-Whitney U test. Panels **A** and **D** were created

using BioRender.

1108

Figure 5: Old Kit^{lo} HSCs Exhibit Enhanced *In Vitro* and *In Vivo* Lymphoid Potential

(A-B) Evaluation of *in vitro* lymphoid progenitor and T-cell differentiation potential of Kit^{hi}

(red) and Kit^{lo} HSCs (blue) from or 24-mo (old) C57BL/6 mice using the S17 lymphoid assay

and murine artificial thymic organoid (mATO) respectively, as shown in Figure 2A.

Enumeration of absolute number of lymphoid progenitor cells following 14 days in S17

lymphoid assay **(A)** and absolute number of T-cell subsets following 8-weeks of culture in

mATOs **(B).** Aggregated data across 4-5 independent experiments. (**C**) Experimental schema for

competitive allogeneic HCT (allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 22–24-mo

(old) C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ

mice transplanted into lethally irradiated young BALB/cJ recipients. **(D-I)** Eight weeks after

competitive HCT enumeration of absolute number of donor-derived T-cells in the PB **(D)**, LSK

cell subsets as defined in Figure 3C (**E**), CLP cells as defined in Figure 3D (**F**), total thymic

cellularity **(G)**, donor-derived T-cell precursor thymocyte subsets as defined in Figure 3F **(H)**,

TEC compartment as defined in Figure 3H **(I).** All data are from n=9-10 mice/group. **(J)**

Experimental schema to evaluate thymic function following competitive allo-HCT using old

HSCs. **(K)** Frequency of donor-derived Recent Thymic Emigrants (RTE: CD3$^+$GFP$^+$) in the PB

at 8 weeks post-HCT. All data are from n=5 mice/group. **(L)** Experimental schema for

competitive allogeneic HCT (allo-HCT) using Kit[hi] (red) and Kit[lo] HSCs (blue) from 22-24-mo

(old) C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ

mice transplanted into lethally irradiated middle-aged BALB/cJ recipients. **(M-O)** Eight weeks

after competitive HCT enumeration of absolute number of donor-derived T cells in PB **(M)**,

donor-derived CLPs in the BM as defined in Figure 3D **(N)**, and thymic cellularity **(O)**. All data

are from n=5-6 mice/group. **(P)** Frequency of T-cell subsets following 6-weeks of culture in

mATO of old Kit[hi] HSCs following *Zbtb1* OE. Aggregated data across 3 independent

experiments. Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001. P values calculated by nonparametric unpaired Mann-Whitney U test. Panels **C,**

J, and **L** were created using BioRender.

1136

Figure 6 KIT[lo] Human BM HSC Subsets show Higher Lymphoid Potential

(A-E) Young and old human BM CITE-seq dataset generated and published by Sommarin et

al.[62] **(A-C)** UMAP of CD34+ young and old human BM annotated with HSC subsets **(A)**, and

age **(B)**. **(C)** HSC cluster composition by age. **(D)** Kit[lo] gene signature (top 50 marker genes

from our mouse data converted to human orthologues) was overlaid on human HSC UMAP. **(E)**

Violin plot for Kit[lo] gene score by age. Statistical analysis was performed using the Wilcoxon

test. **(F-G)** FACS analysis of human BM samples. **(F)** Violin plot of KIT protein expression on

phenotypic HSCs (pHSC: CD34+CD38-CD10-CD45RA-CD90+) by age (young: 20-40yrs;

MA/old: >40yrs). **(G)** Scatterplot showing distribution of KIT[hi] and KIT[lo] HSC subsets within

pHSCs in human BM samples with varying age. **(H)** Experimental schema to evaluate *in vitro* T-

cell and lympho-myeloid potential of human BM HSC subsets based on differential KIT

expression using human artificial thymic organoids (hATO) and MS5 assay respectively. **(I)**

Following 8-10-weeks of culture, enumeration of CD19+ B-cells and T-cell subsets within

CD34+ T-cell precursors (Early Thymic Progenitors, ETP: $CD34^+CD1a^-CD7^-$; $CD1a^{neg}$- ProT:

$CD34^+CD1a^-CD7^+$; $CD1a^{pos}$-ProT: $CD34^+CD1a^+CD7^+$), and mature T-cells (DP: $CD34^-$

$CD5^+CD7^+CD4^+CD8^+$; SP4: $CD34^-CD5^+CD7^+CD4^+CD8^-$; SP8: $CD34^-CD5^+CD7^+CD4^-CD8^+$).

Aggregated data represent mean from 8 independent BM donors ± SEM. **(J)** Following 4-weeks

of culture, enumeration of Lineage output (Neutrophil-Granulocytes: hCD45+CD15+;

monocyte-macrophages: hCD45+CD14+; B-cells: hCD45+CD19+; NK-cells: hCD45+CD56+)

after culturing 200 KIT^{lo} and KIT^{hi} HSC cells on MS-5 stroma, as previously described[63].

Aggregated data represent mean from 5 independent BM donors ± SEM. **(L)** Model illustrating

age-related loss of lymphoid potential is attributed to a decreased in the frequency of Kit^{lo} HSCs

in aged BM.   Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001. P values for panel **G** was based on generalized estimating equation logistic

model adjusted for age, KIT^{lo} and KIT^{hi} groups and their interaction, the matched data structure

captured by an exchangeable working correlation structure. P values calculated either by

nonparametric paired Mann-Whitney U test (Panels **I** and **J**), unpaired Mann-Whitney U test

(Panel **F**). Panels **H** and **L** were created using BioRender.

1165

<u>**Supplemental Figure Legends**</u>

Figure S1: (A) Experimental schema for competitive allogeneic HCT (allo-HCT) using 2-mo

(young) and 24-mo (old) 750 HSCs [Lin^-c-Kit^+Sca1^+ ($L^-S^+K^+$) $CD150^+$ $CD34^-CD48^-$] from

C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ mice

transplanted into lethally irradiated 7-week-old (young) BALB/cJ recipients. **(B-E)** Frequency of

donor-derived chimerism of all hematopoietic cells mature lineages **(B),** T-cell **(C),** Myeloid **(D),**

B-cell **(E)** in the PB at the indicated time points. **(F-G)** Eight weeks after competitive HCT

enumeration of absolute number of LMPP cells (LMPP/MPP4: Lineage$^-$Sca-1$^+$cKit$^+$ Flt3$^+$CD150$^-$

) **(F)**, CLP cells (CLP: Lineage$^-$IL7Ra$^+$Flt3$^+$Sca$^{mid/lo}$ Kit^{lo}) **(G)**. **(H-J)** Post-HCT thymi analysis

for total thymic cellularity **(H)**, enumeration of absolute number of donor-derived cells for T-cell

precursors (ETP: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^+$ CD25$^-$Kit$^+$; DN2: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^+$

CD25$^+$; DN3: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^-$ CD25$^+$) **(I)**, Mature T-cells (DP: Lineage$^-$ CD4$^+$ CD8$^+$;

SP4: Lineage$^-$ CD4$^+$ CD8$^-$; SP8: Lineage$^-$ CD4$^-$ CD8$^+$) **(J)**. All data are from n=9-10 mice/group.

Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. P values

calculated by nonparametric unpaired Mann-Whitney U test. Panel **A** was created using

BioRender.

1182

Figure S2: (A) UMAP of young HSCs after Harmony batch correction annotated by

unsupervised Leiden cluster analysis (**B**), manually annotated HSC subsets (q-HSC: Quiescent

HSCs; Mgk-HSCs: Platelet-biased HSCs; Mlin-HSC: Multilineage HSCs; p-HSC: Proliferative

HSCs; Int-HSC: Intermediate HSCs) (**C**). (**D**) Dot plot showing marker genes for each HSC

subset. Circle size and color indicate percentage and expression level, respectively. (**E**)

Composition of defined HSC subtypes in Kit^{hi} and Kit^{lo} subsets in young HSCs as shown by

scaled change in frequency.

(**F**) UMAP of old HSCs annotated by HSC subsets identified by *scanpy ingest*()[78] using young

HSCs, as reference dataset. **(G)** UMAP of combined young and old HSCs after Harmony batch

correction annotated by sample. (**H**) Composition of defined HSC subtypes in old and young

HSCs as shown by scaled change in frequency. (**I**) A neighborhood graph of the results from

Milo differential abundance testing. Nodes are neighborhoods, colored by their log fold change

across ages. Non-differential abundance neighborhoods (FDR 10%) are colored white, and sizes correspond to the number of cells in each neighborhood. Graph edges depict the number of cells shared between neighborhoods. The layout of nodes is determined by the position of the neighborhood index cell in the UMAP in panel **Figure 1D. (J)** Beeswarm plot of the distribution of log fold change across age in neighborhoods containing cells from different cell type clusters. Differential abundance neighborhoods at FDR 35% are colored. HSCs detected as differentially abundant annotated by HSC subsets (as defined in **Figure 1A**). **(J) (K)** Histogram for magic imputed *Kit* gene expression by age. **(L)** Representative FACS plots showing the gating strategy to define Kit^{hi} and Kit^{lo} HSC subsets in BM from young and old mice.

Figure S3: (A-H) Following allo-HCT (**as shown in Figure 2D**), frequency of donor-derived chimerism of hematopoietic cells **(A)**, mature lineages: Myeloid cells, B-cells, and T-cells **(B)**, in the PB at the indicated time points. **(C-D)** Enumeration of absolute number of donor-derived MP subset cells in the BM as defined in Figure S1H **(C)**, thymic non-TEC compartment (endothelial cells: CD45$^-$ EpCAM$^-$Ter-119 PDGFR◻$^-$ CD31$^+$; fibroblasts: CD45$^-$ EpCAM$^-$Ter-119$^-$CD31$^-$ PDGFR◻$^+$) **(D)**. **(E)** Relative expression of thymic-adhesion molecules (PSGL1, CCR7, and CCR9) on donor-derived CLPs (Statistical analysis was performed using ComBat[85] to perform batch-effect correction for the mean MFI across two experiments), **(F)** Absolute amount of intrathymic thymopoietic ligands, IL-22 (left), RANKL (right) was measured by ELISA after allo-HCT. **(G-H)** Frequency of donor-derived **(G)** CD4+ and **(H)** CD8+ T-cell (naïve: CD62L+CD44- ; Central Memory: CD62L+CD44+; Effector Memory: CD62L-CD44+) subsets in the spleen. All data are from n=10-11 mice/group. **(I-O)** Following allo-HCT (**as shown in Figure 2P),** frequency of donor-derived chimerism of mature lineages, myeloid (**I**) and B-cells

1218　(**J**) in the PB at the indicated timepoints. (**K-O**) Enumeration of absolute number of donor-

1219　derived LSK (**K**) **and** MP subset cells (**M**) in the BM as defined in Figures 2F and S1H,

1220　respectively**,** thymic TEC (**N**) **and** non-TEC (**O**) compartments as defined in Figures 2K and

1221　S3D**.** Analyses were performed at 16 weeks following competitive HCT. All data are from n=9-

1222　10 mice/group.

1223

Figure S4: Competitive allo-HCT using Kit subsets from young mice, as described in Figure 2D.

Twenty weeks after competitive HCT, (**A**) frequency of donor-derived chimerism of mature

lineages in the peripheral blood as described in Figure 2E.Enumeration of absolute number of

donor-derived cells in the BM for LSK cell subsets as defined in Figure 2F (**B**), MP cell subsets

as defined in Figure S1H (**C**), CLP cells as defined in Figure 2G (**D**), total thymic cellularity (**E**),

donor-derived mature T-cells (**F**) and T-cell precursor thymocyte subsets (**G**) as defined in

Figures 3I-3J, thymic stromal compartment (CD45- cells) as defined in Figures 2K and S3D (**H**).

All data are from n=8 mice/group. Error bars represent mean ± SEM. *P<0.05, **P<0.01,

***P<0.001. P values calculated by nonparametric unpaired Mann-Whitney U test.

1233

Figure S5: (A-E) Single-cell ATAC seq pre-processing and QC analysis. (**A**) Comparison of

TSS enrichment score and number of fragments. The dashed lines depict the cutoffs used. (**B**)

Ridge plots showing distribution of TSS enrichment scores per sample (**top**) and distribution of

unique nuclear fragments per sample (**bottom**). (**C**) Plots showing fragment size distribution

(**top**) and TSS enrichment profile (**bottom**) across samples. (**D**) scATAC-seq UMAP without

batch correction colored by sample. (**E**) Volcano plots of differentially accessible (DA) peaks

(fold change> 2, FDR <0.01) for HSC subset Cluster 3 vs. Cluster 6 (Wilcoxon rank-sum test).

1241

Figure S6: Zbtb1 expression assessed by flow cytometric analysis following knock out (KO) in Rosa26Cas9 KI Kit^{lo} HSCs (**A**) and OE of Zbtb1 cDNA in Kit^{hi} HSCs (**B**).

1244

Figure S7: (**A-F**) Following allo-HCT (**as shown in Figure 3C**)**,** frequency of donor-derived chimerism of mature lineages in the PB **(A)**. **(B-F)** Enumeration of absolute number of donor-derived MP subset cells in the BM as defined in Figure S1H **(B),** donor-derived mature T-cells as defined above **(C)**, thymic non-TEC compartment as defined in Figure S3D **(D),** and donor-derived **(E)** CD4+ T-cell and **(F)** CD8+ T-cell subsets in the spleen as defined above. All data are from n=9-10 mice/group. All analyses were performed at 8 weeks following competitive HCT. **(G)** Following allo-HCT (**as shown in Figure 3L**), frequency of donor-derived chimerism of mature lineages in the peripheral blood. Analyses were performed at 8 weeks following allo-HCT. All data are from n=5-6 mice/group. **(H-O)** Competitive allo-HCT using Kit subsets from old mice, as described in Figure 3C. Twenty weeks after competitive HCT, (**H**) frequency of donor-derived chimerism of mature lineages in the peripheral blood as described in Figure 2E.Enumeration of absolute number of donor-derived cells in the BM for LSK cell subsets as defined in Figure 2F (**I**), MP cell subsets as defined in Figure S1H (**J**), CLP cells as defined in Figure 2G (**K**), total thymic cellularity (**L**), donor-derived mature T-cells (**M**) and T-cell precursor thymocyte subsets (**N**) as defined in Figures 3I-3J, thymic stromal compartment (CD45- cells) as defined in Figures 2K and S3D (**O**). All data are from n=6-7 mice/group. Error bars represent mean ± SEM. Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. P values calculated by nonparametric unpaired Mann-Whitney U test.

1263

Figure S8: Equal Competition allo-HCT using Kit^{lo} HSCs from young and old mice. (**A**)

Experimental schema for equal competition allogeneic HCT (allo-HCT) using Kit^{lo} HSCs from

2-mo young (blue) and 22-24-mo old (light-blue) C57BL/6 mice combined with competitor bone

marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ mice transplanted into lethally irradiated 7-

week-old (young) BALB/cJ recipients. Twenty weeks after competitive HCT, (**B**) frequency of

donor-derived chimerism of mature lineages in the peripheral blood as described in Figure 3B.

Enumeration of absolute number of donor-derived cells in the BM for LSK cell subsets as

defined in Figure 2F (**C**), MP cell subsets as defined in Figure S1H (**D**), CLP cells as defined in

Figure 2G (**E**), donor-derived T-cell precursor thymocyte subsets as defined in Figure 3I (**F**) and

mature T-cells as defined in Figure 3J (**G**). All data are from n=9 mice. Error bars represent

mean ± SEM. (**H**) Pathway enrichment analysis performed on differentially expressed genes

comparing old versus young Kit^{lo} HSCs by *gseapy*()[86] using Hallmark_2020 and

GO_Biological_Process_2023 libraries. Bubbleplot showing representative pathways enriched in

young Kit^{lo} HSCs. (**I-K**) Matrix plots showing motif enrichment identified by ChromVAR (I),

gene accessibility score (**J**), and gene expression (**K**) for Kit^{lo} subset in young and old HSCs.

*P<0.05, **P<0.01, ***P<0.001. P values calculated by nonparametric unpaired Mann-Whitney

U test.

1281

1282   **Figure S9:** (**A**) UMAP of CD34+ young and old human BM CITE-seq[62] annotated with 15

1283   clusters. (**B**) Violin plot for CD117 ADT-expression by age. Statistical analysis performed using

1284   Wilcoxon test. (**C**) Representative FACS plots of human BM samples showing gating strategy to

1285   define HSPC subsets. (**D**) Representative FACS plots of human BM samples showing gating

1286   strategy to define KIT^{hi} and KIT^{lo} HSCs in human BM.

1287

Young Donor/ Young Recipients Tx

Young Donor/ MA Recipients Tx

HSC Annotations

Young Donor/ Young Recipients Tx

Young Donor/ MA Recipients Tx

Young Donor/ MA Recipients Tx

A

B

Old Donor/ Young Recipients Tx

A PB chimerism

B MP compartment

C Donor-derived

D

E CD4 T cells

F CD8 T cells

Old Donor/ MA Recipients Tx

G

Old 20-week Post HCT

H PB chimerism

I LSK

J MP

K CLP

L

M Mature Thymocytes

N Thymocyte Precursors

O Stromal analysis

Equal Competition Tx

A Allogeneic HCT — Kit^lo HSC Donors — Young CD45.1/CD45.2 + Old B6 CD45.2 → Host Young BALB/cJ; Young CD45.1 Rescue BM

B PB chimerism — T-cell, B-cell, Myeloid

C LSK — MPP4 (LMPP), MPP3, MPP2, ST HSC, LT HSC

D MP — MEP, GMP, CMP

E CLP — Young Kit^lo HSC, Old Kit^lo HSC

F Mature Thymocytes — SP8, SP4, DP

G Thymocyte Precursors — DN3, DN2, ETP

H Pathways Enriched in Young Kit^lo HSCs — % Genes in set; log₁₀(1/FDR)

I chromVAR Motif Enrichment — Zbtb1_182, Zbtb7a_808, Runx3_858; Z-Score

J Gene Accessibility — Zbtb1, Zbtb7a, Ezh1; Z-Score

K Gene Expression — Zbtb1, Zbtb7a, Ezh1; Z-Score

A

C

B

ADT−CD117

D **Gating Strategy to Fractionate Kit Subsets in Human BM**