An epigenetically distinct HSC subset supports thymic reconstitution

Harold K. Elias[1,2], Sneha Mitra[3], Marina B. da Silva[1,14], Adhithi Rajagopalan[1,14], Brianna Gipson[1], Nicole Lee[1], Anastasia I. Kousa[1], Mohamed A.E. Ali[4], Simon Grassman[1], Xiaoqun Zhang[1], Susan DeWolf [6], Melody Smith[5], Hana Andrlova[1], Kimon V. Argyropoulos[1], Roshan Sharma[7], Teng Fei[8], Joseph C Sun[1], Cynthia E. Dunbar[9], Christopher Y Park[4], Christina S. Leslie[3], Avinash Bhandoola[10], and Marcel R.M. van den Brink[1,11,12,13]

Affiliations
[1]Department of Immunology, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[2]National Institutes of Health (NIH), Bethesda, MD, 20892, USA, USA
[3]Computational and Systems Biology, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[4]Department of Pathology, New York University Grossman School of Medicine, New York, NY, 10016, USA
[5]Division of Blood and Marrow Transplantation and Cellular Therapy, Department of Medicine, Stanford University School of Medicine, Stanford, CA, 94305
[6]Leukemia Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[7]Single-cell Analytics and Innovation Lab, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[8]Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA
[9]Translational Stem Cell Biology Branch, National Heart, Lung, and Blood Institute, NIH, Bethesda, MD, 20892, USA
[10]Laboratory of Genome Integrity, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, MD, 20892, USA
[11]Department of Immunology and Microbial Pathogenesis, Weill Cornell Medical College; New York, NY
[12]Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, 10065, USA.

[13]Lead Contact
[14]These authors contributed equally
Correspondence: eliash@mskcc.org (HKE); mvandenbrink@coh.org (MRMvdB)

ABSTRACT

Hematopoietic stem cells (HSCs) with multilineage potential are critical for effective T cell reconstitution and restoration of the adaptive immune system after allogeneic Hematopoietic Cell Transplantation (allo-HCT). The Kit[lo] subset of HSCs is enriched for multipotential precursors,[1, 2] but their T-cell lineage potential has not been well-characterized. We therefore studied the thymic reconstituting and T-cell potential of Kit[lo] HSCs. Using a preclinical allo-HCT model, we demonstrate that Kit[lo] HSCs support better thymic recovery, and T-cell reconstitution resulting in improved T cell responses to infection post-HCT. Furthermore, Kit[lo] HSCs with augmented BM lymphopoiesis mitigate age-associated thymic alterations, thus enhancing T-cell recovery in middle-aged hosts. We find the frequency of the Kit[lo] subset declines with age, providing one explanation for the reduced frequency of T-competent HSCs and reduced T-lymphopoietic potential in BM precursors of aged mice.[3, 4, 5] Chromatin profiling revealed that Kit[lo] HSCs exhibit higher activity of lymphoid-specifying transcription factors (TFs), including *Zbtb1*. Deletion of *Zbtb1* in Kit[lo] HSCs diminished their T-cell potential, while reinstating *Zbtb1* in megakaryocytic-biased Kit[hi] HSCs rescued T-cell potential, *in vitro* and *in vivo*. Finally, we discover an analogous Kit[lo] HSC subset with enhanced lymphoid potential in human bone marrow. Our results demonstrate that Kit[lo] HSCs with enhanced lymphoid potential have a distinct underlying epigenetic program.

KEYWORDS: Hematopoietic stem cells; lymphoid differentiation; epigenetic priming; thymus; immune reconstitution; immunosenescence.

INTRODUCTION

T lymphocytes play a crucial role in the adaptive immune response. A highly diverse T cell pool is required to recognize and eliminate foreign pathogens while also maintaining self-tolerance. The thymus continuously produces T cells, but with advancing age, its regenerative capacity declines,[6] leading to decreased thymic output and compromised T cell diversity. Furthermore, the thymus is also vulnerable to acute damage from infections, cancer therapies, and conditioning regimens for allogeneic hematopoietic cell transplantation (allo-HCT), which results in prolonged T cell lymphopenia, thereby increasing the risk of infections and cancer relapse, contributing to transplant-related complications and mortality.[7] Likely for these reasons, early T-cell reconstitution is a positive prognostic indicator of allo-HCT outcomes.[8, 9]

Post-transplant T cell reconstitution requires a steady supply of hematopoietic stem cells (HSCs)-derived lymphoid progenitors[10] as well as thymic recovery. Prior research has demonstrated that thymic integrity is dependent on thymocyte-stromal crosstalk, particularly in the thymic epithelial cell (TEC) compartment.[11, 12, 13] Age-associated reduced lymphoid potential originating at the level of HSCs,[14] coupled with thymic decline[6] contribute to delayed immune recovery. Therefore, identifying strategies to augment T lymphopoiesis and promote thymic regeneration is an unmet clinical need.

Single-cell transplantation with tracking and sequencing of progeny[15, 16, 17, 18, 19, 20] has uncovered heterogeneity in self renewal capacity as well as lineage output among reconstituting HSCs-highlighting HSCs with multilineage versus lineage-biased potential. These studies showed that, while only a fraction of HSCs generate "balanced" multilineage output, the majority exhibit a diverse range of lineage potential following transplantation- varying in their contributions and reconstitution kinetics for each lineage, thereby showcasing their inherent biases. These lineage

biases are intrinsically stable and maintained following transplantation.[21, 22, 23] At the molecular

level, these cell autonomous features are mediated by epigenetic configuration and are divergent

from their transcriptional state.[18, 24] Although this suggests the existence of a highly organized

and predictable framework for lineage-restricted fates of long-term self-renewing HSCs, the

molecular identities, gene regulatory networks and functional implications of these differences

governing lymphoid fate decisions remain largely unexplored.

Kit^{lo} HSCs have previously been described to exhibit better self-renewal and multipotency,

including improved T-cell reconstitution in syngeneic transplant models. [1, 2]However, their

thymic reconstituting ability, age-related changes, as well as the molecular basis for their

enhanced T-cell lymphoid potential are unknown. Here, we delineate the pivotal role of Kit^{lo}

HSCs in orchestrating thymic regeneration and immune reconstitution in both young and aged

hosts. We find the frequency of the Kit^{lo} subset declines with age. Leveraging integrated single

cell transcriptional and epigenetic profiling, we uncover the distinct epigenetic signature of Kit^{lo}

HSCs, highlighting the presence of lymphoid-specifying transcription factor *Zbtb1* and its role in

enhancing T-cell lymphoid potential. Additionally, we identify an analogous HSC subset in

human bone marrow and demonstrate their enhanced lymphoid potential.

RESULTS

Reduced lymphoid output in aged HSCs impairs thymic recovery

Previous studies have shown impaired lymphoid reconstitution with aged HSCs in a syngeneic

transplant model, yet this has not been explored using an allogeneic setting. We used a pre-

clinical allo-HCT model to analyze immune reconstitution of young and old donors. To evaluate

if HSCs with differences in lymphoid progenitor production could impact thymic recovery, we

transplanted Lin−Sca-1+Kit+ (LSK) CD34−CD150+CD48− cells LT-HSCs from young adult

(2-mo) or old (24-mo) C57BL/6 mice with rescue BM cells (CD45.1) into lethally irradiated

young BALB/cJ recipients (**Figure S1A**). Because HSC-derived thymic progeny emerge within

5-6 weeks,[25] we established an 8-week harvest timepoint. This interval allowed sufficient time

for HSC-derived progenitors to repopulate the thymus, thereby providing a better understanding

of thymic regeneration dynamics and the efficacy of HSC transplantation in promoting thymic

reconstitution.

Consistent with prior reports, eight weeks post-HCT, we found that young HSCs supported

balanced lineage reconstitution, while old HSCs exhibited significantly impaired T-cell

reconstitution in the peripheral blood (PB), despite comparable reconstitution patterns at four

weeks (**Figures S1B-S1E**). Furthermore, we observed greater multilineage potential of young

HSCs, including lymphoid and myeloid reconstitution (LMPP/MPP4 and CLP, **Figures S1F and**

S1G), when compared to old HSCs. Furthermore, young HSC recipients had significantly greater

thymic cellularity and reconstitution of precursor and mature thymocytes than recipients of old

HSCs (**Figures S1H-S1J**). In conclusion, old HSCs have significantly less multi-lineage

potential, but especially T lymphoid potential than young HSCs in young allo-HCT recipients.

Age-related decline in multilineage HSCs marked by low Kit expression

Recent reports have shown remarkable functional heterogeneity of HSC[26, 27, 28, 29, 30, 31] subsets,

highlighting discordance between current HSC phenotypic definitions to their molecular

identities.[16, 17, 18] To identify HSCs with an enriched multilineage program and gain insights into

their transcriptional and epigenetic framework, we performed multiome single-cell RNA and

ATAC sequencing on hematopoietic stem cells (LT-HSCs: Lineage⁻CD34⁻CD48⁻CD150⁺Sca-

1⁺c-Kit⁺) from 2-mo (young) or 24-mo (old) female mice.

To define HSC subsets, we performed unsupervised Leiden cluster analysis on the multiome

scRNA-seq data from our young cohort (**Figure S2A-S2B**). Using nomenclature and signatures

derived from previous studies, [15, 16, 18, 32, 33, 34, 35] we mapped all unsupervised clusters into five

main subsets (**Figure S2C**): Quiescent HSC (q-HSC), Platelet-biased HSC (Mgk-HSC),

Multilineage HSC (MLin-HSC), Proliferative HSC (p-HSC), and a novel Intermediate HSC (Int-

HSC) that showed overlapping features with Mgk-HSC, MLin-HSC and p-HSC subsets (**Data**

Table 1, Figure S2D). To identify HSCs with multilineage potential, we compared multilineage

enriched MLin-HSC versus low output q-HSC subsets in young HSCs and queried for previously

described bonafide HSC markers (*Hlf, Kit, Neo1, Procr, CD244a, Pdzk1ip1, Ly9, Hoxb5*). *Hlf*

and *Kit* had significantly higher expression in q-HSCs compared to MLin-HSCs (**Figure 1A**).

Hlf has been previously implicated in HSC quiescence,[36, 37, 38] whereas *Kit* enriches for platelet-

biased HSCs.[2] To further analyze the composition of HSCs with varying Kit expression, we

defined Kit high and low HSC populations (Kit^{hi} and Kit^{lo} HSCs) based on the top and bottom

20% of *Kit* gene expression levels. We observed lower Kit expression in MLin-HSCs (**Figure**

1B) and increased representation of q-HSC and MLin-HSCs within the Kit^{lo} HSCs (**Figures 1C**

and S2E), consistent with prior reports of increased self-renewal capacity and multipotency in

Kit^{lo} HSCs.[1, 2]

To assess for age-associated changes, we first compared the transcriptional profile of young and

old HSCs. In our differential gene analysis, we found that HSCs from young mice had higher

expression of genes related to cell cycle (*Ccnd3*, *Cdc25a*, *Cdk1)*, and cell polarization

(*Arhgap15*); conversely, old HSCs were enriched for expression of genes related to quiescence

(*Egr1*, *Junb, Jun*, *Jund*, *Fos*), platelet differentiation (*vWf*, *Clu*, *Itgb3, Fhl1 and Tgm2*), and

myeloid differentiation (*Selp, Neo1*) (**Data Table 2**). Next to investigate age-related shifts in

HSC cluster composition, we employed *ingest()* from scanpy to query the old HSC

transcriptomic data using the "trained" young reference data for HSC subsets (**Figure S2F**).

Subsequently, we generated a UMAP of our young and old datasets (**Figures S2G and 1D**) to

visualize the post-ingestion results (**Figures 1E**). While all HSC subsets were present in young

and old cohorts (**Figure 1F**), we notably observed an increased representation of MLin-HSCs, p-

HSCs, and Int-HSCs in young HSCs. Conversely q-HSCs and Mgk-HSCs were predominantly in

old HSCs (**Figures S2H- S2J**). Consistent with prior work,[18, 27, 39] these findings suggest shifts in

the composition of HSC subsets with aging, indicating a reduction in multilineage potential

accompanied by an increase in platelet bias.

Given the increased frequency of Mgk-HSCs in old HSCs and prior work[2] demonstrating that

Kit^{hi} HSCs are platelet-biased HSCs (**Figures S2F- S2I**), we hypothesized this could be

attributed to an expansion in Kit^{hi} HSCs. We observed significantly higher *Kit* expression

(**Figures 1G-1H and S2K**) in old HSCs, with an age-dependent reduction in the proportion of

Kit^{lo} HSCs accompanied by an increase in the frequency of Kit^{hi} HSCs (**Figure 1I**). Flow

cytometry analysis of young and old bone marrow confirmed a significant expansion of Kit^{hi}

HSCs, accompanied by a concomitant decrease in Kit^{lo} HSCs in old HSCs (**Figures S2L and**

1J), consistent with our prior report.[40] Together, these findings indicate that the frequency of

multilineage Kit^{lo} HSCs declines with age.

Young Kit^{lo} HSCs exhibit enhanced T-cell potential

Next, we sought to determine the lymphoid potential of the HSC Kit subsets. To this end, we

used the S17 lymphoid progenitor assay[41] to assess for lymphoid progenitor output of purified

Kit[hi] and Kit[lo] HSCs from young mice (**Figure 2A**). Prior studies have elucidated the existence of

functional heterogeneity among phenotypic Common Lymphoid Progenitors (CLPs)

characterized by Ly6D expression,[42] resulting in the identification of two distinct subsets: All

Lymphocyte Progenitors (ALPs), also referred to as functional CLPs, and B-cell biased

Lymphocyte Progenitors (BLPs). Following 12-day co-culture, we observed that young Kit[lo]

HSCs generated more CLPs and ALPs than Kit[hi] HSCs (**Figure 2B**). To further examine their T-

cell differentiation potential, we used murine artificial thymic organoids (mATO, BM-ATO)[43, 44,]

[45] (**Figure 2A**). Following an 8-week culture, we noted a substantial increase in overall ATO

output from young Kitl[o] HSCs in comparison to Kit[hi] HSCs (**Figure 2C**).

To further evaluate the reconstituting potential of Kit[lo] HSCs, we competitively transplanted Kit[lo]

or Kit[hi] HSCs from young C57BL/6 (CD45.2) mice with rescue BM cells (CD45.1) into lethally

irradiated young BALB/cJ recipients (**Figure 2D**). Eight weeks after allo-HCT, we found that

Kit[lo] HSCs generated significantly better B- and T-cell reconstitution, while myeloid

reconstitution was comparable (**Figures 2E, S3A-S3B**). Concordant with a prior report of better

self-renewal ability,[2] we found that Kit[lo] HSCs demonstrated increased LT HSC reconstitution

(**Figure 2F**). In support of multilineage potential, we observed that Kit[lo] HSC recipients had

higher lymphoid progenitor reconstitution (including LMPP and CLP, **Figures 2F and 2G**),

along with comparable myeloid precursor output (including MPP3, CMP, GMP, MEP, **Figures**

2F and S3C).

To evaluate whether enhanced lymphoid precursor output could contribute to thymic

reconstitution, we concurrently evaluated the recipient thymi. We found that Kit[lo] recipients

demonstrated enhanced thymic recovery, characterized by significantly higher thymic cellularity

and reconstitution of precursor and mature thymocytes (**Figures 2H-2J**). Notably, in support of a

critical role for thymocyte-stromal crosstalk in preserving thymic architecture,[11, 12, 13] we also

found higher thymic epithelial cell (TEC), specifically medullary TEC, and endothelial cell

recovery in Kit^{lo} reconstituted thymi (**Figures 2K and S3D**). We next examined donor-derived

lymphoid progenitors for differential expression of molecules, involved in homing to the thymus

(CCR7, CCR9, and PSGL1).[46] We observed significantly higher expression of CCR7 on Kit^{lo}-

derived lymphoid progenitors (CLP) (**Figure S3E**). In addition, we analyzed thymic supernatants

and found no differences in levels of thymopoietic ligands, IL22[47] and RANKL[48] (**Figure S3F**).

Finally, we found significantly higher secondary lymphoid organ (spleen) reconstitution in Kit^{lo}

recipients (**Figures S3G and S3H**). To evaluate long-term reconstitution, we performed

additional harvests at 20-weeks following HCT. We observed similar trends in higher BM

lymphopoiesis, thymic reconstitution, and thymic stromal recovery in Kit^{lo} recipients (**Figure

S4**).

To assess thymic output, we used the RAG2-GFP model,[49] which allows the analysis of recent

thymic emigrants (RTEs) (**Figure 2L**). We observed higher numbers of PB GFP+ T cells in Kit^{lo}

recipients (**Figure 2M**). Additionally, to assess whether this increased number PB T cells in Kit^{lo}

recipients results in increased functionality, (**Figure 2N**), we analyzed response to ovalbumin

engineered *Listeria Monocytogenes* (LM-OVA) and observed a significant increase in OT1+

CD8+ T cells (**Figure 2O**).

A key element in the post-HCT recovery of thymic function is the influx of bone marrow-

derived lymphoid progenitors, also known as thymic seeding progenitors (TSPs).[10] Furthermore,

the significance of restoring the bone marrow-thymus axis in mitigating thymic involution has

been underscored through various strategies aimed at augmenting BM-lymphopoiesis, including

sex-steroid ablation[50, 51] and administration of Ghrelin,[52] among other approaches.[53] We therefore

hypothesized that robust BM lymphopoiesis driven by Kit^{lo} HSCs could potentially counteract

age-related alterations within the thymic microenvironment, consequently ameliorating the

decline in T-cell output. To this end, we competitively transplanted Kit^{lo} or Kit^{hi} HSCs from

young C57BL/6 mice (CD45.2) with rescue BM cells (CD45.1) into lethally irradiated 14-

month-old (Middle-aged) BALB/cJ recipients (**Figure 2P**). Sixteen weeks post-transplantation,

we observed in Kit^{lo} vs Kit^{hi} recipients significantly higher numbers of peripheral T– and B-cells

(**Figures 2Q, S3I, and S3J**). This was attributed to increased numbers of Kit^{lo} HSC-derived

lymphoid progenitors (including LMPP/MPP4 and CLPs, **Figures S3K and S3L**), while

numbers of myeloid precursors were comparable (MPP3, CMP, GMP, and MEP, **Figures S3K**

and S3M). In our parallel analysis of the thymi of Kit^{lo} vs Kit^{hi} recipients, we observed increased

numbers of thymocytes and TECs (**Figures 2R-2T and S3N**). Collectively, these findings

establish the potential of Kit^{lo} HSCs for lymphopoiesis and to improve thymic recovery and

peripheral T cell reconstitution in recipients of various ages.

Kit^{lo} HSCs exhibit distinct epigenetic features that facilitate lymphoid potential

Prior work has shown that immunophenotypically defined HSC clones with differential

lymphoid output have distinct epigenetic signatures, with no discernable patterns in terms of

gene expression.[18, 24, 54] To identify the gene regulatory network orchestrating the lymphoid

potential differences in HSCs, we analyzed our single-cell ATAC-seq (scATAC-seq) component

from our multiome dataset (**Figure S5D**). Through unsupervised clustering, we identified 6

clusters, including young and old HSCs (**Figures 3A and 3B**). Distinct HSC subsets identified

via scRNA-seq analyses (**Figure 1B**) are situated in varied regions on the scATAC-seq UMAP

(**Figures 3C and 3D**). Notably, MLin-HSC and Kit[lo] HSC intersect with Cluster 3 from the

scATAC-seq clusters (**Figure 3E**).

We characterized the differentially accessible regions (DARs) of chromatin that defined the HSC

subsets and found 2816 DARs in cluster C3 (MLin-HSC and Kit[lo] enriched) cluster compared to

cluster C6 (q-HSC and Kit[hi] enriched) (**Figure S5F**). Next to determine the differential activity

and accessibility of lymphoid-specifying transcription factors (TFs) in the identified Kit[hi] and

Kit[lo] HSC subsets, we analyzed chromatin accessibility variation within TF motifs using

chromVAR and gene accessibility. Compared to Kit[hi] HSCs, we found increased gene

accessibility and enrichment for Zinc finger motifs, specifically *Zbtb1 and Zbtb7a,* in Kit[lo]

DARs, which are implicated in lymphoid differentiation program[18, 55, 56, 57] (**Figures 3E-3F**).

In summary, we found that Kit[lo] HSCs exhibit an epigenetic program, emphasizing an

enrichment of lymphoid-specifying transcription factors, in support of their heightened lymphoid

potential.

Zbtb1 **regulates T-cell potential in phenotypic HSCs**

Among members of the ZBTB gene family, *Zbtb1* plays a pivotal role in T-cell development[57, 58,]

[59] and lymphoid reconstitution.[57] However, its role in regulating lymphoid potential in HSCs

remains unknown. We hypothesized that Zbtb1 loss impairs the lymphoid potential of Kit[lo]

HSCs. To this end, we used the CRISPR-Cas9 system to generate *Zbtb1*-deficient Kit[lo] HSCs

(**Figure 4A**) and confirmed decrease in Zbtb1 by flow-cytometric analysis (**Figure S6A**). *Zbtb1*-

deficient Kit[lo] HSCs showed significantly reduced *in vitro* T-cell potential (**Figure 4B**) and

impaired lymphoid reconstitution *in vivo* (**Figure 4C**). Next, we questioned whether reinstating

Zbtb1 expression in megakaryocytic-biased, Kit^{hi} HSCs is sufficient to rescue their lymphoid

defect. To test our hypothesis, we overexpressed *Zbtb1* cDNA in Kit^{hi} HSCs (**Figure 4D and**

S6B) and we observed a rescue of their T-cell potential when compared to Kit^{hi} controls, thus

partially phenocopying Kit^{lo} HSCs (**Figures 4E**) as well as improved lymphoid reconstitution *in*

vivo (**Figure 4F**).

Previous studies have identified Zbtb1 targets which facilitate Notch-mediated T-cell

differentiation in lymphoid progenitors[60] and support chromatin remodeling following DNA

damage.[61] Given *Zbtb1* motif enrichment and enhanced T-cell differentiation in Kit^{lo} HSCs, we

hypothesized that Zbtb1 targets mediating lymphoid differentiation gene programs are also

enriched in Kit^{lo} HSCs. To this end, we queried Kit^{lo} enriched genes for Notch1-interacting

proteins.[62] We observed ~50% overlap of Notch1-partners enriched in Kit^{lo} HSCs, of which 60%

were Zbtb1-targets identified in a previous CHIP-seq study[60] (**Figure 4G**). Furthermore, we

identified an additional subset of Zbtb1 targets (33%) that were not associated with the Notch1

interactome (**Data Table 3**). To further query for non-Notch related biological pathways, we

performed pathway analysis and identified a strong enrichment for gene sets related to chromatin

remodeling and RNA stability (**Figure 4H, Data Table 4**).

These data demonstrate that Zbtb1 contributes to an epigenetic regulatory program that facilitates

T-cell differentiation in primitive Kit^{lo} HSCs likely through instructive and permissive

mechanisms.

Old Kit^{lo} HSCs exhibit enhanced T-cell potential

To determine whether Kit^{lo} HSCs retain their lymphoid potential with age, we evaluated the

lymphoid potential of old Kit subsets. Remarkably, even in the case of old mice, we noted a

substantial increase in lymphoid progenitor and ATO output from old Kit^{lo} HSCs compared to

Kit^{hi} HSCs (**Figures 5A-5B**), although modest when compared to the young HSCs. This finding

is consistent with prior reports of proliferative defects in aged T-cell precursors contributing to

decreased thymopoiesis. [3, 4, 5]

Next, to test their *in vivo* potential, we competitively transplanted either Kit^{lo} or Kit^{hi} HSCs from

old C57BL/6 (CD45.2) mice with rescue BM (CD45.1) cells into lethally irradiated young

BALB/cJ recipients (**Figure 5C**). At eight weeks post-transplantation, we noted significantly

higher numbers of T cells in the peripheral blood of Kit^{lo} vs Kit^{hi} recipients. (**Figures 5D and**

S7A). Consistent with our experiments using HSCs from young donors, we observed in the BM

of Kit^{lo} vs Kit^{hi} recipients significantly higher numbers of LT HSCs, LMPPs and CLPs (**Figures**

5E-5F) with comparable myeloid potential (**Figures 5E and S7B**). Furthermore, mirroring our

findings in HSCs from young mice, we observed in recipients of old Kit^{lo} HSCs enhanced

overall thymic cellularity, thymocyte reconstitution, and recovery of TECs and endothelial cells

(**Figures 5G-5I and S7C-S7D**), increased CD4+ and CD8+ T cells in the spleen (**Figures S7E**

and S7F), and higher RTE output (**Figures 5J-5K**). At 20-weeks post-HCT, we observed similar

trends of enhanced reconstitution in old Kit^{lo} HSC recipients, confirming their long-term

reconstituting potential (**Figure S7H-S7O**). Next to examine whether old HSC subsets with

preserved lymphoid could augment aged thymic recovery, we competitively transplanted middle-

aged recipients with old Kit^{lo} or Kit^{hi} donors (**Figure 5L**). We observed increased CLPs in the

BM and peripheral T cells, but a modest, non-significant increase in thymic cellularity in Kit^{lo}

recipients (**Figures 5M-5O, and S7G**). Similar to our *Zbtb1* functional studies with young Kit^{hi}

HSCs, we observed a rescue in T-cell potential in *Zbtb1*-overexpressed old Kit^{hi} HSCs (**Figure**

5P).

Finally, to directly compare the per cell functionality of Kit^{lo} HSCs in the context of aging, we

generated mixed chimeras combining Kit^{lo} HSCs cells from both young and old mice in an equal

competition transplant (**Figure S8A**). Consistent with our *in vitro* findings (**Figures 5A-5B**), we

found an overall decrease in lymphoid reconstitution within the PB, BM, and thymus with old

Kit^{lo} HSCs when compared to their young counterparts (**Figures S8B-S8G**). Pathway analysis

revealed decreased enrichment of chromatin remodeling related gene sets (**Figure S8H, Data**

Table 5). Additionally, we noted decreased accessibility and expression of *Zbtb1 (***Figures S8I-**

S8K*)* in Old Kit^{lo} HSCs, indicating a impaired epigenetic program for lymphoid differentiation.

These findings demonstrate that old Kit^{lo} HSCs exhibit preserved lymphoid potential and

contribute to thymic recovery, albeit with reduced output.

KIT^{lo} human BM HSCs exhibit enhanced lymphoid potential

Informed by a recent study that mapped the molecular profiles of human thymic seeding

progenitors (TSPs) to their transcriptional counterparts within the HSC compartment,[63] we next

sought to identify a comparable human HSC subset with lymphoid potential. Interrogating a

CITE-Seq dataset[64] generated from young and old human bone marrow (BM) samples (**Figures**

6A-6C, S9A), we mapped our mouse Kit^{lo} gene signature onto human HSCs and noted a

significant enrichment in young BM (**Figures 6D and 6E**). We observed lower protein

expression of KIT (CD117) on young HSCs by using antibody-derived tag (ADT) reads (**Figure**

S9B). To validate our in-silico findings, we next assessed KIT expression in human BM samples

(young: 20-40 years; middle-age/ old: >40 years). In our FACS analysis, we found that

phenotypic HSCs (pHSCs: CD34+CD38-CD10-CD45RA-CD90+) showed the lowest KIT

expression (**Figure S9C**), with higher KIT levels on old pHSCs (**Figure 6F**). Moreover, we

noted a statistically significant association between age and the composition of KIT subsets in

pHSCs, indicating an increase in KIT[hi] and a decrease in KIT[lo] HSCs, aligning with our

observations in mice. (**Figure 6G**).

Cord blood (CB) and BM have previously been fractionated based on KIT levels.[65, 66] To

evaluate the T-cell potential of these human BM HSC subsets, we generated artificial human

thymic organoids (huATOs) using purified KIT[hi] or KIT[lo] HSCs from human BM samples

(**Figures S9D and 6H**; **Data Table 6**). In concordance with our mouse *in vitro* studies,

following 8-10 weeks of culture, we observed significantly higher output of precursor and

mature T-cells from KIT[lo] HSCs- derived ATOs compared to KIT[hi] HSCs (**Figure 6I**). To further

evaluate the multilineage potential of KIT HSC subsets, we performed a lympho-myeloid

differentiation assay, as previously described.[67] While we observed comparable myeloid

differentiation potential, Kit[lo] HSCs exhibited significantly more B cell and a non-significant

increase in NK cell differentiation potential (**Figure 6J**).

Collectively, these results identify a previously uncharacterized subset of human BM HSCs,

KIT[lo] HSCs – which exhibit enhanced lymphoid potential.

<u>DISCUSSION</u>

Recent studies have uncovered significant heterogeneity within the HSC compartment and their

role in hematopoiesis. This heterogeneity has implications for disease development and

treatment. Of note, certain aspects of HSC heterogeneity challenge the established hematopoietic

models, with lineage biases and lineage-restricted cells impacting self-renewal dynamics.[68, 69]

Decisions regarding lympho-myeloid fate are established at multiple levels of the hematopoietic

hierarchy: HSCs,[21, 22, 70] MPP, LMPP, and more mature GMPs and CLPs. At the level of HSCs,

the process of epigenetic imprinting confers a distinct advantage for lymphoid priming within

multilineage HSCs, as well as maintaining these cell autonomous features imposed at steady

state, even under conditions of stress.[18, 24]

Here, we report on the enhanced T-cell potential of Kit^{lo} HSCs, observed in both mice and

humans. Through combining multiomic single-cell sequencing and functional analyses utilizing

a preclinical allo-HCT model, we demonstrate augmented potential of Kit^{lo} HSCs for T-cell

lymphopoiesis and to support thymic recovery, independent of age. During the early post-

transplant period, we noted a relative increase in DN3 thymocytes relative to other thymocyte

precursor subsets. This observation is consistent with previous studies reporting that DN3

thymocytes undergo compensatory expansion to support early post-transplant thymopoiesis, with

minimal contribution from T-cell precursor thymocytes (ETP and DN2).[10, 71, 72] Further

investigation into mechanisms driving these compensatory processes and its implications for

immune function is critical to our understanding of post-transplant immune reconstitution.

Mechanistically, Kit^{lo} HSCs exhibit differential expression and activity of lymphoid-specifying

transcription factors (TFs), including *Zbtb1*, indicating their epigenetic program towards a

lymphoid fate. Studies in mouse models have shown that disruptions in *Zbtb1* result in T-cell

developmental defects and severe combined immunodeficiency phenotypes,[57, 58] along with

competitive disadvantages in B- and NK-cell compartments. [57] In our study, we identify *Zbtb1* as

an essential transcription factor governing lymphoid specification in phenotypic HSCs and

facilitating T-cell differentiation by reprogramming lineage biased Kit^{hi} HSCs. Additionally, we

observed functional differences within Kit^{lo} HSCs in the context of aging, specifically a

competitive disadvantage in Old Kit^{lo} HSCs*,* potentially attributable to their differences in

expression of the epigenetic repressor of multipotency, *Ezh1*,[73] and *Zbtb1*. Consistent with prior

research, our study suggests *Zbtb1* modulates Notch-dependent[60] and cellular survival

programs.[29, 74] However, further mechanistic studies are needed to elucidate the precise

transcriptional mechanisms underlying this regulation. While the definitive role of *Zbtb1* in

HSCs is not established, it may not have a direct role at the HSC stage; instead, its expression

could be indicative of epigenetic states that facilitate or permit Zbtb1 expression and function at

later stages of development, in lymphocyte progenitors. Future work will also address if ZBTB1

could be used to augment generation of mature lymphoid lineages from human BM precursors.

Long-term reconstituting CD34+KIT^{lo} human BM cells have been previously reported.[66] In

contrast, fractionating phenotypic HSCs in Cord Blood (CB) based on KIT levels showed

reduced reconstitution of KIT^{int} HSCs compared to KIT^{hi} HSCs.[65] These findings suggest that

differences in Kit signaling have functional relevance depending on the developmental stage. In

our study, combining transcriptional, immunophenotypic, and functional analyses of human BM

samples we identified a KIT^{lo} subset within immunophenotypically defined HSCs with enhanced

lymphoid potential, highlighting its translational relevance in human biology. Further functional

characterization is warranted to definitively establish the clinical significance of these

observations. This understanding is crucial as prospective HSC isolation with subsequent ex vivo

expansion,[75, 76] holds promise for enhancing immune regeneration after bone marrow

transplantation, effectively counteracting treatment-related immunosuppression and age-

associated thymic decline.

ACKNOWLEDGEMENTS

We thank Dr. Andrea Schietinger for the gift of mice; Ronan Chaligné, Kalina Belcheva, and

Anthony Michaels for help; Dr. Gay Crooks, and Julia Gerscheimer (UCLA) for sharing the

mATO and huATO systems and assistance with troubleshooting; Kenneth Dorshkind (UCLA)

for providing S17 cells; Parashar Dhopala and Göran Karlsson (Lund University) for providing

the human BM CITE-seq raw matrices.[64]

FUNDING SOURCES

This research was supported by National Cancer Institute award numbers, R35-CA284024, P01-

CA023766, R01-CA228308, and P30 CA008748 MSK Cancer Center Support Grant/Core

Grant; National Heart, Lung, and Blood Institute (NHLBI) award number R01-HL164902;

National Institute on Aging award number P01-AG052359, and Tri Institutional Stem Cell

Initiative. Additional funding was received from The Lymphoma Foundation, The Susan and

Peter Solomon Family Fund, The Solomon Microbiome Nutrition and Cancer Program, Cycle

for Survival, Parker Institute for Cancer Immunotherapy, Paula and Rodger Riney Multiple

Myeloma Research Initiative, Starr Cancer Consortium, and Seres Therapeutics. HKE receives

research support from NHLBI. SM receives research support from Marie-Josée Kravis

fellowship in quantitative biology. SG receives research support from CRI / Donald J. Gogel

Postdoctoral Fellowship (CRI #3934). SDW reports research funding from the MSK Leukemia

SPORE Career Enhancement Program (NIH/NCI P50 CA254838-01), the MSK Gerstner

Physician Scholar Program, and the Parker Institute for Cancer Immunotherapy. MS receives

research support from the following funding sources: Burroughs Wellcome Fund (PDEP),

Damon Runyon Cancer Research Foundation (Clinical Investigator Award), V Foundation (V

Scholar), Emerson Collective, and the National Institutes of Health (NHLBI K08 HL156082-

01A1). KA receives research support from the Gerstner Physician Scholar program.

AUTHOR CONTRIBUTIONS

HKE, MRMvdB conceived and designed the study. HKE, MRMvdB acquired funding. HKE, MBdS, AR, BG, NL, MAEA, SG planned research activities. HKE curated data and performed initial analysis. AIK and SM performed initial computational analysis of single-cell RNA- seq data, and single-cell ATAC- seq and human CITE-seq data, respectively. RS provided code and assisted HKE with downstream data/sequence computation and statistical analyses. HKE analyzed flow cytometric data. HKE, MBdS, AR, BG, NL, MAEA, XZ, HA, SG, MS, SDW, KV, CYP, CED, and JCS planned and performed experiments. CSL, AB, MRMvdB supervised study. HKE, SM, and MBdS prepared figures and visualized data. HKE, MRMvdB wrote the initial draft. All authors reviewed and approved the manuscript.

DECLARATION OF INTERESTS

H.K.Elias, and M.R.M. van den Brink are co-inventors on patent applications related to this work. MRMvdB has received research support and stock options from Seres Therapeutics and stock options from Notch Therapeutics and Pluto Therapeutics; he has received royalties from Wolters Kluwer; has consulted, received honorarium from or participated in advisory boards for Seres Therapeutics, WindMIL Therapeutics, Rheos Medicines, Merck & Co, Inc., Magenta Therapeutics, Frazier Healthcare Partners, Nektar Therapeutics, Notch Therapeutics, Forty Seven Inc., Priothera, Ceramedix, Lygenesis, Pluto Therapeutics, GlaskoSmithKline, Da Volterra, Vor BioPharma, Novartis (Spouse), Synthekine (Spouse), and Beigene (Spouse); he has IP Licensing with Seres Therapeutics and Juno Therapeutics; and holds a fiduciary role on the Foundation Board of DKMS (a nonprofit organization). The Walter and Eliza Hall Institute of Medical

Research receives milestone and royalty payments related to venetoclax. Employees are entitled

to receive benefits related to these payments.

REFERENCES

1. Grinenko T*, et al.* Clonal expansion capacity defines two consecutive developmental stages of long-term hematopoietic stem cells. *J Exp Med* **211**, 209-215 (2014).

2. Shin JY, Hu W, Naramura M, Park CY. High c-Kit expression identifies hematopoietic stem cells with impaired self-renewal and megakaryocytic bias. *J Exp Med* **211**, 217-231 (2014).

3. Zediak VP, Maillard I, Bhandoola A. Multiple prethymic defects underlie age-related loss of T progenitor competence. *Blood* **110**, 1161-1167 (2007).

4. Min H, Montecino-Rodriguez E, Dorshkind K. Reduction in the developmental potential of intrathymic T cell progenitors with age. *J Immunol* **173**, 245-250 (2004).

5. Srinivasan J*, et al.* The initial age-associated decline in early T-cell progenitors reflects fewer pre-thymic progenitors and altered signals in the bone marrow and thymus microenvironments. *Aging Cell* **22**, e13870 (2023).

6. Hale JS, Boursalian TE, Turk GL, Fink PJ. Thymic output in aged mice. *Proc Natl Acad Sci U S A* **103**, 8447-8452 (2006).

7. van den Brink M, Uhrberg M, Jahn L, DiPersio JF, Pulsipher MA. Selected biological issues affecting relapse after stem cell transplantation: role of T-cell impairment, NK cells and intrinsic tumor resistance. *Bone Marrow Transplant* **53**, 949-959 (2018).

8. de Koning C*, et al.* CD4+ T-cell reconstitution predicts survival outcomes after acute graft-versus-host-disease: a dual-center validation. *Blood* **137**, 848-855 (2021).

9. van Roessel I*, et al.* Early CD4+ T cell reconstitution as predictor of outcomes after allogeneic hematopoietic cell transplantation. *Cytotherapy* **22**, 503-510 (2020).

10. Zlotoff DA*, et al.* Delivery of progenitors to the thymus limits T-lineage reconstitution after bone marrow transplantation. *Blood* **118**, 1962-1970 (2011).

11. van Ewijk W, Shores EW, Singer A. Crosstalk in the mouse thymus. *Immunol Today* **15**, 214-217 (1994).

499 12.   Poliani PL*, et al.* Early defects in human T-cell development severely affect distribution
500   and maturation of thymic stromal cells: possible implications for the pathophysiology of
501   Omenn syndrome. *Blood* **114**, 105-108 (2009).
502
503 13.   Pala F*, et al.* Profound abnormalities in thymic epithelial cells in Rag1 hypomorphic
504   mice: implications for immune reconstitution after stem cell transplantation. *The Journal*
505   *of Immunology* **208**, 167.102-167.102 (2022).
506
507 14.   Elias HK, Bryder D, Park CY. Molecular mechanisms underlying lineage bias in aging
508   hematopoiesis. *Semin Hematol* **54**, 4-11 (2017).
509
510 15.   Rodriguez-Fraticelli AE*, et al.* Single-cell lineage tracing unveils a role for TCF15 in
511   haematopoiesis. *Nature* **583**, 585-589 (2020).
512
513 16.   Dong F*, et al.* Differentiation of transplanted haematopoietic stem cells tracked by single-
514   cell transcriptomic analysis. *Nat Cell Biol* **22**, 630-639 (2020).
515
516 17.   Carrelha J*, et al.* Hierarchically related lineage-restricted fates of multipotent
517   haematopoietic stem cells. *Nature* **554**, 106-111 (2018).
518
519 18.   Meng Y*, et al.* Epigenetic programming defines haematopoietic stem cell fate restriction.
520   *Nat Cell Biol* **25**, 812-822 (2023).
521
522 19.   Morita Y, Ema H, Nakauchi H. Heterogeneity and hierarchy within the most primitive
523   hematopoietic stem cell compartment. *J Exp Med* **207**, 1173-1182 (2010).
524
525 20.   Yamamoto R*, et al.* Large-Scale Clonal Analysis Resolves Aging of the Mouse
526   Hematopoietic Stem Cell Compartment. *Cell Stem Cell* **22**, 600-607 e604 (2018).
527
528 21.   Benz C*, et al.* Hematopoietic stem cell subtypes expand differentially during
529   development and display distinct lymphopoietic programs. *Cell Stem Cell* **10**, 273-283
530   (2012).
531
532 22.   Challen GA, Boles NC, Chambers SM, Goodell MA. Distinct hematopoietic stem cell
533   subtypes are differentially regulated by TGF-beta1. *Cell Stem Cell* **6**, 265-278 (2010).
534
535 23.   Naik SH*, et al.* Diverse and heritable lineage imprinting of early haematopoietic
536   progenitors. *Nature* **496**, 229-232 (2013).
537
538 24.   Yu VWC*, et al.* Epigenetic Memory Underlies Cell-Autonomous Heterogeneous
539   Behavior of Hematopoietic Stem Cells. *Cell* **167**, 1310-1322 e1317 (2016).
540
541 25.   Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid
542   progenitors in mouse bone marrow. *Cell* **91**, 661-672 (1997).
543

544   26.   Flohr Svendsen A*, et al.* A comprehensive transcriptome signature of murine
545   hematopoietic stem cell aging. *Blood* **138**, 439-451 (2021).
546

547   27.   Grover A*, et al.* Single-cell RNA sequencing reveals molecular and functional platelet
548   bias of aged haematopoietic stem cells. *Nat Commun* **7**, 11075 (2016).
549

550   28.   Herault L, Poplineau M, Mazuel A, Platet N, Remy E, Duprez E. Single-cell RNA-seq
551   reveals a concomitant delay in differentiation and cell cycle of aged hematopoietic stem
552   cells. *BMC Biol* **19**, 19 (2021).
553

554   29.   Kim YW*, et al.* Integration of single-cell transcriptomes and biological function reveals
555   distinct behavioral patterns in bone marrow endothelium. *Nat Commun* **13**, 7235 (2022).
556

557   30.   Konturek-Ciesla A*, et al.* Temporal multimodal single-cell profiling of native
558   hematopoiesis illuminates altered differentiation trajectories with age. *Cell Rep* **42**,
559   112304 (2023).
560

561   31.   Safi F*, et al.* Concurrent stem- and lineage-affiliated chromatin programs precede
562   hematopoietic lineage restriction. *Cell Rep* **39**, 110798 (2022).
563

564   32.   Giladi A*, et al.* Single-cell characterization of haematopoietic progenitors and their
565   trajectories in homeostasis and perturbed haematopoiesis. *Nat Cell Biol* **20**, 836-846
566   (2018).
567

568   33.   Cabezas-Wallscheid N*, et al.* Vitamin A-Retinoic Acid Signaling Regulates
569   Hematopoietic Stem Cell Dormancy. *Cell* **169**, 807-823 e819 (2017).
570

571   34.   Lauridsen FKB*, et al.* Differences in Cell Cycle Status Underlie Transcriptional
572   Heterogeneity in the HSC Compartment. *Cell Rep* **24**, 766-780 (2018).
573

574   35.   Cabezas-Wallscheid N*, et al.* Identification of regulatory networks in HSCs and their
575   immediate progeny via integrated proteome, transcriptome, and DNA methylome
576   analysis. *Cell Stem Cell* **15**, 507-522 (2014).
577

578   36.   Wahlestedt M*, et al.* Critical Modulation of Hematopoietic Lineage Fate by Hepatic
579   Leukemia Factor. *Cell Rep* **21**, 2251-2263 (2017).
580

581   37.   Komorowska K*, et al.* Hepatic Leukemia Factor Maintains Quiescence of Hematopoietic
582   Stem Cells and Protects the Stem Cell Pool during Regeneration. *Cell Rep* **21**, 3514-3523
583   (2017).
584

585   38.   Lehnertz B*, et al.* HLF expression defines the human hematopoietic stem cell state. *Blood*
586   **138**, 2642-2654 (2021).
587

588   39.   Verovskaya E*, et al.* Heterogeneity of young and aged murine hematopoietic stem cells
589   revealed by quantitative clonal analysis using cellular barcoding. *Blood* **122**, 523-532
590   (2013).
591

592   40.   Elias HK*, et al.* Identification of Old HSCs with Preserved Self-Renewal and Long-Term
593   Reconstitution Potential. *Blood* **134**, 2479 (2019).
594

595   41.   Montecino-Rodriguez E, Kong Y, Casero D, Rouault A, Dorshkind K, Pioli PD.
596   Lymphoid-Biased Hematopoietic Stem Cells Are Maintained with Age and Efficiently
597   Generate Lymphoid Progeny. *Stem Cell Reports* **12**, 584-596 (2019).
598

599   42.   Inlay MA*, et al.* Ly6d marks the earliest stage of B-cell specification and identifies the
600   branchpoint between B-cell and T-cell development. *Genes Dev* **23**, 2376-2381 (2009).
601

602   43.   Seet CS*, et al.* Generation of mature T cells from human hematopoietic stem and
603   progenitor cells in artificial thymic organoids. *Nat Methods* **14**, 521-530 (2017).
604

605   44.   Montel-Hagen A*, et al.* In Vitro Recapitulation of Murine Thymopoiesis from Single
606   Hematopoietic Stem Cells. *Cell Rep* **33**, 108320 (2020).
607

608   45.   Zhou W, Gao F, Romero-Wolf M, Jo S, Rothenberg EV. Single-cell deletion analyses
609   show control of pro-T cell developmental speed and pathways by Tcf7, Spi1, Gata3,
610   Bcl11a, Erg, and Bcl11b. *Sci Immunol* **7**, eabm1920 (2022).
611

612   46.   Zlotoff DA, Sambandam A, Logan TD, Bell JJ, Schwarz BA, Bhandoola A. CCR7 and
613   CCR9 together recruit hematopoietic progenitors to the adult thymus. *Blood* **115**, 1897-
614   1905 (2010).
615

616   47.   Dudakov JA*, et al.* Interleukin-22 drives endogenous thymic regeneration in mice.
617   *Science* **336**, 91-95 (2012).
618

619   48.   Lopes N, Vachon H, Marie J, Irla M. Administration of RANKL boosts thymic
620   regeneration upon bone marrow transplantation. *EMBO Mol Med* **9**, 835-851 (2017).
621

622   49.   Yu W*, et al.* Continued RAG expression in late stages of B cell development and no
623   apparent re-induction after immunization. *Nature* **400**, 682-687 (1999).
624

625   50.   Goldberg GL*, et al.* Sex steroid ablation enhances lymphoid recovery following
626   autologous hematopoietic stem cell transplantation. *Transplantation* **80**, 1604-1613
627   (2005).
628

629   51.   Goldberg GL*, et al.* Enhanced immune reconstitution by sex steroid ablation following
630   allogeneic hemopoietic stem cell transplantation. *J Immunol* **178**, 7473-7484 (2007).
631

632   52.   Dixit VD*, et al.* Ghrelin promotes thymopoiesis during aging. *J Clin Invest* **117**, 2778-
633   2790 (2007).

634
635 53. Spadaro O, *et al.* Caloric restriction in humans reveals immunometabolic regulators of
636 health span. *Science* **375**, 671-677 (2022).
637
638 54. Lara-Astiaso D, *et al.* Immunogenetics. Chromatin state dynamics during blood
639 formation. *Science* **345**, 943-949 (2014).
640
641 55. Shin B, *et al.* Runx1 and Runx3 drive progenitor to T-lineage transcriptome conversion
642 in mouse T cell commitment via dynamic genomic site switching. *Proc Natl Acad Sci U*
643 *S A* **118**, (2021).
644
645 56. Maeda T, *et al.* Regulation of B versus T lymphoid lineage fate decision by the proto-
646 oncogene LRF. *Science* **316**, 860-866 (2007).
647
648 57. Siggs OM, Li X, Xia Y, Beutler B. ZBTB1 is a determinant of lymphoid development. *J*
649 *Exp Med* **209**, 19-27 (2012).
650
651 58. Punwani D, *et al.* Transcription factor zinc finger and BTB domain 1 is essential for
652 lymphocyte development. *J Immunol* **189**, 1253-1264 (2012).
653
654 59. Zhang X, Lu Y, Cao X, Zhen T, Kovalovsky D. Zbtb1 prevents default myeloid
655 differentiation of lymphoid-primed multipotent progenitors. *Oncotarget* **7**, 58768-58778
656 (2016).
657
658 60. Koizumi M, *et al.* Transcription factor Zbtb1 interacts with bridging factor Lmo2 and
659 maintains the T-lineage differentiation capacity of lymphoid progenitor cells. *J Biol*
660 *Chem* **298**, 102506 (2022).
661
662 61. Kim H, *et al.* Transcriptional repressor ZBTB1 promotes chromatin remodeling and
663 translesion DNA synthesis. *Mol Cell* **54**, 107-118 (2014).
664
665 62. Yatim A, *et al.* NOTCH1 nuclear interactome reveals key regulators of its transcriptional
666 activity and oncogenic function. *Mol Cell* **48**, 445-458 (2012).
667
668 63. Cordes M, *et al.* Single-cell immune profiling reveals thymus-seeding populations, T cell
669 commitment, and multilineage development in the human thymus. *Sci Immunol* **7**,
670 eade0182 (2022).
671
672 64. Sommarin MNE, *et al.* Single-Cell Multiomics Reveals Distinct Cell States at the Top of
673 Human Hematopoietic Hierarchy. *bioRxiv*, 2021.2004.2001.437998 (2021).
674
675 65. Cosgun KN, *et al.* Kit regulates HSC engraftment across the human-mouse species
676 barrier. *Cell Stem Cell* **15**, 227-238 (2014).
677

66. Kawashima I, Zanjani ED, Almaida-Porada G, Flake AW, Zeng H, Ogawa M. CD34+ human marrow cells that express low levels of Kit protein are enriched for long-term marrow-engrafting cells. *Blood* **87**, 4136-4142 (1996).

67. Karamitros D*, et al.* Single-cell analysis reveals the continuum of human lympho-myeloid progenitor cells. *Nat Immunol* **19**, 85-97 (2018).

68. Notta F*, et al.* Distinct routes of lineage development reshape the human blood hierarchy across ontogeny. *Science* **351**, aab2116 (2016).

69. Velten L*, et al.* Human haematopoietic stem cell lineage commitment is a continuous process. *Nat Cell Biol* **19**, 271-281 (2017).

70. Dykstra B*, et al.* Long-term propagation of distinct hematopoietic differentiation programs in vivo. *Cell Stem Cell* **1**, 218-229 (2007).

71. Paiva RA*, et al.* Self-renewal of double-negative 3 early thymocytes enables thymus autonomy but compromises the beta-selection checkpoint. *Cell Rep* **35**, 108967 (2021).

72. Maillard I*, et al.* Notch-dependent T-lineage commitment occurs at extrathymic sites following bone marrow transplantation. *Blood* **107**, 3511-3519 (2006).

73. Vo LT*, et al.* Regulation of embryonic haematopoietic multipotency by EZH1. *Nature* **553**, 506-510 (2018).

74. Cao X, Lu Y, Zhang X, Kovalovsky D. Zbtb1 Safeguards Genome Integrity and Prevents p53-Mediated Apoptosis in Proliferating Lymphoid Progenitors. *J Immunol* **197**, 1199-1211 (2016).

75. Sakurai M*, et al.* Chemically defined cytokine-free expansion of human haematopoietic stem cells. *Nature* **615**, 127-133 (2023).

76. Wilkinson AC*, et al.* Long-term ex vivo haematopoietic-stem-cell expansion allows nonconditioned transplantation. *Nature* **571**, 117-121 (2019).

77. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-connected communities. *Sci Rep* **9**, 5233 (2019).

78. Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289 (2019).

79. Korsunsky I*, et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat Methods* **16**, 1289-1296 (2019).

80. Kharchenko PV, Silberstein L, Scadden DT. Bayesian approach to single-cell differential expression analysis. *Nature methods* **11**, 740-742 (2014).

725 81. van Dijk D, *et al.* Recovering Gene Interactions from Single-Cell Data Using Data
726 Diffusion. *Cell* **174**, 716-729 e727 (2018).
727

728 82. Finak G*, et al.* MAST: a flexible statistical framework for assessing transcriptional
729 changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome*
730 *Biol* **16**, 278 (2015).
731

732 83. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data
733 analysis. *Genome Biol* **19**, 15 (2018).
734

735 84. Dann E, Henderson NC, Teichmann SA, Morgan MD, Marioni JC. Differential
736 abundance testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol* **40**,
737 245-253 (2022).
738

739 85. Granja JM*, et al.* ArchR is a scalable software package for integrative single-cell
740 chromatin accessibility analysis. *Nat Genet* **53**, 403-411 (2021).
741

742 86. Zhang Y*, et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137
743 (2008).
744

745 87. Schep AN, Wu B, Buenrostro JD, Greenleaf WJ. chromVAR: inferring transcription-
746 factor-associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975-
747 978 (2017).
748

749 88. Hao Y*, et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
750 e3529 (2021).
751

752 89. Cunningham F*, et al.* Ensembl 2022. *Nucleic Acids Res* **50**, D988-D995 (2022).
753

754 90. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
755 using empirical Bayes methods. *Biostatistics* **8**, 118-127 (2007).
756

757 91. Fang Z, Liu X, Peltz G. GSEApy: a comprehensive package for performing gene set
758 enrichment analysis in Python. *Bioinformatics* **39**, (2023).
759

<u>FIGURE LEGENDS</u>

Figure 1: Age-related decline in multilineage HSCs marked by low Kit expression

(**A-H**) Multiome single-cell RNA and ATAC sequencing was performed on HSC (Lineage-Sca-

1+cKit+ CD34-CD48-Flt3-CD150+) cells isolated from 2-mo (young) or 24-mo (old) female

C57BL/6 mice. nTOTAL =12,350 cells. (**A**) Volcano plot of DGE analysis between MLin-HSC

vs. q-HSC subsets in young HSCs, highlighting bonafide HSC markers. (**B**) Violin plot for

imputed *Kit* gene expression by annotated HSC subsets (as described in **Fig. S2C**. (**C**) Frequency

of computationally defined HSC subtypes in Kit^{hi} and Kit^{lo} subsets in young HSCs. (**D-E**)

Combined Uniform manifold approximation and projection (UMAP) of young and old HSCs

after Harmony batch correction annotated by age (**D**) and annotated HSC subsets (q-HSC:

Quiescent HSCs; Mgk-HSCs: Platelet-biased HSCs; Mlin-HSC: Multilineage HSCs; p-HSC:

Proliferative HSCs; Int-HSC: Intermediate HSCs) (**E**). (**F**) Frequency of HSC subtypes in young

and old HSCs. (**G**) Heatmap showing gene expression of bonafide HSC markers ordered by

increasing Kit expression. (**H**) Violin plot for imputed Kit gene expression by age. Statistical

analysis was performed using the Wilcoxon test. (**I**) Scaled change in frequency for each Kit

HSC subset with age. (**J**) FACS analysis showing frequency of Kit^{hi} and Kit^{lo} HSCs by age. All

data are from n=6 mice/group. Error bars represent mean ± SEM. **P<0.01, ***P<0.001. P

values calculated by two-way ANOVA.

Figure 2: Young Kit^{lo} HSCs exhibit enhanced T-cell potential

(**A**) Experimental schema to evaluate *in vitro* lymphoid progenitor and T-cell differentiation

potential of Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 2-mo (young) C57BL/6 mice using the S17

lymphoid assay and murine artificial thymic organoid (mATO) respectively. Enumeration of

absolute number of lymphoid progenitor cells following 14 days in S17 lymphoid assay (B) and

absolute number of T-cell subsets following 8-weeks of culture in mATOs (**C**). Aggregated data

across 4-5 independent experiments. **(D)** Experimental schema for competitive allogeneic HCT

(allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 2-mo (young) C57BL/6 mice with

competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ mice transplanted into

lethally irradiated 7-week-old (young) BALB/cJ recipients. **(E-K)** Eight weeks after competitive

HCT, **(E)** frequency of donor-derived chimerism (CD45.2/ H-2Kb) of mature lineages in the

peripheral blood (PB, CD45+, Myeloid cell: Gr1+CD11b+; B-cell: B220+; T-cell: CD3+), **(F-G)**

Enumeration of absolute number of donor-derived cells in the BM for LSK cells (LT HSC:

Lineage$^-$Sca-1$^+$cKit$^+$ CD34$^-$CD48$^-$Flt3$^-$CD150$^+$ **;** ST HSC: Lineage$^-$Sca-1$^+$cKit$^+$ CD48$^-$Flt3$^-$

CD150$^-$**;** MPP2: Lineage$^-$Sca-1$^+$cKit$^+$ CD48$^+$Flt3$^-$CD150$^-$; MPP3: Lineage$^-$Sca-1$^+$cKit$^+$

CD48$^+$Flt3$^-$CD150$^+$; MPP4/ LMPP: Lineage$^-$Sca-1$^+$cKit$^+$ Flt3$^+$CD150$^-$) **(F),** Common Lymphoid

Progenitors cells (CLP: Lineage$^-$IL7Ra$^+$Flt3$^+$Sca$^{mid/lo}$ Kit^{lo}) **(G). (H-K)** Post-HCT thymi analysis

for total thymic cellularity **(H)**, **(I-J)** enumeration of absolute number of donor-derived cells for

T-cell precursors (ETP: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^+$ CD25$^-$Kit$^+$;DN2: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^+$

CD25$^+$; DN3: Lineage$^-$ CD4$^-$ CD8$^-$CD44$^-$ CD25$^+$) **(I)**, Mature T-cells (DP: Lineage$^-$ CD4$^+$ CD8$^+$;

SP4: Lineage$^-$ CD4$^+$ CD8$^-$; SP8: Lineage$^-$ CD4$^-$ CD8$^+$) **(J),** and analysis of thymic CD45-

compartment for Thymic Epithelial Cells (TEC: CD45$^-$ EpCAM$^+$), Cortical TEC (cTEC: CD45$^-$

EpCAM$^+$UEA-1^{lo} 6C3^{hi} MHCII$^{hi/lo}$), and Medullary TEC (mTEC: CD45$^-$ EpCAM$^+$UEA-1^{hi}

6C3^{lo} MHCII$^{hi/lo-}$) **(K)**. All data are from n=10-11 mice/group. **(L)** Experimental schema to

evaluate thymic function following competitive allo-HCT using young HSCs. **(M)** Frequency of

donor-derived Recent Thymic Emigrants (RTE: CD3+GFP+) in the PB at 8 weeks post-HCT.

All data are from n=5-6 mice/group. **(N)** Experimental schema to investigate functional response

of differential Kit-expressing HSC-derived T-cells in secondary recipients to *L. monocytogenes*

expressing chicken ovalbumin (LM-OVA) infection. **(O)** Spleen analysis of secondary recipients

for frequency of OT1-derived chimerism of CD8+ T cells (OT1+CD8+: $Va2^+Vb5^+H2-K^bCD8^+$)

7 days post-infection. All data are from n=9-10 mice/group. **(P)** Experimental schema for

competitive allogeneic HCT (allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 2-mo

(young) C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-

PtprcaPepcb/BoyJ mice transplanted into lethally irradiated 14-mo (Middle-aged) BALB/cJ

recipients. **(Q)** Frequency of donor-derived T-cell chimerism at the indicated time-points. **(R-T)**

Sixteen weeks after competitive HCT enumeration of total thymic cellularity **(R)**, donor-derived

T-cell precursor thymocyte subsets as defined in Figure 3F **(S)**, donor-derived mature T-cell

thymocyte subsets as defined in Figure 3G **(T).** All data are from n=6-7 mice/group. Error bars

represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. P values calculated by

nonparametric unpaired Mann-Whitney U test. Panels **A**, **D**, **L**, **N**, **and P** were created using

BioRender.

Figure 3: Kit^{lo} HSCs exhibit distinct epigenetic features that facilitate lymphoid potential

(A) UMAP of 6 unsupervised HSC clusters identified by tile-based scATAC-seq analysis after

Harmony batch correction. **(B-D)** Annotated by age **(B)**, HSC subsets **(C),** and Kit subsets **(D)** on

the scATAC-seq UMAP. **(E)** scATAC-seq cluster composition by HSC subsets (**left**) and Kit

subsets (**right**). (**F-G**) Matrix plots showing motif enrichment identified by ChromVAR (**F**) and

gene accessibility (**G**) for lymphoid-specifying transcription factors for Kit subsets in young and

old HSCs.

Figure 4: *Zbtb1* regulates T-cell potential in phenotypic HSCs

(A) Experimental schema for T-cell differentiation assay and competitive allogeneic HCT with

Zbtb1-deficient Kit^{lo} HSCs generated with 2-mo young Rosa26Cas9-eGFP KI mice. **(B)**

Frequency of T-cell subsets following 6-weeks of culture in mATOs. Aggregated data across 3

independent experiments. (C) Frequency of donor-derived lineage chimerism of control or *Zbtb1*

KO Kit^{lo} HSCs in the PB at 6 weeks following competitive HCT. All data are from n=5-6

mice/group. (D) Experimental schema for T-cell differentiation assay and competitive allogeneic

HCT with *Zbtb1*-OE Kit^{hi} HSCs from young mice. (E) Frequency of T-cell subsets following 6-

weeks of culture in mATO of young Kit^{hi} HSCs. Aggregated data across 3 independent

experiments. (F) Frequency of donor-derived lineage chimerism of control or *Zbtb1* OE young

Kit^{hi} HSCs in the PB at 6-weeks following competitive HCT. All data are from n=5-6

mice/group. (G) Previously identified Zbtb1-targets[60] and Notch1-interacting proteins[62] enriched

in Kit^{lo} HSCs. (H) Pathway enrichment analysis for non-Notch1 Zbtb1-targets enriched in Kit^{lo}

HSCs by *gseapy*()[91] using GO_Biological_Process_2023 libraries. Bubbleplot showing

representative pathways. Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001. P

values calculated by nonparametric unpaired Mann-Whitney U test. Panels **A** and **D** were created

using BioRender.

Figure 5: Old Kit^{lo} HSCs exhibit enhanced T-cell potential

(A-B) Evaluation of *in vitro* lymphoid progenitor and T-cell differentiation potential of Kit^{hi}

(red) and Kit^{lo} HSCs (blue) from or 24-mo (old) C57BL/6 mice using the S17 lymphoid assay

and murine artificial thymic organoid (mATO) respectively, as shown in Figure 2A.

Enumeration of absolute number of lymphoid progenitor cells following 14 days in S17

lymphoid assay **(A)** and absolute number of T-cell subsets following 8-weeks of culture in

mATOs **(B).** Aggregated data across 4-5 independent experiments. (**C**) Experimental schema for

competitive allogeneic HCT (allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 22–24-mo

(old) C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ

mice transplanted into lethally irradiated young BALB/cJ recipients. **(D-I)** Eight weeks after

competitive HCT enumeration of absolute number of donor-derived T-cells in the PB **(D)**, LSK

cell subsets as defined in Figure 3C (**E**), CLP cells as defined in Figure 3D (**F**), total thymic

cellularity **(G)**, donor-derived T-cell precursor thymocyte subsets as defined in Figure 3F **(H)**,

TEC compartment as defined in Figure 3H **(I).** All data are from n=9-10 mice/group. (**J**)

Experimental schema to evaluate thymic function following competitive allo-HCT using old

HSCs. **(K)** Frequency of donor-derived Recent Thymic Emigrants (RTE: CD3^{+}GFP^{+}) in the PB

at 8 weeks post-HCT. All data are from n=5 mice/group. (**L**) Experimental schema for

competitive allogeneic HCT (allo-HCT) using Kit^{hi} (red) and Kit^{lo} HSCs (blue) from 22-24-mo

(old) C57BL/6 mice with competitor bone marrow (BM) cells from B6.SJL-PtprcaPepcb/BoyJ

mice transplanted into lethally irradiated middle-aged BALB/cJ recipients. **(M-O)** Eight weeks

after competitive HCT enumeration of absolute number of donor-derived T cells in PB **(M)**,

donor-derived CLPs in the BM as defined in Figure 3D **(N)**, and thymic cellularity **(O)**. All data

are from n=5-6 mice/group. (**P**) Frequency of T-cell subsets following 6-weeks of culture in

mATO of old Kit^{hi} HSCs following *Zbtb1* OE. Aggregated data across 3 independent

experiments. Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001. P values calculated by nonparametric unpaired Mann-Whitney U test. Panels **C,**

J, and **L** were created using BioRender.

Figure 6 KIT^{lo} human BM HSCs exhibit enhanced lymphoid potential

(A-E) Young and old human BM CITE-seq dataset generated and published by Sommarin et al.[64] **(A-C)** UMAP of CD34+ young and old human BM annotated with HSC subsets **(A)**, and age **(B)**. **(C)** HSC cluster composition by age. **(D)** Kit^{lo} gene signature (top 50 marker genes from our mouse data converted to human orthologues) was overlaid on human HSC UMAP. **(E)** Violin plot for Kit^{lo} gene score by age. Statistical analysis was performed using the Wilcoxon test. **(F-G)** FACS analysis of human BM samples. **(F)** Violin plot of KIT protein expression on phenotypic HSCs (pHSC: CD34+CD38-CD10-CD45RA-CD90+) by age (young: 20-40yrs; MA/old: >40yrs). **(G)** Scatterplot showing distribution of KIT^{hi} and KIT^{lo} HSC subsets within pHSCs in human BM samples with varying age. **(H)** Experimental schema to evaluate *in vitro* T-cell and lympho-myeloid potential of human BM HSC subsets based on differential KIT expression using human artificial thymic organoids (hATO) and MS5 assay respectively. **(I)** Following 8-10-weeks of culture, enumeration of CD19+ B-cells and T-cell subsets within CD34+ T-cell precursors (Early Thymic Progenitors, ETP: CD34^{+}CD1a^{-}CD7^{-} ; CD1a^{neg}- ProT: CD34^{+}CD1a^{-}CD7^{+}; CD1a^{pos}-ProT: CD34^{+}CD1a^{+}CD7^{+}), and mature T-cells (DP: CD34^{-}CD5^{+}CD7^{+}CD4^{+}CD8^{+}; SP4: CD34^{-}CD5^{+}CD7^{+}CD4^{+}CD8^{-} ; SP8: CD34^{-}CD5^{+}CD7^{+}CD4^{-}CD8^{+}). Aggregated data represent mean from 8 independent BM donors ± SEM. **(J)** Following 4-weeks of culture, enumeration of Lineage output (Neutrophil-Granulocytes: hCD45+CD15+; monocyte-macrophages: hCD45+CD14+; B-cells: hCD45+CD19+; NK-cells: hCD45+CD56+) after culturing 200 KIT^{lo} and KIT^{hi} HSC cells on MS-5 stroma, as previously described[67]. Aggregated data represent mean from 5 independent BM donors ± SEM. **(L)** Model illustrating age-related loss of lymphoid potential is attributed to a decreased in the frequency of Kit^{lo} HSCs in aged BM. Error bars represent mean ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. P values for panel **G** was based on generalized estimating equation logistic

902 model adjusted for age, KIT^{lo} and KIT^{hi} groups and their interaction, the matched data structure

903 captured by an exchangeable working correlation structure. P values calculated either by

904 nonparametric paired Mann-Whitney U test (Panels **I** and **J**), unpaired Mann-Whitney U test

905 (Panel **F**). Panels **H** and **L** were created using BioRender.

Young Donor/ Young Recipients Tx

Young Donor/ MA Recipients Tx

A Young Kit^lo HSC *Zbtb1* KO

B

C

D Young Kit^hi HSC *Zbtb1* OE

E

F

G Zbtb1 targets enriched in Kit^lo HSCs

Notch1 partners: Zbtb1 dependent
Notch1 partners: Zbtb1 independent
Non-Notch1 Zbtb1 targets

H Non-Notch Zbtb1 targets

GO: Biological Process

Old Donor/ Young Recipients Tx

Old Donor/ MA Recipients Tx

HSC Annotations

A Young
B Leiden Clusters
C HSC Subsets
D
E
F Post-ingest HSC Subsets (Old)
G Post Harmony Sample (Young & Old)
H Old Young
I MiloR analysis
J Old Young
K
L Gated on: Lin-

Young Donor/ Young Recipients Tx

A

B

Old Donor/ Young Recipients Tx

A PB chimerism

B MP compartment

C Donor-derived

D

Old Donor/ MA Recipients Tx

E CD4 T cells

F CD8 T cells

G

Old 20-week Post HCT

H PB chimerism

I LSK

J MP

K CLP

L

M Mature Thymocytes

N Thymocyte Precursors

O Stromal analysis

Equal Competition Tx

A Allogeneic HCT

Kit^lo HSC Donors
Young CD45.1/CD45.2 + Old B6 CD45.2

Young CD45.1 — Rescue BM

Host — Young BALB/cJ

B PB chimerism

C LSK

D MP

E CLP

F Mature Thymocytes

G Thymocyte Precursors

H Pathways Enriched in Young Kit^lo HSCs

I chromVAR Motif Enrichment

J Gene Accessibility

K Gene Expression

A

B

ADT−CD117

C

Gated on: Lin⁻CD34⁺CD38⁻

Gated on: Lin⁻CD34⁺CD38⁺

D   **Gating Strategy to Fractionate Kit Subsets in Human BM**