Antarctic geothermal soils exhibit an absence of regional habitat generalist microorganisms

Stephen E. Noell,[a,b] Jaber Abbaszadeh,[a,b] Huw Richards,[a,b] Marie Labat Saint Vincent,[a,b,c] Charles K. Lee,[a,b] Craig W. Herbold,[d] Matthew B. Stott,[d] S. Craig Cary[a,b], Ian R. McDonald[a,b]*

[a] Thermophile Research Unit, Te Aka Mātuatua | School of Science, Te Whare Wānanga o Waikato | University of Waikato, Hamilton, Aotearoa-New Zealand

[b] International Centre for Terrestrial Antarctic Research, Te Whare Wānanga o Waikato | University of Waikato, Hamilton, Aotearoa-New Zealand

[c] University de Lille, Lille, France

[d] Te Kura Pūtaiao Koiora | School of Biological Sciences, Te Whare Wānanga o Waitaha | University of Canterbury, Christchurch, Aotearoa-New Zealand

Running title: Antarctic Geothermal Microorganisms

* Address correspondence to Ian R. McDonald:

Te Aka Mātuatua - School of Science, Te Whare Wānanga o Waikato - University of Waikato, Hamilton, New Zealand

Email: ian.mcdonald@waikato.ac.nz

Abstract

Active geothermal systems are relatively rare in Antarctica and represent

metaphorical islands ideal to study microbial dispersal. In this study, we tested the

macroecological concept that high dispersal rates result in communities being dominated by

either habitat generalists or specialists by investigating the microbial communities on four

geographically separated geothermal sites on three Antarctic volcanoes (Mts. Erebus,

Melbourne and Rittman). We found that the microbial communities at higher temperature

(max 65□) sites (Tramway Ridge on Erebus and Rittmann) were unique from each other and

were dominated by a variety of novel *Archaea* from class *Nitrososphaeria*, while lower

temperature (max 50□) sites (Western Crater on Erebus and Melbourne) had

characteristically mesophilic communities (*Planctomycetes, Acidobacteriota,* etc) that were

highly similar. We found that 97% of the detected microbial taxa were regional habitat

specialists, with no generalists, with community assembly driven by high dispersal rates and

drift (25 and 30% of community assembly, respectively), not environmental selection. Our

results indicate that for microbial communities experiencing high dispersal rates between

isolated communities, habitat specialists may tend to out-compete habitat generalists.

Introduction

In ecology, the niche concept is commonly used to group organisms based on the

range of different habitats they live in. Habitat generalists (which tend to be metabolic

generalists (Barberán *et al.*, 2014; Bell and Bell, 2021; Chen *et al.*, 2021a; Malard and

Guisan, 2023)) are adapted to thrive in a broader range of different ecosystems compared to

habitat specialists, although some organisms may be specialized to a habitat in one area but

have a broader distribution beyond the scope of the current study (Malard and Guisan, 2023).

Habitat generalists that have a global distribution are usually called "cosmopolitan" to

distinguish them from other habitat generalists (Herbold, Lee, *et al.*, 2014). In most

environments, the diversity of habitat specialists is greater than generalists, but habitat

generalists tend to be the most abundant organisms; this is true both for microorganisms

(Barberán *et al.*, 2014; Zhang *et al.*, 2018; Chen *et al.*, 2021b; Gong *et al.*, 2022) and

macroorganisms (Brown, 1984; Magurran and Henderson, 2003). Exceptions to this trend are

few, but in macroecology, high rates of dispersal between communities can lead to habitat

specialists outcompeting habitat generalists, although the opposite has also been observed

(Brown and Pavlovic, 1992). In contrast, at least two reported exceptions to this trend (habitat

generalists being the most abundant) in microbial ecology have been found under situations

of severe environmental selection and/or dispersal limitation: strong salinity gradients

(Logares *et al.*, 2013) and the mammalian gut (Green *et al.*, 2016). Other environmental

factors are well known to be highly important in structuring microbial communities (e.g., pH

(Fierer and Jackson, 2006; Rousk *et al.*, 2010), salinity (Jiang *et al.*, 2022), water availability

(Niederberger *et al.*, 2019)), raising the possibility that environmental factors may be the

most important factor determining the proportion of generalists and specialists in

environments. In microbial ecology, it is unknown how high dispersal rates would influence

the direction of community dynamics: would it result in habitat specialists finding their

optimal niches and outcompeting all generalists, resulting in no generalists? Or, would it lead

to an enrichment in habitat generalists that are capable of filling any niche into which they are

dispersed, given their strong ability to survive dispersal (Sriswasdi *et al.*, 2017; Schiro *et al.*,

2022)? It is challenging to answer this question, however, given the complexity of most

systems (many routes of dispersal, ecological disturbances, the influence of macroorganisms,

etc.).

Antarctic geothermal regions offer a unique opportunity to resolve the outcome of

high dispersal rates on the proportions of habitat generalists and specialists in a microbial

community for several reasons. First, Antarctica is one of the windiest places on earth

(Parish, 1988), which is likely to result in strong dispersal of microorganisms between sites

within Antarctica, as has been found in the McMurdo Dry Valleys (Archer *et al.*, 2019) and

other areas of Antarctica (Hughes *et al.*, 2004; Pearce *et al.*, 2010). Second, aerial dispersal is

the only possible route of dispersal between Antarctic geothermal regions, given the absence

of aquifers and bird traffic connecting regions, which reduces complexity. Third, Antarctic

geothermal regions have relatively simple communities, with little to no macroscopic life

present (Herbold, Mcdonald, *et al.*, 2014). Finally, Antarctic geothermal regions serve as

refugia for thermophiles, given the vast amount of ice and snow separating regions and the

extremely cold air conditions in Antarctica, resulting in little regional influence of non-

thermophiles on the microbial communities present. Interestingly, past studies of

thermophiles, mainly in hot spring environments, have found high niche specialization for

temperature or pH, with environmental selection being the primary factor influencing

community assembly and low dispersal rates between sites (Weltzer and Miller, 2013; Power

et al., 2018a, 2023; Mahato *et al.*, 2019; Fernandes☐Martins *et al.*, 2023). It is unknown

whether the strong winds in Antarctica are capable of overcoming the generally low dispersal

rates for thermophiles at Antarctic geothermal regions.

The southernmost active geothermal features in the world are located in Antarctica,

surrounded by vast swathes of ice and snow: Mt. Erebus, Mt. Melbourne, Mt. Rittmann, and

Deception Island (Herbold, Mcdonald, *et al.*, 2014). Within these geothermal areas, the

interplay between cold air (as low as -60☐) and hot soils results in a large variety of unique

features such as ice caves, ice chimneys, and ice hummocks. Of these regions, Mt. Erebus is

the highest (3794m above sea level, a.s.l.) and most volcanically active. Two of its largest

geothermal sites are Tramway Ridge and Western Crater, which have differing maximal

temperatures (65☐ vs 50☐), geothermal features (many active fumaroles in bare soil

producing large amounts of steam vs. warm soil mostly covered by ice hummocks with less

steam), physicochemistry (e.g., low pH vs. high pH), and productivity levels (visible

phototrophic mats and moss beds vs. primarily chemotrophic driven) (Noell *et al.*, 2022). Mt.

Melbourne (2733m a.s.l.) is located 358 km northeast of Mt. Erebus and only has two

geothermal fields, with maximal temperatures of 50□, limited fumarolic activity, and some

moss beds (Broady *et al.*, 1987). Geothermal activity on Mt. Rittmann (2600m a.s.l.), 450 km

northeast of Mt. Erebus and 103 km north of Mt. Melbourne, consists of one steep wall with

actively steaming fumaroles and hot soil (maximum 65□) that is relatively shallow due to the

steep aspect (Bargagli *et al.*, 1996). Given the strong maritime influence, presence of vascular

plants, and off-continent location of Deception Island north of the Antarctic circle, we

excluded this site from consideration in our study. For this study, we classified the two higher

temperature sites with highly active fumarolic activity (Tramway Ridge on Mt. Erebus and

Mt. Rittmann) as "Active" sites and the two lower temperature sites with less active

fumaroles (Western Crater on Mt. Erebus and Mt. Melbourne) as "Passive" sites.

The microbial communities present at these unique geothermal regions have been

incompletely characterized, especially at Mts. Rittmann and Melbourne. On Mt. Erebus,

culture-independent microbial community analyses have shown that cosmopolitan,

thermophilic microorganisms dominate the surface community of Tramway Ridge, while

novel and endemic members of poorly understood archaeal and bacterial groups are dominant

>2cm below the surface (Soo *et al.*, 2009; Herbold, Lee, *et al.*, 2014; Herbold *et al.*, 2024).

Our study comparing the communities at Tramway Ridge and Western Crater along thermal

transects found distinct microbial communities, primarily driven by strong differences in pH

and water content between the two sites (Noell *et al.*, 2022). Studies of life on Mt. Melbourne

have focused on macroscopic organisms (Broady *et al.*, 1987; Logan *et al.*, 2000; Skotnicki *et*

al., 2001) or were culture-based (Logan *et al.*, 2000; Bargagli *et al.*, 2004; Pepi *et al.*, 2005),

with two metagenomic datasets showing a community comprised of *Proteobacteria*,

Chloroflexota, and candidate phylum AD3 (now *Pseudomonadota*, *Chloroflexota*, and

Dormibacteraeota respectively) (Park *et al.*, 2021; Myeong *et al.*, 2024). Similarly, only

culture-dependent studies of the microorganisms at Mt. Rittmann have been conducted to-

date (Bargagli *et al.*, 1996; Logan *et al.*, 2000; Allan *et al.*, 2005; Poli *et al.*, 2006). These

high-elevation, Antarctic geothermal sites have been theorized to have acted as refugia for

terrestrial Antarctic life during highly glaciated periods (Fraser *et al.*, 2014).

In this study, we used a combination of 16S rRNA gene amplicon sequencing and

metagenomics to resolve the proportion of habitat generalists at these sites (i.e., do we see a

large group of microorganisms common to all sites?), and determine the ecological factors

shaping community assembly. Given that the microbial communities at these sites are largely

uncharacterized, we began by describing the taxonomy and functionality of the microbes

present at these sites before answering our ecological question. We hypothesized that the

lower temperature sites (Western Crater on Erebus and Melbourne) would have different

microbial communities than the higher temperature sites (Tramway Ridge on Erebus and

Rittmann), and that there would be a high proportion of habitat generalists across sites, driven

by high rates of aerial dispersal.

Experimental Procedures

Site Description and Sample Collection

Soil samples were collected from sites on Mt. Erebus, Mt. Rittmann, and Mt.

Melbourne, Antarctica, between 2010 and 2019. Sampling sites with coordinates are given in

Figure 1. On Mt. Erebus, two sites were sampled; the first was Tramway Ridge (TR), which

is the largest geothermal site on the Mt. Erebus summit and lies within Antarctic Specially

Protected Area (ASPA) No. 175. TR is relatively protected from the wind, has large numbers

of actively steaming fumaroles, producing gases enriched in CO_2 and CO (Ilanko *et al.*,

2019), with hot soil reaching 65□ in places, and has extensive moss and lichen mats covering

16% of the site (Management Plan For Antarctic Specially Protected Area No . 175, 2014).

The second site sampled on Mt. Erebus was Western Crater (WC), which is more exposed to

the wind and has very few active fumaroles and no visible moss or lichen mats; soil

temperatures only reach 50□ in ice-free grounds. On the Mt. Melbourne summit, samples

were collected on Cryptogram Ridge (also protected by ASPA No. 175), which, similar to

WC, is comprised of ice-free hot soil spots where temperatures reach 50□. There are some

patches of moss on Melbourne, and the few active fumaroles on Melbourne are covered by

ice hummocks, generating CO_2 and methane-rich gases (Skotnicki *et al.*, 2001; Management

Plan For Antarctic Specially Protected Area No . 175, 2014). On Mt. Rittmann, samples were

collected from the single geothermally active area on the mountain, which is also within

ASPA No. 175. This site is particularly steep (~40 degree slope), with ice free hot soil

reaching 63□ and many actively steaming fumaroles with gases of unknown composition, as

well as a few moss patches (Management Plan For Antarctic Specially Protected Area No .

175, 2014).

Soil temperatures were measured prior to sampling using a Checktemp 1C electronic

temperature sensor (Hanna Instruments, Rhode Island, United States). Mt. Erebus samples

were collected along temperature transects, as detailed in (Noell *et al.*, 2022). Samples were

collected using a sterile spatula after first discarding the top centimetre of soil, except for one

sample each at Rittmann and Melbourne where the top mat-like layer of soil was taken as the

sample (Ritt3M-3 and Mel1M-2). Samples were placed into sterile 50 mL tubes and frozen

for transport to the University of Waikato (Hamilton, New Zealand) for analysis.

Geochemistry

Soil samples were analysed for geochemistry using a variety of analytical techniques.

pH and electroconductivity (EC) were measured by adding MilliQ water to soil samples at a

1:2.5 soil:water ratio. pH and EC measurements were made using a Thermo Scientific Orio 4-

Star Benchtop pH/Conductivity Meter (Thermo Fisher Scientific) using a 3-point calibration.

For gravimetric water content (GWC) measurements, 3 g of soil sample was dried at 105□

until the sample weight remained unchanged. GWC was then calculated as the percentage of

weight loss during drying (Herbold, Lee, *et al.*, 2014). For elemental composition (B, Na,

Mg, P, K, Ca, V, Cr, Fe, Mn, Co, Ni, Cu, Zn, Ga, As, Se, Sr., Ag, Cd, Ba, Tl, Pb, and U),

dried samples (1g) were prepared and analysed as described in (Noell *et al.*, 2022). For total

carbon and nitrogen measurements, dried samples were also ground as for elemental analysis,

then sent to Analytica (Hamilton, New Zealand) for analysis. For total organic carbon (TOC)

and nitrogen (TON) measurements, samples were prepared similarly to (Noell *et al.*, 2022).

Briefly, 1g of samples were weighed out into acidified glass jars with cardboard disks in lids;

jars and lids were washed as for the ICP-MS protocol. 1 mL 6N HCl was added to samples

and dried in an oven in a fume hood for 24 hours at 95□. 100 mg aliquots of samples were

analysed for TOC and TON using an automatic CN analyser.

DNA Extraction, Amplification, and Sequencing

For 16S rRNA gene amplification as well as metagenome sequencing, genomic DNA

was extracted from 0.5 to 0.9 g of each soil sample using a modified CTAB/bead beating

method (Noell *et al.*, 2022). To have enough DNA from low biomass samples for

metagenomes, multiple replicates (up to 10) of samples were extracted and pooled. Genomic

DNA was quantified using a Qubit 2.0 Fluorometer (Thermo Fisher Scientific,

Massachusetts, United States).

The 16S rRNA gene (V4-V5 region) was amplified using PCR with the 515YF

(5'CCATCTCATCCCTGCGTGTCTCCGACTCAGXXXXXXXXXXXXXXGATGTGYCAG

CMGCCGCGGTAA3') and 926R

(5'CCACTACGCCTCCGCTTTCCTCTCTATGGGCAGTCGGTGATCCGYCAATTYMTT

TRAGTTT3') primers (Caporaso *et al.*, 2018; Ul-Hasan *et al.*, 2019), adapted for Ion Torrent

sequencing using fusion primers with a unique tag (Whiteley *et al.*, 2012). PCR reaction

conditions, library prep, and 16S rRNA gene sequencing are the same as described in Noell,

S. E. *et al.* (2022) (Noell *et al.*, 2022). 16S rRNA gene sequencing was conducted on all 21

samples on an Ion Torrent PGM (Thermo Fisher Scientific) at the Thermophile Research

Unit (Hamilton, NZ).

Four samples were chosen for shotgun metagenome sequencing: Ritt1-3, Ritt3M-1

(mat sample from Rittmann), WC-50, and Mel1-3; the mat sample was included to improve

the odds of recovering cyanobacterial genomes. A fifth sample (from Tramway Ridge) was

previously extracted and sequenced using Illumina Hi-Seq; the two surface metagenome

samples from this previous project (SRR6519253 and SRR6519256) were downloaded from

the SRA using SRAtoolkit v3.0.1 and combined into one larger sequence file. For the four

new metagenomic DNA samples, extracted DNA was sent to GENEWIZ Genomics Center

(Suzhou, China). Next generation sequencing library preparations were constructed following

the manufacturer's protocol (NEBNext® Ultra™ DNA Library Prep Kit for Illumina®). For

each sample, 1 μg genomic DNA was randomly fragmented to <500 bp by sonication

(Covaris S220). The fragments were treated with End Prep Enzyme Mix for end repairing, 5'

Phosphorylation and dA-tailing in one reaction, followed by a T-A ligation to add adaptors to

both ends. Size selection of Adaptor-ligated DNA was then performed using AxyPrep Mag

PCR Clean-up (Axygen), and fragments of ~410 bp (with the approximate insert size of 350

bp) were recovered. Each sample was then amplified by PCR for 8 cycles using P5 and P7

primers, with both primers carrying sequences which can anneal with flowcell to perform

bridge PCR and P7 primer carrying a six-base index allowing for multiplexing. The PCR

products were cleaned up using AxyPrep Mag PCR Clean-up (Axygen), validated using an

Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA), and quantified by

Qubit2.0 Fluorometer (Invitrogen, Carlsbad, CA, USA). Then libraries with different indexes

were multiplexed and loaded on an Illumina HiSeq instrument according to manufacturer's

instructions (Illumina, San Diego, CA, USA). Sequencing was carried out using a 2x150

paired-end (PE) configuration; image analysis and base calling were conducted by the HiSeq

Control Software (HCS) + RTA 2.7 (Illumina) on the HiSeq instrument.

Sequence Quality Control and Taxonomic Assignment

For 16S rRNA gene amplicon sequences, the raw reads from the Ion Torrent PGM

were processed using the DADA2 pipeline with recommended Ion Torrent parameters

(Callahan *et al.*, 2016). Chimeras were removed using the function removeBimeraDenovo,

method "consensus." Taxonomy assignment was performed using assignSpecies with the

naïve Bayesian classifier and SILVA database version 138.1. Eukaryotic, mitochondrial, and

chloroplast sequences were removed and an unrooted phylogenetic tree built using the

neighbour-joining method with the decipher (Wright, 2016) and phangorn (Schliep, 2011)

packages in R, generating final amplicon sequence variants (ASVs).

Unfortunately, two samples (one from Tramway Ridge, TR2-64, one from Western

Crater, WC-10) produced no sequence data. As these two samples had also been sequenced in

a previous sequencing run (from (Noell *et al.*, 2022)), we investigated whether sequence data

for these two samples from the previous run could be safely used with the second sequence

run samples. To control for batch effects, we tested whether other samples from the two runs

had similar microbial communities and found that there was no pronounced batch effect for

all samples between the two runs (Figure S1). Thus, we decided to use sequence data for

TR2-64 and WC-10 from the first sequencing run alongside the second run; we denote these

samples as TR2-64-1 and WC-10-1 to distinguish them from samples in the second run (TR1-

32-2, WC-20-2, etc.).

For metagenome sequences, demultiplexing was performed by bcl2fastq 2.17. Raw

data was filtered as follows: (1) Discard pair-end reads with adapter; (2) Discard pair-end

reads when the content of N bases is more than 10% in either read; (3) Discard pair-end reads

when the ration of bases of low quality (Q<20) is more than 0.5 in either read. All four

metagenome samples were quality checked using FastQC v0.11.5 (Andrews) and the reports

visualized using MultiQC (Ewels *et al.*, 2016). Illumina adaptors were trimmed using

trimmomatic v0.39 (Bolger *et al.*, 2014) and sequences were trimmed for quality using the

following command:

trimmomatic PE -phred33 <input files> ILLUMINACLIP:NexteraPE-PE.fa:2:30:10:2:True

MINLEN:70 SLIDINGWINDOW:4:15

In light of the large amount of duplicate sequences observed in some samples

(especially the Rittmann sample) and recent evidence that suggests deduplication is beneficial

(Zhang *et al.*, 2022), we used fastp v0.23.2 (Chen *et al.*, 2018) to deduplicate all metagenome

sequences using the following command:

fastp –in1 <input file forward reads> --out1 <output forward reads> --in2 <input reverse

reads> --out2 <output reverse reads> -A -D -G -Q -L –html <summary file>

Reads with ambiguous bases were removed using prinseq-lite v0.20.4 (Schmieder and

Edwards, 2011) using the -ns_max_n 0 command.

Metagenome analysis, assembly, and binning

We assessed the taxonomy of the quality trimmed metagenomes at the read level

using singleM v0.13.2 (https://github.com/wwood/singlem) with the singleM pipe command;

we saw no differences when we used the raw, non-quality trimmed sequences or the quality

trimmed sequences (data not shown). We used Nonpareil v3.401 (Rodriguez-R and

Konstantinidis, 2014; Rodriguez-R *et al.*, 2018) to assess the fraction of community diversity

sampled by the metagenome.

For metagenome assembled genome (MAG) assembly, binning, and processing, we

used the metaWRAP pipeline v1.3.2 with default parameters (Uritskiy *et al.*, 2018). We used

Metaspades (Nurk *et al.*, 2017) as the assembly program within metaWRAP with default

settings. To minimize the amount of ambiguous bases in the assemblies, we used the non-

scaffolded contigs for binning; these non-scaffolded contigs produced bins that were similar

in number and quality to the scaffolded contigs (for scaffolded vs non-scaffolded, the bins

produced had 94.1% vs 94.0% average completeness, 1.5% vs 1.6% average contamination,

63% vs. 64% average GC, 41309 vs. 40696 average N_{50}). We used prodigal v2.6.3 in

metagenome mode (Hyatt *et al.*, 2010, 2012) to predict coding genes from the non-scaffolded

assemblies; these predicted coding sequences were used for the metagenome functional

analysis. For binning contigs in the metaWRAP pipeline, we used metaBAT2 (Kang *et al.*,

2019), MaxBin2 (Wu *et al.*, 2016), and CONCOCT (Alneberg *et al.*, 2014). For the

bin_refinement module of metaWRAP, we used targets of 80% and 5% for completeness and

contamination, respectively. Bins were then re-assembled and cleaned with MAGpurify

v2.1.2 (Nayfach *et al.*, 2019) using the phylo-markers, clade-markers, gc-content, known-

contam, and clean-bin modules to produce the final MAGs. The final statistics for these

MAGs were produced using checkM v1.0.12 (Parks *et al.*, 2015) with the lineage_wf

command. We used GTDB-Tk v2.1.1 (Chaumeil *et al.*, 2022) for taxonomy assignment of

MAGs (GTDB database R214) and both Prodigal and Prokka v1.11 (Seemann, 2014) for

predicting coding gene regions. We predicted optimal growth rate of MAGs using gRodon

(Weissman *et al.*, 2021) and optimal growth temperature using Tome v1.0.0 (Li *et al.*, 2019).

Finally, we used pfam_scan (https://github.com/aziele/pfam_scan) to predict pfam domains

in MAGs (pfam release 35.0).

We calculated the presence/absence of MAGs across sites by utilising MinHash

sketching techniques implemented with sourmash (Brown and Irber, 2016). Specifically, we

created sourmash signatures of all MAGs and quality filtered metagenomic read sets from all

four sites using a k-mer size of 21 and scaled to 1000. These settings allowed us to query for

the presence of a MAG within a site with greater sensitivity than by using a higher k-mer or

scale value. We selected these metrics to minimize the false negative of not finding a MAG

that is present, though we acknowledge this possibly increases the false positive, something

recently found by other groups (Koslicki *et al.*, 2023). Finally, we included abundance

weighting in the signature creation step of the MAG and metagenome signatures so we could

measure relative abundances of MAGs present. To query presence/absence, we used

sourmash *gather* on each MAG against each metagenome to calculate containment scores.

These scores are a proxy for Average Nucleotide Identity where a containment score of 0.2 is

equivalent to an ANI of 0.95 for a signature created with a k-mer of 21. We therefore

concluded that a MAG was present within a metagenome if its score was greater than 0.2, to

account for the varying completeness levels of the MAGs and to again minimize the false

negative rate. We measured MAG abundances within each site using the abundance weighted

percent match between the MAG and the metagenome. This correlates to the proportion of

metagenomic reads that would map to the MAG, a proxy for abundance. The final set of

MAGs was deduplicated for all analyses using a 95% ANI cutoff. Where duplicate MAGs

were found, the MAG from the site with its highest abundance was retained.

Analysis of 16S rRNA Gene Sequence Data

Using 16S rRNA gene sequences, microbial community structure was assessed using

PcoA of center log-ratio transformed (CLR) sequence abundance on a Unifrac distance

matrix using the vegdist function. Statistical differences between sites were evaluated first by

testing samples did not differ significantly in their dispersion (using an analysis of

multivariate homogeneity, PERMDISP), then by using a permutational analysis of variance

(PERMANOVA) within the R package vegan v2.6-2 (Oksanen *et al.*, 2020). Alpha diversity

was also assessed within vegan via the Shannon index. Venn diagrams were plotted using the

R package ggVennDiagram v1.2.0 (Gao, 2021); a custom modification to the

ggVennDiagram function was used to color the sections of the Venn Diagram by the

abundance of ASVs in those sections. Heatmaps were plotted using the R package pheatmap

v1.0.12 (Kolde, 2019). The random forest differential abundance analysis was conducted

using the R package microeco v0.13.0 with taxa level set to "all" (Liu *et al.*, 2021).

To identify physicochemical parameters that correlated with microbial community

structure, the mantel test function within microeco was used with default parameters. The

factors that had a non-adjusted two-sided p-value < 0.05 were plotted using a distance-based

redundancy analysis (dbRDA). To test for significant differences between physicochemical

parameters, the cal_diff (method = anova) and plot_diff functions within the trans_env

function of microeco were used. To analyse clustering of sites by physicochemical

parameters, principal coordinates analysis (PcoA) of physicochemical parameters in a

Euclidian distance matrix was performed using the vegdist function in the microbiome

package v.1.12.0 (Lahti and Shetty, 2017).

Metagenome Functional Analysis

We estimated the metabolic potential of each site using a method described by

Greening and colleagues (Greening *et al.*, 2019). In short, we used DIAMOND v. 2.1.6

(query cover > 80%) to search quality filtered assembled ORFs of the metagenomes against

custom protein databases of representative metabolic marker genes designed by the authors.

The metabolic marker genes include multiple pathways or energy harvesting systems,

namely: sulfur cycling (AsrA, FCC, Sqr, DsrA, Sor, SoxB), succinate oxidation (SdhA),

selenium cycling (YgfK), rhodopsin (RHO), reductive dehalogenation (RdhA), phototrophy

(PsaA, PsbA, energy-converting microbial rhodopsin), oxidative phosphorylation (AtpA),

nitrogen cycling (AmoA, HzsA, NifH, NarG, NapA, NirS, NirK, NrfA, NosZ, NxrA, NorB),

NADH oxidation (NuoF), methane cycling (McrA, MmoA, PmoA), isoprene oxidation

(IsoA), iron cycling (Cyc2, MtrB, OmcB), hydrogen cycling (catalytic subunit of [NiFe]-

hydrogenases, the catalytic domain of [FeFe]-hydrogenases, and Fe-hydrogenases), fumarate

reduction (FrdA), carbon monoxide oxidation (CoxL, CooS), carbon fixation (RbcL, AcsB,

AclB, Mcr, HbsT, HbsC), arsenic cycling (ARO, ArsC), and aerobic respiration (CoxA,

CcoN, CyoA, CydA). We filtered results based on different identity thresholds: PsaA (80%),

PsbA and IsoA, AtpA, ARO, YgfK (70%), HbsT (75%), group 4 [NiFe]-hydrogenases,

[FeFe]-hydrogenases, CoxL, AmoA, NxrA, NuoF (60%), and all others with an identity

threshold of 50%. We normalized the relative abundance of present markers to RPKM.

RPKM is recommended for relative abundance comparisons within metagenomic

datasets because it normalizes the data based on both sequence depth (per million reads) and

sequence length (in kilobases) (Liao *et al.*, 2023). We used a three-step procedure to calculate

the relative abundance of each metabolic marker in the community. First, we used quality

controlled reads to normalize coverage of ORFs to RPKM using CoverM v0.6.1 contig

(https://github.com/wwood/CoverM) (with parameters: -p bwa-mem -min-read-aligned-

percent 0.75 -min-read-percent-identity 0.95 --min-covered-fraction 0 -m rpkm). In the

second step, we aligned high-quality unassembled reads to 14 universal single-copy

ribosomal marker genes used in SingleM v.0.12.1 (https://wwood.github.io/singlem/) and

PhyloSift (Darling *et al.*, 2014) using DIAMOND (query cover > 80%, bitscore > 40) and

normalized to RPKM as above. Finally, we calculated the average gene copy number of each

marker in the community by dividing the read count to the gene (in RPKM) by the mean of

the read counts of the 14 universal single-copy ribosomal marker genes (in RPKM). Finally,

we visualized the data using ggplot2.

Phylogenomic Tree Analysis

For the *Cyanobacteria* phylogenomic tree, we used GtoTree v1.7.00 (Lee, 2019), with

our 9 cyanobacterial MAGs along with GTDB species cluster representatives from the same

three Cyanobacterial families, resulting in 217 members of *Nostocaceae*, 38 for

Leptolyngbyaceae, and 55 for *Obscuribacteraceae* (accessed on 13 October 2023). We used

the *Cyanobacteria*-specific single copy gene (SCG) set available in the GtoTree package. The

SCG threshold was lowered to 35% to keep the *Obscuribacteraceae* genomes. We used four

members of the Terrabacteria supergroup (*Streptomyces coelicolor*, *Clostridium botulinum*,

Chloroflexus aurantiacus, and *Meiothermus ruber*) as the outgroups to root the tree. The tree

was visualized and edited using the website Interactive Tree of Life

(https://itol.embl.de/upload.cgi).

Community Assembly Analysis

To compare the abundance of cosmopolitan and site-specific microorganisms in the

sites/samples, ASVs were categorized by the number of sites they were present at, with the

threshold for presence at a site being greater than 0 relative abundance within either the

sample or site. MAGs were classified as present or absent as described above.

To calculate niche breadth for ASVs, we first eliminated ASVs only found at one site

(1878 ASVs reduced to 832 ASVs), equivalent to a 5% sample occupancy cutoff (stricter

than (Sriswasdi *et al.*, 2017; Xu *et al.*, 2022) but not as conservative as (Hernandez *et al.*,

2023)). Next, we used the R package MicroNiche v1.0.0 (Finn *et al.*, 2020), with the default

settings for the limit of quantification (LOQ) threshold (q = 1.6). Levin's B_N was used to

detect generalist/specialists across the four sample sites, while Hurlbert's B_N was used to

detect generalist/specialists with regards to temperature or pH. An adjusted p-value cut-off of

0.05 was used, and thresholds for generalist/specialist were determined from the null model

distribution produced by the package during nice breadth calculations. Additionally, we

calculated social niche breadth scores for ASVs using the script calculate_snb.py v0.2 with

default parameters (von Meijenfeldt *et al.*, 2023).

To measure the strength of different ecological factors on shaping community

assembly in the 16S rRNA gene sequence data, we used the following commands within the

microeco package with default settings: trans_nullmodel, cal_ses_betamntd, cal_rcbray, and

cal_process; this approach is based on (Stegen *et al.*, 2013). To generate a distance-decay plot

to look at the influence of geographic distance on the phylogenetic differences between

ASVs, we calculated the geographical difference between sampling sites using the distm

function (distVincentyEllipsoid option) in the geosphere v1.5-18 R package (Hijmans, 2022).

To calculate biological differences in the microbial communities, we chose to use the Chao

index, given the differences in the sample sizes in our data set (Chao *et al.*, 2004),

implemented with the vegdist function in the vegan R package.

MAG Functional Analysis

Functional annotation of MAGs was conducted using METABOLIC-G (Zhou *et al.*,

2022), using the predicted proteins from each MAG as predicted by Prodigal (as above) with

default parameters. Specifically, we used the full KOFAM database and set the cut-off value

to assign the presence of a module to 0.75. We categorized the metabolic categories used by

METABOLIC into either Pathway (sets of genes that take a metabolite from start to end-

point) or Category (contains multiple distinct, not inter-connected pathways). We counted a

Pathway as being present in a MAG if at least 50% of the genes in the pathway were present,

and a Category as being present if there was at least one gene in that Category present.

Statistical Analyses

All statistical analyses were conducted in R (R Core Team, 2020), with plots

produced in ggplot2 v3.3.6(Wickham, 2016) and edited for aesthetics in Inkscape

(https://www.inkscape.org). Data analysis was preformed using tidyverse v1.3.1 (Wickham *et*

al., 2019). The ggpubr v0.6.0 package in R was used to add the results of a one-way ANOVA

test (when comparing all four sites and when data was normally distributed), a one-way

Krustal-Wallis test (when comparing all four sites for non-normally distributed data), or a

400 two-sided Mann-Whitney test (when comparing active vs. passive site types) to plots

401 (Kassambara, 2023).

Results

Prokaryotic Community Diversity and Composition

To characterize the microbial communities inhabiting Antarctic geothermal sites, we

extracted DNA from 21 soil samples from three geothermal areas in Victoria Land,

Antarctica: Mt. Rittmann, Mt. Melbourne, and two sites on Mt. Erebus, Tramway Ridge (TR)

and Western Crater (WC), for a total of four sampling sites (see Figure 1 for maps and site

descriptions). Two of these four sites, Melbourne and WC, have little to no fumarolic activity

(i.e., fumarolic passive) and lower maximal soil temperatures (maximum 50□). The other

two sites, Rittmann and TR, have many actively steaming fumaroles (i.e., fumarolic active)

and higher maximal soil temperatures (maximum 65□). Thus, we categorized WC and

Melbourne as "passive" sites and TR and Rittmann as "active" sites. We used 16S rRNA

gene amplicon sequencing to study the microbial communities present in these samples,

generating 1,871 amplicon sequence variants (ASVs) from 926,649 reads (Table S1). We

also conducted metagenome sequencing on four samples, one from each site, with almost all

metagenomes reaching sequencing saturation (nonpareil C value of 1 for all except TR,

which was 0.93; Figure S2B). Metagenome diversity, based on nonpareil diversity indices,

were 17.8, 15.6, 15.3, and 15.2 for TR, WC, Melbourne, and Rittmann, respectively, all very

low compared to other habitat microbiomes (17.5 for human-associated, 19 for lake water)

(Rodriguez-R *et al.*, 2018). From these metagenomes, we binned 140 metagenome assembled

genomes (MAGs): 42 from Melbourne, 15 from Rittmann, 23 from TR, and 60 from WC, all

with completeness >80% (except for two that dropped to 77% during cleaning) and

contamination <5% (Table S2).

Overall, we found that the microbial communities at WC and Melbourne (hereafter,

passive sites) were more closely related to each other than to TR and Rittmann (hereafter,

active sites) (Figure 2). This was most apparent from the 16S rRNA samples, where passive

sites had significantly higher alpha diversity than the active sites (Figure 2A), clustered very

closely together in a principal coordinates analysis (PCoA) (Figure 2B), and were more

similar to each other based on abundance patterns both at the phylum (Figure S3A) and class

level (Figure S3B). MAGs from passive sites also clustered together based on similarities in

abundance profiles, separate from active sites (Figure S4). However, a pairwise

PERMANOVA test using the 16S rRNA gene sequence data showed no significant

difference between the microbial communities at any of the sites except between TR and WC

($p = 0.05$), which could be due to the small number of samples analysed at Melbourne and

Rittmann. TR and Rittmann also clustered separately from each other in the PCoA plot,

indicating distinct microbial communities at the two active sites (Figure 2B).

Based on the 16S rRNA gene sequence-based and metagenome-based community

composition of the sites, we found that differences between site types were primarily

attributable to greater abundances of members of *Crenarchaeota* (class *Nitrososphaeria*) at

active sites, while passive sites had significantly greater abundances of phylum

Planctomycetota (family *Gemmataceae*) and class *Acidobacteriae* within *Acidobacteriota*

(Figure 2C-D, Figure S3). When looking for distinguishing taxa at each of the four sites, we

found that TR was the easiest to distinguish from all other sites based on greater abundances

of phylum *Bacteroidota* and the archaeal family *Nitrososphaeraceae* present (Figure 2E).

Interestingly, both the Rittmann and WC metagenomes had much greater relative abundances

of *Nitrososphaeria* compared to the corresponding 16S rRNA gene sequence-based

community profile (85% vs. 20% for Rittmann, 25% vs. 5% for WC for metagenome vs. 16S

rRNA, respectively), which could indicate primer selection against this group.

We used our complete set of non-dereplicated MAGs, categorized by the site the

MAG was binned from, to examine genomic characteristics of microbes from each site. We

found that MAGs binned from active sites were distinguishable by their significantly smaller

average genome size (average: 3.1 Mbp for active, 4.8 Mbp for passive) and significantly

higher predicted optimal growth temperature (OGT) (average: 50.8□ for active, 38.0□ for

passive); MAGs from Melbourne had the largest average genome size and the lowest

predicted OGT of the four sites (Figure 3A-B). We did not see significant differences in

%GC, coding density, or predicted maximal growth rate between MAGs from the two site

types or the four sites (Figure S5A-B).

When we explored the physicochemical parameters that could be influencing the

microbial communities at these sites, we first found that the three volcanoes clustered

separately from each other in the physicochemical data (Figure S6A), with elemental

composition playing a large role in this separation (Figure S6B), reflecting the unique

geological origin of each volcano. When we looked for correlations between physicochemical

parameters and biological communities (16S rRNA gene sequence-based), although many

parameters were significantly correlated (Figure S6C), only magnesium remained significant

($p = 0.033$) after Benjamini and Hochberg adjustment. Values for selected parameters are

presented in Figure S6D and all physicochemical parameter values are given in Table S3.

Exploration of Cyanobacterial Diversity and Functions

We used the *Cyanobacteria* group as a case study to explore the influence of

physicochemistry on the distribution and functioning of these important primary producers

and to investigate the global uniqueness of these microorganisms. We binned an additional

five cyanobacterial MAGs from the metagenome of Ritt3M-1, a cyanobacterial mat sample

from Rittmann; this was necessary for our analysis as there were no cyanobacterial MAGs

recovered from the other Rittmann metagenome sample. In total, we retrieved 50

cyanobacterial ASVs and 9 cyanobacterial MAGs from our data set.

Overall, we found organisms from three families (*Nostocaceae*, *Leptolyngbyaceae*,

and *Obscuribacteraceae*), but could only identify a few instances of clear correlations

between physicochemical parameters and the distribution and functioning of these groups

(Figure 4A, B). Supporting earlier findings at the entire microbial community scale,

Tramway Ridge had a distinct *Cyanobacteria* community, with species from

Leptolyngbyaceae and the genus *Fischerella* (family *Nostocaceae*) dominant and an absence

of *Obscuribacteraceae*. *Leptolyngbyaceae* was positively correlated with gravimetric water

content (GWC) and negatively correlated with nickel (Figure 4C). In general, *Nostocaceae*

was the most widely distributed and had a positive relationship with nickel and sulfur and

was negatively correlated with GWC (Figure 4C). Only MAGs from *Nostocaceae* had the

capacity to fix nitrogen (Figure S7), which matches previously measured nitrogen fixation by

this group at Melbourne (Broady *et al.*, 1987).We found that *Obscuribacteraceae* MAGs

were non-photosynthetic, as previously observed for this group (Soo *et al.*, 2014), with

pathways for ethanol fermentation and fatty acid degradation instead (Figure S7B).

To illuminate potential evolutionary relationships between the *Cyanobacteria* MAGs

from this study and globally distributed representatives of these families from GTDB (GTDB

species cluster representatives, 217 *Nostocaceae*, 38 *Leptolyngbyaceae*, and 55

Obscuribacteraceae), we used a phylogenomic approach (Figure S8). We observed that all

MAGs from our study clustered with globally distributed members of these families, from the

Mojave Desert to China to the continental subsurface. The short branch lengths we observed

for our MAGs also confirming the cosmopolitan distribution of these families, as has been

found previously at Tramway Ridge (Herbold *et al.*, 2024). These results are dependent on

the completeness of the database used and should not be treated as conclusive evidence.

Functional Comparison of Sites

We analyzed the functional potential of potentially representative microbial

communities at the four sites using the community metagenome sequences, post-assembly,

looking at the abundances of 52 conserved marker genes that characterize various key

metabolic pathways (Figure 3C). At all sites, hydrogenases involved in atmospheric H_2

scavenging (Group 1h and 1l NiFe hydrogenases) (Greening *et al.*, 2016) were present at low

abundances.

Several key differences were notable between sites. First, TR was distinguishable by

an enrichment in the number of genes involved in sulfur (SoxB, Sqr, and FCC) and nitrogen

(NosZ, NxrA, NirS, NirK, NorB, and NarG) metabolisms (Figure 3C). These differences are

expected given the relatively large concentrations of TN and sulfur at TR (Figure S6). A

notable exception to the suite of nitrogen metabolising genes detected at TR is the lack of

AMO (ammonia monooxygenase) encoding genes, an exception which has recently been

accounted for (Herbold *et al.*, 2024). Melbourne and WC were highly similar to each other,

except for the presence of methane monooxygenase genes at Melbourne and not WC (Figure

3C). Several MAGs in the genus *Methylocella* were found at Melbourne, confirming the

potential for methane metabolism there, which aligns with large measured concentrations of

methane at Melbourne (Geyer, 2021) (Table S2). Notably, Rittmann was distinguishable by

an enrichment in marker genes for anaerobic metabolism (fumarate reductase, FrdA) and CO

oxidation (CoxL), as well as much lower abundance of genes related to aerobic metabolism

(cytochrome aa3 oxidase, CoxA; NADH-ubiquinone oxidoreductase, NuoF; and F-type ATP

synthase, AtpA) (Figure 3C). The gas composition of Rittmann fumaroles has yet to be

studied, but based on these results, is likely to be enriched in CO and depleted in oxygen.

Habitat Specialization and Community Assembly Factors

Following our taxonomic and functional analysis of these Antarctic geothermal sites,

we asked whether there was a large community of shared microorganisms between these sites

(habitat generalists) and, if so, whether dispersal between sites was an important factor in

structuring these communities. To do this, we first classified 16S rRNA gene sequence ASVs

and MAGs by the number of sites or (for ASVs) samples at which they were present, after

confirming that all samples were similarly sampled using rarefaction (Figure S2). Based on

this classification, we observed that most ASVs (90%) and MAGs (77%) in our study were

only found at either one or two sites/samples (Figure S9A-C, F, I). However, these

site/sample-specific ASVs and MAGs comprised only a small fraction of the total read count

at all samples. Instead, it was the ASVs and MAGs that were found across all four sites (36

ASVs, 3% of all ASVs; 2 MAGs, 2% of all MAGs) that were generally the most abundant

(>30% of all reads for both ASVs and MAGs) (Figure S9A, B, D, G, J). Melbourne and WC

shared the most ASVs and MAGs (120 ASVs, 9% of all ASVs; 36 MAGs, 32% of all MAGs)

(Figure S9A, B), consistent with their similar overall microbial community composition. TR

shared very few ASVs with any other site (Figure S9A, B).

Next, we asked whether these common microorganisms were adapted to thrive at all

of the sites (habitat generalists), or if their presence at all sites was a product of stochastic

processes. We used a robust implementation of Levin's niche breadth (Finn *et al.*, 2020) with

our 16S rRNA gene sequence data (reduced to 832 ASVs by eliminating ASVs only present

in one sample) to identify specialists and generalists. Levin's niche breadth (B_N) is calculated

by:

$$Levins' B_N = \frac{1}{R \sum_{i=1} p_i^2}$$

where R is the number of differing environments and p_i is what proportion of all instances of

taxon j is found in environment i. B_N values close to 1 indicate broad habitat occupancy,

while values close to 1/*R* indicate habitat specialization. In our data set, we defined each of

our four sites as different environments and used ASV read counts for the taxon proportions.

In contrast to the theoretical distribution of niche breadth scores (Figure S10A),

surprisingly, we did not find any habitat generalists, with 95% of ASVs instead identified as

specialists and the rest as intermediate (Figure 5A). Because two of our sample sets (TR and

WC) were collected along temperature gradients, we re-ran the analysis with only high

temperature samples (>35☐), but again found 96% of ASVs were specialists; only one ASV

(ASV_5, GAL15) was returned as a generalist (Figure S10B). This absence of habitat

generalists held true both with a relaxed limit of quantification (LOQ) for including ASVs in

the niche breadth analysis, as well as when we randomly subsampled within each site to bring

the sample number from all sites to the same amount (data not shown). A recently developed

social niche breadth metric (SNB (von Meijenfeldt *et al.*, 2023)) identifies specialists or

generalists based on the compositional similarity of the communities that microbes inhabit.

Using this metric, we found a distinct skew towards specialization (lower SNB score) in our

data set, with a much longer tail on the specialist side both in raw SNB scores and in

modified z-scores at all taxonomic ranks, supporting our findings of high habitat

specialization (Figure S10).

A refinement to Levins' niche breadth adds in a measure of the resource availability

across environments, called Hulbert's niche breadth (Finn *et al.*, 2020). This allows

measurement of habitat specialism or generalism in reference to environmental gradients; in

our case, we chose to examine temperature and pH, as these were identified as having the

strongest correlations with community composition (Figure S6C). Again, we found a strong

skew towards specialization for temperature and pH (95% specialists for temperature, 73%

for pH) (Figure 5B, C). However, there were two ASVs identified as generalists for

temperature alone (ASV 5, phylum GAL15) or both temperature and pH (ASV 42, phylum

Acidobacteriota, genus *Bryobacter*). When examining the abundance of these ASVs across

temperature and pH, they clearly had a broader distribution across different temperature/pH

values than highly specialized ASVs (Figure S10E). However, their distribution within sites

is more variable (Figure S10E).

Next, we explored what ecological factors might be driving the lack of habitat

generalists in our data set. First, we used a common method, beta nearest taxon index and

Bray-Curtis-based Raup-Crick, to calculate contribution of different factors to community

assembly in the 16S rRNA gene sequence data (Stegen *et al.*, 2013; Zhu *et al.*, 2022; DePoy

and King, 2023). We found that, across all sites, ecological drift was the strongest factor

(30.5%), with homogenous dispersal (i.e., similar community structure caused by high

dispersal rates) was the second strongest factor (25.5%) (Figure 5D). On the other hand, 20%

of the community assembly could be explained by dispersal limitation (i.e., dissimilar

community structure caused by low dispersal rates) (Figure 5D). Environmental selection

(either variable selection, i.e. dissimilar community structure caused by different

environmental conditions, or homogenous selection, i.e. similar community structure caused

by similar environmental conditions) was the weakest factor (11.4% variable selection,

12.9% homogenous selection) (Figure 5D). Similar results were found when comparing only

active sites or only passive sites, although homogeneous environmental selection was much

stronger within passive sites than active (Figure 5D). The importance of ecological drift and

dispersal in structuring these communities was again apparent when we examined the

distance-decay plot for the 16S rRNA gene sequence data, finding a relatively flat slope of

the linear regression line ($m = -0.0003$), which points to strong dispersal between sites

(Figure 5E). This shallow slope held true when Active and Passive site samples were

considered separately as well ($m = -0.0007$ and -0.0003 for Active and Passive, respectively)

(Figure 5F, G). Slopes for similarly large-scale studies have shown much larger slopes (e.g.,

(Whitaker *et al.*, 2003; Green *et al.*, 2004; Locey *et al.*, 2020)).

Functional Imprint of Habitat Specialization

Finally, we explored the impact of high levels of habitat specialization on the

microbial communities from these sites by exploring the genomic and functional profiles of

MAGs based on site distribution. Generally, habitat generalists tend to be metabolic

generalists (Barberán *et al.*, 2014; Chen *et al.*, 2021a); given the absence of habitat

generalists in our data set, we predicted that the number of sites that a microbe is present at

should have little to no correlation with genome size, number of unique protein domains

(used as a proxy for metabolic breadth), or predicted metabolic pathways present. Rather, we

would expect that most microorganisms would likely be metabolic specialists.

Indeed, we found very few significant differences (two-sided Wilcox test, Benjamini

& Hochberg-adjusted p-value < 0.05) between different MAG site occupancy categories in

genome size, number of unique pfam domains, or predicted metabolic pathways/categories

present (Figure 6A-B). We did find that MAGs occupying three sites had significantly more

metabolic pathways/categories than MAGs only occupying one site. Additionally, we

observed significantly larger genome sizes and numbers of unique pfam domains in MAGs

present at both WC and Melbourne compared with those occupying only WC (Figure 6B).

Correlations between other genome characteristics and site occupancy were non-significant

(Figure S11A-B). When we explored the ASVs that were identified previously as generalists

for temperature and pH, ASVs 5 and 42, we were able to match ASV 5 to the MAG

Ritt_bin.6.strict_final with 100% identity. This MAG had some hallmarks of generalism,

with 20.9 predicted metabolic pathways/categories but a genome size of only 3.4 Mbp. Two

MAGs (Erebus-WC_bin.32.strict_final and Mel_bin.5.orig_final) were classified to the

Bryobacteraceae family, as was ASV 42; neither MAG contained any 16S sequence, so we

cannot be sure if they match ASV 42. Still, both MAGs have genome sizes and numbers of

predicted metabolic pathways/categories on the upper end of the distribution (6.2 Mbp and

10.1 Mbp, respectively for genome size; 18.2 and 26.2, respectively, for metabolic

pathways/categories).

We investigated the different kind of metabolisms present in MAGs based on site

occupancy patterns and found few patterns (Figure 6C, Table S4). We did observe a trend

towards MAGs present at two or three sites being more likely to have the capacity for nitrile

hydration (6%, 17%, or 11% of MAGs present at one, two, or three sites, respectively)

(Figure 6C). Nitrile hydration provides microorganisms with another avenue of acquiring

carbon and nitrogen needed for growth (Fang *et al.*, 2015). Notably, the two MAGs that were

present at all four sites lack many pathways/categories that were found in other MAGs, such

as ethanol fermentation, nitric and nitrous oxide reduction, urea utilization, hydrogenases,

and halogenated compound utilization, among others (Figure 6C), indicating that presence at

all four sites does not require metabolic generalism.

As we initially hypothesized, there was no observable trend towards metabolic

specialization, rather, relatively even numbers of metabolic specialists and generalists across

sites and site occupancy patterns (Figure 6C). Rather, there were relatively even numbers of

metabolic generalists (many types of pathways, aerobic/anaerobic, multiple energy sources,

etc.) and metabolic specialists between the different site occupancy patterns. One example of

a metabolic generalist, the MAG with the second largest number of metabolic pathways

present, Ritt_bin.7.strict_final (*Gammaproteobacteria*, genus *Paraburkholderia*), appears to

be capable of, among other things, urea utilization, thiosulfate and sulfur oxidation, nitrite

reduction to ammonia, fatty acid degradation, ethanol fermentation, some types of complex

carbon degradation and C1 metabolism, carbon fixation (via RuBisCo), and aerobic CO

oxidation, yet was only present at Rittmann (Figure 6C). On the other hand, we observed

multiple metabolic generalist MAGs that also had broad habitat ranges, such as Erebus-

WC_bin.32.strict_final (*Acidobacteriota*, order *Bryobacterales*) which was found at all sites

except TR and is capable of both organic and inorganic metabolisms, and both aerobic and

anaerobic metabolism (among other things, sulfur oxidation, nitrile hydration, nitrite

reduction to ammonia, iron oxidation, fatty acid degradation, ethanol fermentation, some

level of complex carbon degradation, chlorite reduction, and aerobic CO oxidation) (Figure

6C).

One factor that seems to be largely negating the trends towards metabolic generalism

in MAGs with higher site occupancy is the presence in our dataset of a large number of

archaeal MAGs with broad habitat ranges (present at 3+ sites) with few predicted metabolic

pathways and very small genomes (mean archaeal genome size: 2 Mbp). For example,

Erebus-TR_bin.23.permissive_final (class *Nitrososphaeria*), present at all four sites, has a

limited metabolic repertoire (genes for aerobic CO oxidation, chlorite reduction, nitrite

reduction, and sulfur oxidation), with mostly aerobic pathways (Figure 6C). Similarly,

Ritt_bin.9.permissive_final (class *Nitrososphaeria*, order *Conexivisphaerales*), present at all

sites except Melbourne, is capable of the same set of metabolic pathways minus nitrite

reduction (Figure 6C).

Discussion

In this research, we have characterized the microbial communities inhabiting remote,

isolated Antarctic geothermal sites and investigated the impacts of high dispersal rates on the

abundance and distribution of habitat specialists and generalists in a patchy environment. We

found that lower temperature, passive sites (WC, and Melbourne) share very similar

microbial communities that were dominated by *Plancomycetota* and *Acidobacteriota*, while

the hotter, active sites (TR, and Rittmann), each harbored distinct communities with high

abundances of *Nitrososphaeria*. We also discovered an absence of habitat generalists in our

data set, with no microbes thriving at all four sites, driven by high dispersal between sites and

high ecological drift. It should be noted that "habitat generalists" here refers to generalists

across Antarctic geothermal sites, which is different from "globally cosmopolitan" microbes.

We also note that only a single sample from each site was sequenced for metagenomics,

which may not be sufficient to capture the community heterogeneity present at these sites.

Site Characterizations

Both taxonomically (Figure 2, Figure S3, Figure S4, Figure S9A, B) and functionally

(Figure 3), the microbial communities at active and passive sites are distinct from each other.

The higher temperatures at active sites appear to be a major barrier preventing

microorganisms from passive sites colonizing active sites. This is apparent from the

significantly lower predicted optimal growth temperatures (OGT) of MAGs from passive vs.

active sites as well as that of MAGs from the Rittmann mat sample (32□) vs. the Rittman

surface sample (60□), which was collected directly underneath the Rittmann mat sample. It is

likely that the steep temperature gradient between the air (generally below -30□) and hot soil

(60-65□) is a large factor in facilitating microorganisms with lower temperature optima, such

as *Cyanobacteria* and *Bacteroidota*, to survive at hot spots at Rittmann and TR.[103,104] MAGs

present at both active sites and at least one passive site tended to have higher predicted OGTs

than all other MAGs (Figure 6B), which has been observed before for thermophiles (i.e.,

there is a positive correlation between maximal growth temperature and temperature growth

range (Miller, Williams, *et al.*, 2009; Weltzer and Miller, 2013)).

TR and Rittmann have additional, unique physicochemical properties (nutrient

availability and soil depth) that appear to influence the microbial community composition

present. The extensive phototrophic mats and moss beds present at TR that are not found at

the other sites(Skotnicki *et al.*, 2001) likely provide the complex carbohydrates required for

microorganisms from *Bacteroidota* to thrive there, despite their low predicted OGT (35□).

The high levels of total nitrogen (TN) at TR (likely, in part, due to the extensive moss beds

observed) could explain the presence of *Leptolyngbyaceae* there and its absence from other

sites, given the inability of MAGs from this family in our data set to fix nitrogen. Finally,

Rittmann has lower richness than other sites, both in terms of 16S rRNA gene sequence data

(Figure 2A) and number of MAGs binned (Rittmann was lowest at 15), perhaps due to the

shallow layer of topsoil present caused by the steep aspect. Shallow depth to bedrock has

been previously associated with lower bacterial counts and diversity (Cartwright and Hui,

2015; Cartwright *et al.*, 2016). Rittmann has previously been found to have a lower diversity

of *Cyanobacteria* than Melbourne and TR (Bargagli *et al.*, 1996).

It is challenging to put the functional characteristics of Antarctic geothermal

microbial communities in a global context, given the scarcity of metagenomics studies

examining microbial communities inhabiting geothermally heated soils (not hot spring

aqueous or sediment) in other parts of the world. However, compared to globally distributed

hot springs (Shu and Huang, 2021), we found Antarctic geothermal systems to be notably low

in sulphur and associated metabolisms (Figure S6D, Table S3). For instance, in Yellowstone

hot springs, genes associated with sulphur metabolism (sulfur and sulfate reduction, reduced

sulfur oxidation, and sulfite oxidation) were found to be among the most abundant metabolic

genes (Inskeep *et al.*, 2010), with similar results found in other geothermal systems globally

(Poddar and Das, 2018; Power *et al.*, 2018b; Prieto-Barajas *et al.*, 2018; Wilkins *et al.*, 2019).

Erebus has long been known to be a low-sulphur volcano, but the reason behind this

phenomenon have yet to be untangled (Sweeney *et al.*, 2008; Sims *et al.*, 2021). On the other

hand, we observed large abundances of hydrogenases and carbon monoxide dehydrogenase

genes at all our sites. Other geothermal systems globally have been observed to have

similarly high abundances of genes related to metabolizing geothermally produced gases,

especially hydrogen (Boyd *et al.*, 2010; Lindsay *et al.*, 2019; Shu and Huang, 2021).

Interestingly, non-geothermally heated soils in Antarctica are known to be populated by a

number of microbes that use trace gas (H_2, CO, and CO_2) metabolism for survival, forming

the base of the food chain along with photosynthesis, resulting in similarly high abundances

of these phenotypes as observed in our samples (Ji *et al.*, 2017), but with a different origin for

the gases (volcanic vs. atmospheric). In fact, some of the same, broad microbial groups

conducting trace gas metabolism in cold Antarctic soils are present in our samples (*Ca.*

Dormibacterota and *Ca.* Eremiobacterota).

Evidence of Dispersal Leading to High Habitat Specialization

We had hypothesized that high rates of dispersal between Antarctic geothermal sites

would result in a large number of shared microbes (habitat generalists) across all sites.

Although we did, indeed, find evidence of high dispersal between sites (25% of all variation

in our 16S rRNA gene sequence data can be explained by homogenous dispersal), we found

support for the opposite of our hypothesis: there were no habitat generalists between sites,

with the only shared microbial taxa present being clearly due to mass effects, not adaptation

to multiple sites. This lack of habitat generalists cannot be explained by environmental

selection, as variable environmental selection only accounted for 11% of community

differences between all sites and there were very few environmental parameters significantly

correlating with the overall biological community, and *Cyanobacteria* in particular (Figure

6C). Our results stand in stark contrast to past studies of thermophiles(Sharp *et al.*, 2014;

Power *et al.*, 2018a; Louca, 2022; DePoy and King, 2023) and habitat specialists(Pandit *et*

al., 2009; Xu *et al.*, 2022) that have found environmental selection to be the primary factor

influencing distribution, not dispersal, although one study did find a large influence of

dispersal on the abundant biosphere of Tibetan hot springs (Zhang *et al.*, 2018). We note that

many thermophilic habitat generalists have been identified before, even along strong

environmental gradients, e.g. (Weltzer and Miller, 2013; Wall *et al.*, 2014; He *et al.*, 2023;

Sriaporn *et al.*, 2023), so an absence of habitat generalists in our data set is surprising,

although many thermophiles tend to be habitat specialists for temperature (Miller, Strong, *et*

al., 2009; Mahato *et al.*, 2019; DePoy and King, 2023).

Our findings provide key insights into how microbial communities assemble, with

high dispersal rates and ecological drift leading to a lack of habitat generalists. These results

are similar to past macroecological studies that have found specialist species are favored

when dispersal is high and conditions are stable (i.e., low short-term habitat disturbance)

(Odum, 1969; Wilson and Yoshimura, 1994; Marvier *et al.*, 2004; Büchi and Vuilleumier,

2014; Koffel *et al.*, 2022). High dispersal rates allow specialist species to reach optimal

niches and outcompete generalist species. This does not preclude the existence of globally

cosmopolitan microbes at these sites, as has been postulated previously (Herbold, Lee, *et al.*,

2014), given that cosmopolitan species may be specialists for certain environments.

Additionally, highly stable conditions like those experienced at these sites (Soo *et al.*, 2009),

partially a result of their isolation and the absence of outside disturbances, results in few

opportunities for habitat generalists to take over niches vacated by specialists during a

disturbance. High environmental stability has been found in macroecology to result in a

selection towards K-strategists (lower growth rate, few offspring) (Odum, 1969), which may

be a contributing factor in the slow predicted growth rates of MAGs in our data set (average

across all four sites: 9.2 d^{-1}, fastest: 1.9 d^{-1}). The other factor influencing the lack of habitat

generalists in our data set is the large effect of ecological drift (30% of community assembly),

likely due to the small population size at these sites. In macroecology, ecological drift can

override niche effects in small populations, resulting in a co-occurrence of two species even

when one is a superior competitor (Orrock and Watling, 2010).

The high level of habitat specialization at these sites may also be indicative of high

levels of endemism, as these two have been found to be correlated in macroecological studies

(Cowling and Holmes, 1992; Ibarra and Martin, 2015). Antarctica is home to an unusually

high level of endemic organisms due to its geographic isolation (Vyverman *et al.*, 2010).

Tramway Ridge has previously been found to harbor a number of endemic microbes

(Herbold *et al.*, 2024). Additionally, the most abundant organism across all sites was a highly

divergent member of the class *Nitrososphaeria* that lacks genes for ammonia oxidation,

despite this class being primarily known for its ammonia oxidation activities. The functional

potential of this MAG has previously been inferred, with the proposed name *Candidatus*

Australlarchaeum erebusii (Herbold *et al.*, 2024). Despite its high abundance, this MAG

(Erebus-TR_bin.23.permissive_final) had the smallest genome size (1.4 mbp) in our data set.

Culturing efforts are needed to better understand the mechanisms by which this novel

archaeon is capable of wide habitat range despite limited metabolic flexibility, especially

given that *Archaea* are generally much more dispersal limited than *Bacteria* (Louca, 2022).

Our hope is that further microbial explorations at these unique sites, with broader sampling at

more areas and further metagenomic, transcriptomic, and culturing work, will help uncover

the wealth of novel microorganisms present.

Acknowledgments

The author(s) wish to acknowledge the contribution of NeSI to the results of this

research. New Zealand's national compute and analytics services and team are supported by

the New Zealand eScience Infrastructure (NeSI) and funded jointly by NeSI's collaborator

institutions and through the Ministry of Business, Innovation and Employment. URL

http://www.nesi.org.nz. We especially recognize the logistical support from Antarctica New

Zealand for fieldwork on Mt. Erebus and support in the field from Jon Tyler. We would like

to give a big thanks to Roanna Richards-Babbage for DNA extractions and preparing samples

for ICP-MS, TC/TN, and TOC/TON.

Data Availability

All R scripts used to analyze the data are available on Github at

https://github.com/ThermophileResearchUnit/Antarctic_volcano_comparison_manuscript.

16S rRNA gene sequence data has been deposited at DDBJ/EMBL/GenBank under the

accession KIGX00000000. The version described in this paper is the first version,

KIGX01000000. Metagenome reads are available on SRA under BioProject ID

PRJNA1113340.

Funding

This work was supported by the Royal Society of New Zealand (Marsden Grant 18-

UOW-028 to SC, MS, IM, and CL).

References

Allan, R.N., Lebbe, L., Heyrman, J., De Vos, P., Buchanan, C.J., and Logan, N.A. (2005)
Brevibacillus levickii sp. nov. and Aneurinibacillus terranovensis sp. nov., two novel
thermoacidophiles isolated from geothermal soils of northern Victoria Land, Antarctica.
Int J Syst Evol Microbiol **55**: 1039–1050.

Alneberg, J., Bjarnason, B.S., de Bruijn, I., Schirmer, M., Quick, J., Ijaz, U.Z., et al. (2014)
Binning metagenomic contigs by coverage and composition. *Nat Methods* **11**: 1144–
1146.

Andrews, S. FastQC.

Archer, S.D.J., Lee, K.C., Caruso, T., Maki, T., Lee, C.K., Cary, S.C., et al. (2019) Airborne
microbial transport limitation to isolated Antarctic soil habitats. *Nat Microbiol* **4**: 925–
932.

Barberán, A., Ramirez, K.S., Leff, J.W., Bradford, M.A., Wall, D.H., and Fierer, N. (2014)
Why are some microbes more ubiquitous than others? Predicting the habitat breadth of
soil bacteria. *Ecol Lett* **17**: 794–802.

Bargagli, R., Broady, P.A., and Walton, D.W.H. (1996) Preliminary investigation of the
thermal biosystem of Mount Rittmann fumaroles (northern Victoria Land, Antarctica).
Antarct Sci **8**: 121–126.

Bargagli, R., Skotnicki, M.L., Marri, L., Pepi, M., Mackenzie, A., and Agnorelli, C. (2004)
New record of moss and thermophilic bacteria species and physico-chemical properties
of geothermal soils on the northwest slope of Mt. Melbourne (Antarctica). *Polar Biol*
27: 423–431.

Bell, T.H. and Bell, T. (2021) Many roads to bacterial generalism. *FEMS Microbiol Ecol* **97**:
1–10.

Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Boyd, E.S., Hamilton, T.L., Spear, J.R., Lavin, M., and Peters, J.W. (2010) [FeFe]-
hydrogenase in Yellowstone National Park: evidence for dispersal limitation and
phylogenetic niche conservatism. *ISME J* **4**: 1485–1495.

Broady, P., Given, D., Greenfield, L., and Thompson, K. (1987) The biota and environment
of fumaroles on Mt Melbourne, northern Victoria Land. *Polar Biol* **7**: 97–113.

Brown, C.T. and Irber, L. (2016) sourmash: a library for MinHash sketching of DNA. *The*
Journal of Open Source Software **1**: 27.

Brown, J.H. (1984) On the Relationship between Abundance and Distribution of Species. *Am*
Nat **124**: 255–279.

Brown, J.S. and Pavlovic, N.B. (1992) Evolution in heterogeneous environments: Effects of
migration on habitat specialization. *Evol Ecol* **6**: 360–382.

Büchi, L. and Vuilleumier, S. (2014) Coexistence of specialist and generalist species is
shaped by dispersal and environmental factors. *American Naturalist* **183**: 612–624.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P.
(2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nat*
Methods **13**: 581–583.

Caporaso, J.G., Ackermann, G., Apprill, A., Bauer, M., Berg-Lyons, D., Betley, J., et al.
(2018) Earth microbiome project: EMP 16S Illumina Amplicon Protocol. *ProtocolsIo* 1–
7.

Cartwright, J., Dzantor, E.K., and Momen, B. (2016) Soil microbial community profiles and
functional diversity in limestone cedar glades. *Catena (Amst)* **147**: 216–224.

Cartwright, J. and Hui, D. (2015) Soil respiration patterns and controls in limestone cedar
glades. *Plant Soil* **389**: 157–169.

Chao, A., Chazdon, R.L., Colwell, R.K., and Shen, T.-J. (2004) A new statistical approach
for assessing similarity of species composition with incidence and abundance data. *Ecol*
Lett **8**: 148–159.

Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., and Parks, D.H. (2022) GTDB-Tk v2:
memory friendly classification with the genome taxonomy database. *Bioinformatics* **38**:
5315–5316.

Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018) fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**: i884–i890.

Chen, Y.J., Leung, P.M., Wood, J.L., Bay, S.K., Hugenholtz, P., Kessler, A.J., et al. (2021a)
Metabolic flexibility allows bacterial habitat generalists to become dominant in a
frequently disturbed ecosystem. *ISME Journal* **15**: 2986–3004.

Chen, Y.J., Leung, P.M., Wood, J.L., Bay, S.K., Hugenholtz, P., Kessler, A.J., et al. (2021b)
Metabolic flexibility allows bacterial habitat generalists to become dominant in a
frequently disturbed ecosystem. *ISME Journal* **15**: 2986–3004.

Cowling, R.M. and Holmes, P.M. (1992) Endemism and speciation in a lowland flora from
the Cape Floristic Region. *Biological Journal of the Linnean Society* **47**: 367–383.

Darling, A.E., Jospin, G., Lowe, E., Matsen, F.A., Bik, H.M., and Eisen, J.A. (2014)
PhyloSift: phylogenetic analysis of genomes and metagenomes. *PeerJ* **2**: e243.

DePoy, A.N. and King, G.M. (2023) Distribution and diversity of anaerobic thermophiles and
putative anaerobic nickel-dependent carbon monoxide-oxidizing thermophiles in
mesothermal soils and sediments. *Front Microbiol* **13**: 1–14.

Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016) MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**: 3047–3048.

Fang, S., An, X., Liu, H., Cheng, Y., Hou, N., Feng, L., et al. (2015) Enzymatic degradation
of aliphatic nitriles by Rhodococcus rhodochrous BX2, a versatile nitrile-degrading
bacterium. *Bioresour Technol* **185**: 28–34.

Fernandes￼Martins, M.C., Colman, D.R., and Boyd, E.S. (2023) Relationships between fluid
mixing, biodiversity, and chemosynthetic primary productivity in Yellowstone hot
springs. *Environ Microbiol* **25**: 1022–1040.

Fierer, N. and Jackson, R.B. (2006) The diversity and biogeography of soil bacterial
communities. *Proceedings of the National Academy of Sciences* **103**: 626–631.

Finn, D.R., Yu, J., Ilhan, Z.E., Fernandes, V.M.C., Penton, C.R., Krajmalnik-Brown, R., et al.
(2020) MicroNiche: An R package for assessing microbial niche breadth and overlap
from amplicon sequencing data. *FEMS Microbiol Ecol* **96**: 1–12.

Fraser, C.I., Terauds, A., Smellie, J., Convey, P., and Chown, S.L. (2014) Geothermal
activity helps life survive glacial cycles. *Proc Natl Acad Sci U S A* **111**: 5634–5639.

Gao, C. (2021) ggVennDiagram: A "ggplot2" Implement of Venn Diagram.

Geyer, A. (2021) Antarctic volcanism: Active volcanism overview. *Geological Society*
Memoir **55**: 55–72.

Gong, X., del Río, Á.R., Xu, L., Chen, Z., Langwig, M. V., Su, L., et al. (2022) New globally
distributed bacterial phyla within the FCB superphylum. *Nat Commun* **13**:.

Green, H.C., Fisher, J.C., McLellan, S.L., Sogin, M.L., and Shanks, O.C. (2016)
Identification of specialists and abundance-occupancy relationships among intestinal
bacteria of Aves, Mammalia, and Actinopterygii. *Appl Environ Microbiol* **82**: 1496–
1503.

Green, J.L., Holmes, A.J., Westoby, M., Oliver, I., Briscoe, D., Dangerfield, M., et al. (2004)
Spatial scaling of microbial eukaryote diversity. *Nature* **432**: 747–750.

Greening, C., Biswas, A., Carere, C.R., Jackson, C.J., Taylor, M.C., Stott, M.B., et al. (2016)
Genomic and metagenomic surveys of hydrogenase distribution indicate H2 is a widely
utilised energy source for microbial growth and survival. *ISME J* **10**: 761–777.

Greening, C., Geier, R., Wang, C., Woods, L.C., Morales, S.E., McDonald, M.J., et al. (2019)
Diverse hydrogen production and consumption pathways influence methane production
in ruminants. *ISME J* **13**: 2617–2632.

He, Q., Wang, S., Feng, K., Michaletz, S.T., Hou, W., Zhang, W., et al. (2023) High
speciation rate of niche specialists in hot springs. *ISME J*.

Herbold, C.W., Lee, C.K., McDonald, I.R., and Cary, S.C. (2014) Evidence of global-scale
aeolian dispersal and endemism in isolated geothermal microbial communities of
Antarctica. *Nat Commun* **5**:.

Herbold, C.W., Mcdonald, I.R., and Cary, S.C. (2014) Microbial Ecology of Geothermal
Habitats in Antarctica. In *Antarctic Terrestrial Microbiology: Physical and Biological*
Properties of Antarctic Soils. Cowan, D. (ed). Berlin, Heidelberg: Springer, pp. 1–328.

Herbold, C.W., Noell, S.E., Lee, C.K., Vickers, C.J., Stott, M.B., Eisen, J.A., et al. (2024)
Nutritional niches of endemic, aerobic heterotrophs from an isolated Antarctic terrestrial
hydrothermal refugium elucidated through metagenomics. *in review*.

Hernandez, D.J., Kiesewetter, K.N., Almeida, B.K., Revillini, D., and Afkhami, M.E. (2023)
Multidimensional specialization and generalization are pervasive in soil prokaryotes.
Nat Ecol Evol.

Hijmans, R.J. (2022) geosphere: Spherical Trigonometry.

Hughes, K.A., McCartney, H.A., Lachlan-Cope, T.A., and Pearce, D.A. (2004) A preliminary
study of airborne microbial biodiversity over Peninsular Antarctica. *Cell Mol Biol*
(Noisy-le-grand) **50**: 537–42.

Hyatt, D., Chen, G.L., LoCascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010)
Prodigal: Prokaryotic gene recognition and translation initiation site identification. *BMC*
Bioinformatics **11**:.

Hyatt, D., LoCascio, P.F., Hauser, L.J., and Uberbacher, E.C. (2012) Gene and translation
initiation site prediction in metagenomic sequences. *Bioinformatics* **28**: 2223–2230.

Ibarra, J.T. and Martin, K. (2015) Beyond species richness: an empirical test of top predators
as surrogates for functional diversity and endemism. *Ecosphere* **6**: art142.

Ilanko, T., Fischer, T.P., Kyle, P., Curtis, A., Lee, H., and Sano, Y. (2019) Modification of
fumarolic gases by the ice-covered edifice of Erebus volcano, Antarctica. *Journal of*
Volcanology and Geothermal Research **381**: 119–139.

Inskeep, W.P., Rusch, D.B., Jay, Z.J., Herrgard, M.J., Kozubal, M.A., Richardson, T.H., et al.
(2010) Metagenomes from High-Temperature Chemotrophic Systems Reveal
Geochemical Controls on Microbial Community Structure and Function. *PLoS One* **5**:
e9773.

Ji, M., Greening, C., Vanwonterghem, I., Carere, C.R., Bay, S.K., Steen, J.A., et al. (2017)
Atmospheric trace gases support primary production in Antarctic desert surface soil.
Nature **552**: 400–403.

Jiang, X., Van Horn, D.J., Okie, J.G., Buelow, H.N., Schwartz, E., Colman, D.R., et al.
(2022) Limits to the three domains of life: lessons from community assembly along an
Antarctic salinity gradient. *Extremophiles* **26**: 15.

Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., and Wang, Z. (2019) MetaBAT
2: an adaptive binning algorithm for robust and efficient genome reconstruction from
metagenome assemblies. *PeerJ* **7**: e7359.

Kassambara, A. (2023) ggpubr: "ggplot2" Based Publication Ready Plots.

Koffel, T., Umemura, K., Litchman, E., and Klausmeier, C.A. (2022) A general framework
for species☐abundance distributions: Linking traits and dispersal to explain
commonness and rarity. *Ecol Lett* **25**: 2359–2371.

Kolde, R. (2019) pheatmap: Pretty Heatmaps.

Koslicki, D., White, S., Ma, C., and Novikov, A. (2023) YACHT: an ANI-based statistical
test to detect microbial presence/absence in a metagenomic sample. *bioRxiv*.

Lahti, L. and Shetty, S. (2017) Tools for microbiome analysis in R.

Lee, M.D. (2019) GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* **35**:
4162–4164.

Li, G., Rabe, K.S., Nielsen, J., and Engqvist, M.K.M. (2019) Machine Learning Applied to
Predicting Microorganism Growth Temperatures and Enzyme Catalytic Optima. *ACS*
Synth Biol **8**: 1411–1420.

Liao, H., Liu, C., Ai, C., Gao, T., Yang, Q.-E., Yu, Z., et al. (2023) Mesophilic and
thermophilic viruses are associated with nutrient cycling during hyperthermophilic
composting. *ISME J* **17**: 916–930.

Lindsay, M.R., Colman, D.R., Amenabar, M.J., Fristad, K.E., Fecteau, K.M., Debes, R. V., et
al. (2019) Probing the geological source and biological fate of hydrogen in Yellowstone
hot springs. *Environ Microbiol* **21**: 3816–3830.

Liu, C., Cui, Y., Li, X., and Yao, M. (2021) microeco☐: an R package for data mining in
microbial community ecology. *FEMS Microbiol Ecol* **97**:.

Locey, K.J., Muscarella, M.E., Larsen, M.L., Bray, S.R., Jones, S.E., and Lennon, J.T. (2020)
Dormancy dampens the microbial distance–decay relationship. *Philosophical*
Transactions of the Royal Society B: Biological Sciences **375**: 20190243.

Logan, N.A., Lebbe, L., Hoste, B., Goris, J., Forsyth, G., Heyndrickx, M., et al. (2000)
Aerobic endospore-forming bacteria from geothermal environments in northern Victoria
Land, Antarctica, and Candlemas Island, South Sandwich archipelago, with the proposal
of Bacillus fumarioli sp. nov. *Int J Syst Evol Microbiol* **50**: 1741–1753.

Logares, R., Lindström, E.S., Langenheder, S., Logue, J.B., Paterson, H., Laybourn-Parry, J.,
et al. (2013) Biogeography of bacterial communities exposed to progressive long-term
environmental change. *ISME J* **7**: 937–948.

Louca, S. (2022) The rates of global bacterial and archaeal dispersal. *ISME Journal* **16**: 159–
167.

Magurran, A.E. and Henderson, P.A. (2003) Explaining the excess of rare species in natural
species abundance distributions. *Nature* **422**: 714–716.

Mahato, N.K., Sharma, A., Singh, Y., and Lal, R. (2019) Comparative metagenomic analyses
of a high-altitude Himalayan geothermal spring revealed temperature-constrained
habitat-specific microbial community and metabolic dynamics. *Arch Microbiol* **201**:
377–388.

Malard, L.A. and Guisan, A. (2023) Into the microbial niche. *Trends Ecol Evol* **xx**: 1–10.

Management Plan For Antarctic Specially Protected Area No . 175 (2014) **13**: 199–225.

Marvier, M., Kareiva, P., and Neubert, M.G. (2004) Habitat Destruction, Fragmentation, and
Disturbance Promote Invasion by Habitat Generalists in a Multispecies Metapopulation.
Risk Analysis **24**: 869–878.

von Meijenfeldt, F.A.B., Hogeweg, P., and Dutilh, B.E. (2023) A social niche breadth score
reveals niche range strategies of generalists and specialists. *Nat Ecol Evol*.

Miller, S.R., Strong, A.L., Jones, K.L., and Ungerer, M.C. (2009) Bar-coded pyrosequencing
reveals shared bacterial community properties along the temperature gradients of two
alkaline hot springs in Yellowstone National Park. *Appl Environ Microbiol* **75**: 4565–
4572.

Miller, S.R., Williams, C., Strong, A.L., and Carvey, D. (2009) Ecological Specialization in a
Spatially Structured Population of the Thermophilic Cyanobacterium Mastigocladus
laminosus. *Appl Environ Microbiol* **75**: 729–734.

Myeong, N.R., Choe, Y.-H., Shin, S.C., Kim, J., Sul, W.J., and Kim, M. (2024) Genomic
profiling of Antarctic geothermal microbiomes using long-read, Hi-C, and single-cell
techniques. *Sci Data* **11**: 1023.

Nayfach, S., Shi, Z.J., Seshadri, R., Pollard, K.S., and Kyrpides, N.C. (2019) New insights
from uncultivated genomes of the global human gut microbiome. *Nature* **568**: 505–510.

Niederberger, T.D., Bottos, E.M., Sohm, J.A., Gunderson, T., Parker, A., Coyne, K.J., et al.
(2019) Rapid microbial dynamics in response to an induced wetting event in Antarctic
dry valley soils. *Front Microbiol* **10**: 1–12.

Noell, S.E., Baptista, M.S., Smith, E., McDonald, I.R., Lee, C.K., Stott, M.B., et al. (2022)
Unique Geothermal Chemistry Shapes Microbial Communities on Mt. Erebus,
Antarctica. *Front Microbiol* **13**:.

Nurk, S., Meleshko, D., Korobeynikov, A., and Pevzner, P.A. (2017) metaSPAdes: a new
versatile metagenomic assembler. *Genome Res* **27**: 824–834.

Odum, E.P. (1969) The Strategy of Ecosystem Development: An understanding of ecological
succession provides a basis for resolving man's conflict with nature. *Science (1979)* **164**:
262–270.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2020)
vegan: Community Ecology Package.

Orrock, J.L. and Watling, J.I. (2010) Local community size mediates ecological drift and
competition in metacommunities. *Proceedings of the Royal Society B: Biological*
Sciences **277**: 2185–2191.

Pandit, S.N., Kolasa, J., and Cottenie, K. (2009) Contrasts between habitat generalists and
specialists: An empirical extension to the basic metacommunity framework. *Ecology* **90**:
2253–2262.

Parish, T.R. (1988) Surface winds over the Antarctic continent: A review. *Reviews of*
Geophysics **26**: 169.

Park, M., Choe, Y.-H., and Kim, M. (2021) Soil Metagenome Sequences from a Geothermal
Site at Mount Melbourne in Antarctica. *Microbiol Resour Announc* **10**: 7–8.

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., and Tyson, G.W. (2015)
CheckM: assessing the quality of microbial genomes recovered from isolates, single
cells, and metagenomes. *Genome Res* **25**: 1043–1055.

Pearce, D.A., Hughes, K.A., Lachlan-Cope, T., Harangozo, S.A., and Jones, A.E. (2010)
Biodiversity of air-borne microorganisms at Halley station, Antarctica. *Extremophiles*
14: 145–159.

Pepi, M., Agnorelli, C., and Bargagli, R. (2005) Iron demand by thermophilic and mesophilic
bacteria isolated from an antarctic geothermal soil. *BioMetals* **18**: 529–536.

Poddar, A. and Das, S.K. (2018) Microbiological studies of hot springs in India: a review.
Arch Microbiol **200**: 1–18.

Poli, A., Esposito, E., Lama, L., Orlando, P., Nicolaus, G., de Appolonia, F., et al. (2006)
Anoxybacillus amylolyticus sp. nov., a thermophilic amylase producing bacterium
isolated from Mount Rittmann (Antarctica). *Syst Appl Microbiol* **29**: 300–307.

Power, J.F., Carere, C.R., Lee, C.K., Wakerley, G.L.J., Evans, D.W., Button, M., et al.
(2018a) Microbial biogeography of 925 geothermal springs in New Zealand. *Nat*
Commun **9**:.

Power, J.F., Carere, C.R., Lee, C.K., Wakerley, G.L.J., Evans, D.W., Button, M., et al.
(2018b) Microbial biogeography of 925 geothermal springs in New Zealand. *Nat*
Commun **9**:.

Power, J.F., Lowe, C.L., Carere, C.R., McDonald, I.R., Cary, S.C., and Stott, M.B. (2023)
Temporal dynamics of geothermal microbial communities in Aotearoa-New Zealand.
Front Microbiol **14**:.

Prieto-Barajas, C.M., Alcaraz, L.D., Valencia-Cantero, E., and Santoyo, G. (2018) Life in
Hot Spring Microbial Mats Located in the Trans-Mexican Volcanic Belt: A 16S/18S
rRNA Gene and Metagenomic Analysis. *Geomicrobiol J* **35**: 704–712.

R Core Team (2020) R: A language and environment for statistical computing.

Rodriguez-R, L.M., Gunturu, S., Tiedje, J.M., Cole, J.R., and Konstantinidis, K.T. (2018)
Nonpareil 3: Fast Estimation of Metagenomic Coverage and Sequence Diversity.
mSystems **3**:.

Rodriguez-R, L.M. and Konstantinidis, K.T. (2014) Nonpareil: a redundancy-based approach
to assess the level of coverage in metagenomic datasets. *Bioinformatics* **30**: 629–635.

Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., et al. (2010)
Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* **4**:
1340–1351.

Schiro, G., Chen, Y., Blankinship, J.C., and Barberán, A. (2022) Ride the dust: linking dust
dispersal and spatial distribution of microorganisms across an arid landscape. *Environ
Microbiol* **24**: 4094–4107.

Schliep, K.P. (2011) phangorn: phylogenetic analysis in R. *Bioinformatics* **27**: 592–593.

Schmieder, R. and Edwards, R. (2011) Quality control and preprocessing of metagenomic
datasets. *Bioinformatics* **27**: 863–864.

Seemann, T. (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**: 2068–
2069.

Sharp, C.E., Brady, A.L., Sharp, G.H., Grasby, S.E., Stott, M.B., and Dunfield, P.F. (2014)
Humboldt's spa: Microbial diversity is controlled by temperature in geothermal
environments. *ISME Journal* **8**: 1166–1174.

Shu, W. and Huang, L. (2021) Microbial diversity in extreme environments. *Nat Rev
Microbiol*.

Sims, K.W.W., Aster, R.C., Gaetani, G., Blichert-Toft, J., Phillips, E.H., Wallace, P.J., et al.
(2021) Chapter 7.2 Mount Erebus. *Geological Society, London, Memoirs* **55**: 695–739.

Skotnicki, M.L., Selkirk, P.M., Broady, P., Adam, K.D., and Ninham, J.A. (2001) Dispersal
of the moss Campylopus pyriformis on geothermal ground near the summits of Mount
Erebus and Mount Melbourne, Victoria Land, Antarctica. *Antarct Sci* **13**: 280–285.

Soo, R.M., Skennerton, C.T., Sekiguchi, Y., Imelfort, M., Paech, S.J., Dennis, P.G., et al.
(2014) An Expanded Genomic Representation of the Phylum Cyanobacteria. *Genome
Biol Evol* **6**: 1031–1045.

Soo, R.M., Wood, S.A., Grzymski, J.J., McDonald, I.R., and Cary, S.C. (2009) Microbial
biodiversity of thermophilic communities in hot mineral soils of Tramway Ridge, Mount
Erebus, Antarctica. *Environ Microbiol* **11**: 715–728.

Sriaporn, C., Campbell, K.A., Van Kranendonk, M.J., and Handley, K.M. (2023) Bacterial
and archaeal community distributions and cosmopolitanism across physicochemically
diverse hot springs. *ISME Communications* **3**: 80.

Sriswasdi, S., Yang, C.C., and Iwasaki, W. (2017) Generalist species drive microbial
dispersion and evolution. *Nat Commun* **8**:.

Stegen, J.C., Lin, X., Fredrickson, J.K., Chen, X., Kennedy, D.W., Murray, C.J., et al. (2013)
Quantifying community assembly processes and identifying features that impose them.
ISME Journal **7**: 2069–2079.

Sweeney, D., Kyle, P.R., and Oppenheimer, C. (2008) Sulfur dioxide emissions and
degassing behavior of Erebus volcano, Antarctica. *Journal of Volcanology and
Geothermal Research* **177**: 725–733.

Ul-Hasan, S., Bowers, R.M., Figueroa-Montiel, A., Licea-Navarro, A.F., Beman, J.M.,
Woyke, T., and Nobile, C.J. (2019) Community ecology across bacteria, archaea and
microbial eukaryotes in the sediment and seawater of coastal Puerto Nuevo, Baja
California. *PLoS One* **14**: e0212355.

Uritskiy, G. V., DiRuggiero, J., and Taylor, J. (2018) MetaWRAP—a flexible pipeline for
genome-resolved metagenomic data analysis. *Microbiome* **6**: 158.

Vyverman, W., Verleyen, E., Wilmotte, A., Hodgson, D.A., Willems, A., Peeters, K., et al.
(2010) Evidence for widespread endemism among Antarctic micro-organisms. *Polar Sci*
4: 103–113.

Wall, C.A., Koniges, G.J., and Miller, S.R. (2014) Divergence with gene flow in a population
of thermophilic bacteria: a potential role for spatially varying selection. *Mol Ecol* **23**:
3371–3383.

Weissman, J.L., Hou, S., and Fuhrman, J.A. (2021) Estimating maximal microbial growth
rates from cultures, metagenomes, and single cells via codon usage patterns. *Proc Natl
Acad Sci U S A* **118**: 1–10.

Weltzer, M.L. and Miller, S.R. (2013) Ecological divergence of a novel group of
Chloroflexus strains along a geothermal gradient. *Appl Environ Microbiol* **79**: 1353–
1358.

Whitaker, R.J., Grogan, D.W., and Taylor, J.W. (2003) Geographic Barriers Isolate Endemic
Populations of Hyperthermophilic Archaea. *Science (1979)* **301**: 976–978.

Whiteley, A.S., Jenkins, S., Waite, I., Kresoje, N., Payne, H., Mullan, B., et al. (2012)
Microbial 16S rRNA Ion Tag and community metagenome sequencing using the Ion
Torrent (PGM) Platform. *J Microbiol Methods* **91**: 80–88.

Wickham, H. (2016) Ggplot2: Elegant Graphics For Data Analysis, New York: Springer-
Verlag.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., et al. (2019)
Welcome to the Tidyverse. *J Open Source Softw* **4**: 1686.

Wilkins, L.G.E., Ettinger, C.L., Jospin, G., and Eisen, J.A. (2019) Metagenome-assembled
genomes provide new insight into the microbial diversity of two thermal pools in
Kamchatka, Russia. *Sci Rep* **9**: 3059.

Wilson, D.S. and Yoshimura, J. (1994) On the Coexistence of Specialists and Generalists. *Am*
Nat **144**: 692–707.

Wright, E.S. (2016) Using DECIPHER v2.0 to analyze big biological sequence data in R. *R*
Journal **8**: 352–359.

Wu, Y.-W., Simmons, B.A., and Singer, S.W. (2016) MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**:
605–607.

Xu, Q., Luo, G., Guo, J., Xiao, Y., Zhang, F., Guo, S., et al. (2022) Microbial generalist or
specialist: Intraspecific variation and dormancy potential matter. *Mol Ecol* **31**: 161–173.

Zhang, Y., Wu, G., Jiang, H., Yang, J., She, W., Khan, I., and Li, W. (2018) Abundant and
rare microbial biospheres respond differently to environmental and spatial factors in
tibetan hot springs. *Front Microbiol* **9**: 1–16.

Zhang, Z., Zhang, L., Zhao, Z., Wang, H., and Ju, F. (2022) Deduplication Improves Cost-
Efficiency and Yields of De novo Assembly and Binning of
Shot-Gun Metagenomes in Microbiome Research. *bioRxiv* 2022.10.12.512008.

Zhou, Z., Tran, P.Q., Breister, A.M., Liu, Y., Kieft, K., Cowley, E.S., et al. (2022)
METABOLIC: high-throughput profiling of microbial genomes for functional traits,
metabolism, biogeochemistry, and community-scale functional networks. *Microbiome*
10: 1–22.

Zhu, J., Ma, Y., Huang, L., and Zhang, W. (2022) Homogeneous selection is not always
important in bacterial community in the eutrophic enclosed bay. *Ecol Process* **11**: 27.

1147

 Figure Legends

Figure 1 Map and images of sampling sites. (A) Satellite image of Antarctica (top right inset,

red circle indicates Victoria Land); main image shows Victoria Land and Ross Island. (B)

Aerial picture of Mt. Erebus main crater. (C) Picture of the hot soil site sampled at Western

Crater on Mt. Erebus. (D) Satellite image showing the summit of Mt. Erebus and the two

sampling sites in this study, Tramway Ridge (-77.519444, 167.116389 DD) and Western

Crater (-77.519167, 167.187778 DD). (E) Image of sampling site on Tramway Ridge. (F)

Image of sampling on the side of Mt. Rittmann (-73.448445, 165.497444 DD). (G) Image of

Mt. Rittmann. (H) Image of Mt. Melbourne. (I) Image of sampling at a hot soil site on Mt.

Melbourne (-74.350016, 164.700139 DD). Satellite image in (A) is from the Antarctic Digital

Database Map Viewer https://www.add.scar.org/, Open Source, using hillshade and

bathymetry layers. Satellite image in (D) was purchased from DigitalGlobe Incorporated,

Longmont CO USA (2019). All other images were taken by Ian McDonald and S. Craig

Cary.

Figure 2 Diversity and composition of microbial communities across the four sites. (A) Alpha

diversity of microbial communities based on 16S rRNA gene sequences across the four sites

(Tramway Ridge (TR), Western Crater (WC), Melbourne, or Rittmann, n = 8, 6, 4, or 3

respectively) using the Shannon index of diversity. Paired significance test used an

undirected Wilcox test. (B) Principal coordinates analysis (PCoA) of the microbial

communities based on Unifrac similarities between 16S rRNA gene sequences. The

percentages in brackets for each axis indicate the percentage of the variation in community

composition explained along that axis. (C) Microbial community composition at the phylum

level for both 16S rRNA gene sequencing samples (16S) and metagenome samples at the

read level (Metagenome reads); singleM was used for determining taxonomy for the

metagenome reads. For the 16S rRNA gene sequence data, the sequencing run used for

Tramway Ridge (TR) and Western Crater (WC) samples are denoted by the last digit, either 1

for the first (data from (Noell *et al.*, 2022)) or second sequencing run. For this plot, phylum

names for metagenome reads were changed to match SILVA 16S rRNA gene taxonomy to

allow for ease of comparison between the two methods (*Thermoproteota* to *Crenarchaeota*,

Chloroflexota to *Chloroflexi*, CSP1-3 to GAL15, *Dormibacterota* to *Chloroflexi* (class AD3),

Eremiobacterota to WPS-2). (D, E) Results of a differential abundance random forest

analysis to identify taxa at any level that are most important for distinguishing between (D)

active and passive sites, or (E) all four sites in the 16S rRNA gene sequence data. The plots

on the left give the mean decrease in Gini value, which is a measure of the importance of that

taxa to distinguishing the sites. The plot on the right shows the relative abundance of those

taxa at the respective sites. A threshold of either 0.2 or 0.1 relative abundance was used for

plotting ASVs for the active vs. passive comparison or all four sites comparison, respectively.

Figure 3 Genomic and functional characteristics of MAGs and metagenomes from each site.

(A, B) Comparison of genome characteristics for MAGs recovered from different site

categories (A) or sites (B). (A) n = 32 or 79 for active or passive, respectively (*p*-values: two-

sided Wilcox test). (B) n = 19, 37, 42, or 13 for Tramway Ridge (TR), Western Crater (WC),

Melbourne, or Rittmann, respectively. p-values given for the pairwise comparisons are for

two-sided Wilcox tests. (C) Analysis of the functional potential of the microbial communities

at the four site samples with metagenomes, based on the abundance of marker genes for

different metabolic pathways. Abundance within community was calculated using RPKM

values, which were normalized by the abundance of single copy marker genes within each

metagenome. TR: Tramway Ridge. WC: Western Crater.

Figure 4 Analysis of Antarctic geothermal *Cyanobacteria* diversity and factors shaping it.

Abundance heatmaps of *Cyanobacteria* communities using (A) 16S rRNA gene sequence and

(B) metagenome assembled genome (MAG) data. For (A), relative abundances of

Cyanobacteria families are presented as relative to the abundance of other Cyanobacteria.

For (B), MAG abundances were marked as zero if the MAG was not present at the site. Then,

the abundances were normalized: $\log_{10}(\text{abundance} + 0.001) + 4$. (C) Correlation heatmap,

using the Pearson coefficient, of the cyanobacterial families to the relevant environmental

variables (selected for having Mantel test $p < 0.05$). GWC: gravitational water content; TN:

total nitrogen.

Figure 5 Ecological predictions for microorganisms at Antarctic geothermal sites. (A-C)

Frequency plots identifying the number of either specialists or generalists for (A) habitat

(using Levin's niche breadth, B_n), (B) temperature (using Hulbert's niche breadth, B_n), or (C)

pH (using Hulbert's niche breadth, B_n) in the 16S rRNA gene amplicon sequence data.

Arrows point to generalists where difficult to visualize. (D) Contribution of different

ecological factors to shaping the community assembly of microorganisms in the 16S rRNA

gene amplicon data. (E, F, G) Distance-decay plots for the 16S rRNA gene amplicon data

with a linear regression best fit model for (E) all samples, (F) only samples from Active sites,

or (G) only samples from Passive sites.

Figure 6 (A-B) Exploring evidence for genomic markers of metabolic generalism in

metagenome assembled genomes (MAGs) from the four sites. For (A), n = 35, 52, 27, or 2

for the categories 1 site, 2 sites, 3 sites, or 4 sites, respectively. p-values for ANOVA tests

conducted in (A) are, from top left clockwise order: 0.08, 0.07, 0.04, and 0.02. p-value shown

on plot is the result of an undirect Wilcox test, Benjamini-Hochburg corrected. No

differences for pfam domains remained significant after BH correction. For (B) and (C), n =

11, 2, or 7 for Tramway Ridge (TR) only, Rittmann only, or TR/Rittmann only MAGs,

respectively; n = 5, 3, 8, 15, or 2 for TR/Western Crater (WC) only, WC/Rittmann only,

TR/WC/Rittmann only, WC/Melbourne/Rittmann only, or MAGs at all four sites,

respectively; n = 10, 11, or 36 for WC only, Melbourne only, or WC/Melbourne only MAGs,

respectively. For (B), * represents a statistically significant ($p < 0.05$) result of an undirected

Wilcox test after Benjamini-Hochburg correction. See Methods for how Metabolic

Pathways/Categories were marked as present or absent in MAGs. (C) Heatmap showing

functional predictions for MAGs based on METABOLIC predictions, separated by site

occupancy (number and identity of sites occupied) and whether the metabolic predictions are

categories (containing multiple, non-sequential metabolic reactions) or pathways. MAGs that

are specifically mentioned in the text are indicated by an arrow. Pink boxes differentiate

different site occupancy patterns. X-axis MAGs are clustered first by site occupancy pattern,

then alphabetically by Phylum-Class. TR: Tramway Ridge. WC: Western Crater. RT:

Rittmann. ML: Melbourne.

B

C

A

Mt. Erebus

D

0.5 km

Western
Crater

Tramway
Ridge

Mt. Melbourne

Mt. Rittmann

100 km

50 mi

N

E

F

G

H

I